Long-read sequencing to interrogate strain-level variation among

adherent-invasive *Escherichia coli* isolated from human intestinal tissue

Jeremy Wang[1,¶], Rachel Bleich[2,¶], Sandra Zarmer[3], Shiying Zhang[4], Belgin Dogan[4], Kenneth W

Simpson[4], Janelle Arthur[2,5,6]*

[1]Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[2]Department of Microbiology & Immunology, University of North Carolina at Chapel Hill,

Chapel Hill, NC, USA

[3]Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC,

USA

[4]Department of Clinical Sciences, College of Veterinary Medicine, Cornell University, Ithaca,

New York, USA.

[5]Center for Gastrointestinal Biology & Disease, University of North Carolina at Chapel Hill,

Chapel Hill, NC, USA

[6]Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel

Hill, NC, USA

[¶] These authors contributed equally to this work.

* Corresponding author

Email: janelle_arthur@med.unc.edu

Abstract

Adherent-invasive *Escherichia coli* (AIEC) is a pathovar linked to inflammatory bowel diseases

(IBD), especially Crohn's disease, and colorectal cancer. AIEC are genetically diverse, and in

the absence of a universal molecular signature, are defined by *in vitro* functional attributes. The

relative ability of difference AIEC strains to colonize, persist, and induce inflammation in an

IBD-susceptible host is unresolved. To evaluate strain-level variation among tissue-associated *E.*

coli in the intestines, we develop a long-read sequencing approach to identify AIEC by strain that

excludes host DNA. We use this approach to distinguish genetically similar strains and assess

their fitness in colonizing the intestine. Here we have assembled complete genomes using long-

read nanopore sequencing for a model AIEC strain, NC101, and seven strains isolated from the

intestinal mucosa of Crohn's disease and non-Crohn's tissues. We show these strains can

colonize the intestine of IBD susceptible mice and induce inflammatory cytokines from cultured

macrophages. We demonstrate that these strains can be quantified and distinguished in the

presence of 99.5% mammalian DNA and from within a fecal population. Analysis of global

genomic structure and specific sequence variation within the ribosomal RNA operon provides a

framework for efficiently tracking strain-level variation of closely-related *E. coli* and likely other

commensal/pathogenic bacteria impacting intestinal inflammation in experimental settings and

IBD patients.

Introduction

Crohn's disease (CD) is a type of Inflammatory bowel disease (IBD), a chronic inflammatory

condition that is the result of an inappropriate immune response towards elements of the

intestinal microbiota [1]. Due to chronic inflammation and exposure to pro-carcinogenic

microbes, CD patients are at a greater risk of developing colorectal cancer (CRC) [2-4]. CD

patients harbor microbiomes that differ from non-CD individuals in both composition and

function, with reduced diversity and an increase of adherent/invasive *E. coli* (AIEC) [5-11].

AIEC are closely associated with the intestinal mucosa, termed the mucosal niche, and induce

inflammation in murine models of IBD, including the well-established, inflammation-susceptibe

Il10$^{-/-}$ mouse model [3, 12-14]. AIEC strains lack genes associated with virulence in

diarrheagenic *E. coli* and are defined through the *in vitro* ability to adhere to and invade

epithelial cells and survive in macrophages [5, 7, 15]. However, as AIEC are genetically diverse,

there is no universal genomic feature that distinguishes the pathotype [16], limiting our ability to

detect these pro-inflammatory strains from among patient microbiota.

Escherichia coli make up a diverse species that is known to variously have beneficial,

commensal, or pathogenic impact in the mammalian gut. However, most commonly used

methods for characterizing the composition of gut microbiota from fecal or tissue samples,

notably sequencing of selected variable regions of the 16S rRNA operon, do not differentiate

among these closely-related but functionally divergent types. Emerging single-molecule (third-

generation/long-read) sequencing technologies including Pacific Biosciences ("PacBio") and

Oxford Nanopore Technologies ("nanopore") produce reads dozens to hundreds of kilobases in

length, often without prior amplification. Recent work [17-20] has demonstrated the utility of

long reads for sequencing the entire 16S gene or entire rRNA operon, with corresponding

increase in taxonomic resolution by capturing more variable sequence, including all nine variable

regions of 16S, the internal transcribed spacers (ITS), and 23S. Traditional metataxonomic

analysis using next-generation sequencing of one or two adjacent hypervariable regions of the

16S operon are unable to discriminate among closely-related species and genera [21], much less

capture strain-level variation. We propose a full-length rRNA sequencing approach – enabled by

longer reads produced by nanopore sequencing – that is capable of discriminating among

closely-related strains of *E. coli*. To support the extension of these approaches to characterize

strain-level variation of tissue-associated (mucosa-associated) microbiota contributing to IBD

and in models of experimental colitis in mice, with a particular focus on mechanistic studies of

AIEC, we produced complete genome assemblies for eight AIEC and non-AIEC *E. coli*, describe

the genomic variation among these strains, particularly within the rRNA operon, and

demonstrate the accurate identification of these strains in mixed *in vitro* and *in vivo* microbiota.

Methods

DNA extraction and nanopore sequencing

We extracted ultra-high-molecular-weight genomic DNA from liquid *E. coli* cultures using a

modified phenol:chloroform protocol [22]. Briefly, 1-1.5 ml of stationary-phase liquid culture

was pelleted and resuspended in TE/NaCl/Triton buffer. Cells were lysed by addition of SDS to a

final concentration of 2% and Proteinase K to a final concentration of 300 mg/ml and incubated

at 55°C for 30 minutes. DNA was purified twice by addition of 1x volume phenol:chloroform

and phase separation. DNA was precipitated by addition of 0.1x volume sodium acetate and 2x

volume isopropanol. Ultra-high-molecular weight DNA was "hooked" out with a melted glass

capillary and washed in 70% ethanol, dried, and resuspended in nuclease-free water (NFW) by

incubation overnight at 4°C. Multiplexed sequencing libraries were prepared for nanopore

sequencing using Oxford Nanopore's Ligation Sequencing Kit (LSK109) and Barcoding

Expansion (NBD104) per the manufacturer's recommended protocol with the following

modifications [23]. Instead of SPRI bead cleanup after each ligation reaction, we added 1 volume

of "clumping buffer" (9% PEG 8000, 1 M NaCl, 10 mM Tris), incubated at room temperature

for 30 mins, pelleted by centrifugation at 13K rpm, washed twice with 70% ethanol, then

allowed to dry and resuspended in NFW by incubating at 4°C overnight. This "bead-free"

cleanup both avoids shearing and loss of high-molecular-weight DNA and selects for larger

fragments in the PEG precipitations. Multiplexed samples were sequenced on R9.4.1 flow cells

on either MinION or GridION with real-time basecalling using Guppy v3.2.2 in high-accuracy

mode. The distribution of read lengths is shown in Figure 1.

Fig 1. Read length distribution across samples using high-molecular weight extraction and

bead-free library prep.

Illumina sequencing

Strains NC101 and HM670 were sequenced by the Microbial Whole Genome Sequencing Center

(www.migscenter.com) on the NextSeq 2000 platform, yielding paired end reads (2x151bp)

delivered to us as fastq files. Strains denoted "CU" were Illumina whole-genome sequenced by

Enterome (Paris, Ile-de-France, France).

Genome assembly

Basecalled reads for each sample were assembled using Miniasm v0.2 [24], followed by four

rounds of polishing with Racon v1.3.1 [25], and Medaka v0.10.0 [26]. Each assembly consisted

of one contig representing the full-length genome. Assembled genomes were normalized to start

at the origin of replication and re-polished across the previous breakpoint. Assemblies were

further polished with Illumina sequencing data using Pilon v1.22-1 [27]. Annotation was

performed using the NCBI Prokaryotic Genome Annotation Pipeline [PGAP release 2020-02-

06.build4373; 28]. Predicted genes were subsequently assigned putative function by aligning to

the RefSeq non-redundant protein database with Diamond v0.9.22 [29] allowing for frame-shift

sensitive alignment (diamond blastp --more-sensitive --frameshift 15). Sequence data and

assembled genomes have been made publicly available under NCBI BioProject

ID PRJNA759208.

Serotype, MLST, virulence, and antibiotic resistance genes

Polished assemblies were aligned against EcOH database [30] for serotyping, multi-locus

sequence typing (MLST), and VirulenceFinder databases [31], and NCBI antimicrobial

resistance database using Minimap2 [32] with parameter '-cx asm5' to allow for indel-sensitive

alignment appropriate for hybrid-assembled genomes [33]. For serotype and MLST, the single

best-aligning type gene (minimum edit distance) is given as the predicted type. Predicted

serotypes, MLSTs, AMR, and virulence genes were further validated using corresponding web-

based tools provided by the Center for Genomic Epidemiology ([31],

http://www.genomicepidemiology.org/services/), which showed complete agreement.

rRNA operon analysis

All copies of the rRNA operon in each assembly were identified by aligning to the *E. coli* K12

reference rRNA using Minimap2 [32]. Variants relative to the K12 reference were identified for

each assembled rRNA copy across all assemblies based on the alignments. Total polymorphism

distance was computed between every pair of alignments across each region of the rRNA operon

(16S, ITS, 23S).

Bacterial Growth Conditions

Bacteria were grown in Luria broth (LB) at 37°C and 250 rpm unless otherwise indicated.

Animal care

Germ-free mice were reared in the National Gnotobiotic Rodent Resource Center at UNC Chapel

Hill. All animal experiments and procedures were approved by UNC's Institutional Animal Care

and Use Committee (IACUC).

Murine stool samples

To generate an in vivo gut microbial community containing the seven *E. coli* strains of interest

and endogenous mouse microbiota, we inoculated mice as follows. Three (2m/1f) interleukin-10-

deficient (*Il10$^{-/-}$*) mice (129Sv/Ev background) were reared germ-free to adulthood (8-10) weeks

and colonzied by oral gavage with an even mixture of a total of 10^7 CFU clinical *E. coli* strains

isolated from the intestinal tissue of Crohn's disease and non-Crohn's disease patients [7]. All

strains have been classified as AIEC and non-AIEC using standard *in vitro* assays to evaluate

adhesion/invasion to Caco2 epithelial cells and uptake/survival in J774 macrophages [5,7]. To

validate our results by tracking individual strains with PCR, each strain was marked with an

antibiotic resistance cassette and molecular barcode inserted into a neutral chromosomal region

using Tn7 transposon insertion [34-35]. During experimentation, these strains were referenced

by their blinded laboratory designation and barcode: JA0018/A1, JA0019/A3, JA0022/B6,

JA0036/C5, JA0044/D2, JA0048/D5, and JA0091/C2. The identity of these strains is described

in Table 1. Five strains were originally isolated by KW Simpson at Cornell University as the

strains CU39ES-1, CU532-9, CU568-3, CU37RT-2, CU42ET-1; HM670 was isolated previously

[6] and gifted by Barry Campbell from Liverpool University; and LF82 is the prototypic AIEC

strain gifted to KW Simpson by Arlette Darfeuille-Michaud [5].

After colonization, mice were maintained in specific pathogen free (SPF) housing, where they

acquired a simplified mouse microbiota. After 2 weeks of colonization, we gave kanamycin

water *ad libetum* for 2 weeks to suppress this microbiota and ensure that all strains could persist

to some extent. Mice were sacrificed 6 weeks later (by CO_2 asphyxiation), for a total of 10 weeks

colonization. A stool sample was removed from the lumen of the distal colon for our analysis.

DNA was extracted and purified as described in [3,36].

Strain designation	Phylogroup	AIEC	Tissue origin	Blinded strain code	Barcode
CU39ES-1	B2	N	Ileal Crohn's disease	JA0018	A1
CU532-9	B2	N	Ileal Crohn's disease	JA019	A3
LF82	B2	Y	Ileal Crohn's disease [5]	JA0022	B6
CU568-3	B2	Y	Colonic Crohn's disease	JA0036	C5
CU37RT-2	B2	N	non-IBD	JA0044	D2
CU42ET-1	B2	Y	non-IBD	JA0048	D5
HM670	B2	N	Colon cancer [6]	JA0091	C2
NC101	B2	Y	WT Murine stool [3,12]	JA0058	-

Table 1: Strains used in study.

In vivo colonization and persistence studies

To demonstrate that the seven human-derived strains and murine *E. coli* NC101 could colonize

germ-free *Il10*$^{-/-}$ mice throughout the gastrointestinal tract, we singly housed 1 male and 1 female

mouse per *E. coli* strain, for a total of 16 cages. Mice were gavaged with 10^8 CFU of a single

strain and moved to SPF housing. After 1 week, each mouse received a fecal transplant (prepared

anaerobically from a pool of 7 C57BL/6 WT mice) via gavage to provide niche competition.

Stool samples were collected almost daily and CFUs were quantified on kanamycin plates to

monitor fecal *E. coli* colonization. Mice were sacrificed by CO_2 asphyxiation after 5 weeks, and

the following tissues were harvested and CFUs quantified on kanamycin plates to measure viable

colonizing bacteria: ileal tissue, cecal content, colon content, colon tissue and colon mucus layer.

E. coli CFUs were also quantified from stomach content, duodenal content, and jejunal content,

but only 1-3 samples of each tissue had detectable bacterial growth of more than 10^2 CFUs/10

mg.

188   To demonstrate that consistent colonizing strain, CU42ET-1-D5, and inconsistent

colonizing strain, HM670-C2, colonized germ-free *Il10*$^{-/-}$ mice in a similar manner across a

larger cohort, we housed 7 (3M/4F) or 8 (4M/4F) mice, respectively, in two cages for each

cohort. Mice were gavaged with 10^8 CFU of a single strain and moved to SPF housing. After 1

week, each mouse received the same fecal transplant as used previously (prepared anaerobically

from a pool of 7 C57BL/6 WT mice that were *Helicobacter spp.* free) via gavage to provide

niche competition. After 3 weeks of colonization (2 weeks post-FMT), we administered

kanamycin water *ad libetum* for 2 weeks to suppress this microbiota and ensure that all strains

could persist to some extent. Mice were sacrificed 6 weeks later (by CO_2 asphyxiation), for a

total of 10 weeks colonization. The following tissues were harvested and CFUs quantified on

kanamycin plates to measure viable colonizing bacteria: colon content, colon tissue and colon

mucus layer.

In vitro co-culture assays to measure pro-inflammatory cytokine production

To assess the inflammatory capacity of the clinical strains, J774 macrophages grown in DMEM

complete media (supplemented with 10% heat-inactivated FBS and 100 U/ml penicillin-

100 mg/ml streptomycin) were transferred to 12-well plates at $5x10^5$ cells/well. Cells were

grown overnight at 5% CO_2 at 37°C, and washed twice with PBS before adding fresh culture

media without antibiotics. Cells were infected at an MOI of 1 for 4 hours before removing the

bacteria, washing with PBS, and adding fresh media with 100 μg/mL gentamycin. Cells were

grown another 20 hours at which point the supernatants were collected for ELISA and the cells

were collected for qPCR analysis.

ELISA

Supernatants from stimulated J774 macrophages (above) were analyzed for cytokine IL12p40

production through ELISA following manufacturer's protocol (BD Bioscience Opt EIA, catalog

#555165). They were assayed in 4 independent experiments with 2-3 technical replicate wells.

Quantification of cytokine expression by qPCR

Stimulated J774 macrophages were washed with PBS and transferred into 1 mL Trizol

(Invitrogen) and RNA was extracted following the manufacturer's protocol. Isolated RNA was

subjected to DnaseI treatment (Invitrogen) prior to cDNA synthesis. cDNA synthesis was

completed using qScript cDNA SuperMix (Quantabio). qPCR amplification was performed in

221 triplicate with SYBR green qPCR chemistry (Bioline) using primers for *Tnfa* (F-5'-

222 ACCTCACACTCAGATCATCTTCTC-3', R-5'-TGAGATCCATGCCGTTGG-3'), *Il1b* (F-5'-

223 ACAGAATATCAACCAACAAGTGATATTCTC-3', R-5'-GATTCTTTCCTTTGAGGCCCA-

224 3'), *Ifnγ* (F-5'-CGGCACAGTCATTGAAAGCC-3', R-5'-TGTCACCATCCTTTTGCCAGT-3'),

225 *Csf2* (F-5'-ATGCCTGTCACGTTGAATGAAG-3', R-5'-GCGGGTCTGCACACATGTTA-3'),

226 *Il12b(p40)* (F-5'-CGCAAGAAAGAAAAGATGAAGGAG-3', R-5'-

227 TTGCATTGGACTTCGGTAGATG-3'), and *Gapdh* (F-5'-

228 GGTGAAGGTCGGAGTCAACGGA-3', R-5'-GAGGGATCTCGCTCCTGGAAGA-3') on a

229 QuantStudio 6 Real-Time PCR System. C_t values were normalized to *Gapdh* to generate ΔC_t

230 values, and fold changes were calculated by $\Delta\Delta C_t$ to the ΔC_t of unstimulated controls. These

231 were assayed in 3 independent experiments in technical triplicates.

233 *Quantification of strains by qPCR*

234 Stool DNA (6 ng each) from three mice was subjected to qPCR with primer pairs targeting each

235 barcode to quantify relative amounts of each barcoded strain from within a complex population.

236 Amplification was performed in duplicate using SYBR green qPCR chemistry (Bioline) using a

237 universal barcode forward primer (F-5'-GCTTGGTTAGAATGGGTAACTAGTTTGCAG-3')

238 and barcode-specific reverse primers for A1 (R-5'-TTCCCGAGCGCACCACAAA-3'), A3 (R-5'-

239 ACACATACATCTCGCACGCAAACG-3'), B6 (R-5'-AAACCAACATCTCCCTCGCCC-3'),

240 C2 (R-5'-GGTGATGTTTGGGCGTGGTAGAA-3'), C5 (R-5'-

241 ATAACACTCCCGCCCACGAGAA-3'), D2 (R-5'-TTCGAACTCGACCGCCAACCAAA-3'),

242 and D5 (R-5'-CCACTCAATCACGCAACACCC-3') with *E. coli* 16S primers (F-5'-

243 ATTGACGTTACCCGCAGAAGA-3', R-5'-GGGATTTCACATCCGACTTGA-3') [34-35] on a

QuantStudio 6 Real-Time PCR System. C_t values were normalized to *E. coli* 16S rRNA to

generate ΔC_t values, and fold changes were calculated by $\Delta\Delta C_t$ to the ΔC_t of the mouse

inoculum.

Mock tissue-associated microbiome

Aliquots of purified genomic DNA isolated from independent strain cultures were mixed at equal

abundance by weight. HMW DNA was isolated from a frozen aliquot of 10^7 HEK293 cells using

the Circulomics Nanobind kit (https://www.circulomics.com/nanobind) per the manufacturer's

recommended protocol. Human DNA was mixed with microbial mixture at a ratio of 99.5 : 0.5%

to represent a realistic tissue-associated microbiota composition.

Full-length rRNA amplicon sequencing

For stool and *in vitro* host/microbiota samples, primers for proximal 16S (27F:

AGRGTTTGATYHTGGCTCAG) and distal 23S (2241R: ACCRCCCCAGTHAAACT) were

used to amplify the full-length rRNA operon (4,500bp). Starting with ~5 ng expected microbial

DNA, 25 ul PCR reactions were prepared with LongAmp Taq 2x Master Mix (New England

Biolabs, Inc; M0287L) and 0.4 uM of each primer. We ran 20 cycles consisting of denaturation

at 94°C for 10s, annealing at 51°C for 30s, and extension at 65°C for 225s, followed by final

extension at 65°C for 10min. Typical yield is 700ng of full-length amplicons, which were

verified by agarose gel electrophoresis and fluorescent quantification (Qubit). Amplicon libraries

were prepared for nanopore sequencing using the Oxford Nanopore Ligation Sequencing Kit

(LSK109) per the manufacturer's protocol and sequenced on R9.4.1 flow cell on GridION with

real-time basecalling using Guppy v3.2.2 in high-accuracy mode.

Results

Genome assembly and annotation

To establish a baseline for variation among known enteric *E. coli*, we produced finished genomes

for eight AIEC and non-AIEC strains (NC101 [3,12], CU39ES-1-A1, CU532-9-A3, CU568-3-

C5, CU37RT-2-D2, CU42ET-1-D5, HM670-C2, and LF82-B6). These genomes were

sequenced, assembled, polished, and annotated as described in the Methods. Sequence data and

assembled genomes have been made publicly available under NCBI BioProject

ID PRJNA759208. Each genome was assembled into a single complete, circular contig

representing the entire genome (Table 2). Table 3 describes the serotype (O and H), multi-locus

sequence type (MLST), and virulence and antibiotic genes found in each. Figure 2 illustrates

these eight genomes, annotated genes, and approximate homology between them.

Strain	Genome size	Genes
HM670-C2	5,141,716	4,762
CU39ES-1-A1	5,124,020	4,630
CU532-9-A3	4,858,260	4,433
LF82-B6	4,772,884	4,399
CU568-3-C5	5,024,072	4,681
CU37RT-2-D2	5,015,239	4,632
CU42ET-1-D5	4,903,963	4,438
NC101	5,030,087	4,630

282 *Table 2. Genome size and number of genes identified in assembled E. coli genomes.*

Strain	Serotype	MLST	Virulence genes	Antibiotic resistance
HM670-C2	O2:H1	73	clbB*, pic*, iha, yfcV*, sat, ompT*, kpsMII, sitA, usp, fyuA*, papA_F43*, vat, tcpC, terC*, papA_F7-1, gad, iucC*, iutA, kpsE, chuA, iss, papC*, irp2, ireA	mdf(A)*, sitABCD*
CU39ES-1-A1	O25:H1	73	clbB, papA_F14, kpsMII_K5, mchB, mchC*, mchF, iha, yfcV*, sat, ompT*, mcmA, sitA, usp, fyuA*, iutA, hra, vat*, pic, cea, terC*, focI, iroN, gad, iucC, focG*, kpsE, focCsfaE, chuA*, iss, papA_F43*, hlyA, papC*, irp2	mdf(A)*, sitABCD*
CU532-9-A3	O4:H5	12	ompT*, clbB, chuA, iss, sitA, usp*, terC*, tcpC, yfcV*, irp2, fyuA*, gad, vat*, kpsE*	mdf(A)*, sitABCD*
LF82-B6	O83:H1	135	ompT*, kpsE*, chuA, iss, sitA, usp*, terC*, ibeA, yfcV*, irp2*, fyuA*, kpsMII_K5, gad, vat	mdf(A)*, sitABCD*
CU568-3-C5	O25:H4	131	ompT*, kpsE, gad, chuA, iss, sitA*, usp*, terC*, iutA, papA_F43*, irp2, fyuA*, kpsMII_K5, iha, iucC, yfcV*, sat	mdf(A)*, sitABCD*
CU37RT-2-D2	O16:H6	144	ompT*, kpsE, gad*, chuA*, sitA*, usp*, terC*, papA_F7-1*, iutA, neuC, tcpC, kpsMII_K1, papC*, yfcV*, irp2*, fyuA*, iha*, iucC*, vat*, sat*	mdf(A)*, sitABCD*
CU42ET-1-D5	O99:H6	1859	papA_F12*, kpsMII_K1*, mchB, mchC*, mchF*, ibeA*, yfcV*, ompT*, sitA*, usp*, fyuA*, hra, vat*, pic*, terC*, focI*, sfaE*, iroN*, gad*, focG*, kpsE, chuA*, papC*, neuC, irp2*	mdf(A)*, sitABCD*
NC101	O2:H6	998	clbB, kpsMII_K1, pic*, ibeA, yfcV*, ompT*, sitA, usp*, fyuA*, vat, tcpC, sfaS, terC*, focI*, iroN*, gad*, kpsE, focCsfaE*, chuA, iss, neuC ,irp2	mdf(A)*, sitABCD*

Table 3. Characterization of each strain by serotype and MLST, including virulence, and antibiotic resistance genes.

Fig 2. Assembled genomes, illustrating relative GC content (red) and gene content (blue) for

each strain, and approximate synteny between assemblies represented in gray, where darker

bars represent higher identity (darkest gray is ~100% identity, white is <70%).

Strain-level variation

As illustrated in Figure 2, we observe significant genome-wide structural variation while broadly

conserving GC content and annotated gene density within homologous blocks, even among these

closely-related strains with similar adherent-invasive (AI) or non-AI phenotype. We computed

the pairwise hamming distance [37] between sequences across each strain and each copy of the

rRNA operon for the 16S, ITS, 23S, and entire rRNA region (Figure 3).

Fig 3. Difference between 16S (a), 23S (b), ITS (c), and entire rRNA (d) sequences across strains

and copies. Heatmap shows the relative number of discriminating sites (alleles) between

sequences. Sequences are clustered hierarchically using Ward's variance minimization and are

labeled by strain.

We evaluated several regions independently for informative variation among these strains *in*

silico by aligning our whole-genome sequence to either 1) the V1-V2 hypervariable region of

16S, 2) V3-V4 hypervariable region, 3) the entire 16S sequence containing all nine hypervariable

regions and conserved spacers, 4) the entire rRNA operon, including 16S, ITS, and 23S, and 5)

the entire genome. Very long nanopore reads allow us to use this as a proxy for PCR

amplification of the various rRNA amplicon analyses since these reads average 10-20 Kbp (see

Table 4), and thereby most often cover the entire region of interest.

Strain	# reads	Total Gbp	Average read length	N50
HM670-C2	129,071	3.26	25,252	53,682
CU39ES-1-A1	33,058	1.30	39,369	76,514
CU532-9-A3	94,599	1.46	15,384	54,555
LF82-B6	126,340	3.19	25,252	55,067
CU568-3-C5	239,866	3.73	15,547	48,832
CU37RT-2-D2	61,344	2.14	34,912	58,392
CU42ET-1-D5	112,086	2.90	25,871	56,055
NC101	22,886	0.44	19,341	78,877

Table 4. Nanopore sequencing statistics for each strain, after demultiplexing.

In general, and not unexpectedly, the larger the region used for classification, the greater the

accuracy in detecting the correct strain (Table 5). Hypervariable regions V1-V2 (27F-338R) and

V3-V4 (343F-806R) produces poor results (averaging 35% and 27% accuracy, respectively),

little better than chance since there is little or no discriminating variation within that region. The

entire 16S gene performs better (70%). Notably, there is a huge amount of variation in the

accuracy across strains owing to their relative dissimilarity in a particular region. Full-length

rRNA achieves an average 87.3% accuracy at the strain level, followed closely by all reads

aligned against the entire genome (i.e. shotgun metagenomics) at 91%. Additional benefits of

whole-metagenome sequencing include the observation of genes that may imply the functional

capacity of the community without explicitly characterizing its taxonomic structure, and the

ability to assemble the metagenome into an approximation of its component microbial genomes.

These are important, tangible benefits, but are beyond the scope of this paper.

Strain	V1-V2	V3-V4	16S	rRNA	Genomic
HM670-C2	14.48	29.66	61.95	71.75	74.85
CU39ES-1-A1	14.72	12.50	47.39	70.73	84.79
CU532-9-A3	17.52	19.61	96.93	93.44	90.44
LF82-B6	93.18	18.66	97.62	87.07	94.78
CU568-3-C5	89.62	87.39	99.06	99.19	97.09
CU37RT-2-D2	14.68	31.25	59.16	96.83	97.57
CU42ET-1-D5	17.60	2.90	55.26	99.07	96.13
NC101	16.67	10.77	45.71	80.18	93.43
Average	*34.81*	*26.59*	*70.39*	*87.28*	*91.14*

Table 5. Classification accuracy using commonly amplified 16S hypervariable regions, V1-V2

and V3-V4, the entire 16S and rRNA operons, and shotgun sequencing of the entire genome.

While short hypervariable regions perform poorly and inconsistently across these samples,

classification based on the entire rRNA operon (~4,500bp) approaches classification accuracy of

whole-genome sequencing.

There is a relatively small improvement in classification accuracy using the entire genomes

compared to the full-length rRNA operon. Like commonly used 16S primers that target subsets

of the hypervariable regions, the primer pair used for the full operon are well-conserved across

bacteria, but presents a much larger sequence with which to accurately classify sequences. This

coupled with cost-effective multiplexing and long-read sequencing, despite much higher error

rates than standard Illumina sequencing, should make this a viable approach for characterizing

complex microbiome samples.

In vivo colonization studies

We performed *in vivo* colonization studies using a well-established IBD mouse model [3, 12-13,

36, 38-39]. We first determined if these human-derived strains and murine-derived NC101 could

colonize and persist in mice in the presence of a competing microbiota. We colonized two

singly-housed, adult, germ-free, *Il10*[-/-] mice each with a single *E. coli* strain, then after one week

provided niche competition by gavaging with a murine fecal transplant as outlined in Figure S2.

After five weeks total, gastrointestinal tissues, including ileal tissue, cecal content, colon content,

colon tissue, and colon mucus layer, were harvested and viable *E. coli* were quantified by serial

plating. Table S1 reveals that almost all *E. coli* strains could colonize the lumen and intestinal

tissue with complex community competition, even without kanamycin to supress the competing

microbiota. Two strains of *E. coli* colonized inconsistently across both mice, with high levels of

E. coli in one mouse and low or undetected levels of *E. coli* in the other. To verify if this would

persist across a larger cohort, consistent colonizing strain, CU37RT-2-D5, and inconsistent

colonizing strain, HM670-C2, were given by gavage to seven or eight germ-free, *Il10*[-/-] mice,

respectively. After one week we provided niche competition by gavaging with a murine fecal

transplant, and then after three weeks we provided kanamycin water *ad libitum* for two weeks to

supress the competing microbiota as outlined in Figure S3. After ten weeks total, gastrointestinal

tissues, including colon content, colon tissue, and colon mucus layer, were harvested and viable

E. coli were quantified by serial plating. Table S2 reveals that consistent colonizing strain

CU37RT-2-D5 colonized all seven mice to high levels, while inconsistent colonizing strain

HM670-C2 colonized five of eight mice at the lumen and intestinal tissue. Future studies are

needed to validate these results and draw strong conclusions, including colonization in inflamed

and non-inflamed environments.

To validate the pro-inflammatory potential of these strains, each strain was co-cultured with J774

macrophages and inflammatory colitis-inducing cytokine transcription and secretion was

measured [40-43]. The results in Figure 4 demonstrate that all strains induced inflammatory

cytokine production. Inducing inflammatory cytokine production is an indicator of inflammatory

potential, but it is not a known predictor of tissue inflammation or colonization *in vivo*. Further

effort is needed to compare *in vitro* cytokine production with *in vivo* inflammation elicited by

colonizing AIEC isolates.

Figure 4: Inflammatory cytokine production by macrophages stimulated with clinical E. coli

strains. (a-e) Relative mRNA abundance of each cytokine from stimulated J774 macrophages

determined by qRT- PCR. Values normalized to gapdh, and fold increase relative to

unstimulated controls. Each bar represents the mean across 3 independent experiments in

*triplicate and error bars show standard deviation (*p<0.05, Tukey's multiple comparisons test).*

If not otherwise noted, there is no significance. (f) Protein abundance of IL12(p40) from

stimulated J774 macrophages determined by ELISA. Values normalized to NC101 stimulated

macrophages. Each bar represents the mean across 4 independent experiments in duplicate and

*error bars show standard deviation (*p<0.05, Tukey's multiple comparisons test). If not*

otherwise noted, there is no significance

Full-length rRNA metataxonomics

To further evaluate the utility of performing strain-level metataxonomics using full-length rRNA

sequencing on an Oxford Nanopore device, we prepared and sequenced one stool sample from

in vivo Il10$^{-/-}$ colonization studies depicted in Figure 5. *Il10$^{-/-}$* mice, a well-established model of

microbially driven intestinal inflammation, were colonized with an equally-proportioned pool of the

seven *E. coli* clinical isolates, and then housed in an SPF facility where they become colonized over

time with a natural mouse microbiome [3, 36]. After ten weeks, the mice were sacrificed and

stool samples were collected for nanopore sequencing and quantification of each strain by

targeted qPCR. We extracted DNA from the resulting stool samples. Full-length rRNA amplification

and sequencing using universal 16S 27F and 23S 2241R primers (see Methods) produced

>600,000 reads with a median length of 4,149 bp (Table 6, Figure 6). We assigned these reads to

individual rRNA copies by aligning to our database of known sequences in the assembled

genomes (40 copies, five in each of eight strains). Figure 7 illustrates the distribution of

alignment identity (number of matching nucleotides) over the full-length rRNA (4,245bp).

Sequences derived from the stool sample show a bimodal distribution with peaks at 3.9Kbp

(92% identity) and 2.9Kbp (68% identity), representing the expected divergence and sequencing

error matching *E. coli* and non-*Escherichia* genera, respectively. The proportion of reads

assigned to each strain is shown in Figure 5(b). There is a skewed representation of strains in the

stool sample due to natural variation in their ability to colonize and thrive and due to interstrain

competition between isolates in the *Il10*$^{-/-}$ mouse gut. We confirmed the relative abundance of

each strain in the stool sample through qPCR of the molecular barcode inserted in the neutral

Tn7 site of each *E. coli* genome [34]. The qPCR showed a similar result, as there was skewing

among the *E. coli* isolates in the mouse gut as shown in Figure 5 with the CU568-3-C5 strain

most abundant. Both consistent and inconsistent colonizing strains showed lower abundance in

this stool sample as there were multiple *E. coli* isolates providing competition upon initial

infection, and this data is a measure of relative abundance across *E. coli* strains and not total load

with the stool. Additionally, the inconsistent colonizing strain still colonized in 5 of 8 mice

(Table S2).

Sample	# reads	Total bp	Average read length	Median read length	Read N50
Mouse stool sample	657,234	2.78 Gbp	4,229.65	4,149	4,194
0.5% *E.coli* mix, 99.5% HEK293	240,691	0.92 Gbp	3,829.50	4,136	4,173

Table 6. Sequencing statistics for full-length rRNA amplicon sequencing.

Figure 5. Strain distribution of seven E. coli isolates in murine stool sample. (a) Timeline for

mouse experiment. (b) Relative percent abundance of each strain determined by full-length rRNA

sequencing. Off-target hits to NC101 (4%) are expected to happen by chance given the

classification error rate for the whole rRNA sequence (~13%, see Table 5). (c) Relative percent

abundance of each strain determined by quantitative PCR of molecular barcode.   Values

normalized to E. coli *16S rRNA, and fold increase relative to pooled* E. coli *inoculum.   Each*

bar represents the mean across 3 murine stool samples in duplicate and error bars show

standard deviation.

Figure 6. Read length distribution for stool and mock mixture full-length rRNA amplicon

sequencing. The secondary peak at ~8Kbp represents chimeric reads resulting from nanopore

sequencing that are separated prior to alignment.

Figure 7. Distribution of nucleotide matches of the best alignment from amplified rRNA

sequencing. Match/Mismatch represent the aligned rRNA segments from the barcoded whole-

genome sequencing where we know the ground truth. Mismatches have a nearly identical

distribution to matches, indicating sequencing errors caused miscategorization as a very closely-

related rRNA. Mock shows a very similar distribution, but sequences from stool are bimodal

indicating a (smaller) proportion of reads originated from these or similar E.coli rRNA, and a

majority (~75%) originating from highly divergent sequences (70-80% identity) likely

originating from non-Escherichia genera and phyla.

We additionally prepared a mock mixture consisting of all eight strains at even abundance mixed

0.5 : 99.5% with human DNA (HEK293) to simulate a realistic tissue-associated microbiota

sample. Full-length rRNA sequences were successfully amplified and sequenced as done with

the stool sample. We observed relatively uniform representation of the eight strains in the mock

mixture, demonstrating our ability to amplify and properly classify closely-related rRNA

sequences in a host-tissue-like context, as illustrated in Figure 8.

Figure 8. Strain representation based on full-length rRNA sequencing from mock host:microbial

mixture. Mock mixture contained an approximately uniform quantify of all eight strains before

amplification.

Discussion

Alterations in the composition and function of gut microbial communities are implicated in both

IBD and CRC, with CD patient microbiomes harboring an increase in mucosa-associated

Enterobacteriaceae, including AIEC [7]. After excluding genes associated with diarrheagenic *E.*

coli, AIEC are defined by their *in vitro* adherent-invasive behavior. There is a clear need to

extend our knowledge of AIEC to pathoadaptive determinants of colonization and virulence *in*

vivo. Recent work highlights mechanisms AIEC have evolved to more effectively colonize the

host mucosa including hypermotility and metabolic changes [44] and T4SS-linked biofilm

formation [45]. These efforts involved either long *in vivo* host adaptation experiments or

genome-wide screens with a limited number of *E. coli* strains. Other molecular mechanisms

including the type 1 pili-CEACAM6 interaction that promotes adhesion of AIEC to intestinal

epithelial cells [46-47] do not offer a genetic marker to distinguish AIEC. Competitive

colonization with pools of known AIEC and non-AIEC strains as a parallel strategy could help

elucidate traits and genes important for colonization, but requires methods to discriminate

between closely-related microbes. To relate specific traits of different AIEC to pre-clinical

models and patients, we need to be able to distinguish individual strains from within a complex

community. As AIEC are mucosa-associated, the ability to evaluate colonization of inflamed or

cancerous lesions and normal mucosa would be optimal.

The commonly used metataxonomic approach of sequencing one or two amplified

hypervariable regions of the 16S gene is unable to distinguish among closely-related, but

functionally diverse, species and strains [21]. Consistent with previous work [17-21], we

demonstrate that sequencing of longer regions of the 16S, ITS, and 23S operons, enabled by

modern long-read sequencing technologies, dramatically increases taxonomic specificity. We

illustrated the increase in sensitivity and specificity of strain detection using full-length rRNA

amplification and whole-metagenome shotgun sequencing in *in silico* models, mock mixtures,

and *in vivo* colonized murine microbiome .

This focused study highlights specifically the challenges in resolving closely-related but

functionally divergent AIEC and non-adherent/invasive *E. coli* strains within mixed microbial

populations. We demonstrated accurate identification and discrimination of these strains in

complex *in vitro* and *in vivo* models relevant to colitis. Our novel nanopore sequencing approach

illustrates an efficient and effective method for discrimination of known pathogenic or

specifically colonized microbiota, including tissue-associated bacteria, to enable investigation of

spatiotemporal dynamics of microbial colonization without relying on genetic engineering,

molecular barcoding [34], or *in vivo* fluorescence markers [48].

While these results suggest a method for sensitive identification of known strains, this

approach does not suggest a general taxonomic analysis protocol for microbiome studies since

the genomes of the strains of interest must be known *a priori* in order to properly identify them.

Full-length rRNA sequencing following PCR amplification permits accurate strain identification

even when the microbial abundance - for example in tissue-associated microbial samples - is

very low. We demonstrated high concordance among relative abundance between sequenced

full-length rRNA sequences and qPCR results targeting the inserted molecular barcodes,

suggesting that abundances are reliably preserved through amplification and nanopore

sequencing. However, even higher accuracy and explicit identification of microbial genic content

can be captured by shotgun metagenome sequencing when there exists enough material to extract

high-quality microbial DNA.

Future work includes optimization of the full-length rRNA protocol for tissue samples

with adherent microbes. Our analysis of *in vitro* mixtures indicate this should be feasible with

microbial:host DNA content as low as 0.5% - consistent with expected abundances [49-50] in

many human mucosa, including the gut - but we have so far failed to amplify full-length rRNA

from resected mouse gut. Additionally, given the extraordinary utility of shotgun metagenomic

sequence, it is worthwhile to explore the several existing methods for selective extraction and

sequencing of microbial DNA to expand the availability of non-PCR-based approaches in the

context of tissue-associated microbial communities.

Acknowledgements

This work was supported by the following grants: NIH/NIDDK K01 DK119582 (JW),

SPIRE (NIH IRACDA program) NIH/NIGMS K12 GM000678-21 (RMB), NIH/NIDDK K01

DK103952 (JCA), NIH/NIDDK R01 DK124617 (JCA), NIH/NIAID R21 AI159786 (JCA),

American Gastroenterological Association Augustyn Award in Digestive Cancer (JCA),

Lineberger Comprehensive Cancer Center Pilot Grant (JCA).

We acknowledge the Gnotobiotic Core at the UNC Center for Gastrointestinal Biology

and Disease (CGIBD: supported by NIH P30DK34987), the National Gnotobiotic Rodent

Resource Center (supported by NIH P40 OD01995), and UNC's High-Throughput Sequencing

Facility. We also acknowledge Lacey Lopez, Christopher Broberg, Adrienne Franks, and

Cassandra Barlogio for experimental assistance.

References

1.   Sartor RB, Wu GD. Roles for Intestinal Bacteria, Viruses, and Fungi in Pathogenesis
519   of Inflammatory Bowel Diseases and Therapeutic Approaches. Gastroenterology.
520   2017;152(2):327-339.e4.
521   2.   Kappelman MD, Farkas DK, Long MD, Erichsen R, Sandler RS, Sørensen HT, et al.
522   Risk of cancer in patients with inflammatory bowel diseases: A nationwide
523   population-based cohort study with 30 years of follow-up evaluation. Clin
524   Gastroenterol Hepatol [Internet]. 2014;12(2):265-273.e1. Available from:
525   http://dx.doi.org/10.1016/j.cgh.2013.03.034
526   3.   Arthur JC, Perez-Chanona E, Mühlbauer M, Tomkovich S, Uronis JM, Fan T-J, et al.
527   Intestinal Inflammation Targets Cancer-Inducing Activity of the Microbiota. Science.
528   2012;338(6103):120-123.

529   4.   Dejea CM, Fathi P, Craig JM, Boleij A, Taddese R, Geis AL, et al. Patients with
530   familial adenomatous polyposis harbor colonic biofilms containing tumorigenic
531   bacteria. Science. 2018;359(6375):592–7.
532   5.   Darfeuille-Michaud A, Boudeau J, Bulois P, Neut C, Glasser A-L, Barnich N et al.
533   High Prevalence of Adherent-Invasive Escherichia Coli Associated With Ileal Mucosa
534   in Crohn's Disease. Gastroenterology. 2004;127(2):412-21.
535   6.   Martin HM, Campbell BJ, Hart CA, Mpofu C, Nayar M, Singh R, et al. Enhanced
536   Escherichia coli adherence and invasion in Crohn's disease and colon cancer.
537   Gastroenterology. 2004;127(1):80–93.
538   7.   Baumgart M, Dogan B, Rishniw M, Weitzman G, Bosworth B, Yantiss R, et al.
539   Culture independent analysis of ileal mucosa reveals a selective increase in invasive
540   Escherichia coli of novel phylogeny relative to depletion of Clostridiales in Crohn's
541   disease involving the ileum. ISME J. 2007;1(5):403–18.
542   8.   Knights D, Lassen KG, Xavier RJ. Advances in inflammatory bowel disease
543   pathogenesis: Linking host genetics and the microbiome. Gut. 2013;62(10):1505–10.
544   9.   Frank DN, St. Amand AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR.
545   Molecular-phylogenetic characterization of microbial community imbalances in
546   human inflammatory bowel diseases. Proc Natl Acad Sci U S A. 2007;104(34):13780–
547   5.
548   10.   Rooks MG, Veiga P, Wardwell-Scott LH, Tickle T, Segata N, Michaud M, et al. Gut
549   microbiome composition and function in experimental colitis during active disease and
550   treatment-induced remission. ISME J. 2014;8:1403–1417.
551   11.   Vila AV, Imhann F, Collij V, Jankipersadsing SA, Gurry T, Mujagic A, et al. Gut
552   microbiota composition and functional changes in inflammatory bowel disease and
553   irritable bowel syndrome. Sci Trans Med. 2018;10(472):eaap8914.
554   12.   Kim SC, Tonkonogy SL, Albright CA, Tsang J, Balish EJ, Braun J, et al. Variable
555   phenotypes of enterocolitis in interleukin 10-deficient mice monoassociated with two
556   different commensal bacteria. Gastroenterology. 2005;128(4):891–906.
557   13.   Kim SC, Tonkonogy SL, Karrasch T, Jobin C, Balfour Sartor R. Dual-association of
558   gnotobiotic IL-10-/- mice with 2 nonpathogenic commensal bacteria induces
559   aggressive pancolitis. Inflamm Bowel Dis. 2007;13(12):1457–66.
560   14.   Carvalho FA, Koren O, Goodrich JK, Johansson MEV, Nalbantoglu I, Aitken JD, et
561   al. Transient inability to manage proteobacteria promotes chronic gut inflammation in
562   TLR5-deficient mice. Cell Host Microbe. 2012;12(2):139–52.
563   15.   O'Brien CL, Bringer MA, Holt KE, Gordon DM, Dubois AL, Barnich N, et al.
564   Comparative genomics of Crohn's disease-Associated adherent-invasive Escherichia
565   coli. Gut. 2017;66(8):1382–9.
566   16.   Dogan B, Scherl E, Bosworth B, Yantiss R, Altier C, McDonough PL, et al. Multidrug
567   Resistance Is Common in Escherichia coli Associated with Ileal Crohn's Disease.
568   Inflamm Bowel Dis. 2013;19(1):141–150.
569   17.   Benítez-Páez A, Portune KJ, Sanz Y. Species-level resolution of 16S rRNA gene
570   amplicons sequenced through the MinION™ portable nanopore sequencer.
571   GigaScience. 2016;5(1). doi:10.1186/s13742-016-0111-z.
572   18.   Cuscó A, Catozzi C, Viñes J, Sanchez A, Francino O. Microbiota profiling with long
573   amplicons using Nanopore sequencing: full-length 16S rRNA gene and the 16S-ITS-

23S of the rrn operon [version 2; peer review: 2 approved, 3 approved with
reservations]. F1000Research 2019, 7:1755.

19.   Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS et al. High-throughput
amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide
resolution. Nucleic Acids Research. 2019;47(18):e103. doi:10.1093/nar/gkz569

20.   Karst SM, Ziels RM, Kirkegaard RH, Sørensen EA, McDonald D, Zhu Q, et al. High-
accuracy long-read amplicon sequences using unique molecular identifiers with
Nanopore or PacBio sequencing. Nat Methods. 2021;**18:**165–169.
https://doi.org/10.1038/s41592-020-01041-y

21.   Johnson JS, Spakowicz DJ, Hong B-Y, Petersen LM, Demkowicz P, Chen L, et al.
Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome
analysis. Nat Commun. 2019;10(5029). doi:10.1038/s41467-019-13036-1.

22.   Sambrook J, Russell DW. Purification of Nucleic Acids by Extraction With
Phenol:chloroform. CSH Protoc, 2006 (1).

23.   Tyson J. Rocky Mountain adventures in Genomic DNA sample preparation, ligation
protocol optimisation / simplification and Ultra long read generation. protocols.io.
2020. dx.doi.org/10.17504/protocols.io.7euhjew.

24.   Li H. Minimap and miniasm: fast mapping and de novo assembly for noisy long
sequences. Bioinformatics. 2016;32(14):2103–2110.

25.   Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly
from long uncorrected reads. Genome Res. 2017;27:737-46.
doi:10.1101/gr.214270.116.

26.   Oxford Nanopore Technologies. Medaka; 2018 [cited 2020 March 3]. Available
from: https://nanoporetech.github.io/medaka/

27.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, et al. Pilon: An Integrated Tool
for Comprehensive Microbial Variant Detection and Genome Assembly Improvement.
PLOS ONE. 2014;9(11):e112963. doi:10.1371/journal.pone.0112963.

28.   Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L et al.
NCBI prokaryotic genome annotation pipeline. Nucleic Acids Res. 2016;44(14):6614-
24. doi: 10.1093/nar/gkw569.

29.   Buchfink B, Xie C. Huson DH. Fast and sensitive protein alignment using
DIAMOND. Nature Methods. 2015;12:59–60.

30.   Inouye M, Dashnow H, Raven L-A, Schultz MB, Pope BJ, Tomita T et al. SRST2:
Rapid genomic surveillance for public health and hospital microbiology labs. Genome
Med. 2014;6(11):90.

31.   Center for Genomic Epidemiology; 2011 [cited 2020 March 3]. [Internet]. Available
from: http://www.genomicepidemiology.org/

32.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094–3100.

33.   Tegha G, Ciccone EJ, Krysiak R, Kaphatika J, Chikaonda T, Ndhlovu I, et al.
Genomic epidemiology of Escherichia coli isolates from a tertiary referral center in
Lilongwe, Malawi. Microbial Genomics. 2021;7(1):mgen000490.
doi:10.1099/mgen.0.000490.

34.   Walters MS, Lane MC, Vigil PD, Smith SN, Walk SR, Mobley HLT. Kinetics of
Uropathogenic Escherichia Coli Metapopulation Movement During Urinary Tract
Infection. mBio. 2012;3(1):e00303-11.

35. Gonzalez RJ, Lane MC, Wagner NJ, Weening EH, Miller VL. Dissemination of a
highly virulent pathogen: tracking the early events that define infection. PLoS Pathog.
2015;11(1):e1004587.
36. Arthur JC, Gharaibeh RZ, Mühlbauer M, Perez-Chanona E, Uronis JM, McCafferty J
et al. Microbial genomic analysis reveals the essential role of inflammation in
bacteria-induced colorectal cancer. Nature Communications. 2014;5(4724).
doi:10.1038/ncomms5724
37. Hamming RW. Error detecting and error correcting codes. The Bell System Technical
Journal. 1950;29(2):147-160. doi: 10.1002/j.1538-7305.1950.tb00463.x.
38. Kühn R, Löhler J, Rennick D, Rajewsky K, Müller W. Interleukin-10-deficient mice
develop chronic enterocolitis. Cell. 1993;75(2):263–74.
39. Ellermann M, Gharaibeh RZ, Fulbright L, Dogan B, Moore LN, Broberg CA et al.
Yersiniabactin-Producing Adherent/Invasive *Escherichia coli* Promotes Inflammation-
Associated Fibrosis in Gnotobiotic *Il10*$^{-/-}$ Mice. Infection and Immunity.
2019;87(11):1–18.
40. Sartor RB. Cytokines in intestinal inflammation: Pathophysiological and clinical
considerations. Gastroenterology. 1994;106(2):533–9.
41. Patwa LG, Fan TJ, Tchaptchet S, Liu Y, Lussier YA, Sartor RB, et al. Chronic
intestinal inflammation induces stress-response genes in commensal Escherichia coli.
Gastroenterology. 2011;141(5):1842-1851.e10. doi:10.1053/j.gastro.2011.06.064
42. Ellermann M, Huh EY, Liu B, Carroll IM, Tamayo R, Sartor RB. Adherent-invasive
Escherichia coli production of cellulose influences iron-induced bacterial aggregation,
phagocytosis, and induction of colitis. Infect Immun. 2015;83(10):4068–80.
43. Segain JP, Galmiche JP, Raingeard De La Blétière D, Bourreille A, Leray V, Gervois
N, et al. Butyrate inhibits inflammatory responses through NFκB inhibition:
Implications for Crohn's disease. Gut. 2000;47(3):397–403.
44. Elhenawy W, Tsai CN, Coombes BK, Elhenawy W, Tsai CN, Coombes BK. Host-
Specific Adaptive Diversification of Crohn's Escherichia coli Article Host-Specific
Adaptive Diversification of Crohn's Disease-Associated Adherent-Invasive
Escherichia coli. Cell Host Microbe. 2019;25(2):301-312.e5.
45. Elhenawy W, Hordienko S, Gould S, Oberc AM, Tsai CN, Hubbard TP, et al. High-
throughput fitness screening and transcriptomics identify a role for a type IV secretion
system in the pathogenesis of Crohn’s disease-associated Escherichia coli. Nat
Commun. (2021):12(2032). doi:10.1038/s41467-021-22306-w
46. Boudeau Ã, Barnich N, Intestinale Â, De L, Pharmacie De. Type 1 pili-mediated
adherence of Escherichia coli strain LF82 isolated from Crohn's disease is involved in
bacterial invasion of intestinal epithelial cells. Mol Micro. 2001;39(5):1272-1284.
47. Barnich N, Colombel J, Barnich N, Carvalho FA, Glasser A, Darcha C, et al.
CEACAM6 acts as a receptor for adherent-invasive E. coli, supporting ileal mucosa
colonization in Crohn disease. J Clin Invest. 2007;117(6):1566-1574.
48. Tropini C, Earle KA, Huang KC, Sonnenburg JL. The Gut Microbiome: Connecting
Spatial Organization to Function. Cell Host Microbe. 2017;21(4):433–42.
doi:10.1016/j.chom.2017.03.010
49. Human Microbiome Project Consortium. Structure, function and diversity of the
healthy human microbiome. Nature. 2012;486:207–214. doi:10.1038/nature11234.

50.   Lloyd-Price J, Mahurkar A, Rahnavard G, Crabtree J, Orvis J, Hall AB, et al. Strains,
666   functions and dynamics in the expanded human microbiome project. Nature.
667   2017;550:61–66. doi:10.1038/nature23889.

A

Il10⁻ᐟ⁻

E. coli pool

Kan

70 days
Harvest Stool
and Tissue

Day: 0   14   28

B

Strain Relative Abundance in Stool

Percent Abundance

NC 101 | CU39 ES-1 | CU 532-9 | LF 82 | CU 568-3 | CU37 RT-2 | CU42 ET-1 | HM 670

C

Strain Relative Abundance in Stool

qPCR Percent Abundance

NC 101 | CU39 ES-1 | CU 532-9 | LF 82 | CU 568-3 | CU37 RT-2 | CU42 ET-1 | HM 670

S1 Figure. Timeline for mouse experiment to determine colonize of individual *E. coli* isolates.

Strain	Mouse Gender	Ileal Tissue	Cecal Content	Colon Content	Mucus Layer	Colon Tissue
CU39ES-1/A1	M	65957	3916667	8588235	3556	12889
CU39ES-1/A1	F	611	5882353	34166667	6500	10833
CU532-9/A3	M	0	5806452	7058824	6667	455556
LF82/B6	M	1875	232609	257732	200	1800
LF82/B6	F	500	111111	74684	417	417
CU568-3/C5	M	1250	40909091	5306122	24545	681818
CU568-3/C5	F	0	0	71	0	0
CU37RT-2/D2	M	0	0	0	0	0
CU42ET-1/D5	F	28947	167857	320513	200	5133
CU42ET-1/D5	M	107857	1333333	ND	1500	32308
HM670/C2	F	0	34	0	0	188
HM670/C2	M	13810	460317	422018	6583	25000
NC101	M	500	39333	24615	0	70000

*Mice were all singly-housed.
* CU532-9/A3 F (dead), CU37RT-2/D2 F (dead), NC101 F (dead)

S1 Table. *E. coli* isolates colonize the lower gastrointestinal tract of *Il10*[-/-] mice with FMT competition. Values indicate CFUs/10 mg of tissue.

A

B

S2 Figure. A. Timeline for mouse experiment for colonization of CU42ET-1/D5 or HM670/C2. B. Levels of *E. coli* in stool over time (n=7 in CU42ET-1/D5 and n=8 in HM670/C2). Data point indicates the mean and error bars indicates the standard deviation (**p<0.01 Mann-Whitney test). If not otherwise noted, there is no significance.

Mouse	Cage	Colon Content	Mucus Layer	Colon Tissue
CU42ET-1/D5 1	1	19,083,969	400,000	1,181,818
CU42ET-1/D5 2	1	7,565,217	177,778	722,222
CU42ET-1/D5 3	1	3,802,817	5,455	150,000
CU42ET-1/D5 4	2	49,494,949	210,526	894,737
CU42ET-1/D5 5	2	37,419,355	67,308	730,769
CU42ET-1/D5 6	2	33,750,000	225,000	616,667
CU42ET-1/D5 7	2	95,890,411	45,000	4,000,000
HM670/C2 1	3	553,191	333	66,667
HM670/C2 2	3	1,193,277	7,250	300,000
HM670/C2 3	3	2,019,231	0	17,000
HM670/C2 4	3	2,179,487	1,125	13,125
HM670/C2 5	4	204,301	1,000	10,778
HM670/C2 6	4	0	0	0
HM670/C2 7	4	139	0	0
HM670/C2 8	4	0	0	0

S2 Table. Consistent *E. coli* colonizer (CU42ET-1/D5) and inconsistent *E. coli* colonizer (HM670/C2) colonize the lower gastrointestinal tract of *Il10*[-/-] mice with FMT competition.