In silico Design of novel Multi-epitope recombinant Vaccine based on Coronavirus surface glycoprotein

Mandana Behbahani[1]*

1 Department of Biotechnology, Faculty of Biological Science and Technology, University of Isfahan, Hezar Jareeb St., Isfahan 81746-73441, Iran.

* Corresponding author: E-mail: Ma_behbahani@yahoo.com, Ma.behbahani@ast.ui.ac.ir, Tel (Office): 0098-311-7934327, Fax: 0098-313-7932342.

Abstract

It is of special significance to find a safe and effective vaccine against coronavirus disease 2019 (COVID-19) that can induce T cell and B cell -mediated immune responses. There is currently no vaccine to prevent COVID-19. In this project, a novel multi-epitope vaccine for COVID-19 virus based on surface glycoprotein was designed through application of bioinformatics methods. At the first, seventeen potent linear B-cell and T-cell binding epitopes from surface glycoprotein were predicted in silico, then the epitopes were joined together via different linkers. The ability of the selected epitopes to induce interferon-gamma was evaluate using IFNepitope web server. One final vaccine was constructed which composed of 398 amino acids and attached to 50S ribosomal protein L7/L12 as adjuvant. Physicochemical properties, as well as antigenicity in the proposed vaccines, were checked for defining the vaccine stability and its ability to induce cell-mediated immune responses. Three-dimensional structure of the mentioned vaccine was subjected to the molecular docking studies with MHC-I and MHC-II molecules. The results proposed that the multi-epitope vaccine with 50S ribosomal protein L7/L12 was a stable construct with high aliphatic content and high antigenicity.

Keyboards: Vaccine, Multi-epitope, Coronavirus, Surface glycoprotein

1. Introduction

In early 2020, COVID-19 began generating headlines all over the world because of the extraordinary speed of its transmission. So, there are rising concerns about community infections. Vaccination is one of the most effective tools to prevent infectious diseases [1][2]. As far as our knowledge concerns, there is no report about developing COVID-19 multi epitope vaccine. Therefore, we became eager to design potent multiepitope vaccines from antigenic sites of coronavirus surface glycoprotein. The multiepitope vaccines have advantageous over conventional vaccines with regards to safety profile and high immunogenicity [3]. Multiepitope vaccines have the potential to induce responses restricted by a wide variety of HLA molecules and generate a balanced CD4+ and CD8+ cellular immune response. Another molecule that contribute to innate immunity contains TLR3 that activates antiviral mechanism during infection. Recently, in silico design of epitope-based vaccines has been done for vaccine developing against many infectious diseases. Some bioinformatics tools could facilitate the development of multi epitope-based vaccines. The computational tools can optimize the extensive immunological data such as antigen presentation and processing to achieve specific interpretations [4]. In recent decades, several vaccines were

established based on in silico methods that include efficient vaccines against Toxoplasma gondii
[5], Brucella abortus [6], Escherichia coli [7], Vibrio cholera [8], Human immunodeficiency
virus-1 [9], Hepatitis C virus [10] and many others. In several experimental studies, the efficacy
of computationally designed vaccines has been recently approved for use in defined human
vaccines [11][13]. In this study, in silico analysis were performed to determine exclusive B cell
and T-cell epitopes from coronavirus surface glycoprotein that are antigenically most significant
for coronavirus. In our research, some unique exclusive B cell and T-cell epitopes from
coronavirus surface glycoprotein were selected based on their antigenicity, stability and length.
The selected epitopes were merged into each other using suitable linkers for organization of final
vaccine construct. Consequently, the stability and efficacy of the vaccines were predicted by a
set of bioinformatics methods.

2. Material and methods

2.1 Data collection

At the first step of our study, the reference amino acid sequences of coronavirus surface glycoprotein
(YP-001856243.1), five   HLA-1 (NP_001229971.1, NP_001229687.1, NP_002118.1, NP_061823.2,
NP_005507.3) and six HLA-2 protein (NP_001229454.1, NP_006111.2, NP_001230891.1,
NP_002110.1, NP_061984.2, NP_001020330.1) were retrieved from NCBI
(https://www.ncbi.nlm.nih.gov). SWISS-MODEL Server (https://swissmodel.expasy.org/) was utilized
for modelling of 3-D structures of HLA class I and HLA class II, But for TLR-3 the data in PDB bank
was used and optimized by chimera 1.12 [14.]

2.2 Multiple sequence alignment and antigen selection

To determine exclusive conserved sequence of the coronavirus surface glycoprotein, NCBI BLAST was
performed (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Also, for defining the conserved region (s) in the
protein sequences, multiple sequence alignment was done by Multalin server
(https://www.multalin.toulouse.inra.fr/multalin). Additionally, the antigenicity of the coronavirus surface
glycoprotein was evaluated using VaxiJen 2.0 server (http://www.ddg-
pharmfac.net/vaxijen/VaxiJen/VaxiJen.html[15]. Finally, the most particular conserved sequence and
antigenic peptide were selected for further analysis.

2.3 B-cell epitope prediction and selection

Linear B-cell epitopes in the vaccine model were predicted using ElliPro
(http://crdd.osdd.net/raghava/bcepred) [16] and IEDB analysis Resource (http://tools.iedb.org/population)
[17].

2.4 T-cell epitope prediction and selection

MHC-I restricted epitopes were predicted through ProPred-1 server
(http://tools.immuneepitope.org/analyze/html/mhc_binding.html) [18]. The server uses special patterns
for HLA-A*03:01 allele. Similarly, MHC-II restricted epitopes were predicted using ProPred server
(http://tools.immuneepitope.org/mhcii). The server uses special patterns for DRB1*01:07 allele. Finally,
the conserved sequence and antigenic peptide were selected for further analysis.

2.5 Construction of final vaccine

Seventeen suitable common B-cell and T-cell epitopes (9-16 amino acids) from coronavirus surface
glycoprotein were selected and organized in the final vaccine construct. Then, these epitopes were merged
together with AAY, KK linkers and considered as a multi-epitope vaccine. One adjuvant "50S ribosomal
protein L7/L12
(MSDINKLAETLVNLKIVEVNDLAKILKEKYGLDPSANLAIPSLPKAEILDKSKEKTSFDLILKGAG
SAKLTVVKRIKDLIGLGLKESKDLVDNVPKHLKKGLSKEEAESLKKQLEEVGAEVELK)

with 124 amino acids were incorporated with **EAAAK** linker at N-terminal portion of the constructs.
The sequence of the designed vaccine structures with their adjuvant are depicted in Table 3. The final
vaccines (IV1) stretch was found to be 398 amino acids.

2.6 Physicochemical properties analysis

In this research, five characteristics (molecular weight, theoretical pI, extinction coefficient, aliphatic
index and grand average of hydropathicity) of the constructed vaccine was evaluated using ProtParam
server (http://web.expasy.org/protparam).

2.7 Secondary structure analysis

The frequency of the secondary structure of the constructed vaccines (alpha helix, extended strand and
random coil) were computed using GOR IV web server (http://npsa-pbil.ibcp.fr/cgi-
bin/npsa_automat.pl?page=npsa_gor4.html).

2.8 Molecular docking study

To confirm the binding affinity of the best vaccine to MHC-I and MHC-II molecules, molecular docking
was done between the selected vaccines and five HLA-1 structures (NP_001229971.1, NP_001229687.1 ,
NP_002118.1 , NP_061823.2 , NP_005507.3), also six HLA-2 proteins (accession numbers:
NP_001229454.1 , NP_006111.2 , NP_001230891.1 , NP_002110.1 , NP_061984.2 , NP_001020330.1)
and TLR-3 (PDB ID: 2A0Z) separately. Molecular docking studies were done using H-dock server
(https://bioinfo3d.cs.tau.ac.il/ PatchDock/) with default complex type and clustering RMSD of 4Å. The
binding sites of final construct and TLR3 were studied using fully automated protein-ligand interaction
profiler server (PLIP; https://projects.biotec.tu-dresden.de/plip-web/plip/index). The outputs of PLIP were
in XML format, flat text, and visualization files [19]. The visualization files were visualized using
PyMOL software (windows version 2.0.7). Then, seventeen epitopes were analysed by IFNepitope server.

.

2.10 IFNγ analysis

IFNepitope web server is established for users working in the field of vaccine design. This server allows
users to predict and design IFN-gamma inducing peptides. The ability of the selected epitopes to induce
interferon-gamma was evaluate using IFN epitope server.

3. Results

3.1 Multiple sequence alignment and antigen selection

Initially, coronavirus surface glycoprotein was studied for determining specific conserved part of protein
between the virus serotypes. Results of protein BLAST are shown in Table 1. The results demonstrated
that coronavirus surface glycoprotein had the most conservancy levels between 97. 8 and 100. Also,
VaxiJen score of the protein showed high antigenicity. Due to having good antigenicity, high exposure
probability to the immune system and high conservancy, this protein was selected for vaccine design.

3.2 T-cell and B-cell epitope prediction

In the current study, some appropriate common B-cell and T-cell epitopes were designed. The predicted
MHC-I and MHC-II restricted epitopes were compared to B-cell epitopes to determine shared epitopes
(Table 2). Finally, 17 epitopes with 9-16 amino acids were selected. These epitopes are located between
residues 14-642. These epitopes exhibited a relatively high aliphatic index (>60), high antigenicity (>1.6)
and low instability index (less than 20). The most potent epitope was repeated three times and merged
into the other epitopes using suitable linkers (KK, AYY) for organization of final vaccine construct.

3.3 Antigen selectivity of constructed vaccines

Final construct vaccine was composed of 398 amino acids which were respectively attached to 50S
ribosomal protein L7/L12 as adjuvant. The antigenicity score of constructed vaccine are shown in
Table 3. The result demonstrated that IV1 has 1.2110 antigenicity (Table 3).

3.4 Physicochemical properties

Physicochemical properties of the constructed vaccine were predicted using Protparam server. The results
revealed that this multi-epitope vaccine have low instability "as a value below 40" predicts that the
protein is stable. The IV1 construct showed the highest Isoelectric point with 8.52 value. From the
aliphatic Index of view, this construct showed aliphatic index more than 90 % (Table 4). The results of
gor4 demonstrated that the random coil values of the IV1 were high compared to Alpha helix and
extended structure.

3.5 Analysis of docking results

The results of docking the constructions with HLA-1 and HLA-2 confirmed the high values with IV1
construct. Also, the results of TLR-3 analysis showed values of 1680.569 for IV1 which demonstrate the
high efficacy of vaccine construct (Tables 5). Figure 2 indicate docking results of the Vac1 construct
with the TLR-3 as the examples of vaccines potential for interaction. The results showed that TLR-3 have
interaction to Glu115, Gly 118 residues of AV1.

2.10 IFNγ analysis

The result of IFNγ analysis was described in Table 6. The result demonstrates that among 17 epitopes ,16
epitopes have potential to produce IFN-γ. Epitope 6 showed the highest score with value of 2.

4. Discussion

Due to the nature of coronavirus and high infectious rate, the progress of a vaccine against coronavirus is
very challenging. Though, with the development of computational methods, these limitations are reduced.
By the way, using computational methods, the design of recombinant vaccines and the estimation of
physicochemical properties as well as vaccines efficacy could be available [20] [21]. Then, this research

was intended to design an effective multi-epitope recombinant vaccine against coronavirus using a unique
multi-step bioinformatics approach. Our potent multi-epitope vaccine is contained seventeen epitopes in
surface glycoprotein of coronavirus along with AAY and KK linkers and 50S ribosomal protein L7/L12
as adjuvant. Based on our knowledge, there is no report about computational design of epitope-based
vaccine for coronavirus. Recently, in silico, design of epitope-based vaccine was used for vaccine
development against several infectious diseases. Several bioinformatics tools have been established that
accelerate the growth of multi epitope-based vaccines. In recent decade, several multiepitope vaccines for
pathogenic viruses have been reported. Multi epitope vaccines could provide an effective immunization
against different serotypes of a pathogen. Despite mentioned advantages of Multi epitope vaccines, poor
immunogenicity is considered as a major drawback to growth of these vaccines [22], [23]. The in-silico
results proposed that our multi-epitope vaccine was very stable with high aliphatic index and it was
potentially antigenic. As reported earlier, high aliphatic index shows the higher thermos-stability of the
constructed vaccine. At the present research, the aliphatic index was high, and instability was low and
Gravy indices were negative. Also, the higher PI, as the case of this study, shows the higher potential for
cell wall attachment. However, the proposed vaccine has high antigenicity. This was chosen for docking
studies. The results demonstrated that our mentioned vaccine could be a right candidate for experimental
research [24].

Conclusion

174 This study introduced designing novel multi epitope vaccines against Coronavirus which could cover
175 conserve sequence of the virus. The multi-epitope vaccine presented by this study showed promising
176 result through in silico step, which could be followed by in vitro and in vivo studies.

Conflict of interest

179 Authors declare no conflict of interest.

Acknowledgment

181 We wish to thank the University of Isfahan for their supports.

References

1. Li Q., Guan X., Wu P., Wang X., Zhou L (2020) Tong Y., Feng Z. Early transmission dynamics in
Wuhan, China, of novel coronavirus-infected pneumonia. New England Journal of Medicine.

2. World Health Organization Novel coronavirus (2019-nCoV) situation reports. 2020.
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports

3. Cicala C, Nawaz F, Jelicic K, et al. (2016) HIV-1 gp120: A Target for Therapeutics and Vaccine
Design. Current Drug Targets. 17(1):122-35.

4. Kumar M, Thakur V, Raghava GP (2008) "COPid: composition based protein identification. In silico
b10.Soria-Guerra RE, Nieto-Gomez R, Govea-Alonso DO, et al. An overview of bioinformatics tools for
epitope prediction: implications on vaccine Development. J Biomed Inform. 2015;53, 405−414. doi:
10.1016/j.jbi.2014.11.003.iology. 8:121-128.

5. Soria-Guerra RE, Nieto-Gomez R, Govea-Alonso DO (2015) t al. An overview of bioinformatics tools
for epitope prediction: implications on vaccine Development. J Biomed Inform. 2015;53, 405‒414. doi:
10.1016/j.jbi.2014.11.003.

6.Bissati KE, Chentoufi AA, Krishack PA (2016) Adjuvanted multi-epitope vaccines protect HLA-A* 11:
01 transgenic mice against Toxoplasma gondii. JCI Insight. 1(15): doi: 10.1172/jci.insight.85955.

7.Escalona E, Saez D, Onate A (2017) Immunogenicity of a multi-epitope dna vaccine encoding epitopes
from Cu‒Zn superoxide dismutase and open reading Frames of Brucella abortus in mice. Frontiers
Immunology. 8, 125

8.Rodrigues-da-Silva RN, Martins da Silva JH, Singh B (2016) In silico identification and validation of a
linear and naturally immunogenic B-cell epitope of the Plasmodium vivax malaria vaccine candidate
merozoite surface protein-9. Plos One;11(1): e 0146951. doi: 10.1371
204

9.Nezafat N, Karimi Z, Eslami M (2016) Designing an efficient multi-epitope peptide vaccine against
Vibrio cholerae via combined immunoinformatics and protein interaction based approaches.
Computational Biology and Chemistry. 62: 82‒95. doi:10.1016/j.compbiolchem.2016.04.006

10.Yang Y, Sun W, Guo J, et al. In silico design of a DNA based HIV-1 multi-epitope vaccine for
Chinese populations. Human Vaccines Immunother. 2015;11(3): 795‒805. doi:
10.1080/21645515.2015.1012017

11.Nosrati M, Mohabatkar H, Behbahani M (2017) A novel multi-epitope vaccine for cross protection
against Hepatitis C Virus (HCV): an immunoinformatics. Approach Research in Molecular Medicine
5(1): 17‒26.

12.Rahjerdi AK, AmaniJ, Rad I (2016) Designing and structure evaluation of multi-epitope vaccine
against ETEC and EHEC, an in silico approach.Protein Peptide Letters. 23(1): 33‒42.

13.Oscherwitz J (2016) The promise and challenge of epitope-focused vaccines. Human Vaccines
Immunother.; 12(8): 2113‒2116.

14. Conrad C. Huang, Elaine C. Meng, John H. Morris (2014) Enhancing UCSF Chimera through web
services. Nucleic Acids Research. 42(1):478-484. https://doi.org/10.1093/nar/gku377

15.Doytchinova IA, Flower DR (2007) VaxiJen: a server for prediction of protective antigens, tumour
antigens and subunit vaccines. BMC Bioinformatics.

16.Duquesnoy R, Marrari M (2017) Usefulness of the ElliPro epitope predictor program in defining the
repertoire of HLA-ABC eplets. Hum .78(7-8): 481-488. doi:10.1016/j.humimm.03.005

17.Beaver JE, Bourne PE, Ponomarenko JV (2007) Epitope Viewer: a Java application for the
visualization and analysis of immune epitopes in the Immune Epitope Database and Analysis Resource
(IEDB). Immunome   Res   DOI:10.1186/1745-7580-3-3.

18. Patronov AI. Doytchinov I (2013) T-cell epitope vaccine design by immunoinformatics. Open Biol.
2013; 3(1):120139. doi: 10.1098/rsob.120139.

19.Salentin S, Schreiber S, Haupt VJ (2015) PLIP: fullyautomated protein–ligand interaction profiler.
Nucleic Acids Research ;43: W443–W447.https://doi.org/10.1093/nar/gkv315
231
232
233   20. Roosa K. , Lee Y., Luo R., Kirpich A., Rothenberg R., et al (2020) Hyman J.M., Yan P., and G.
234   Chowell, Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th,
235   2020, Infect Dis Model. 2020; 5: 256–263.
236
237
238   21. Ai S., Zhu G., Tian F., Li H., Gao Y., Wu Y., Lin H (2020) Population movement, city closure and
239   spatial transmission of the 2019-nCoV infection in China.
240
241

22.Paul S, Piontkivska H (2010) Frequent associations between CTL and T-Helper epitopes in HIV-1
genomes and implications for multi-epitope vaccine designs. BMC Microbiology. 10:212. doi:
10.1186/1471-2180-10-212

23.Hajighahramani N, Nezafat N, Eslami M, et al. (2017) Immunoinformatics analysis and in silico
designing of a novel multiepitope peptide vaccine against Staphylococcus aureus. Infection Genetics and
Evolution. 48: 83–94. doi: 10.1016/j.meegid

24.Solanki V, Tiwari M, Tiwari V. (2019) Prioritization of potential vaccine targets using comparative
proteomics and designing of the chimeric multi-epitope vaccine against Pseudomonas aeruginosa.
Scientific Reports. 9.

261   Fig 1: Molecular docking analysis of IV1construct with TLR-3. The results showed that TLR-3
262   have interaction to Glu 115 and Gly 118 residues of AV1

274   Table 1: Results of the antigenicity and BLAST of coronavirus surface glycoprotein and
275   antigenicity prediction

Protein	Average antigenicity	VaxiJen score	Minimum identity (%)	Maximum identity (%)
Coronavirus surface glycoprotein	0.4646	0.4	97.8	100

278   Table 2: Result of final T-cell and B-cell epitope prediction screening from coronavirus surface
279   glycoprotein

	Sequence	Length	Start-End	Vaxijen	Isoelectric point	Aliphatic index	Instability	GRAVY
1	KLNDLCFTN	9	386-394	2.90	5.83	86.67	17.24	-0.24
2	KLNDLCFTNV	10	386-395	2.69	5.83	88	-14.52	-0.2
3	KLNDLCFTNVY	11	386-396	2.22	5.83	97.27	-19.15	-0.06
4	PTKLNDLCFTN	11	384-394	2.19	6.22	70.91	-12.29	-0.4
5	SPTKLNDLCFTN	12	383-394	2.01	5.55	65	26.18	-0.4
6	VSPTKLNDLCFTN	13	382-394	2.17	5.80	82.32	24.94	-0.08
7	KLNDLCFTNVYA	12	386-397	1.92	5.83	97.5	3	0.2
8	KLNDLCFTNVYAD	13	386-398	1.67	4.21	90	-2.98	-0.07
9	LNDLCFTNV	9	387-395	2.01	3.80	118.89	-7.81	0.6
10	GVSPTKLNDLCFTN	14	381-384	2.21	5.83	76.43	23.87	-0.1
11	YGVSPTKLNDLCFTN	15	380-384	2.06	5.83	71.33	17.29	-0.18
12	CYGVSPTKLNDLCFTN	16	379-384	2.01	5.82	66.88	16.83	-0.01
13	CVNLTTRTQ	9	15-23	1.87	8.25	75.56	-7.87	-0.34
14	QCVNLTTRTQ	10	14-23	1.78	8.25	68	-13.62	-0.66
15	LDITPCSFGGVSV	13	585-697	1.88	3.80	104.62	12.92	1.06
16	LDITPCSFGGVSVI	14	585-698	1.61	3.80	125	12.71	1.03
17	VKNKCVNFN	9	534-642	2.05	9.31	64.44	6.92	-0.51

Vaccine	Sequence	Vaxijen
IV1	**MSDINKLAETLVNLKIVEVNDLAKILKEKYGLDPSANLAIPSLPKAEILDKSKEKTSF DLILKGAGSAKLTVVKRIKDLIGLGLKESKDLVDNVPKHLKKGLSKEEAESLKKQL EEVGAEVELKEAAAKKLNDLCFTNAAYKLNDLCFTNAAYKLNDLCFTNAAYKLND LCFTNVAAYKLNDLCFTNVYAAYPTKLNDLCFTNAAYSPTKLNDLCFTNAAYVSPT KLNDLCFTNAAYLNDLCFTNVAAYKLNDLCFTNVYAAAYKLNDLCFTNVYADAAY GVSPTKLNDLCFTNAAYYGVSPTKLNDLCFTNAAYCYGVSPTKLNDLCFTNAAYCV NLTTRTQAAYQCVNLTTRTQKKLDITPCSFGGVSVKKLDITPCSFGGVSVIKKVKN KCVNFN**	1.2110

289 **Table 3:** Average antigenicity of constructed vaccine using Vaxijen

297 **Table 4:** Physico-properties of constructed vaccines

Protein	Molecular weight	Isoelectric point	Aliphatic index	GRAVY	Instability	Alpha	Extended	Random coil
IV1	43679.43	8.52	92.71	-0.046	11.85	23.37%	8.54%	68.09

301 **Table 5.** HDOCK scores of HLA1, HLA2 and interaction residues of selected sequences

Protein	Hla1-A1	Hla1-Chain G	Hla1-Chain E	Hla1-Chain F	Hla1-Chain CW-1	TLR3	Hla2 -antigen gama chain	Hla2-DM	Hla2-DO	Hla2-DP	Hla2-DQ	Hla2- DR
IV1	951.2	957.2	951.2	1215.7	1220.8	1680.5	1215	1214	1215.7	1215.7	1215.7	1215.7

Table 6: IFNγ analysis score of 17 selected epitopes using IFNepitope web server

Donor	EP1	EP2	EP3	EP4	EP5	EP6	EP7	EP8	EP9	EP10	EP11	EP12	EP13	EP14	EP15	EP16	EP17
1	0.45	0.46	0.47	1	1	2	-	0.47	0.45	1	1	1	0.46	0.46	1	1	0.45