1 **In silico Design of novel Multi-epitope recombinant Vaccine based on Coronavirus surface**
2 **glycoprotein**
3

Mandana Behbahani*

1 Department of Biotechnology, Faculty of Biological Science and Technology, University of Isfahan,
Hezar Jareeb St., Isfahan 81746-73441, Iran.

* Corresponding author: E-mail: Ma_behbahani@yahoo.com, Ma.behbahani@ast.ui.ac.ir, Tel (Office):
0098-311-7934327, Fax: 0098-313-7932342.

Abstract

It is of special significance to find a safe and effective vaccine against coronavirus disease 2019 (COVID-
19) that can induce T cell and B cell -mediated immune responses. There is currently no vaccine to prevent
COVID-19. In this project, a novel multi-epitope vaccine for COVID-19 virus based on surface
glycoprotein was designed through application of bioinformatics methods. At the first, seventeen potent
linear B-cell and T-cell binding epitopes from surface glycoprotein were predicted in silico, then the
epitopes were joined together via different linkers. The immunogenicity of these epitopes was identified
using IFN-γ ELIspot assays. The IFN-γ producing T cell variation ranged from 11.1 ± 1.2 SFU to 38.2 ± 2.1
SFU per 10^6 PBMCs. One final vaccine was constructed which composed of 398 amino acids and attached
to 50S ribosomal protein L7/L12 as adjuvant. Physicochemical properties, as well as antigenicity in the
proposed vaccines, were checked for defining the vaccine stability and its ability to induce cell-mediated
immune responses. Three-dimensional structure of the mentioned vaccine was subjected to the molecular
docking studies with MHC-I and MHC-II molecules. The results proposed that the multi-epitope vaccine
with 50S ribosomal protein L7/L12 was very stable with high aliphatic content and high antigenicity.

Keyboards: Vaccine, Multi-epitope, Coronavirus, Surface glycoprotein

1. Introduction

In early 2020, COVID-19 began generating headlines all over the world because of the extraordinary speed
of its transmission. So, there are rising concerns about community infections. Vaccination is one of the
most effective tools to prevent infectious diseases [1][2]. As far as our knowledge concerns, there is no
report about developing COVID-19 multi epitope vaccine. Therefore, we became eager to design potent
multiepitope vaccines from antigenic sites of coronavirus surface glycoprotein. The multiepitope
vaccines have advantageous over conventional vaccines with regards to safety profile and high
immunogenicity [3]. Multiepitope vaccines have the potential to induce responses restricted by a
wide variety of HLA molecules and generate a balanced CD4+ and CD8+ cellular immune
response. Another molecule that contribute to innate immunity contains TLR3 that activates
antiviral mechanism during infection. Recently, in silico design of epitope-based vaccines has
been done for vaccine developing against many infectious diseases. Some bioinformatics tools
could facilitate the development of multi epitope-based vaccines. The computational tools can
optimize the extensive immunological data such as antigen presentation and processing to achieve
specific interpretations [4]. In recent decades, several vaccines were established based on in silico

methods that include efficient vaccines against Toxoplasma gondii [5], Brucella abortus [6],
Escherichia coli [7], Vibrio cholera [8], Human immunodeficiency virus-1 [9], Hepatitis C virus
[10] and many others. In several experimental studies, the efficacy of computationally designed
vaccines has been recently approved for use in defined human vaccines [11][13]. In this study, in
silico analysis were performed to determine exclusive B cell and T-cell epitopes from coronavirus
surface glycoprotein that are antigenically most significant for coronavirus. In our research, some
unique exclusive B cell and T-cell epitopes from coronavirus surface glycoprotein were selected
based on their antigenicity, stability and length. The selected epitopes were merged into each other
using suitable linkers for organization of final vaccine construct. Consequently, the stability and
efficacy of the vaccines were predicted by a set of bioinformatics methods.

2. Material and methods

2.1 Data collection

At the first step of our study, the reference amino acid sequences of coronavirus surface glycoprotein (YP-
001856243.1), five   HLA-1 (NP_001229971.1, NP_001229687.1, NP_002118.1, NP_061823.2,
NP_005507.3) and six HLA-2 protein (NP_001229454.1, NP_006111.2, NP_001230891.1, NP_002110.1,
NP_061984.2, NP_001020330.1) were retrieved from NCBI (https://www.ncbi.nlm.nih.gov). SWISS-
MODEL Server (https://swissmodel.expasy.org/) was utilized for modelling of 3-D structures of HLA class
I and HLA class II, But for TLR-3 the data in PDB bank was used and optimized by chimera 1.12 [14.]

2.2 Multiple sequence alignment and antigen selection

To determine exclusive conserved sequence of the coronavirus surface glycoprotein, NCBI BLAST was
performed (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Also, for defining the conserved region (s) in the
protein   sequences,   multiple   sequence   alignment   was   done   by   Multalin   server
(https://www.multalin.toulouse.inra.fr/multalin). Additionally, the antigenicity of the coronavirus surface
glycoprotein   was   evaluated   using   VaxiJen   2.0   server   (http://www.ddg-
pharmfac.net/vaxijen/VaxiJen/VaxiJen.html[15]. Finally, the most particular conserved sequence and
antigenic peptide were selected for further analysis.

2.3 B-cell epitope prediction and selection

Linear B-cell epitopes in the vaccine model were predicted using ElliPro
(http://crdd.osdd.net/raghava/bcepred) [16] and IEDB analysis Resource (http://tools.iedb.org/population)
[17].

2.4 T-cell epitope prediction and selection

MHC-I restricted epitopes were predicted through ProPred-1 server
(http://tools.immuneepitope.org/analyze/html/mhc_binding.html) [18]. The server uses special patterns
for HLA-A*03:01 allele. Similarly, MHC-II restricted epitopes were predicted using ProPred server
(http://tools.immuneepitope.org/mhcii). The server uses special patterns for DRB1*01:07 allele. Finally,
the conserved sequence and antigenic peptide were selected for further analysis.

2.5 Construction of final vaccine

Seventeen suitable common B-cell and T-cell epitopes (9-16 amino acids) from coronavirus surface
glycoprotein were selected and organized in the final vaccine construct. Then, these epitopes were merged
together with AAY, KK linkers and considered as a multi-epitope vaccine. One adjuvant "50S ribosomal
protein L7/L12
(MSDINKLAETLVNLKIVEVNDLAKILKEKYGLDPSANLAIPSLPKAEILDKSKEKTSFDLILKGAG
SAKLTVVKRIKDLIGLGLKESKDLVDNVPKHLKKGLSKEEAESLKKQLEEVGAEVELK)

with 124 amino acids were incorporated with **EAAAK** linker at N-terminal portion of the constructs.
The sequence of the designed vaccine structures with their adjuvant are depicted in Table 3. The final
vaccines (IV1) stretch was found to be 398 amino acids.

87 2.6 Physicochemical properties analysis

88 In this research, five characteristics (molecular weight, theoretical pI, extinction coefficient, aliphatic
89 index and grand average of hydropathicity) of the constructed vaccine was evaluated using ProtParam
90 server (http://web.expasy.org/protparam).

91 2.7 Secondary structure analysis

92 The frequency of the secondary structure of the constructed vaccines (alpha helix, extended strand and
93 random coil) were computed using GOR IV web server (http://npsa-pbil.ibcp.fr/cgi-
94 bin/npsa_automat.pl?page=npsa_gor4.html).

96 2.8 Molecular docking study

To confirm the binding affinity of the best vaccine to MHC-I and MHC-II molecules, molecular docking
was done between the selected vaccines and five HLA-1 structures (NP_001229971.1, NP_001229687.1 ,
NP_002118.1 , NP_061823.2 , NP_005507.3), also six HLA-2 proteins (accession numbers:
NP_001229454.1 , NP_006111.2 , NP_001230891.1 , NP_002110.1 , NP_061984.2 , NP_001020330.1)
and TLR-3 (PDB ID: 2A0Z) separately. Molecular docking studies were done using H-dock server
(https://bioinfo3d.cs.tau.ac.il/ PatchDock/) with default complex type and clustering RMSD of 4Å. The
binding sites of final construct and TLR3 were studied using fully automated protein-ligand interaction
profiler server (PLIP; https://projects.biotec.tu-dresden.de/plip-web/plip/index). The outputs of PLIP were
in XML format, flat text, and visualization files [19]. The visualization files were visualized using PyMOL
software (windows version 2.0.7). Then, seventeen epitopes were synthetized by Zist Mobin company
(Isfahan, Iran) at the recommended purity for ELISPOT assay.

108 2.9 Human PBMC preparation

109 Peripheral Blood Mononuclear Cells (PBMCs) from five healthy volunteers were separated on density
110 gradient centrifugation (lymphodex). The PBMCs were grown in RPMI medium supplemented with 10%
111 (v/v) heat inactivated Fetal Calf Serum (FCS; Gibco-BRL, Grand Island, NY, USA), 100 U/ml penicillin,
112 100 mg/ml streptomycin, 2 mM glutamine and 1 mM Na-pyruvate and activated with 5 μg/ml
113 phytohemagglutinin (PHA) and IL-2. .

114 2.10 IFNγ-ELISPOT assay

IFNγ-ELISPOT assay was used to identify the prevalence of pre-existing CD4+ Tcell immunity. Human IFN-γ ELISPOTPRO kits from Mabtech, Nacka Strand, Sweden were used to quantify the frequency of epitope-specific IFN-γ secreting T cells in PBMCs. PBMCs (10^6) of each healthy donor were incubated with each of the synthetic epitopes in ELISPOT plates. The epitope-specific CD4+ T cell responses were determined after 24 hours. Each spot-forming unit (SFU) corresponds to one IFNγ-secreting T cell. Results are expressed as numbers of SFU per 10^6 PBMC. The negative control was PBMCs in medium without epitope stimulation and was used to evaluate the spontaneous secretion of IFNγ. The positive control was polyclonal activator anti-CD3 monoclonal antibody, which determine T cell numbers and viability of the immunoassay.

3. Results

3.1 Multiple sequence alignment and antigen selection

Initially, coronavirus surface glycoprotein was studied for determining specific conserved part of protein between the virus serotypes. Results of protein BLAST are shown in Table 1. The results demonstrated that coronavirus surface glycoprotein had the most conservancy levels between 97. 8 and 100. Also, VaxiJen score of the protein showed high antigenicity. Due to having good antigenicity, high exposure probability to the immune system and high conservancy, this protein was selected for vaccine design.

3.2 T-cell and B-cell epitope prediction

In the current study, some appropriate common B-cell and T-cell epitopes were designed. The predicted MHC-I and MHC-II restricted epitopes were compared to B-cell epitopes to determine shared epitopes (Table 2). Finally, 17 epitopes with 9-16 amino acids were selected. These epitopes are located between residues 14-642. These epitopes exhibited a relatively high aliphatic index (>60), high antigenicity (>1.6) and low instability index (less than 20). The most potent epitope was repeated three times and merged into the other epitopes using suitable linkers (KK, AYY) for organization of final vaccine construct.

3.3 Antigen selectivity of constructed vaccines

Final construct vaccine was composed of 398 amino acids which were respectively attached to 50S ribosomal protein L7/L12 as adjuvant. The antigenicity score of constructed vaccine are shown in Table 3. The result demonstrated that IV1 has 1.2110 antigenicity (Table 3).

3.4 Physicochemical properties

Physicochemical properties of the constructed vaccine was predicted using Protparam server. The results revealed that this multi-epitope vaccine have low instability "as a value below 40" predicts that the protein is stable. The IV1 construct showed the highest Isoelectric point with 8.52 value. From the aliphatic Index of view, this construct showed aliphatic index more than 90 % (Table 4). The results of gor4 demonstrated that the random coil values of the IV1 were high compared to Alpha helix and extended structure.

3.5 Analysis of docking results

The results of docking the constructions with HLA-1 and HLA-2 confirmed the high values with IV1 construct. Also, the results of TLR-3 analysis showed values of 1680.569 for IV1 which demonstrate the high efficacy of vaccine construct (Tables 5). Figure 2 indicate docking results of the Vac1 construct with

the TLR-3 as the examples of vaccines potential for interaction. The results showed that TLR-3 have
interaction to Glu115, Gly 118 residues of AV1.

2.10 IFNγ-ELISPOT assay

We examined the numbers of primed CD4+ cells that produced IFN-γ in response to epitopes, using
ELISPOT assays described in Table 5. The CD4+ T cellular memory responses of highest magnitude were
obtained for epitopes 10, 11 and 12 with mean SFU across all responding donors as: 35.0, 37.3 and 38.2
respectively. The results demonstrated that these donors showed very low response to epitopes 17 and 1
just above the cut-off (11 and 15 SFU respectively). The IFN-γ producing T cell variation ranged from
11.1 ±1.2 SFU to 38.2 ± 2.1 SFU per 10^6 PBMCs among all five donors. Only four donors out of five
donors (D5, D10, D11, and D14) responded to Epitope 1, 2, 4 and 17 conserved epitopes and the responses
were of a low magnitude (Table 6).

4. Discussion

Due to the nature of coronavirus and high infectious rate, the progress of a vaccine against coronavirus is
very challenging. Though, with the development of computational methods, these limitations are reduced.
By the way, using computational methods, the design of recombinant vaccines and the estimation of
physicochemical properties as well as vaccines efficacy could be available [20] [21]. Then, this research
was intended to design an effective multi-epitope recombinant vaccine against coronavirus using a unique
multi-step bioinformatics approach. Our potent multi-epitope vaccine is contained seventeen epitopes in
surface glycoprotein of coronavirus along with AAY and KK linkers and 50S ribosomal protein L7/L12 as
adjuvant. Based on our knowledge, there is no report about computational design of epitope-based vaccine
for coronavirus. Recently, in silico, design of epitope-based vaccine was used for vaccine development
against several infectious diseases. Several bioinformatics tools have been established that accelerate the
growth of multi epitope-based vaccines. In recent decade, several multiepitope vaccines for pathogenic
viruses have been reported. Multi epitope vaccines could provide an effective immunization against
different serotypes of a pathogen. Despite mentioned advantages of Multi epitope vaccines, poor
immunogenicity is considered as a major drawback to growth of these vaccines [22], [23]. The in-silico
results proposed that our multi-epitope vaccine was very stable with high aliphatic index and it was
potentially antigenic. As reported earlier, high aliphatic index shows the higher thermos-stability of the
constructed vaccine. At the present research, the aliphatic index was high, and instability was low and
Gravy indices were negative. Also, the higher PI, as the case of this study, shows the higher potential for
cell wall attachment. However, the proposed vaccine has high antigenicity. This was chosen for docking
studies. The results demonstrated that our mentioned vaccine could be a right candidate for experimental
research [24].

Conclusion

This study introduced designing novel multi epitope vaccines against Coronavirus which could cover
conserve sequence of the virus. The multi-epitope vaccine presented by this study showed promising
result through in silico step, which could be followed by in vitro and in vivo studies.

Conflict of interest

Authors declare no conflict of interest.

Acknowledgment

We wish to thank the University of Isfahan for their supports.

References

1. Li Q., Guan X., Wu P., Wang X., Zhou L (2020) Tong Y., Feng Z. Early transmission dynamics in
Wuhan, China, of novel coronavirus-infected pneumonia. New England Journal of Medicine.

2. World Health Organization Novel coronavirus (2019-nCoV) situation reports. 2020.
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports

3. Cicala C, Nawaz F, Jelicic K, et al. (2016) HIV-1 gp120: A Target for Therapeutics and Vaccine
Design. Current Drug Targets. 17(1):122-35.

4. Kumar M, Thakur V, Raghava GP (2008) "COPid: composition based protein identification. In silico
b10.Soria-Guerra RE, Nieto-Gomez R, Govea-Alonso DO, et al. An overview of bioinformatics tools for
epitope prediction: implications on vaccine Development. J Biomed Inform. 2015;53, 405–414. doi:
10.1016/j.jbi.2014.11.003.iology. 8:121-128.

5. Soria-Guerra RE, Nieto-Gomez R, Govea-Alonso DO (2015) t al. An overview of bioinformatics tools
for epitope prediction: implications on vaccine Development. J Biomed Inform. 2015;53, 405–414. doi:
10.1016/j.jbi.2014.11.003.

6.Bissati KE, Chentoufi AA, Krishack PA (2016) Adjuvanted multi-epitope vaccines protect HLA-A* 11:
01 transgenic mice against Toxoplasma gondii. JCI Insight. 1(15): doi: 10.1172/jci.insight.85955.

7.Escalona E, Saez D, Onate A (2017) Immunogenicity of a multi-epitope dna vaccine encoding epitopes
from Cu–Zn superoxide dismutase and open reading Frames of Brucella abortus in mice. Frontiers
Immunology. 8, 125

8.Rodrigues-da-Silva RN, Martins da Silva JH, Singh B (2016) In silico identification and validation of a
linear and naturally immunogenic B-cell epitope of the Plasmodium vivax malaria vaccine candidate
merozoite surface protein-9. Plos One;11(1): e 0146951. doi: 10.1371
221

9.Nezafat N, Karimi Z, Eslami M (2016) Designing an efficient multi-epitope peptide vaccine against
Vibrio cholerae via combined immunoinformatics and protein interaction based approaches.
Computational Biology and Chemistry. 62: 82–95. doi:10.1016/j.compbiolchem.2016.04.006

10.Yang Y, Sun W, Guo J, et al. In silico design of a DNA based HIV-1 multi-epitope vaccine for
Chinese populations. Human Vaccines Immunother. 2015;11(3): 795–805. doi:
10.1080/21645515.2015.1012017

11.Nosrati M, Mohabatkar H, Behbahani M (2017) A novel multi-epitope vaccine for cross protection
against Hepatitis C Virus (HCV): an immunoinformatics. Approach Research in Molecular Medicine
5(1): 17–26.

12.Rahjerdi AK, AmaniJ, Rad I (2016) Designing and structure evaluation of multi-epitope vaccine
against ETEC and EHEC, an in silico approach.Protein Peptide Letters. 23(1): 33–42.

13.Oscherwitz J (2016) The promise and challenge of epitope-focused vaccines. Human Vaccines
Immunother.; 12(8): 2113–2116.

14. Conrad C. Huang, Elaine C. Meng, John H. Morris (2014) Enhancing UCSF Chimera through web
services. Nucleic Acids Research. 42(1):478-484. https://doi.org/10.1093/nar/gku377

15.Doytchinova IA, Flower DR (2007) VaxiJen: a server for prediction of protective antigens, tumour
antigens and subunit vaccines. BMC Bioinformatics.

16.Duquesnoy R, Marrari M (2017) Usefulness of the ElliPro epitope predictor program in defining the
repertoire of HLA-ABC eplets. Hum .78(7-8): 481-488. doi:10.1016/j.humimm.03.005

17.Beaver JE, Bourne PE, Ponomarenko JV (2007) Epitope Viewer: a Java application for the
visualization and analysis of immune epitopes in the Immune Epitope Database and Analysis Resource
(IEDB). Immunome   Res   DOI:10.1186/1745-7580-3-3.

18. Patronov AI. Doytchinov I (2013) T-cell epitope vaccine design by immunoinformatics. Open Biol.
2013; 3(1):120139. doi: 10.1098/rsob.120139.

19.Salentin S, Schreiber S, Haupt VJ (2015) PLIP: fullyautomated protein–ligand interaction profiler.
Nucleic Acids Research ;43: W443–W447.https://doi.org/10.1093/nar/gkv315
248
249
250   20. Roosa K. , Lee Y., Luo R., Kirpich A., Rothenberg R., et al (2020) Hyman J.M., Yan P., and G.
251   Chowell, Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th,
252   2020, Infect Dis Model. 2020; 5: 256–263.
253
254
255   21. Ai S., Zhu G., Tian F., Li H., Gao Y., Wu Y., Lin H (2020) Population movement, city closure and
256   spatial transmission of the 2019-nCoV infection in China.
257
258

22.Paul S, Piontkivska H (2010) Frequent associations between CTL and T-Helper epitopes in HIV-1
genomes and implications for multi-epitope vaccine designs. BMC Microbiology. 10:212. doi:
10.1186/1471-2180-10-212

23.Hajighahramani N, Nezafat N, Eslami M, et al. (2017) Immunoinformatics analysis and in silico
designing of a novel multiepitope peptide vaccine against Staphylococcus aureus. Infection Genetics and
Evolution. 48: 83–94. doi: 10.1016/j.meegid

24.Solanki V, Tiwari M, Tiwari V. (2019) Prioritization of potential vaccine targets using comparative
proteomics and designing of the chimeric multi-epitope vaccine against Pseudomonas aeruginosa.
Scientific Reports. 9.

278   Fig 1: Molecular docking analysis of IV1construct with TLR-3. The results showed that TLR-3
279   have interaction to Glu 115 and Gly 118 residues of AV1

287 Table 1: Results of the antigenicity and BLAST of coronavirus surface glycoprotein and
288 antigenicity prediction

Protein	Average antigenicity	VaxiJen score	Minimum identity (%)	Maximum identity (%)
Coronavirus surface glycoprotein	0.4646	0.4	97.8	100

291 Table 2: Result of final T-cell and B-cell epitope prediction screening from coronavirus surface
292 glycoprotein

	Sequence	Length	Start-End	Vaxijen	Isoelectric point	Aliphatic index	Instability	GRAVY
1	KLNDLCFTN	9	386-394	2.90	5.83	86.67	17.24	-0.24
2	KLNDLCFTNV	10	386-395	2.69	5.83	88	-14.52	-0.2
3	KLNDLCFTNVY	11	386-396	2.22	5.83	97.27	-19.15	-0.06
4	PTKLNDLCFTN	11	384-394	2.19	6.22	70.91	-12.29	-0.4
5	SPTKLNDLCFTN	12	383-394	2.01	5.55	65	26.18	-0.4
6	VSPTKLNDLCFTN	13	382-394	2.17	5.80	82.32	24.94	-0.08
7	KLNDLCFTNVYA	12	386-397	1.92	5.83	97.5	3	0.2
8	KLNDLCFTNVYAD	13	386-398	1.67	4.21	90	-2.98	-0.07
9	LNDLCFTNV	9	387-395	2.01	3.80	118.89	-7.81	0.6
10	GVSPTKLNDLCFTN	14	381-384	2.21	5.83	76.43	23.87	-0.1
11	YGVSPTKLNDLCFTN	15	380-384	2.06	5.83	71.33	17.29	-0.18
12	CYGVSPTKLNDLCFTN	16	379-384	2.01	5.82	66.88	16.83	-0.01
13	CVNLTTRTQ	9	15-23	1.87	8.25	75.56	-7.87	-0.34
14	QCVNLTTRTQ	10	14-23	1.78	8.25	68	-13.62	-0.66
15	LDITPCSFGGVSV	13	585-697	1.88	3.80	104.62	12.92	1.06
16	LDITPCSFGGVSVI	14	585-698	1.61	3.80	125	12.71	1.03
17	VKNKCVNFN	9	534-642	2.05	9.31	64.44	6.92	-0.51

300 **Table 3:** Average antigenicity of constructed vaccine using Vaxijen

Vaccine	Sequence	Vaxijen
IV1	**MSDINKLAETLVNLKIVEVNDLAKILKEKYGLDPSANLAIPSLPKAEILDKSKEKTSF DLILKGAGSAKLTVVKRIKDLIGLGLKESKDLVDNVPKHLKKGLSKEEAESLKKQL EEVGAEVELK****EAAAK****KLNDLCFTN****AAY****KLNDLCFTN****AAY****KLNDLCFTN****AAY****KLND LCFTNV****AAY****KLNDLCFTNVY****AAY****PTKLNDLCFTN****AAY****SPTKLNDLCFTN****AAY****VSPT KLNDLCFTN****AAY****LNDLCFTNV****AAY****KLNDLCFTNVYA****AAY****KLNDLCFTNVYAD****AAY** **GVSPTKLNDLCFTN****AAY****YGVSPTKLNDLCFTN****AAY****CYGVSPTKLNDLCFTN****AAY****CV NLTTRTQ****AAY****QCVNLTTRTQ****KKL****DITPCSFGGVSV****KKL****DITPCSFGGVSVI****KKV****KN KCVNFN**	1.2110

308 **Table 4:** Physico-properties of constructed vaccines
309

Protein	Molecular weight	Isoelectric point	Aliphatic index	GRAVY	Instability	Alpha	Extended	Random coil
IV1	43679.43	8.52	92.71	-0.046	11.85	23.37%	8.54%	68.09

312 **Table 5.** HDOCK scores of HLA1, HLA2 and interaction residues of selected sequences

Protein	Hla1-A1	Hla1-Chain G	Hla1-Chain E	Hla1-Chain F	Hla1-Chain CW-1	TLR3	Hla2 - antigen gama chain	Hla2-DM	Hla2-DO	Hla2-DP	Hla2-DQ	Hla2- DR
IV1	951.2	957.2	951.2	1215.7	1220.8	1680.5	1215	1214	1215.7	1215.7	1215.7	1215.7

316
317
318
319

320
321

Table 6: Mean number of responding T cells (SFU) of donors to predicted and conserved T cell epitopes encompassing entire HA protein

Donor	EP1	EP2	EP3	EP4	EP5	EP6	EP7	EP8	EP9	EP10	EP11	EP12	EP13	EP14	EP15	EP16	EP17
1	15.1 ±2	16.4 ±1.9	22.2 ±1.8	17.4 ±2	27.1 ±2.2	24.9 ±2	23.4 ±1.6	2± 1.5	23.2± 1.6	35.0± 2.9	37.3 ±2.6	38.2 ±2.1	25.3 ±2	24.4 ±1.7	24.1 ±1.7	22.1 ±1.8	11.1 ±1.2