ZENITH: A flow cytometry based method for functional profiling energy metabolism with single cell resolution

Rafael J. Argüello[1∞*], Alexis J. Combes[2,3∞], Remy Char[1], Evens Bousiquot[1], Julien-Paul Gigan[1], Voahirana Camosseto[1,7], Bushra Samad[2,3], Jessica Tsui[2,3], Peter Yan[2,3], Sebastien Boissoneau[4], Dominique Figarella-Branger[5], Emeline Tabouret[6], Evelina Gatti[1,7,8], Matthew F. Krummel[2,3,§] and Philippe Pierre[1,7,8,§]

[1] Aix Marseille Univ, CNRS, INSERM, CIML, Centre d'Immunologie de Marseille-Luminy, Marseille, France.
[2] Department of Pathology, University of California, San Francisco, San Francisco, CA, USA.
[3] ImmunoProfiler Initiative, University of California, San Francisco, San Francisco, CA, USA.
[4] Aix-Marseille Univ, Institut de Neurosciences des Systems, Faculté de Medecine, Marseille France.
[5] Aix-Marseille Univ, APHM, CNRS, INP, Inst Neurophysiopathol, CHU Timone, Service d'Anatomie Pathologique et de Neuropathologie, Marseille, France.
[6] Aix-Marseille Univ, APHM, CNRS, INP, Inst Neurophysiopathol, CHU Timone, Service de Neurooncologie, Marseille, France.
[7] Institute for Research in Biomedicine (iBiMED) and Ilidio Pinho Foundation, Department of Medical Sciences, University of Aveiro, 3810-193 Aveiro, Portugal.
[8] International associated laboratory (LIA) CNRS "Mistra", 13288 Marseille cedex 9, France

[∞] [§] These authors contributed equally to this work.
*To whom correspondence should be addressed: RJA arguello@ciml.univ-mrs.fr

Abstract

Energetic metabolism (EM) reprogramming is critical for both cancer and immune response initiation. Current methods for EM monitoring are performed in bulk and do not have the sensitivity nor the resolution to dissect EM at single cell scale. We designed a novel EM profiling method called ZENITH, that provides single-cell resolution using multiparametric flow cytometry. ZENITH highlights the dichotomy among EM displayed respectively by naïve, effector and memory T cells, demonstrating its ability to discriminate individual cell subsets on the basis on their specific metabolic status. Tumor-associated myeloid cells isolated from biopsies are also shown to display highly variable EM profiles, in ways that are not linked to their lineage nor their activation phenotype. ZENITH ability to determine complex and linked metabolic activities in discrete cell subpopulations within tissue or blood draws will contribute to the functional information needed for evaluating therapeutic responses or patient stratification.

Introduction

Energetic metabolism (EM) comprises a series of interconnected biochemical pathways capable of using energy rich molecules to produce ATP. In the presence of enough oxygen, cells can produce ATP either by oxidative phosphorylation (OXPHOS) and/or by performing glycolysis. Glycolytic metabolism in the presence of non-limiting concentration of oxygen is called aerobic glycolysis and is characteristic of proliferating cells. Aerobic glycolytic metabolism not only supports proliferation but also cell survival in hypoxic conditions. Immune cells are specially adapted to migrate into peripheral tissues and change of microenvironment. Their energetic metabolism profile is known to correlate with the microanatomical localization, activation, proliferation or functional state[1,2]. Activation and proliferation drives the differentiation of long lived quiescent naïve into effector T cells. This process is tighly linked to a metabolic switch from OXPHOS to aerobic glycolysis[3,4]. Interestingly, naïve cells re-circulate between the blood and lymph nodes while blood circulating effector cells need to reach peripheral tissues (i.e. infection site or tumors). Supporting the idea that metabolic profile is pre-adapted and is related to cell function, post-mitotic blood immune cells whose function is to rapidly migrate into hypoxic/damaged tissues, such as circulating neutrophils and monocytes, are already engaged in aerobic glycolysis in the blood. Altogether, the current paradigm suggests that a particular metabolic profile of the cells in the blood and tissues is associated to their competence to migrate, proliferate and exert their effector functions in target tissues. However, due to technical limitations and the heterogeneity of cell types, it is not clear which subsets of immune cells rapidly adapt to changing environments or which have predefined metabolic profile.

Competition for glucose within the tumor microenvironment influences cancer progression and the anti-tumoral immune response by regulating EM in both tumoral cells and tumor-

infiltrating lymphocytes (TILs), consequently influencing their function[5]. The presence of effector immune cells into target tissue has clinical relevance and is a well accepted predictive parameter to determine response to immunotherapy[6]. The success of immunotherapies (e.g. checkpoint inhibitors) is restricted to a relatively small proportion of patients that require a functional and metabolically competent immune system to respond to treament[7,8]. Consequenly, the need of better and accessible tools for patient immuno-metabolic profiling as a strategy for stratification, and monitoring responses to immunotherapies is strongly needed. The metabolic profile of innate and adaptive immune cells correlates with the type of cytokines they produce , and from this point of view metabolic profiles represent a read-out for functional states.

Current methods to determine EM profiles can be classified in three groups. The first group epitomized by Seahorse® (Agilent technologies), uses metabolic inhibitors (i.e. 2-DeoxyGlucose/"DG" and Oligomycin A/"O") during monitoring of the extracellular acidification rate (ECAR), as well as oxygen consumption rate (OCR) of a large amount of cells in culture, to establish the dependency and metabolic capacities of the cells on different EM pathways. The second group, also a bulk measurement, quantifies the activity of enzymes implicated in particular metabolic pathways in fixed cells or lysates[9]. Finally, the third group uses mass spectometry and mass-spectrometry imaging to measure the levels of different metabolites produced by EM pathways[10]. Very recently, CyTOF-based quantification of different metabolic enzymes was used in parallel to histology to correlate expression of metabolic enzymes with the microanatomical localization of mouse and human T cells[11–13].

Current methods need large numbers of highly purified cells[14], and dedicated equipments, and thus cannot be applied to establish functional EM profiles of heterogenous and scarce living cell populations obtained from human samples or biopsies[15] (Table 1). Given the unpredictability of functional metabolism by phenotyping the levels of a relatively small subset of enzymes in different cell types, we focused on developing a method that functionally quantifies energetic metabolism. Approximatively half of the total energy that mammalian cells produce by degrading glucose, aminoacids and/or lipids is immediately consumed by the protein synthesis machinery[16–19]. The tremendous energetic cost associated with mRNA translation offers a methodological opportunity of determining the protein synthesis levels as a measure of global EM activity, and its response to environmental cues or different metabolic inhibitors. We took advantage of the drug puromycin, whose incorporation is a highly reliable readout for measuring protein synthesis levels[20–23], combined to a novel anti-puromycin monoclonal antibody, to develop a reliable method to perform EM profiling with single cell level resolution based on protein synthesis intensity as read-out. We termed this method ZENITH, (Zelle Energie von Translation Histoanalyse) meaning "histoanalysis of cell energy via translation", with reference to our previous SUnSET[24] and SunRiSE[25] methods for studying translation. ZENITH was used to deconvolve the complex energetics of blood T cells and human tumors-associated myeloid cells, with a much-needed resolution for analysing these highly heterogenous samples, from which the circuitry for metabolism, particularly amongst immune cell subsets, has remained inaccessible.

Results

Characterizing the energetic metabolism profile by monitoring changes in protein synthesis levels in response to metabolic inhibitors.

To test whether the level of protein synthesis (PS) and the pool of ATP are tightly kinetically coupled, we measured in mouse embryonic fibroblasts (MEF), on the one hand ATP levels using the CellTiter-Glo® luminescence based system, and on the other hand PS levels via puromycin incorporation and staining, after blocking ATP production (Fig. 1a). To completely inhibit ATP production, we treated cells with a mix of inhibitors that block both glycolysis and oxidative phosphorylation (OXPHOS); 100mM 2-Deoxy-D-Glucose (DG), 1µM FCCP and 1µM Oligomycin A (O) (Fig. 1a). To further optimize the signal to noise ratio of puromycin intracellular detection, we developed a novel monoclonal anti-puromycin antibody (clone R4743L-E8, Rat IgG2a) specifically adapted to intracellular flow cytometry. Both protein synthesis levels (Fig. 1b,d) and ATP levels (Fig. 1c) dropped within 5-10 minutes after blocking ATP synthesis, with a strikingly similar slope, showing that changes in ATP levels and PS levels are tighly correlated (Fig. 1e; r 0.985; P<0.0001). To test the relationship between ATP consumption and translational and transcriptional activities, we treated cells with the same inhibitors to block de novo ATP synthesis, together with

Table 1. Comparative table of different methods to profile metabolism.

Method	CyTOF	MSI	Seahorse®	ZENITH
Functional profile of the cells (# of treatments)	NO	NO	YES (4)	YES (non-limited)
Cell purification required	NO	NO	YES	NO
Single cell resolution	YES	YES	NO	YES
Fixation compatible	Yes	YES	NO	YES
Stability of the readout	High	Variable	Low	High
Phenotypic analysis	YES	NO	NO	YES
Metabolite profiling	NO	YES	NO	NO
Compatible with cell sorting	NO	NO	NO	YES*
***In vivo* application**	NO	NO	NO	YES$
***Ex vivo* application**	YES	YES	NO	YES
***In-vitro* application**	YES	YES	YES	YES
Quantitative & Quantitative	NO	YES	YES	YES
Tissue spatial resolution	NO	YES	NO	NO
Metabolic Readout	Enzyme expression levels	Metabolite levels	Changes in extracellular pH and $[O_2]$	Changes in Protein synthesis levels
Time (Hs) from sampling to metabolic profiling	NA	NA	24	1
Total time	6	6	24	7
# cells required in subsets	500	200	1.000.000	2000
Accessibility	Low	Low	Low	High
Equipment needed	CyTOF cytometer	MIBIscope® IONpath	Seahorse® Analyser	Any Flow cytometer#

* Not shown
$ Potential application
# ZENITH has also the potential to be analyzed by CyTOF, MSI, Microscopy, not done.

Figure 1.

a, Blocking ATP synthesis and kinetic analysis of ATP and protein synthesis levels.

b, Histogram showing protein synthesis levels by puromycin incorporation by flow cytometry in MEFs after blocking both mitochondrial respiration and glucose oxidation (Oligomycin 1uM and 2-DG 0,1M) for different amounts of time.

c, ATP levels vs. time after blocking ATP synthesis (n=3). ATP concentration in cell lysates was calculated from ATP standard curve.

d, Protein synthesis levels by flow cytometry vs. time after blocking ATP synthesis in MEFs (n=3).

e, Correlation between changes in ATP levels and protein synthesis levels (Pearson R=0,985, P<0,0001).

f, ZENITH is based on the principle that if a cell depends on a particular pathway to produce NTPs, then an inhibitor of that pathway will induce a drop in ATP and protein synthesis. The glucose dependence and the Fatty and amino-acids oxidation capacity; and the mitochondrial dependence and glycolytic capacity can be calculated from the MFI of Puromycin in the different treatments following the formulas (for more details see materials and methods).

g, ZENITH procedure starts by extracting the tissue, dividing the sample and treat with the inhibitors and the puromycin after. After staining and FCM, the profile of response of the different cells subsets present in a sample can be analyzed in parallel without the need of cells purification. The profile of response to the inhibitors of cells with different metabolism is different.

translation and/or transcription inhibitors. Altogether, our results confirmed that protein synthesis is one of the most energy consuming metabolic activities (Supplementary fig. 1), and most importantly, it represents a stable and reliable readout to evaluate rapidly the impact of metabolic pathways inhibition on the cell.

As both ATP and PS levels are kinetically coupled, puromycin fluorescence (measuring PS) can act as a surrogate for energetic status in the context of flow cytometry. The principle of ZENITH is thus to incubate a given sample in parallel with specific inhibitors of known EM pathways. If a cell population is energetically dependent on a particular pathway, its ATP content will immediately drop, and so will its protein synthesis levels. The latter will be established using puromycin incorporation and its detection with fluorescent anti-puromycin antibody (Fig. 1f and Supplementary fig. 1b). ZENITH allows EM profiles to be measured in heterogenous cell populations at single cell resolution by combining cell identification and puromycilation detection by multiparametric flow cytometry (FCM) (Fig. 1g). Using PS levels, the level of glucose dependence (Gluc. Dep.) can be calculated to quantify the proportion of the PS, and therefore of ATP/GTP production, dependent on glucose oxidation (Fig. 1f, see Materials and Methods). The mitochondrial dependence (Mitoc. Dep), namely the proportion of PS dependent on oxidative phosphorylation, can be similarly established. Two additional derived parameters, "Glycolytic capacity" (Glyc. Cap.) and "Fatty acids and amino acids oxidation capacity" (FAAO Cap.) can also be calculated. Glycolytic capacity is defined as the maximum capacity to sustain PS when mitochondrial OXPHOS is inhibited (Fig. 1f, see statistic in Materials and Methods section). Conversely, FAAO Capacity is defined as the capacity to use fatty acids and aminoacids as sources for ATP production in the mitochondria, when glucose oxidation is inhibited (Glycolysis and Glucose derived Acetyl-CoA by OXPHOS) (Fig. 1f and Supplementary fig. 1b).

ZENITH recapitulates Seahorse® EM profiling of steady state and activated T cells.

The metabolic switch of T cells to aerobic glycolysis upon activation was originally documented in the 1970s[3] and more recently confirmed using the Seahorse® technology [26,27]. To benchmark our method, we monitored the variations in EM observed in isolated bulk human blood T cells at steady state or upon activation by

Seahorse® in parallel to ZENITH (Fig. 2a). Upon activation, an increase in the glycolytic capacity of T cells was measured with both methods (Fig. 2b and 2c, respectively) in excellent agreement (correlation Spearman r squared 0.85, P<0.01) (Fig. 2d). We observed a significant decrease in the spare respiratory capacity in bulk T cells upon activation with Seahorse® (Supplementary fig. 2a and 2b). Interestingly, an increase in oxygen consumption rate (OCR) by Seahorse® , was paralleled with a strong increase in the global level of PS measured by ZENITH although to a larger extent (Fig. 2e and 2f, respectively). Overall, the EM profiles of T cells upon activation obtained by Seahorse® and by ZENITH were therefore very consistent. The level of translation (Fig. 2f) correlated with the global metabolic activity of the cells, and changes in the response to inhibitors confirmed the metabolic switch towards aerobic glycolysis that occurs upon T cell activation. However, ZENITH showed two main advantages over Seahorse® measurements. First, the magnitude of the change in the glycolytic capacity and the standard error of the measurements with ZENITH were superior (Fig. 1b vs 1c, 1e vs 1f). Second, ZENITH analysis was performed with 10 fold less T cells ($1,2.10^5$ in triplicates vs. $1,2.10^6$ cells, respectively). Moreover, ZENITH could incorporate a full spectrum of T cell markers in the analysis allowing to study in parallel the different CD3[+] T cells subpopulations present into the bulk sample (Fig. 2), encompassing naïve, memory and effector CD4[+] or CD8[+] T cells subsets.

Metabolic deconvolution of blood T cell subsets by ZENITH identifies a memory CD8[+] T cells subset constitutively displaying high glycolytic capacity.

To expand upon the ability to deconvolve T cell subpopulations, we next applied ZENITH to mixed populations that due to heterogeneity are inaccessible to EM monitoring by Seahorse®. We took advantage of CD45RA, IL7RA (CD127), CCR7, CD45RO, CD57, PD1, and Perforin expression to identify and analyse the naïve and memory CD4+ and CD8+ T cell subsets present in total human blood draws. Immunodetection of these nine markers yielded six phenotypically distinct clusters/subpopulations with different abundances (Fig. 3a and 3b). The metabolic profiles of non-activated naïve T cells (CD8 or CD4), as well as memory (EM and CM) CD4 and highly differentiated CD8 (HDE) showed a medium-high degree of mitochondrial dependence (Fig. 3c), consistet with what was

Figure 2.

a, Parallel Seahorse and ZENITH metabolic analysis of resting and activated T cells.

b, Metabolic profile of $1,2.10^6$ T cells from different subjects (P1, P2, P3) with Seahorse. Profile of ECAR of P1 is shown (N=3, each in triplicates, *P<0,05; **P<0,01).

c, Metabolic profiling of $1,2.10^5$ T cells from different subjects (P1, P2, P3) with ZENITH. Profile of translation of P1 is shown (N=3, each in triplicates, ***P<0,001).

d, Correlation between the changes in glycolytic capacity of steady state and activated (aCD3/CD28) T cells from three different subjects measured by Seahorse and ZENITH (P1, P2, and P3, Pearson r=0,92; R^2=0,85; P<0.01).

e, Basal Oxygen Consumption Rate (OCR) in non-activated (non-Act) and activated T cells (aCD3/aCD28). Each bar represents the mean of P1, P2, and P3 (in triplicates).

f, Basal translation levels (anti-Puro, Geometric Mean Fluorescence intensity) in non-activated (non-Act) and activated T cells (aCD3/CD28). Each bar represents the mean of P1, P2, and P3 (in triplicates)

Figure 3.

Metabolism profile of resting human blood T cells by ZENITH identifies different metabolic profile of human T cells subsets.

a, ZENITH analysis pipeline of T cells purified from human blood (95% pure). Dimensionality reduction (t-SNE) based on phenotypic markers is performed to the concatenated FCM file containing treated cells (Co, DG, O, Z). Metabolic deconvolution is performed to extract the level protein synthesis of each phenotypic cluster of T cells in response to the inhibitors.

b, List of markers and relative levels of expression (Geometric Mean fluorescence intensity heatmap) in each cluster from the t-SNE after dimensionality reduction.

c, Metabolic profile of the different T cells subsets identified in the t-SNE reveals heterogeneous metabolism and the presence of subsets with high glycolytic capacity in non-activated circulating blood T cells. Anti-Puromycin MFI (translation) from P1 are shown and representative of all three healthy donors assessed.

d, T cells treated with mitochondrial inhibitor (left panel) containing glycolytic and mitochondrial dependent subsets. Histogram (middle panel), showing the level of translation in all T cells (light grey line) upon mitochondrial inhibition, indicating the presence mitochondrial independent ("Glycolytic" cells subsets, red) and mitochondrial dependent cells (blue). Gating on Glycolytic and Mitochondrial dependent cells and shown into the t-SNE plot (right panel) to identify the phenotype of "Glycolytic" T cells gate (red) and phenotype of the "Mitochondrial dependent" cells (blue). One marker that correlates with the metabolic profile is the marker of antigen experience CD45RA, that is lost in cells that have been previously exposed to TCR stimulations.

previously reported on their metabolic activity[4]. In contrast, the less abundant cell subsets such as CD8 early effector memory (EEM) and Natural Killer (NK) cells (that co-purfied with T cells and represented just 5% of the cells) showed higher glycolytic capacity. To determine if similar metabolic trends are observed in other species and preparations, we performed ZENITH on resting and activated mouse splenic T cells (Supplementary fig. 3a and 3b) and human blood central memory CD4 T cell subsets (Supplementary fig. 3c), demonstrating a consistent switch towards high glycolytic capacity and high glucose dependence in both mouse and human T cells upon activation (Supplementary fig. 3). During bulk analysis, naïve cells represented 42%, (the majority out of 78%) of the T cells and thus likely dominate the EM monitoring performed by Seahorse® (Fig. 2e and Fig. 3c). Consequently Seahorse® measurements indicate a rather low "mean" glycolytic rate/capacity and high "mean" oxygen consumtion rate (Fig. 2b) and is thus in accordance with the metabolism of naïve T cells determined by ZENITH (Fig. 3c, 3d). However, the presence of CD8[+] EEM that display high glycolytic capacity, but represent no more that 5% of the T cells present in the sample (2000 cells) remained completely masked during Seahorse® analysis.

Another important feature of the multiparametric ZENITH resolutive power is the possibility to feature single cell behaviors according to their sensitivity to metabolic inhibitors independently of their phenotype. This allows to identify functional metabolic heterogeneity first, and to determine the phenotype or sort different cells afterwards. As a proof of concept, resting purified T cells were treated with Oligomycin to inhibit mitochondrial respiration prior to translation monitoring. The histogram plotting translation levels showed two T cell subpopulations, one with high and one with low levels of translation (Fig. 3d). The population that showed high level of translation upon mitochondrial inhibition were labeled as "Glycolytic" (Fig. 3d, in red) and the cells with blocked translation as "Mitochondrial dependent" (in, blue). As shown using t-distributed stochastic neighbor embedding (t-SNE), the phenotype of Glycolytic and Respiratory T cells recapitulated our previous results (Fig. 3a-c) and showed that the expression of CD45RA, mostly present in naïve T and NK cells, correlated well with "Mitochondrial dependence" (Fig. 3d). In conclusion, we found that ZENITH allows for both the measurement of the EM profile of known non-abundant cell subsets of interest, but also the sorting and identification of "unknown" cells with specific metabolic profile present within an heterogenous sample.

Metabolic profiling of mouse and human myeloid cell subsets.

Compared to T cell subsets, the metabolism profile of myeloid cell subsets from human and mouse tissue origins has been far less studied[28]. Among myeloid cell subsets, dendritic cells (i.e. DC1, DC2, DC3 and pDCs) are non-abundant professional antigen presenting cells (APCs) which serve as sentinels for the immune system. Each subset expresses a particular set of microbial pattern recognition receptors and are specialized in activation of CD8[+] (i.e. DC1), CD4[+] (i.e. DC2 and DC3) T cells and antiviral cytokine production (pDCs). For instance, Lipopolysaccharide (LPS) detection by TLR4 on DC2 and DC3 promotes activation by triggering a series of signaling cascades resulting in changes in gene expression, in membrane traffic and in energetic metabolism[29–31]. Dendritic cells patrol all tissues emanating from the blood stream where they represent a very small fraction of the PBMC, making the isolation of millions of DC1, DC2 or pDC from the same donor very challenging. We therefore decided to use ZENITH to perform EM profiling of blood myeloid cell subsets from heathly donors as well as of mouse bone marrow- and spleen-derived DCs stimulated or not to generate a detailed metabolic atlas of the myeloid cell populations present in mouse or human blood[24].

Following deconvolution, we first ranked myeloid cells by glucose dependency, finding that classical monocytes (Mono1, CD14[+]CD16[-]) were the most glucose dependent, whereas the DC precursors (DC5) were the least (Fig. 4a). These populations then lay near the extremes of mitochondrial dependence, where DC5 display the highest and Mono1 the lowest. However, we also observed examples of cell subsets (e.g. pDC and Mono2) that were highly glucose dependent as well as mitochondrial dependent and had only moderate glycolytic capacity. In contrast, some (DC1 and DC2) displayed relatively high glycolytic capacity and moderate glucose dependency, suggesting a certain degree of metabolic plasticity. Cells ranked in opposite order for FAAO capacity compared to glucose dependency, consistent with the idea that cells with low glucose dependence can sustain translation and energy production by free-fatty and amino acids oxidation.

To test whether a rapid metabolic switch occurs upon TLR4 activation in human myeloid cells, we

Figure 4.

a, Metabolic profile of human blood monocyte and dendritic cells subsets by ZENITH. (N=5 independent healthy donors) Statistical comparisons are: for Glucose dependence and FAAO capacity, Mono1, Mono2 and pDC vs. DC5 and DC1; and for Mitochondrial dependence and Gylcolytic capacity, DC5 vs. pDC and vs. DC1). Statistical significance * p<0.05; ** p<0.005; ****p<0.0001.

b, Metabolic profile of mouse bone marrow derived (FLT3L) by ZENITH (N=3) in non-treated vs LPS (100ng/ml) treated cells. Statistical comparisons are: DC1, DC2 and pDC, non-treated vs. LPS. Statistical significance * p<0.05; ** p<0.005; ****p<0.0001.

used an antibody panel for analyzing Mono1, Mono2, DC1, DC2 and pDCs in PBMCs treated with LPS. Most subsets did change their metabolic profile, while Mono1 maintained their glycolytic profile but increased their global level of translation (Supplementary fig. 4a). The only subset that changed its metabolic profile were the DC1 that showed a mild increase in glucose dependence and a moderate decrease in mitochondrial dependence (Supplementary fig. 4a).

To gain insight into the metabolic profile of mouse DC populations, we analyzed the metabolism profile of mouse bone marrow-derived DC1, DC2 and pDC (FLT3L-BMDC) at steady state and upon activation with LPS (Fig. 4b). In accordance to what was observed in steady state human blood DCs, mouse DC2 subset showed the highest glycolytic capacity, followed by DC1 and pDC. While Glucose dependence was relatively high in DC1 and DC2 from human and mouse (40-55% vs. 80-100%, respectively), an important difference was observed in the glucose dependency of mouse versus human pDCs (15% vs 60%). LPS treatment shifted EM profile towards lower mitochondrial dependence of DC1 and DC2. However only DC2, showed an increased global level of protein synthesis, probably reflecting the abundant TLR4 expression in this DC subset.. Considering that we analyzed DC isolated from human blood and derived in-vitro from mouse bone marrow, ZENITH EM profiles are surprisingly consistant. These results confirm that ZENITH allows to identify different cell populations sharing similar EM profile and that EM in DCs varies according to their state of activation.

Profiling the metabolic state of human tumor-associated myeloid cells.

Immunotherapies are a game changer in oncology yet only a fraction of patients shows complete immune-mediated rejection of the tumor. The variations observed in patients responses to treatment have created a strong need for understanding the functional state of tumor-associated immune cells (immunoprofiling)[6]. We thus used ZENITH to perform paralleled phenotypic and metabolic profiling of human tumor samples, and investigate the heterogeity of immune cell subsets by comparing tumors of diverse origins with tumor-free adjacent tissue. We analyzed PBMCs from healthy donors, explanted meningioma, brain metastasis (originated from a breast cancer), as well as renal

carcinoma tumors and renal juxtatumoral tissue. In the case of renal cancer, both ZENITH and single cell RNA seq analysis were performed in parallel on the same sample.

We observed 8 different myeloid populations in meningioma and 6 different subset in renal carcinoma (Fig. 5a and 5b, respectively; Supplementary fig. 5a and 5b)), that were all profiled by ZENITH (Fig. 5c). Upon clustering of the different cell subsets based on EM profiling, two groups emerged, a "Glycolytic cluster" and a "Respiratory cluster" (Fig. 5c and Supplementary fig. 5c). Mono1 and Neutrophils displayed glycolytic metabolism profiles in all blood samples and tumors tested (Fig. 5d). In contrast, Mono2, DC1 and DC2 showed relatively high glycolytic capacity when isolated from kidney tumor and juxtatumoral tissues, while these subsets showed high respiratory metabolism profile in the two brain tumors. Conversely, tumor-associated macrophages (TAM), showed high mitochondrial dependence, while juxta-tumoral macrophages displayed high glycolytic capacity (Fig. 5d), suggesting that tumor microenvironment modifies TAM EM. The decrease of glycolytic capacity in TAM as compared to juxta-tumoral macrophages was previously associated with increased immunosuppression in the tumor environment, tumor progression via both nutritional and immunological cues, and poor patient survival[32]. These results demonstrate again the analytical capacity and descriminative power of ZENITH, that reveals that in addition to the cancer type, the tumor anatomical origin could influence the metabolism of immune subsets, introducing an additional layer of heterogeneity in the tumor environment.

Linking scRNA-seq and energetic metabolism profile in tumor-associated myeloid cells.

We decided to link the descriminative power of ZENITH with single cell RNA-seq (scRNAseq) to support our results regarding the cellular complexity of the tumor environment and correlate EM profiling with gene expression. We compared in each individual population, the functional EM profile obtained by ZENITH with a metabolic gene expression pattern obtained by scRNAseq. We first identified expression signatures of specific glycolytic and respiratory genes that correlated with the functional metabolism profiles of different blood myeloid cells (Supplementary Fig. 5d). We then tested the expression of these glycolytic and respiratory metabolic genes (mRNA levels) in the

Figure 5.

a, Clusters and Composition (% in each color gate) of myeloid cell subsets (gated on CD45[+]/CD3[-]CD20[-]CD19[-]CD56[-]/Live-dead[-]/singlets; 19 markers t-SNE) using the myeloid panel observed in the human meningioma tumor sample.

b, Clusters and Composition of Renal carcinoma in the same gate as **a**; for this sample and juxta-tumoral tissue, myeloid cells were sorted and analyzed in parallel by single cell-RNA-seq.

c, Metabolic profile of each myeloid cell subset from each type of tissue is shown in the heatmap (columns). Clustering of subsets by their metabolic profile identifies two clusters of cells, Respiratory (blue bar, on top) and Glycolytic (red bar) cell clusters.

d, Reordering of the rows by cell type (and not metabolism profile) to identify changes in metabolism profile in the same cell subset in the blood, the different tumors and juxta-tumoral tissue.

e, Clusters of myeloid cells identified in the renal carcinoma and juxta-tumoral tissue by single cell RNA-seq (left panel). Expression of glycolytic and respiratory gene signatures in all cells extracted from the tumor. Table summarizing the results obtained by ZENITH and scRNA-seq in tumor and juxta-tumoral myeloid cells.

different tumor-associated myeloid populations (CD45[+]Lin[-]HLA-DR[+]). Sorted cells from the renal carcinoma and juxtatumoral tissue were subjected to scRNAseq using 10X Genomics Chromium paired with deep sequencing (Fig. 5e and Supplementary Fig. 5e). Analysis of 12,801 cells for the tumor and 2,080 for the juxta tumoral tissue yielded 6 and 5 high quality population clusters respectively. To rigorously identify the myeloid populations, we verified the expression of characteristic markers of these populations[33] and assigned cellular identities in the t-SNE representations (Supplementary Fig. 5e). We focused on 5 monocytes and macrophages clusters expressing MAFB and/or CSF1R, that were present both in tumor and juxta-tumoral tissue (Supplementary Fig. 5f). Expression of classical surface markers like FCGR3A/CD16 and CD14 (Supplementary Fig. 5g), confirmed that clusters 0 and 1 represent CD14[+]CD16[-] classical monocytes. Cluster 2 represents CD14[-]CD16[+] non classical monocytes (Mono2), while co-expression of CD14 and CD16 for the clusters 3 and 4 corresponds to macrophage-like populations. We performed differential expression (DE) unsupervised analysis for each of the myeloid cluster versus all the other clusters identified in the tumor bed and generated heatmaps for the top 5 most differentially expressed genes in the tumor (Supplementary Fig. 5h) and the juxta tumoral tissue (Supplementary Fig. 5h, right). In addition to highlighting key genes that contributed to the unbiased segregation of these populations in both tissue, we confirmed the high expression of macrophage specific genes by cluster 3 and 4, such as APOE, C1QC and RGS1. We next overlayed on the t-SNE plots our two EM gene signatures. Strongly correlating with ZENITH profiles, monocytes clusters (0,1,2) presented an enrichment in glycolytic genes signature both in tumor and juxta tumoral tissue. Conversely macrophages (cluster 3) showed high expression of the respiratory signature in the tumor, while, as predicted by ZENITH, this was not detectable in juxta-tumoral tissue (Fig. 5e). Monocytes-derived dendritic cells differently from macrophages, presented an enrichment in glycolytic genes signature both in tumor and juxta tumoral tissue (Supplementary Fig. 5i).

Altogether, these results indicate that tumor micro-environment specifically modifies functional EM of Macrophages by durably affecting metabolic gene expression. By correlating the results of scRNAseq analysis and ZENITH profiling on different blood myeloid cell subsets (Fig. 5f), we identify a functional gene signature (Supplementary Fig. 5d) that can be used to metabolically profile a variety of cell types and tissues, using gene expression.

Discussion

ZENITH is a flow cytometry-based method to establish simultaneously the phenotypic and energetic metabolism profiles of multiple cell types in parallel. The method is rapid, highly sensitive and consistent with other established techniques, including Seahorse®. As the treatments with different inhibitory drugs are performed in parallel, ZENITH can be used to monitor cellular responses to a variety of metabolites, inhibitory compounds and their combinations. Given that flow cytometers are available in most research institutes and hospitals, ZENITH represents an accessible method to perform functional EM profiling. Compared to other methods (see Table 1), its sensitivity, accessibility, single cell resolution, stability of the readout, short experiment time, compatibility with fixation and sorting, makes of ZENITH, an unrivalled aproach for studying energy metabolism in tissues and complex populations.

The resolutive power of ZENITH highlighted important variations amongst subsets of the same cell population. During our analysis of human blood myeloid subsets, we monitored myeloid subpopulations that are known to be precursors of DC1, pDC and DC3. The precursor of DC1 and pDC, called DC5, showed an EM with intermediate characteristics between DC1 and pDC, that are both metabolically polarized. Mono1 are precursors of DC3 and even if the metabolic profile is similar, Mono1 EM is more glycolytic. Our results suggest that changes in metabolism is embedded in the myeloid differentiation program and indicate that modulating metabolism in dendritic cells precursors might influence the generation of different dendritic cells subsets with potential clinical interest. The activation of human and mouse myeloid cells with LPS, showed that only cells expressing the LPS receptor/co-receptor (TLR4, CD14) show an increase in the global level of protein synthesis (human Mono1 and mouse DC2). In contrast, DC1 that do not respond to LPS, also showed a change in their metabolic profile, suggesting that bystander activation of DCs by Mono1 or DC2 derived cytokines can induce a change in EM profile, independently of TLR triggering.

Importantly, ZENITH can establish the metabolic profile of very scarce cells, as exemplified by the analysis of early effector T cells that represent around 5% of the total T cells isolated from blood (500 cells), thus representing a gain of sensitivity of aproximatively 800 fold compared to Seahorse® measurements. This gain was even more dramatic, when the metabolism profile of DC cell subsets was established from human tumor biopsies, in which they constituted aproximately 0,5% of the myeloid cell population. These results demonstrate the analytical capacity and discriminative power of ZENITH and its potential in analysing how the anatomical and tissue context could influence energetic metabolism of immune cell subsets, thus introducing an additional layer of heterogeneity in the healthy tissues and tumor environment. Thus, ZENITH is a powerful technique that: (1) allows to rapidly establish metabolic profile of several populations of cells in an heteregenous sample by flow cytometry, (2) does not require a large amount of cells and thus avoids their expansion in culture and (3) can be combined with scRNA-seq to establish a full map of EM profile of cells. Given the direct relationship between EM and the functionality of lymphoid effector cells and myeloid cells, ZENITH analysis could be used to define the 'immune EM contexture' and complement the establishment of an immunoscore that's defines immune fitness of tumours and predicts and stratifies patients for tailored therapies, aiming at manipulating metabolic pathways to improve anti-tumoral immune effector functions.

Materials and Methods

Cells and Cell Culture

Mouse splenocytes from WT C57BL/6J (Jackson) or PERK KO C57BL/6J background mice (Zhang et al., 2002) were cultured in DMEM containing 5% of Fetal Calf Serum (FCS) and 50 µM of 2-Mercaptoethanol (Mouse cells culture media, MCCM) at 37 ºC 5% of CO2. GM-CSF BM-derived dendritic cells (GM-bmDCs) were differentiated in vitro from the bone marrow of 6–8-week-old male mice, using GM-CSF, produced by J558L cells. Bone marrow progenitors were plated at $0.8.10^6$ cells/ml, 5 ml/well in 6-well plates, and cultivated with RPMI (GIBCO), 5% FCS (Sigma-Aldrich), 20 µg/ml gentamycin (Sigma-Aldrich), 50 µM β-mercaptoethanol (VWR), and GM-CSF. The medium was replaced every 2 days; BM-derived DCs were used for experiments at day 6.

Similarly, FLT3L BM-derived dendritic cells (FLT3L-bmDCs) were differentiated by adding FLT3L to RPMI, 10% of Fetal Calf Serum (FCS) and 50 µM of 2-Mercaptoethanol (Mouse cells culture media, MCCM) during 6 days at 37 ºC 5% of CO2. To obtain splenocytes, eight weeks old wild type C57BL/6J mice were sacrificed by cervical dislocation and splenectomized. Single cells suspentions from the spleens were generated and cultured in MCCM as previously described. Mononuclear cell enriched from blood of healthy donors was submitted to Ficoll-paque plus (PBL Biomedical Laboratories). PBMCs and Whole blood were cultured in the absense (non stimulated) or in the presence of LPS for 4hs. Immune cell stimulations were performed in the absence (Control) or presence of 0,1 µg/ml of extrapure Lipopolysacharide (Invivogen LPS, Cat. tlrl-3pelps), 10 µg/ml Poly I:C (Invivogen, Cat. No. tlrl-pic), CpG-A ODN 2216 (Invivogen, Cat. No. tlrl-2216) or PMA (5 ng/ml; Sigma, Cat. no. P-8139) and ionomycin (500 ng/ml; Sigma, cat. no. I-0634) over night for T cell stimulations and 4 hours for Dendritic cells. T cells from different human donors (P1, P2, P3) were isolated using the RosetteSep™ negative isolation method and activated (using BD Human T cell activator beads coated with anti-CD3 and anti-CD28) or not.

ATP measurement

2.10^4 MEFs were seeded in 100ul of 5% FCS DMEM culture media ON in opaque 96 well plates. Cells were incubated with the inhibitors for the times indicated in the Figure. After, 100ul of Cell titer-Glo luminiscence ATP reconstituted buffer and substrate (Promega, Cat. No. G7570) was added to each well and Luminiscence was measured after 10 minutes following manufacturer instructions. A standard curve with ATP was performed using the same kit and following manufacturer instructions.

Metabolic flux analysis (Seahorse®)

OCR and ECAR were measured with the XF24 Extracellular Flux Analyzer (Seahorse Bioscience). 4.10^5 cells with αCD3/αCD28 beads or not, were placed in triplicates in XF medium (nonbuffered Dulbecco's modified Eagle's medium containing 2.5 mM glucose, 2 mM L-glutamine, and 1 mM sodium pyruvate) and monitored 25 min under basal conditions and in response to 10mM Glucose, 1 µM oligomycin, 100 mM 2-Deoxy-Glucose. Glycolytic capacity was measured by the difference between ECAR level after add oligomycin and before add glucose. OCR, ECAR and SRC parameters was analyzed

and extract from Agilent Seahorse Wave Desktop software. Glycolytic capacity was obtained by the difference between ECAR level after add Oligomycin and before add Glucose.

ZENITH

Cells were plated at 1.10^6 cells/ml, 0,5 ml/well in 48-well plates. Experimental duplicates/triplicates were performed in all conditions. After differentiation, activation or harvesting of human of cells, wells were treated during 30 minutes with Control, 2-Deoxy-D-Glucose (DG, final concentration 100mM; Sigma-Aldrich Cat. No. D6134), Oligomycin (Oligo, final concentration 1μM; Sigma-Aldrich Cat. No.75351), or a combination of the drugs at the final concentrations before mentioned. As negative control, the translation initiation inhibitor Harringtonine was added 15 minutes before addition of Puromycin (Harringtonine, 2 μg/ml; Abcam, cat. ab141941). Puromycin (Puro, final concentration 10 μg/ml; Sigma-Aldrich, Cat. No. P7255) is added during the last 15 minutes of the metabolic inhibitors treatment. After puro treatment, cells were washed in cold PBS and stained with a combination of Fc receptors blockade and fluorescent cell viability marker, then primary conjugated antibodies against different surface markers (see FACS material and methods) during 30 minutes at 4°C in PBS 1X 5% FCS, 2mM EDTA (FACS wash buffer). After washing with FACS wash buffer, cells were fixed and permeabilized using BD Cytofix/Cytoperm™ (Catalog No. 554714) following manufacturer instructions. Intracellular staining of Puro using our fluorescently labeled anti-Puro monoclonal antibody was performed by incubating cells during 1 hour (1:800, Clone R4743L-E8, in house produced and conjugated with Alexa Fluor 647, to obtain 10 times better signal to noise ratio than commercially available monoclonal antibodies) at 4°C diluted in Permwash. For ZENITH troubleshooting see Table 2.

Patients and samples.

The renal carcinoma patient enrolled in this study provided written and informed consent to tissue collection under a University of California, San Francisco (UCSF) institutional review board (IRB)-approved protocol (UCSF Committee on Human Research (CHR) no. 13-12246). The meningioma and brain meastasis patients enroled in this study provided written and informed consent in accordance with institutional, national guidelines and the Declaration of Helsinki. This protocol was approved by institutional review board (AP-HM

CRB-TBM tumor bank: authorization number AC-2018-31053, B-0033-00097).

Processing of and mouse solid tumors ZENITH

0,2-0,4 grams of solid tumor tissue was partially dissociated using chirurgical scissors or tissue chopper (McIlwain Tissue Chopper® Standard plate) to generate "tumor explant suspention". Tissue explants suspention, containing tissue cubes of approximately 400μm of cross section, were put in suspension in complete RPMI media and incubated directly with control or metabolic inhibitors, and with Puromycin following the ZENITH protocol. Next, tumor explants were dissociated using Tissue Liberase and DNAseI with the help of a Gentle Macs (Miltenyi) following manufacturers instructions. Cell suspentions were washed, counted and $2-5.10^6$ total cells were seed in triplicates before proceeding with lived dead and FC block staining. Next, cells were stained for surface makers, fixed and permeabilized (ThermoFisher, FOXP3 fixation kit) and stained for nuclear and cytoplasmic markers as mentioned above.

Human single cell RNA-sequencing

Live CD3-CD19/20-CD56- cells were sorted from renal carcinoma tumor and juxta tumoral tissue using a BD FACSAria Fusion. After sorting, cells were pelleted and resuspended at 1.10^3 cells/μl in 0.04%BSA/PBA and loaded onto the Chromium Controller (10X Genomics). Samples were processed for single-cell encapsulation and cDNA library generation using the Chromium Single Cell 3' v2 Reagent Kits (10X Genomics). The library was subsequently sequenced on an Illumina HiSeq 4000 (Illumina).

Single cell data processing

Sequencing data was processed using 10X Genomics Cell Ranger V1.2 pipeline. The Cell Ranger subroutine mkfastq converted raw, Illumina bcl files to fastqs which were then passed to Cell Ranger's count, which aligned all reads using the aligner STAR (Dobin et al., 2013)ref against GRCh38 genomes for human cells. After filtering reads with redundant unique molecular identifiers (UMI), count generated a final gene-cellular barcode matrix. Both mkfastq and count were run with default parameters.

Cellular Identification and Clustering

For each sample, the gene - barcode matrix was passed to the R (v. 3.6.0) software package

March 10, 2020

Seurat[34] (http://satijalab.org/seurat) (v3.1.1) for all downstream analyses. We then filtered on cells that expressed a minimum of 200 genes and required that all genes be expressed in at least 3 cells. We also removed cells that contained > 5% reads associated with cell cycle genes[35,36]. Count data was then log2 transformed and scaled using each cell's proportion of cell cycle genes as a nuisance factor (implemented in Seurat's ScaleData function) to correct for any remaining cell cycle effect in downstream clustering and differential expression analyses. For each sample, principal component (PC) analysis was performed on a set of highly variable genes defined by Seurat's FindVariableGenes function. Genes associated with the resulting PCs (chosen by visual inspection of scree plots) were then used for graph-based cluster identification and subsequent dimensionality reduction using t-distributed stochastic neighbor embedding (tSNE). Cluster-based marker identification and differential expression were performed using Seurat's FindAllMarkers for all between-cluster comparisons.

Flow cytometry

Flow cytometry was conducted using BD Symphony and BD LSR Fortessa X-20 machine (BD Biosciences™) and data were analyzed with FlowJo (Tree Star™) or FLOWR-software (Guillaume VOISSINE, https://github.com/VoisinneG/flowR). The antibodies used to stain mouse splenocytes were anti-Puro-Clone R4743L-E8, rat IgG2A in house produced and (conjugated with Alexa Fluor 647 or Alexa-Fluor 488), anti-Ki67 PE-eFluor-610 (eBioscience™, Catalog No. 61-5698-82) CD4-APC-eF780 (eBioscience™, Catalog No. 47-0042-82), CD8-APC (eBioscience™, Catalog No. 17-0081-83), CD80-PercPCy5.5 (Biolegend™, Catalog No. 104722), anti-B220-BV421 (Biolegend™, Catalog No. 103251), anti-MHC-II-AF700 (eBioscience™, Catalog No. 56-5321-82), LIVE/DEAD™ Fixable Aqua Dead Cell Stain (Invitrogen™, Catalog No. L34957). The following anti-Human antigens antibodies were used for staining whole blood and PBMCs upon ZENITH protocol application. Alexa Fluor-488 Mouse Anti-Human Axl (Clone 108724, R&D Biosystems, Cat. No. FAB154G), BUV395 Mouse Anti-Human CD11c (Clone B-ly6, BD Bioscience, Cat. No. 563787), BUV737 Mouse Anti-Human CD86 (Clone FUN-1, BD Bioscience, Cat. No. 564428), BV510 Mouse Anti-Human CD19 (Clone HIB19, BD Bioscience, Cat. No. 740164), BV510 Mouse Anti-Human CD3 (Clone HIT3a, BD Bioscience,

Cat. No. 564713), BV510 Mouse Anti-Human CD56 (Clone B159, BD Bioscience, Cat. No. 740171), BV605 Anti-Human HLA-DR (Clone L243, BioLegend, Cat. No. BLE307640), BV650 Mouse Anti-Human CD16 (Clone 3G8, BD Bioscience, Cat. No. 563692), BV711 Mouse Anti-Human CD14 (Clone M5E2, BD bioscience, Cat. No. 740773), BV785 Mouse Anti-Human CD45RA (Clone HI100, BioLegend , Cat. No. BLE304140), Live Dead Fixable Aqua Dead Cell Stain Kit (Life Technologies, Cat. No. L34957), PE Rat Anti-Human Clec9A/CD370 (Clone 3A4, BD Bioscience, Cat. No. 563488), PE-Cy7 Mouse Anti-Human CD22 (Clone HIB22, BD Bioscience, Cat. No. 563941), AF488 Mouse Anti-Human CD38 (Clone HIT-2, BioLegend, Cat. No. BLE303512).

Animal studies

Wild type C57BL/6 mice were purchased from Charles River and maintained in the animal facility of CIML under specific pathogen-free conditions. This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals the French Ministry of Agriculture and of the European Union. Animals were housed in the CIML animal facilities accredited by the French Ministry of Agriculture to perform experiments on alive mice. All animal experiments were approved by Direction Départementale des Services Vétérinaires des Bouches du Rhône (Approval number A13-543). All efforts were made to minimize animal suffering.

Statistical analysis

Statistical analysis was performed with GraphPad Prism software. When several conditions were to compare, we performed a one-way ANOVA, followed by Tukey range test to assess the significance among pairs of conditions. When only two conditions were to test, we performed Student's t-test or Welch t-test, according the validity of homoscedasticity hypothesis (* $P<0.05$, ** $P<0.01$, *** $P<0.005$).

Calculation and meaning of ZENITH derived parameters

To quantify the different energetic metabolism parameters that constitute the metabolic profile of a cell, such as pathways dependency, we used simple algorithms that quantifiy the relative impact of inhibiting a given pathway compared to a complete inhibition of ATP synthesis (Fig. 1f). While ZENITH allow the use of any combination of metabolic or signalling inhibitors, herein we

focused on inhibitors of glycolysis and of mitochondrial respiration to derive metabolic parameters. The percentual of glucose dependence (Gluc. Dep.) quantifies how much the translation levels are dependent on glucose oxidation. Gluc. Dep. is calculated as the difference between PS levels in 2-Deoxy-D-Glucose (DG) treated cells compared to control (Co), divided by the difference in PS upon complete inhibition of ATP synthesis (DG, FCCP and Oligomycin A, combined; treatment Z) compared to control cells (Fig. 1f). In a similar fashion, percentual mitochondrial dependence (Mitoc. Dep) quantifies how much translation is dependent on oxydative phosphorylation. Mitoc. Dep. is defined as the difference in PS levels in Oligomycin A ("O", mitochondrial inhibitor) treated cells compared to control relative to the decreased in PS levels upon full inhibition of ATP synthesis inhibition (treatment Z) also compared to control cells (Fig. 1f). Two additional derived parameters, "Glycolytic capacity" (Glyc. Cap.) and "Fatty acids and amino acids oxidation capacity" (FAAO Cap.) were also calculated. Glycolytic capacity is

defined as the maximum capacity to sustain protein synthesis levels when mitochondrial OXPHOS is inhibited (Fig. 1f, see statistic in Materials and Methods section). Converserly, FAAO Capacity is defined as the capacity to use fatty acids and aminoacids as sources for ATP production in the mitochondria when glucose oxidation is inhibited (Glycolysis and Glucose derived Acetyl-CoA by OXPHOS) (Fig. 1f and Supplementary fig. 1b). While the total level of translation correlates with the global metabolic activity of the cells, the dependency parameters underline essential cellular pathways that cannot be compensated, while "capacity"; as the inverse of dependency, shows the maximun compensatory capacity of a subpopulation of cells to exploit alternative pathway/s when a particular one is inhibited (Fig. 1f and Supplemenary Fig. 1c).

For standard deviation calculation of ZENITH, we followed the propagation of error that is required when the means of means are used into a formula:

For error calculation:

Co= GeoMFI of anti-Puromycin-Fluorochrome upon Control treatment

DG= GeoMFI of anti-Puromycin-Fluorochrome upon 2-Deoxy-D-Glucose treatment

O= GeoMFI of anti-Puromycin-Fluorochrome upon Oligomycin A treatment

Z= GeoMFI of anti-Puromycin-Fluorochrome upon DG+O treatment

$$Glucose\ Dependence\ (Gluc.\ Dep.) = \frac{100\,(Co - DG)}{(Co - Z)}$$

$$SD_{Gluc\ dep} = \sqrt{\left(\frac{\partial}{\partial Co}[Gluc.\ dep.]\right)^2 \times SD_{Co}{}^2} + \sqrt{\left(\frac{\partial}{\partial DG}[Gluc.\ dep.]\right)^2 \times SD_{DG}{}^2} + \sqrt{\left(\frac{\partial}{\partial Z}[Gluc.\ dep.]\right)^2 \times SD_{Z}{}^2}$$

$$SD_{Gluc\ dep} = \sqrt{\left(\left(\frac{-100\,(Z-DG)}{(Co-Z)^2}\right)^2 \times SD_{Co}{}^2\right)} + \sqrt{\left(\left(\frac{100}{Z-Co}\right)^2 \times SD_{DG}{}^2\right)} + \sqrt{\left(\left(\frac{100\,(Co-DG)}{(Co-Z)^2}\right)^2 \times SD_{Z}{}^2\right)}$$

$$Mitochondrial\ dependence\ (Mitoc.\ Dep.) = \frac{100\,(Co - O)}{(Co - Z)}$$

$$SD_{Mito\ dep} = \sqrt{\left(\left(\frac{-100\,(Z-DG)}{(Co-Z)^2}\right)^2 \times SD_{Co}{}^2\right)} + \sqrt{\left(\left(\frac{100}{Z-O}\right)^2 \times SD_{O}{}^2\right)} + \sqrt{\left(\left(\frac{100\,(Co-DG)}{(Co-Z)^2}\right)^2 \times SD_{Z}{}^2\right)}$$

$$Glycolytic\ capacity\ (Glyco.\ Cap.) = 100 - \left(\frac{100\,(Co - O)}{(Co - Z)}\right)$$

$$SD_{Glyc.\ cap.} = \sqrt{\left(\left(\frac{100\,(Z-DG)}{(Co-Z)^2}\right)^2 \times SD_{Co}{}^2\right)} + \sqrt{\left(\left(\frac{100}{Co-Z}\right)^2 \times SD_{O}{}^2\right)} + \sqrt{\left(\left(\frac{100\,(DG-Co)}{(Z-Co)^2}\right)^2 \times SD_{Z}{}^2\right)}$$

$$FAO\ and\ AAO\ capacity\ (FAAO\ Cap.) = 100 - \left(\frac{100\,(A - DG)}{(A - Z)}\right)$$

$$SD_{FAO\ and\ AAO\ cap.} = \sqrt{\left(\left(\frac{100\,(Z-DG)}{(Co-Z)^2}\right)^2 \times SD_{Co}{}^2\right)} + \sqrt{\left(\left(\frac{100}{Co-Z}\right)^2 \times SD_{O}{}^2\right)} + \sqrt{\left(\left(\frac{100\,(DG-Co)}{(Z-Co)^2}\right)^2 \times SD_{Z}{}^2\right)}$$

March 10, 2020

Acknowledgments

The PP laboratory is "Equipe de la Fondation de la Recherche Médicale" (FRM) sponsored by the grant DEQ20140329536. The project was supported by grants from l'Agence Nationale de la Recherche (ANR), « ANR-FCT 12-ISV3-0002-01», A*MIDEX project "CSI" (ANR-11-IDEX-0001-02), «DCBIOL Labex ANR-11-LABEX-0043 », « INFORM Labex ANR-11-LABEX-0054 » funded by the "Investissements d'Avenir" French government program and initiated through Contrats Proof of concept CoPoc 2019 INSERM "ZIPT R19011AS" to RJA, Bourse Mobilité Cancéropôle (to UCSF) to RJA and Cancéropôle PACA "Emergence" grant "HiDi-Glio" to RJA and Projet Fondation ARC 2016 to PP. The research was also supported by the Ilídio Pinho foundation and FCT - Fundação para a Ciência e a Tecnologia - and Programa Operacional Competitividade e Internacionalização - Compete2020 (FEDER) – references POCI-01-0145-FEDER-016768 and POCI-01-0145-FEDER-030882. We acknowledge financial support from n° ANR-10-INBS-04-01 France Bio Imaging and the ImagImm CIML imaging. We thank Noella LOPES-PAPPALARDO and Alexandre BOISSONNAS for reading the manuscript and suggestions and Lionel SPINELLI for usefull statistical advices. We thank for the help of Marc BARAD and Sylvain BIGOT from the CIML Cytometry core facility and Alice CARRIER and Laurence BORGE from the CRCM metabolomics core facility. I thank Pierre GOLSTEIN for mentoring and correcting the manuscript. The authors declare to have no competing interest.

Authors' contribution:

Conceptualization, R.J.A.
Method development, R.J.A., A.C.
Validation, R.J.A., A.C, R.C, E.B., P.C., V.C., J.P.G.
Formal Analysis, R.J.A., A.C, B.S.
Investigation, R.J.A., A.C, M.F.K. and P.P.
Tissue Resources, E.T., D.F., S.B., M.F.K and P.P.
Writing – Original Draft, R.J.A.
Writing – R.J.A., A.C, E.G., M.F.K., and P.P.
Review & Editing, R.J.A., A.C, E.G, R.C., M.F.K. and P.P.
Visualization, R.J.A., A.C, E.G, M.F.K and P.P.
Supervision, R.J.A., M.F.K and P.P.
Funding Acquisition, R.J.A., M.F.K and P.P.

References

1. O'Sullivan, D., Sanin, D. E., Pearce, E. J. & Pearce, E. L. Metabolic interventions in the immune response to cancer. *Nat. Rev. Immunol.* **19**, 324–335 (2019).
2. Russell, D. G., Huang, L. & VanderVen, B. C. Immunometabolism at the interface between macrophages and pathogens. *Nature Reviews Immunology* **19**, 291–304 (2019).
3. Roos, D. & Loos, J. A. Changes in the carbohydrate metabolism of mitogenically stimulated human peripheral lymphocytes. II. Relative importance of glycolysis and oxidative phosphorylation on phytohaemagglutinin stimulation. *Exp. Cell Res.* **77**, 127–135 (1973).
4. Pearce, E. L. *et al.* Enhancing CD8 T-cell memory by modulating fatty acid metabolism. *Nature* **460**, 103–7 (2009).
5. Chang, C. H. *et al.* Metabolic Competition in the Tumor Microenvironment Is a Driver of Cancer Progression. *Cell* **162**, 1229–1241 (2015).
6. Galon, J. *et al.* Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science (80-.).* **313**, 1960–1964 (2006).
7. Antonia, S. J. *et al.* Overall survival with durvalumab after chemoradiotherapy in stage III NSCLC. *N. Engl. J. Med.* **379**, 2342–2350 (2018).
8. Wolchok, J. D. *et al.* Overall Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N. Engl. J. Med.* **377**, 1345–1356 (2017).

March 10, 2020

9. Miller, A. *et al.* Exploring Metabolic Configurations of Single Cells within Complex Tissue Microenvironments. *Cell Metab.* **26**, 788-800.e6 (2017).

10. Palmer, A. *et al.* FDR-controlled metabolite annotation for high-resolution imaging mass spectrometry. *Nat. Methods* **14**, 57–60 (2016).

11. Levine, L. S. *et al.* Single-cell metabolic analysis by mass cytometry reveals distinct transitional states of CD8 T cell differentiation. *bioRxiv* (2020). doi:10.1101/2020.01.21.911545

12. Ahl, P. J. *et al.* A novel strategy for single-cell metabolic analysis highlights dynamic changes in immune subpopulations. *bioRxiv* 2020.01.21.914663 (2020). doi:10.1101/2020.01.21.914663

13. Hartmann, F. J. *et al.* Multiplexed Single-cell Metabolic Profiles Organize the Spectrum of Cytotoxic Human T Cells. *bioRxiv* 2020.01.17.909796 (2020). doi:10.1101/2020.01.17.909796

14. Zhang, J. *et al.* Measuring energy metabolism in cultured cells, including human pluripotent stem cells and differentiated cells. *Nat. Protoc.* **7**, 1068–85 (2012).

15. Llufrio, E. M., Wang, L., Naser, F. J. & Patti, G. J. Sorting cells alters their redox state and cellular metabolome. *Redox Biol.* **16**, 381–387 (2018).

16. Neijssel, O. M., Teixeira De Mattos, M. J. & Tempest, D. W. Growth yield and energy distribution. (1996).

17. Buttgereit, F. & Brand, M. D. A hierarchy of ATP-consuming processes in mammalian cells. *Biochem. J.* **312**, 163–167 (1995).

18. Lindqvist, L. M., Tandoc, K., Topisirovic, I. & Furic, L. Cross-talk between protein synthesis, energy metabolism and autophagy in cancer. *Curr. Opin. Genet. Dev.* **48**, 104–111 (2018).

19. Schimmel, P. GTP hydrolysis in protein synthesis: two for Tu? *Science (80-.).* **259**, 1264–1265 (1993).

20. Miyamoto-Sato, E., Nemoto, N., Kobayashi, K. & Yanagawa, H. Specific bonding of puromycin to full-length protein at the C-terminus. *Nucleic Acids Res.* **28**, 1176–82 (2000).

21. Nemoto, N., Miyamoto-Sato, E. & Yanagawa, H. Fluorescence labeling of the C-terminus of proteins with a puromycin analogue in cell-free translation systems. *FEBS Lett.* **462**, 43–6 (1999).

22. Wool, I. G. & Kurihara, K. Determination of the number of active muscle ribosomes: effect of diabetes and insulin. *Proc. Natl. Acad. Sci. U. S. A.* **58**, 2401–2407 (1967).

23. Andrews, T. M. & Tata, J. R. Protein synthesis by membrane-bound and free ribosomes of secretory and non-secretory tissues. *Biochem. J.* **121**, 683–694 (1971).

24. Schmidt, E. K., Clavarino, G., Ceppi, M. & Pierre, P. SUnSET, a nonradioactive method to monitor protein synthesis. *Nat. Methods* **6**, 275–277 (2009).

25. Argüello, R. J. *et al.* SunRiSE - measuring translation elongation at single-cell resolution by means of flow cytometry. *J. Cell Sci.* **131**, jcs214346 (2018).

26. Van Der Windt, G. J. W. *et al.* CD8 memory T cells have a bioenergetic advantage that underlies their rapid recall ability. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 14336–14341 (2013).

27. van der Windt, G. J. W. *et al.* Mitochondrial Respiratory Capacity Is a Critical Regulator of CD8 + T Cell Memory Development. *Immunity* **36**, 68–78 (2012).

28. Saha, S., Shalova, I. N. & Biswas, S. K. Metabolic regulation of macrophage phenotype and function. *Immunol. Rev.* **280**, 102–111 (2017).

29. Everts, B. *et al.* Commitment to glycolysis sustains survival of NO-producing inflammatory dendritic cells. *Blood* **120**, 1422–1431 (2012).

30. Krawczyk, C. M. *et al.* Toll-like receptor-induced changes in glycolytic metabolism regulate dendritic cell activation. *Blood* **115**, 4742–4749 (2010).

31. Amiel, E. *et al.* Inhibition of Mechanistic Target of Rapamycin Promotes Dendritic Cell Activation and Enhances Therapeutic Autologous Vaccination in Mice. *J. Immunol.* **189**, 2151–2158 (2012).

32. Vitale, I., Manic, G., Coussens, L. M., Kroemer, G. & Galluzzi, L. Macrophages and Metabolism in the Tumor Microenvironment. *Cell Metabolism* **30**, 36–50 (2019).

33. Villani, A.-C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science* **356**, eaah4573 (2017).

34. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nat. Biotechnol.* **33**, 495–502 (2015).

35. Kowalczyk, M. S. *et al.* Single-cell RNA-seq reveals changes in cell cycle and differentiation programs upon aging of

March 10, 2020

hematopoietic stem cells. *Genome Res.* **25**, 1860–1872 (2015).

36.   Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).

March 10, 2020

a

b

Supplementary figure 1.

a, Impact of simultaneous block of ATP synthesis and translation (CHX) or transcription (Actinomycin D, ActD) or both translation and transcription in the levels of ATP (n=3), showing that blocking translation induce an increase in ATP levels.

b, Schematic representation of a sample that contains three cell types (each of different color) with three different metabolism profiles (Aerobic glycolysis, Glycolysis/OXPHOS, and FAAO). Treating the mix of cells with DG, or O, will impact each cell subset in a different way.

c, Example of ZENITH metabolic profile (left) and five different metabolic parameters/measurements that can be extracted from the graph (right) using DG and O as inhibitors.

Supplementary figure 2.
Seahorse and ZENITH metabolic parameters in non-activated and activated purified human T cells.
a, Seahorse Maximal oxygen consumption rate (FCCP) in non-activated or T cells stimulated over night with CD3/CD28 activator beads (n=3).
b, Seahorse Spare Respiratory Capacity (calculated using the seahorse software) in non-activated or T cells stimulated over night with CD3/CD28 activator beads (n=3).

Supplementary figure 3. Changes in ZENITH EM profile in mouse and human T cells upon activation.
a, Mouse splenic T cells where analyzed using ZENITH. Total Puro MFI is indicated in CD3+CD8+CD4-non-activated, left or PMA/Ionomycin-treated right.
b, EM profiles calculated from the values shown in a).
c, EM profile of non-stimulated or stimulated human blood central memory CD4 T cells (CD3+CD4+CD45RO+CCR7+) from two different subjects (P4 and P5) using the ZENITH method. N=3, ANOVA on at least 2 independent experiments (* p<0.05; ** p<0.005; ***p<0.0005).

Supplementary figure 4.

a, Profile of protein synthesis levels of human blood mononuclear cells treated (4hs) or not with LPS (100ng/ml) and subjected to ZENITH.

b, Mouse Bone marrow FLT3-L derived dendritic cells at day 7 were activated or not with LPS (100ng/ml) and analyzed with ZENITH (A= Control; B=2DG, C=Oligomycin, Z=2DG+Oligomycin) by FACS using a panel to identify DC1 (CD11c+MHC-II+SIRPa-CD24+SiglecH-), DC2 (CD11c+MHC-II+SIRPa+XCR1-CD24-SiglecH-) and pDC (CD11c+MHC-II+SIRPa-CD24-SiglecH+); the Mean fluorescence of anti-puromycin in each DC subset from each mice (1, 2, 3) in triplicates is shown (ANOVA in at least two independent experiments; * p<0.05; ** p<0.005; ***p<0.0005)

c, Mouse Bone marrow FLT3-L derived dendritic cells at day 7 were activated or not with LPS (100ng/ml) and analyzed with ZENITH by FACS using a panel to identify DC1 (CD11c+MHC-II+SIRPa-CD24+SiglecH-), DC2 (CD11c+MHC-II+SIRPa+XCR1-CD24-SiglecH-) and pDC (CD11c+MHC-II+SIRPa-CD24-SiglecH+) (N=3 different mice, ANOVA in at least two independent experiments ; * p<0.05; ** p<0.005; ***p<0.0005) **d**, Clustering of metabolic profile of dendritic cells subsets from three independent mice (rows) treated or not with LPS. **e**, Splenocytes were treated for 4 hours with or without different TLR ligands (i.e. LPS, 100ng/ml; Poly (I:C) 10ug/ml) or PMA/ionomycin (10 ng/ml PMA; 2.5 μM Ionomycin). Protein synthesis levels in DC2 (left) or DC1 (right); show different metabolic profile of splenic dendritic cells (N=3 independent spleens) (* p<0.05; ** p<0.005; ****p<0.0001).

Supplementary figure 5 (part 1 of 2)

Composition, phenotype and metabolic profile of multiple immune cells populations can be analyzed in parallel in human tumors, ex-vivo both by ZENITH and Single cell-RNA sequencing

a, t-SNE graph showing the different cell clusters analyzed by ZENITH in the meningioma, the percentage (in each gate) and the phenotype of each subpopulation (each subpop in one color).

b, Same as a, but for Myeloid cell from the renal carcinoma patient.

c, The metabolic profile of each subset from each tissue was used to cluster cells. Two clusters were observed with different metabolic profile, and were name accordingly as Glycolytic cluster and Respiratory cluster of cells.

d, Heatmap of metabolic gene expression and ZENITH functional metabolic profile in myeloid cells from PBMC. * The identiy of the genes is not revealed because the gene signature is patent pending.

e, Far left, t-SNE display and graph-based clustering of CD3e-,CD19-,CD20-,CD56-, HLA-DR+/- myeloid cells sorted from Renal carcinoma (top) or Juxta tumoral tissue (bottom) biopsies and processed for scRNA-seq. Each dot represents a single cell. From left to right, classical monocytes, non classical monocytes, type 1 dendritic cell, type 2 dendritic cell and plasmacytoid dendritic cell specific gene signature identified in PBMC from heathly donors (Villani et al. 2017). distributed across t-SNE plot of myeloid cells sorted from renal carcinoma (top) or juxta-tumoral tissue (bottom).

f, Left, t-SNE display and graph-based clustering of CD3e-,CD19-,CD20-,CD56-, HLA-DR+/- monocytes and macrophages sorted from Renal carcinoma (top) or Juxta tumoral tissue (bottom) biopsies and processed for scRNA-seq. Each dot represents a single cell. CSF1R (middle) and MAFB (right) gene expression distributed across t-SNE plot of monocytes and macrophages sorted from renal carcinoma (top) or juxta-tumoral tissue (bottom).

g,CD14 (left) and FCGR3A (right) gene expression distributed across t-SNE plot of monocytes and macrophages sorted from renal carcinoma (top) or juxta-tumoral tissue (bottom).

h, Heatmap displaying top 5 DE genes for each cluster of monocytes and macrophages present in renal carcinoma (left) or in juxta-tumoral tissue (right) when comparing cluster 0 through 4 in renal carcinoma (ranked by log fold change).

i, Glycolytic (left) and Respiratory (right) specific gene signature expression distributed across t-SNE plot of myeloid cells sorted from renal carcinoma (top) or juxta-tumoral tissue (bottom) with DC clusters circled.

Table 2. ZENITH Troubleshooting

Problem	Potential cause	Suggestion
Signal for translation (Puro MFI) is too low or lower signal in control treatment	Cells were not kept at 37°C during treatments.	Prepare al reagents and equilibrate the temperature. Neither put cells on ice, nor keep them at room temperature.
	Puromycin stock was not correctly stored.	Avoid exposure to light and freeze thawing cycles. Use small aliquots of puromycin and discard after each use.
	An inappropriate concentration of puromycin was used	Puromycin concentration can be cell dependent and should be determined experimentally for each sample being tested. Too high, or too low concentrations result in low staining.
	The incubation time with puromycin was insufficient	Keep Puromycin during at least 30 min
	Pathogen infection	Obtain fresh culture with fresh medium to prevent contamination. 1% Penicillin (not streptomycin) can be used
	Puromycin resistance in your cells	Change antibiotic resistance into the construct to be compatible with ZENITH.
	Cell culture medium was depleted of nutrients.	Change media 2 ~ 8 hr before the experiment.
	Too many cells during staining	Do not stain in each tube more than 5 million cells. Adapt the volume of antibody.
Signal for translation (Puro MFI) is too high	Commercially available clones can show substantial background.	Order or ask for the ZENITH kit containing the R4743L-E8 mAb clone.
Signal for translation (Puro MFI) is higher in my cells treated with inhibitor than the control	Some metabolic inhibitors make cells to switch to other sources to compensate. The activation of this compensatory mechanism might result in higher levels of translation than the control.	If cells show increased translation upon treatment when calculating the metabolic dependency, the formula would result in a negative dependency value. Conceptually a negative dependency is equal to no dependency (0).
		As the inhibitor induce an increase in the translation levels and cells are only relying on the compensatory pathway, then the formula will reveal a percentual metabolic capacity that is over 100%. Meaning that cells have the metabolic capacity to sustain even higher levels of translation than in non-treated cells.
Signal of cell surface markers decrease after treatment with inhibitors	An inappropriate time of treatment with inhibitors was performed	Time of treatment with inhibitors depends on the metabolic activity of the cells and have to be determined experimentally. Reduce the time of Z treatment (i.e. 20min).
		Perform intracellular staining to detect the internalized markers.
Signal is present in Z (all inhibitors)	Oligomycin acts faster than DG (competitive inhibitor of glucose).	Treat cells 10-15´min with 2DG and then add the Oligomycin 20-15´, use that measure as Z treatment.
	High background from cells	Use Fc block, Fcs 2% or BSA 1-3% during staining can reduce background
Signal doesn't change with any kind of treatment	Inhibitor doesn't work	Use inhibitors at 37°C to have powerful effect
	The concentration and/or time of treatment are not optimized	Inhibitors concentration can be cell dependent and should be determined experimentally for each cell line being tested.
Lot of doublet cells during Facs acquisition	Cells were aggregated	Obtain a single cell suspension by using 2mM EDTA in facs buffer