Title: The Demographic History of Micro-endemics: Have Rare Species Always Been Rare?

Authors: Andrew J. Helmstetter[1,2], Stuart Cable[1,3], Franck Rakotonasolo[3], Romer Rabarijaona[3], Mijoro Rakotoarinivo[4], Wolf L. Eiserhardt[1,5], William J. Baker[1], Alexander S.T. Papadopulos[1,6]

Affiliations: [1]Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK

[2]Institut de Recherche pour le Développement (IRD), UMR-DIADE, 911 Avenue Agropolis, BP 64501, 34394 Montpellier, France.

[3]Kew Madagascar Conservation Centre, Lot II J 131 B Ambodivoanjo, Ivandry, Antananarivo 101, Madagascar

[4]Mention Biologie et Ecologie Végétales, Faculté des Sciences, Université d'Antananarivo, BP 906101, Antananarivo, Madagascar

[5]Department of Biology, Aarhus University, Aarhus, Denmark

[6]Molecular Ecology and Fisheries Genetics Laboratory, Environment Centre Wales, School of Natural Sciences, Bangor University, Bangor, LL57 2UW, UK

Corresponding Authors: Alexander S.T. Papadopulos (a.papadopulos@bangor.ac.uk) and Andrew J. Helmstetter (andrew.j.helmstetter@gmail.com)

Keywords: Conservation genomics; Demographic inference; Effective population size; Bayesian skyline;

Summary

Current IUCN conservation assessment methods are reliant on observed declines in species population and range sizes over the last one hundred years, but for the majority of species this information is not available. We used a population genetic appraoch to reveal historical demographic decline in the rare endemic flora of Madagascar. These results show that it is possible to predict extinction risk from demographic patterns inferred from genetic data and that destructive human influence is likely to have resulted in the very high frequency of narrow endemics present on the island. Our approach will act as an important tool for rapidly assessing the threatened status of poorly known species in need of further study and conservation, particularly for tropical flora and fauna.

Abstract

Extinction has increased as human activities impact ecosystems. Conservation assessments for the IUCN red list are a fundamental tool in aiding the prevention of further extinction, yet, relatively few species have been thoroughly assessed. To increase the efficiency of assessments, novel approaches are needed to highlight threatened species that are currently data deficient. Many Madagascan plant species currently have extremely narrow ranges, but this may not have always been the case. To assess this, we used high-throughput DNA sequencing for 2-5 individuals of each species - reflecting the paucity of samples available for rare species. We estimated effective population size (N_e) for each species and compared this to census population (N_c) sizes when known. In each case, N_e was an order of magnitude larger than N_c – a signature of rapid, recent population decline. We then estimated the demographic history of each species, tracking changes in N_e over time. Five out of ten species displayed significant population declines towards the present (68–90% decreases). Our results for palm trees indicate that it is possible to predict extinction risk, particularly in the most threatened species. Similar declines to those in palms were observed in data deficient species or those assessed as of least concern. These analyses reveal that Madagascar's narrow endemics were not always rare, having experienced rapid decline in their recent history. Our approach offers the opportunity to target species in need of conservation assessment with little prior information, particularly in regions where human modification of the environment has been rapid.

Introduction

The existence of a mass extinction during modern times has become well accepted (Ceballos *et al.* 2017). According to the International Union for Conservation of Nature (IUCN) red list, approximately 1.2% of assessed species are extinct in the wild and 29.7% are threatened by extinction (Pimm *et al.* 2015). With fewer than 100k species assessed, we have poor knowledge of the extinction risk in many taxa, including those that are vital to human wellbeing. For example, only 27k plant species have IUCN assessments (~6% of all plants), but the loss of plant diversity is expected to have a greater effect on humans than the loss of diversity in any other group (Schatz 2009). Therefore, it is critical that we assess and continue to monitor plant species. Despite the rigour with which assessments are made, it is possible that current methods substantially underestimate the extent to which species may be at risk of extinction. For example, population decline within the last 100 years is a fundamental component of how IUCN red listing is performed (Maes *et al.* 2015; Collen *et al.* 2016), but recent estimates suggest that 30% of all vertebrates species are in decline and as much as 55% of declining bird species are not classified as threatened (Ceballos *et al.* 2017). Furthermore, substantial population decline in many species may have pre-dated historical records or may not have been observed directly at all. As a result, it is unclear whether species that are rare today have experienced population decline or simply have a low carrying capacity (e.g. after becoming highly specialised on spatially-restricted habitats). Without taking into account unobserved declines in population sizes we may be drastically underestimating the number of threatened species and the severity of the extinction threat they face. The unique and highly endemic flora of Madagascar provide an opportunity to investigate the link between historical demographic change and extinction risk.

Levels of endemism on Madagascar are some of the highest worldwide; all of the island's amphibians and terrestrial mammals, 83% of vascular plant species and 86% of macroinvertebrates are found nowhere else (Allnutt *et al.* 2008; Vences *et al.* 2009; Goyder *et al.* 2017). A striking feature of the Madagascan flora and fauna is the abundance of micro-endemic species that have small ranges and are relatively rare (Yoder *et al.* 2000; Wilme *et al.* 2006; Vences *et al.* 2009). For example, more than half of the island's endemic palms are known to be extremely rare – many are found in just a single population or with fewer than 100 individuals in the wild (Rakotoarinivo *et al.* 2014). The processes that gave rise to the incredible diversity and micro-endemism of Madagascar are hotly debated (Wilme *et al.* 2006; Pearson & Raxworthy 2009; Vences *et al.* 2009; Eberhart-Phillips *et al.* 2015; Federman *et al.* 2015; Yoder *et al.* 2016; Hackel *et al.* 2018), but proposed models put little emphasis on the role human beings have played in shaping the distribution of species. Humans arrived on Madagascar recently, around 2500 years ago, and preceded the rapid extinction of the island's megafauna (Crowley 2010). Although the exact causes of megafaunal extinction are less clear, increasing human activities (including hunting, slash and burn agriculture, illegal logging and introduction of invasive species) are likely to have played a major role (Godfrey *et al.* 2019). Extinction of large lemuriforms is also thought

to have had a negative impact on dispersal and fitness of large-seeded plants (Federman *et al.* 2016). Deforestation has also advanced rapidly so that only 10% of the island's forest remains (Harper *et al.* 2019). This combination of recent human influence and abundant micro-endemism invites the question: have Madagascar's micro-endemics always been rare, or is this pattern driven by exceptional population decline across the island's flora and fauna? In this study, our aims were to (i) investigate whether Madagascan plants have suffered population declines in the recent past and (ii) determine whether inference of past demographic changes from genetic data can predict extinction risk.

Results and Discussion

We investigated population decline in 10 Madagascan plant species using genome-wide genetic markers. Our sample included a range of endemic species from the humid and littoral forests of eastern Madagascar (Fig. 1). These forests have decreased by up to a third since the 1970s and are severely under threat (Moat & Smith 2007). The diverse palm (Arecaceae) flora of Madagascar is reasonably well studied and more than 90% of the 204 Madagascan palms have been placed in one of the threatened categories following red list assessment – an indication that these all have declining or small populations. For this reason, we included four palm species (all of which have decreasing populations) as our training data set. Two of these have been assessed as near threatened - *Ravenea robustior* (Rakotoarinivo & Dransfield 2012b) is estimated to have <1000 mature individuals in the wild, while *Dypsis procumbens* is more common but restricted to 32 locations and is exploited for timber (Rakotoarinivo & Dransfield 2012a). *Dypsis rabepierrei* is critically endangered with no more than 20 individuals known from a single location (Eiserhardt *et al.* 2018) and *Satranala decussilvae* is endangered with fewer than 200 mature trees in the wild (Rakotoarinivo & Dransfield 2012c).

The effective population size (N_e) of a species is normally substantially smaller than the census population size, such as in humans where N_e is in the order of 10,000 (Gazave *et al.* 2013). When populations are experiencing rapid decline it is possible for N_e to substantially exceed census population size, as has been observed in the critically endangered black rhinoceros (Braude & Templeton 2009). If the studied palm species have always been rare, we expected that N_e would be equal or less than the census population sizes. Conversely, if rare species were more abundant prior to human influences in Madagascar N_e would be substantially higher than contemporary census sizes. We performed double digest restriction site associated DNA (ddRAD) sequencing (Peterson *et al.* 2012) on 2-5 individuals of each species and genotyped these at 24-78k loci using the Stacks (Catchen *et al.* 2013) pipeline (see table S1 and Supplementary materials and methods for details). We then estimated Θ, the population scaled mutation rate ($\Theta = 4N_e\mu$), using the Bayesian method implemented in the R package *thetamateR*, which allows for rate variation between loci. The estimated Θ value for each species was converted to N_e assuming the average mutation rate in angiosperms (De La Torre *et al.* 2017) and a generation time

of 10 years. In each the four palm species, estimated N_e substantially exceeds the current census population sizes (Fig. 2, table S1), a strong indication that these rare species have not always been rare.

To provide further evidence that these palm populations have experienced declines, we used a subset of loci to investigate the demographic history through time using the Extended Bayesian Skyline Plot approach (EBSP; Heled & Drummond 2008) implemented in BEAST v2.4.2 (Bouckaert *et al.* 2014) following the methodology outlined by Trucchi et al. (Trucchi *et al.* 2014). These methods struggle to estimate N_e in the very recent past due to a paucity of coalescent events and maintenance of variation in diminished populations, so declines very close the present may not be inferred. Given that our N_e estimates and IUCN assessment suggest our four palm species have rapidly decreasing populations, we expected that species population sizes over time (estimated using EBSP) would decrease sharply towards the present. The endangered palm species have very small census sizes, so we expected that any decline would be more pronounced in these species. If so, this approach could provide a useful indicator of extinction risk, the efficacy of which can be demonstrated using our palm dataset.

For each species, we assessed whether we could reject a hypothesis of constant size through time by assessing the "sum(indicators.alltrees)" statistic, which is analogous to the number of population size change events. When 95% highest posterior density (HPD) values do not include zero this indicates that N_e has not been constant through time. The results of the EBSP analysis for palms (Fig. 2) do not contain evidence of a recent decline in the near threatened species, each with a single population size change event, corresponding to early population expansion (Table S1). It is possible that these species at less risk are more abundant than thought. However, given the large difference between N_e and census sizes (Table S1), this could also indicate a very rapid a steep decline in recent history that could not be detected with EBSP. On the other hand, the endangered species have experienced very steep population declines towards the present, as much as a 90% reduction in population size, and multiple demographic events in their history. These results are expected for an endangered species, so we suggest that the approach used here is a useful tool for predicting more extreme levels of extinction risk.

To validate whether our approach has the power to detect anthropogenic population decline, we simulated sequence datasets using *fastsimcoal2* (Excoffier et al. 2013) with ancestral populations sizes set to *thetamateR* Ne estimates and a range of population declines of varying severity that took place 2,500 years ago. We then ran our pipeline, sampling loci for the simulated data in the same way as from the empirical data for each of the palm species (i.e., the same number of loci with n SNPs per locus). We found that we were able to detect significant population decline in each species except *R. robustior*, likely due to the low number of individuals (n = 2). In those species where empirical data suggested a decline, we found that all simulated runs also estimated a decline (Fig. 2). The severity of empirical decline matched the simulated severity, though in *S. decussilvae* there was a significant drop even when no decline was simulated. This is likely to be a consequence of using the same sampling of loci between

empirical and simulated data. The types and frequency of loci with different numbers of SNPs that are available for sampling in declining species are a consequence of population decline. As a result, when the same sampling regime is applied to a simulated dataset with no decline, the resulting skyline plot also captures the signal of a population reduction. These results show that our approach has the power to detect population decline in the Anthropocene given enough individuals, but the capacity to estimate the magnitude of the decline may be more limited.

To investigate the prevalence of these patterns in less well-studied Madagascan species, we repeated the above analyses for the remaining six species in our dataset – only one of which has known census population sizes. Among this group, just three species have been red list assessed and are categorised as of least concern: *Psiadia altisimma* and *Dichaetanthera cordifolia* have stable populations (*Psiadia altisimma* is a slash and burn successor), while *Dilobeia thouarsii* is decreasing due to agriculture, harvesting for timber and traditional medicine (Lemmens *et al.* 2012; Letsara 2018). *Chysophyllum boivinianum* is also exploited for timber and malarial medicine, and is suspected to be declining rapidly (Lemmens 2007). Little is known about the extent of exploitation and biology of the other two species in our sample; *Adenia perrieri* and an undescribed species of *Psychotria*. In these six species, N_e ranged from 7,253 in *Psychotria sp.* to 25,702 in *D. cordifolia*. Although census sizes are not available for these populations, it seems unlikely that any of these species would have always been rare, indeed *P. altisimma* is relatively common (Danthu *et al.* 2008). However, the EBSP analyses are more revealing. Two species showed no evidence of recent decline; *A. perrieri* and, as expected, *D. cordifolia*. *Psiadia altissima* showed some evidence of decline (↓43%) but we could not reject constant size for this species. The remaining three species showed clear decreases in population size towards the present, which is consistent with the suspected exploitation of *C. boivinianum* (↓83%) and *D. thouarsii* (↓82%). Madagascar is a centre on endemism for *Psychotria* (↓68%), which are often small understory trees and shrubs with limited ranges so these species may be particularly susceptible to deforestation. Given that demographic decline in the EBSP was only detected in the most severe cases in the palms, the decline observed in these species suggests that they may be at more severe risk of extinction than current knowledge suggests. It is important to note, however, that coalescent estimation of population sizes calculated in this way can only determine demographic patterns for the population that the sampled individuals represent, which may not reflect the species as a whole. As the majority of species were collected in the fragmented forests of Andasibe (Fig. 1), it is possible that these declines are only pertinent to the Andasibe populations. This region experienced very rapid deforestation that has recently been limited by community-based conservation efforts (Dolch *et al.* 2015) and our results highlight the high levels of population decline that have happened even in currently well-managed areas.

Our ddRADseq approach is a relatively cheap and accessible way to assess past demographic changes with a handful of specimens rapidly. It holds great potential to screen species at a much higher rate than is possible with observation-based red listing assessments. A single protocol was successful for species

from seven different plant families, highlighting the universality of our approach. Demographic trends can be inferred using just a few individuals with large numbers of markers (Heled & Drummond 2008), where previous approaches with small numbers of markers (e.g., microsatellites or chloroplast DNA) fall short. This makes our approach ideal for studying rare and difficult to access species, many of which may be the most threatened and in need of assessment. Effective population sizes estimated using coalescent methods are not proxies for the variance effective size often used in conservation efforts (Lauterbur 2019), but changes in N_e are informative regarding recent demographic changes, as is shown here. Further work is required to establish the nature of the relationship between the population declines observed in EBSP analyses and the extinction risk of these species, and to determine what role these types of analyses can play in red listing efforts. Our results suggest that rapid decline has affected 70% of the species we assessed and it seems likely that many of the micro-endemic species in Madagascar have not always been so rare. Instead, we propose that the radical environmental changes that have taken place in Madagascar since the arrival of humans have driven rapid population decline in Madagascan plants.

Materials and Methods

Sample Collection

43 individuals across 10 species and seven families were sampled (Table S1). Leaf tissue was collected from wild individuals and desiccated using silica gel to preserve DNA. Approximately 20mg of dried tissue was used for DNA extractions. The majority of samples were collected in Andasibe, Madagascar and palm species were collected across the island (Fig. 1). Location data are detailed in Table S1.

Library preparation

Genomic DNA was extracted using a hexadecyltrimethylammonium bromide (CTAB) mini-extraction protocol (Doyle & Doyle 1987), purified using spin columns from the Qiagen DNeasy Plant Mini Kit and then eluted in 60µl water. Double-digest Restriction-site Associated DNA (ddRAD) libraries were constructed following the protocol of Peterson et al. (Peterson *et al.* 2012). Briefly, 1 µg of high molecular weight DNA was digested using two rare-cutting enzymes; EcoRI-HF (NEB) and SphI-HF (NEB). Barcoded flex adapters (Peterson *et al.* 2012) were ligated to 400 ng digested DNA for multiplexing and samples were pooled. Size selection was carried out with a Pippin Prep (Sage Biosciences) with a tight window of 468 to 546bp. Eight PCR reactions per library were conducted to minimize the effect of PCR bias.

Sequencing and genotyping

Libraries were sequenced on three Illumina MiSeq runs, using 2x75bpb (V3), 2x250bp (V2) and 2x300bp (V3) kits respectively. Read pairs were concatenated to produce a single sequence of either 147bp or 380bp after barcode removal and trimming. Reads were demultiplexed using *process_radtags*

in the software Stacks (v1.40) (Catchen *et al.* 2013). Reads were aligned and SNPs detected using *ustacks*. A separate catalogue was built for each species (*cstacks*), to which loci were matched from each individual of a species (*sstacks*). *Populations* was used to filter and output data by only retaining those loci that were present in all individuals within a species.

Inference of N_e and demographic history

For each species, the fasta format output of from the STACKS pipeline containing all aligned loci, was used to estimate effective population size. We first estimated the population scaled mutation rate ($\Theta = 4N_e\mu$) using a Bayesian method, which permits for rate variation between loci, implemented in the R package *thetamateR*. The estimated Θ value for each species was then converted to N_e assuming an average silent site mutation rate in angiosperms of 5.35×10^{-9} sites/year (De La Torre *et al.* 2017) and a generation time of 10 years.

To infer changes if population size through time, we used the EBSP (Heled & Drummond 2008) approach in BEAST v2.4.2 (Bouckaert *et al.* 2014), following the methodology outlined in Trucchi et al. (Trucchi *et al.* 2014) for each species separately. A custom R script (Trucchi *et al.* 2014) was used to select only those loci with high numbers of SNPs (3 or more per locus) to maximise the amount of information per marker to increase the accuracy and reliability of our inference. From these 50 loci were selected randomly to use in the EBSP analysis. One haplotype was chosen at random for each individual at heterozygous loci to minimize bias in haplotype frequencies and reduce bias caused by undiscovered heterozygote samples (Trucchi *et al.* 2014). A separate substitution model was assigned for each SNP class with kappa linked between models, trees were unlinked across loci and a single strict clock was assigned to all loci. Operators were modified to increase mixing efficiency (Trucchi *et al.* 2014). Each MCMC ran for 500,000,000 generations, sampling every 50,000 and the level of stationarity (*R. robustior* was run for twice this length). Stationarity was verified using Tracer v1.6 (Drummond & Rambaut 2007). Multiple replicate runs were conducted to ensure the same stationarity point was reached and runs combined to increase ESS values to suitable levels (ESS > 100 for all important parameters). EBSPanalyser in BEAST was applied to perform a linear reconstruction with data from combined runs. If an additional, unused set of 50 highly-variable loci was available analyses were repeated and results compared across datasets.

Simulated data

We used fastsimcoal2 (v2.6)(Excoffier et al. 2013) to simulate sequence data for a genetic bottleneck 250 generations (2500 years) ago. For each species we simulated four strengths of bottleneck; a 90%, 50%, 10% and 0% reduction in population size with ancestral population set to our thetamateR N_e estimates for each species. We simulated 40,000-60,000 DNA sequences matching the length of the corresponding empirical RAD tags and selected loci with the same distribution of SNPs in the empirical

dataset (i.e., each SNP category contained the same number of tags). We then used these loci as input for our pipeline (described above). We repeated simulations three times to ensure consistency in demographic patterns across different simulations.

Acknowledgments: We thank the Howard Lloyd Davies legacy and NERC for funding. Rhian J. Smith, Simon Creer, Andrew Foote and Kathy Willis for support and comments.

Author Contributions: ASTP conceived and designed the study with input from all authors. AJH generated and analysed the data with contributions from ASTP. SC coordinated sample collection by RR and FR. WLE and WJB provided samples. AJH and ASTP wrote the manuscript with contributions from all authors.

References

Allnutt, T.F., Ferrier, S., Manion, G., Powell, G.V.N., Ricketts, T.H., Fisher, B.L., *et al.* (2008). A method for quantifying biodiversity loss and its application to a 50-year record of deforestation across Madagascar. *Conserv. Lett.*, 1, 173–181.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.H., Xie, D., *et al.* (2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS Comput. Biol.*, 10, 1–6.

Braude, S. & Templeton, A.R. (2009). Understanding the multiple meanings of 'inbreeding' and 'effective size' for genetic management of African rhinoceros populations. *Afr. J. Ecol.*, 47, 546–555.

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. (2013). Stacks: An analysis tool set for population genomics. *Mol. Ecol.*, 22, 3124–3140.

Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci.*, 114, 6089–6096.

Collen, B., Dulvy, N.K., Gaston, K.J., Gärdenfors, U., Keith, D.A., Punt, A.E., *et al.* (2016). Clarifying misconceptions of extinction risk assessment with the IUCN Red List. *Biol. Lett.*, 12, 20150843.

Crowley, B.E. (2010). A refined chronology of prehistoric Madagascar and the demise of the megafauna. *Quat. Sci. Rev.*, 29, 2591–2603.

Danthu, P., Rakotobe, M., Mauclere, P., Andrianoelisoa, H., Behra, O., Rahajanirina, V., *et al.* (2008). Essential oil production increases value of Psiadia altissima fallows in Madagascar ' s eastern

forests. *Agroforest Syst*, 72, 127–135.

Dolch, R., Ndriamiary, J., Ratolojanahary, T., Randrianasolo, M. & Ramanantenasoa, I. (2015). Improving livelihoods, training para-ecologists, enthralling children: Earning trust for effective community- based biodiversity conservation in Andasibe, eastern Madagascar. *Madagascar Conserv. Dev.*, 10, 21.

Doyle, J.J. & Doyle, J.L. (1987). A rapid DNA isolation procedure for small amounts of fresh leaf tissue. *Phytochem. Bull.*, 19, 11–15.

Drummond, A.J. & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.*, 7, 214.

Eberhart-Phillips, L.J., Hoffman, J.I., Brede, E.G., Zefania, S. & Martina, J. (2015). Contrasting genetic diversity and population structure among three sympatric Madagascan shorebirds: parallels with rarity, endemism and dispersal. *Ecol. Evol.*, 5, 997–1010.

Eiserhardt, W.L., Dransfield, J., Rakotoarinivo, M. & Baker, W.J. (2018). Four new species of Dypsis (Arecaceae: Arecoideae) from Madagascar. *Kew Bull.*, 73, 1–12.

Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V.C. & Foll, M. (2013). Robust Demographic Inference from Genomic and SNP Data. *PLoS Genet.*, 9.

Federman, S., Dornburg, A., Daly, D.C., Downie, A., Perry, G.H., Yoder, A.D., *et al.* (2016). Implications of lemuriform extinctions for the Malagasy flora. *Proc. Natl. Acad. Sci.*, 113, 5041–5046.

Federman, S., Dornburg, A., Downie, A., Richard, A.F., Daly, D.C. & Donoghue, M.J. (2015). The biogeographic origin of a radiation of trees in Madagascar : implications for the assembly of a tropical forest biome. *BMC Evol. Biol.*, 1–11.

Gazave, E., Ma, L., Chang, D., Coventry, A., Gao, F., Muzny, D., *et al.* (2013). Neutral genomic regions refine models of recent rapid human population growth. *Proc. Natl. Acad. Sci.*, 111, 757–762.

Godfrey, L.R., Scroxton, N., Crowley, B.E., Burns, S.J., Sutherland, M.R., Pérez, V.R., *et al.* (2019). A new interpretation of Madagascar's megafaunal decline: The "Subsistence Shift Hypothesis." *J. Hum. Evol.*, 130, 126–140.

Goyder, D., Baker, W., Besnard, G., Dransfield, J., Gardiner, L., Moat, J., *et al.* (2017). Country focus – status of knowledge of Madagascan plants. In: *State of the world's plants*. Royal Botanic Gardens, Kew.

Hackel, J., Vorontsova, M.S., Nanjarisoa, O.P., Hall, R.C., Razanatsoa, J., Malakasi, P., *et al.* (2018).

Grass diversification in Madagascar: In situ radiation of two large C3 shade clades and support for a Miocene to Pliocene origin of C4 grassy biomes. *J. Biogeogr.*, 45, 750–761.

Harper, G.J., Steininger, M.K., Tucker, C.J., Juhn, D. & Hawkins, F. (2019). Fifty years of deforestation and forest fragmentation in Madagascar. *Environ. Conserv.*, 34, 325–333.

Heled, J. & Drummond, A.J. (2008). Bayesian inference of population size history from multiple loci. *BMC Evol. Biol.*, 8, 289.

De La Torre, A.R., Li, Z., Van De Peer, Y. & Ingvarsson, P.K. (2017). Contrasting rates of molecular evolution and patterns of selection among gymnosperms and flowering plants. *Mol. Biol. Evol.*, 34, 1363–1377.

Lauterbur, M.E. (2019). Coalescent models at small effective population sizes and population declines are positively misleading. *bioRxiv*, 705335.

Lemmens, R.H.M.J. (2007). *Chrysophyllum boivinianum (Pierre) Baehni. PROTA (Plant Resour. Trop. Africa / Ressources végétales l'Afrique Trop.* Available at: https://uses.plantnet-project.org/en/Chrysophyllum_boivinianum_(PROTA). Last accessed 14 May 2019.

Lemmens, R.H.M.J., Louppe, D. & Oteng-Amoako, A.A. (2012). *Plant resources of tropical Africa 7 (2). Timbers 2.* PROTA Foundation, Wageningen, Netherlands.

Letsara, R. (2018). *Dilobeia thouarsii. IUCN Red List Threat. Species*. Available at: http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T70102219A70134142.en. Last accessed 22 May 2019.

Maes, D., Isaac, J.B.N., Harrower, A.C., Collen, B., Strien, J.A., Roy, B.D., *et al.* (2015). The use of opportunistic data for IUCN Red List assessments BT - Biological Journal of the Linnean Society. *Biol. J. Linn. Soc.*, 115, 690–706.

Moat, J. & Smith, P.P. (2007). *Atlas of the Vegetation of Madagascar*. Royal Botanic Gardens, Kew, London.

Pearson, R.G. & Raxworthy, C.J. (2009). The evolution of local endemism in Madagascar : Watershed versus climatic gradient hypotheses evaluated by null biogeographic models. *Evolution (N. Y).*, 63, 959–967.

Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S. & Hoekstra, H.E. (2012). Double digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One*, 7, e37135.

Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N., *et al.* (2015). The biodiversity of species and their rates of extinction, distribution, and protection. *Science (80-.).*,

344.

Rakotoarinivo, M. & Dransfield, J. (2012a). *Dypsis procumbens. IUCN Red List Threat. Species*. Available at: http://dx.doi.org/10.2305/IUCN.UK.2012.RLTS.T195965A2437132.en. Last accessed 22 May 2019.

Rakotoarinivo, M. & Dransfield, J. (2012b). *Ravenea robustior. IUCN Red List Threat. Species*. Available at: http://dx.doi.org/10.2305/IUCN.UK.2012.RLTS.T38678A2883023.en. Last accessed 22 May 2019.

Rakotoarinivo, M. & Dransfield, J. (2012c). *Satranala decussilvae. IUCN Red List Threat. Species*. Available at: http://dx.doi.org/10.2305/IUCN.UK.2012.RLTS.T38696A2883367.en. Last accessed 21 May 2019.

Rakotoarinivo, M., Dransfield, J., Bachman, S.P., Moat, J. & Baker, W.J. (2014). Comprehensive red list assessment reveals exceptionally high extinction risk to Madagascar palms. *PLoS One*, 9.

Trucchi, E., Gratton, P., Whittington, J.D., Cristofari, R., Le Maho, Y., Stenseth, N.C., *et al.* (2014). King penguin demography since the last glaciation inferred from genome-wide data. *Proc. Biol. Sci.*, 281, 20140528-.

Vences, M., Wollenberg, K.C., Vieites, D.R. & Lees, D.C. (2009). Madagascar as a model region of species diversification. *Trends Ecol. Evol.*, 24, 456–465.

Wilme, L., Goodman, S. & Ganzhorn, J. (2006). Biogeographic Evolution of Madagascar's Microendemic Biota, 312, 1063–1066.

Yoder, A.D., Campbell, C.R., Blanco, M.B., Ganzhorn, J.U. & Goodman, S.M. (2016). Geogenetic patterns in mouse lemurs (genus Microcebus) reveal the ghosts of Madagascar ' s forests past. *Proc. Natl. Acad. Sci.*, 113, 8049–8056.

Yoder, A.D., Rasoloarison, R.M., Goodman, S.M., Irwin, J.A., Atsalis, S., Ravosa, M.J., *et al.* (2000). Remarkable species diversity in Malagasy mouse lemurs (primates , Microcebus). *Proc. Natl. Acad. Sci.*, 97, 11325–11330.

Figure Legends

Figure 1 A map of collection sites for the 10 species used in this study. Stars represent approximate collection locations and are colour coded for each palm species sampled. The remaining six species were collected from a single location shown as a white star. Original map of Madagascar by Maky and shared under the CC BY-SA 3.0 license (https://creativecommons.org/licenses/by-sa/3.0/deed)

Figure 2 Effective population size through time for four species of palm that have conservation assessments. The present day is on the left side of each plot. Black lines represent the median population size and shaded polygons show the 95% highest posterior density (HPD). To the right of each plot is a picture of the study species and the current IUCN red list assessment. Values in red arrows indicate the percentage decline from maximum population size to the minimum population size inferred after the maximum. The increase in the median value of population size for *D. rabepierrei* in the very recent past is the result of a lack of coalescent events, which is reflected by extremely wide 95% HPD values during this period, and should not be interpreted as population size increase. Results of skyline plots for simulated data are shown as coloured Photo of *R. robustior* by Henk Beentje. All other photos by WJB.

Figure 3 Effective population size through time for six species from six different plant families. The present day is on the left side of each plot. Black lines represent the median population size and shaded polygons show the 95% highest posterior density (HPD). To the right of each plot is a picture of the study species and the current IUCN red list assessment, where available. Values in red arrows indicate the percentage decline from maximum population size to the minimum population size inferred after the maximum. As described in in figure 1, rapid increases towards the present in *C. boivinianum* and *Psychotria sp.* are due to lack of signal. Photos of *A. perrieri*, *C. boivinianum* and *D. thouarsii* by FR, Solofo Eric Rakotoarisoa and Rahaingoson Fabien respectively and shared under the CC BY-NC 4.0 license (https://creativecommons.org/licenses/by-nc/4.0/). Photo of *D. cordifolia* by Frank Vassen and shared under the CC BY 2.0 license (https://creativecommons.org/licenses/by/2.0/deed.en). Photo of *P. altissima* by B.navez shared under the CC BY-SA 3.0 license (https://creativecommons.org/licenses/by-sa/3.0/). Small modifications have been made to each photo.

☆ *Dypsis procumbens*
☆ *Dypsis rabepierrei*
☆ *Ravenea robustior*
☆ *Satranala decussilvae*
☆ All other species

a Census = ?
N_e = 16594

Arecaceae **NEAR THREATENED**
Dypsis procumbens

0%
10%
50%
90%

b Census < 20
N_e = 4272

Arecaceae **CRITICALLY ENDANGERED 90%**
Dypsis rabepierrei

c Census = 1000
N_e = 9405

Arecaceae **NEAR THREATENED**
Ravenea robustior

d Census = 200
N_e = 18093

Arecaceae **ENDANGERED 78%**
Satranala decussilvae

Population size (N_e*μ)

Time (mutations/site)

a Pop. status = ?
N_e = 15523
Passifloraceae
Adenia perrieri
NOT ASSESSED

b Pop. status = ?
N_e = 11673
Sapotaceae
Chrysophyllum boivinianum
NOT ASSESSED 83%

c Census > 1000
N_e = 25702
Melastomataceae
Dichaetanthera cordifolia
LEAST CONCERN

d Pop. status = decreasing
N_e = 10672
Proteaceae
Dilobeia thouarsii
LEAST CONCERN 82%

e Pop. status = stable
N_e = 16950
Asteraceae
Psiadia altissima
LEAST CONCERN 43%

f Pop. status = ?
N_e = 7254
Rubiaceae
Psychotria sp.
NOT ASSESSED 68%
?

Population size (N_e*μ)

Time (mutations/site)