Title:

SUMOylation of rice DELLA SLR1 modulates transcriptional responses and improves yield under salt stress

Authors:

Nuno M. Gonçalves[a*], Telma Fernandes[a], Cátia Nunes[b**], Margarida T. G. Rosa[a], Cleverson C. Matiolli[a], Mafalda A. A. Rodrigues[a], Pedro M. Barros[a], M. Margarida Oliveira[a], Isabel A. Abreu[a1]

Affiliations:

[a] Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa (ITQB NOVA), 2780-157, Oeiras, Portugal

[b] Instituto de Biologia Experimental Tecnológica (iBET), 2780-157, Oeiras, Portugal

Author Contributions:

NMG participated in the conception of research, executed most experiments, and wrote the article with contributions from all authors. TF did protein interaction experiments. MTGR characterized SUMO-protease rice lines. CCM analysed data and completed the writing. MAAR purified recombinant proteins. PMB analysed RNA seq. MMO supervised experiments and analysed results. IAA conceived original project, supervised experiments, and finalised the article.

Contact:

[1] Corresponding Author:

Isabel A. Abreu

Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa (ITQB NOVA), 2780-157, Oeiras, Portugal, Tel.: +351 214469647, E-mail: abreu@itqb.unl.pt

* Currently at Leibniz Institute of Plant Biochemistry (IPB), Weinberg 3 | 06120 Halle | Germany (nunomiguel.goncalves@ipb-halle.de)

** Currently at Instituto Nacional de Investigação Agrária e Veterinária, Avenida da República, Oeiras, Portugal (catia.soares@iniav.pt)

Short title (50 characters): SUMOylation of rice SLR1 increases salt tolerance

ABSTRACT

DELLA proteins modulate GA signalling and are major regulators of plant plasticity to endure stress. DELLAs are mostly regulated at the post-translational level, and their activity relies on the interaction with upstream regulators and transcription factors (TFs). SUMOylation is a post-translational modification (PTM) capable of changing protein interaction and found to influence DELLA activity in Arabidopsis. We determined that SUMOylation of the single rice DELLA SLENDER RICE1 (SLR1) occurs in a lysine residue different from the one identified in Arabidopsis REPRESSOR OF GA (RGA), which suggests differential DELLA regulation by SUMOylation in monocots and dicots. Remarkably, artificially increasing SUMOylated SLR1 (SUMO1SLR1) levels attenuated the penalty of salt stress on plant yield. Gene expression analysis revealed that the overexpression of SUMOylated SLR1 regulates key dioxygenases that modulate active GA levels, namely *GA20ox2* and *GA2ox3*, which could partially explain the sustained productivity upon salt stress imposition. Besides, SLR1 SUMOylation blocked the interaction with the growth regulator YAB4, which may fine-tune *GA20ox2* expression. Mechanistically, we propose that SLR1 SUMOylation disrupts the interaction with members of several transcription factor families to modulate gene expression. We found that SLR1 SUMOylation represents a novel mechanism modulating DELLA activity, which attenuates the impact of stress on plant performance.

Keywords: DELLA, SUMOylation, stress, productivity, SLR1, YAB4, BZIP23, gibberellin

One sentence summary: Rice plants show increased yield under salt stress when its gibberellin transcriptional regulator DELLA protein is artificially SUMOylated.

INTRODUCTION

Salinity is one of the most serious threats to food security, affecting 20% of irrigated soils worldwide. Rice (*Oryza sativa* L.) is a crop with high sensitivity to salinity compared to other monocot crops, such as wheat and barley (Munns and Tester, 2008). Plant sessile nature requires rapid physiological responses to cope with stress, which is partly achieved through a complex crosstalk between hormone signalling pathways affecting growth, flowering, and defence against environmental challenges (Vanstraelen and Benková, 2012). Gibberellins (GA) are diterpenoid hormones involved in the regulation of several developmental processes, including seed germination, stem elongation, and flowering (Hirano et al., 2008). GA responses are modulated by DELLA proteins, which act as transcriptional regulators (Hedden and Thomas, 2012). In the presence of GA, DELLAs are targeted to ubiquitin-mediated proteasome degradation by the GA-receptor GA-INSENSITIVE DWARF1 (GID1) through binding to specific DELLA domains (DELLA/TVHYNP), followed by DELLA polyubiquitination mediated by the E3-ligase GA-INSENSITIVE DWARF2/SLEEPY1 (GID2/SLY1). When GA levels decrease, DELLA proteins accumulate and interact with key transcriptional regulators to restrain growth (Hirano et al., 2010; 2008; Ueguchi-Tanaka et al., 2008). For instance, DELLAs can sequester the transcription factors PHYTOCHROME INTERACTING FACTOR3 and 4 (PIF3 and PIF4), in addition to negatively regulate PIF3 protein abundance, mediating the crosstalk of light and GA signalling to modulate cell growth (de Lucas et al., 2008; Li et al., 2016). DELLA transcriptional regulation of GA biosynthetic genes responsible for hormonal homeostasis also potentiates growth arrest. The single rice DELLA SLENDER RICE1 (SLR1) interacts with YAB4 transcription factor to prevent the induction of *GA20ox2*, which encodes a dioxygenase that is essential for the synthesis of bioactive GA (Yang et al., 2016). In Arabidopsis, DELLA interaction with GAI-ASSOCIATED FACTOR1 (GAF1) TF triggers a negative-feedback loop that activates GA biosynthesis and subsequent DELLA degradation (Fukazawa et al., 2017). The increasing number of validated DELLA interactors showcase an overlap with most hormone pathways, impacting development, fertility, and seed number (Davière and Achard, 2015; Van De Velde et al., 2017).

DELLA activity is mostly modulated through post-translational modifications (PTMs), which impact its ability to interact with transcriptional regulators and modulate gene expression (Van De Velde et al., 2017). Besides ubiquitination and

subsequent proteasome-mediated degradation, there are other PTMs that influence

the activity of the DELLA's in response to GA signalling. For instance, the

phosphorylation by kinase EARLY FLOWERING 1 (EL1) regulates SLR1 protein

stability and activity (Dai and Xue, 2010; Itoh et al., 2005). Recently, a mechanism of

antagonistic PTMs was proposed to regulate Arabidopsis DELLA's interaction with

specific protein partners. *O*-fucosylation by SPINDLY (SPY) promotes DELLA

interaction with essential growth regulators – such as BRASSINOLE-RESISTANT1

(BZR1), PIF3 and PIF4 – leading to growth arrest (Zentella et al., 2017). Conversely,

SLR1 *O*-GlcNAcylation by SECRET AGENT (SEC) blocks these interactions and

promotes GA-induced growth resumption (Camut et al., 2017; Zentella et al., 2016).

DELLA SUMOylation has also been indicated as a PTM controlling the stability of

REPRESSOR OF GA (RGA), one of the Arabidopsis five DELLA proteins. SUMO

attachment to RGA hinders its ubiquitin-mediated proteasome degradation through

blockage of GID1-RGA interaction in a GA-independent manner (Conti et al., 2014).

The stabilization of Arabidopsis DELLA by SUMOylation also positively affected salt

stress response, in line with observations that DELLA accumulation leads to

increased salinity tolerance (Achard et al., 2006; Conti et al., 2014; Nelis et al.,

2015). SUMOylation is a eukaryotic ubiquitination-like process that involves the

processing, activation, conjugation, and ligation of mature SMALL UBIQUITIN-LIKE

MODIFIER (SUMO) to acceptor lysines (K) in the target protein (Miura et al., 2007).

Unlike ubiquitination, SUMO-modifications often occur in a canonical consensus

motif (ψ-$\underline{K}$-X-E/D), which can be used for *in silico* prediction of SUMOylation targets

(Xue et al., 2006). SUMOylation regulates protein stability, activity, and interactions

with specific protein partners (Cremona et al., 2012). SUMO deconjugation

(deSUMOylation) from target proteins is performed by deSUMOylating proteases

(DSPs), with some members recently reported to be specific for poly-SUMO chain

processing (Eckhoff and Dohmen, 2015; Yates et al., 2016; Augustine et al., 2018).

In rice, reduced levels of SUMO conjugates due to DSP activity of OVERTOLERANT

TO SALT1/2 (OTS1/OTS2) have been connected to increased salt tolerance

(Srivastava et al., 2016a; 2016b). SUMOylation of specific TFs, such as the stress-

induced BASIC LEUCINE ZIPPER 23 (BZIP23), was suggested to be crucial to

drought tolerance and fertility (Srivastava et al., 2017).

Few studies have addressed the biological function of SUMOylation of stress-
induced targets. Moreover, most of the studies investigating the effects of PTMs on
DELLA function were done in Arabidopsis. This dicot model plant encodes five
DELLAs in the genome and essentially differs from monocots, which harbour a
single DELLA gene. This study demonstrates that SUMOylation of the single rice
DELLA SLR1 impacts the transcriptional output in response to salt stress, possibly
through modulation of SLR1 interaction with transcription factors regulating
transcriptional patterns governing plant growth and stress responses. We suggest
that SLR1 SUMOylation affects its interaction with key transcription factors that
impacts GA homeostasis (YAB4), ABA signalling (BZIP23), and JA signalling
(bHLH089/94). Additionally, the evidence provided suggests that it is possible to
uncouple stress response from decreased plant performance by manipulating the
levels of SUMOylated SLR1.
142
143

RESULTS

SLR1 SUMOylation *in vitro* analysis suggests diversification of DELLA regulation by SUMO in plants

SUMOylation of lysine 65 (K65) in Arabidopsis DELLA RGA modulates its stability in a GA-independent manner (Conti et al., 2014). To determine rice SLR1 SUMOylation, we started by performing an *in silico* prediction of SUMOylation motifs using SUMOsp2 software (Xue et al., 2006) in rice DELLA SLR1 and Arabidopsis DELLA RGA protein sequences. We sought for putative conserved SUMOylation motifs in both proteins and found a low probability SUMOylation site which corresponds to Arabidopsis RGA (Figure 1A). The SUMO acceptor lysine (K65) in RGA is part of a canonical SUMOylation motif (LKLE). In contrast, its counterpart in SLR1(K60) is not part of a canonical motif (QKLE), nor are the equivalent ones in another Arabidopsis or monocot DELLAs (Supplemental Figure S1). Nevertheless, both SLR1 and RGA have a high probability SUMOylation site on lysine 2 (K2).

To verify if the *in silico* prediction of SUMO attachment to SLR1 holds true, we performed an *in vitro* SUMOylation assay targeting both full-length SLR1 and a truncated isoform containing the leading 190 amino acids (SLR1t). Importantly, SLR1t includes the N-terminal domains (DELLA and TVHYNP) responsible for GA-mediated modulation of DELLA stability and SLR1 transactivation properties (Davière and Achard, 2015; Marín-de la Rosa et al., 2015; Hirano et al., 2012), as well as the predicted SUMOylation sites depicted in the Figure 1A. We observed a mobility-shift of both full-length SLR1 (Figure 1B) and truncated SLR1t (Figure 1C; with both SUMO1 and SUMO2 - Supplemental Figure S2A-B), consistent with ATP-dependent SUMO-conjugation in the presence of the rice SUMOylation machinery. The band resulting from *in vitro* SLR1 and SLR1t SUMOylation disappeared upon treatment with recombinant SUMO-specific protease ULP1 that cleaves SUMO from its conjugates (Li and Hochstrasser, 2003), confirming that the observed mobility-shift was caused by attached SUMO (Figure 1B-C). To analyse the contribution of each predicted SUMO acceptor lysine in SLR1, we performed directed mutagenesis changing the lysines K2 and K60 to arginine residues, which cannot be conjugated to SUMO (K2R and K60R). The mutated SLR1t K60R isoform retained the ability to be SUMOylated – which was confirmed by the band disappearance after SUMO proteolytic removal by ULP1 treatment – while K2R did not (Figure 1C). The result shows that K2 is the preferential SUMOylated residue in SLR1, suggesting that

Figure 1. SLR1 is SUMOylated *in vitro* in a novel SUMO-conjugation canonical motif. (A) *in silico* prediction of SLR1 canonical motifs (ψ-K-X-E) in the N-terminal region as scored by SUMOsp2 software (Xue et al., 2006), compared to *Arabidopsis thaliana* RGA corresponding region. Motifs showing high SUMOylation probability are marked in red, and low probability motifs are marked in blue. SLR1 K2 and K60 are shown in green and orange, respectively. **(B-C)** *in vitro* SUMOylation assays using rice SUMOylation machinery (OsSCE1, OsSAE1/2, and OsSUMO1 = ~ 11 kDa) with recombinant SLR1 (~ 65 kDa) and SLR1t (~ 19 kDa) proteins, treated with ULP1 SUMO protease, as indicated. SLR1 and SLR1t were detected by immunoblotting using custom made anti-SLR1 antibody, where **(B)** corresponds to the full-length SLR1, and **(C)** to the SLR1 N-terminal truncated (SLR1t) coding the leading 190 first SLR1 residues, as well as the mutated isoforms SLR1t K2R and SLR1t K60R.

SUMO attachment to DELLA occurs in distinct lysine residues of Arabidopsis RGA

(K65) and rice SLR1 (K2). SUMOylation in different sites potentially results in

different outcomes regarding protein interactions and activity of SUMOylated SLR1

compared to SUMOylated RGA.

SUMO protease activity analyses provide evidence of *in vivo* SLR1

SUMOylation

To demonstrate the *in vivo* SUMOylation of SLR1, we treated non-denatured

rice protein extracts with the ULP1 SUMO protease. The procedure would allow for

Figure 2. SUMO-specific protease activity reveal SLR1 *in vivo* regulation by SUMO.
(A) *Oryza sativa* L. cv Nipponbare non-denaturing protein extracts treated with *Saccharomyces cerevisiae* Ulp1 (ULP1) for SLR1 deSUMOylation analysis as detected by immunoblot using anti-SLR1 antibody. **(B)** Loading-normalized relative quantification of total SLR1 protein in *els1* and *fug1* mutants, and the respective negative segregants (NS) after deSUMOylation by ULP1 treatment; and **(C)** total SUMOylated SLR1, as shown by ratio comparison to non-SUMOylated protein after ULP1 treatment. Error bars correspond to standard error (SE; n=4). Statistically significant values are represented as **(p<0.01).

indirect detection of SUMOylated SLR1 by analysing the increase in non-
SUMOylated SLR1 protein amount resulting from ULP1 treatment, in comparison to
SUMOylated SLR1. We observed an increase of non-SUMOylated SLR1 upon the
addition of ULP1, as detected by the anti-SLR1 antibody in protein extracts (Figure
2A). The result is in line with the previous *in silico* and *in vitro* analyses indicating
that SLR1 could be SUMOylated. Our results also suggest that SLR1 mono-
SUMOylation may be much less abundant than other forms of SUMOylation since
we could not detect an SLR1 band with the expected mobility shift for mono-
SUMOylation (Figure 1B-C). To further support the evidence of *in vivo* SLR1
SUMOylation, we tested the levels of SUMOylated SLR1 in knockout mutant lines for
genes encoding rice SUMO proteases, namely *els1* and *fug1* (Figure 2B-C). ELS1
and FUG1 are active rice SUMO proteases, as both knockout lines contain
increased levels of SUMOylated proteins (Rosa et al., 2018). We treated protein
extracts from *els1* and *fug1* homozygous lines with ULP1 and compared their total

SUMOylated SLR1 levels with those present in negative segregant lines (NS). The
results show that although the overall level of non-SUMOylated SLR1 is not
significantly affected in either one of the KO lines when compared to their negative
segregant lines (Figure 2B), the levels of SUMOylated SLR1 are increased in *fug1*
mutant (Figure 2C). These results demonstrate SUMOylation of SLR1 occurs *in vivo*,
as indicated by the processing (*i.e.,* switch from SUMO-SLR1 to SLR1) mediated by
FUG1. Interestingly, it also suggests that there is a certain degree of specificity for
SLR1 deSUMOylation, as suggested by SUMO cleavage by FUG1 and not ELS1
(Figure 2C). Altogether, the evidence supports that SLR1 is SUMOylated *in planta*
and that SUMO specific proteases could regulate the levels of SUMOylated SLR1.
211

Overexpression of SUMOylated SLR1 increases salt tolerance

Because DELLA protein accumulation was previously connected with stress
tolerance (Achard et al., 2006), we assessed SLR1 expression in response to salt
stress. Wild-type seedlings accumulated SLR1 protein in shoots during the first 8
hours of stress (Supplemental Figure S3A). On the other hand, *SLR1* transcript
levels decreased, which is consistent with the previously reported DELLA negative
feedback-loop regulation on its own transcript abundance (Hedden & Thomas,
2012).

To gain insight on the role of SLR1 SUMOylation in rice development and
responses to salt stress, we generated transgenic lines overexpressing SLR1
(SLR1-OX), as well as lines constitutively expressing a mimicry of the SUMOylated
form of SRL1 (SUMO1SLR1-OX) originated from the genetic fusion of mature
SUMO1 to the K2 residue, which is the SUMOylated lysine of SLR1
($SUMO1_{GG}::SLR1_{K2}$) (Figure 3A). The strategy of using SUMOylated mimetic
isoforms originated by gene fusion with the SUMO coding sequence has been used
in several biological systems, including plants (Crozet et al., 2015; Ulrich, 2008). We
confirmed the overexpression of non-SUMOylated SLR1 isoforms in both SLR1-OX
and SUMO1SLR1, with the latter also showing a band with the expected mobility-
shift of SLR1 corresponding to mono-SUMOylation (Figure 3B). We also observed
an increase in the amount of non-SUMOylated SLR1 protein in SUMO1SLR1-OX
compared to WT and NS lines, which could be arising from the processing by
endogenous SUMO proteases because the fusion protein retains the SUMO
diglycine protease cleavage motif (-GGK-).

Figure 3. SLR1 SUMOylation confers salt tolerance at the seedling stage. (A) Depiction of gene constructs used to generate rice SLR1-OX and SUMO1SLR1-OX transgenic lines. *Zea mays* Ubi1 (ZmUbi1) promoter drives the expression of both OsSLR1 (SLR1-OX) and OsSLR1-OsSUMO1 gene fusion (SUMO1SLR1-OX). **(B)** SLR1 and SUMO1SLR1 protein accumulation in T2 independent transgenic lines (#1 and #2) of SUMO1SLR1-OX and SLR1-OX as detected by immunoblot using custom made anti-SLR1 antibody. **(C)** Stress severity as evaluated by IRRI salinity trial SES score for all lines in 6 and 8 days of stress and after 8 days of recovery period.

Rice is particularly sensitive to salt at both seedling and booting stages

(Gregorio et al., 1997). Therefore, we analysed stress marker symptoms of SLR1-

OX and SUMO1SLR1-OX lines subjected to high salt concentrations at seedling

stage, as well as yield parameters on salt treatments at booting stage. Evaluation of

symptoms in seedlings subjected to severe salinity (120mM NaCl) for 8 days showed

that both SUMO1SLR1-OX lines performed better under salt stress, displaying

decreased symptoms severity according to SES score (Standard Evaluation System

for Rice) by IRRI (International Rice Research Institute) (Gregorio et al., 1997),
including minimal leaf tip rolling and increased survival (Figure 3C). Importantly, salt
tolerance could be correlated with SUMOylated SLR1 levels, as SUMO1SLR1-OX
#1 – showing higher SLR1 SUMOylation levels compared to SUMO1SLR1-OX #2 –
performed better than the latter (Figure 3B-C). Interestingly, salt-induced
SUMOylated SLR1 increased alongside with the non-SUMOylated SLR1 in
SUMO1SLR1-OX lines (Supplemental Figure S3B), suggesting that SUMO cleavage
processing could be occurring alongside the duration of salt stress. Plants
overexpressing SUMO1SLR1 and SLR1-OX also recovered better than the wild type
Nipponbare in the 8 following days to salt stress cessation. Thus, mild
overexpression of SLR1 can increase rice tolerance to salinity, and this effect is
increased by its SUMOylation.

We next tested rice performance under salt stress at the booting stage.
Panicle initiation was the cue for mild salt stress imposition (80 mM NaCl), as at this
developmental stage pollen viability is most affected by high salinity (Negrão et al.,
2011; Sarhadi et al., 2012). Upon measurement of agronomical parameters, we did
not detect stress-induced changes in plant height (Figure 4A), tillering or panicle
length (Supplemental Figure S4) when compared to the control. However, two
critically important parameters for rice yield – namely fertility (filled grains) and
productivity (grain weight) – were significantly improved in SUMO1SLR1-OX lines
under mild salt stress. Both SLR1-OX and Nipponbare presented around 10%
decrease in fertility when comparing to mock conditions, while SUMO1SLR1-OX
showed the same fertility in both salt and mock treatments (Figure 4B). This
tendency was also observed in grain weight, where SUMO1SLR1-OX presented
higher tolerance to mild salt stress regarding productivity compared to Nipponbare
and SLR1-OX (Figure 4C). Although the SUMO1SLR1-OX lines had lower
productivity under mock conditions compared to Nipponbare, they showed improved
performance under salt stress. These results substantiate the significance of
SUMOylation of SLR1 in salt stress tolerance acquisition and yield protection
observed in two sensitive rice developmental stages.

SLR1 SUMOylation regulates GA biosynthetic pathway transcripts in response
to salt stress

Figure 4. SLR1 SUMOylation reduces the impact of salt stress on yield. Agronomic parameters as measured at the end of rice life cycle: **(A)** plant height (N=6) **(B)** fertility as evaluated by the number of filled grains ratio to total panicle flowers (n>10) and **(C)** productivity as evaluated by the weight of 1000 grains (n>10). Error bars correspond to standard error (SE; 2 independent biological replicates with N=6 each) and statistically significant p-values represented as *($p<0.05$).

The salt tolerance observed in SUMO1SLR1-OX lines could not be attributed

to the increased accumulation of SLR1, as SLR1 protein levels were similar in the

SLR1-OX and SUMO1SLR1-OX transgenic lines (Figure 3B). Thus, we inferred that

the observed phenotypes in SUMO1SLR1-OX lines could be a consequence of the

increased levels of SUMOylated SLR1. Because DELLA proteins regulate

transcription, we performed a genome-wide expression analysis to identify

differentially regulated genes in SLR1-OX and SUMO1SLR1-OX that could explain

the phenotypical differences observed in these lines in response to salt stress. We

exposed SLR1-OX, SUMO1SLR1-OX, and Nipponbare genetic background to high

salt concentrations (120 mM NaCl) during 8 hours at the seedling stage (2 weeks

old), and collected samples for subsequent RNA-seq analysis (Supplemental Table

S1). We selected the lines SLR1-OX #2 and SUMO1SLR1-OX #2 for the RNA-seq

assay because they showed similar non-SUMOylated SLR1 levels, as we aimed to

tackle the effect of SLR1 SUMOylation on gene expression. Both overexpressor

lines showed an approximately 1.8-fold increase in SLR1 transcript levels when

compared to Nipponbare, which is in line with the levels validated by real-time qPCR

for the same samples, *ca.* 1.5-fold increase (Supplemental Figure S5), and with

protein levels assessed by immunoblotting (Figure 3B). The induction of salt-induced

genes *Rab21* and *P5CS* was used as a transcriptional readout of the salt stress

treatment efficiency (Supplemental Figure S5).

Principal Component Analysis (PCA) revealed a clear separation of samples

by treatment, this factor being responsible for 47% of the variability in gene

expression, while genotype-specific transcriptomic changes explained 21% of the

observed variability (Supplemental Figure S6). The constitutive overexpression of

SLR1 and SUMO1SLR1 in absence of salt stress resulted in differential regulation of

247 and 207 genes, respectively (Supplemental Tables S1.6 to S1.8). Functional

annotation of SLR1-OX DEGs showed enrichment of gene ontology (GO) terms

associated with diterpenoid biosynthesis, highlighting the impact of SLR1 on GA

metabolism (Supplemental Figure 7A and Supplemental Tables S1.6 to S1.7).

Notably, we observed overall downregulation in genes coding for enzymes essential

in non-gibberellin diterpenoid synthesis, namely syn-copalyl diphosphate synthase

(*CPS4*) and ent-cassa-12,15-diene synthase (*KSL7*) (Supplemental Figures S7A and

S7C). These enzymes are part of a diterpenoid pathway branching point that feeds

the phytoalexins biosynthetic pathway (Otomo et al., 2004). Concurrently, other

genes related to phytoalexin and phytocassane synthesis, such as cytochrome P450

family members *CYP701A8/KO1, CYP99A3, CYP71Z6, CYP76M8,* and *CYP71Z7*,

were also downregulated in SLR1-OX.

Phenotypical discrimination of SLR-OX and SUMO1SLR1-OX was observed

under salt stress but not under control conditions (Figure 3C). Interestingly, a closer

look into the distribution of SLR-OX and SUMO1SLR1-OX samples in the PCA,

under control and stress conditions highlights a similar tendency (Supplemental Figure S6). Hence, we focused on the analysis on the 236 DEGs specific to SUMO1SLR1-OX exposed to salt stress. After functional annotation of DEGs specific to SUMO1SLR1-OX, we found a significant over-representation of GO terms associated with defence response and gibberellin dioxygenase activity (Supplemental Figures S7B and S7C). We also found a salt stress-dependent induction of *GA20ox2* and *GA2ox3* genes in SUMO1SLR1-OX (Supplemental Table S1.4). *GA20ox2* and *GA2ox3* encode enzymes involved in the regulation of active GA (GA_4) levels. Therefore, it is reasonable to hypothesize that SLR1 SUMOylation could adjust active GA homeostasis upon exposure to salt stress (Supplemental Figure S7C).

SUMOylation disrupts SLR1 interaction with stress-associated transcription factors

The increased salt tolerance and sustained productivity of SUMO1SLR1-OX under salt stress likely arise from the presence of increased levels of SUMOylated SLR1 (Figure 3B-C; Figure 4A-B), which seems to be affecting the gene expression profile of plants exposed to stress. The *GA20ox2* transcript was induced around 2-fold in SUMO1SLR1-OX exposed to salt stress, while it showed only a marginal induction in SLR1-OX under the same conditions (Supplemental Table S1.2). One possible mechanism by which SLR1 could affect *GA20ox2* transcription is through the interaction with the transcription factor YAB4, thereby repressing the expression of *GA20ox2* (Yang et al., 2016). We did not detect *YAB4* transcript changes after 8 hours of salt stress exposure, but a search in public transcriptome data revealed that *YAB4* is induced in rice shoots after 3h of salt stress imposition (Genevestigator - Hruz et al., 2008; Supplemental Figure S8A-B), which could lead to increased YAB4 protein accumulation during later stages of plant adaptation to salt stress. In an attempt to dissect the molecular pathways involved in *GA20ox2* upregulation in SUMO1SLR1-OX, we verified the interaction of SLR1 and SUMOylated SLR1 with YAB4 TF using a yeast two-hybrid system and bimolecular fluorescence complementation (BiFC). The aim was to test the effect of SUMOylation on SLR1-YAB4 interaction, which could affect *GA20ox2* transcription. We used a variant of the SUMO1SLR1 fusion protein lacking the SUMO protease recognition domain to avoid possible processing by yeast SUMO proteases (SUMO1$_{(GG-AA)}$::SLR1). We

Figure 5. SLR1 SUMOylation affects its interaction with key transcription factors
(A) Yeast-two hybrid (Y2H) analysis of both SLR1 and SUMO1SLR1. SUMO1::SLR1 fusion carries two mutations in the SUMO-protease cleavage motif (GG → AA) to avoid its removal by endogenous yeast proteases and guarantee constitutive SUMO attachment to SLR1 during the assay. Positive controls marked as (SD/-Leu/-Trip) and positive interactions marked with (SD/-Leu/-Trip/-Ade/-His) with C+ as an confirmed positive interaction control. **(B)** Bimolecular fluorescent complementation (BiFC) assays showing positive interactions between nYFP: SLR1 with cYFP:BHLH089, cYFP:BHLH094, and cYFP:YAB4. The presence of SUMO1 attached to lysine 2 of SLR1 (nYFP:SUMO1$_{(GG-AA)}$::SLR1) disrupts SLR1 interaction with these transcription factors. Negative controls for Y2H and BiFC assays can be found in Supplementary Figure S9.

confirmed the previously demonstrated interaction between SLR1 and YAB4 (Yang

et al., 2016), and also found that the presence of SUMO in lysine 2 of SLR1 disrupts

SLR1-YAB4 interaction in both yeast and *in planta* (Figure 5A-B; negative controls in

Supplementary Figure S9). The blocking of the SLR1 interaction with YAB4 by

SUMO correlates with the induction of *GA20ox2* in salt-stressed SUMO1SLR1-OX
lines.

To further evaluate the molecular aspects of SUMO1SLR1-OX tolerance to
salt stress, we explored the notion that SLR1 exerts its function mainly by
transcriptional regulation through interaction with transcription factors. We designed
a strategy that allowed us to identify SLR1 TF interactors that could explain, at least
partially, the differences in gene expression observed in the transgenic lines upon
exposure to salt stress. Briefly, we submitted the promoter sequences (1000 bp
upstream to the Transcription Start Site – TSS) of 417 selected DEGs (SLR1- and
SUMO1SLR1-dependent salt stress responsive genes; Supplemental Tables S1.3
and S1.4) to an *in silico* search for transcription factor binding sites (TFBS) in the
PlantPAN online database (Chow et al., 2016). Next, we selected only TFBS and
corresponding binding TFs that have been experimentally validated. This allowed us
to identify TFs from the TCP, bZIP, bHLH, and B3 families that could be regulating
the output gene expression of SLR1-OX and SUMO1SLR1-OX transgenic lines
under salt stress (Table 1). To cross-validate the TF identification, we made sure that
a substantial overlap between the known target genes of those TFs and the SLR1-
and SUMO1SLR1-dependent DEGs were observed (Supplemental Tables S1 and
S2).

The interaction between SLR1 with the selected candidates (Table 1) and its
regulation by SUMOylation was assessed through Y2H assays and BiFC analyses
(Figure 5A-B). The basic helix-loop-helix transcription factors bHLH089 and
bHLH094, as well as the homeobox TF OSH1, showed a strong interaction with
SLR1 but not with SUMO1$_{(GG-AA)}$::SLR1. This finding suggests that SUMOylation
could modulate the binding of SLR1 to these transcription factors, potentially
affecting their transcriptional activities. We also found that BZIP23 can interact with
both SUMOylated and non-SUMOylated SLR1 (Figure 5A). The interaction of SLR1
with BHLH089 and BHLH094, as well as its disruption by SLR1 SUMOylation, was
validated through BiFC in *Nicotiana benthamiana* leaves, supporting the *in planta*
SUMO-dependent regulation of these interactions (Figure 5B). Neither IDEF1 nor
PCF1 showed interactions with SLR1 or SUMO1$_{(GG-AA)}$::SLR1.

DISCUSSION

DELLA proteins are central regulators of plant productivity and stress responses, and the hormonal crosstalk mediated by these proteins is essential for plant plasticity to endure stress. As other hormone modulators, DELLA activity is regulated by post-translational modifications, such as ubiquitination, phosphorylation, *O*-fucosylation, *O*-GlcNAcylation, and SUMOylation (Yano et al., 2019; Zentella et al., 2016; Zentella et al., 2017; Conti et al., 2014; Dai and Xue, 2010; Itoh et al., 2003). Recently, SUMO proteases were linked to salt stress response in rice (Srivastava et al., 2016b; 2017). Although deSUMOylation was proposed as a general mechanism of tolerance acquisition in response to salt, we show that artificially SUMOylated SLR1 could alleviate the salt stress impact on rice productivity, also improving recovery after stress imposition is released.

Following the evidence that SUMO proteases are important for DELLA activity (Campanaro et al., 2016), the knockout mutant of SUMO protease *fug1* showed a higher ratio of overall SUMOylated SLR1. Interestingly, the FUG subclass is phylogenetically close to the SPF subclass, which contains Ulp2-like SUMO proteases (Castro et al., 2018). A Ulp2 from *Saccharomyces cerevisiae* (ScUlp2) was previously reported to be the key protease responsible for poly-SUMO chain processing (Eckhoff and Dohmen, 2015). If FUG1 is indeed related to Ulp2 proteases and it processes SUMO-SLR1, SLR1 deSUMOylation may be occurring in poly-SUMO chain high molecular weight conjugates.

DELLA proteins control several plant processes through interaction with a myriad of transcriptional factors (Van De Velde et al., 2017). By affecting protein-protein interaction, SLR1 SUMOylation has the potential to modulate plant processes by altering the transcriptional outcome of SLR1-TFs interacting proteins. Indeed, transcriptomic analysis of SUMOylated SLR1 overexpressing lines suggested that this isoform plays a role in early transcriptional responses to salt stress, affecting the expression of key genes coding for enzymes controlling the accumulation of active GA, such as *GA20ox2* and *GA2ox3*, involved in activation and inactivation of the active GA_4. We were able to determine that SLR1 SUMOylation occurs in lysine 2 (Figure 1A-C), distant from the conserved GA regulatory domains DELLA and TVHYNP. This finding suggests that SLR1 SUMOylation contrasts with Arabidopsis RGA, in which the amino acid residue responsible for SUMO modification is the

lysine 65 (Conti et al., 2014), located very close to the GA-regulatory domain

TVHYNP. The location of SUMO attachment in SLR1 protein could exert quite

different effects on protein-protein interactions compared to its well-described

Arabidopsis counterpart RGA. An interesting possibility is that SUMOylation of these

residues could have a different outcome regarding the transcriptional activation

properties of SLR1 DELLA/TVHYNP motifs (Hirano et al., 2012). Both lysines are

within SUMOylation canonical motifs, but while the SUMOylation canonical motif

surrounding K2 is conserved in all analysed DELLAs, the one containing K65 is

exclusive to RGA. The different SUMOylation sites of SLR1 and RGA might account

for alterations in DELLA regulation amongst species that probably determine

different interactome networks.

The SUMOylation of SLR1 could also be affecting the interaction with

upstream regulatory proteins. For instance, SUMO attachment could affect SLR1

interaction with SPY, which enhances SLR1 activity through *O*-fucosylation and is a

major determinant of rice plant architecture (Yano et al., 2019). Crosstalk between

SUMOylation and *O*-fucosylation could play a role in GA-promoted growth regulation

by decreasing TF-interactions that result in growth repression (Camut et al., 2017).

This control of the expression of GA enzymes is demonstrated by our transcriptomic

results of SUMO1SLR1-OX lines under salt stress, which shows upregulation of both

GA20ox2 and *GA2ox3*, suggesting that SLR1 SUMOylation could also regulate GA

homeostasis.

SUMOylation of specific targets was suggested to be crucial for stress

tolerance, fertility and yield maintenance in rice. For instance, BZIP63 SUMOylation

levels are modulated by OTS1, which affects its stability and the plant's ability to deal

with drought and salt stress (Srivastava et al., 2017). Interestingly, we found that

BZIP23 interacts with both SLR1 and SUMO1SLR1. Another novel interaction with

both bHLH089 and bHLH094 shows upregulation of JAZ-like repressors like JAZ8,

consistent with findings that this TF is regulating genes from the JA pathway as

observed in our transcriptomic analysis. Both these interactions may partly explain

the JA-induced stress response promoted by SLR1 accumulation (Supplemental

Figures S7A and S7B).

It was already shown that YAB4 interacts with SLR1 to repress *GA20ox2*

gene expression and restrain development in rice (Yang et al., 2016). We confirmed

the interaction between SLR1 and YAB4 by Y2H assays and BiFC (Figure 5B-C).

We additionally show that SUMOylation of SLR1 abolished its interaction with YAB4.
Accordingly, we found a stress-induced upregulation of *GA20ox2* in SUMO1SLR1-
OX (Supplemental Table S1). Moreover, *GA20ox2* was also found to be
downregulated by KNOTTED1-like homeobox (KNOX) proteins like OSH1
(Sakamoto et al., 2006). SLR1 SUMOylation also disrupted its interaction with
OSH1, suggesting that SLR1 SUMOylation mechanistic action is partly achieved
through SLR1 dissociation from transcriptional regulators involved in GA signalling
and stress signalling. It could represent a dissociation of DELLA function during salt
stress tolerance from its impact on productivity by means of SUMOylation.

The emergence of SUMOylation of SLR1 as an effector of rice tolerance to
salt stress, through the growth-maintaining modulation of protein-protein interaction
with transcriptional regulators such as YAB4, is evidence of the tight regulation of the
GA pathway by rice DELLA during stress. Recent studies have demonstrated that
DELLA proteins are targets of different post-translational modifications, which could
adjust their interactions with transcriptional regulators in a dynamic fashion, ensuring
the best developmental decisions when facing stress. Further studies need to be
performed to unveil upstream effectors that perceive stress and the underlying signal
transduction pathway that leads to SLR1 SUMOylation. By establishing the trigger
for SLR1 SUMOylation and its effect on stress-induced SLR1 protein-protein
interactions, new strategies can be developed involving the transient post-
translational modification of DELLA, as mutagenesis and selection of growth-
favourable SLR1 alleles. This can be used to activate the SUMO-regulated tolerance
mechanism when stress is forthcoming and protect crops from yield losses due to
abiotic stress.
478
479

METHODS

SLR1 genomic DNA cloning and vectors construction

The intronless gene SLR1 entire coding sequence was amplified from Nipponbare wild-type genomic DNA, using the primers SLR1-CDS-Fw and SLR1-CDS-Rv (Supplemental Table S4). The fragment was cloned into the pJET 1.2/blunt-end cloning vector (ThermoScientific), and the SLR1 full sequence confirmed by Sanger sequencing. SLR1 CDS was then amplified from pJET-SLR1 using the primers pET-Fw-*Eco*RI and pET-Rv-*Psp*OMI (Supplemental Table S4) to insert the restriction sites for *Eco*RI and *Psp*OMI to clone SLR1 in the pET28a expression vector (Novagen). For the generation of rice transgenic lines overexpressing SLR1, the SLR1 coding sequence was amplified from pJET-SLR1 using OsSLR1-GW-Fw and OsSLR1-GW-Fw primers to insert the attB1 and attB2 recombination sites. The amplified PCR-product was recombined into the pDONR221 (ThermoFisher).The SUMO1SLR1 gene fusion was generated using high-fidelity PCR (Phusion DNA Polymerase; NEB) essentially as described in Atanassov et al. (2009). Primers overlapping both the OsSUMO1 3'-end and OsSLR1 5'-end OL-SLR1+SUMO1-3'-Fw and OL SUMO1+SLR1-5'-Rv, were used in combination with OsSLR1-Gw-Rv and OsSUMO1-GW-Fw, respectively, to generate the fragments to be fused. In the next step, the PCR products were purified and used in the fusion PCR reaction containing the OsSUMO1-GW-Fw and OsSLR1-Gw-Rv primers for the final amplification of the final fusion product (Supplemental Table S4). The gene fusions were confirmed by Sanger sequencing. To obtain rice transgenic plant lines overexpressing SLR1 (SLR1-OX) and SLR1 fused to SUMO1 (SUMO1SLR1-OX), pDONR221 containing the inserts described above was used as entry vector for the pANIC6B destination vector (Mann et al., 2011). The expression of SLR1 and SUMO1SLR1 is controlled by the maize *ZmUbi1* promoter (Figure 3A).

Site-directed mutagenesis

For the production of truncated recombinant SLR1 protein (SLR1t) containing the leading 190 amino acid residues of SLR1, a stop codon was introduced by site-directed mutagenesis (SDM) in pET28a-SLR1 using *Pfu*Turbo DNA polymerase (Agilent) and the primers (SLR1t-STOP-Fw and SLR1t-STOP-Rv) designed with QuickChangeII online tool (Agilent) (Supplemental Table S4). Unmodified plasmids were digested with methylation-sensitive enzyme *Dpn*I (New England Biolabs). The

mutated plasmids were transformed in *E. coli* DH5α and the mutagenized site

confirmed by sequencing. Mutated pET28a coding the truncated SLR1 isoform

(SLR1t) was subjected to further mutagenesis procedures to perform the amino acid

substitutions K2R and K60R (primers SLR1t-K2R-Fw, SLR1t-K2R-Rv, SLR1t-K60R-

Fw and SLR1t-K60R-Rv, all listed in Supplemental Table S4) following the same

protocol.

Recombinant Protein Production and Purification

pET28a vectors containing the full-length SLR1, N-terminus truncated SLR1 (SLR1t),

SLR1t (K2R) and SLR1t (K60R) were transformed into *E. coli* strain BL21 Star (DE3,

Sigma-Aldrich) for recombinant protein expression. Cells were grown at 37ºC until

OD_{600} = 0.6, and protein production was induced by addition of 200 µM isopropyl-d-

1-thiogalactopyranoside (IPTG). After three hours, the cells were harvested by

centrifugation in a J2MI centrifuge (Beckman) at 7000 RPM during 20 minutes at

4ºC. The cells were resuspended in lysis buffer (phosphate buffer 20 mM pH 7.5,

500 mM NaCl, 10 mM Imidazole) containing 5 µg/mL of DNAse, 50 µg/mL of

lysozyme, PMSF 100 mM and $MgCl_2$ 25 mM. After freezing at -20ºC and thawing,

extracts were loaded into a HisTrap chelating column (Amersham Biosciences) and

proteins eluted in phosphate buffer containing 500 mM imidazole, following buffer

exchange dialysis (phosphate buffer 20 mM, NaCl 150 mM, DTT 1 M, glycerol 2%),

for increased protein stability. Recombinant proteins were analysed by Western Blot

using custom-made SLR1 specific polyclonal antibody with Mini-Protean II system

(Biorad).

In vitro **SUMOylation assays and ULP1 treatment**

The *in vitro* SUMOylation assays were performed in 20 µL of SUMOylation reaction

buffer (Werner et al., 2009) containing 3 µg of SLR1 recombinant protein, 1 µg of E1-

activating enzyme (OsSAE1/ OsSAE2), 0.8 µg of E2-conjugating enzyme

(OsSCE1a), 0.5 µg of OsSUMO1/2, 0.4 units of creatine phosphokinase, and 10 mM

creatine phosphate. Reactions were initiated by adding 1 mM ATP and then

incubated at 37ºC for 1 h to 3 h. To confirm SUMOylation specificity, 3 µg of purified

SUMO protease ULP1 from *Saccharomyces cerevisiae* (Li and Hochstrasser, 2003)

was used for deSUMOylation analysis. SUMO conjugates were analysed by Western

Blot with custom anti-SLR1 antibody. For ULP1 treatments, whole plants were

ground in liquid N_2 and 100 mg of ground tissue powder were mixed with non-

denaturing buffer (Tris-HCl pH 8.0 50 mM, sodium chloride 150 mM, NP-40 1%,

sodium deoxycholate 0.5%, SDS 0.1%, EDTA pH 8.0 1 mM, NEM 50 mM), followed

by homogenisation by vortexing, and subsequent centrifugation at 4 ºC for 20

minutes to recover the supernatant containing the proteins. The non-denaturing

protein extracts were treated with 7 µg of ULP1, incubated at 37 ºC with collection

points taken at 5, 30, 60, and 120 seconds. Proteins were denatured at 90ºC for 10

minutes with Laemmli Buffer with 5 mM DTT and anti-SLR1 immunoblot proceeded

as described above.

SLR1 protein extraction, detection and quantification analysis

Plant shoot samples were ground in liquid N_2 and 100 mg of ground tissue used for

protein extraction with denaturing TCA/Acetone precipitation, followed by

resuspension in lysis buffer (7 M Urea, 2 M Thiourea, 30 mM Tris, 4% (w/v) CHAPS,

4 % (v/v), cOmplete protease inhibitor cocktail EDTA-Free (Sigma-Aldrich), 0.1%

(v/v) Nuclease Mix 100x, and NEM 20 mM. Protein quantification was performed with

2-D Quant kit (GE Healthcare) following manufacturer's instructions. Anti-SLR1

immunoblots were performed in a Mini-Protean II System (Biorad), using custom-

made anti-SLR1 polyclonal specific antibody [1:5.000, overnight incubation] and anti-

rabbit (Novex) [1:20.000, 90 min incubation] in 5% non-fat dry milk in TBS 1X.

Quantification of native SLR1 bands provided by immunoblotting analysis,

normalized to protein loading control using stripped membrane stained with

Coomassie G-250 (Fluka), was performed with ImageJ software as indicated in the

official ImageJ website online tutorial.

Plant material and Rice transformation

Calluses induced from wild-type *Oryza sativa* (japonica spp.) cv. Nipponbare mature

seeds were used for rice transformations according to (Almadanim et al., 2017), and

selection for transformants asserted by hygromycin resistance and standard GUS-

staining with X-Gluc solution (Roth) of a section of seminal roots. Proteins were

extracted, detected and quantified using a modified TCA/Acetone method (Luís et

al., 2016). Due to growth arrest phenotype and fertility decrease in homozygous lines

of SUMO1SLR1-OX lines, all assays continued with positively-selected

heterozygous plants. Moreover, lines #2 for both SUMO1SLR1-OX and SLR1-OX

were selected for further analysis at rice booting stage and transcriptomic evaluation

according to non-SUMOylated SLR1 levels, in order to avoid DELLA pleiotropic

effects (Davière and Achard, 2015). Wild-type Nipponbare was used as a control for

SLR1-OX and SUMO1SLR1-OX obtained by transformation as described above.

N<egative segregant lines of Zhonghua and Hwayoung varieties were used as

controls for T-DNA insertion lines knock-out mutants for rice SUMO proteases ELS1

(04Z11JY66) and FUG1 (2A-20225 obtained from RMD (www.rmd.ncpgr.cn),

developed by the National Centre of Plant Gene Research) and Postech

(www.postech.ac.kr/life/pfg/risd, developed by Gynheung An), respectively. These

insertion line seeds were germinated, self-crossed and screened using PCR analysis

for homozygosity (Rosa et al., 2018). The T-DNA insertion position was determined

by sequencing. Gene expression was analysed by RT-qPCR (Rosa et al., 2018).

Salinity Stress Assays

For the assays performed at the seedling stage, heterozygous transgenic seeds

were germinated and selected through screening for hygromycin resistance and

GUS staining. Seedlings were transferred to hydroponics for salinity resistance

screening (Almeida et al., 2016) with half-strength Murashige & Skoog Medium and

MES buffer 250 mg/L (Duchefa) with pH adjusted to 5.2. Seedlings were grown for

14 days before the salt stress assay. Then, the seedlings were exposed to salt

stress, with electric conductivity (EC) set and maintained at 12 dS/m with sodium

chloride solution for the extension of the stress treatment. After 8 days, the seedlings

were transferred to regular medium for recovery. Stress severity (SES score) – 1 for

lower severity and 9 for higher severity - was evaluated as set by IRRI (Gregorio et

al., 1997) at 6 and 8 days after stress and 8 days after recovery. Samples for

transcriptomic analysis – 4 replicates of a pool of 3 plants - were collected at 8 hours

after the initiation of the salt stress, corresponding to 10 hours after the lights went

on, for both control and salt treatments, and frozen immediately in liquid nitrogen for

grinding and analysis. Samples from 0h, 2h, 4h, and 8h treatments were obtained

from shoots and roots collected separately, and frozen immediately in liquid nitrogen.

Protein was extracted, detected and quantified as described above. For the booting

stage assay, previously germinated and selected heterozygous transgenic seeds

were sown in pots with 2-parts soil, 2-parts peat, and 1-part vermiculite. The

experiment was set up in the greenhouse from June to September, according to IRRI

standard screening procedures (Gregorio et al., 1997). Genotypes were maintained

in separate containers, for both stress and control pots, to control for discrepancies

in panicle initiation times. Panicle initiation was evaluated by the emergence of the

flag-leaf with a distance between 1 and 2 cm to the adjacent leaf and salinity applied

individually for each genotype at that point. Electric conductivity (EC) was set and

maintained at 8 dS/m with sodium chloride solution for the extension of the life cycle.

Upon complete maturation of the seed, agronomical parameters were measured

(plant height, tillering, and panicle length) and panicles collected for further

characterization of fertility (analysing the number of full matured seeds to total

number of flowers ratio) and productivity (weight of 1000 seeds as measured in a

Pfeuffer seed counter). The significance of statistics was asserted by t-test with

Holm-Sidak method correction for multiple comparisons and a p<0.05 threshold.

RNA Extraction and expression analysis

RNA was extracted from a total of 100 mg of ground frozen shoot tissue with Direct-

zol kit (Zymo Research) according to manufacturer's protocol with in-column DNAse

treatment. The first-strand cDNA was synthesized from 2 µg of total RNA with an

anchored-oligo (dT)18 primer using the Transcriptor High Fidelity cDNA Synthesis

Kit (Roche) according to the manufacturer's instructions. Reverse Transcription

quantitative PCR (RT- qPCR) analysis was performed using a LightCycler 480

system (Roche) and the SYBR Green I Master mix (Roche). The RT-qPCR reactions

were carried out with the primers listed in Supplemental Table S4. PCR running

conditions were: one cycle at 95ºC for 5 min and 45 cycles of amplification at 95ºC

for 10 s, 56–60ºC for 10 s and 72ºC /10 s. The CT values were calculated using

three technical replicates, and gene expression was assessed relative to described

internal reference transcript controls, rice *UBIQUITIN CONJUGASE2* (OsUBC2) and

UBIQUITIN10 (OsUBQ10) (Moraes et al., 2015; Pabuayon et al., 2016). For the

RNA-seq procedure, the small RNAs in the extracted total RNA samples were

cleaned by using the RNeasy spin column from RNeasy Mini Kit (Qiagen) following

RNA Cleanup protocol. RNA quality was assessed by Fragment Analyzer

(ThermoFisher) with RNA Quality Number over 5. A total of 18 cDNA libraries were

prepared, using three biological replicates for each genotype (SLR1-OX,

SUMO1SLR1-OX, Nipponbare) and each condition (mock and salt). Enzymatic

fragmentation with Pico Nextera kit and Smart-Seq2 multiplexed-library

preparation were performed[49] and single-end sequencing on NextSeq500 (Illumina)

with High Output Kit (75 cycles).

RNA-seq Data Analysis

Raw data FASTQ files were inputted on R/Bioconductor-based user interface

analysis software Chipster for RNA-seq. Single-end reads were trimmed with

PRINSEQ for 55 bp total with 18 bp trimmed on 5' end and quality control of 28% set

for the 3' end for adapter removal. Read alignment was performed with TOPHAT2

against the *Oryza sativa* genome (version IRGSP-1.0.36), selecting Sanger-Phred

+33 as quality score format. An average of 25.000.000 and 21.000.000 aligned and

uniquely mapped reads, respectively, were obtained per library (Supplemental Table

S2). For differential expression analysis, BAM files resulting from alignment were

submitted to pairwise comparisons with Cufflinks2 tool Cuffdiff for a set 0.05 False

Discovery Rate (5% FDR) with sequence-specific bias and multi-mapped read

corrections enabled. Differentially expressed genes (DEGs) with a 1.5 fold-change

were selected for data visualization and functional annotation using ClueGO plug-in

(Bindea et al., 2009) for Cytoscape (Su et al., 2014).

Promoter Analysis and Interaction Validation by Y2H and BiFC

The 1000 bp promoter region of differential expressed genes DEGs were

downloaded from Rice Annotation Project Database (RAP-DB) (Sakai et al., 2013)

and submitted to PlantPAN 2.0 (Chow et al., 2016) for *cis*-elements for transcription

factor binding site (TFBS) analysis. Gene Group Analysis was used to search for

common (90%) *cis*-element binding (TFBS) sites and the transcription factors (TFs)

associated with DEG promoters. Rice TFs were examined individually for

experimentally verified data concerning the interaction to binding sites and

downstream target genes listing (Supplemental Table S2). Resulting target genes

were then overlapped with previously verified DEGs from RNA-seq analysis. For the

Y2H assays, coding sequences of SLR1 and OsSUMO1$_{(GG-AA)}$::SLR1 were cloned

into the pGADT7 vector (Stratagene, USA) while OsBHLH089, OsBHLH094,

OsBZIP23, OsIDEF1, OsOSH1, OsPCF1 and OsYAB4 were cloned into the

pGBKT7 (Stratagene, USA) vector. These constructs were transformed into

Saccharomyces cerevisiae strain Y2HGold (Clontech, US) applying the LiAc method.

Yeast transformed cells were plated on synthetic defined (SD) medium lacking

leucine and tryptophan for plasmid transformation control and in SD lacking leucine

(-Leu), tryptophan (-Trip), adenine (-Ade), and histidine (-His) for interaction

screening. The interactions were evaluated in three individual colonies transformed

with both plasmids. pAD-WT and pBD-WT from the HybriZAP 2.1 kit (Stratagene,

USA) were used as a control of positive interaction (C+). pGBKT7 and pGADT7

empty vectors were also co-transformed with pGADT7-SLR1, pGADT7-SUMO1$_{GG-AA}$)::SLR1 and pGBKT7-BHLH089, pGBKT7-BZIP23, pGBKT7-IDEF1, and pGBKT7-

AA)::SLR1 and pGBKT7-BHLH089, pGBKT7-BZIP23, pGBKT7-IDEF1, and pGBKT7-

YAB4, respectively, as negative controls. For bimolecular fluorescence

complementation (BiFC), OsSLR1, OsSUMO1$_{GG-AA}$)::SLR1 were cloned in YFN43,

and OsBHLH089, OsBHLH094 and OsYAB4 were cloned into YFC43 (Belda-

Palazón et al., 2012). *Nicotiana benthamiana* leaves were co-infiltrated with

Agrobacterium tumefaciens strain EH105 harbouring the appropriate vectors.

Fluorescence was visualized using the Leica DM6 B fluorescence microscope (40x)

with L5 filter cube (Leica, Germany) 48-72h following infiltration. As negative control,

YFN43 (empty vector) was also co-infiltrated with YFC43-BHLH089, YFC43-

BHLH094 and YFC43-YAB4. *Agrobacterium* expressing the viral silencing

suppressor P19 was included in all infiltrations to increase the expression in *N.*

benthamiana leaves. BiFC analyses were performed essentially as described in

Belda-Palazón et al., 2012.

Accession Numbers

Sequence data from this article can be found in the Rice Annotation Project

Database (RAP-DB) under the following accession numbers: OsSLR1

(Os03g0707600), OsSUMO1 (Os01g0918300), OsSUMO2 (Os01g0918200),

OsSAE1 (Os11g0497000), OsSCE1a (Os03g0123100), OsELS1 (Os01g0355900),

OsFUG1 (Os03g0344300), OsUBC2 (Os02g0634800), OsUBQ10 (Os06g0681400),

OsP5CS (Os05g0455500), OsRAB21 (Os11g0454300), RGA (At2g01570), OsCPS4

(Os04g0178300), OsKSL7 (Os02g0570400), OsCYP701A8/KO1 (Os06g0569500),

OsCYP99A3 (Os04g0178400), OsCYP71Z6 (Os02g0570500), OsCYP76M8

(Os02g0569400), OsCYP71Z7 (Os02g0570700), OsCKX2 (Os01g0197700),

OsCKX5 (Os01g0775400), OsLOX2 (Os08g0509100), OsCHT3 (Os06g0726100),

OsCHT8 (Os10g054290), OsCHT11 (Os03g0132900), OsGA20ox2

(Os01g0883800), OsGA2ox3 (Os01g0757200), OsPCF1 (Os04g0194600),

OsRISBZ3   (Os02g0266800),   OsBHLH089   (Os03g0802900),   OsBHLH094
(Os07g0193800), OsBZIP23 (Os02g0766700), OsOSH1 (Os03g0727000), OsPCF1
(Os03g0785800), OsIDEF1 (Os11g0175700), OsYAB4 (Os02g0643200).
720
721

Supplementary Materials

Figure S1. Alignment of DELLAs protein sequences of selected monocots and dicots.

Figure S2. SRL1 can be SUMOylated by both OsSUMO1 and OsSUMO2 *in vitro.*

Figure S3. Accumulation dynamics of SRL1 transcript and protein in response to salt stress.

Figure S4. Tillering and panicle length were not affected by mild salt stress imposition.

Figure S5. Analysis of salt treatment efficiency by salt-stress marker gene induction.

Figure S6. Salt treatment had a major contribution to the differences in gene expression among Nipponbare, SLR1-OX and SUMO1SLR1-OX.

Figure S7. Transcriptomics reveal differential stress-induced regulation by SUMOylated SLR1.

Figure S8. Genevestigator Results for Gene Expression of YABBY4 (LOC_Os02g4950) in Salinity Assays.

Figure S9. Protein-protein interaction (PPI) negative controls for Y2H and BiFC assays.

Table S1. RNA seq data showing >1.5 fold Differentially Expressed Genes in Nipponbare, SLR1-OX and SUMOSLR1-OX, in mock and salt stress conditions.

Table S2. Expression changes of selected putative SLR1 TF interactors (A) and the respective downstream target genes (B). Fold-change shown as $\log_2$ FC. Comparison of condition A with condition B corresponding to respective selected analysis in Cufflinks (FPKM1 x FPKM2, for condition A and B, respectively). Nipp-Nipponbare; (m)-mock; (s)-stress

Table S3. RNA-seq read count for each of the sequenced libraries and respective legend

Table S4. List of primers used for gene cloning, site directed mutagenesis (SDM) and gene expression quantification (RT-qPCR). Details for the use of specific primers are given in the text.

Acknowledgments

Portuguese Fundação para a Ciência e a Tecnologia (FCT) for fellowships for NMG (SFRH/BD/85799/2012), TF (PD/BD/135584/2018), MTGR (SFRH/BD/84219/2012), MAAR (SFRH/BD/87420/2012), PMB (SFRH/BPD/86742/2012). FCT Investigator IF/00764/2014 (POPH-QREN) for IAA and PTDC/BIA-FBT/31211/2017 for CM contract. Work was supported by P-KBBE/0001/2012 and also by the FCT research unit GREEN-it "Bioresources4sustainability" (UID/Multi/04551/2013 and UIDB/04551/2020). The funding sources had no involvement in study design, analyses, and interpretation of data, writing, or in the submission decision.

Table 1. Enriched binding sites and associated transcription factors in SLR1 DEGs promoters. Promoter sequence analysis output from PlantPAN with Support referencing the percentage of transcription factor binding elements (TFBS/Sequence) found in SLR1- and SUMO1SLR1-dependent stress-responsive DEGs. Resulting TF overlap with DEGs detailed in Supplementary Table S4.

TFBS	Support	TF Family	TF	Annotation
GGTCCCAC	91.4	TCP	PCF1/2	essential for meristematic tissue-specificity expression of the PCNA gene promoter
TACGTA	91.4	(Others)	RISBZ3	found in seed storage protein gene promoters
GCACGTGC	94.7	bHLH	BHLH089/094	forms a ternary complex with RSS3 and TIFY11A/JAZ9 to negatively regulate jasmonate-responsive genes
ACACGTGT	97.4	bZIP	BZIP23/46	binds specifically to the abiotic stress ABA-responsive elements (ABRE) in the promoter of target genes
TNCGTACAA	98.0	SBP	SPL1	trans-acting factor that binds specifically to the consensus nucleotide sequence 5'-TNCGTACAA-3'
TGAC	98.7	Homeobox	OSH1	involved in shoot formation during embryogenesis
CATGC	100.0	B3	IDEF1	involved in iron deficiency response and tolerance
CCAAT	100.0	NF-YB	NFBY1/2/3/4	component of the NF-Y/HAP transcription factor complex regulating the expression of photosynthetic genes

Parsed Citations

Achard, P., Cheng, H., De Grauwe, L., Decat, J., Schoutteten, H., Moritz, T., Van Der Straeten, D., Peng, J., and Harberd, N.P. (2006). Integration of plant responses to environmentally activated phytohormonal signals. Science 311, 91–94.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Almadanim, M.C., Alexandre, B.M., Rosa, M.T.G., Sapeta, H., Leitão, A.E., Ramalho, J.C., Lam, T.T., Negrão, S., Abreu, I.A, and Oliveira, M.M. (2017). Rice calcium-dependent protein kinase OsCPK17 targets plasma membrane intrinsic protein and sucrose-phosphate synthase and is required for a proper cold stress response. Plant Cell Environ. 40, 1197–1213.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Almeida, D.M., Almadanim, M.C., Lourenço, T., Abreu, I.A, Saibo, N.J.M., and Oliveira, M.M. (2016). Screening for Abiotic Stress Tolerance in Rice: Salt, Cold, and Drought. In Environmental Responses in Plants, (New York, NY: Humana Press, New York, NY), pp. 155–182.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Atanassov, I.I., Atanassov, I.I., Etchells, J., and Turner, S.R. (2009). A simple, flexible and efficient PCR-fusion/Gateway cloning procedure for gene fusion, site-directed mutagenesis, short sequence insertion and domain deletions and swaps. Plant Methods 2009 5:1 5, 14.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Augustine, R.C. and Vierstra, R.D. (2018). SUMOylation: re-wiring the plant nucleus during stress and development. Current opinion in plant biology, 45, pp.143-154.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Belda-Palazón, B., Ruiz, L., Martí, E., Tárraga, S., Tiburcio, A.F., Culiáñez, F., Farràs, R., Carrasco, P. and Ferrando, A. (2012). Aminopropyltransferases involved in polyamine biosynthesis localize preferentially in the nucleus of plant cells. PLoS One, 7(10), p.e46907.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A, Fridman, W.-H., Pagès, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. Bioinformatics 25, 1091–1093.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Campanaro, A, Battaglia, R., Galbiati, M., Sadanandom, A, Tonelli, C., and Conti, L. (2016). SUMO proteases OTS1 and 2 control filament elongation through a DELLA-dependent mechanism. Plant Reprod 29, 1–4.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Camut, L., Davière, J.-M., and Achard, P. (2017). Dynamic Regulation of DELLA Protein Activity: SPINDLY and SECRET AGENT Unmasked! Mol Plant 10, 785–787.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Castro, P.H., Bachmair, A, Bejarano, E.R., Coupland, G., Lois, L.M., Sadanandom, A, van den Burg, H.A., Vierstra, R.D., and Azevedo, H. (2018). Revised nomenclature and functional overview of the ULP gene family of plant deSUMOylating proteases. J. Exp. Bot. 69, 4505–4509.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Chow, C.-N., Zheng, H.-Q., Wu, N.-Y., Chien, C.-H., Huang, H.-D., Lee, T.-Y., Chiang-Hsieh, Y.-F., Hou, P.-F., Yang, T.-Y., and Chang, W.-C. (2016). PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants. Nucleic Acids Research 44, D1154–D1160.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Conti, L., Nelis, S., Zhang, C., Woodcock, A, Swarup, R., Galbiati, M., Tonelli, C., Napier, R., Hedden, P., Bennett, M., et al. (2014). Small Ubiquitin-like Modifier protein SUMO enables plants to control growth independently of the phytohormone gibberellin. Developmental Cell 28, 102–110.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cremona, C.A, Sarangi, P., and Zhao, X. (2012). Sumoylation and the DNA Damage Response. Biomolecules 2012, Vol. 2, Pages 376-388 2, 376–388.
Pubmed: Author and Title

Google Scholar: Author Only Title Only Author and Title

Crozet, P., Margalha, L., Butowt, R., Fernandes, N., Elias, C.A, Orosa, B., Tomanov, K., Teige, M., Bachmair, A, Sadanandom, A, et al. (2015). SUMOylation represses SnRK1 signaling in Arabidopsis. The Plant Journal 85, 120–133.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dai, C., and Xue, H.-W. (2010). Rice early flowering1, a CKI, phosphorylates DELLA protein SLR1 to negatively regulate gibberellin signalling. The EMBO Journal 29, 1916–1927.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Davière, J.-M., and Achard, P. (2015). A pivotal role of DELLAs in regulating multiple hormone signals. Mol Plant 9, 10–20.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

de Lucas, M., Daviere, J.M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S., Fankhauser, C., Blazquez, M.A, Titarenko, E., and Prat, S. (2008). A molecular framework for light and gibberellin control of cell elongation. Nature 451, 480-484.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Eckhoff, J., and Dohmen, R.J. (2015). In Vitro Studies Reveal a Sequential Mode of Chain Processing by the Yeast SUMO (Small Ubiquitin-related Modifier)-specific Protease Ulp2. J. Biol. Chem. 290, 12268–12281.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fukazawa, J., Mori, M., Watanabe, S., Miyamoto, C., Ito, T., and Takahashi, Y. (2017). DELLA-GAF1 Complex Is a Main Component in Gibberellin Feedback Regulation of GA20 Oxidase 2. Plant Physiology 175, 1395–1406.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gregorio, G.B., Senadhira, D., and Mendoza, R.D. (1997). Screening rice for salinity tolerance. Irri.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hedden, P., and Thomas, S.G. (2012). Gibberellin biosynthesis and its regulation. Biochemical Journal 444, 11–25.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hirano, K., Asano, K., Tsuji, H., Kawamura, M., Mori, H., Kitano, H., Ueguchi-Tanaka, M., and Matsuoka, M. (2010). Characterization of the molecular mechanism underlying gibberellin perception complex formation in rice. The Plant Cell Online 22, 2680–2696.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hirano, K., Ueguchi-Tanaka, M., and Matsuoka, M. (2008). GID1-mediated gibberellin signaling in plants. Trends Plant Sci 13, 192–199.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hirano, K., Kouketu, E., Katoh, H., Aya, K., Ueguchi-Tanaka, M. and Matsuoka, M. (2012). The suppressive function of the rice DELLA protein SLR1 is dependent on its transcriptional activation activity. The Plant Journal, 71(3), pp.443-453.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W. and Zimmermann, P. (2008). Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. Advances in bioinformatics.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Itoh, H., Matsuoka, M., and Steber, C.M. (2003). A role for the ubiquitin-26S-proteasome pathway in gibberellin signaling. Trends Plant Sci 8, 492–497.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Itoh, H., Sasaki, A, Ueguchi-Tanaka, M., Ishiyama, K., Kobayashi, M., Hasegawa, Y., Minami, E., Ashikari, M., and Matsuoka, M. (2005). Dissection of the phosphorylation of rice DELLA protein, SLENDER RICE1. Plant Cell Physiol. 46, 1392–1399.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Li, K., Yu, R., Fan, L.M., Wei, N., Chen, H. and Deng, X.W. (2016). DELLA-mediated PIF degradation contributes to coordination of light and gibberellin signalling in Arabidopsis. Nature communications, 7, p.11868.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Li, S.-J., and Hochstrasser, M. (2003). The Ulp1 SUMO isopeptidase: distinct domains required for viability, nuclear envelope localization, and substrate specificity. 160, 1069.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Luís, I.M., Alexandre, B.M., Oliveira, M.M., and Abreu, I.A. (2016). Selection of an Appropriate Protein Extraction Method to Study the Phosphoproteome of Maize Photosynthetic Tissue. Plos One 11, e0164387.

Mann, D.G.J., LaFayette, P.R., Abercrombie, L.L., King, Z.R., Mazarei, M., Halter, M.C., Poovaiah, C.R., Baxter, H., Shen, H., Dixon, R.A., et al. (2011). Gateway-compatible vectors for high-throughput gene functional analysis in switchgrass (Panicum virgatum L.) and other monocot species. - PubMed - NCBI. Plant Biotechnol. J. 10, 226–236.

Marín-de la Rosa, N., Pfeiffer, A., Hill, K., Locascio, A., Bhalerao, R.P., Miskolczi, P., Grønlund, A.L., Wanchoo-Kohli, A., Thomas, S.G., Bennett, M.J. and Lohmann, J.U. (2015). Genome wide binding site analysis reveals transcriptional coactivation of cytokinin-responsive genes by DELLA proteins. PLoS genetics, 11(7), p.e1005337.

Miura, K., Jin, J.B., and Hasegawa, P.M. (2007). Sumoylation, a post-translational regulatory process in plants. Current Opinion in Plant Biology 10, 495–502.

Moraes, G.P., Benitez, L.C., do Amaral, M.N., Vighi, I.L., Auler, P.A., da Maia, L.C., Bianchi, V.J., and Braga, E.J.B. (2015). Evaluation of reference genes for RT-qPCR studies in the leaves of rice seedlings under salt stress. Genet. Mol. Res. 14, 2384–2398.

Munns, R., and Tester, M. (2008). Mechanisms of salinity tolerance. Annu. Rev. Plant Biol. 59, 651–681.

Negrão, S., Courtois, B., Ahmadi, N., Abreu, I., Saibo, N., and Oliveira, M.M. (2011). Recent Updates on Salinity Stress in Rice: From Physiological to Molecular Responses. Critical Reviews in Plant Sciences 30, 329–377.

Nelis, S., Conti, L., Zhang, C., and Sadanandom, A. (2015). A functional Small Ubiquitin-like Modifier (SUMO) interacting motif (SIM) in the gibberellin hormone receptor GID1 is conserved in cereal crops and disrupting this motif does not abolish hormone dependency of the DELLA-GID1 interaction. Plant Signaling & Behavior 10, e987528.

Otomo, K., Kenmoku, H., Oikawa, H., König, W.A., Toshima, H., Mitsuhashi, W., Yamane, H., Sassa, T., and Toyomasu, T. (2004). Biological functions of ent- and syn-copalyl diphosphate synthases in rice: key enzymes for the branch point of gibberellin and phytoalexin biosynthesis. Plant J. 39, 886–893.

Pabuayon, I.M., Yamamoto, N., Trinidad, J.L., Longkumer, T., Raorane, M.L., and Kohli, A. (2016). Reference genes for accurate gene expression analyses across different tissues, developmental stages and genotypes in rice for drought tolerance. Rice 9, 32.

Rosa, M.T.G., Almeida, D.M., Pires, I.S., da Rosa Farias, D., Martins, A.G., da Maia, L.C., de Oliveira, A.C., Saibo, N.J.M., Oliveira, M.M., and Abreu, I.A. (2018). Insights into the transcriptional and post-transcriptional regulation of the rice SUMOylation machinery and into the role of two rice SUMO proteases. BMC Plant Biol 18, 349.

Sakai, H., Lee, S.S., Tanaka, T., Numa, H., Kim, J., Kawahara, Y., Wakimoto, H., Yang, C.-C., Iwamoto, M., Abe, T., et al. (2013). Rice Annotation Project Database (RAP-DB): an integrative and interactive database for rice genomics. - PubMed - NCBI. Plant Cell Physiol. 54, e6–e6.

Sakamoto, T., Sakakibara, H., Kojima, M., Yamamoto, Y., Nagasaki, H., Inukai, Y., Sato, Y., and Matsuoka, M. (2006). Ectopic Expression of KNOTTED1-Like Homeobox Protein Induces Expression of Cytokinin Biosynthesis Genes in Rice. Plant Physiology 142, 54–62.

Sarhadi, E., Bazargani, M.M., Sajise, A.G., Abdolahi, S., Vispo, N.A., Arceta, M., Nejad, G.M., Singh, R.K., and Salekdeh, G.H. (2012). Proteomic analysis of rice anthers under salt stress. Plant Physiology and Biochemistry 58, 280–287.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Srivastava, A.K., Zhang, C., and Sadanandom, A. (2016a). Rice OVERLY TOLERANT TO SALT 1 (OTS1) SUMO protease is a positive regulator of seed germination and root development. Plant Signaling & Behavior 11, e1173301.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Srivastava, A.K., Zhang, C., Caine, R.S., Gray, J., and Sadanandom, A. (2017). Rice SUMO protease Overly Tolerant to Salt 1 targets the transcription factor, OsbZIP23 to promote drought tolerance in rice. The Plant Journal 92, 1031–1043.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Srivastava, A.K., Zhang, C., Yates, G., Bailey, M., brown, A., and Sadanandom, A. (2016b). SUMO is a critical regulator of salt stress responses in rice. Plant Physiology 170, pp.01530.2015–pp.01530.2391.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Su, G., Morris, J.H., Demchak, B., and Bader, G.D. (2014). Biological network exploration with Cytoscape 3. - PubMed - NCBI. Current Protocols in Bioinformatics 47, 8.13.1–8.13.24.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ueguchi-Tanaka, M., Hirano, K., Hasegawa, Y., Kitano, H., and Matsuoka, M. (2008). Release of the repressive activity of rice DELLA protein SLR1 by gibberellin does not require SLR1 degradation in the gid2 mutant. Plant Cell 20, 2437–2446.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ulrich, H.D. (2008). The fast-growing business of SUMO chains. Mol. Cell 32, 301–305.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Van De Velde, K., Ruelens, P., Geuten, K., Rohde, A., and Van Der Straeten, D. (2017). Exploiting DELLA Signaling in Cereals. Trends Plant Sci.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vanstraelen, M., and Benková, E. (2012). Hormonal interactions in the regulation of plant development. Annu. Rev. Cell Dev. Biol. 28, 463–487.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Werner, A., Moutty, M.-C., Möller, U., and Melchior, F. (2009). Performing in vitro sumoylation reactions using recombinant enzymes. Methods Mol. Biol. 497, 187–199.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xue, Y., Zhou, F., Fu, C., Xu, Y., and Yao, X. (2006). SUMOsp: a web server for sumoylation site prediction. Nucleic Acids Research 34, W254–W257.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang, C., Ma, Y., and Li, J. (2016). The rice YABBY4 gene regulates plant growth and development through modulating the gibberellin pathway. J. Exp. Bot. 67, erw319–erw5556.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yano, K., Morinaka, Y., Wang, F., Huang, P., Takehara, S., Hirai, T., Ito, A., Koketsu, E., Kawamura, M., Kotake, K. and Yoshida, S., 2019. GWAS with principal component analysis identifies a gene comprehensively controlling rice architecture. Proceedings of the National Academy of Sciences, 116(42), pp.21262-21267.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yates, G., Srivastava, A.K., and Sadanandom, A. (2016). SUMO proteases: uncovering the roles of deSUMOylation in plants. J. Exp. Bot. 67, 2541–2548.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zentella, R., Hu, J., Hsieh, W.P., Matsumoto, P.A, Dawdy, A, Barnhill, B., Oldenhof, H., Hartweck, L.M., Maitra, S., Thomas, S.G. and Cockrell, S. (2016). O-GlcNAcylation of master growth repressor DELLA by SECRET AGENT modulates multiple signaling pathways in Arabidopsis. Genes & development, 30(2), pp.164-176.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zentella, R., Sui, N., Barnhill, B., Hsieh, W.-P., Hu, J., Shabanowitz, J., Boyce, M., Olszewski, N.E., Zhou, P., Hunt, D.F., et al. (2017). The

Arabidopsis O-fucosyltransferase SPINDLY activates nuclear growth repressor DELLA. Nat. Chem. Biol. 21, 985–485.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title