Title:

SUMOylation of rice DELLA SLR1 modulates transcriptional responses and improves

yield under salt stress

Authors:

Nuno M. Gonçalves[a*], Telma Fernandes[a], Cátia Nunes[b**], Margarida T. G. Rosa[a],

Cleverson C. Matiolli[a], Mafalda A. A. Rodrigues[a], Pedro M. Barros[a], M. Margarida

Oliveira[a], Isabel A. Abreu[a1]

Affiliations:

[a] Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa (ITQB

NOVA), 2780-157, Oeiras, Portugal

[b] Instituto de Biologia Experimental Tecnológica (iBET), 2780-157, Oeiras, Portugal

Author Contributions:

NMG participated in the conception of research, executed most experiments, and

wrote the article with contributions from all authors. TF did protein interaction

experiments. MTGR characterized SUMO-protease rice lines. CCM analysed data

and completed the writing. MAAR purified recombinant proteins. PMB analysed RNA

seq. MMO supervised experiments and analysed results. IAA conceived original

project, supervised experiments, and finalized the article.

Contact:

[1] Corresponding Author:

Isabel A. Abreu

Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa (ITQB

NOVA), 2780-157, Oeiras, Portugal, Tel.: +351 214469647, E-mail: abreu@itqb.unl.pt

* Currently at Leibniz Institute of Plant Biochemistry (IPB), Weinberg 3 | 06120 Halle |

Germany (nunomiguel.goncalves@ipb-halle.de)

** Currently at Instituto Nacional de Investigação Agrária e Veterinária, Avenida da

República, Oeiras, Portugal (catia.soares@iniav.pt)

Short title (50 characters): SUMOylation of rice SLR1 increases salt tolerance

ABSTRACT

DELLA proteins modulate GA signalling and are major regulators of plant plasticity to

endure stress. DELLAs are mostly regulated at the post-translational level, and their

activity relies on the interaction with upstream regulators and transcription factors

(TFs). SUMOylation is a post-translational modification (PTM) capable of changing

protein interaction and found to influence DELLA activity in Arabidopsis. We

determined that SUMOylation of the single rice DELLA SLENDER RICE1 (SLR1)

occurs in a lysine residue different from the one previously identified in Arabidopsis

REPRESSOR OF GA (RGA). Remarkably, artificially increasing SUMOylated SLR1

(SUMO1SLR1) levels attenuated the penalty of salt stress on plant yield. Gene

expression analysis revealed that the overexpression of SUMOylated SLR1 regulates

key dioxygenases that modulate active GA levels, namely *GA20ox2* and *GA2ox3*,

which could partially explain the sustained productivity upon salt stress imposition.

Besides, SLR1 SUMOylation blocked the interaction with the growth regulator YAB4,

which may fine-tune *GA20ox2* expression. Mechanistically, we propose that SLR1

SUMOylation disrupts the interaction with members of several transcription factor

families to modulate gene expression. We found that SLR1 SUMOylation represents

a novel mechanism modulating DELLA activity, which attenuates the impact of stress

on plant performance.

Keywords: DELLA, SUMOylation, stress, productivity, SLR1, YAB4, bZIP23,

gibberellin

One sentence summary: Rice plants show increased yield under salt stress when its

gibberellin transcriptional regulator DELLA protein is artificially SUMOylated.

INTRODUCTION

Salinity is one of the most serious threats to food security, affecting 20% of irrigated soils worldwide. Rice (*Oryza sativa* L.) is a crop with high sensitivity to salinity compared to other monocot crops, such as wheat and barley (Munns and Tester, 2008). Plant sessile nature requires rapid physiological responses to cope with stress, which is partly achieved through a complex crosstalk between hormone signalling pathways affecting growth, flowering, and defence against environmental challenges (Vanstraelen and Benková, 2012). Gibberellins (GA) are diterpenoid hormones involved in the regulation of several developmental processes, including seed germination, stem elongation, and flowering (Hirano et al., 2008). GA responses are modulated by DELLA proteins, which act as transcriptional regulators (Hedden and Thomas, 2012). In the presence of GA, DELLAs are targeted to ubiquitin-mediated proteasome degradation by the GA-receptor GA-INSENSITIVE DWARF1 (GID1) through binding to specific DELLA domains (DELLA/TVHYNP), followed by DELLA polyubiquitination mediated by the E3-ligase GA-INSENSITIVE DWARF2/SLEEPY1 (GID2/SLY1). When GA levels decrease, DELLA proteins accumulate and interact with key transcriptional regulators to restrain growth (Hirano et al., 2010; 2008; Ueguchi-Tanaka et al., 2008). For instance, DELLAs can sequester the transcription factors PHYTOCHROME INTERACTING FACTOR3 and 4 (PIF3 and PIF4), in addition to negatively regulate PIF3 protein abundance, mediating the crosstalk of light and GA signalling to modulate cell growth (de Lucas et al., 2008; Li et al., 2016). DELLA transcriptional regulation of GA biosynthetic genes responsible for hormonal homeostasis also potentiates growth arrest. The single rice DELLA SLENDER RICE1 (SLR1) interacts with YAB4 transcription factor to prevent the induction of *GA20ox2*, which encodes a dioxygenase that is essential for the synthesis of bioactive GA (Yang et al., 2016). In Arabidopsis, DELLA interaction with GAI-ASSOCIATED FACTOR1 (GAF1) TF triggers a negative-feedback loop that activates GA biosynthesis and subsequent DELLA degradation (Fukazawa et al., 2017). The increasing number of validated DELLA interactors showcase an overlap with most hormone pathways, impacting development, fertility, and seed number (Davière and Achard, 2015; Van De Velde et al., 2017).

DELLA activity is mostly modulated through post-translational modifications (PTMs), which impact its ability to interact with transcriptional regulators and modulate gene expression (Van De Velde et al., 2017). Besides ubiquitination and subsequent

proteasome-mediated degradation, there are other PTMs that influence the activity of the DELLA's in response to GA signalling. For instance, the phosphorylation by kinase EARLY FLOWERING 1 (EL1) regulates SLR1 protein stability and activity (Dai and Xue, 2010; Itoh et al., 2005). Recently, a mechanism of antagonistic PTMs was proposed to regulate Arabidopsis DELLA's interaction with specific protein partners. *O*-fucosylation by SPINDLY (SPY) promotes DELLA interaction with essential growth regulators – such as BRASSINOLE-RESISTANT1 (BZR1), PIF3 and PIF4 – leading to growth arrest (Zentella et al., 2017). Conversely, RGA *O*-GlcNAcylation by SECRET AGENT (SEC) blocks these interactions and promotes GA-induced growth resumption (Camut et al., 2017; Zentella et al., 2016). DELLA SUMOylation has also been indicated as a PTM controlling the stability of REPRESSOR OF GA (RGA), one of the Arabidopsis five DELLA proteins. SUMO attachment to RGA hinders its ubiquitin-mediated proteasome degradation through blockage of GID1-RGA interaction in a GA-independent manner (Conti et al., 2014). The stabilization of Arabidopsis DELLA by SUMOylation also positively affected salt stress response, in line with observations that DELLA accumulation leads to increased salinity tolerance (Achard et al., 2006; Conti et al., 2014; Nelis et al., 2015). SUMOylation is a eukaryotic ubiquitination-like process that involves the processing, activation, conjugation, and ligation of mature SMALL UBIQUITIN-LIKE MODIFIER (SUMO) to acceptor lysine's (K) in the target protein (Miura et al., 2007). Unlike ubiquitination, SUMO-modifications often occur in a canonical consensus motif (ψ-$\underline{K}$-X-E/D), which can be used for *in silico* prediction of SUMOylation targets (Xue et al., 2006). SUMOylation regulates protein stability, activity, and interactions with specific protein partners (Cremona et al., 2012). SUMO deconjugation (deSUMOylation) from target proteins is performed by deSUMOylating proteases (DSPs), with some members recently reported to be specific for poly-SUMO chain processing (Eckhoff and Dohmen, 2015; Yates et al., 2016; Augustine et al., 2018). In rice, reduced levels of SUMO conjugates due to DSP activity of OVERTOLERANT TO SALT1/2 (OTS1/OTS2) have been connected to increased salt tolerance (Srivastava et al., 2016a; 2016b). SUMOylation of specific TFs, such as the stress-induced basic LEUCINE ZIPPER 23 (bZIP23), was suggested to be crucial to drought tolerance and fertility (Srivastava et al., 2017).

Few studies have addressed the biological function of SUMOylation of stress-induced targets. Moreover, most of the studies investigating the effects of PTMs on

DELLA function were done in Arabidopsis. This dicot model plant encodes five
DELLAs in the genome and essentially differs from monocots, which harbour a single
DELLA gene. This study demonstrates that SUMOylation of the single rice DELLA
SLR1 impacts the transcriptional output in response to salt stress, possibly through
modulation of SLR1 interaction with transcription factors regulating transcriptional
patterns governing plant growth and stress responses. We suggest that SLR1
SUMOylation affects its interaction with key transcription factors that impacts GA
homeostasis (YAB4), ABA signalling (bZIP23), and JA signalling (bHLH089/94).
Additionally, the evidence provided suggests that it is possible to uncouple stress
response from decreased plant performance by manipulating the levels of
SUMOylated SLR1.
138
RESULTS
SLR1 SUMOylation *in vitro* analysis suggests diversification of DELLA
regulation by SUMO in plants
SUMOylation of lysine 65 (K65) in Arabidopsis DELLA RGA modulates its
stability in a GA-independent manner (Conti et al., 2014). To determine rice SLR1
SUMOylation, we started by performing an *in silico* prediction of SUMOylation motifs
using SUMOsp2 software (Xue et al., 2006) in rice DELLA SLR1 and Arabidopsis
DELLA RGA protein sequences. We sought for putative conserved SUMOylation
motifs in both proteins and found a low probability SUMOylation site which
corresponds to Arabidopsis RGA (Figure 1A). The SUMO acceptor lysine (K65) in
RGA is part of a canonical SUMOylation motif (LKLE). In contrast, its counterpart in
SLR1(K60) is not part of a canonical motif (QKLE), nor are the equivalent ones in
another Arabidopsis or monocot DELLAs (Supplemental Figure S1). Nevertheless,
both SLR1 and RGA have a high probability SUMOylation site on lysine 2 (K2).
To verify if the *in silico* prediction of SUMO attachment to SLR1 holds true, we
performed an *in vitro* SUMOylation assay targeting both full-length SLR1 and a
truncated isoform containing the leading 190 amino acids (SLR1t). Importantly, SLR1t
includes the N-terminal domains (DELLA and TVHYNP) responsible for GA-mediated
modulation of DELLA stability and SLR1 transactivation properties (Davière and
Achard, 2015; Marín-de la Rosa et al., 2015; Hirano et al., 2012), as well as the
predicted SUMOylation sites depicted in the Figure 1A. We observed a mobility-shift
of both full-length SLR1 (Figure 1B) and truncated SLR1t (Figure 1C; with both

SUMO1 and SUMO2 - Supplemental Figure S2A-B), consistent with ATP-dependent

SUMO-conjugation in the presence of the rice SUMOylation machinery. The band

resulting from *in vitro* SLR1 and SLR1t SUMOylation disappeared upon treatment with

recombinant SUMO-specific protease ULP1 that cleaves SUMO from its conjugates

(Li and Hochstrasser, 2003), confirming that the observed mobility-shift was caused

by attached SUMO (Figure 1B-C). To analyse the contribution of each predicted

SUMO acceptor lysine in SLR1, we performed directed mutagenesis changing the

lysines K2 and K60 to arginine residues, which cannot be conjugated to SUMO (K2R

and K60R). The mutated SLR1t K60R isoform retained the ability to be SUMOylated

– which was confirmed by the band disappearance after SUMO proteolytic removal by

ULP1 treatment – while K2R did not (Figure 1C). The result shows that K2 is the

preferential SUMOylated residue in SLR1, suggesting that SUMO attachment to

DELLA may occurs in distinct lysine residues of Arabidopsis RGA (K65) and rice SLR1

(K2). SUMOylation in different sites potentially results in different outcomes regarding

protein interactions and activity of SUMOylated SLR1 compared to SUMOylated RGA.

However, the *in silico* prediction of SUMOylation sites suggests that RGA K2 could

also be SUMOylated (Figure 1A). We need further studies to clarify if K2 SUMOylation

occurs in Arabidopsis RGA, or even in DELLA proteins from other species.

SUMO protease activity analyses provide evidence of *in vivo* SLR1 SUMOylation

To demonstrate the *in vivo* SUMOylation of SLR1, we treated non-denatured

rice protein extracts with the ULP1 SUMO protease. The procedure would allow for

indirect detection of SUMOylated SLR1 by analysing the increase in non-SUMOylated

SLR1 protein amount resulting from ULP1 treatment, in comparison to SUMOylated

SLR1. We observed an increase of non-SUMOylated SLR1 upon the addition of ULP1,

as detected by the anti-SLR1 antibody in protein extracts (Figure 2A). The result is in

line with the previous *in silico* and *in vitro* analyses indicating that SLR1 could be

SUMOylated. Our results also suggest that SLR1 mono-SUMOylation may be much

less abundant than other forms of SUMOylation since we could not detect an SLR1

band with the expected mobility shift for mono-SUMOylation (Figure 1B-C). To further

support the evidence of *in vivo* SLR1 SUMOylation, we tested the levels of

SUMOylated SLR1 in knockout mutant lines for genes encoding rice SUMO

proteases, namely *els1* and *fug1* (Figure 2B-C). ELS1 and FUG1 are active rice SUMO

proteases, as both knockout lines contain increased levels of SUMOylated proteins

(Rosa et al., 2018). We treated protein extracts from *els1* and *fug1* homozygous lines
with ULP1 and compared their total SUMOylated SLR1 levels with those present in
negative segregant lines (NS). The results show that although the overall level of non-
SUMOylated SLR1 is not significantly affected in either one of the KO lines when
compared to their negative segregant lines (Figure 2B), the levels of SUMOylated
SLR1 are increased in *fug1* mutant (Figure 2C). These results demonstrate
SUMOylation of SLR1 occurs *in vivo*, as indicated by the processing (*i.e.,* switch from
SUMO-SLR1 to SLR1) mediated by FUG1. Interestingly, it also suggests that there is
a certain degree of specificity for SLR1 deSUMOylation, as suggested by SUMO
cleavage by FUG1 and not ELS1 (Figure 2C). Altogether, the evidence supports that
SLR1 is SUMOylated *in planta* and that SUMO specific proteases could regulate the
levels of SUMOylated SLR1.
207
Overexpression of SUMOylated SLR1 increases salt tolerance
Because DELLA protein accumulation was previously connected with stress
tolerance (Achard et al., 2006), we assessed SLR1 expression in response to salt
stress. Wild-type seedlings accumulated SLR1 protein in shoots during the first 8
hours of stress (Supplemental Figure S3A). On the other hand, *SLR1* transcript levels
decreased, which is consistent with the previously reported DELLA negative feedback-
loop regulation on its own transcript abundance (Hedden & Thomas, 2012).
To gain insight on the role of SLR1 SUMOylation in rice development and
responses to salt stress, we generated transgenic lines overexpressing SLR1 (SLR1-
OX), as well as lines constitutively expressing a mimicry of the SUMOylated form of
SRL1 (SUMO1SLR1-OX) originated from the genetic fusion of mature SUMO1 to the
K2 residue, which is the SUMOylated lysine of SLR1 ($SUMO1_{GG}::SLR1_{K2}$) (Figure 3A).
The strategy of using SUMOylated mimetic isoforms originated by gene fusion with
the SUMO coding sequence has been used in several biological systems, including
plants (Crozet et al., 2015; Ulrich, 2008). We confirmed the overexpression of non-
SUMOylated SLR1 isoforms in both SLR1-OX and SUMO1SLR1, with the latter also
showing a band with the expected mobility-shift of SLR1 corresponding to mono-
SUMOylation (Figure 3B). We also observed an increase in the amount of non-
SUMOylated SLR1 protein in SUMO1SLR1-OX compared to WT and NS lines, which
could be arising from the processing by endogenous SUMO proteases because the
fusion protein retains the SUMO diglycine protease cleavage motif (-GGK-).

Rice is particularly sensitive to salt at both seedling and booting stages
(Gregorio et al., 1997). Therefore, we analysed stress marker symptoms of SLR1-OX
and SUMO1SLR1-OX lines subjected to high salt concentrations at seedling stage, as
well as yield parameters on salt treatments at booting stage. Evaluation of symptoms
in seedlings subjected to severe salinity (120mM NaCl) for 8 days showed that both
SUMO1SLR1-OX lines performed better under salt stress, displaying decreased
symptoms severity according to SES score (Standard Evaluation System for Rice) by
IRRI (International Rice Research Institute) (Gregorio et al., 1997), including minimal
leaf tip rolling and increased survival (Figure 3C). Importantly, salt tolerance could be
correlated with SUMOylated SLR1 levels, as SUMO1SLR1-OX #1 – showing higher
SLR1 SUMOylation levels compared to SUMO1SLR1-OX #2 – performed better than
the latter (Figure 3B-C). Interestingly, salt-induced SUMOylated SLR1 increased
alongside with the non-SUMOylated SLR1 in SUMO1SLR1-OX lines (Supplemental
Figure S3B), suggesting that SUMO cleavage processing could be occurring
alongside the duration of salt stress. Plants overexpressing SUMO1SLR1 and SLR1-
OX also recovered better than the wild type Nipponbare in the 8 following days to salt
stress cessation. Thus, mild overexpression of SLR1 can increase rice tolerance to
salinity, and this effect is increased by its SUMOylation.
We next tested rice performance under salt stress at the booting stage. Panicle
initiation was the cue for mild salt stress imposition (80 mM NaCl), as at this
developmental stage pollen viability is most affected by high salinity (Negrão et al.,
2011; Sarhadi et al., 2012). Upon measurement of agronomical parameters, we did
not detect stress-induced changes in plant height (Figure 4A), tillering or panicle length
(Supplemental Figure S4) when compared to the control. However, two critically
important parameters for rice yield – namely fertility (filled grains) and productivity
(grain weight) – were significantly improved in SUMO1SLR1-OX lines under mild salt
stress. Both SLR1-OX and Nipponbare presented around 10% decrease in fertility
when comparing to mock conditions, while SUMO1SLR1-OX showed the same fertility
in both salt and mock treatments (Figure 4B). This tendency was also observed in
grain weight, where SUMO1SLR1-OX presented higher tolerance to mild salt stress
regarding productivity compared to Nipponbare and SLR1-OX (Figure 4C). Although
the SUMO1SLR1-OX lines had lower productivity under mock conditions compared to
Nipponbare, they showed improved performance under salt stress. These results

substantiate the significance of SUMOylation of SLR1 in salt stress tolerance
acquisition and yield protection observed in two sensitive rice developmental stages.
264

SLR1 SUMOylation regulates GA biosynthetic pathway transcripts in response
to salt stress

The salt tolerance observed in SUMO1SLR1-OX lines could not be attributed
to the increased accumulation of SLR1, as SLR1 protein levels were similar in the
SLR1-OX and SUMO1SLR1-OX transgenic lines (Figure 3B). Thus, we inferred that
the observed phenotypes in SUMO1SLR1-OX lines could be a consequence of the
increased levels of SUMOylated SLR1. Because DELLA proteins regulate
transcription, we performed a genome-wide expression analysis to identify
differentially regulated genes in SLR1-OX and SUMO1SLR1-OX that could explain
the phenotypical differences observed in these lines in response to salt stress. We
exposed SLR1-OX, SUMO1SLR1-OX, and Nipponbare genetic background to high
salt concentrations (120 mM NaCl) during 8 hours at the seedling stage (2 weeks old),
and collected samples for subsequent RNA-seq analysis (Supplemental Table S1).
We selected the lines SLR1-OX #2 and SUMO1SLR1-OX #2 for the RNA-seq assay
because they showed similar non-SUMOylated SLR1 levels, as we aimed to tackle
the effect of SLR1 SUMOylation on gene expression. Both overexpressor lines
showed an approximately 1.8-fold increase in SLR1 transcript levels when compared
to Nipponbare, which is in line with the levels validated by real-time qPCR for the same
samples, *ca.* 1.5-fold increase (Supplemental Figure S5), and with protein levels
assessed by immunoblotting (Figure 3B). The induction of salt-induced genes *Rab21*
and *P5CS* was used as a transcriptional readout of the salt stress treatment efficiency
(Supplemental Figure S5).

Principal Component Analysis (PCA) revealed a clear separation of samples by
treatment, this factor being responsible for 47% of the variability in gene expression,
while genotype-specific transcriptomic changes explained 21% of the observed
variability (Supplemental Figure S6). The constitutive overexpression of SLR1 and
SUMO1SLR1 in absence of salt stress resulted in differential regulation of 247 and
207 genes, respectively (Supplemental Tables S1.6 to S1.8). Functional annotation of
SLR1-OX DEGs showed enrichment of gene ontology (GO) terms associated with
diterpenoid biosynthesis, highlighting the impact of SLR1 on GA metabolism
(Supplemental Figure 7A and Supplemental Tables S1.6 to S1.7). Notably, we

observed overall downregulation in genes coding for enzymes essential in non-
gibberellin diterpenoid synthesis, namely syn-copalyl diphosphate synthase (*CPS4*)
and ent-cassa-12,15-diene synthase (*KSL7*) (Supplemental Figures S7A and S7C).
These enzymes are part of a diterpenoid pathway branching point that feeds the
phytoalexins biosynthetic pathway (Otomo et al., 2004). Concurrently, other genes
related to phytoalexin and phytocassane synthesis, such as cytochrome P450 family
members *CYP701A8/KO1, CYP99A3, CYP71Z6, CYP76M8,* and *CYP71Z7*, were
also downregulated in SLR1-OX.

Phenotypical discrimination of SLR1-OX and SUMO1SLR1-OX was observed
under salt stress but not under control conditions (Figure 3C). Interestingly, a closer
look into the distribution of SLR1-OX and SUMO1SLR1-OX samples in the PCA, under
control and stress conditions highlights a similar tendency (Supplemental Figure S6).
Hence, we focused on the analysis on the 236 DEGs specific to SUMO1SLR1-OX
exposed to salt stress. After functional annotation of DEGs specific to SUMO1SLR1-
OX, we found a significant over-representation of GO terms associated with defence
response and gibberellin dioxygenase activity (Supplemental Figures S7B and S7C).
We also found a salt stress-dependent induction of *GA20ox2* and *GA2ox3* genes in
SUMO1SLR1-OX (Supplemental Table S1.4). *GA20ox2* and *GA2ox3* encode
enzymes involved in the regulation of active GA (GA_4) levels. Therefore, it is
reasonable to hypothesize that SLR1 SUMOylation could adjust active GA
homeostasis upon exposure to salt stress (Supplemental Figure S7C).

318 SUMOylation disrupts SLR1 interaction with stress-associated transcription
319 factors

The increased salt tolerance and sustained productivity of SUMO1SLR1-OX
under salt stress likely arise from the presence of increased levels of SUMOylated
SLR1 (Figure 3B-C; Figure 4A-B), which seems to be affecting the gene expression
profile of plants exposed to stress. The *GA20ox2* transcript was induced around 2-fold
in SUMO1SLR1-OX exposed to salt stress, while it showed only a marginal induction
in SLR1-OX under the same conditions (Supplemental Table S1.2). One possible
mechanism by which SLR1 could affect *GA20ox2* transcription is through the
interaction with the transcription factor YAB4, thereby repressing the expression of
GA20ox2 (Yang et al., 2016). We did not detect *YAB4* transcript changes after 8 hours
of salt stress exposure, but a search in public transcriptome data revealed that *YAB4*

is induced in rice shoots after 3h of salt stress imposition (Genevestigator - Hruz et al.,

2008; Supplemental Figure S9A-B), which could lead to increased YAB4 protein

accumulation during later stages of plant adaptation to salt stress. In an attempt to

dissect the molecular pathways involved in *GA20ox2* upregulation in SUMO1SLR1-

OX, we verified the interaction of SLR1 and SUMOylated SLR1 with YAB4 TF using a

yeast two-hybrid system and bimolecular fluorescence complementation (BiFC). The

aim was to test the effect of SUMOylation on SLR1-YAB4 interaction, which could

affect *GA20ox2* transcription. We used a variant of the SUMO1SLR1 fusion protein

lacking the SUMO protease recognition domain to avoid possible processing by yeast

SUMO proteases (SUMO1$_{(GG-AA)}$::SLR1). We confirmed the previously demonstrated

interaction between SLR1 and YAB4 (Yang et al., 2016), and also found that the

mimicking SUMOylation in lysine 2 of SLR1 disrupts SLR1-YAB4 interaction in both

yeast and *in planta* (Figure 5A-B; negative controls in Supplementary Figure S8). The

blocking of the SLR1 interaction with YAB4 by SUMO correlates with the induction of

GA20ox2 in salt-stressed SUMO1SLR1-OX lines.

To further evaluate the molecular aspects of SUMO1SLR1-OX tolerance to salt

stress, we explored the notion that SLR1 exerts its function mainly by transcriptional

regulation through interaction with transcription factors. We designed a strategy that

allowed us to identify SLR1 TF interactors that could explain, at least partially, the

differences in gene expression observed in the transgenic lines upon exposure to salt

stress. Briefly, we submitted the promoter sequences (1000 bp upstream to the

Transcription Start Site – TSS) of 417 selected DEGs (SLR1- and SUMO1SLR1-

dependent salt stress responsive genes; Supplemental Tables S1.3 and S1.4) to an

in silico search for transcription factor binding sites (TFBS) in the PlantPAN online

database (Chow et al., 2016). Next, we selected only TFBS and corresponding binding

TFs that have been experimentally validated. This allowed us to identify TFs from the

TCP, bZIP, bHLH, and B3 families that could be regulating the output gene expression

of SLR1-OX and SUMO1SLR1-OX transgenic lines under salt stress (Table 1). To

cross-validate the TF identification, we made sure that a substantial overlap between

the known target genes of those TFs and the SLR1- and SUMO1SLR1-dependent

DEGs were observed (Supplemental Tables S1 and S2).

The interaction between SLR1 with the selected candidates (Table 1) and its

regulation by SUMOylation was assessed through Y2H assays and BiFC analyses

(Figure 5A-B). The basic helix-loop-helix transcription factors bHLH089 and bHLH094,

as well as the homeobox TF OSH1, showed a strong interaction with SLR1 but not
with SUMO1$_{(GG-AA)}$::SLR1. This finding suggests that SUMOylation could modulate
the binding of SLR1 to these transcription factors, potentially affecting their
transcriptional activities. We also found that bZIP23 can interact with both
SUMOylated and non-SUMOylated SLR1 (Figure 5A). The interaction of SLR1 with
bHLH089 and bHLH094, as well as its disruption by SLR1 SUMOylation, was validated
through BiFC in *Nicotiana benthamiana* leaves, supporting the *in planta* SUMO-
dependent regulation of these interactions (Figure 5B). Neither IDEF1 nor PCF1
showed interactions with SLR1 or SUMO1$_{(GG-AA)}$::SLR1.
373
DISCUSSION
DELLA proteins are central regulators of plant productivity and stress
responses, and the hormonal crosstalk mediated by these proteins is essential for
plant plasticity to endure stress. As other hormone modulators, DELLA activity is
regulated by post-translational modifications, such as ubiquitination, phosphorylation,
O-fucosylation, *O*-GlcNAcylation, and SUMOylation (Yano et al., 2019; Zentella et al.,
2016; Zentella et al., 2017; Conti et al., 2014; Dai and Xue, 2010; Itoh et al., 2003).
Recently, SUMO proteases were linked to salt stress response in rice (Srivastava et
al., 2016b; 2017). Although deSUMOylation was proposed as a general mechanism
of tolerance acquisition in response to salt, we show that artificially SUMOylated SLR1
could alleviate the salt stress impact on rice productivity, also improving recovery after
stress imposition is released.
Following the evidence that SUMO proteases are important for DELLA activity
(Campanaro et al., 2016), the knockout mutant of SUMO protease *fug1* showed a
higher ratio of overall SUMOylated SLR1. Interestingly, the FUG subclass is
phylogenetically close to the SPF subclass, which contains Ulp2-like SUMO proteases
(Castro et al., 2018). A Ulp2 from *Saccharomyces cerevisiae* (ScUlp2) was previously
reported to be the key protease responsible for poly-SUMO chain processing (Eckhoff
and Dohmen, 2015). If FUG1 is indeed related to Ulp2 proteases and it processes
SUMO-SLR1, SLR1 deSUMOylation may be occurring in poly-SUMO chain high
molecular weight conjugates.
DELLA proteins control several plant processes through interaction with a
myriad of transcriptional factors (Van De Velde et al., 2017). By affecting protein-
protein interaction, SLR1 SUMOylation has the potential to modulate plant processes

by altering the transcriptional outcome of SLR1-TFs interacting proteins. Indeed,

transcriptomic analysis of SUMOylated SLR1 overexpressing lines suggested that this

isoform plays a role in early transcriptional responses to salt stress, affecting the

expression of key genes coding for enzymes controlling the accumulation of active

GA, such as *GA20ox2* and *GA2ox3*, involved in activation and inactivation of the active

GA_4. We were able to determine that SLR1 SUMOylation occurs in lysine 2 (Figure

1A-C), distant from the conserved GA regulatory domains DELLA and TVHYNP. This

finding suggests that SLR1 SUMOylation contrasts with Arabidopsis RGA, in which

the amino acid residue responsible for SUMO modification is the lysine 65 (Conti et

al., 2014), located very close to the GA-regulatory domain TVHYNP. The location of

SUMO attachment in SLR1 protein could exert quite different effects on protein-protein

interactions compared to its well-described Arabidopsis counterpart RGA. An

interesting possibility is that SUMOylation of these residues could have a different

outcome regarding the transcriptional activation properties of SLR1 DELLA/TVHYNP

motifs (Hirano et al., 2012). Both lysines are within SUMOylation canonical motifs, but

while the SUMOylation canonical motif surrounding K2 is conserved in all analysed

DELLAs, the one containing K65 is exclusive to RGA. The different SUMOylation sites

of SLR1 and RGA might account for alterations in DELLA regulation amongst species

that probably determine different interactome networks.

The SUMOylation of SLR1 could also be affecting the interaction with upstream

regulatory proteins. For instance, SUMO attachment could affect SLR1 interaction with

SPY, which enhances SLR1 activity through *O*-fucosylation and is a major determinant

of rice plant architecture (Yano et al., 2019). Crosstalk between SUMOylation and *O*-

fucosylation could play a role in GA-promoted growth regulation by decreasing TF-

interactions that result in growth repression (Camut et al., 2017). This control of the

expression of GA enzymes is demonstrated by our transcriptomic results of

SUMO1SLR1-OX lines under salt stress, which shows upregulation of both *GA20ox2*

and *GA2ox3*, suggesting that SLR1 SUMOylation could also regulate GA

homeostasis. The induction of genes involved in both anabolism and catabolism of

bioactive GA, catalized by GA20ox2 and GA2ox3, respectivelly, seems paradoxical.

However, each gene could be activated in different tissues under salt stress,

determining tissue specific expression and regulating plant plasticity. These

differences in spatial expression are lost in whole plant transcriptomic analysis.

Further experiments evaluating *GA20ox2* and *GA2ox3* spatial expression under salt

stress have to be done to address the issue.

SUMOylation of specific targets was suggested to be crucial for stress

tolerance, fertility and yield maintenance in rice. For instance, bZIP23 SUMOylation

levels are modulated by OTS1, which affects its stability and the plant's ability to deal

with drought and salt stress (Srivastava et al., 2017). Interestingly, we found that

bZIP23 interacts with both SLR1 and SUMO1SLR1. Another novel interaction with

both bHLH089 and bHLH094 shows upregulation of JAZ-like repressors like JAZ8,

consistent with findings that this TF is regulating genes from the JA pathway as

observed in our transcriptomic analysis. Both these interactions may partly explain the

JA-induced stress response promoted by SLR1 accumulation (Supplemental Figures

S7A and S7B).

It was already shown that YAB4 interacts with SLR1 to repress *GA20ox2* gene

expression and restrain development in rice (Yang et al., 2016). We confirmed the

interaction between SLR1 and YAB4 by Y2H assays and BiFC (Figure 5B-C). We

additionally show that SUMOylation of SLR1 abolished its interaction with YAB4.

Accordingly, we found a stress-induced upregulation of *GA20ox2* in SUMO1SLR1-OX

(Supplemental Table S1). Moreover, *GA20ox2* was also found to be downregulated

by KNOTTED1-like homeobox (KNOX) proteins like OSH1 (Sakamoto et al., 2006).

SLR1 SUMOylation also disrupted its interaction with OSH1, suggesting that SLR1

SUMOylation mechanistic action is partly achieved through SLR1 dissociation from

transcriptional regulators involved in GA signalling and stress signalling. It could

represent a dissociation of DELLA function during salt stress tolerance from its impact

on productivity by means of SUMOylation.

The emergence of SUMOylation of SLR1 as an effector of rice tolerance to salt

stress, through the growth-maintaining modulation of protein-protein interaction with

transcriptional regulators such as YAB4, is evidence of the tight regulation of the GA

pathway by rice DELLA during stress. Recent studies have demonstrated that DELLA

proteins are targets of different post-translational modifications, which could adjust

their interactions with transcriptional regulators in a dynamic fashion, ensuring the best

developmental decisions when facing stress. Further studies need to be performed to

unveil upstream effectors that perceive stress and the underlying signal transduction

pathway that leads to SLR1 SUMOylation. By establishing the trigger for SLR1

SUMOylation and its effect on stress-induced SLR1 protein-protein interactions, new

strategies can be developed involving the transient post-translational modification of
DELLA, as mutagenesis and selection of growth-favourable SLR1 alleles. This can be
used to activate the SUMO-regulated tolerance mechanism when stress is
forthcoming and protect crops from yield losses due to abiotic stress.
469
METHODS
SLR1 genomic DNA cloning and vectors construction
The intronless gene SLR1 entire coding sequence was amplified from Nipponbare
wild-type genomic DNA, using the primers SLR1-CDS-Fw and SLR1-CDS-Rv
(Supplemental Table S4). The fragment was cloned into the pJET 1.2/blunt-end
cloning vector (ThermoScientific), and the SLR1 full sequence confirmed by Sanger
sequencing. SLR1 CDS was then amplified from pJET-SLR1 using the primers pET-
Fw-*Eco*RI and pET-Rv-*Psp*OMI (Supplemental Table S4) to insert the restriction sites
for *Eco*RI and *Psp*OMI to clone SLR1 in the pET28a expression vector (Novagen). For
the generation of rice transgenic lines overexpressing SLR1, the SLR1 coding
sequence was amplified from pJET-SLR1 using OsSLR1-GW-Fw and OsSLR1-GW-
Fw primers to insert the attB1 and attB2 recombination sites. The amplified PCR-
product was recombined into the pDONR221 (ThermoFisher).The SUMO1SLR1 gene
fusion was generated using high-fidelity PCR (Phusion DNA Polymerase; NEB)
essentially as described in Atanassov et al. (2009). Primers overlapping both the
OsSUMO1 3'-end and OsSLR1 5'-end OL-SLR1+SUMO1-3'-Fw and OL
SUMO1+SLR1-5'-Rv, were used in combination with OsSLR1-Gw-Rv and OsSUMO1-
GW-Fw, respectively, to generate the fragments to be fused. In the next step, the PCR
products were purified and used in the fusion PCR reaction containing the OsSUMO1-
GW-Fw and OsSLR1-Gw-Rv primers for the final amplification of the final fusion
product (Supplemental Table S4). The gene fusions were confirmed by Sanger
sequencing. To obtain rice transgenic plant lines overexpressing SLR1 (SLR1-OX)
and SLR1 fused to SUMO1 (SUMO1SLR1-OX), pDONR221 containing the inserts
described above was used as entry vector for the pANIC6B destination vector (Mann
et al., 2011). The expression of SLR1 and SUMO1SLR1 is controlled by the maize
ZmUbi1 promoter (Figure 3A).
496
Site-directed mutagenesis

For the production of truncated recombinant SLR1 protein (SLR1t) containing the
leading 190 amino acid residues of SLR1, a stop codon was introduced by site-
directed mutagenesis (SDM) in pET28a-SLR1 using *Pfu*Turbo DNA polymerase
(Agilent) and the primers (SLR1t-STOP-Fw and SLR1t-STOP-Rv) designed with
QuickChangeII online tool (Agilent) (Supplemental Table S4). Unmodified plasmids
were digested with methylation-sensitive enzyme *Dpn*I (New England Biolabs). The
mutated plasmids were transformed in *E. coli* DH5α and the mutagenized site
confirmed by sequencing. Mutated pET28a coding the truncated SLR1 isoform
(SLR1t) was subjected to further mutagenesis procedures to perform the amino acid
substitutions K2R and K60R (primers SLR1t-K2R-Fw, SLR1t-K2R-Rv, SLR1t-K60R-
Fw and SLR1t-K60R-Rv, all listed in Supplemental Table S4) following the same
protocol.
510

Recombinant Protein Production and Purification
pET28a vectors containing the full-length SLR1, N-terminus truncated SLR1 (SLR1t),
SLR1t (K2R) and SLR1t (K60R) were transformed into *E. coli* strain BL21 Star (DE3,
Sigma-Aldrich) for recombinant protein expression. Cells were grown at 37ºC until
OD_{600} = 0.6, and protein production was induced by addition of 200 µM isopropyl-d-1-
thiogalactopyranoside (IPTG). After three hours, the cells were harvested by
centrifugation in a J2MI centrifuge (Beckman) at 7000 RPM during 20 minutes at 4ºC.
The cells were resuspended in lysis buffer (phosphate buffer 20 mM pH 7.5, 500 mM
NaCl, 10 mM Imidazole) containing 5 µg/mL of DNAse, 50 µg/mL of lysozyme, PMSF
100 mM and $MgCl_2$ 25 mM. After freezing at -20ºC and thawing, extracts were loaded
into a HisTrap chelating column (Amersham Biosciences) and proteins eluted in
phosphate buffer containing 500 mM imidazole, following buffer exchange dialysis
(phosphate buffer 20 mM, NaCl 150 mM, DTT 1 M, glycerol 2%), for increased protein
stability. Recombinant proteins were analysed by Western Blot using custom-made
SLR1 specific polyclonal antibody with Mini-Protean II system (Biorad).
526

***In vitro* SUMOylation assays and ULP1 treatment**
The *in vitro* SUMOylation assays were performed in 20 µL of SUMOylation reaction
buffer (Werner et al., 2009) containing 3 µg of SLR1 recombinant protein, 1 µg of E1-
activating enzyme (OsSAE1/ OsSAE2), 0.8 µg of E2-conjugating enzyme (OsSCE1a),

0.5 µg of OsSUMO1/2, 0.4 units of creatine phosphokinase, and 10 mM creatine

phosphate. Reactions were initiated by adding 1 mM ATP and then incubated at 37ºC

for 1 h to 3 h. To confirm SUMOylation specificity, 3 µg of purified SUMO protease

ULP1 from *Saccharomyces cerevisiae* (Li and Hochstrasser, 2003) was used for

deSUMOylation analysis. SUMO conjugates were analysed by Western Blot with

custom anti-SLR1 antibody. For ULP1 treatments, whole plants were ground in liquid

N_2 and 100 mg of ground tissue powder were mixed with non-denaturing buffer (Tris-

HCl pH 8.0 50 mM, sodium chloride 150 mM, NP-40 1%, sodium deoxycholate 0.5%,

SDS 0.1%, EDTA pH 8.0 1 mM, NEM 50 mM), followed by homogenisation by

vortexing, and subsequent centrifugation at 4 ºC for 20 minutes to recover the

supernatant containing the proteins. The non-denaturing protein extracts were treated

with 7 µg of ULP1, incubated at 37 ºC with collection points taken at 5, 30, 60, and

120 seconds. Proteins were denatured at 90ºC for 10 minutes with Laemmli Buffer

with 5 mM DTT and anti-SLR1 immunoblot proceeded as described above.

SLR1 protein extraction, detection and quantification analysis

Plant shoot samples were ground in liquid N_2 and 100 mg of ground tissue used for

protein extraction with denaturing TCA/Acetone precipitation, followed by

resuspension in lysis buffer (7 M Urea, 2 M Thiourea, 30 mM Tris, 4% (w/v) CHAPS,

4 % (v/v), cOmplete protease inhibitor cocktail EDTA-Free (Sigma-Aldrich), 0.1% (v/v)

Nuclease Mix 100x, and NEM 20 mM. Protein quantification was performed with 2-D

Quant kit (GE Healthcare) following manufacturer's instructions. Anti-SLR1

immunoblots were performed in a Mini-Protean II System (Biorad), using custom-

made anti-SLR1 polyclonal specific antibody [1:5.000, overnight incubation] and anti-

rabbit (Novex) [1:20.000, 90 min incubation] in 5% non-fat dry milk in TBS 1X.

Quantification of native SLR1 bands provided by immunoblotting analysis, normalized

to protein loading control using stripped membrane stained with Coomassie G-250

(Fluka), was performed with ImageJ software as indicated in the official ImageJ

website online tutorial.

Plant material and Rice transformation

Calluses induced from wild-type *Oryza sativa* (japonica spp.) cv. Nipponbare mature

seeds were used for rice transformations according to (Almadanim et al., 2017), and

selection for transformants asserted by hygromycin resistance and standard GUS-

staining with X-Gluc solution (Roth) of a section of seminal roots. Proteins were
extracted, detected and quantified using a modified TCA/Acetone method (Luís et al.,
2016). Due to growth arrest phenotype and fertility decrease in homozygous lines of
SUMO1SLR1-OX lines, all assays continued with positively-selected heterozygous
plants. Moreover, lines #2 for both SUMO1SLR1-OX and SLR1-OX were selected for
further analysis at rice booting stage and transcriptomic evaluation according to non-
SUMOylated SLR1 levels, in order to avoid DELLA pleiotropic effects (Davière and
Achard, 2015). Wild-type Nipponbare was used as a control for SLR1-OX and
SUMO1SLR1-OX obtained by transformation as described above. Negative segregant
lines of Zhonghua and Hwayoung varieties were used as controls for T-DNA insertion
lines knock-out mutants for rice SUMO proteases ELS1 (04Z11JY66) and FUG1 (2A-
20225 obtained from RMD (www.rmd.ncpgr.cn), developed by the National Centre of
Plant Gene Research) and Postech (www.postech.ac.kr/life/pfg/risd, developed by
Gynheung An), respectively. These insertion line seeds were germinated, self-crossed
and screened using PCR analysis for homozygosity (Rosa et al., 2018). The T-DNA
insertion position was determined by sequencing. Gene expression was analysed by
RT-qPCR (Rosa et al., 2018).
582
Salinity Stress Assays
For the assays performed at the seedling stage, heterozygous transgenic seeds were
germinated and selected through screening for hygromycin resistance and GUS
staining. Seedlings were transferred to hydroponics for salinity resistance screening
(Almeida et al., 2016) with half-strength Murashige & Skoog Medium and MES buffer
250 mg/L (Duchefa) with pH adjusted to 5.2. Seedlings were grown for 14 days before
the salt stress assay. Then, the seedlings were exposed to salt stress, with electric
conductivity (EC) set and maintained at 12 dS/m with sodium chloride solution for the
extension of the stress treatment. After 8 days, the seedlings were transferred to
regular medium for recovery. Stress severity (SES score) – 1 for lower severity and 9
for higher severity - was evaluated as set by IRRI (Gregorio et al., 1997) at 6 and 8
days after stress and 8 days after recovery. Samples for transcriptomic analysis – 4
replicates of a pool of 3 plants - were collected at 8 hours after the initiation of the salt
stress, corresponding to 10 hours after the lights went on, for both control and salt
treatments, and frozen immediately in liquid nitrogen for grinding and analysis.
Samples from 0h, 2h, 4h, and 8h treatments were obtained from shoots and roots

collected separately, and frozen immediately in liquid nitrogen. Protein was extracted,
detected and quantified as described above. For the booting stage assay, previously
germinated and selected heterozygous transgenic seeds were sown in pots with 2-
parts soil, 2-parts peat, and 1-part vermiculite. The experiment was set up in the
greenhouse from June to September, according to IRRI standard screening
procedures (Gregorio et al., 1997). Genotypes were maintained in separate
containers, for both stress and control pots, to control for discrepancies in panicle
initiation times. Panicle initiation was evaluated by the emergence of the flag-leaf with
a distance between 1 and 2 cm to the adjacent leaf and salinity applied individually for
each genotype at that point. Electric conductivity (EC) was set and maintained at 8
dS/m with sodium chloride solution for the extension of the life cycle. Upon complete
maturation of the seed, agronomical parameters were measured (plant height, tillering,
and panicle length) and panicles collected for further characterization of fertility
(analysing the number of full matured seeds to total number of flowers ratio) and
productivity (weight of 1000 seeds as measured in a Pfeuffer seed counter). The
significance of statistics was asserted by t-test with Holm-Sidak method correction for
multiple comparisons and a $p<0.05$ threshold.
616

RNA Extraction and expression analysis

RNA was extracted from a total of 100 mg of ground frozen shoot tissue with Direct-
zol kit (Zymo Research) according to manufacturer's protocol with in-column DNAse
treatment. The first-strand cDNA was synthesized from 2 µg of total RNA with an
anchored-oligo (dT)18 primer using the Transcriptor High Fidelity cDNA Synthesis Kit
(Roche) according to the manufacturer's instructions. Reverse Transcription
quantitative PCR (RT- qPCR) analysis was performed using a LightCycler 480 system
(Roche) and the SYBR Green I Master mix (Roche). The RT-qPCR reactions were
carried out with the primers listed in Supplemental Table S4. PCR running conditions
were: one cycle at 95ºC for 5 min and 45 cycles of amplification at 95ºC for 10 s, 56–
60ºC for 10 s and 72ºC /10 s. The CT values were calculated using three technical
replicates, and gene expression was assessed relative to described internal reference
transcript controls, rice *UBIQUITIN CONJUGASE2* (OsUBC2) and *UBIQUITIN10*
(OsUBQ10) (Moraes et al., 2015; Pabuayon et al., 2016). For the RNA-seq procedure,
the small RNAs in the extracted total RNA samples were cleaned by using the RNeasy
spin column from RNeasy Mini Kit (Qiagen) following RNA Cleanup protocol. RNA

quality was assessed by Fragment Analyzer (ThermoFisher) with RNA Quality Number over 5. A total of 18 cDNA libraries were prepared, using three biological replicates for each genotype (SLR1-OX, SUMO1SLR1-OX, Nipponbare) and each condition (mock and salt). Enzymatic fragmentation with Pico Nextera kit and Smart-Seq2 multiplexed-library preparation were performed[49] and single-end sequencing on NextSeq500 (Illumina) with High Output Kit (75 cycles).

RNA-seq Data Analysis

Raw data FASTQ files were inputted on R/Bioconductor-based user interface analysis software Chipster for RNA-seq. Single-end reads were trimmed with PRINSEQ for 55 bp total with 18 bp trimmed on 5' end and quality control of 28% set for the 3' end for adapter removal. Read alignment was performed with TOPHAT2 against the *Oryza sativa* genome (version IRGSP-1.0.36), selecting Sanger-Phred +33 as quality score format. An average of 25.000.000 and 21.000.000 aligned and uniquely mapped reads, respectively, were obtained per library (Supplemental Table S2). For differential expression analysis, BAM files resulting from alignment were submitted to pairwise comparisons with Cufflinks2 tool Cuffdiff for a set 0.05 False Discovery Rate (5% FDR) with sequence-specific bias and multi-mapped read corrections enabled. Differentially expressed genes (DEGs) with a 1.5 fold-change were selected for data visualization and functional annotation using ClueGO plug-in (Bindea et al., 2009) for Cytoscape (Su et al., 2014).

Promoter Analysis and Interaction Validation by Y2H and BiFC

The 1000 bp promoter region of differential expressed genes DEGs were downloaded from Rice Annotation Project Database (RAP-DB) (Sakai et al., 2013) and submitted to PlantPAN 2.0 (Chow et al., 2016) for *cis*-elements for transcription factor binding site (TFBS) analysis. Gene Group Analysis was used to search for common (90%) *cis*-element binding (TFBS) sites and the transcription factors (TFs) associated with DEG promoters. Rice TFs were examined individually for experimentally verified data concerning the interaction to binding sites and downstream target genes listing (Supplemental Table S2). Resulting target genes were then overlapped with previously verified DEGs from RNA-seq analysis. For the Y2H assays, coding sequences of SLR1 and OsSUMO1(GG-AA)::SLR1 were cloned into the pGADT7 vector (Stratagene, USA) while OsbHLH089, OsbHLH094, OsbZIP23, OsIDEF1, OsOSH1,

OsPCF1 and OsYAB4 were cloned into the pGBKT7 (Stratagene, USA) vector. These
constructs were transformed into *Saccharomyces cerevisiae* strain Y2HGold
(Clontech, US) applying the LiAc method. Yeast transformed cells were plated on
synthetic defined (SD) medium lacking leucine and tryptophan for plasmid
transformation control and in SD lacking leucine (-Leu), tryptophan (-Trip), adenine (-
Ade), and histidine (-His) for interaction screening. The interactions were evaluated in
three individual colonies transformed with both plasmids. pAD-WT and pBD-WT from
the HybriZAP 2.1 kit (Stratagene, USA) were used as a control of positive interaction
(C+). pGBKT7 and pGADT7 empty vectors were also co-transformed with pGADT7-
SLR1, pGADT7-SUMO1($_{GG-AA}$)::SLR1 and pGBKT7-bHLH089, pGBKT7-bZIP23,
pGBKT7-IDEF1, and pGBKT7-YAB4, respectively, as negative controls. For
bimolecular fluorescence complementation (BiFC), OsSLR1, OsSUMO1($_{GG-AA}$)::SLR1
were cloned in YFN43, and OsbHLH089, OsbHLH094 and OsYAB4 were cloned into
YFC43 (Belda-Palazón et al., 2012). *Nicotiana benthamiana* leaves were co-infiltrated
with *Agrobacterium tumefaciens* strain EH105 harbouring the appropriate vectors.
Fluorescence was visualized using the Leica DM6 B fluorescence microscope (40x)
with L5 filter cube (Leica, Germany) 48-72h following infiltration. As negative control,
YFN43 (empty vector) was also co-infiltrated with YFC43-bHLH089, YFC43-bHLH094
and YFC43-YAB4. *Agrobacterium* expressing the viral silencing suppressor P19 was
included in all infiltrations to increase the expression in *N. benthamiana* leaves. BiFC
analyses were performed essentially as described in Belda-Palazón et al., 2012.
688

Accession Numbers
Sequence data from this article can be found in the Rice Annotation Project Database
(RAP-DB) under the following accession numbers: OsSLR1 (Os03g0707600),
OsSUMO1 (Os01g0918300), OsSUMO2 (Os01g0918200), OsSAE1
(Os11g0497000), OsSCE1a (Os03g0123100), OsELS1 (Os01g0355900), OsFUG1
(Os03g0344300), OsUBC2 (Os02g0634800), OsUBQ10 (Os06g0681400), OsP5CS
(Os05g0455500), OsRAB21 (Os11g0454300), RGA (At2g01570), OsCPS4
(Os04g0178300), OsKSL7 (Os02g0570400), OsCYP701A8/KO1 (Os06g0569500),
OsCYP99A3 (Os04g0178400), OsCYP71Z6 (Os02g0570500), OsCYP76M8
(Os02g0569400), OsCYP71Z7 (Os02g0570700), OsCKX2 (Os01g0197700),
OsCKX5 (Os01g0775400), OsLOX2 (Os08g0509100), OsCHT3 (Os06g0726100),
OsCHT8 (Os10g054290), OsCHT11 (Os03g0132900), OsGA20ox2

(Os01g0883800), OsGA2ox3 (Os01g0757200), OsPCF1 (Os04g0194600),

OsRISBZ3 (Os02g0266800), OsbHLH089 (Os03g0802900), OsbHLH094

(Os07g0193800), OsbZIP23 (Os02g0766700), OsOSH1 (Os03g0727000), OsPCF1

(Os03g0785800), OsIDEF1 (Os11g0175700), OsYAB4 (Os02g0643200).

Supplementary Materials

Figure S1. Alignment of DELLAs protein sequences of selected monocots and dicots.

Figure S2. SRL1 can be SUMOylated by both OsSUMO1 and OsSUMO2 *in vitro.*

Figure S3. Accumulation dynamics of SRL1 transcript and protein in response to salt stress.

Figure S4. Tillering and panicle length were not affected by mild salt stress imposition.

Figure S5. Analysis of salt treatment efficiency by salt-stress marker gene induction.

Figure S6. Salt treatment had a major contribution to the differences in gene expression among Nipponbare, SLR1-OX and SUMO1SLR1-OX.

Figure S7. Transcriptomics reveal differential stress-induced regulation by SUMOylated SLR1.

Figure S8. Protein-protein interaction (PPI) negative controls for Y2H and BiFC assays.

Figure S9. Genevestigator Results for Gene Expression of YABBY4 (LOC_Os02g4950) in Salinity Assays.

Table S1. RNA seq data showing >1.5 fold Differentially Expressed Genes in Nipponbare, SLR1-OX and SUMOSLR1-OX, in mock and salt stress conditions.

Table S2. Expression changes of selected putative SLR1 TF interactors (A) and the respective downstream target genes (B). Fold-change shown as $\log_2$ FC. Comparison of condition A with condition B corresponding to respective selected analysis in Cufflinks (FPKM1 x FPKM2, for condition A and B, respectively). Nipp-Nipponbare; (m)-mock; (s)-stress

Table S3. RNA-seq read count for each of the sequenced libraries and respective legend

Table S4. List of primers used for gene cloning, site directed mutagenesis (SDM) and gene expression quantification (RT-qPCR). Details for the use of specific primers are given in the text.

Acknowledgments

Portuguese Fundação para a Ciência e a Tecnologia (FCT) for fellowships for NMG (SFRH/BD/85799/2012), TF (PD/BD/135584/2018), MTGR (SFRH/BD/84219/2012), MAAR (SFRH/BD/87420/2012), PMB (SFRH/BPD/86742/2012). FCT Investigator IF/00764/2014 (POPH-QREN) for IAA and PTDC/BIA-FBT/31211/2017 for CM contract. Work was supported by P-KBBE/0001/2012 and also by the FCT research unit GREEN-it "Bioresources4sustainability" (UID/Multi/04551/2013 and UIDB/04551/2020). The funding sources had no involvement in study design, analyses, and interpretation of data, writing, or in the submission decision.

Table 1. Enriched binding sites and associated transcription factors in SLR1 DEGs promoters. Promoter sequence analysis output from PlantPAN with Support referencing the percentage of transcription factor binding elements (TFBS/Sequence) found in SLR1- and SUMO1SLR1-dependent stress-responsive DEGs. Resulting TF overlap with DEGs detailed in Supplementary Table S4.

TFBS	Support	TF Family	TF	Annotation
GGTCCCAC	91.4	**TCP**	**PCF1/2**	essential for meristematic tissue-specificity expression of the PCNA gene promoter
TACGTA	91.4	**(Others)**	**RISBZ3**	found in seed storage protein gene promoters
GCACGTGC	94.7	**bHLH**	**BHLH089/094**	forms a ternary complex with RSS3 and TIFY11A/JAZ9 to negatively regulate jasmonate-responsive genes
ACACGTGT	97.4	**bZIP**	**BZIP23/46**	binds specifically to the abiotic stress ABA-responsive elements (ABRE) in the promoter of target genes
TNCGTACAA	98.0	**SBP**	**SPL1**	trans-acting factor that binds specifically to the consensus nucleotide sequence 5'-TNCGTACAA-3'
TGAC	98.7	Homeobox	**OSH1**	involved in shoot formation during embryogenesis
CATGC	100.0	**B3**	**IDEF1**	involved in iron deficiency response and tolerance
CCAAT	100.0	**NF-YB**	**NFBY1/2/3/4**	component of the NF-Y/HAP transcription factor complex regulating the expression of photosynthetic genes

REFERENCES

o Achard, P., Cheng, H., De Grauwe, L., Decat, J., Schoutteten, H., Moritz, T., Van Der Straeten, D., Peng, J., and Harberd, N.P. (2006). Integration of plant responses to environmentally activated phytohormonal signals. Science *311*, 91–94.

o Almadanim, M.C., Alexandre, B.M., Rosa, M.T.G., Sapeta, H., Leitão, A.E., Ramalho, J.C., Lam, T.T., Negrão, S., Abreu, I.A., and Oliveira, M.M. (2017). Rice calcium-dependent protein kinase OsCPK17 targets plasma membrane intrinsic protein and sucrose-phosphate synthase and is required for a proper cold stress response. Plant Cell Environ. *40*, 1197–1213.

o Almeida, D.M., Almadanim, M.C., Lourenço, T., Abreu, I.A., Saibo, N.J.M., and Oliveira, M.M. (2016). Screening for Abiotic Stress Tolerance in Rice: Salt, Cold, and Drought. In Environmental Responses in Plants, (New York, NY: Humana Press, New York, NY), pp. 155–182.

o Atanassov, I.I., Atanassov, I.I., Etchells, J., and Turner, S.R. (2009). A simple, flexible and efficient PCR-fusion/Gateway cloning procedure for gene fusion, site-directed mutagenesis, short sequence insertion and domain deletions and swaps. Plant Methods 2009 5:1 *5*, 14.

o Augustine, R.C. and Vierstra, R.D. (2018). SUMOylation: re-wiring the plant nucleus during stress and development. Current opinion in plant biology, 45, pp.143-154.

o Belda-Palazón, B., Ruiz, L., Martí, E., Tárraga, S., Tiburcio, A.F., Culiáñez, F., Farràs, R., Carrasco, P. and Ferrando, A. (2012). Aminopropyltransferases involved in polyamine biosynthesis localize preferentially in the nucleus of plant cells. PLoS One, 7(10), p.e46907.

o Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H., Pagès, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. Bioinformatics *25*, 1091–1093.

o Campanaro, A., Battaglia, R., Galbiati, M., Sadanandom, A., Tonelli, C., and Conti, L. (2016). SUMO proteases OTS1 and 2 control filament elongation through a DELLA-dependent mechanism. Plant Reprod *29*, 1–4.

o Camut, L., Davière, J.-M., and Achard, P. (2017). Dynamic Regulation of DELLA Protein Activity: SPINDLY and SECRET AGENT Unmasked! Mol Plant *10*, 785–787.

o Castro, P.H., Bachmair, A., Bejarano, E.R., Coupland, G., Lois, L.M., Sadanandom, A., van den Burg, H.A., Vierstra, R.D., and Azevedo, H. (2018). Revised nomenclature and functional overview of the ULP gene family of plant deSUMOylating proteases. J. Exp. Bot. *69*, 4505–4509.

o Chow, C.-N., Zheng, H.-Q., Wu, N.-Y., Chien, C.-H., Huang, H.-D., Lee, T.-Y., Chiang-Hsieh, Y.-F., Hou, P.-F., Yang, T.-Y., and Chang, W.-C. (2016). PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants. Nucleic Acids Research *44*, D1154–D1160.

o Conti, L., Nelis, S., Zhang, C., Woodcock, A., Swarup, R., Galbiati, M., Tonelli, C., Napier, R., Hedden, P., Bennett, M., et al. (2014). Small Ubiquitin-like Modifier protein SUMO enables plants to control growth independently of the phytohormone gibberellin. Developmental Cell *28*, 102–110.

o Cremona, C.A., Sarangi, P., and Zhao, X. (2012). Sumoylation and the DNA Damage Response. Biomolecules 2012, Vol. 2, Pages 376-388 *2*, 376–388.

o Crozet, P., Margalha, L., Butowt, R., Fernandes, N., Elias, C.A., Orosa, B., Tomanov, K., Teige,
M., Bachmair, A., Sadanandom, A., et al. (2015). SUMOylation represses SnRK1 signaling in
Arabidopsis. The Plant Journal *85*, 120–133.
o Dai, C., and Xue, H.-W. (2010). Rice early flowering1, a CKI, phosphorylates DELLA protein
SLR1 to negatively regulate gibberellin signalling. The EMBO Journal *29*, 1916–1927.
o Davière, J.-M., and Achard, P. (2015). A pivotal role of DELLAs in regulating multiple hormone
signals. Mol Plant *9*, 10–20.
o de Lucas, M., Daviere, J.M., Rodriguez-Falcon, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S.,
Fankhauser, C., Blazquez, M.A., Titarenko, E., and Prat, S. (2008). A molecular framework for
light and gibberellin control of cell elongation. Nature 451, 480-484.
o Eckhoff, J., and Dohmen, R.J. (2015). In Vitro Studies Reveal a Sequential Mode of Chain
Processing by the Yeast SUMO (Small Ubiquitin-related Modifier)-specific Protease Ulp2. J.
Biol. Chem. *290*, 12268–12281.
o Fukazawa, J., Mori, M., Watanabe, S., Miyamoto, C., Ito, T., and Takahashi, Y. (2017). DELLA-
GAF1 Complex Is a Main Component in Gibberellin Feedback Regulation of GA20 Oxidase 2.
Plant Physiology *175*, 1395–1406.
o Gregorio, G.B., Senadhira, D., and Mendoza, R.D. (1997). Screening rice for salinity tolerance.
Irri.
o Hedden, P., and Thomas, S.G. (2012). Gibberellin biosynthesis and its regulation. Biochemical
Journal *444*, 11–25.
o Hirano, K., Asano, K., Tsuji, H., Kawamura, M., Mori, H., Kitano, H., Ueguchi-Tanaka, M., and
Matsuoka, M. (2010). Characterization of the molecular mechanism underlying gibberellin
perception complex formation in rice. The Plant Cell Online *22*, 2680–2696.
o Hirano, K., Ueguchi-Tanaka, M., and Matsuoka, M. (2008). GID1-mediated gibberellin signaling
in plants. Trends Plant Sci *13*, 192–199.
o Hirano, K., Kouketu, E., Katoh, H., Aya, K., Ueguchi-Tanaka, M. and Matsuoka, M. (2012). The
suppressive function of the rice DELLA protein SLR1 is dependent on its transcriptional
activation activity. The Plant Journal, 71(3), pp.443-453.
o Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem,
W. and Zimmermann, P. (2008). Genevestigator v3: a reference expression database for the
meta-analysis of transcriptomes. Advances in bioinformatics.
o Itoh, H., Matsuoka, M., and Steber, C.M. (2003). A role for the ubiquitin-26S-proteasome pathway
in gibberellin signaling. Trends Plant Sci *8*, 492–497.
o Itoh, H., Sasaki, A., Ueguchi-Tanaka, M., Ishiyama, K., Kobayashi, M., Hasegawa, Y., Minami,
E., Ashikari, M., and Matsuoka, M. (2005). Dissection of the phosphorylation of rice DELLA
protein, SLENDER RICE1. Plant Cell Physiol. *46*, 1392–1399.
o Li, K., Yu, R., Fan, L.M., Wei, N., Chen, H. and Deng, X.W. (2016). DELLA-mediated PIF
degradation contributes to coordination of light and gibberellin signalling in Arabidopsis. Nature
communications, 7, p.11868.
o Li, S.-J., and Hochstrasser, M. (2003). The Ulp1 SUMO isopeptidase: distinct domains required
for viability, nuclear envelope localization, and substrate specificity. *160*, 1069.
o Luís, I.M., Alexandre, B.M., Oliveira, M.M., and Abreu, I.A. (2016). Selection of an Appropriate
Protein Extraction Method to Study the Phosphoproteome of Maize Photosynthetic Tissue. Plos
One *11*, e0164387.
o Mann, D.G.J., LaFayette, P.R., Abercrombie, L.L., King, Z.R., Mazarei, M., Halter, M.C.,
Poovaiah, C.R., Baxter, H., Shen, H., Dixon, R.A., et al. (2011). Gateway-compatible vectors
for high-throughput gene functional analysis in switchgrass (Panicum virgatum L.) and other
monocot species. - PubMed - NCBI. Plant Biotechnol. J. *10*, 226–236.
o Marín-de la Rosa, N., Pfeiffer, A., Hill, K., Locascio, A., Bhalerao, R.P., Miskolczi, P., Grønlund,
A.L., Wanchoo-Kohli, A., Thomas, S.G., Bennett, M.J. and Lohmann, J.U. (2015). Genome
wide binding site analysis reveals transcriptional coactivation of cytokinin-responsive genes by
DELLA proteins. PLoS genetics, 11(7), p.e1005337.
o Miura, K., Jin, J.B., and Hasegawa, P.M. (2007). Sumoylation, a post-translational regulatory
process in plants. Current Opinion in Plant Biology *10*, 495–502.
o Moraes, G.P., Benitez, L.C., do Amaral, M.N., Vighi, I.L., Auler, P.A., da Maia, L.C., Bianchi, V.J.,
and Braga, E.J.B. (2015). Evaluation of reference genes for RT-qPCR studies in the leaves of
rice seedlings under salt stress. Genet. Mol. Res. *14*, 2384–2398.
o Munns, R., and Tester, M. (2008). Mechanisms of salinity tolerance. Annu. Rev. Plant Biol. *59*,
651–681.

o Negrão, S., Courtois, B., Ahmadi, N., Abreu, I., Saibo, N., and Oliveira, M.M. (2011). Recent
Updates on Salinity Stress in Rice: From Physiological to Molecular Responses. Critical
Reviews in Plant Sciences *30*, 329–377.
o Nelis, S., Conti, L., Zhang, C., and Sadanandom, A. (2015). A functional Small Ubiquitin-like
Modifier (SUMO) interacting motif (SIM) in the gibberellin hormone receptor GID1 is conserved
in cereal crops and disrupting this motif does not abolish hormone dependency of the DELLA-
GID1 interaction. Plant Signaling & Behavior *10*, e987528.
o Otomo, K., Kenmoku, H., Oikawa, H., König, W.A., Toshima, H., Mitsuhashi, W., Yamane, H.,
Sassa, T., and Toyomasu, T. (2004). Biological functions of ent- and syn-copalyl diphosphate
synthases in rice: key enzymes for the branch point of gibberellin and phytoalexin biosynthesis.
Plant J. *39*, 886–893.
o Pabuayon, I.M., Yamamoto, N., Trinidad, J.L., Longkumer, T., Raorane, M.L., and Kohli, A.
(2016). Reference genes for accurate gene expression analyses across different tissues,
developmental stages and genotypes in rice for drought tolerance. Rice *9*, 32.
o Rosa, M.T.G., Almeida, D.M., Pires, I.S., da Rosa Farias, D., Martins, A.G., da Maia, L.C., de
Oliveira, A.C., Saibo, N.J.M., Oliveira, M.M., and Abreu, I.A. (2018). Insights into the
transcriptional and post-transcriptional regulation of the rice SUMOylation machinery and into
the role of two rice SUMO proteases. BMC Plant Biol *18*, 349.
o Sakai, H., Lee, S.S., Tanaka, T., Numa, H., Kim, J., Kawahara, Y., Wakimoto, H., Yang, C.-C.,
Iwamoto, M., Abe, T., et al. (2013). Rice Annotation Project Database (RAP-DB): an integrative
and interactive database for rice genomics. - PubMed - NCBI. Plant Cell Physiol. *54*, e6–e6.
o Sakamoto, T., Sakakibara, H., Kojima, M., Yamamoto, Y., Nagasaki, H., Inukai, Y., Sato, Y., and
Matsuoka, M. (2006). Ectopic Expression of KNOTTED1-Like Homeobox Protein Induces
Expression of Cytokinin Biosynthesis Genes in Rice. Plant Physiology *142*, 54–62.
o Sarhadi, E., Bazargani, M.M., Sajise, A.G., Abdolahi, S., Vispo, N.A., Arceta, M., Nejad, G.M.,
Singh, R.K., and Salekdeh, G.H. (2012). Proteomic analysis of rice anthers under salt stress.
Plant Physiology and Biochemistry *58*, 280–287.
o Srivastava, A.K., Zhang, C., and Sadanandom, A. (2016a). Rice OVERLY TOLERANT TO SALT
1 (OTS1) SUMO protease is a positive regulator of seed germination and root development.
Plant Signaling & Behavior *11*, e1173301.
o Srivastava, A.K., Zhang, C., Caine, R.S., Gray, J., and Sadanandom, A. (2017). Rice SUMO
protease Overly Tolerant to Salt 1 targets the transcription factor, OsbZIP23 to promote drought
tolerance in rice. The Plant Journal *92*, 1031–1043.
o Srivastava, A.K., Zhang, C., Yates, G., Bailey, M., brown, A., and Sadanandom, A. (2016b).
SUMO is a critical regulator of salt stress responses in rice. Plant Physiology *170*,
pp.01530.2015–pp.01530.2391.
o Su, G., Morris, J.H., Demchak, B., and Bader, G.D. (2014). Biological network exploration with
Cytoscape 3. - PubMed - NCBI. Current Protocols in Bioinformatics *47*, 8.13.1–8.13.24.
o Ueguchi-Tanaka, M., Hirano, K., Hasegawa, Y., Kitano, H., and Matsuoka, M. (2008). Release
of the repressive activity of rice DELLA protein SLR1 by gibberellin does not require SLR1
degradation in the gid2 mutant. Plant Cell *20*, 2437–2446.
o Ulrich, H.D. (2008). The fast-growing business of SUMO chains. Mol. Cell *32*, 301–305.
o Van De Velde, K., Ruelens, P., Geuten, K., Rohde, A., and Van Der Straeten, D. (2017).
Exploiting DELLA Signaling in Cereals. Trends Plant Sci.
o Vanstraelen, M., and Benková, E. (2012). Hormonal interactions in the regulation of plant
development. Annu. Rev. Cell Dev. Biol. *28*, 463–487.
o Werner, A., Moutty, M.-C., Möller, U., and Melchior, F. (2009). Performing in vitro sumoylation
reactions using recombinant enzymes. Methods Mol. Biol. *497*, 187–199.
o Xue, Y., Zhou, F., Fu, C., Xu, Y., and Yao, X. (2006). SUMOsp: a web server for sumoylation site
prediction. Nucleic Acids Research *34*, W254–W257.
o Yang, C., Ma, Y., and Li, J. (2016). The rice YABBY4 gene regulates plant growth and
development through modulating the gibberellin pathway. J. Exp. Bot. *67*, erw319–erw5556.
o Yano, K., Morinaka, Y., Wang, F., Huang, P., Takehara, S., Hirai, T., Ito, A., Koketsu, E.,
Kawamura, M., Kotake, K. and Yoshida, S., 2019. GWAS with principal component analysis
identifies a gene comprehensively controlling rice architecture. Proceedings of the National
Academy of Sciences, 116(42), pp.21262-21267.
o Yates, G., Srivastava, A.K., and Sadanandom, A. (2016). SUMO proteases: uncovering the roles
of deSUMOylation in plants. J. Exp. Bot. *67*, 2541–2548.
o Zentella, R., Hu, J., Hsieh, W.P., Matsumoto, P.A., Dawdy, A., Barnhill, B., Oldenhof, H.,
Hartweck, L.M., Maitra, S., Thomas, S.G. and Cockrell, S. (2016). O-GlcNAcylation of master

growth repressor DELLA by SECRET AGENT modulates multiple signaling pathways in
Arabidopsis. Genes & development, 30(2), pp.164-176.
o Zentella, R., Sui, N., Barnhill, B., Hsieh, W.-P., Hu, J., Shabanowitz, J., Boyce, M., Olszewski,
N.E., Zhou, P., Hunt, D.F., et al. (2017). The Arabidopsis O-fucosyltransferase SPINDLY
activates nuclear growth repressor DELLA. Nat. Chem. Biol. *21*, 985–485.

Figure 1. SLR1 is SUMOylated *in vitro* in a novel SUMO-conjugation canonical motif. **(A)** *in silico* prediction of SLR1 canonical motifs (ψ-K-X-E) in the N-terminal region as scored by SUMOsp2 software (Xue et al., 2006), compared to *Arabidopsis thaliana* RGA corresponding region. Motifs showing high SUMOylation probability are marked in red, and low probability motifs are marked in blue. SLR1 K2 and K60 are shown in green and orange, respectively. **(B-C)** *in vitro* SUMOylation assays using rice SUMOylation machinery (OsSCE1, OsSAE1/2, and OsSUMO1) with recombinant SLR1 and SLR1t (SLR1 N-terminal truncated) proteins, treated with ULP1 SUMO protease, as indicated. SLR1 and SLR1t were detected by immunoblotting using custom made anti-SLR1 antibody, where **(B)** corresponds to the full-length SLR1, and **(C)** to the SLR1t coding the leading 190 first SLR1 residues, as well as the mutated isoforms SLR1t K2R and SLR1t K60R.

Figure 2. SUMO-specific protease activity reveal SLR1 *in vivo* regulation by SUMO.
(A) *Oryza sativa* L. cv Nipponbare non-denaturing protein extracts treated with *Saccharomyces cerevisiae* Ulp1 (ULP1) for SLR1 deSUMOylation analysis as detected by immunoblot using anti-SLR1 antibody. **(B)** Loading-normalized relative quantification of total SLR1 protein in *els1* and *fug1* mutants, and the respective negative segregants (NS) after deSUMOylation by ULP1 treatment; and **(C)** total SUMOylated SLR1, as shown by ratio comparison to non-SUMOylated protein after ULP1 treatment. Error bars correspond to standard error (SE; n=4). Statistically significant values are represented as **(p<0.01). The significance of statistics was asserted by t-test with Holm-Sidak method correction for multiple comparisons.

Figure 3. SLR1 SUMOylation confers salt tolerance at the seedling stage. (A) Depiction of gene constructs used to generate rice SLR1-OX and SUMO1SLR1-OX transgenic lines. *Zea mays* Ubi1 (ZmUbi1) promoter drives the expression of both OsSLR1 (SLR1-OX) and OsSLR1-OsSUMO1 gene fusion (SUMO1SLR1-OX). **(B)** SLR1 and SUMO1SLR1 protein accumulation in T2 independent transgenic lines (#1 and #2) of SUMO1SLR1-OX and SLR1-OX as detected by immunoblot using custom made anti-SLR1 antibody. **(C)** Stress severity as evaluated by IRRI salinity trial SES score for all lines in 6 and 8 days of stress and after 8 days of recovery period.

Figure 4. SLR1 SUMOylation reduces the impact of salt stress on yield. Agronomic parameters as measured at the end of rice life cycle: **(A)** plant height (N=6) **(B)** fertility as evaluated by the number of filled grains ratio to total panicle flowers (n>10) and **(C)** productivity as evaluated by the weight of 1000 grains (n>10). Error bars correspond to standard error (SE; 2 independent biological replicates with N=6 each) and statistically significant p-values represented as *(p<0.05). The significance of statistics was asserted by t-test with Holm-Sidak method correction for multiple comparisons.

Figure 5. SLR1 SUMOylation affects its interaction with key transcription factors **(A)** Yeast-two hybrid (Y2H) analysis of both SLR1 and SUMO1SLR1. SUMO1::SLR1 fusion carries two mutations in the SUMO-protease cleavage motif (GG → AA) to avoid its removal by endogenous yeast proteases and guarantee constitutive SUMO attachment to SLR1 during the assay. Positive controls marked as (SD/-Leu/-Trip) and positive interactions marked with (SD/-Leu/-Trip/-Ade/-His). The vectors pAD-WT and pBD-WT (Stratagene) were used as positive control. **(B)** Bimolecular fluorescent complementation (BiFC) assays showing positive interactions between nYFP: SLR1 with cYFP:BHLH089, cYFP:BHLH094, and cYFP:YAB4. The presence of SUMO1 attached to lysine 2 of SLR1 (nYFP:SUMO1(GG-AA)::SLR1) disrupts SLR1 interaction with these transcription factors. Negative controls for Y2H and BiFC assays can be found in Supplementary Figure S8.