Crystal structures of the sheeppoxvirus encoded inhibitor of apoptosis SPPV14 bound to Hrk and Bax BH3 peptides

Chathura D. Suraweera[1], Denis R. Burton[1#], Mark G. Hinds[2&] and Marc Kvansakul[1&*]

[1]La Trobe Institute for Molecular Science, Department of Biochemistry and Genetics, La Trobe University, VIC 3086, Australia

[2]Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Australia

#current address:

&Co-senior author

*Correspondence: m.kvansakul@latrobe.edu.au (M.K.)

Running title: Crystal structures of SPPV14:Hrk and SPPV14:Bax BH3 complexes

Keywords: Poxvirus, sheeppoxvirus, apoptosis, X-ray crystallography, isothermal titration calorimetry, Bcl-2

Abstract

Programmed death of infected cells is used by multicellular organisms to counter viral infections. Sheeppoxvirus encodes for SPPV14, a potent inhibitor of Bcl-2 mediated apoptosis. We reveal the structural basis of apoptosis inhibition by determining crystal structures of SPPV14 bound to BH3 motifs of proapoptotic Bax and Hrk. The structures reveal that SPPV14 engages BH3 peptides using the canonical ligand binding groove. Unexpectedly, Arg84 from SPPV14 forms an ionic interaction with the conserved Asp in the BH3 motif in a manner that replaces the canonical ionic interaction seen in almost all host Bcl-2:BH3 motif complexes. These results reveal the flexibility of virus encoded Bcl-2 proteins to mimic key interactions from endogenous host signalling pathways to retain BH3-binding and pro-survival functionality.

Introduction

The altruistic death of infected host cells via programmed cell death or apoptosis is a potent feature of the first line of defence against invading pathogens [1]. The ability of multicellular hosts to eliminate infected cell via apoptosis forced viruses to evolve sophisticated strategies to counter host cell apoptotic defences, including the use of virus encoded homologs of the B-cell lymphoma 2 or Bcl-2 family [2]. As primary modulators of the mitochondrially mediated or intrinsic apoptosis pathway the Bcl-2 family can be divided into two factions; the prosurvival members and the proapoptotic members [3]. All mammalian Bcl-2 family members are characterized by the presence of one or more of the Bcl-2 homology or BH sequence motifs. The prosurvival members, which include Bcl-2, Bcl-w, Bcl-x_L, A1, Mcl-1 and Bcl-b all feature multiple BH motifs as well as a transmembrane region that enables localization to the outer mitochondrial membrane [4]. The proapoptotic family members can be further separated into two groups, the multi BH motif proteins Bak, Bax and Bok, and the

BH3-only proteins Bim, Bid, Puma, Bad, Bik, Bmf, Hrk and Noxa that only feature the BH3 motif [5]. Mechanistically, the BH3 only proteins act by either relieving the ability of prosurvival Bcl-2 to hold Bak and Bax in an inactive state, or may directly activate Bak and Bax [6]. The interplay between the members is mediated by a helix-in-groove interaction, where a helical BH3 motif is bound in the canonical hydrophobic ligand binding groove on prosurvival Bcl-2 proteins [7]. Ultimately, it is the balance of the prosurvival and proapoptotic Bcl-2 members that determines the fate of a given cell.

Large DNA viruses encode a number of sequence and structural homologs of Bcl-2, including examples among the *adenoviridae* [8], *asfarviridae* [9, 10], *herpesviridae* [11, 12] and *iridoviridae* [13, 14]. Among the *poxviridae*, the majority of genera have been shown to harbor one or more examples, including the *orthopoxviridae* [15, 16], *leporipoxviridae* [17, 18], *cervidpoxviridae* [19, 20], *avipoxviridae* [21, 22], *parapoxviridae* [23] and *chordopoxviridae [24],* with the vaccinia virus F1L and myxomavirus M11L the prototypical members of the family. Amongst the *capripoxviridae*, sheeppoxvirus has been shown to encode the potent apoptosis inhibitory protein SPPV14, a Bcl-2 homolog [25]. SPPV14 has been shown to inhibit apoptosis in transfected cells after treatment with etoposide, arabinose-C or UV irradiation. Furthermore, interaction studies revealed that SPPV14 is able directly bind BH3 motif peptides of several mammalian proapoptotic Bcl-2 family members including, and functionally replaces F1L in the context of a recombinant vaccinia virus infection. However, the structural and biochemical basis for apoptosis inhibition by SPPV14 remained unclear. Here we report the crystal structures and affinity data of SPPV14 bound to BH3 motifs of proapoptotic Bcl-2 family members. Our findings provide a structural and biochemical platform to understand sheeppoxvirus mediated inhibition of apoptosis.

Materials and methods

Protein expression and purification

Recombinant SPPV14ΔC31, (uniprot accession number A0A2P1A8X8; SPPV14 residues 1-145 with 31 residue C-terminal deletion and hereafter referred to as SPPV14) was expressed and purified as previously described [26]. SPPV14 mutants R84A and Y46A were synthesized as codon optimized cDNA (Genscript), cloned into the bacterial expression vector pGEX-6P3 and purified as previously described [26]. Purified protein obtained using glutathione affinity chromatography was then subjected to size-exclusion chromatography using a Superdex S75 10/300 column mounted on an ÄKTA Pure system (GE Healthcare) equilibrated in 25 mM HEPES pH 7.5, 150 mM NaCl and 10 mM TCEP (Tris(2-carboxyethyl)phosphine hydrochloride), and fractions analysed using SDS-PAGE. The final sample purity was estimated to be higher than 95% based on SDS–PAGE analysis and appropriate fractions were pooled and concentrated using a centrifugal concentrator with 3 kDa molecular weight cut-off (Amicon ® Ultra 15) to final concentration of 10.0 mg/ml.

Measurement of dissociation constants

Binding affinities were measured using a MicroCal iTC200 system (GE Healthcare) at 25°C using wild type SPPV14 as well as two mutants SPPV14 Y46A and R84A in 25 mM HEPES pH 7.5, 150 mM NaCl, 10 mM TCEP at a final concentration of 30 μM. BH3 motif peptides were used at a concentration of 300 μM and titrated using 19 injections of 2.0 μl of ligand solution. All affinity measurements were performed in triplicate. Protein concentrations were measured using a Nanodrop UV spectrophotometer (Thermo Scientific) at a wavelength of 280 nm. Peptide concentrations were calculated based on the dry peptide weight after synthesis. The BH3-motif peptides used were commercially synthesized and were purified to a final purity of 95% (GenScript) and based on the human sequences previously described [27] except for Bok BH3: VPGRLAEVCAVLLRLGDELEMIRPSV (accession code

Q9UMX3, residues 59-84).

Crystallization and structure determination

Crystals for SPPV14:Hrk BH3 and SPPV14:Bax BH3 complexes were obtained by mixing SPPV14 with human Hrk 26-mer or Bax 28-mer peptide using a 1:1.25 molar ratio as described [28] and concentrated using a centrifugal concentrator with 3 kDa molecular weight cut-off (Amicon ® Ultra 0.5) to 10.0 mg/ml. Concentrated protein was immediately used for crystallization trials. Initial high throughput sparse matrix screening was performed using 96 well sitting drop trays (swissic, Neuheim, Switzerland).

SPPV14:Hrk BH3 crystals were grown by the sitting drop vapour diffusion method at 20 °C in 0.2 M Sodium potassium tartrate, 0.1 M Bis-Tris propane pH 6.5, 20% PEG 3350. The crystals were flash cooled at -173 °C in mother liquor supplemented with 25% glucose. The SPPV14: Hrk BH3 complex formed single cuboidal crystals belong to space group $C222_1$ with a=43.28 Å, b=70.66 Å, c=115.08 Å, α=90.00°, β=90.00°, γ=90.00° in the orthorhombic crystal system.

All diffraction data were collected at the Australian Synchrotron MX2 [29] beamline using an Eiger detector with an oscillation range 0.1° per frame at a wavelength of 0.9537 Å. The diffraction data were integrated using XDS [30] and scaled using AIMLESS [31]. A molecular replacement solution was obtained using the Balbes [32] automated pipeline and identified M11L (PDB code: 2BJY)[33] as the optimal model. SPPV14:Hrk BH3 crystals contained one molecule of SPPV14 and one Hrk BH3 peptide in the asymmetric unit, with a 43% solvent content and gave final TFZ and LLG values of 32.2 and 1539, respectively. The final model of SPPV14:Hrk BH3 was built manually over several cycles using Coot [34] and refined using PHENIX [35] giving final R_{work}/R_{free} values of 0.221/0.235, with 99% of residues in the favoured region of the Ramachandran plot and no outliers.

SPPV14:Bax BH3 crystals were obtained in 1.0 M Lithium chloride, 0.1 M citrate pH 5.0, 30% PEG 6000. The crystals were flash cooled at -173 °C in mother liquor. The SPPV14: Bax BH3 complex formed single rod-shaped crystals belong to space group $P2_1$ with cell dimensions a=100.27 Å, b=78.56 Å, c=107.76 Å, α=90.00°, β=110.96°, γ=90.00° in the monoclinic crystal system. Diffraction data collection, integration and scaling were performed as described above. The molecular replacement was performed using PHASER [36] with the previously solved structure of SPPV14:Hrk BH3 as a search model. SPPV14:Bax BH3 crystals contain eight molecules of SPPV14 and eight Bax BH3 peptides, with a 48.9% solvent content and final TFZ and LLG values of 11.2 and 1720 respectively. The final model of SPPV14:Bax BH3 was built manually as described above giving final R_{work}/R_{free} values of 0.238/ 0.288, with 99% of residues in Ramachandran favoured region and no outliers. All images for SPPV14:Hrk and SPPV14:Bax complexes were generated using PyMOL molecular graphic system version 1.8.6.0 (Schrödinger, LLC, New York, USA). The coordinates for the structures were deposited at the Protein Data Bank (PDB) using accession codes 6XY4 and 6XY6. All raw images were deposited at the SBGridDB using their PDB accession codes [37]. All software were accessed through the SBGrid suite [38].

Results

We previously showed sheeppoxvirus encoded SPPV14 binds peptides spanning the BH3 motif of the proapoptotic Bcl-2 family members Bak and Bax as well as Bim, Bid, Bmf, Hrk and Puma as determined by surface plasmon resonance (SPR) [25]. Here we performed isothermal titration calorimetry (ITC) on recombinant SPPV14 with BH3 motif peptides of human proapoptotic Bcl-2 to determine the binding specificity of SPPV14 (Table 1). ITC confirmed Bak and Bax as well as Bim, Bid, Bmf, Hrk and Puma as high affinity interactors,

as previously reported [25]. In addition, we observed weaker affinities for Bad and Bok, with K_D values of 5.2 μM and 7.6 μM, respectively. To understand the structural basis for BH3 motif binding by SPPV14 we expressed and purified recombinant SPPV14 lacking the C-terminal 31 residues (SPPV14ΔC31, herein referred to as SPPV14) and reconstituted SPPV14:Hrk BH3 and SPPV14:Bax BH3 complexes. Crystals of the SPPV14:Hrk BH3 complex diffracted to 2.05 Å (Table 2). After molecular replacement using M11L as a search model (PDB ID 2BJY) [18] clear and continuous electron density was observed for SPPV14 residues 5-139 and Hrk residues 26-50 (Figure 2A), with the remaining residues presumed disordered. SPPV14 adopts a globular helical bundle structure comprising a central helix α5 around which 6 additional helices are scaffolded to form the classical Bcl-2 fold (Figure 1A). A Dali analysis identified deerpoxvirus DPV022 (PDB ID: 4UF3) [20] as the closest structural homolog of SPPV14 with an rmsd of 2.1 Å over 106 Cα atoms, with M11L the second closest match with an rmsd of 2.3 Å over 111 Cα atoms (Figure 1C).

SPPV14 harbors the canonical ligand binding groove formed by helices α2-5, which engages Hrk BH3 (Figure 1A, 2A). The mode of binding of BH3 motif peptides observed for SPPV14 is nearly identical to what has been previously observed for M11L:Bak (Figure 1C), with the SPPV14:Hrk complex superimposing onto the M11L:Bak complex with an rmsd of 1.8 Å over the entire SPPV14 chain. The three conserved hydrophobic residues L37, I40 and L44 as well as T35 from Hrk protrude into the SPPV14 binding groove and engage the resident four hydrophobic pockets (Figure 3A). In addition to these hydrophobic interactions, SPPV14 forms an ionic interaction with Hrk via SPPV14 R84 and Hrk D42 carboxyl group. Furthermore, two hydrogen bonds are observed between SPPV14 Y46 hydroxyl and Hrk E43 carboxyl SPPV14 and T78 hydroxyl with Hrk A34 amide.

The SPPV14:Bax complex was solved by molecular replacement using the SPPV14 model obtained from the SPPV14:Hrk complex and refined to a resolution of 2.9 Å (Figure

1B, Table 2). Clear and continuous density was observed for SPPV14 5-139 and Bax 52-74 (Figure 2 B). The four conserved hydrophobic residues L59, L63, I66 and L70 from Bax are nestled in the corresponding four hydrophobic pockets of the SPPV14 binding groove (Figure 3B). Furthermore, Bax M74 also makes contact with the SPPV14 binding groove in a pocket formed by residues C38, V41, I42, F133 and N137. Additional interactions are observed as ionic interactions between SPPV14 R84 and the carboxyl groups of Hrk D68 and D71 as well as hydrogen bonds between SPPV14 S81 hydroxyl and Bax K64 lysyl, and the main chain carbonyl of SPPV14 I74 and Bax S60 hydroxyl group (Figure 3B).

To validate our crystal structures we performed structure-guided mutagenesis to modulate the affinity of SPPV14 for BH3 motif ligands (Table 1). Indeed, the SPPV14 mutants Y46A and R84A display attenuated affinities, with SPPV14 Y46A binding Bak, Bax, Bim, Bmf, Hrk and Puma with lower affinity and losing binding to Bad, Bid, Bik and Bok whilst SPPV14 R84A bound Bax, Bak, Bid, Bik and Hrk with lower affinities and lost binding to Bok compared to wildtype protein.

Discussion

Subversion of premature host cell apoptosis is a frequently used strategy for many large DNA viruses, including the *poxviridae,* to evade the innate immune response [39]. It has previously been shown that sheeppoxvirus harbors the potent apoptosis inhibitor SPPV14, which acts by sequestering BH3-only proteins including Bim, Bid, Bmf, Hrk and Puma as well as Bak and Bax [25]. In addition to the known interactors, we observed previously unreported low micromolar affinities for Bad and Bok BH3 motifs. Previous studies had relied on the use of competition surface plasmon resonance spectroscopy measurements, which could not detect binding affinities weaker than $IC_{50} = 2$ μm. These findings implicate Bad and Bok inhibition as a potential anti-apoptotic strategy of SPPV14, however considering the comparatively low

affinity of the interaction the functional relevance of Bad and Bok inhibition remains to be established.

Having identified potentially new BH3 ligands we now defined the structural basis for BH3 motif interactions with SPPV14 by determining the crystal structures of the SPPV14:Hrk BH3 and SPPV14:Bax BH3 complexes. SPPV14 binds both Hrk and Bax tightly with K_D values of 39 and 26 nM, respectively, and the interactions are mediated by the canonical hydrophobic interactions involving the four hydrophobic pockets in SPPV14 as well as several ionic interactions and hydrogen bonds. A comparison with other virus encoded Bcl-2 proteins reveals that unlike the myxomavirus M11L and vaccinia virus F1L prosurvival Bcl-2 proteins, which both feature low sequence identity to mammalian Bcl-2 [7], SPPV14 forms an ionic interaction using R84 with Hrk D42 or Bax D68 that is reminiscent of the highly conserved ionic interaction between a conserved Arg in the BH1 region of mammalian prosurvival Bcl-2 proteins and the conserved Asp of the BH3 motif (Figure 4A). Such an ionic interaction is seen in all mammalian prosurvival Bcl-2 protein : BH3 peptide complexes exemplified by Bcl-x_L R139-Bax D68 (Figure 4A). This Arg:Asp interaction is also observed in the complexes with BH3 peptides of the virus-encoded Bcl-2 homologs A179L [10, 40], BHRF1 [12] and FPV039 [22]. Unlike the canonical ionic interaction [41, 42], the geometric configuration of the ionic bond in SPPV14 is different, allowing SPPV14 to engage BH3 motif in manner that resembles the use of the BH1 motif in mammalian prosurvival Bcl-2 interactions despite the lack of a recognizable BH1 sequence motif (Figure 4A,B). Furthermore, loss of this ionic interaction substantially impacts SPPV14 affinities for BH3 peptide of proapoptotic host Bcl-2 proteins (Table 1), epitomising the behaviour of endogenous mammalian prosurvival Bcl-2 interactions [43].

The SPPV14:Bax BH3 complex revealed that an additional residue, M74, engages a fifth pocket in the canonical ligand binding groove besides the four canonical hydrophobic

residues that are already accommodated in their respective pockets. This observation mirrors findings previous findings for Bcl-x_L and Mcl-1 [44]. A further example of additional hydrophobic residues from a BH3 motif peptide engaging a pocket is the variola virus F1L:Bid BH3 complex, where I101 from Bid protrudes into a fifth pocket [45]. A comparison of the detailed architecture of these additional pockets (Figure 4C-F) reveals that there is little conservation of residues that form the pockets. In Mcl-1, V216, F318 and F319 are involved, whereas in Bcl-x_L it is A93 and Y195. In variola virus F1L, I101 from the Bid BH3 motif engages a fifth pocket formed by Y111, L154, L158 and I200. Interestingly, mutations affecting the binding of Bax M74 to Bcl-x_L or Mcl-1 resulted in increased sensitivity to the BH3 mimetic drug ABT-737 [44]. However, the functional consequences of disruption of M74 mediated binding of Bax to SPPV14 remains to be established.

Previously several residues were predicted to be located in the binding groove of SPPV14 based on the M11L structure, and thus designated as potential mediators of interactions with BH3 motif ligands [25]. A comparison of the predicted structure with our experimentally determined structure reveals that out of the six predicted residues, all six were indeed found in the SPPV14 binding groove, revealing the high degree of similarity of between the SPPV14 and M11L canonical ligand binding grooves despite an overall sequence identity of 22% [25]. However, whilst six key residues in the binding groove are conserved, the remaining additional BH3 motif contacting residues influence the selectivity or affinity of the viral Bcl-2 for its mammalian targets, as seen by the differential ligand binding profiles and affinities of SPPV14 and M11L.

In summary we show that SPPV14 is a monomeric prosurvival Bcl-2 protein that utilizes the canonical ligand binding groove to engage BH3 motif peptides of proapoptotic Bcl-2 proteins. Our structures provide a mechanistic basis to delineate detailed intermolecular

interactions and the importance of neutralizing different proapoptotic Bcl-2 proteins for sheeppoxvirus infectivity and proliferation.

Funding

This research was funded by the Australian Research Council (Fellowship FT130101349 to MK) and La Trobe University (scholarship to CDS).

Author contributions

CDS designed and performed experiments, analyzed data and wrote the manuscript. DRB performed experiments and commented on the manuscript. MGH conceived the study, analyzed data and wrote the manuscript. MK conceived the study, designed and performed experiments, analyzed data and wrote the manuscript.

Acknowledgments

We thank staff at the MX beamlines at the Australian Synchrotron for help with X-ray data collection. We thank the ACRF for their support of the Eiger MX detector at the Australian Synchrotron MX2 beamline and the Comprehensive Proteomics Platform at La Trobe University for core instrument support.

Conflicts of Interest

The authors declare no conflict of interest

Tables

Table 1: Interactions of wild-type sheeppoxvirus SPPV14 and two mutants R84A and Y46A with human pro-apoptotic BH3 motif peptides. Measurements were performed using isothermal titration calorimetry. All K_D values (in nM) are the means of three independent measurements with standard error. "NB" denotes no binding.

Peptide	WT KD (nM)	R84A KD (nM)	Y46A KD (nM)
Bak	48 ± 1	139 ± 10	2553 ± 353
Bax	26 ± 3	182 ± 15	4645 ± 393
Bok	7580± 643	NB	NB
Bad	5197± 533	6700 ± 806	NB
Bid	136± 23	1698 ± 16	NB
Bik	1766± 280	9245 ± 701	NB
Bim	19± 3	44 ± 3	165 ± 18
Bmf	44± 7	97 ± 2	151 ± 18
Hrk	39± 5	132 ± 3	2250 ± 318
Noxa	NB	NB	NB
Puma	56± 8	95 ± 3	988 ± 142

Table 2: X-ray data collection and refinement statistics.

	SPPV14: Hrk BH3	SPPV14: Bax BH3
Data collection		
Space group	$C222_1$	$P2_1$
Cell dimensions		
$\quad$a, b, c (Å)	43.28, 70.66, 115.08	100.28, 78.56, 107.77
$\quad\alpha, \beta, \gamma$ (°)	90, 90, 90	90, 110.97, 90
Wavelength (Å)	0.9537	0.9537
Resolution (Å)	31.07 - 2.05 (2.12 - 2.05) *	37.61 - 2.91 (3.02 - 2.9) *
R_{sym} or R_{merge}	0.066 (1.12)	0.15 (1.24)
$I / \sigma I$	8.7 (0.9)	5.2 (0.9)
Completeness (%)	98.2 (90.3)	99.6 (98.3)
$CC_{1/2}$	0.99 (0.59)	0.99 (0.54)
Redundancy	5.0 (4.7)	3.5 (3.6)
Refinement		
Resolution (Å)	31.07 - 2.05 (2.12 - 2.05)	37.61 - 2.91 (3.02 - 2.9)
No. reflections	11232	34323
R_{work} / R_{free}	0.221/0.235	0.238/0.288
Clashscore	0.40	5.14
No. atoms		
$\quad$Protein	1235	10281
$\quad$Ligand/ion	0	0
$\quad$Water	28	44
B-factors		
$\quad$Protein	59.40	66.91
$\quad$Ligand/ion	0	0
$\quad$Water	52.66	46.66
R.m.s. deviations		
$\quad$Bond lengths (Å)	0.003	0.004
$\quad$Bond angles (°)	0.45	0.63

$*$ Values in parentheses are for the highest resolution shell.

References

[1] Galluzzi L, Brenner C, Morselli E, Touat Z, Kroemer G. Viral control of mitochondrial apoptosis. PLoS Pathog. 2008;4:e1000018.

[2] Kvansakul M, Caria S, Hinds MG. The Bcl-2 Family in Host-Virus Interactions. Viruses. 2017;9:290.

[3] Adams JM, Cory S. The Bcl-2 protein family: arbiters of cell survival. Science. 1998;281:1322-6.

[4] Banjara SS, C.D.; Hinds, M.G.; Kvansakul, M. The Bcl-2 Family: Ancient Origins, Conserved Structures, and Divergent Mechanisms. Biomolecules. 2020;10:128.

[5] Kvansakul M, Hinds MG. The structural biology of BH3-only proteins. Methods Enzymol. 2014;544:49-74.

[6] Shamas-Din A, Kale J, Leber B, Andrews DW. Mechanisms of action of bcl-2 family proteins. Cold Spring Harb Perspect Biol. 2013;5:a008714.

[7] Kvansakul M, Hinds MG. Structural biology of the Bcl-2 family and its mimicry by viral proteins. Cell Death Dis. 2013;4:e909.

[8] White E, Sabbatini P, Debbas M, Wold WS, Kusher DI, Gooding LR. The 19-kilodalton adenovirus E1B transforming protein inhibits programmed cell death and prevents cytolysis by tumor necrosis factor alpha. Mol Cell Biol. 1992;12:2570-80.

[9] Neilan JG, Lu Z, Afonso CL, Kutish GF, Sussman MD, Rock DL. An African swine fever virus gene with similarity to the proto-oncogene bcl-2 and the Epstein-Barr virus gene BHRF1. J Virol. 1993;67:4391-4.

[10] Banjara S, Caria S, Dixon LK, Hinds MG, Kvansakul M. Structural Insight into African Swine Fever Virus A179L-Mediated Inhibition of Apoptosis. J Virol. 2017;91:e02228-16.

[11] Henderson S, Huen D, Rowe M, Dawson C, Johnson G, Rickinson A. Epstein-Barr virus-coded BHRF1 protein, a viral homologue of Bcl-2, protects human B cells from programmed cell death. Proc Natl Acad Sci U S A. 1993;90:8479-83.

[12] Kvansakul M, Wei AH, Fletcher JI, Willis SN, Chen L, Roberts AW, et al. Structural basis for apoptosis inhibition by Epstein-Barr virus BHRF1. PLoS Pathog. 2010;6:e1001236.

[13] Lin PW, Huang YJ, John JA, Chang YN, Yuan CH, Chen WY, et al. Iridovirus Bcl-2 protein inhibits apoptosis in the early stage of viral infection. Apoptosis. 2008;13:165-76.

[14] Banjara S, Mao J, Ryan TM, Caria S, Kvansakul M. Grouper iridovirus GIV66 is a Bcl-2 protein that inhibits apoptosis by exclusively sequestering Bim. J Biol Chem. 2018;293:5464-77.

[15] Wasilenko ST, Stewart TL, Meyers AF, Barry M. Vaccinia virus encodes a previously uncharacterized mitochondrial-associated inhibitor of apoptosis. Proc Natl Acad Sci U S A. 2003;100:14345-50.

[16] Campbell S, Thibault J, Mehta N, Colman PM, Barry M, Kvansakul M. Structural insight into BH3 domain binding of vaccinia virus antiapoptotic F1L. J Virol. 2014;88:8667-77.

[17] Graham KA, Opgenorth A, Upton C, McFadden G. Myxoma virus M11L ORF encodes a protein for which cell surface localization is critical in manifestation of viral virulence. Virology. 1992;191:112-24.

[18] Kvansakul M, van Delft MF, Lee EF, Gulbis JM, Fairlie WD, Huang DC, et al. A structural viral mimic of prosurvival Bcl-2: a pivotal role for sequestering proapoptotic Bax and Bak. Mol Cell. 2007;25:933-42.

[19] Banadyga L, Lam SC, Okamoto T, Kvansakul M, Huang DC, Barry M. Deerpox virus encodes an inhibitor of apoptosis that regulates Bak and Bax. J Virol. 2011;85:1922-34.

[20] Burton DR, Caria S, Marshall B, Barry M, Kvansakul M. Structural basis of Deerpox virus-mediated inhibition of apoptosis. Acta Crystallogr D Biol Crystallogr. 2015;71:1593-603.

[21] Banadyga L, Gerig J, Stewart T, Barry M. Fowlpox virus encodes a Bcl-2 homologue that protects cells from apoptotic death through interaction with the proapoptotic protein Bak. J Virol. 2007;81:11032-45.

[22] Anasir MI, Caria S, Skinner MA, Kvansakul M. Structural basis of apoptosis inhibition by the fowlpox virus protein FPV039. J Biol Chem. 2017;292:9010-21.

[23] Westphal D, Ledgerwood EC, Hibma MH, Fleming SB, Whelan EM, Mercer AA. A novel Bcl-2-like inhibitor of apoptosis is encoded by the parapoxvirus ORF virus. J Virol. 2007;81:7178-88.

[24] Suraweera CD, Ishtiaq Anasir M, Chugh S, Javorsky A, Impey RA, Hasan Zadeh M, et al. Structural insight into tanapoxvirus mediated inhibition of apoptosis. BioRxiv. 2020:https://doi.org/10.1101/2020.01.21.914671.

[25] Okamoto T, Campbell S, Mehta N, Thibault J, Colman PM, Barry M, et al. Sheeppox virus SPPV14 encodes a Bcl-2-like cell death inhibitor that counters a distinct set of mammalian proapoptotic proteins. J Virol. 2012;86:11501-11.

[26] Okamoto T, Campbell S, Mehta N, Thibault J, Colman PM, Barry M, et al. Sheeppox Virus SPPV14 Encodes a Bcl-2-Like Cell Death Inhibitor That Counters a Distinct Set of Mammalian Proapoptotic Proteins. Journal of Virology. 2012;86:11501-11.

[27] Caria S, Hinds MG, Kvansakul M. Structural insight into an evolutionarily ancient programmed cell death regulator - the crystal structure of marine sponge BHP2 bound to LB-Bak-2. Cell Death Dis. 2017;8:e2543.

[28] Kvansakul M, Czabotar PE. Preparing Samples for Crystallization of Bcl-2 Family Complexes. In: Puthalakath H, Hawkins CJ, editors. Programmed Cell Death: Methods and Protocols. New York, NY: Springer New York; 2016. p. 213-29.

[29] Aragao D, Aishima J, Cherukuvada H, Clarken R, Clift M, Cowieson NP, et al. MX2: a high-flux undulator microfocus beamline serving both the chemical and macromolecular crystallography communities at the Australian Synchrotron. Journal of Synchrotron Radiation. 2018;25:885-91.

[30] Kabsch W. XDS. Acta Crystallographica Section D. 2010;66:125-32.

[31] Evans P. Scaling and assessment of data quality. Acta Crystallographica Section D. 2006;62:72-82.

[32] Long F, Vagin AA, Young P, Murshudov GN. BALBES: a molecular-replacement pipeline. Acta Crystallogr D Biol Crystallogr. 2008;64:125-32.

[33] Kvansakul M, van Delft MF, Lee EF, Gulbis JM, Fairlie WD, Huang David CS, et al. A Structural Viral Mimic of Prosurvival Bcl-2: A Pivotal Role for Sequestering Proapoptotic Bax and Bak. Molecular Cell. 2007;25:933-42.

[34] Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta Crystallographica Section D. 2010;66:486-501.

[35] Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, et al. Towards automated crystallographic structure refinement with phenix.refine. Acta Crystallographica Section D. 2012;68:352-67.

[36] McCoy A. Solving structures of protein complexes by molecular replacement with Phaser. Acta Crystallographica Section D. 2007;63:32-41.

[37] Meyer PA, Socias S, Key J, Ransey E, Tjon EC, Buschiazzo A, et al. Data publication with the structural biology data grid supports live analysis. Nat Commun. 2016;7:10882.

[38] Morin A, Eisenbraun B, Key J, Sanschagrin PC, Timony MA, Ottaviano M, et al. Collaboration gets the most out of software. eLife. 2013;2:e01456.

[39] Nichols DB, De Martini W, Cottrell J. Poxviruses Utilize Multiple Strategies to Inhibit Apoptosis. Viruses. 2017;9.

[40] Banjara S, Shimmon GL, Dixon LK, Netherton CL, Hinds MG, Kvansakul M. Crystal Structure of African Swine Fever Virus A179L with the Autophagy Regulator Beclin. Viruses. 2019;11:789.

[41] Liu X, Dai S, Zhu Y, Marrack P, Kappler JW. The structure of a Bcl-xL/Bim fragment complex: implications for Bim function. Immunity. 2003;19:341-52.

[42] Kvansakul M, Hinds MG. The Bcl-2 family: structures, interactions and targets for drug discovery. Apoptosis. 2015;20:136-50.

[43] Sattler M, Liang H, Nettesheim D, Meadows RP, Harlan JE, Eberstadt M, et al. Structure of Bcl-xL-Bak peptide complex: recognition between regulators of apoptosis. Science. 1997;275:983-6.

[44] Czabotar PE, Lee EF, Thompson GV, Wardak AZ, Fairlie WD, Colman PM. Mutation to Bax beyond the BH3 domain disrupts interactions with pro-survival proteins and promotes apoptosis. J Biol Chem. 2011;286:7123-31.

[45] Marshall B, Puthalakath H, Caria S, Chugh S, Doerflinger M, Colman PM, et al. Variola virus F1L is a Bcl-2-like protein that unlike its vaccinia virus counterpart inhibits apoptosis independent of Bim. Cell Death Dis. 2015;6:e1680.

Figures and figure legends

SPPV14: Hrk BH3

SPPV14: Bax BH3

M11L: Bak BH3

Bcl-x_L: Bax BH3

Figure 1: SPPV14 binds to BH3 motif peptides of proapoptotic Bcl-2 proteins using the canonical ligand binding groove (A) Crystal structure of SPPV14:Hrk BH3 complex. SPPV14 is shown as green ribbon, Hrk is shown in pink, with SPPV14 helices labeled α1-7. The view is into the hydrophobic binding groove formed by helices α2-5. **(B)** Crystal structure of SPPV14:Bax BH3 complex. SPPV14 is shown as in (A), Bax BH3 is shown in slate. The view is as in (A). **(C)** Cartoon of M11L:Bak BH3 complex (PDB ID 2JBY) [18]. M11L is shown in light blue, Bak is shown in red. **(D)** Cartoon of Bcl-x$_L$:Bax BH3 complex (PDB ID 1BXL) [43]. Bcl-x$_L$ is shown as sand coloured ribbon, Bax is shown in slate. The views in C-D are as in (A).

Figure 2: Electron density maps of SPPV14:Hrk and Bax BH3 complexes (A) 2Fo-Fc electron density maps of SPPV14:Hrk BH3 complex interface contoured at 1.5 σ. SPPV14 is shown as green sticks and Hrk as pink coloured sticks. **(B)** 2Fo-Fc electron density maps of SPPV14:Bax BH3 complex interface contoured at 1.5 σ. SPPV14 is shown as green sticks and Bax as slate coloured sticks. The N and C-termini of the BH3 peptide are labelled.

Figure 3: Detailed view of the SPPV14: Hrk BH3, SPPV14: Bax BH3, M11L: Bak BH3 and Bcl-x$_L$: Bax BH3 interfaces. (**A**) The SPPV14 backbone, floor of the binding groove and surface are shown in green, yellow and grey respectively, whilst the Hrk BH3 ribbon is shown in hot pink. The three hydrophobic residues of Hrk (L37, L40 and L44) as well as T35 are protruding into the binding groove, and conserved ionic interaction formed by SPPV14 R84 and Hrk BH3 D42 are labelled, as well as all other residues involved in hydrogen bonds between protein and peptide. (**B**) SPPV14 is shown as in (A) whilst the Bax BH3 ribbon is shown in slate. The five hydrophobic residues of Bax BH3 (L59, L63, I66, L70 and M74) are protruding into the binding groove, and conserved salt bridge interaction formed by SPPV14 R84 and Bax BH3 D68 are labelled, as well as other residues involved in additional ionic interactions and hydrogen bonds. (**C**) M11L (light blue) is shown as in (C) whilst the Bak BH3 ribbon is shown in red (PDB ID 1JBY). The four hydrophobic residues of Bak BH3 (L74, L78, I81 and I85) are protruding into the binding groove, and the ionic interaction formed by M11L K126 and Bak BH3 D90 is labelled, as well as other residues involved in

protein:peptide hydrogen bonds. **(D)** Bcl-x$_L$ (wheat) is shown as in (A) whilst the Bax BH3 is shown as slate coloured sticks (PDB ID 1BXL). The five hydrophobic residues of Bax BH3 (L59, L63, I66, L70 and M74) are protruding into the binding groove, and the conserved ionic interaction formed by Bcl-x$_L$ R139 and Bax BH3 D68 is labelled, as well as other residues involved in additional ionic interactions and hydrogen bonds. Ionic interactions and hydrogen bonds are shown as dotted black lines and the participating residues are labelled.

Figure 4: Non-canonical interactions in SPPV14:Bax BH3 complex. (A) SPPV14 R84 (green) forms an ionic bond with Bax BH3 D68 (slate) that resembles the canonical ionic interaction between Bcl-x$_L$ R139 (sand) and Bax BH3 D68 (slate). For clarity only Bax BH3 from the SPPV14:Bax BH complex is shown. (B) Sequence alignment of SPPV14, M11L and Bcl-x$_L$ BH1 motifs. Arg residues involved in the ionic interactions with Bax BH3 shown in (A) are coloured. (C) Bax M74 protrudes into a fifth pocket in the SPPV14 binding groove formed by C38, V41, I42, F133 and N137. (D) Bax M74 bound in the equivalent pocket in Bcl-x$_L$ [44]. (E) Bax M74 bound in the equivalent pocket in Mcl-1 [44]. (F) Bid I101 bound in the equivalent pocket in variola virus F1L [45].