Microbial function and genital inflammation in young South African women at high risk of HIV infection

Arghavan Alisoltani[1,2], Monalisa T. Manhanzva[1], Matthys Potgieter[3,4], Christina Balle[5], Liam Bell[6], Elizabeth Ross[6], Arash Iranzadeh[3], Michelle du Plessis[6], Nina Radzey[1], Zac McDonald[6], Bridget Calder[4], Imane Allali[3,7], Nicola Mulder[3,8,9], Smritee Dabee[1,10], Shaun Barnabas[1], Hoyam Gamieldien[1], Adam Godzik[2], Jonathan M. Blackburn[4,8], David L. Tabb[8,11,12], Linda-Gail Bekker[8,13], Heather B. Jaspan[5,8,10], Jo-Ann S. Passmore[1,8,14,15], Lindi Masson[1,8,14,16, 17*]

[1]Division of Medical Virology, Department of Pathology, University of Cape Town, Cape Town 7925, South Africa; [2]Division of Biomedical Sciences, University of California Riverside School of Medicine, Riverside, CA 92521, USA; [3]Computational Biology Division, Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town 7925, South Africa; [4]Division of Chemical and Systems Biology, Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town 7925, South Africa. [5]Division of Immunology, Department of Pathology, University of Cape Town, Cape Town 7925, South Africa; [6]Centre for Proteomic and Genomic Research, Cape Town 7925, South Africa; [7]Laboratory of Human Pathologies Biology, Department of Biology and Genomic Center of Human Pathologies, Mohammed V University in Rabat, Morocco; [8]Institute of Infectious Disease and Molecular Medicine (IDM), University of Cape Town, Cape Town 7925, South Africa; [9]Centre for Infectious Diseases Research (CIDRI) in Africa Wellcome Trust Centre, University of Cape Town, Cape Town 7925, South Africa; [10]Seattle Children's Research Institute, University of Washington, WA 98101, USA; [11]Bioinformatics Unit, South African Tuberculosis Bioinformatics Initiative, Stellenbosch University, Stellenbosch 7602, South Africa; [12]DST–NRF Centre of Excellence for Biomedical Tuberculosis Research, Stellenbosch University, Stellenbosch 7602, South Africa; [13]Desmond Tutu HIV Centre, Cape Town 7925, University of Cape Town, South Africa; [14]Centre for the AIDS Programme of Research in South Africa, Durban 4013, South Africa; [15]National Health Laboratory Service, Cape Town 7925, South Africa; [16]Disease Elimination Program, Life Sciences Discipline, Burnet Institute, Melbourne 3004, Australia; [17]Central Clinical School, Monash University, Melbourne 3004, Australia

*Corresponding author: Dr Lindi Masson; 85 Commercial Road, Melbourne, Victoria, Australia 3004; GPO Box 2284, Melbourne, Victoria, Australia 3001; Email: Lindi.Masson@burnet.edu.au

Abstract

Background: Female genital tract (FGT) inflammation is an important risk factor for HIV acquisition.

The FGT microbiome is closely associated with inflammatory profile, however, the relative importance

of microbial activities has not been established. Since proteins are key elements representing actual

microbial functions, this study utilized metaproteomics to evaluate the relationship between FGT

microbial function and inflammation in 113 young and adolescent South African women at high risk of

HIV infection. Women were grouped as having low, medium or high FGT inflammation by K-means

clustering according to pro-inflammatory cytokine concentrations.

Results: A total of 3,186 microbial and human proteins were identified in lateral vaginal wall swabs

using liquid chromatography-tandem mass spectrometry, while 94 microbial taxa were included in the

taxonomic analysis. Both metaproteomics and 16S rRNA gene sequencing analyses showed

increased non-optimal bacteria and decreased lactobacilli in women with FGT inflammatory profiles.

However, differences in the predicted relative abundance of most bacteria were observed between

16S rRNA gene sequencing and metaproteomics analyses. Bacterial protein functional annotations

(gene ontology) predicted inflammatory cytokine profiles more accurately than bacterial relative

abundance determined by 16S rRNA gene sequence analysis, as well as functional predictions based

on 16S rRNA gene sequence data ($p<0.0001$). The majority of microbial biological processes were

underrepresented in women with high inflammation compared to those with low inflammation,

including a *Lactobacillus*-associated signature of reduced cell wall organization and peptidoglycan

biosynthesis. This signature remained associated with high FGT inflammation in a subset of 74

women nine weeks later, was upheld after adjusting for *Lactobacillus* relative abundance, and was

associated with *in vitro* inflammatory cytokine responses to *Lactobacillus* isolates from the same

women. Reduced cell wall organization and peptidoglycan biosynthesis were also associated with

high FGT inflammation in an independent sample of ten women.

Conclusions: Both the presence of specific microbial taxa in the FGT and their properties and

activities are critical determinants of FGT inflammation. Our findings support those of previous studies

suggesting that peptidoglycan is directly immunosuppressive, and identify a possible avenue for

biotherapeutic development to reduce inflammation in the FGT. To facilitate further investigations of

microbial activities, we have developed the FGT-METAP application that is available at

(http://immunodb.org/FGTMetap/).

Keywords

Background

Despite the large reduction in AIDS-related deaths as a result of the expansion of HIV antiretroviral

programs, global HIV incidence has declined by only 16% since 2010 [1]. Of particular concern are

the extremely high rates of HIV infection in young South African women [1]. The behavioral and

biological factors underlying this increased risk are not fully understood, however, a key predisposing

factor that has been identified in this population is genital inflammation [2–4]. We have previously

shown that women with elevated concentrations of pro-inflammatory cytokines and chemokines in

their genital tracts were at higher risk of becoming infected with HIV [2]. Furthermore, the protection

offered by a topical antiretroviral tenofovir microbicide was compromised in women with evidence of

genital inflammation [3].

The mechanisms by which female genital tract (FGT) inflammation increases HIV risk are likely

multifactorial, and increased genital inflammatory cytokine concentrations may facilitate the

establishment of a productive HIV infection by recruiting and activating HIV target cells, directly

promoting HIV transcription and reducing epithelial barrier integrity [5–7]. Utilizing proteomics, Arnold,

et al. showed that proteins associated with tissue remodeling processes were up-regulated in the

FGT, while protein biomarkers of mucosal barrier integrity were down-regulated in individuals with an

inflammatory profile [7]. This suggests that inflammation increases tissue remodeling at the expense

of mucosal barrier integrity, which may in turn increase HIV acquisition risk [7]. Furthermore,

neutrophil-associated proteins, especially certain proteases, were positively associated with

inflammation and may be involved in disrupting the mucosal barrier [7]. This theory was further

validated by the finding that antiproteases associated with HIV resistance were increased in a cohort

of sex workers from Kenya [8].

Bacterial vaginosis (BV), non-optimal cervicovaginal bacteria, and sexually transmitted infections

(STIs), which are highly prevalent in South African women, are likely important drivers of genital

inflammation in this population [9,10]. BV and non-optimal bacteria have been consistently associated

with a marked increase in pro-inflammatory cytokine concentrations [9–12]. Conversely, women who

have "optimal" vaginal microbiota, primarily consisting of *Lactobacillus* spp., have low levels of genital

inflammation and a reduced risk of acquiring HIV [13]. Lactobacilli and the lactic acid that they

produce may actively suppress inflammatory responses and thus may play a significant role in modulating immune profiles in the FGT [14–16]. However, partly due to the complexity and diversity of the microbiome, the immunomodulatory mechanisms of specific vaginal bacterial species are not fully understood [17]. Further adding to this complexity is the fact that substantial differences in the cervicovaginal microbiota exist by geographical location and ethnicity and that the properties of different strains within particular microbial species are also highly variable [15,18].

In addition to providing insight into host mucosal barrier function, metaproteomic studies of the FGT can evaluate microbial activities and functions that may influence inflammatory profiles [19]. A recent study highlighted the importance of vaginal microbial function by demonstrating that FGT bacteria, such as *Gardnerella vaginalis,* modulate the efficacy of the tenofovir microbicide by active metabolism of the drug [20]. Additionally, Zevin et al. found that bacterial functional profiles were associated with epithelial barrier integrity and wound healing in the FGT [21], suggesting that microbial function may also be closely linked to genital inflammation. The aim of the present study was to utilize both metaproteomics and 16S rRNA gene sequencing to improve our understanding of the relationship between microbial function and inflammatory profiles in the FGTs of South African women.

Results

This study included 113 young and adolescent HIV-uninfected women (aged 16-22 years) residing in Cape Town, South Africa [22]. Seventy-four of these women had samples and metadata available for analysis at a second time-point nine weeks later. Ten women had samples available at the second time-point, but not the first, and were included to validate the functional signature identified in the primary analysis of this study. STI and BV prevalence were high in this cohort, with 48/113 (42%) of the women having at least one STI and 56/111 (50%) of the women being BV positive by Nugent scoring (Additional file 1: Table S1). The use of injectable contraceptives, BV, and chlamydia were associated with increased genital inflammatory cytokines, as previously described [10,23].

FGT metaproteome associates with BV and inflammatory cytokine profiles

Liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis was conducted on the total protein acquired from lateral vaginal wall swabs for all participants. Using principal component analysis (PCA), it was found that women with BV clustered separately from women who were BV negative according to the relative abundance of all host and microbial proteins identified (Additional file 1: Fig. S1a). Women who were considered to have high levels of genital inflammation based on pro-inflammatory cytokine concentrations (Additional file 1: Fig. S2) tended to cluster separately from women with low inflammation, while no clear grouping was observed by STI status or chemokine profiles (Additional file 1: Fig. S1b-d). After adjusting for potentially confounding variables including age, contraceptives, prostate specific antigen (PSA) and co-infections, BV was associated with the largest number of differentially abundant proteins, followed by inflammatory cytokine profiles, while fewer associations were observed between chemokine profiles and protein relative abundance and none between STI status and protein relative abundance (Additional file 1: Fig. S1e). Individual STIs were also not associated with marked changes in the metaproteome, with *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *Trichomonas vaginalis* and *Mycoplasma genitalium* associated with significant changes in zero, five, seven and eleven proteins, respectively (after adjustment for potential confounders and multiple comparisons). However, this stratified analysis is limited by the small number of STI cases for some infections and the prevalence of co-infections.

Differences in taxonomic assignment using metaproteomics and 16S rRNA gene sequencing

Of the proteins identified in lateral vaginal wall swab samples, 38.8% were human, 55.8% were

bacterial, 2.7% fungal, 0.4% archaeal, 0.09% of viral origin, and 2% grouped as "other" (not shown).

However, as the majority of taxa identified had ≤2 proteins detected, a more stringent cut-off was

applied to include only taxa with ≥3 detected proteins, or 2 proteins detected in multiple taxa. The final

curated dataset included 44% human, 55% bacterial (n=81 taxa), and 1% fungal proteins (n=13 taxa)

(Fig. 1a-d; Additional file 2: Table S2). When the relative abundance of the most abundant bacterial

taxa identified using metaproteomics was compared to the relative abundance of the most abundant

bacteria identified using 16S rRNA gene sequencing, a large degree of similarity was found at the

genus level. A total of 6/9 of the genera identified using 16S rRNA gene sequence analysis were also

identified using metaproteomics (Fig. 1e, f; Additional file 1: Fig. S3). As expected, both approaches

showed that multiple *Lactobacillus* species were less abundant in women with high versus low

inflammation, while BV-associated bacteria (including *G. vaginalis*, *Prevotella* species, *Megasphaera*

species, *Sneathia amnii* and *Atopobium vaginae*) were more abundant in women with high

inflammation compared to those with low inflammation (Additional file 2: Table S3**)**.   However,

species-level annotation differed between the 16S rRNA gene sequencing and metaproteomics

analyses, with metaproteomics identifying more lactobacilli at the species level, while

Lachnovaginosum genomospecies (previously known as BVAB1) was not identified using

metaproteomics. Additionally, the relative abundance of the taxa differed between 16S rRNA gene

sequencing and metaproteomics analyses.

Protein profiles differ between women defined as having high, medium or low FGT

inflammation

A total of 449, 165, and 39 host and microbial proteins were differentially abundant between women

with high versus low inflammation, medium versus high inflammation and medium versus low

inflammation, respectively (Additional file 2: Table S4). Microbial proteins that were more abundant in

women with high or medium inflammation versus low inflammation were mostly assigned to non-

optimal bacterial taxa (including *G. vaginalis*, *Prevotella* species, *Megasphaera* species, *S. amnii* and

A. vaginae). Less abundant microbial proteins in women with high inflammation were primarily of

Lactobacillus origin (Additional file 1: Fig. S4). Of the human proteins that were differentially abundant

according to inflammation status, 93 were less abundant and 132 were more abundant in women with high compared to those with low inflammation. Human biological process gene ontologies (GOs) that were significantly underrepresented in women with high versus low inflammation included positive regulation of apoptotic signaling, establishment of endothelial intestinal barrier, ectoderm development and cornification [false discovery rate (FDR) adj. p<0.0001 for all; Additional file 1: Fig. S5]. Significantly overrepresented pathways in women with high versus low inflammation included multiple inflammatory processes - such as chronic response to antigenic stimulus, positive regulation of IL-6 production and inflammatory response (FDR adj. p<0.0001 for all; Additional file 1: Fig. S5).

Weighted correlation network analysis of microbial and host proteins identified five modules (clusters) representing co-correlations between microbial and host proteins (Fig. 2a, b). The yellow module included primarily *L. iners* proteins, while the turquoise module primarily consisted of *L. crispatus* and host proteins. The grey and brown modules consisted entirely of host proteins and the blue module included non-optimal bacteria and host proteins. Pro-inflammatory cytokines correlated inversely with the *Lactobacillus* modules (yellow and turquoise) and positively with the grey, brown and blue modules. Chemokines were significantly positively correlated with the brown and grey modules and inversely with the turquoise module (Fig. 2c). Blue, yellow, turquoise and grey modules correlated with BV Nugent score, while BV showed no significant correlation with the brown module (Fig. 2d). The finding that the brown module did not include any microbial proteins and was not associated with BV or STIs, suggests the presence of inflammatory processes independent of the microbiota. For STIs, no significant correlations (p-value <0.05) with any of the five modules were detected (Fig. 2d). To determine whether the co-correlations of microbial proteins with host proteins had functional meaning, we profiled the top biological processes of proteins as shown in the row sidebar in Fig. 2a. Host proteins that correlated negatively with the blue module (consisting of non-optimal bacteria) were mostly involved in cell adhesion, cell-cell adhesion, cell-cell junction assembly, and regulation of cell-adhesion pathways. This suggests reduced epithelial barrier function associated with *G. vaginalis*, *Prevotella* spp., *Megasphaera,* and *A. vaginae*. Host proteins involved in immune system and inflammatory response pathways were positively correlated with brown and grey modules and inversely associated with the turquoise module including *L. crispatus*. Despite variability in the microbial proteins and taxa, redundant microbial functional pathways were identified across different

modules, indicating that microbiota share a set of common metabolic functions (e.g. glucose

metabolism, protein translation and synthesis), as expected. Although most lactobacilli proteins were

negatively associated with BV and pro-inflammatory cytokines (see turquoise and yellow modules in

Fig. 2a), almost all *L. iners* proteins were clustered in the yellow module with slightly different

functional pathways from lactobacilli proteins in the turquoise module.

Microbial function predicts genital inflammation with greater accuracy than taxa relative

abundance

The accuracies of (i) microbial relative abundance (determined using 16S rRNA gene sequencing), (ii)

functional prediction based on 16S rRNA gene sequence data, (iii) microbial protein relative

abundance (determined using metaproteomics) and (iv) microbial molecular function (determined by

aggregation of GOs of proteins identified using metaproteomics) for prediction of genital inflammation

status (low, medium and high groups) were evaluated using random forest analysis. The three most

important species predicting genital inflammation grouping by 16S rRNA gene sequence data were

Sneathia sanguinegens, *A. vaginae,* and *Lactobacillus iners* (Fig. 3a). The top functional pathways

based on 16S rRNA gene sequence data for the identification of women with inflammation were

prolactin signaling, biofilm formation, and tyrosine metabolism (Fig. 3b). Similarly, the most important

proteins predicting inflammation grouping by metaproteomics included two *L. iners* proteins

(chaperone protein DnaK and pyrophosphate phospho-hydrolase), two *A. vaginae* proteins

(glyceraldehyde-3-phosphate dehydrogenase and phosphoglycerate kinase), *Megasphaera*

genomosp. type 1 cold-shock DNA-binding domain protein, and *L. crispatus* ATP synthase subunit

alpha (Fig. 3c). The top molecular functions associated with genital inflammation included

oxidoreductase activity (of multiple proteins expressed primarily by lactobacilli, but also *Prevotella* and

Atopobium species), beta-phosphoglucomutase activity (expressed by lactobacilli) and GTPase

activity (of multiple proteins produced by mainly *Prevotella, G. vaginalis*, and some lactobacilli) as

illustrated in Fig. 3d. These findings suggest that the functions of key BV-associated bacteria and

lactobacilli are critical for determining the level of genital inflammation. It was also found that the out-

of-bag (OOB) error rate distribution was significantly lower for molecular function, followed by protein

relative abundance, functional prediction based on 16S rRNA gene sequence data and then taxa

relative abundance (Fig. 3e). This suggests that the activities of the bacteria present in the FGT play a

critical role in driving genital inflammation, in addition to the presence and relative abundance of

particular taxa.

Differences in microbial stress response, lactate dehydrogenase and metabolic pathways

between women with high versus low FGT inflammation

Microbial functional profiles were further investigated by comparing molecular functions, biological

processes and cellular component GO enrichment for detected proteins between women with low

versus high inflammation using differential expression analysis, adjusting for potentially confounding

variables including age, contraceptives, PSA and STIs (Fig. 4; Additional file 2: Table S5). The

majority of microbial molecular functions (90/107), biological processes (57/83) and cellular

components (14/23) were underrepresented in women with high compared to low genital

inflammation, with most of the differentially abundant metabolic processes, including peptide,

nucleoside, glutamine, and glycerophospholipid metabolic processes, decreased in women with

inflammation. The phosphoenolpyruvate-dependent sugar phosphotransferase system (a major

microbial carbohydrate uptake system including proteins assigned to *Lactobacillus* species, *S. amnii*,

A. vaginae, and *G. vaginalis*) was also underrepresented in women with evidence of genital

inflammation compared to women with low inflammation (FDR adj. p<0.0001; Fig. 4a). Additionally, L-

lactate dehydrogenase activity was inversely associated with inflammation (Additional file 2: Table S5)

and both large and small ribosomal subunit cellular components were underrepresented in those with

high FGT inflammation (Fig. 4b). The only metabolic processes that were overabundant in women

with high versus low inflammation were malate and ATP metabolic processes. Overrepresented

microbial biological processes in women with genital inflammation also included response to oxidative

stress and cellular components associated with cell division (e.g. mitotic spindle midzone) and

ubiquitination (e.g. Doa10p ubiquitin ligase complex, Hrd1p ubiquitin ligase ERAD-L complex, RQC

complex; Fig. 4**)**.

Underrepresented *Lactobacillus* peptidoglycan, cell wall and membrane pathways in women

with high versus low FGT inflammation

Among the top biological processes inversely associated with inflammation were cell wall

organization, regulation of cell shape and peptidoglycan biosynthetic process (all adj. p≤0.0001; Fig.

4a). The GO terms, regulation of cell shape, and peptidoglycan biosynthetic process included identical proteins (e.g. D-alanylalanine synthetase, D-alanyl-D-alanine-adding enzyme and Bifunctional protein GlmU), while cell wall organization included the same proteins plus D-alanyl carrier protein. Although proteins involved in these processes were only detected for *Lactobacillus* species and may thus be biomarkers of the increased relative abundance of lactobacilli, each remained significantly associated with inflammation after adjusting for the relative abundance of *L. iners* and non-*iners Lactobacillus* species (determined using 16S rRNA gene sequencing), as well as STI status, PSA detection and hormonal contraceptive use [beta-coefficient: -1.68; 95% confidence interval (CI): -2.84 to -0.52; p=0.004 for cell wall organization; beta-coefficient: -1.71; 95% CI: -2.91 to -0.53; p=0.005 for both regulation of cell shape and peptidoglycan biosynthetic process]. This suggests that the relationship between these processes and inflammation is independent of the relative abundance of *Lactobacillus* species and that the cell wall and membrane properties of lactobacilli may play a role in modulating FGT inflammation. Furthermore, we found that multiple cell wall and membrane cellular components, including the S-layer, peptidoglycan-based cell wall, cell wall and membrane, were similarly underrepresented in women with high versus low FGT inflammation (Fig. 4b). Using metaproteomic data from an independent group of 10 women, we similarly found that peptidoglycan biosynthetic process, cell wall organization, and regulation of cell shape GOs were underrepresented in women with high inflammation (Fig. 5a).

To further investigate whether the associations between FGT inflammation and *Lactobacillus* biological processes and cellular components were independent of *Lactobacillus* relative abundance, we compared these GOs in BV negative women with *Lactobacillus* dominant communities who were grouped (median splitting of samples based on the first principal component of nine pro-inflammatory cytokine concentrations) as having low (n=20) versus high (n=20) inflammation (Fig. S6). Interestingly, it was found that, even though only non-significant trends towards decreased relative abundance of lactobacilli determined by 16S rRNA gene sequencing (p=0.73) and metaproteomics (p=0.14) were observed, several previously identified GOs, including peptidoglycan-based cell wall (p=0.01), cell wall organization (p=0.03), and S-layer (p=0.01), remained associated with the level of inflammation (Fig. S6). This suggests an association between these biological processes and cellular components that is not fully explained by the relative abundance of the lactobacilli themselves.

***In vitro* confirmation of the role of *Lactobacillus* cell wall and membrane pathways in**

regulating genital inflammation

As previous studies have suggested that the cell wall, as well as the cell membrane, may influence

the immunomodulatory properties of *Lactobacillus* species and that peptidoglycan may be directly

immunosuppressive [14,24], we further investigated the relationships between *Lactobacillus* cell wall

and membrane properties and inflammation using 64 *Lactobacillus* species that were isolated from

the same women [25]. From these lactobacilli, the 22 isolates that induced the lowest levels of pro-

inflammatory cytokine production by vaginal epithelial cells and the 22 isolates that induced the

greatest cytokine responses were selected and their protein profiles evaluated using LC-MS/MS. Cell

wall organization, regulation of cell shape, and peptidoglycan biosynthetic process that were

associated with genital inflammation in the FGT metaproteomic analysis of vaginal swab samples

were also significantly overrepresented in isolates that induced low compared to high levels of

inflammatory cytokines *in vitro* (Fig. 5b-d). Furthermore, plasma membrane, cell wall and membrane

cellular components were similarly overrepresented in isolates that induced low versus high levels of

inflammation (Fig. 5e-i). The cell wall cellular component included multiple proteins involved in

adhesion (adhesion exoprotein, LPXTG-motif cell wall anchor domain protein, collagen-binding

protein, mannose-specific adhesin, putative mucin binding protein, sortase-anchored surface protein

and surface anchor protein). The relative abundance of this cell wall cellular component correlated

positively with the level of adhesion of the isolates to vaginal epithelial cells *in vitro* (rho=0.30,

p=0.0476), which in turn correlated inversely with *in vitro* inflammatory cytokine production by these

cells in response to the isolates [25]. Importantly, the cell wall organization biological process included

multiple proteins involved in peptidoglycan biosynthesis, which has the potential to dampen immune

responses in a strain-dependent manner [14,26].

Changes in FGT metaproteomic profiles over time

To evaluate variations in FGT metaproteomic profiles associated with inflammatory profiles over time,

we compared proteins, taxa and biological process and cellular component GOs between two time-

points nine weeks apart (n=74). The grouping according to inflammatory cytokine profile was

consistent at both visits for 41/74 (55%) participants, including 7/12 (58%) women who received

antibiotic treatment between visits (Fig. 6a). The findings based on the second visit also confirmed the

role of inflammation as one of the drivers of variation in complex metaproteome data, and similar

patterns of data distribution according to inflammation status were observed at both visits (Fig. 6b).

When the proteins, taxa and functions that were most closely associated with inflammation grouping

(high vs. low) at the first visit, were evaluated at the second visit, overall profiles remained similar

between visits (Fig. 6c-e). Significantly lower relative abundance of *Lactobacillus* species and

increased non-optimal bacteria were observed in women with inflammation at the second visit (Fig.

6d). Similarly, cell wall organization, regulation of cell shape, and peptidoglycan biosynthetic process

were among the top biological processes associated with low versus high inflammation at both visits

(for other top GO terms see Fig. 6e). Among women assigned to the same inflammatory profile group

at both visits, the relative abundance of cell wall-related GO terms, malate metabolic process and

response to oxidative stress were highly consistent (Additional file 1: Fig. S7a-c). However, decreased

levels of inflammation between visits were associated with an increase in the relative abundance of

cell wall GOs and decreased response to oxidative stress and malate metabolic process GOs

(Additional file 1: Fig. S7d-f). On the other hand, increased levels of inflammation were associated

with decreased relative abundance of cell wall GOs and increased response to oxidative stress and

malate metabolic process GOs (Additional file1: Fig. S7g-i).

Discussion

Understanding biological risk factors for HIV acquisition is critical for the development of effective HIV prevention strategies. FGT inflammation, defined by elevated inflammatory cytokine levels, has been identified as a key factor associated with HIV infection risk. However, the causes and underlying mechanisms are not fully understood. In this study, we demonstrate that, while the presence of particular microorganisms, including non-optimal bacteria and *Lactobacillus* species, is important for modulating FGT inflammation, their activities and functions are likely of greater importance. We showed that the molecular functions of bacterial proteins predicted FGT inflammation status more accurately than taxa relative abundance determined using 16S rRNA gene sequencing, as well as functional predictions based on 16S rRNA gene sequencing analysis. The majority of microbial biological processes were underrepresented in women with high compared to low inflammation (57/83), including metabolic pathways and a strong signature of reduced *Lactobacillus*-associated cell wall organization and peptidoglycan biosynthesis. This signature remained associated with FGT inflammation after adjusting for the relative abundance of lactobacilli, as well as other potential confounders (such as hormone contraceptive use, semen exposure, STIs), and was also associated with inflammatory cytokine induction in vaginal epithelial cells in response to *Lactobacillus* isolates *in vitro*. We observed a high level of consistency between FGT metaproteomics profiles of the same women at two time points nine weeks apart, particularly among those with the same level of inflammation at both visits. However, cell wall organization and peptidoglycan biosynthesis increased between visits in women who moved from a higher to a lower inflammation grouping and decreased in women who moved from a lower to a higher inflammation grouping.

While we found a large degree of similarity between metaproteomics and 16S rRNA gene sequence taxonomic assignment determined for the same women [27], important differences in the predicted relative abundance of most bacteria were observed. Differences in the relative abundance of certain taxa observed using metaproteomics and 16S rRNA gene sequence data is not surprising as taxonomic assignment from metaproteomic approaches also indicates functional activity (measured by the amount of expressed protein), rather than simply the abundance of a particular bacterial taxon. Additionally, others have noted that 16S rRNA gene sequencing does not have the ability to distinguish between live bacteria and transient DNA [28]. We also observed differences that are likely

due to database limitations. For example, *Lachnovaginosum* genomospecies (previously known as

BVAB1) was not identified in the metaproteomics analysis. This is due to the fact that this species

was not present in the UniProt or NCBI databases [29]. However, the *Clostridium* and *Ruminococcus*

species which were identified are likely to be *Lachnovaginosum* genomospecies since these taxa fall

into the same Clostridiales order. Additionally, *Clostridium* and *Ruminococcus* species were more

abundant in women with high compared to low inflammation, as expected for

Lachnovaginosum genomospecies [10]. Metaproteomic analysis was able to identify lactobacilli to

species level, while sequencing of the V4 region of the 16S rRNA gene had more limited resolution for

this genus, as expected.

The finding that fungal protein relative abundance was substantially lower than bacterial protein

relative abundance is not surprising as it has been estimated that the total number of fungal cells is

orders of magnitude lower than the number of bacterial cells in the human body [30]. Although

Candida proteins were detected, the curated dataset (following removal of taxa that had only 1 protein

detected or 2 proteins detected in only a single sample) did not include any *Candida* species. This

may be due to the relative rarity of fungal cells in these samples and it would be interesting to

evaluate alternative methodology for sample processing to better resolve this population in the future

[31]. A limitation of microbial functional analysis using metaproteomics is that, at present, there is

generally sparse population of functional information in the databases for the majority of the

microbiome. Thus, the findings of this study will be biased toward the microbes that are better

annotated. Nonetheless, microbial function was closely associated with genital inflammatory profiles

and it is possible that this relationship may be even stronger should better annotation exist. Another

limitation of this study is that the *in vitro Lactobacillus* analysis was conducted using a transformed

cell line system that is a greatly simplified model and cannot recapitulate the complexity of the FGT

and the microbiome. It is however significant to note that similar proteomic signatures were noted

both *in vitro* and *in vivo*, regardless of this limitation. To enable researchers to further study

associations between microbial composition and function with BV, STIs, and FGT inflammation by

conducting detailed analysis of specific proteins, taxa and GO terms of interest, we have developed

the "FGT-METAP" online application. The application is available at (http://immunodb.org/FGTMetap/)

and allows users to repeat the analyses presented here, as well as obtain additional information by

analyzing individual proteins, species, and pathways.

A significant functional signature associated with FGT inflammation grouping included *Lactobacillus*-

associated cell wall, peptidoglycan and cell membrane biological processes and cellular components.

Although these functions were linked exclusively to *Lactobacillus* species, the associations with FGT

inflammation were upheld after adjusting for *Lactobacillus* relative abundance determined by 16S

rRNA gene sequencing and were also evident in an analysis including only women with *Lactobacillus*-

dominant microbiota. This profile was confirmed when these GOs were compared between

Lactobacillus isolates that induced high versus low levels of cytokine production *in vitro*. Previous

studies have suggested that the peptidoglycan structure, the proteins present in the cell wall, as well

as the cell membrane, may influence the immunomodulatory properties of *Lactobacillus* species

[14,24]. In a murine model, administration of peptidoglycan extracted from gut lactobacilli was able to

rescue the mice from colitis [26]. This activity was dependent on the NOD2 pathway and correlated

with an upregulation of the indoleamine 2,3-dioxygenase immunosuppressive pathway. The

immunomodulatory properties of peptidoglycan were furthermore dependent on the strain of

lactobacilli from which the peptidoglycan was isolated [26]. The increase in cell wall organization and

peptidoglycan biosynthesis that accompanied a reduction in local inflammation suggests an increase

in *Lactobacillus* strains producing relatively large amounts of proteins involved in these processes. In

a previous study, we showed that adhesion of lactobacilli to vaginal epithelial cells *in vitro* was

inversely associated with cytokine responses, suggesting that direct interaction between the isolates

and vaginal epithelial cells is critical for immunoregulation [25]. In addition to cell wall and membrane

properties, L-lactate dehydrogenase activity was identified as a molecular function inversely

associated with inflammation. Similarly, D-lactate dehydrogenase relative abundance and D-lactate

production by the *Lactobacillus* isolates included in this study were also associated with inflammatory

profiles in an *in vitro* analysis [25]. These findings support the results of previous studies showing that

lactic acid has immunoregulatory properties [16]. Taken together, these findings suggest that

lactobacilli may modulate the inflammatory environment in the FGT through multiple mechanisms. It

was further found that *L. crispatus* was more strongly associated with low inflammation in the

differential abundance analysis and that *L. iners* was the *Lactobacillus* sp. most frequently detected in

women with high inflammation. The co-expression analysis revealed that the majority of *L. crispatus* proteins grouped separately to *L. iners* proteins. However, both the *L. crispatus* and the *L. iners* modules were strongly inversely associated with inflammatory cytokine and chemokine concentrations. This is interesting since *L. crispatus* dominance is considered optimal, while the role of *L. iners,* the most prevalent *Lactobacillus* species in African women [10,11], is poorly understood and it has been associated with compositional instability and transition to a non-optimal microbiota, as well as increased risk of STI acquisition [32,33].

In this study, we observed a similar host proteome profile associated with high FGT inflammation compared to previous studies [7]. Multiple inflammatory pathways were overrepresented in women with high versus low FGT inflammation, while signatures of reduced barrier function were observed in women with high inflammation, with underabundant endothelial, ectoderm and tight junction biological processes. These findings similarly suggest that genital inflammation may be associated with epithelial barrier function.

Conclusions

The link between FGT microbial function and local inflammatory responses described in this study suggests that both the presence of specific microbial taxa in the FGT and their properties and activities likely play a critical role in modulating inflammation. Currently, the annotation of microbial functions is sparse, but with ever-increasing amounts of high-quality, high-throughput data, the available information will improve steadily. The findings of the present study contribute to our understanding of the mechanisms by which the microbiota may influence local immunity, and in turn alter the risk of HIV infection. Additionally, the analyses described herein identify specific microbial properties that may be harnessed for biotherapeutic development.

Methods

Participants

This study included sexually experienced HIV-negative adolescent girls and young women (16-22

years) from the Women's Initiative in Sexual Health (WISH) study in Cape Town, South Africa [22].

While the parent study cohort comprised 149 women, the present sub-study included 113 women who

had both vaginal swabs available for metaproteomics analysis and menstrual cup (MC) cervicovaginal

secretions available for cytokine profiling. Seventy-four of these women had specimens available at a

second time-point 9 weeks later (interquartile range: 9-11 weeks) for metaproteomics analysis. Of

these women, 12 received antibiotic treatment for laboratory-diagnosed or symptomatic STIs and/or

BV (doxycycline, metronidazole, azithromycin, cefixime, ceftriaxone and/or amoxicillin) between study

visits, while 2 received topical antifungal treatment (clotrimazole vaginal cream). Ten additional

women who did not have visit 1 samples available for analysis were included for validation purposes.

The University of Cape Town Faculty of Health Sciences Human Research Ethics Committee (HREC

REF: 267/2013) approved this study. Women ≥18 years provided written informed consent, those <18

years provided informed assent and informed consent was obtained from parents/guardians.

Definition of genital inflammation based on cytokine concentrations

For cytokine measurement, MCs (Softcup®, Evofem Inc, USA) were inserted by the clinician and kept

in place for an hour, after which they were transported to the lab at 4°C and processed within 4 hours

of removal from the participants. MCs were centrifuged and the cervicovaginal secretions were re-

suspended in phosphate buffered saline (PBS) at a ratio of 1ml of mucus: 4 ml of PBS and stored at -

80°C until cytokine measurement. Prior to cytokine measurement, MC secretions were pre-filtered

using 0.2 μm cellulose acetate filters (Sigma-Aldrich, MO, USA). The concentrations of 48 cytokines

were measured in MC samples using Luminex (Bio-Rad Laboratories Inc®, CA, USA) [23]. K-means

clustering was used to identify women with low, medium and high pro-inflammatory cytokine

[interleukin (IL)-1α, IL-1β, IL-6, IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-

related apoptosis-inducing ligand (TRAIL), interferon (IFN)-γ] and chemokine profiles [cutaneous T-

cell-attracting chemokine (CTACK), eotaxin, growth regulated oncogene (GRO)-α, IL-8, IL-16, IFN-γ-

induced protein (IP)-10, monocyte chemoattractant protein (MCP)-1, MCP-3, monokine induced by

IFN-γ (MIG), macrophage inflammatory protein (MIP)-1α, MIP-1β, regulated on activation, normal T

cell expressed and secreted (RANTES)] (Additional file 1: Fig. S2).

STI and BV diagnosis and evaluation of semen contamination

Vulvovaginal swab samples were screened for common STIs, including *Chlamydia trachomatis*,

Neisseria gonorrhoeae, *Trichomonas vaginalis*, *Mycoplasma genitalium,* herpes simplex virus (HSV)

types 1 and 2, *Treponema pallidum* and *Haemophilus ducreyi*, using real-time multiplex PCR assays

[22]. Lateral vaginal wall swabs were collected for BV assessment by Nugent scoring. Blood was

collected for HIV rapid testing (Alere Determine™ HIV-1/2 Ag/Ab Combo, Alere, USA). PSA was

measured in lateral vaginal wall swabs using Human Kallikrein 3/PSA Quantikine ELISA kits (R&D

Systems, MN, USA).

Discovery metaproteomics and analysis

For shotgun LC-MS/MS, lateral vaginal wall swabs were collected, placed in 1 ml PBS, transported to

the laboratory at 4°C and stored at -80°C. Samples were computationally randomized for processing

and analysis. The stored swabs were thawed overnight at 4°C before each swab sample was

vortexed for 30 seconds, the mucus scraped off the swab on the side of the tube and vortexed for an

additional 10 seconds. The samples were then clarified by centrifugation and the protein content of

each supernatant was determined using the Quanti-Pro bicinchoninic acid (BCA) assay kit (Sigma-

Aldrich, MO, USA). Equal protein amounts (100 µg) were denatured with urea exchange buffer,

filtered, reduced with dithiothreitol, alkylated with iodoacetamide and washed with hydroxyethyl

piperazineethanesulfonic acid (HEPES). Acetone precipitation/formic acid (FA; Sigma-Aldrich, MO,

USA) solubilisation was added as a further sample clean-up procedure. The samples were then

incubated overnight with trypsin (Promega, WI, USA) and the peptides were eluted with HEPES and

dried via vacuum centrifugation. Reversed-phase liquid chromatography was used for desalting using

a step-function gradient. The eluted fractions were dried via vacuum centrifugation and kept at -80°C

until analysis. LC-MS/MS analysis was conducted on a Q-Exactive quadrupole-Orbitrap MS (Thermo

Fisher Scientific, MA, USA) coupled with a Dionex UltiMate 3000 nano-UPLC system (120min per

sample). The solvent system included Solvent A: 0.1% Formic Acid (FA) in LC grade water (Burdick

and Jackson, NJ, USA); and Solvent B: 0.1% FA in Acetonitrile (ACN; Burdick & Jackson, NJ, USA).

The MS was operated in positive ion mode with a capillary temperature of 320°C, and 1.95 kV

electrospray voltage was applied. All 187 samples were evaluated in 11 batches over a period of 3

months. A quality control consisting of pooled lateral vaginal swab eluants from the study participants

was run at least twice with every batch. Preliminary quality control analysis was conducted, and

flagged samples/batches were rerun.

Due to the important role of database in metaproteomic analysis, we compared two different

databases to identify proteins in our dataset: (i) UniProt database restricted to human and microbial

entries (73,910,451, release August-2017) and filtered using the MetaNovo pipeline [34]; (ii) vaginal

metagenome-based database with human proteins obtained from the study of Afiuni-Zadeh et al. [35].

The raw data were processed using MaxQuant version 1.5.7.4 against each database, separately.

The detailed parameters are provided in Additional file 1: Table S6. In brief, methionine oxidation and

acetylation of the protein N-terminal amino acid were considered as variable modifications and

carbamidomethyl (C) as a fixed modification. The digestion enzyme was trypsin with a maximum of

two missed cleavages. The taxonomic analysis of the proteins identified using both of the databases

was comparable, however, the first database (filtered UniProt human and microbial entries) resulted

in the identification of a greater number of proteins compared to the vaginal metagenome database,

whilst exhibiting a high degree of similarity compared to the 16S rRNA gene sequence data.

Therefore, the data generated using the first database was used for downstream analysis.

For quality control analysis, raw protein intensities and $\log_{10}$-transformed iBAQ intensities of the

quality control pool, as well as the raw protein intensities of the clinical samples, were compared

between batches (not shown). As variation in raw intensity was observed, all data were adjusted for

batch number prior to downstream analysis. The cleaned and transformed iBAQ values were used to

identify significantly differentially abundant proteins according to inflammatory cytokine profiles, BV

status or STIs using the limma R package [36]. The p-values were obtained from the moderated t-test

after adjusting for confounding variables including age, contraceptives, PSA, and STIs. Proteins with

FDR adjusted p-values ≤ 0.05 and $\log_2$-transformed fold change ≥ 1.2 or ≤ -1.2 were considered as

significantly differentially abundant. To define significantly differentially abundant taxa and GOs,

aggregated iBAQ values of proteins of the same species or GO IDs were compared between women

with low, medium and high inflammation using the limma R package. The aggregation was performed separately for host and microbial proteins. The p-values along with GO IDs were uploaded to REVIGO (http://revigo.irb.hr/) to visualize the top 50 biological processes with the lowest p-values.

The effects of BV, STIs and chemokine and pro-inflammatory cytokine profiles on the metaproteome were investigated by PCA using the mixOmics R package [37]. The ComplexHeatmap package [38] was used to generate heatmaps and cluster the samples and metaproteomes based on the hierarchical and k-means clustering methods. Additionally, R packages EnhancedVolcano, weighted correlation network analysis (WGCNA) [39], and FlipPlots were applied for plotting volcano plots, co-correlation analysis, and Sankey diagrams, respectively. Basic R functions and the ggplot2 R package were used for data manipulation, transformation, normalization and generation of graphics. To identify the key factors distinguishing women defined as having low, medium and high inflammation in their FGTs, we used random forest analysis using the R package randomForest [40] using the following settings: (i) the type of random forest was classification, (ii) the number of trees was 2000, and (iii) the number of variables evaluated at each split was one-third of the total number of variables [proteins/molecular function GO IDs/16S rRNA gene-based operational taxonomic units (OTUs)]. To compare the accuracy of the protein-, function- and species-based models, we iterated each random forest model 100 times for each of the input datasets separately. The distributions of the OOBs for 100 models for each data type were then compared using t-tests. For the WGCNA analysis, a microbial-host weighted co-correlation network was built using iBAQ values for all proteins and modules were identified using dynamic tree cut. Correlations between proteins and modules were then determined and the top taxa and biological processes assigned to each of the proteins identified. Spearman's rank correlation was used to determine correlation coefficients between individual cytokines/chemokines and module eigengenes (the first principal component of the expression matrix of the corresponding module) for all samples. Average Spearman's correlation coefficients for all pro-inflammatory cytokines and chemokines were then determined. Similarly, Spearman's correlation coefficients were calculated between module eigengenes and Nugent scores and STIs (including *C. trachomatis*, *N. gonorrhoeae*, *T. vaginalis*, *M. genitalium* and active HSV-2).

16S rRNA gene sequencing and analysis

Bacterial 16S rRNA gene sequencing and analysis was conducted as previously described [27]. Briefly, the V4 hypervariable region of the 16S rRNA gene was amplified using modified universal primers [41]. Duplicate samples were pooled, purified using Agencourt AMPure XP beads (Beckman Coulter, CA, USA) and quantified using the Qubit dsDNA HS Assay (Life Technologies, CA, USA). Illumina sequencing adapters and dual-index barcodes were added to the purified amplicon products using limited cycle PCR and the Nextera XT Index Kit (Illumina, CA, USA). Amplicons from 96 samples and controls were pooled in equimolar amounts and the resultant libraries were purified by gel extraction and quantified (Qiagen, Germany). The libraries were sequenced on an Illumina MiSeq platform (300 bp paired-end with v3 chemistry). Following de-multiplexing, raw reads were preprocessed, merged and filtered using DADA2 [42]. Primer sequences were removed using a custom python script and the reads truncated at 250 bp. Taxonomic annotation was based on a customized version of SILVA reference database. Only samples with $\geq$ 2000 reads were included in further analyses. The representative sequences for each amplicon sequence variant were BLAST searched against NCBI non-redundant protein and 16S rRNA gene sequence databases for further validation, as well as enrichment of the taxonomic classifications. Microbial functional predictions were performed using Piphillin server [43] against the Kyoto Encyclopedia of Genes and Genomes (KEGG) database (99% identity cutoff). The relative abundance of functional pathways (KEGG level 3) was used for RF analysis.

Lactobacillus **isolation**

Lactobacilli were isolated from cervicovaginal secretions collected using MCs by culturing in de Man Rogosa and Sharpe (MRS) broth for 48 hours at 37°C under anaerobic conditions. The cultures were streaked onto MRS agar plates under the same culture conditions, single colonies were picked and then pre-screened microscopically by Gram staining. Matrix Assisted Laser Desorption Ionization Time of Flight (MALDI-TOF) was conducted to identify the bacteria to species level. A total of 115 lactobacilli were isolated and stored at -80°C in a final concentration of 60% glycerol. From these, 64 isolates from 25 of the study participants were selected for evaluation of inflammatory profiles.

Vaginal epithelial cell stimulation and measurement of cytokine concentrations

As described previously [25], vaginal epithelial cells (VK2/E6E7 ATCC®CRL-2616™; RRID:CVCL_6471)

were maintained in complete keratinocyte serum free media (KSFM) supplemented with 0.4 mM

calcium chloride, 0.05 mg/ml of bovine pituitary extract, 0.1 ng/ml human recombinant epithelial

growth factor and 50 U/ml penicillin and 50 U/ml streptomycin (Sigma-Aldrich, MO, USA). The VK2

cells were seeded into 24-well tissue culture plates, incubated at 37°C in the presence of 5% carbon

dioxide and grown to confluency. Lactobacilli, adjusted to 4.18 x 10^6 colony forming units (CFU)/ml in

antibiotic-free KSFM, were used to stimulate VK2 cells for 24 hours at 37°C in the presence of 5%

carbon dioxide. As previously described, *G. vaginalis* ATCC 14018 cultures standardized to 1 x 10^7

CFU/ml in antibiotic-free KSFM were used as a positive control [25]. Production of IL-6, IL-8, IL-1α, IL-

1β, IP-10, MIP-3α, MIP-1α, MIP-1β and IL-1RA were measured using a Magnetic Luminex Screening

Assay kit (R&D Systems, MN, USA) and a Bio-Plex™ Suspension Array Reader (Bio-Rad

Laboratories Inc®, CA, USA). VK2 cell viability following bacterial stimulation was confirmed using the

Trypan blue exclusion assay. PCA was used to compare the overall inflammatory profiles of the

isolates and to select the 22 most inflammatory and 22 least inflammatory isolates for characterization

using proteomics.

Characterization of *Lactobacillus* protein expression *in vitro* using mass spectrometry

Forty-four *Lactobacillus* isolates were adjusted to 4.18 x 10^6 CFU/ml in MRS and incubated for 24

hours under anaerobic conditions. Following incubation, the cultures were centrifuged and the pellets

washed 3x with PBS. Protein was extracted by resuspending the pellets in 100 mM triethylammonium

bicarbonate (TEAB; Sigma-Aldrich, MO, USA) 4% sodium dodecyl sulfate (SDS; Sigma-Aldrich, MO,

USA). Samples were sonicated in a sonicating water bath and subsequently incubated at 95°C for

10min. Nucleic acids were degraded using benzonase nuclease (Sigma-Aldrich, MO, USA) and

samples were clarified by centrifugation at 10 000xg for 10 min. Quantification was performed using

the Quanti-Pro BCA assay kit (Sigma-Aldrich, MO, USA). HILIC beads (ReSyn Biosciences, South

Africa) were washed with 250 µl wash buffer [15% ACN, 100 mM Ammonium acetate (Sigma-Aldrich,

MO, USA) pH 4.5]. The beads were then resuspended in loading buffer (30% ACN, 200mM

Ammonium acetate pH 4.5). A total of 50 µg of protein from each sample was transferred to a protein

LoBind plate (Merck, NJ, USA). Protein was reduced with tris (2-carboxyethyl) phosphine (Sigma-

Aldrich, MO, USA) and alkylated with methylmethanethiosulphonate (MMTS; Sigma-Aldrich, MO,

USA). HILIC magnetic beads were added at an equal volume to that of the sample and a ratio of 5:1

total protein and incubated on the shaker at 900 rpm for 30 min. After binding, the beads were

washed four times with 95% ACN. Protein was digested by incubation with trypsin for four hours and

the supernatant containing peptides was removed and dried down in a vacuum centrifuge. LC-MS/MS

analysis was conducted with a Q-Exactive quadrupole-Orbitrap MS as described above. Raw files

were processed with MaxQuant version 1.5.7.4 against a database including the *Lactobacillus* genus

and common contaminants. Log_{10}-transformed iBAQ intensities were compared using Mann-Whitney

U test and p-values were adjusted for multiple comparisons using an FDR step-down procedure.

FGT-METAP application

We have developed FGT-METAP (Female Genital Tract Metaproteomics), an online application for

exploring and mining the FGT microbial composition and function of South African women at high risk

of HIV infection. The app is available at (http://immunodb.org/FGTMetap/) and allows users to repeat

the analyses presented here, as well as perform additional analyses and investigation using the

metaproteomic data generated in this study.

Abbreviations

BP: biological processes

BV: Bacterial vaginosis

CC: cellular component

FDR: false discovery rate

FGT: Female genital tract

GO: gene ontology

iBAQ: intensity-based absolute quantification

LC-MS/MS: liquid chromatography-tandem mass spectrometry

MF: molecular function

OOB: out-of-bag

PCA: principal component analysis

PSA: prostate specific antigen

RF: random forest

STIs: sexually transmitted infections

WGCNA: Weighted correlation network analysis

Declarations

Ethics approval and consent to participate

The University of Cape Town Faculty of Health Sciences Human Research Ethics Committee (HREC

REF: 267/2013) approved this study. Women ≥18 years provided written informed consent, those <18

years provided informed assent and informed consent was obtained from parents/guardians.

Consent for publication

Not applicable.

Availability of data and materials

The dataset supporting the conclusions of this article is available in the FGT-METAP application,

http://immunodb.org/FGTMetap/. Data are available from the corresponding author upon reasonable

request. The protein accession IDs provided at http://immunodb.org/FGTMetap/ reflect UniProt IDs of

each protein and additional information for each protein is provided based on UniProt and NCBI

databases.

Competing interests

The authors declare that they have no competing interests.

Funding

The laboratory work, data analysis and writing were supported by the Carnegie Corporation of New

York, South African National Research Foundation (NRF), the Poliomyelitis Research Foundation,

and the South African Medical Research Council (PI: L. Masson). The WISH cohort was supported by

the European and Developing Countries Clinical Trials Partnership Strategic Primer grant

(SP.2011.41304.038; PI J.S. Passmore) and the South African Department of Science and

Technology (DST/CON 0260/2012; PI J.S. Passmore). J.M.B. thanks the NRF for a SARChI Chair.

D.L.T. was supported by the South African Tuberculosis Bioinformatics Initiative (SATBBI), a Strategic

Health Innovation Partnership grant from the South African Medical Research Council and South

African Department of Science and Technology.

Authors' contributions

AA analyzed the data, developed the FGT-METAP application, and wrote the manuscript; MTM, CB, SD, and SB conducted some of the laboratory experiments, analyzed the data and contributed to manuscript preparation; MP, AI, NR, IA, NM, BC and DLT analyzed the data and contributed to manuscript preparation; LB, EW, MP, ZM, HG conducted some of the laboratory experiments and contributed to manuscript preparation; NM and JMB supervised the data analysis and contributed to manuscript preparation; AG supervised the development of FGT-METAP and hosting the application and contributed to manuscript preparation; LGB managed the clinical site for the WISH study, collected some of the clinical data and contributed to manuscript preparation; HJ supervised the data collection, analyzed the data and contributed to manuscript preparation; JSP was Principal Investigator of the WISH cohort, supervised the collection of some of the data and contributed to manuscript preparation; LM conceptualized the study, supervised the data collection, analyzed some of the data and wrote the manuscript.

Acknowledgements

We would like to acknowledge the participants of the Women's Initiative in Health Study.

Figure legends

Figure 1 Bacterial relative abundance determined using metaproteomics versus 16S rRNA

gene sequencing by inflammation cytokine profile. Liquid chromatography-tandem mass

spectrometry was used to evaluate the metaproteome in lateral vaginal wall swabs from 113 women

from Cape Town, South Africa. Proteins were identified using MaxQuant and a custom database

generated using *de novo* sequencing to filter the UniProt database. Taxonomy was assigned using

UniProt and relative abundance of each taxon was determined by aggregating the intensity-based

absolute quantification (iBAQ) values of all proteins identified for each taxon. (**a**) Proteins identified

were assigned to the Eukaryota and Bacteria domains. Eukaryota proteins included those assigned to

the fungi kingdom and metazoan subkingdom, while the bacteria domain included actinobacteria,

firmicutes, fusobacteria, bacteriodetes and gammaproteobacter phyla. (**b**) Distribution of taxa

identified is shown as a pie chart. (**c**) Number of proteins detected for taxa for which the greatest

number of proteins were identified. (**d**) Protein relative abundance per taxon for taxa with the highest

relative abundance. The relative abundance of the top 20 most abundant bacterial taxa identified

using (**e**) 16S rRNA gene sequencing and (**f**) metaproteomics is shown for all participants for whom

both 16S rRNA gene sequence data and metaproteomics data were generated (n=74). For 16S rRNA

gene sequence analysis, the V4 region was amplified and libraries sequenced on an Illumina MiSeq

platform. Inflammation groups were defined based on hierarchical followed by K-means clustering of

all women according to the concentrations of nine pro-inflammatory cytokine concentrations

[interleukin (IL)-1α, IL-1β, IL-6, IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-

related apoptosis-inducing ligand (TRAIL), interferon (IFN)-γ]. OTU: Operational taxonomic unit.

Figure 2 Weighted co-correlation network analysis of microbial, and human proteins. The

weighted correlation network analysis (WGCNA) R package was used to build a microbial-host

functional weighted co-correlation network using intensity-based absolute quantification (iBAQ)

values. **a**) Correlations between proteins and modules are shown by the heatmap, with positive

correlations shown in red and negative correlations shown in blue. Row sidebars represent the top

taxa and biological processes assigned to each of the proteins (no separate correlation coefficients

were calculated for taxa and biological processes). **b**) The protein dendrogram and module

assignment are displayed, with five modules identified using dynamic tree cut. **c**) Spearman's rank

correlation was used to determine correlation coefficients between individual cytokines/chemokines

and module eigengenes (the first principal component of the expression matrix of the corresponding

module) for all samples. Finally, we reported the average Spearman's correlation coefficients for all

pro-inflammatory cytokines and chemokines. **d**) Similarly, Spearman's correlation coefficients were

calculated between module eigengenes and Nugent scores [bacterial vaginosis (BV)] and sexually

transmitted infections (STIs) as a categorical variable.

Figure 3 Comparison of bacterial, bacterial protein and bacterial function relative abundance

for prediction of genital inflammation. Random forest analysis was used to evaluate the accuracy

of (**a**) bacterial relative abundance (determined using 16S rRNA gene sequencing; n=74), (**b**)

bacterial functional predictions based on 16 rRNA data (n=74), (**c**) bacterial protein relative

abundance (determined using metaproteomics; n=74) and (**d**) bacterial protein molecular function

relative abundance (determined by metaproteomics and aggregation of protein values assigned to the

same gene ontology term; n=74) for determining the presence of genital inflammation (low, medium

and high groups). Inflammation groups were defined based on hierarchical followed by K-means

clustering of women according to the concentrations of nine pro-inflammatory cytokine concentrations

[interleukin (IL)-1α, IL-1β, IL-6, IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-

related apoptosis-inducing ligand (TRAIL), interferon (IFN)-γ]. The bars and numbers within the bars

indicate the relative importance of each taxon, protein or function based on the Mean Decrease in

Gini Value. The sizes of bars in each panel differ based on the length of the labels. (**e**) Each random

forest model was iterated 100 times for each of the input datasets separately and the distribution of

the out-of-bag error rates for the 100 models were then compared using t-tests. OTU: Operational

taxonomic unit.

Figure 4 Microbial biological process and cellular component gene ontologies associated with

genital inflammatory cytokine profiles. Unsupervised hierarchical clustering of aggregated

intensity-based absolute quantification (iBAQ) values for microbial protein (**a**) biological process (BP)

or (**b**) cellular component (CC) gene ontology (GO) relative abundance (n=113). GO relative

abundance was determined by metaproteomics and aggregation of microbial protein iBAQ values

assigned to the same GO term. The heatmaps show aggregated microbial GO relative abundance

and the bar graphs show fold changes in aggregated $\log_2$-transformed iBAQ values (LOGFC) for each

microbial protein with the same (**a**) BP or (**b**) CC GO in women with high versus low inflammation.

Inflammation groups were defined based on hierarchical followed by K-means clustering of women

according to the concentrations of nine pro-inflammatory cytokine concentrations [interleukin (IL)-1α,

IL-1β, IL-6, IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-related apoptosis-

inducing ligand (TRAIL), interferon (IFN)-γ]. Red bars indicate positive and blue bars indicate negative

fold changes in women with high versus low inflammation. False discovery rate adjusted p-values are

shown by dots, with red dots indicating low p-values. Red arrows indicate cell wall and membrane

processes and components. BV: Bacterial vaginosis. Proinflam cyt: Pro-inflammatory cytokine

Figure 5 Relative abundance of gene ontologies in independent samples and in inflammatory

versus non-inflammatory *Lactobacillus* isolates. (a) The top 14 microbial biological process (BP)

and cellular component (CC) gene ontology (GO) terms that distinguished women with low versus

high inflammation in the full cohort (n=113) were validated in an independent sample of ten women

from Cape Town, South Africa. Liquid chromatography-tandem mass spectrometry was used to

evaluate the metaproteome in lateral vaginal wall swabs from these women. Inflammation groups

were defined based on hierarchical followed by K-means clustering of these ten women according to

the concentrations of nine pro-inflammatory cytokine concentrations [interleukin (IL)-1α, IL-1β, IL-6,

IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-related apoptosis-inducing ligand

(TRAIL), interferon (IFN)-γ]. GO relative abundance was determined by aggregation of microbial

protein intensity-based absolute quantification (iBAQ) values assigned to a same GO term. The

relative abundance of the top BPs and CCs (except ATP binding cassette transporter complex which

was not detected in these samples) are shown as bar graphs, with blue indicating women with low

inflammation (n=5) and red indicating women with high inflammation (n=5). Welch's t-test was used

for comparisons. *P<0.05. **(b-i)** Twenty-two *Lactobacillus* isolates that induced relatively high

inflammatory responses and 22 isolates that induced lower inflammatory responses were adjusted to

4.18×10^6 CFU/ml in bacterial culture medium and incubated for 24 hours under anaerobic

conditions. Following incubation, protein was extracted, digested and liquid chromatography-tandem

mass spectrometry was conducted. Raw files were processed with MaxQuant against a database

including the *Lactobacillus* genus and common contaminants. The iBAQ values for proteins with the

same gene ontologies were aggregated, $\log_{10}$-transformed and compared using Mann-Whitney U test.

Box-and-whisker plots show $\log_{10}$-transformed iBAQ values, with lines indicating medians and

whiskers extending to 1.5 times the interquartile range from the box. A false discovery rate step-down

procedure was used to adjust for multiple comparisons and adjusted p-values <0.05 were considered

statistically significant.

Figure 6 Longitudinal changes in FGT metaproteomic profiles. Liquid chromatography-tandem

mass spectrometry was used to evaluate the metaproteome in lateral vaginal wall swabs from 74

women from Cape Town, South Africa at two visits nine weeks (interquartile range: 9-11 weeks)

apart. (**a**) Sankey diagram was used to visualize changes in inflammatory cytokine profiles between

visits. (**b**) Principal component analysis (mixOmics R package) was used to group women based on

the $\log_2$-transformed intensity-based absolute quantification (iBAQ) values of all proteins identified at

both visits. Grouping is based on vaginal pro-inflammatory cytokine concentrations and each point

represents an individual woman. (**c**) Top 20 proteins (UniProt IDs) distinguishing women with and

without inflammation at both visits determined by moderated t-test (limma R package) and random

forest analysis (randomForest R package). (**d**) Top 10 taxa distinguishing women with and without

inflammation at both visits determined by moderated t-test (limma R package) and random forest

algorithm (randomForest R package). (**e**) Top 14 biological process and cellular component gene

ontology terms distinguishing women with and without inflammation at both visits determined by

moderated t-test (limma R package) and random forest algorithm (randomForest R package).

Positions of participants in each heatmap are fixed and the inflammation status of each participant

across the visits can be tracked using the row sidebars. Inflammation groups were defined based on

hierarchical followed by K-means clustering of nine pro-inflammatory cytokine concentrations

[interleukin (IL)-1α, IL-1β, IL-6, IL-12p40, IL-12p70, tumor necrosis factor (TNF)-α, TNF-β, TNF-

related apoptosis-inducing ligand (TRAIL), interferon (IFN)-γ]. ABC: ATP-binding cassette. Agg:

aggregated. BP: biological process. CC: cellular component. Infla: inflammation. PC: principal

component. PEP-PTS: phosphoenolpyruvate-dependent phosphotransferase system. RNDP

complex: ribonucleoside-diphosphate reductase complex.

Supplementary table titles and legends

Additional file 1: Table S1 Demographic and clinical characteristics of study participants stratified by

female genital tract pro-inflammatory cytokine levels

Additional file 2: Table S2 List of protein BLAST results against UniProt and NCBI nr databases

(DBs). Liquid chromatography-tandem mass spectrometry was used to evaluate the metaproteome in

lateral vaginal wall swabs from 113 women from Cape Town, South Africa. Primary assigned taxa,

obtained based on MaxQuant version 1.5.7.4, are presented in the second column. The curated taxa

(third column) were obtained after considering the top similar and frequent BLAST hits for each

protein. IDs: UniProt Identity codes

Additional file 2: Table S3 List of differentially abundant taxa distinguishing women with low and

high inflammation. Liquid chromatography-tandem mass spectrometry was used to evaluate the

metaproteome in lateral vaginal wall swabs from 113 women from Cape Town, South Africa. The

false discovery rate (FDR) corrected p-values were obtained based on the aggregated $\log_2$-

transformed intensity-based absolute quantification of all proteins assigned to the same taxa applying

the moderated t-test (limma R package), after adjusting for confounding variables including age,

contraceptives, prostate specific antigen (PSA) and sexually transmitted infections (STIs). Taxa with

FDR adjusted p-values $\leq$0.05 and $\log_2$-transformed fold change $\geq$ 1.2 or $\leq$ -1.2 were considered as

differentially abundant taxa.

Additional file 2: Table S4 List of differentially abundant proteins distinguishing women with low,

medium and high inflammation. Liquid chromatography-tandem mass spectrometry was used to

evaluate the metaproteome in lateral vaginal wall swabs from 113 women from Cape Town, South

Africa. The false discovery rate (FDR) corrected p-values were obtained based on the $\log_2$-

transformed intensity-based absolute quantification of each protein applying the moderated t-test

(limma R package), after adjusting for confounding variables including age, contraceptives, prostate

specific antigen (PSA) and sexually transmitted infections (STIs). Proteins with FDR adjusted p-

values $\leq$0.05 and log$_2$-transformed fold change $\geq$ 1.2 or $\leq$ -1.2 were considered as differentially

abundant proteins. IDs: UniProt Identity codes

Additional file 2: Table S5 List of differentially abundant molecular function (MF) gene ontology

terms distinguishing women with low and high inflammation. Liquid chromatography-tandem mass

spectrometry was used to evaluate the metaproteome in lateral vaginal wall swabs from 113 women

from Cape Town, South Africa. The false discovery rate (FDR) corrected p-values were obtained

based on the aggregated log$_2$-transformed intensity-based absolute quantification of all proteins

assigned to the same MF term applying the moderated t-test (limma R package), after adjusting for

confounding variables including age, contraceptives, prostate specific antigen (PSA) and sexually

transmitted infections (STIs). Taxa with FDR adjusted p-values $\leq$0.05 and log$_2$-transformed fold

change $\geq$ 1.2 or $\leq$ -1.2 were considered as significantly abundant MFs.

Additional file 1: Table S6 Details of MaxQuant version 1.5.7.4 parameters used for analysis of

metaproteomic data obtained from lateral vaginal wall swabs from 113 women from Cape Town,

South Africa.

References

1. UNAIDS Data 2019. Available at: https://www.unaids.org/sites/default/files/media_asset/2019-UNAIDS-data_en.pdf. Accessed 20/02/20.

2. Masson L, Passmore JAS, Liebenberg LJ, Werner L, Baxter C, Arnold KB, et al. Genital Inflammation and the Risk of HIV Acquisition in Women. Clin Infect Dis. 2015;61:260-269.

3. McKinnon LR, Liebenberg LJ, Yende-Zuma N, Archary D, Ngcapu S, Sivro A, et al. Genital inflammation undermines the effectiveness of tenofovir gel in preventing HIV acquisition in women. Nat Med. 2018;24:491.

4. Morrison C, Fichorova RN, Mauck C, Chen PL, Kwok C, Chipato T, et al. Cervical inflammation and immunity associated with hormonal contraception, pregnancy, and HIV-1 seroconversion. J Acquir Immune Defic Syndr. 2014;66:109-117.

5. Li Q, Estes JD, Schlievert PM, Duan L, Brosnahan AJ, Southern PJ, et al. Glycerol monolaurate prevents mucosal SIV transmission. Nature. 2009;458:1034-1038.

6. Osborn L, Kunkel S, Nabel GJ. Tumor necrosis factor α and interleukin 1 stimulate the human immunodeficiency virus enhancer by activation of the nuclear factor κB. Proc Natl Acad Sci U S A. 1989;86:2336-2340.

7. Arnold KB, Burgener A, Birse K, Romas L, Dunphy LJ, Shahabi K, et al. Increased levels of inflammatory cytokines in the female reproductive tract are associated with altered expression of proteases, mucosal barrier proteins, and an influx of HIV-susceptible target cells. Mucosal Immunol. 2016;9:194-205.

8. Burgener A, Rahman S, Ahmad R, Lajoie J, Ramdahin S, Mesa C, et al. Comprehensive proteomic study identifies serpin and cystatin antiproteases as novel correlates of HIV-1 resistance in the cervicovaginal mucosa of female Sex workers. J Proteome Res. 2011;11:5139-5149.

9. Masson L, Mlisana K, Little F, Werner L, Mkhize NN, Ronacher K, et al. Defining genital tract cytokine signatures of sexually transmitted infections and bacterial vaginosis in women at high risk of HIV infection: A cross-sectional study. Sex Transm Infect. 2014;90:580-587.

10. Lennard K, Dabee S, Barnabas SL, Havyarimana E, Blakney A, Jaumdally SZ, et al. Microbial composition predicts genital tract inflammation and persistent bacterial vaginosis in South African adolescent females. Infect Immun. 2018;86:e00410-17.

11. Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M, et al. Cervicovaginal bacteria are a major modulator of host inflammatory responses in the female genital tract. Immunity. 2015;42: 965-976.

12. Deese J, Masson L, Miller W, Cohen M, Morrison C, Wang M, et al. Injectable progestin-only contraception is associated with increased levels of pro-inflammatory cytokines in the female genital tract. Am J Reprod Immunol. 2015;74:357-367.

13. Gosmann C, Anahtar MN, Handley SA, Farcasanu M, Abu-Ali G, Bowman BA, et al. Lactobacillus-deficient cervicovaginal bacterial communities are associated with increased HIV acquisition in young South African women. Immunity. 2017;46:29-37.

14. Chapot-Chartier MP, Kulakauskas S. Cell wall structure and function in lactic acid bacteria. Microb Cell Fact. 2014;13:S9.

15. Chetwin E, Manhanzva MT, Abrahams AG, Froissart R, Gamieldien H, Jaspan H, et al. Antimicrobial and inflammatory properties of South African clinical Lactobacillus isolates and vaginal probiotics. Sci Rep. 2019; 9:1-15.

16. Hearps AC, Tyssen D, Srbinovski D, Bayigga L, Diaz DJD, Aldunate M, et al. Vaginal lactic acid elicits an anti-inflammatory response from human cervicovaginal epithelial cells and inhibits production of pro-inflammatory mediators associated with HIV acquisition. Mucosal Immunol. 2017;10:1480-1490.

17. McKinnon LR, Achilles SL, Bradshaw CS, Burgener A, Crucitti T, Fredricks DN, et al. The Evolving Facets of Bacterial Vaginosis: Implications for HIV Transmission. AIDS Res. Hum. Retroviruses. 2019.

18. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, et al. Vaginal microbiome of reproductive-age women. Proc Natl Acad Sci U S A. 2011;35:219-228.

19. Xiong W, Abraham PE, Li Z, Pan C, Hettich RL. Microbial metaproteomics for characterizing the range of metabolic functions and activities of human gut microbiota. Proteomics. 2015;15:3424-38.

20. Klatt NR, Cheu R, Birse K, Zevin AS, Perner M, Noël-Romas L, et al. Vaginal bacteria modify HIV tenofovir microbicide efficacy in African women. Science. 2017;356:938-945.

21. Zevin AS, Xie IY, Birse K, Arnold K, Romas L, Westmacott G, et al. Microbiome Composition and Function Drives Wound-Healing Impairment in the Female Genital Tract. PLoS Pathog. 2016;12: e1005889.

22. Barnabas SL, Dabee S, Passmore JAS, Jaspan HB, Lewis DA, Jaumdally SZ, et al. Converging

epidemics of sexually transmitted infections and bacterial vaginosis in southern African female

adolescents at risk of HIV. Int J STD AIDS. 2018;29:531-539.

23. Dabee S, Barnabas SL, Lennard KS, Jaumdally SZ, Gamieldien H, Balle C, et al. Defining

characteristics of genital health in South African adolescent girls and young women at high risk for

HIV infection. PLoS One. 2019;14:e0213975.

24. Fernandez EM, Valenti V, Rockel C, Hermann C, Pot B, Boneca IG, et al. Anti-inflammatory

capacity of selected lactobacilli in experimental colitis is driven by NOD2-mediated recognition of a

specific peptidoglycan-derived muropeptide. Gut. 2011;60:1050-1059.

25. Manhanzva MT, Abrahams AG, Gamieldien H, Froissart R, Jaspan H, Jaumdally SZ, et al.

Inflammatory and antimicrobial properties differ between vaginal Lactobacillus isolates from South

African women with non-optimal versus optimal microbiota. Sci Rep. 2020;10:1-13.

26. Macho Fernandez E, Pot B, Grangette C. Beneficial effect of probiotics in IBD: Are peptidogycan

and NOD2 the molecular key effectors? Gut Microbes. 2011;2:280-6.

27. Balle C, Lennard K, Dabee S, Barnabas SL, Jaumdally SZ, Gasper MA, et al. Endocervical and

vaginal microbiota in South African adolescents with asymptomatic Chlamydia trachomatis infection.

Sci Rep. 2018;8:1-9.

28. Lagier JC, Dubourg G, Million M, Cadoret F, Bilen M, Fenollar F, et al. Culturing the human

microbiota and culturomics. Nat. Rev. Microbiol. 2018;16:540-550.

29. Holm JB, France M, Ma B, Robinson CK, McComb E, Brotman RM, et al. Comparative

metagenome-assembled genome analysis of Lachnovaginosum genomospecies, formerly known as

BVAB1. bioRxiv. 2019;657197.

30. Huffnagle GB, Noverr MC. The emerging world of the fungal microbiome. Trends Microbiol. 2013.

31. Bianco L, Perrotta G. Methodologies and perspectives of proteomics applied to filamentous fungi:

From sample preparation to secretome analysis. Int. J. Mol. Sci. 2015;21:334-41.

32. Verstraelen H, Verhelst R, Claeys G, De Backer E, Temmerman M, Vaneechoutte M. Longitudinal

analysis of the vaginal microflora in pregnancy suggests that L. crispatus promotes the stability of the

normal vaginal microflora and that L. gasseri and/or L. iners are more conducive to the occurrence of

abnormal vaginal microflora. BMC Microbiol. 2009;9:116.

33. van Houdt R, Ma B, Bruisten SM, Ravel AGCLSJ, Vries HJC de. Lactobacillus iners-dominated vaginal microbiota is associated with increased susceptibility to Chlamydia trachomatis infection in Dutch women: a case–control study. Sex Transm Infect. 2018;94(2):117-23.

34. Potgieter MG, Nel AJM, Tabb DL, Fortuin S, Garnett S, Blackburn JM, et al. MetaNovo: a probabilistic approach to peptide and polymorphism discovery in complex mass spectrometry datasets. bioRxiv. 2019;605550.

35. Afiuni-Zadeh S, Boylan KLM, Jagtap PD, Griffin TJ, Rudney JD, Peterson ML, et al. Evaluating the potential of residual Pap test fluid as a resource for the metaproteomic analysis of the cervical-vaginal microbiome. Sci Rep. 2018;8:10868.

36. Smyth GK. limma: Linear Models for Microarray Data. Bioinforma Comput Biol Solut Using R Bioconductor. 2005.

37. Rohart F, Gautier B, Singh A, Lê Cao KA. mixOmics: An R package for 'omics feature selection and multiple data integration. PLoS Comput Biol. 2017;13:e1005752.

38. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics. 2016;32:2847-9.

39. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network analysis. BMC Bioinformatics. 2008;9:559.

40. Liaw A, Wiener M. Classification and Regression by randomForest. R News. 2002;2:18-22.

41. Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, et al. The female urinary microbiome: A comparison of women with and without urgency urinary incontinence. MBio. 2014;5, e01283-14.

42. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13:581.

43. Narayan NR, Weinmaier T, Laserna-Mendieta EJ, Claesson MJ, Shanahan F, Dabbagh K, et al. Piphillin predicts metagenomic composition and dynamics from DADA2-corrected 16S rDNA sequences. BMC Genomics. 2020;21:1-12.

a

Prevotella	0.5
Mycoplasma hominis	0.51
Fastidiosipila	0.52
Prevotella(OTU336)	0.52
Megasphaera	0.53
Dialister micraerophilus	0.54
Peptostreptococcus anaerobius	0.54
Dialister	0.62
Ureaplasma parvum	0.65
Gardnerella(OTU7)	0.67
Gemella asaccharolytica	0.67
Lactobacillus	0.67
Corynebacterium	0.75
Lachnovaginosum BVAB1	1.04
Aerococcus christensenii	1.18
Sneathia amnii	1.71
Gardnerella vaginalis	1.89
Lactobacillus iners	2.23
Atopobium vaginae	2.4
Sneathia sanguinegens	2.64

b

Sphingolipid metabolism	0.39
Biosynthesis of ansamycins	0.42
Neomycin kanamycin and gentamicin biosynthesis	0.43
Primary immunodeficiency	0.44
Prodigiosin biosynthesis	0.45
Chloroalkane and chloroalkene degradation	0.46
Tuberculosis	0.47
PPAR signaling	0.49
Epithelial cell signaling	0.52
Alanine aspartate and glutamate metabolism	0.52
Photosynthesis	0.52
GABAergic synapse	0.53
Oxidative phosphorylation	0.53
Sulfur relay system	0.63
Secondary bile acid biosynthesis	0.64
Primary bile acid biosynthesis	0.65
Carbon metabolism	0.67
Tyrosine metabolism	1.03
Biofilm formation	1.32
Prolactin signaling	1.63

c

30S ribosomal protein S13	0.44
60 kDa chaperonin	0.46
Formate C−acetyltransferase	0.47
Transcription elongation factor GreA	0.47
Elongation factor Tu	0.48
Threonine−−tRNA ligase	0.49
Fructose−1,6−bisphosphate aldolase, class II	0.53
Phosphoglycerate kinase	0.56
2,3−bisphosphoglycerate	0.57
Glutamine synthetase	0.59
Triosephosphate isomerase	0.75
Elongation factor Tu(C8PAS6)	0.9
Actin, cytoplasmic 2	0.9
Phosphocarrier, HPr family	0.91
ATP synthase subunit alpha	0.92
Pyrophosphate phospho−hydrolase	1.05
Cold−shock DNA−binding domain protein	1.28
Phosphoglycerate kinase(A0A0E9DXN4)	1.72
Glyceraldehyde−3−phosphate dehydrogenase	1.87
Chaperone protein DnaK	2.05

d

ATP binding	0.47
D−alanine−poly(phosphoribitol) ligase activity	0.47
rRNA binding	0.52
aminoacyl−tRNA editing activity	0.53
adenylate kinase activity	0.54
5S rRNA binding	0.62
GTP binding	0.63
ribosome binding	0.63
phosphoglucosamine mutase activity	0.64
translation elongation factor activity	0.64
structural constituent of ribosome	0.65
phosphoglycerate kinase activity	0.68
amidase activity	0.7
DNA binding	0.77
adenylosuccinate synthase activity	0.84
hydrolase activity	1.18
glutamate−ammonia ligase activity	1.22
GTPase activity	1.51
beta−phosphoglucomutase activity	2.92
oxidoreductase activity	3.27

e

a

CC
- ribonucleoside-diphosphate reductase complex
- S-layer
- small ribosomal subunit
- large ribosomal subunit
- peptidoglycan-based cell wall

BP
- peptidoglycan biosynthetic process *
- cell wall organization *
- transcription, DNA-templated
- PEP-PTS *
- ATP metabolic process *
- response to oxidative stress
- regulation of cell shape *
- malate metabolic process

Low (n=5)
High (n=5)

Log2-transformed GO iBAQ value

b p=0.0172* Cell wall organization BP

c p=0.0272 Regulation of cell shape BP

d p=0.0327 Peptidoglycan biosynthetic process BP

e p=0.0002* Plasma membrane CC

f p=0.0092* Cell wall CC

g p=0.0327 Membrane CC

h p=0.1372 S-layer CC

i p=0.1372 Peptidoglycan-based cell wall CC

Log10-transformed GO iBAQ value

Lactobacillus isolates inducing low inflammatory responses (n=22)
Lactobacillus isolates inducing high inflammatory responses (n=22)