Genomic signals of admixture and reinforcement between two closely related species of

European sepsid flies

Athene Giesen[1], Wolf U. Blanckenhorn[1], Martin A. Schäfer[1], Kentaro K. Shimizu[1], Rie

Shimizu-Inatsugi[1], Bernhard Misof[2], Oliver Niehuis[3], Lars Podsiadlowski[2], Heidi E. L.

Lischer[1,4], Simon Aeschbacher[1], Martin Kapun[1,5]

[1]Department of Evolutionary Biology and Environmental Studies, University of Zürich,

Switzerland

[2]Zoological Research Museum Alexander Koenig, Bonn, Germany

[3]Department of Evolutionary Biology and Ecology, Institute of Biology I (Zoology), Albert

Ludwig University, Freiburg, Germany

[4]Interfaculty Bioinformatics Unit, University of Bern, Switzerland

[5]Division of Cell & Developmental Biology, Medical University of Vienna, Austria

Corresponding author: Martin Kapun; Winterthurerstrasse 190, CH-8057 Zürich,

Switzerland; Tel.: +41 44 635 47 55; martin.kapun@ieu.uzh.ch

ABSTRACT

Interspecific gene flow by hybridization may weaken species barriers and adaptive

divergence, but can also initiate reinforcement of reproductive isolation trough natural and

sexual selection. The extent of interspecific gene flow and its consequences for the initiation

and maintenance of species barriers in natural systems remain poorly understood, however.

To assess genome-wide patterns of gene flow between the two closely related European dung

fly species *Sepsis cynipsea* and *Sepsis neocynipsea* (Diptera: Sepsidae), we tested for

historical gene flow with the aid of ABBA-BABA test using whole-genome resequencing

data from pooled DNA of male specimens originating from natural and laboratory

populations. We contrasted genome-wide variation in DNA sequence differences between

samples from sympatric populations of the two species in France and Switzerland with that of

interspecific differences between pairs of samples involving allopatric populations from

Estonia and Italy. In the French Cevennes, we detected a relative excess of DNA sequence

identity, suggesting interspecific gene flow in sympatry. In contrast, at two sites in

Switzerland, we observed a relative depletion of DNA sequence identity compatible with

reinforcement of species boundaries in sympatry. Our results suggest that the species

boundaries between *S. cynipsea* and *S. neocynipsea* in Europe depend on the eco-geographic

context.

KEYWORDS: ABBA-BABA test, gene flow, hybridization, introgression, reproductive

isolation, speciation, sepsid flies

Running title: Patterns of Gene flow in European Sepsid flies

INTRODUCTION

According to the biological species concept, species represent groups of individuals that are reproductively isolated from other groups of individuals (Mayr 1942). Speciation entails the evolution of reproductive isolation among lineages derived from a common ancestral population and is considered completed if the diverged populations remain reproductively isolated, for example, after coming into secondary contact (Coyne & Orr, 2004; Dobzhansky, 1951; Mayr, 1963). However, many animal and plant species remain distinct entities in nature even if they occasionally hybridize and exchange genes in parapatry or sympatry (Anderson, 1949; Barton & Bengtsson, 1986; cf. DeMarais *et al.*, 1992; Gante *et al.*, 2016; Mallet, 2007; Nolte & Tautz, 2010; Rieseberg *et al.*, 2003; Trier *et al.*, 2014). Research during the past decades has indicated that hybridization not only has deleterious effects due to hybrid inferiority and negative epistasis in admixed genomes, but that it may also fuel adaptive diversification and speciation by facilitating novel combinations of alleles that become targets of divergent selection (Arnold & Meyer, 2006; Berner & Salzburger, 2015; Fontaine *et al.* 2015; Seehausen, 2004; Saetre, 2013). Together, these findings corroborate the view that genomes of recently diverged species may represent mosaics, consisting of genomic regions with significant differentiation (i.e., low to no admixture) interspersed by genomic regions that are more free to be exchanged (Nosil *et al.*, 2009; Wu, 2001).

Sepsid flies (Diptera: Sepsidae) have become a model group for the study of sexual selection and ecological adaptation (Baur *et al.*, 2020; Blanckenhorn, 1999; Blanckenhorn *et al.*, 2000; Eberhard, 1999; Kraushaar & Blanckenhorn, 2002; Parker, 1972a,b; Puniamoorthy *et al.*, 2009; Pont & Meier, 2002; Rohner *et al.*, 2015; Rohner, Blanckenhorn, & Puniamoorthy, 2016; Ward, 1983; Ward, Hemmi, & Rösli, 1992). The phylogeny of sepsid flies is well resolved (Su *et al.*, 2008, 2016) and entails multiple pairs of closely related

species that occupy similar ecological niches. These species pairs provide excellent

opportunities to study the genomic consequences of hybridization and introgression during

early stages of speciation. One of these pairs comprises *Sepsis cynipsea* and *Sepsis*

neocynipsea, sister species with a wide geographic distribution that occur in sympatry across

major parts of their natural range. While *S. cynipsea* is the most abundant sepsid species in

Central and Northern Europe and deposits its eggs into fresh cow dung, *S. neocynipsea* is

common throughout North America, where it occupies a niche similar to that of *S. cynipsea* in

Europe. While overall very rare in Europe, *S. neocynipsea* can be locally common at higher

altitudes, such as the Alps, where the species occurs in sympatry with *S. cynipsea* (Ozerov,

2005; Pont & Meier, 2002; Rohner, Blanckenhorn, & Puniamoorthy, 2016). Despite strong

similarities in morphology and behavior (Giesen, Blanckenhorn, & Schäfer, 2017), the two

species are genetically distinct. Previous studies showed that *S. cynipsea* and *S. neocynipsea*

produce fertile hybrid offspring under laboratory conditions despite strong pre- and post-

mating isolating barriers. These barriers are mediated by assortative mating behaviors that are

partly reinforced in areas where the two species occur in sympatry (Giesen *et al.*, 2017, 2019).

While these findings imply that interspecific gene flow may occur and vary with eco-

geographic context in nature, we know little about actual levels of historical and ongoing gene

flow between the two species in areas where they occur in sympatry or parapatry. We

therefore investigated the extent of gene flow between *S. cynipsea* and *S. neocynipsea* in

nature by comparative population genomic analyses.

In recent years, multiple approaches have been employed to explore the importance of

interspecific gene flow in sympatry vs. allopatry (e.g. Nakazato, Warren, & Moyle, 2010;

Nadeau *et al.*, 2013; Brandvain *et al.*, 2014; Bouchemousse *et al.*, 2016; Feulner &

Seehausen, 2019; Kastally, Trasoletti, & Mardulyn, 2019; Moran *et al.*, 2019). The objective

of our study is to assess the extent of genome-wide admixture between *S. cynipsea* and *S.*

neocynipsea in Europe by applying a version of the ABBA-BABA test for historical gene

flow (Green *et al*., 2010; Durand *et al*., 2011; Soraggi et al., 2018) that can exploit genome-

wide allele frequency data of single nucleotide polymorphisms (SNPs). To this end, we

sequenced the pooled genomic DNA of males from wild-caught populations of *S. cynipsea*

and *S. neocynipsea* collected at multiple sites ranging from Southern France to Estonia. Some

of these populations occur in sympatry (e.g., in the Swiss Alps and the French Cevennes),

others in allopatry (Pont & Meier, 2002). Based on hybridization opportunity alone, we

expected to find higher genome-wide levels of gene flow between *S. cynipsea* and *S.*

neocynipsea in geographic areas where both species occur in sympatry (e.g. Nadeau *et al*.,

2013, or Martin *et al*., 2014, for *Heliconius* butterflies). In contrast, less gene flow in

sympatry than in allopatry would indicate selection against gene flow and thus suggest

reinforcement of reproductive barriers at sites of co-occurrence (Butlin, 1995; Noor, 1999;

Coyne & Orr, 2004; e.g. Kulathinal & Singh, 2000; Massie & Makow, 2005; Giesen *et al*.

2017, 2019).

MATERIALS & METHODS

Sample collection and treatment of fly cultures

We studied inter- and intraspecific gene flow in S. *cynipsea* and *S. neocynipsea* using

genomic data of sympatric *S. cynipsea* and *S. neocynipsea* populations from two high altitude

sampling sites in the Swiss Alps (Sörenberg) and the French Cevennes (Le Mourier) (Table

1). To complement these two pairs of sympatric populations with geographically distant

allopatric populations in Europe (Table 1), we further included *S. cynipsea* samples from the

Swiss lowlands (Zürich), from Tuscany, Italy (Petroia) and Estonia (Pehka), and from two

additional high-altitude *S. neocynipsea* samples collected in Switzerland (Geschinen and

Hospental) (Table 1). In line with the previous observation that *S. neocynipsea* tends to be

rare at low altitudes (Pont & Meier, 2002), we were not able to collect sufficient numbers of

flies of this species outside the Alps, except for the sample collected near Le Mourier. For

each of the wild-caught populations, we randomly selected 20 males for pooled DNA

resequencing (Pool-Seq).

In addition to wild-caught populations, we compiled genomic data from two laboratory

populations that were established from sympatric *S. cynipsea* and *S. neocynipsea* populations

collected near Zürich. We started each of these populations from a single gravid, wild-caught

female and subsequently propagated them in the laboratory for at least two years at census

population sizes ranging from 10 to 100. We then sampled and pooled the DNA of 50 males

from each population for whole-genome sequencing (see Supporting Information for more

details).

DNA library preparation and next generation sequencing

Genomic DNA was extracted from the pooled DNA of males using the UltraPure

Phenol:Chloroform:Isoamyl alcohol (25:24:1, v/v) extraction kit (Thermo Fischer Scientific,

Waltham, USA) according to the manufacturer's protocol. Quantification of genomic DNA

was performed with a Qubit Fluorometer (Thermo Fischer Scientific; Table 1). Library

preparation was carried out with the TruSeq DNA PCR-Free Library Preparation (Illumina,

San Diego, USA) kit according to the manufacturer's protocol. Fragment-size distributions of

all libraries were validated on a TapeStation 2200 (Agilent Technologies, Waldbronn,

Germany). Sequencing on Illumina HiSeq 2500 version 4 was conducted after labeling and

pooling the barcoded DNA representing the four laboratory populations onto one lane to

achieve a genome-wide coverage of ca. 60x per DNA pool library. All sequencing data are

available at the short-read archive (SRA; https://www.ncbi.nlm.nih.gov/sra) under the

accession number PRJNA612154.

Table 1: Sampling sites, sample IDs, percentages of DNA sequence reads mapping to the reference genome and read-depths of pool-sequenced field-collected (a) and laboratory (b) populations. Sites where *S. cynipsea* and *S. neocynipsea* occur in sympatry are in regular font. Sites where *S. neocynipsea* does not occur to our knowledge are highlighted in italics.

Species	ID	Location	Country	Coordinates	Altitude (m)	Alignment rate to *S. thoracica* reference [%]	Average Read-depth
a) Wild-caught							
S. cynipsea	PhC	*Pehka*	*Estonia*	59°28'45.1''N,25°44'52.3''E	5	50.6%	19.8
	PtC	*Petroia*	*Italy*	43°15'06.4''N,12°32'41.2''E	570	48.1%	29.2
	SoC	Sörenberg	Switzerland	46°52'12''N, 8°16'12''E	1200	51.3%	35.1
	ZuC	Zürich	Switzerland	47°24'0.6''N, 8°34'24.0''E	450	51.3%	37.3
	MoC	Le Mourier	France	44° 3'30.5"N, 3°25'38.8"E	900	49.5%	62.5
S. neocynipsea	GeN	Geschinen	Switzerland	46°49'23.7''N, 8°1'54.6''E	1350	50.5%	32.1
	HoN	Hospenthal	Switzerland	46° 37'12''N, 8°34'12''E	1500	52.3%	25.4
	SoN	Sörenberg	Switzerland	46°24'8.2''N, 8°21'28.4''E	1200	50.7%	23.0
	MoN	Le Mourier	France	44° 3'30.5"N, 3°25'38.8"E	900	48.6%	74.2
b) Laboratory							
S. cynipsea	IZuC	Zürich	Switzerland	47°24'0.6''N, 8°34'24.0''E	402	50.1%	19.2
S. neocynipsea	IZuN	Zürich	Switzerland	47°24'0.6''N, 8°34'24.0''E	402	50.0%	67.2
c) *S. orthocnemis*		Lenzerheide	Switzerland	46°43'45''N, 9°33'25''E	1470	46.9%	13.0

Raw read processing

Qualitative validation of sequence data before and after trimming was done with FastQC (v. 0.11.4; Andrews *et al*., 2011). After removal of Illumina-specific adapters and trimming with *Trimmomatic* (v. 0.36; Bolger, Lohse & Usadel, 2014), the *S. cynipsea* and *S. neocynipsea* reads were mapped to the draft genome of *S. thoracica* with *bwa mem* (v. 0.7.12;

Li & Durbin, 2009) using default parameters. The *S. thoracica* sample used for genome

sequencing was collected near Capriasca, Ticino, Switzerland. The draft genome, v. 0.1 was

built from Oxford Nanopore long reads (app. 25x read depth), assembled with Canu (Korner

et al., 2017), and polished with Illumina short reads (app. 20x read depth). The assembly used

in this study is available under the rules of the Fort Lauderdale agreement from

www.cgae.de/seto_01_genome.fasta and www.cgae.de/seto_01_genome_masked.fasta.

PCR duplicates were removed with Picard (v. 1.109; http://broadinstitute.github.io/picard/)

and reads were realigned around indels in each raw alignment file using

RealignerTargetCreator and IndelRealigner from GATK (v. 3.4-46; McKenna et al. 2010).

Only reads with a PHRED-scaled mapping quality of 20 and more were retained. We applied

the same procedure to map reads of a pool of ten inbred *S. orthocnemis* males from near

Lenzerheide, Switzerland, to the reference genome (Table 1). We compiled the aligned reads

from all population samples into a single *mpileup* file using *samtools* (v. 1.3.1; Li *et al.*,

2009).

Using the variant-caller Pool-SNP (Kapun *et al.*, 2018), we identified high-confidence

SNPs with the following combination of heuristic SNP-calling parameters: coverage of each

Pool-Seq sample ≥ 10x; coverage of each Pool-Seq sample ≤ 95 % percentile of coverage

distribution across contigs and samples; minimum allele count of a minor allele at a SNP

across all combined Pool-Seq samples > 20x; minor allele frequency at a SNP across all

combined Pool-Seq samples > 0.01. We only retained SNPs for which all samples fulfilled

the above coverage threshold criteria. To avoid paralogous SNPs due to mis-mapping around

indel polymorphisms, we removed SNPs located within a 5-bp distance to indels that

occurred in more than 20 copies at a given site across all pooled samples. In addition, we

ignored SNPs located within repetitive regions. The resulting VCF file was converted to the

SYNC file format (Kofler *et al*., 2011), and a custom Python script was used to calculate

sample-specific allele frequencies for major alleles at each SNP (*sync2AF.py*;

https://github.com/capoony/DrosEU_pipeline).

Testing for introgression

To test for introgression, we used the ABBA-BABA test for historical gene flow in the

presence of incomplete lineage sorting (Green *et al*., 2010; Durand *et al*., 2011). We provide a

brief motivation of the approach in the following and refer to the Supporting Information for

further details. Imagine a rooted phylogeny with four species (P1 to P4) and a topology of

(((P1, P2), P3), P4) as illustrated in Fig. 1B. In an alignment of one haploid genome from

each species, incomplete lineage sorting under neutral evolution leads to two mutational

configurations of the ancestral (A) and derived (B) allele, ABBA and BABA, that are

incompatible with the species topology (Fig. 1B) under the infinite-sites model of mutation

(Kimura, 1969). In the absence of gene flow between P2 and P3, ABBA and BABA

configurations occur with equal probability (Hudson, 1983; Tajima, 1983). In contrast,

historical gene flow between P2 and P3 causes an excess of ABBA configurations (Green *et*

al., 2010; Durand *et al*., 2011). This logic is captured by the *D*-statistic, a scaled difference

across a set of bases between counts of ABBA and BABA configurations bounded by –1 and

1 (Green *et al*., 2010). The ABBA-BABA test examines significant deviations of *D* from 0. A

significant excess of ABBA suggests evidence for gene flow between P2 and P3. A

significant depletion of ABBA is equivalent to an excess of ABBA if the positions of P1 and

P2 in the species topology are swapped, and hence either suggests gene flow between P1 and

P3, or a reduction in gene flow between P2 and P3 relative to gene flow between P1 and P3.

The original version of the ABBA-BABA test and modifications of it have been applied to

different types of genome-scale polymorphism data obtained with various sequencing

strategies, including whole genome sequencing (e.g. Green *et al*., 2010), RAD sequencing

(e.g., Eaton & Ree, 2013; Meier *et al*., 2017; Streicher *et al*., 2014), and exon capture data

(e.g., Heliconius Genome Consortium, 2012). More recently, Durand *et al*. (2011) and

Soraggi *et al.* (2018) extended the original test to allele frequency data, hence to unphased

sequencing data (including Pool-Seq data; Schlötterer *et al.* 2014). We implemented the

extensions by Durand *et al.* (2011) and Soraggi *et al.* (2018) in a Python script

(https://github.com/capoony/ABBABABA-4AF) and refer to the respective test statistics as

D_D and D_S. Our script also computes jack-knifed *z*-scores based on a matrix of allele

frequencies for previously defined high-confidence SNPs (see Supporting Information for

details). We adopted the commonly used significance threshold of $|z| > 3$ (Reich *et al.*, 2011;

Jeong *et al.*, 2016; Novikova *et al.*, 2016) to identify significant deviations of D_S and D_D from

zero.

We were concerned that using the same genome assembly both as the reference for read

mapping and as the outgroup (P4) in the ABBA-BABA tests could lead to biased estimates of

gene flow. We therefore mapped the three ingroup and the outgroup species against the

reference genome of yet another species, *S. thoracica*. A phylogenetic analysis based on CO-

II sequences by Su *et al.* (2008) indicated that *S. thoracica* is approximately equally related to

S. cynipsea and *S. neocynipsea* as it is from the outgroup species *S. orthocnemis* (Fig. 1A).

We therefore expected that reads of the ingroup and the outgroup species would map equally

well to the reference species, which is indeed what we observed (Table 1).

Fig. 1. Putative phylogeny of sepsid species studied. (A) Species phylogeny inferred from
alignments of nucleotide sequences at the cytochrome oxidase subunit II (CO-II)
mitochondrial locus (Genbank sequences from Su *et al*., 2016) with the Neighbor-Joining
method implemented in CLC Main Workbench (v. 8.1.2; https://
https://www.qiagenbioinformatics.com/products/clc-main-workbench/). The tree topology is
consistent with the phylogeny presented by Su *et al.'s* (2016) combined phylogenetic analysis
of multiple nuclear and mitochondrial markers. *Sepsis orthocnemis* was used as the outgroup
for the focal species *S. cynipsea* and *S. neocynipsea*. Whole-genome sequencing reads from
all three species were aligned to the *S. thoracica* reference genome (dashed red branches).
Branch lengths are proportional to the mean number of pairwise sequence differences. (B)
Generic species tree assumed in all our ABBA-BABA tests for gene flow among triplets of *S.*
cynipsea and *S. neocynipsea* populations from various sites in Europe.

We focused our analysis of historical interspecific gene flow on the three sampling sites

Sörenberg, Zürich and Le Mourier (Table 1). At all these sites, *S. cynipsea* and *S. neocynipsea*

occur in sympatry. In all ABBA-BABA tests, we positioned the two focal populations as P2

and P3 ingroups in the phylogeny (in both possible orders) and used various P1 ingroup

populations (Fig. 1B) assumed to occur in allopatry with the P2 populations. We used various

P1 ingroups, because Durand *et al*. (2011) showed that the choice of the ingroup P1 can

influence the test results.

RESULTS

Signals of reinforcement in Swiss alpine and sub-alpine regions

The majority of our tests for gene flow at the Sörenberg site revealed a genome-wide

deficiency of derived-allele sharing (i.e., ABBA patterns) among the local *S. cynipsea* and *S.*

neocynipsea populations (Table 2), suggesting that interspecific gene flow in sympatry

(among P2 and P3) is lower than interspecific gene flow among putatively allopatric

populations (P1 and P3). This observation is robust to our choice of the P1 ingroup

population, but sensitive to the version of D-statistic used (D_S vs. D_D). We first conducted

tests involving two *S. neocynipsea* populations from the Swiss Alps (*S. neocynipsea* from

Geschinen, GeN; *S. neocynipsea* from Hospental, HoN; Table 1) as putatively allopatric P1,

and the sympatric interspecific pair from Sörenberg as P2 (*S. neocynipsea*) and P3 (*S.*

cynipsea). We observed a significant deficiency of ABBA patterns in all configurations ($D_S >$

0; Table 2, upper part), suggesting overall evidence for higher gene flow among putatively

allopatric than sympatric interspecific population pairs. While GeN and HoN are both

separated by high mountains (altitudes ranging from ca. 600–3,200 m asl) from the Sörenberg

site, they are geographically close (ca. 67.5 km and ca. 46.7 km Euclidean distance for GeN

and HoN, respectively) and thy occur in sympatry with other sepsid species, including *S.*

cynipsea. We were therefore concerned that our assumption of no gene flow between P1 and

P3 was violated, because GeN and HoN are parapatric rather than allopatric to our focal

populations in Sörenberg.

To address this concern, we conducted additional ABBA-BABA tests involving P1

populations from geographically more remote sites in Europe. Specifically, we performed a

second series of tests with two alternative *S. cynipsea* populations as allopatric P1 ingroups

(Fig. 1B), one from Pehka, Estonia (PhC), and one from Petroia, Italy (PtC; Table 1).

Contrary to our expectation based on the much larger geographical distance between P1 and

P3 (ca. 1,850 km and ca. 550 km for PhC and PtC, respectively) than between P2 and P3 in

these configurations, we still found a significant deficiency of ABBA patterns in both tests

($D_S > 0$; Table 2, lower part). In agreement with our previous tests involving GeN and HoN as

P1, these additional tests thus equally suggest higher interspecific gene flow among allopatric

(P1 and P3) than among sympatric populations (P2 and P3). As we are much more confident

that PhC and PtC represent truly allopatric P1 ingroups relative to P2 and P3, we think that

the deficiency of ABBA patterns at the Sörenberg site indicates a local reduction in

interspecific gene flow in sympatry, likely due to reinforcement.

When repeating the ABBA-BABA tests with the D-statistic (D_D) suggested by Durand *et*

al. (2011), we obtained results that qualitatively agree with those obtained with the D_S statistic

only when using PhC and PtC as P1 ($D_D < 0$; z scores -8.4 and -7.7, respectively; Table S1

lower part), but not with GeN and HoN as P1 (z-scores 1.4 and 1.9, respectively; Table S1,

upper part). Besides the above qualitative differences between tests based on D_S vs. D_D, we

observed that variances of D_D estimated by jackknifing were about one order of magnitude

larger than the corresponding variances of D_S, which might indicate differences in the

robustness of the two measures. Nonetheless, the qualitatively similar results obtained with

the two versions of D-statistics makes us confident that our findings are robust. We further

observed that windows of elevated D_S (Fig. 2, top) were not homogeneously distributed along

the genome, but aggregated in clusters in several contigs. Such clustering might suggest that

the respective genomic regions contain variation involved in reinforcement caused by

divergent selection against gene flow between the two species.

Since both species commonly co-occur in alpine and sub-alpine regions of Switzerland, we

were curious to see if a signal of reinforcement was also evident at other sites of sympatry in

this geographic area. We therefore investigated a nearby lowland sampling site in Zürich

where we also found *S. cynipsea* and *S. neocynipsea* in sympatry. In contrast to the

sequencing data from Sörenberg, which were generated from wild-caught specimens only, the

sequencing data from the Zürich site derive from a combination of samples from natural and

laboratory populations. Specifically, for *S. cynipsea*, we sequenced a pool of flies sampled

directly from the natural population as well as a pool of flies from a laboratory population that

was derived from the same natural population (see Material and Methods for details); for *S.*

neocynipsea, we only had a pool of flies sampled from a laboratory population derived from

the natural population (Table 1). We found that the ABBA-BABA tests were qualitatively

robust to whether we used *S. cynipsea* data from natural or laboratory populations. However,

results depended on the choice of the D statistic (D_S vs. D_D) and of the P1 ingroup

(Supplementary Information, Supplementary Table S2). All ABBA-BABA tests based on D_S

that included GeN, HoN, or PtC as P1 were significant and revealed a deficiency of ABBA

patterns, whereas tests with PhC as P1 were not significant (Supplementary Table S2). All

significant tests were therefore consistent with our tests for the Sörenberg site in that they

suggested reduced interspecific gene flow in sympatry relative to allopatry. In contrast, only

two out of eight ABBA-BABA tests based on D_S were significant, one suggesting an excess

and one a deficiency of ABBA patterns. Overall, our results for the Zurich site are compatible

with reinforcement, although the signal seems to be weaker compared to the Sörenberg site.

Table 2: Results of ABBA-BABA tests applied to samples originating from sympatric
populations of *S. cynipsea* and *S. neocynipsea* in Sörenberg as P2 and P3 (or vice versa) based
on D_S. The upper half of the table shows tests with two European *S. neocynipsea* populations
from Geschinen (GeN) and Hospental (HoN) as P1 ingroups, and the bottom half shows the
reciprocal approach with two *S. cynipsea* populations from Pehka, Estonia (PhC) and Petroia,
Italy (PtC) as P1 ingroups. The first three columns indicate the phylogeny assumed for the
test, with P4 always being *S. orthocnemis*; columns 4 to 6 give the genome-wide average D_S,
its jackknifed standard deviation SD(D_S), and the corresponding z-score. Significant tests are
shown in bold.
312

P1	P2	P3	D_S	SD(D_S)	z
GeN	**SoN**	**SoC**	**0.008**	**1.35E−03**	**5.8**
HoN	**SoN**	**SoC**	**0.006**	**1.34E−03**	**4.2**
PhC	**SoC**	**SoN**	**0.026**	**1.92E−03**	**13.4**
PtC	**SoC**	**SoN**	**0.039**	**1.97E−03**	**19.8**

Signals of introgression in high altitude populations from Southern France

Complementary to the alpine location in Sörenberg (Switzerland), we further investigated

a pair of sympatric *S. cynipsea* and *S. neocynipsea* populations at high altitude site around Le

Mourier in the Cevennes, Southern France (790 m asl), approximately 450 km southwest of

the sampling site in Switzerland. Interestingly, we found opposite patterns to those observed

at the Sörenberg site (Table 3). Specifically, all tests based on D_S showed a highly significant

excess of ABBA patterns, and hence more interspecific gene flow in sympatry than allopatry.

The corresponding tests based on D_D also indicated an excess of ABBA patterns (recall the

difference in sign between D_S and D_D), but were not significant (Supporting Table S3). In

contrast to the patterns in Sörenberg, we did not observe genomic clustering of extreme

values of D_S (Fig. 2, bottom).

Table 3: Results of ABBA-BABA tests applied to samples originating from sympatric
populations of *S. cynipsea* and *S. neocynipsea* in the French Cevennes as P2 and P3 (or *vice*
versa) based on D_S. As in Tables 2 and 3, the upper half of the table summarizes tests
involving two *S. neocynipsea* populations from Geschinen (GeN) and Hospental (HoN) as P1
ingroups, and the bottom half summarizes tests involving *S. cynipsea* populations from Pehka,
Estonia (PhC) and Petroia, Italy (PtC) as P1.

P1	P2	P3	D_S	SD(D_S)	z
GeN	**MoN**	**MoC**	**–0.014**	**1.30E–03**	**–11.0**
HoN	**MoN**	**MoC**	**–0.014**	**1.36E–03**	**–10.6**
PhC	**MoC**	**MoN**	**–0.030**	**1.84E–03**	**–16.2**
PtC	**MoC**	**MoN**	**–0.016**	**1.93E–03**	**–8.3**

Fig. 2. Variation of D_S along the genome. Dots indicate D_S (Soraggi *et al*. 2018) for genomic windows of 500 consecutive SNPs, oriented by contigs of the *S. thoracica* reference genome with a length ≥ 50,000 bp (highlighted by alternating black and grey colors). The top and bottom panels show results for the population tree topologies used to test for interspecific gene flow at the focal sites in Sörenberg (SoC-SoN; deficiency of ABBA patterns) and Le Mourier (MoC-MoN; excess of ABBA patterns), respectively. The horizontal dashed black and green lines indicate the neutral expectation ($D_S = 0$) and the mean genome-wide jackknife estimate, respectively. Green shading delimits mean $D_S \pm 2$ standard deviations (based on windows of 500 SNPs).

DISCUSSION

We applied the ABBA-BABA approach (Green *et al*., 2010; Durand *et al*., 2011; Soraggi

et al., 2018) to infer historical gene flow between a pair of closely related species of sepsid

flies for which we have detailed information on ecology, morphology, life history, and

behavior (Blanckenhorn, 1999; Blanckenhorn *et al*., 2000; Eberhard, 1999; Ozerov, 2005;

Parker, 1972a,b; Pont & Meier, 2002; Puniamoorthy *et al.*, 2009; Rohner, Blanckenhorn &

Puniamoorthy, 2016; Ward, 1983; Ward, Hemmi & Rösli, 1992; Giesen *et al.*, 2017, 2019).

Previous population genetic analyses of microsatellite markers (Baur *et al.*, 2020) and

hybridization experiments in the laboratory (Giesen *et al.*, 2017, 2019) suggested that *S.*

cynipsea and *S. neocynipsea* are genetically distinct, but may occasionally hybridize in nature

when occurring in sympatry. Here, we tested this hypothesis by analyzing patterns of DNA

sequence variation among pools of field-caught and laboratory specimens from multiple

sampling sites across Europe at which either both species occur in sympatry or only one

species (*S. cynipsea*) occurs. As discussed in detail below, depending on the focal site of

sympatry, we found two qualitatively opposite patterns of interspecific gene flow in sympatry

versus allopatry: a relative reduction of gene flow in sympatry at two sampling sites in

Switzerland, and a relative excess of gene flow in sympatry at a sampling site in Southern

France. Previous studies comparing levels of interspecific gene flow in sympatry and

allopatry in other taxa found the full spectrum of results that we here obtained for one species

pair. Martin *et al.* (2014; see also Nadeau *et al.,* 2013) and Brandvain *et al.* (2014; see also

Grossenbacher & Whittall, 2011) found evidence for higher levels of interspecific gene flow

in sympatry than in allopatry when studying *Heliconius* butterflies and monkey flowers

(*Mimulus guttatus/nasutus*), respectively. In contrast, less gene flow in sympatry than

allopatry was observed in studies on *Drosophila arizonae/mojavensis* (Massie & Makow,

2005) and on two species of sea squirt (Tunicata: Bouchemousse *et al.,* 2016), while no

geographic variation in levels of gene flow between sympatric and allopatric populations was

found in studies on wild tomatoes (Nakazato *et al.,* 2010). A reduction in average gene flow

in sympatry compared to allopatry may suggest reinforcement by natural or sexual selection

(Butlin, 1995; Noor, 1999; Coyne & Orr, 2004; cf. Giesen *et al.*, 2017, 2019). However, we

caution that comparing average levels of gene flow between sympatric and allopatric

population pairs may fall short of detecting selection against interspecific gene flow. In

monkey flowers, there is evidence for divergent selection against gene flow in sympatry (in

terms of a negative correlation between nucleotide divergence and recombination rate) even

though average levels of introgression are higher in sympatry than allopatry (Brandvain *et al.*

2014; Aeschbacher *et al.* 2017).

Similar to research on *Heliconius* (Martin *et al.*, 2014), we found evidence for an excess of

average levels of interspecific gene flow in sympatry compared to allopatry in flies from Le

Mourier based on pool-sequenced field-caught specimens (Table 3; Table S3). In contrast, at

two Swiss sites where the two species also occur in sympatry, we observed reduced average

levels of interspecific gene flow relative to interspecific pairs of parapatric and allopatric

populations, similar to patterns found in *Drosophila arizonae/mojavensis* (Massie & Makow,

2005) and sea squirts (Bouchemousse *et al.*, 2016). Consistent with our previous study

showing behavioral reinforcement after enforced hybridization in the laboratory (Giesen *et*

al., 2017), the genomic data analyzed here suggest that selection against interspecific gene

flow (Noor, 1999) might also occur in nature in the two sympatric Swiss populations. The

reduction of interspecific gene flow was more pronounced at the Sörenberg site in the Swiss

Alps (Table 2; Table S1) than at the lowland site in Zürich (Tables 3; Table S2), but

qualitatively consistent. However, as evidenced by our contrasting findings for the site in Le

Mourier, reinforcement may not be a necessary outcome whenever *S. cynipsea* and *S.*

neocynipsea occur in sympatry. The extent of gene flow between these two dung fly sister

species thus appears to covary with local environmental conditions. Well-documented

differences in climate between Southern France and the Swiss Alps, and between the Swiss

Alps and the Swiss lowlands, might provide an ecological explanation for the qualitative and

quantitative differences in interspecific gene flow we observed at the different locations

(Doebeli & Dieckmann, 2003; Rohner *et al.*, 2015). While average minimum temperatures are

similar in the French Cevennes and the Swiss Alps, average maximum temperatures are up to

5 °C higher at the French site compared to Sörenberg (Figure 3). In addition, both sampling

sites are characterized by very different precipitation regimes. During the main growing

season from May until October, Sörenberg (a peat bog site) has on average twice as much

rainfall than the location in Le Mourier. However, the mechanisms by which climate affects

behavior and/or fitness remain to be understood (Giesen *et al.*, 2017, 2019).

We found a weaker signal of reinforcement between *S. cynipsea* and *S. neocynipsea* at the

lowland Zürich site than at the high altitude Sörenberg site in the Swiss Alps. Differences in

the strength of reinforcement may also result from climatic variation between these two sites

in Switzerland to ultimately influence the strength of selection on natural hybrids. While

precipitation patterns of both locations are comparable, mean temperature is approximately

4 °C lower in Sörenberg than in Zürich (Supplementary Figure 1). Alternatively, differences

between the two Swiss sites might relate to *S. neocynipsea* being rare around Zürich, in fact in

European lowlands in general (Pont & Meier, 2002), while this species is common at higher

alpine altitudes. If based primarily on behavioral mate choice, reinforcement is expected to be

stronger, and hence its evolution more likely, wherever interspecific encounter frequencies

are higher. Moreover, given that the evidence for the Zürich site is based on both wild-caught

and laboratory specimens, the weaker patterns of reinforcement observed in Zürich might be

partially explained by purifying selection in the laboratory against introgressed variants.

Nevertheless, the results from our analyses performed with both wild-caught and laboratory *S.*

cynipsea samples from Zürich yielded consistent results (Supplementary Table 2), suggesting

that whether flies came from a natural or laboratory population did not affect the outcome.

Overall, we therefore do not think that purifying selection in the laboratory confounded our

results.

Fig. 3. Climatic differences between the Sörenberg (Swiss Alps; blue) and Le Mourier
(French Cevennes; red); polygons and dashed lines show monthly average temperature
ranges and precipitation at the two sampling sites, respectively. The data from the WorldClim
dataset (Hijmans *et al*. 2011) represent 50-year averages of observations in quadratic grid
cells with and edge of length 2.5' ($\sim$ 5 km^2 in size) around the coordinates of the two sites
(Table1). The two locations are characterized by strong differences in precipitation, especially
during the main reproductive season of sepsid flies from May to October.

Laboratory hybridization experiments have shown that interspecific mating between *S.*

cynipsea and *S. neocynipsea* can result in viable and fertile F$_1$ hybrid females and offspring in

backcrosses with the parental species (Giesen *et al*., 2017, 2019). Together with

morphological and behavioral similarities among the two species, these laboratory

experiments suggest that hybridization may well take place in areas of co-occurrence, such as

the French Cevennes. In other areas, including our sampling site in the Swiss Alps, subtle

premating behavioral barriers and the reductions in fecundity and fertility also previously

documented in the laboratory by Giesen *et al*. (2017, 2019) may combine with micro-

ecological niche differences to mediate spatio-temporally divergent reproductive timing. Such

a divergence may effectively prevent hybridization in nature and ultimately reinforce species

boundaries. This interpretation is strengthened by a study on Central American *Archisepsis*

diversiformis flies showing that mating between two disjunct populations is only evident

under forced laboratory conditions, while under conditions of free mate choice flies from the

different populations do not interbreed (Puniamoorthy 2014). Behavioral mating barriers

therefore seem to evolve comparatively fast (Gleason & Ritchie, 1998; Puniamoorthy *et al.*,

2009; Puniamoorthy, 2014), especially if species occur in sympatry and reinforcement by

natural or sexual selection can operate (Coyne & Orr, 2004; Seehausen, 2004; Ritchie, 2007).

Our study of introgression patterns among sympatric *S. cynipsea* and *S. neocynipsea*

populations relied on the ABBA-BABA test, which was initially designed for the analysis of

haploid genotypes of one individual from each of four lineages (Green *et al.*, 2010). Recent

extensions of this original approach to allele frequency data by Durand *et al.* (2011) and

Soraggi *et al.* (2018), which incidentally were here shown to not be equally sensitive in

picking up introgression patterns, enabled us to apply the method equivalently to pool-

sequenced laboratory as well as to field-caught specimens (see also Deitz *et al.* 2016). As a

technical innovation, we have here provided a Python implementation of the ABBA-BABA

approaches by Durand *et al.* (2011) and Soraggi *et al.* (2018) that takes a simple 2D matrix of

allele frequencies as input. This implementation allows restricting the ABBA-BABA test to

high-confidence SNPs that pass a set of user-defined filters. Such filtering can reduce

artefacts of pool-sequencing (Schlötterer *et al.* 2014) that may produce false polymorphisms

due to sequencing errors (Futschik & Schlötterer 2011; Cutler & Jensen 2012). In addition,

and in contrast to the software package ANGSD (Korneliussen *et al.* 2014), our

implementation accommodates an outgroup (P4) taxon different from the one used as a

reference to call the SNPs. This feature reduces a potential reference bias if the ingroup taxa

(P1, P2, P3) strongly differ from an outgroup that also serves as reference (Ballouz *et al.*

2019). Our script and detailed instructions for how to use it are available on GitHub

(https://github.com/capoony/ABBA-BABA-4-AF; see also Supporting Information).

In summary, we found evidence that the extent of interspecific gene flow between two closely related sepsid fly species in sympatry relative to allopatry varies with the geographic location of the site. Interspecific gene flow in sympatry appears to be reduced at two Swiss sites, where the two species have likely locally adapted to different (micro-)ecological niches, such as different breeding times or phenologies, and/or where pre- or postmating reproductive barriers (Giesen *et al*., 2017) have been and are subject to stronger reinforcement. Future studies are needed to determine if ecological and/or sexual selection are indeed driving the gene flow patterns we identified, and future genomic research should analyze patterns of genomic differentiation across allopatric populations from Europe as well as North America (Baur *et al*., 2020). It remains unclear why *S. neocynipsea* is rare in Europe, but common in North America, and why *S. cynipsea* abounds around fresh dung all over Europe north of the Alps and apparently outcompetes and ultimately relegates *S. neocynipsea* towards presumably marginal, high altitude habitats (Pont & Meier, 2002). Such analyses would provide further insights into introgression patterns across the species' entire natural range, as well as into the role of local (e.g., longitudinal or latitudinal) adaptation in shaping genome-wide patterns of differentiation.

ACKNOWLEDGEMENTS

Genome resequencing was performed at the Functional Genomics Center Zürich (FGCZ), and we particularly thank FGCZ staff member Jelena Kühn-Georgijevic for her support. We thank the Wolfpack at the University of Zürich for their support and help in maintaining fly cultures, and Patrick Rohner, Juan Pablo Busso, Nalini Puniamoorthy, Anders Kjaersgaard, Toomas Esperk and Cait Dmitriew for contributing samples. B.M., O.N. and L.P. thank the Leibniz association for granting funds to establish the Leibniz graduate school for Genomic Biodiversity Research (GBR), in which context the reference draft genomes of several *Sepsis*

species explored in this study were sequenced and assembled. We particularly thank the GBR

members Jeanne Wilbrandt, Jan Philip Øyen, and Malte Petersen for their contribution to the

assembly pipeline. The University Research Priority Program 'Evolution in Action' of the

University of Zürich, the Swiss National Science Foundation, and the Georges and Antoine

Claraz-Donation funded the work of A.G., the Austrian Science Foundation (grant no. FWF

P32275) the work of M.K.

DATA ACCESSIBILITY

All sequencing data have been deposited at the short-read archive (SRA;

https://www.ncbi.nlm.nih.gov/sra) under the accession number PRJNA612154. Novel

software is available at GitHub (https://github.com/capoony/ABBA-BABA-4-AF).

AUTHORS CONTRIBUTION

AG, WUB, KKS, HELL, SA and MK conceived and designed the study. MAS and AG

gathered and prepared the samples and generated the genomic data. BM, ON and LP

sequenced and constructed the reference genome. MK, AG, SA and HELL analyzed the

genomic data. MK developed new software. AG, MK, SA and WUB wrote the manuscript,

with ON, KKS, RSI, LP, BM, MAS and HELL contributing to the interpretation of the data,

and editing and commenting on various versions of the manuscript. WUB, KKS, RSI, and

HELL were members of AG's PhD committee, providing guidance and input at all stages.

508
LITERATURE CITED

Aeschbacher, S., Selby, J.P., Willis, J.H. & Coop, G. 2017. Population-genomic inference of the
strength and timing of selection against gene flow. *PNAS* 114: 7061–7066.

Anderson M.B. (1949). Sexual selection. *Princeton University Press*.

Andrews, S., Lindenbaum, P., Howard, B., & Ewels, P. (2011). FastQC High throughput
sequence QC report v. 0.10. *Babraham Bioinformatics, Cambridge, UK*.

Arnold M.L. & Meyer A. (2006). Natural hybridization in primates: one evolutionary
mechanism. *Zoology*, 109(4): 261–276.

Ballouz, S., Dobin, A. & Gillis, J.A. (2019). Is it time to change the reference genome? *Genome*
Biology 20: 159.

Barton N., & Bengtsson B.O. (1986). The barrier to genetic exchange between hybridising
populations. *Heredity*, 57(3): 357–376.

Baur J., Schäfer M.A., Blanckenhorn W.U., & Giesen A. (2020). Patterns of genetic
differentiation in sexual and neutral morphological traits among species and populations of dung
flies (Diptera: Sepsidae). *Journal of Zoological Systematics and Evolutionary Research* 58: 159–
173.

Berner D. & Salzburger W. (2015). The genomics of organismal diversification illuminated by
adaptive radiations. *Trends in Genetics*, 31: 491–499.

Blanckenhorn W.U. (1999). Different growth responses to temperature and resource limitation
in three fly species with similar life histories. *Evolutionary Ecology*, 13: 395–409.

Blanckenhorn W.U., Mühlhäuser C., Morf C., *et al*. (2000). Female choice, female reluctance to
mate and sexual selection on body size in the dung fly *Sepsis cynipsea*. *Ethology*, 106: 577–593.

Bolger A.M., Lohse M., & Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*: btu170.

Bouchemousse S., Liautard-Haag C., Bierne N., & Viard F. (2016) Distinguishing
contemporary hybridization from past introgression with postgenomic ancestry-informative SNPs
in strongly differentiated *Ciona* species. *Molecular Ecology*, 25: 5527-5542.

Brandvain Y., Kenney A.M., Flgel L., Coop G., & Sweigart A.L. (2014). Speciation and
introgression between *Mimulus nasutus* and *M*. *guttatus*. *Plos Genetics*, 10:
doi:10.1371/journal.pgen.1004410

Butlin R. K. 1995. Reinforcement: an idea evolving. *Trends in Ecology & Evolution*, 10, 432-
434.

Coyne J.A., & Orr H.A. (2004). Speciation. *Sinauer Associates*.

Cutler, D.J. & Jensen, J.D. (2010). To Pool, or Not to Pool? *Genetics* 186: 41–43.

Deitz, K.C., Athrey, G.A., Jawara, M., Overgaard, H.J., Matias, A. & Slotman, M.A. (2016).
Genome-Wide Divergence in the West-African Malaria Vector Anopheles melas. *G3: Genes,*
Genomes, Genetics 6: 2867–2879.

DeMarais B.D., Dowling T.E., Douglas M.E., *et al*. (1992). Origin of Gila seminude (Teleostei:
Cyprinidae) through introgressive hybridization: implications for evolution and conservation.
Proceedings of the National Academy of Sciences 89(7): 2747–2751.

Dobzhansky T. (1951). Genetics and the origin of species (3rd ed.). *Columbia University Press*.

Doebeli M, & Dieckmann U. (2003) Speciation along thermal gradients. Nature 421: 259–264.
Durand E.Y., Patterson N., Reich D., & Slatkin M. (2011). Testing for ancient admixture
between closely related populations. *Molecular Biology and Evolution*, 28(8): 2239–2252.

Eaton D.A. & Ree R.H. (2013). Inferring phylogeny and introgression using RADseq data: an
example from flowering plants (Pedicularis: Orobanchaceae). *Systematic Biology*, 62(5): 689–706.

Eberhard W.G. (1999). Mating systems of sepsid flies and sexual behavior away from
oviposition sites by *Sepsis neocynipsea* (Diptera: Sepsidae). *Journal of the Kansas Entomological
Society,* 72(1): 129–130.

Feulner P.G.D & Seehausen O. (2018). Genomic insights into the vulnerability of sympatric
whitefish species flocks. *Molecular Ecology*, 28: 615-629.

Futschik A. & Schlötterer C. (2010). The next generation of molecular markers from passively
parallel sequencing of pooled DNA samples. *Genetics*, 186(1): 207–218.

Gante H.F., Matschiner M., Malmstrom M., Jakobsen K.S., Jentoft S., & Salzburger W. (2016).
Genomics of speciation and introgression in Princess cichlid fishes from Lake Tanganyika.
Molecular Ecology, 25: 6143–6161.

Giesen A, Blanckenhorn W.U., & Schäfer M.A. (2017). Behavioural mechanisms of
reproductive isolation between two hybridizing dung fly species. *Animal Behaviour*, 132: 155–166.

Giesen A., Schäfer M.A., & Blanckenhorn W.U. (2019). Geographic patterns of postzygotic
isolation between two closely related widespread dung fly species (*Sepsis cynipsea* and *Sepsis
neocynipsea*; Diptera: Sepsidae). *Journal of* Zoological Systematics and Evolutionary Research,
57: 80–90

Gleason, J.M. & Ritchie, M.G. 1998. Evolution of courtship song and reproductive isolation in
the *Drosophila willistoni* species complex: Do sexual signals diverge the most quickly? *Evolution*,
52: 1493–1500.

Green R.E., Krause J., Briggs A.W., *et al*. (2010). A draft sequence of the Neanderthal genome.
Science, 328(5979): 710–722.

Grossenbacher D.L. & Whittall J.B. (2011). Increased floral divergence in sympatric
monkeyflowers. *Evolution*, 65: 2712-2718.

Haldane J.B.S. (1922). Sex ratio and the unisexual sterility in animal hybrids. *Journal of
Genetics* 12: 101–109.

Heliconius Genome Consortium (2012). Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. *Nature*, 487(7405): 94–98.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. 2005. Very high resolution
interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**: 1965–1978.

Hudson R.R. (1983). Testing the Constant-Rate Neutral Allele Model with Protein Sequence
Data. *Evolution* 37: 203–217.

Jeong C., Nakagome S., Di Rienzo A. (2016). Deep History of East Asian Populations Revealed
Through Genetic Analysis of the Ainu. *Genetics*, 202: 261–272.

Kapun, M., Barron Aduriz, M.G., Staubach, F., *et al*. (2018). Genomic analysis of
European *Drosophila* populations reveals major longitudinal structure, continent-wide selection,
and unknown DNA viruses. *bioRxiv*, DOI: 10.1101/313759.

591  Kastally Ch., Trasoletti M., & Mardulyn P. (2019). Limited gene exchange between two sister
592  species of leaf beetles within a hybrid zone in the Alps. *Journal of evolutionary Biology,* 32: 1406-
593  1417.

594  Kimura, M. (1969). The Number of Heterozygous Nucleotide Sites Maintained in a Finite
595  Population Due to Steady Flux of Mutations. *Genetics* 61: 893–903.

596  Kofler, R., Pandey, R.V. & Schlötterer, C. (2011). PoPoolation2: identifying differentiation
597  between populations using sequencing of pooled DNA samples (pool-seq). *Bioinformatics* 27:
598  3435–3436.

599  Koren S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H. & Phillippy, A.M. (2017).
600  Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat
601  separation. *Genome Research* 27: 722–736.

602  Kulathinal R.J. & Singh R.S. (2000). A biogeographic genetic approach for testing the role of
603  reinforcement: the case of *Drosophila pseudoobscura* and *D. persimilis*. *Evolution,* 54: 210-217.

604  Kraushaar U. & Blanckenhorn W.U. (2002). Population variation in sexual selection and its
605  effect on size allometry in two dung fly species with contrasting sexual size dimorphism.
606  *Evolution*, 56(2): 307–321.

607  Kraushaar U., Goudet J., & Blanckenhorn W.U. (2002). Geographical and altitudinal population
608  genetic structure of two dung fly species with contrasting mobility and temperature preference.
609  *Heredity*, 89: 99-106.
610  Korneliussen T.S., Albrechtsen A., & Nielsen R. (2014). ANGSD: analysis of next generation
611  sequencing data. *BMC bioinformatics*, 15(1): 356.

612  Li H., Handsaker B., Wysoker A., *et al*. (2009). The sequence alignment/map format and
613  SAMtools. *Bioinformatics*, 25(16): 2078–2079.

614  Li H. & Durbin R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
615  transform. *Bioinformatics*, 25(14): 1754–1760.

616  Mallet J. (2007). Hybrid speciation. *Nature*, 446: 279–283.

617  Martin S.H., Dasmahapatra K.K., Nadeau N.J., *et al*. (2014). Genome-wide evidence for
618  speciation with gene flow in *Hiliconius* butterflies. *Genome Research*, 23: 1817-1828.

619  Massie K.R. & Markow T.A. (2005). Sympatry, allopatry and sexual isolation between
620  *Drosophila mojavensis* and *D. arizonae*. *Hereditas*, 142: 51-55.

621  Mayr E. (1942). Systematics and the origin of species from the viewpoint of a zoologist.
622  *Columbia University Press*.

623  Mayr E. (1963). Animal and species evolution. In: Animal species and their evolution, *Harvard
624  University Press*.

625  McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., et al. 2010.
626  The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
627  sequencing data. *Genome Research* 20: 1297–1303.

628  Meier J.I., Marques D.A., Selz O.M., *et al*. (2017). Ancient hybridization fuels rapid cichlid fish
629  adaptive radiations. *Nature Communications*, 8: 14363.

Moran, P.A., Hunt J., Mitchell Ch., Ritchie M.G., & Bailey N.W. 2019. Behavioural
mechanisms of sexual isolation involving multiple modalities and their inheritance. *Journal of*
evolutionary Biology, 32: 243-258.

Nadeau N.J., Martin S.H., Kozak K.M., *et al*. (2013). Genome-wide patterns of divergence in
gene flow across a butterfly radiation. *Molecular Ecology*, 22: 814-826.

Nolte A. W., & Tautz D. (2010). Understanding the onset of hybrid speciation. *Trends in*
Genetics, 26(2): 54–58.

Nakazato T., Warren D.L., & Moyle L.C. (2010). Ecological and geographic modes of species
divergence in wild tomatoes. *American Journal of Botany*, 97: 680-693.

Nei M. (1987). *Molecular Evolutionary Genetics*. Columbia University Press.

Noor M.A.F. (1999). Reinforcement and other consequences of sympatry. *Heredity*, 83: 503-
508.

Nosil P. (2009). Adaptive population divergence in cryptic color-pattern following a reduction
in gene flow. *Evolution*, 63(7): 1902–1912.

Novikova Y.P. *et al*. (2016). Sequencing of the genus *Arabidopsis* identifies a complex history
of non-bifurcating speciation and abundant trans-specific polymorphism. Nature Genetics 48:
1077–1082.

Ozerov A.L. (2005). World catalogue of the family Sepsidae (insect: Diptera).*Zoologicheskie*
Issledovania (Zoological Studies), 8: 1–74.

Parker G.A. (1972a). Reproductive behaviour of Sepsis cynipsea (L.) (Diptera: Sepsidae) I. a
preliminary analysis of the reproductive strategy and its associated behaviour patterns. *Behaviour*,
41(1): 172–205.

Parker G.A. (1972b). Rerpductive behaviorur of Sepsis cynipsea (L.) (Diptera: Sepsidae) II. the
significance of the precopulatory passive phase and emigration. *Behaviour*, 41(3): 242–250.

Pont A.C., & Meier R. (2002). *The Sepsidae (Diptera) of Europe*. Leiden, N.E.: *Fauna*
Entomologica Scandinavia, 37.

Puniamoorthy N, Ismail M.R.B., Tan D.S.H., & Meier R. (2009). From kissing to belly
stridulation: Comparative analysis reveals surprising diversity, rapid evolution, and much
homoplasy in the mating behaviour of 27 species of sepsid flies (Diptera: Sepsidae). *Journal of*
Evolutionary Biology, 22(11): 2146–2156.

Puniamoorthy N. (2014). Behavioural barriers to reproduction may evolve faster than sexual
morphology among populations of a dung fly (Sepsidae). *Animal Behaviour*, 98: 139–148.

Reich, D., N. Patterson, M. Kircher, F. Delfin, M. R. Nandineni et al., 2011 Denisova admixture
and the first modern human dis- persals into Southeast Asia and Oceania. *American Journal of*
Human Genetics, 89: 516–528.

Rieseberg L.H., Raymond O., Rosenthal D.M., *et al*. (2003). Major ecologocial transitions in
wild sunflowers facilitated by hybridization. *Science*, 301(5637): 1211–1216.

Ritchie M.G. (2007). Sexual selection and speciation. *Annual Review of Ecology, Evolution and*
Systematics, 38: 79–102.

Rohner P.T., Bächli, G., Paltrinieri, L.P., Duelli, P., Obrist, M.K., Jochmann, R., &
Blanckenhorn, W.U. (2015). Distribution, diversity gradients and Rapoport's elevational rule in the
black scavenger flies of the Swiss Alps (Diptera: Sepsidae). *Insect Conservation and Diversity*, 8,
367–376.

Rohner P.T., Blanckenhorn W.U., & Puniamoorthy N. (2016). Sexual selection on male size
drives the evolution of male-biased sexual size dimorphism via theprolongation of male
development. *Evolution*, 70(6): 1189–1199.

Dafu R., Kangshan M., Lei Z., *et al*. (2016). Genomic evidence for polyphyletic origins and
interlineage gene flow within complex taxa: a case study of *Picea brachytyla* in the Qinghai-Tibet
Plateau. *Molecular Ecology*, 25: 2373–2386.

Saetre G.P. (2013). Hybridization is important in evolution, but is speciation? *Journal of
Evolutionary Biology*, 26(2): 256–258.

Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. (2014). Sequencing pools of individuals —
mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics* 15: 749–
763.

Seehausen O. (2004). Hybridization and adaptive radiation. *Trends in Ecology and Evolution*,
19(4): 198–207.

Soraggi, S., Wiuf, C. & Albrechtsen, A. (2018). Powerful Inference with the D-Statistic on
Low-Coverage Whole-Genome Data. *G3: Genes, Genomes, Genetics* **8**: 551–566.

Streicher J.W., Devitt T.J., Goldberg C.S., *et al*. (2014). Diversification and asymmetrical gene
flow across time and space: lineage sorting and hybridization in polytypic barking frogs. *Molecular
Ecology*, 23(13): 3273–3291.

Su, K.F.Y., S. N. Kutty, & R. Meier (2008). Morphology versus molecules: the
phylogenetic relationships of Sepsidae (Dptera: Cyclorrhapha) based on morphology and
DNA sequence data from ten genes. *Cladistics* 24:902-916.

Su, K.F.Y., Puniamoorthy, J., Özsu, N., A. Srivathsan, & R. Meier (2016). Evolutionary
analysis identifies multiple genome expansions and contractions in Sepsidae (Diptera) and suggests
targets for future genomic research. *Cladistics* 32: 308–316.

Tajima, F. (1983). Evolutionary Relationship of DNA Sequences in Finite Populations.
Genetics 105: 437–460.

Trier C. N., Hermansen J. S., Sætre G. P., & Bailey R. I. (2014). Evidence for mitonuclear and
sex-linked reproductive barriers between the hybrid Italian sparrow and its parent species. *PLoS
Genet*, 10(1): e1004075.

Via S. & West J. (2008). The genetic mosaic suggests a new role for hitchhiking in ecological
speciation. *Molecular Ecology*, 17(19): 4334–4345.

Ward P.I. (1983). The effects of size on the mating behaviour of the dung fly *Sepsis cynipsea*.
Behavioural Ecology and Sociobiology, 13(1): 75–80.

Ward P.I., Hemmi J., & Rösli T. (1992). Sexual conflict on the dung fly Sepsis cynipsea.
Functional Ecology, 6(6): 649–653.

Wood H.M., Grahame J.W., Humphray S., *et al*. (2008). Sequence differentiation in regions
identified by a genome scan for local adaptation. *Molecular Ecology*, 17(13): 3123–3135.

Wu C.I. (2001). The genic view of the process of speciation. *Journal of Evolutionary Biology*,
14(6): 851–856.