Atypical chemoreceptor arrays accommodate high membrane curvature

Alise R. Muok[1], Davi R. Ortega[2,3], Kurni Kurniyati[4], Wen Yang[1], Adam Sidi Mabrouk[1],

Chunhao Li[4], Brian R. Crane[5], Ariane Briegel[1*]

[1]Institute for Biology, Leiden University, Sylviusweg 72, 2333 BE Leiden

[2]Department of Biology, California Institute of Technology, 1200 E. California Blvd., Pasadena, CA 91125, USA

[3]Laboratory for Research in Complex Systems, 425 California St., Suite 1200, San Francisco, CA 94104, USA

[4]Department of Oral and Craniofacial Molecular Biology, Philips Research Institute for Oral Health, Virginia

Commonwealth University, Richmond, VA 23298

[5]Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14850

Abstract/Summary

The prokaryotic chemotaxis system is arguably the best-understood signaling pathway in

biology, but most insights have been obtained from only a few model organisms and many

studies have relied on artificial systems that alter membrane curvature[1–3]. In all previously

described species, chemoreceptors organize with the histidine kinase (CheA) and coupling

protein (CheW) into a hexagonal (P6 symmetry) extended array that is considered universal

among archaea and bacteria[4,5]. Here, for the first time, we report an alternative symmetry

(P2) of the chemotaxis apparatus that emerges from a strict linear organization of CheA in

Treponema denticola cells, which possesses arrays with the highest native curvature

investigated thus far. Using cryo-ET, we reveal that the *Td* chemoreceptor arrays assume a

truly unusual arrangement of the supra-molecular protein assembly that has likely evolved to

accommodate the high membrane curvature. The arrays have several additional atypical

features, such as an extended dimerization domain of CheA and a variant CheW-CheR-like

fusion protein that is critical for maintaining an ordered chemosensory apparatus in an

extremely curved cell. Furthermore, the previously characterized *Td* oxygen sensor ODP

influences array integrity and its loss substantially orders CheA. These results suggest a

greater diversity of the chemotaxis signaling system than previously thought and

demonstrate the importance of examining transmembrane systems *in vivo* to retain native

membrane curvature.

Introduction

Chemotaxis is a behavior most motile bacteria employ to sense their chemical environment

and navigate toward favorable conditions. The main components of the system are

transmembrane chemotaxis receptors called methyl-accepting chemotaxis proteins (MCPs),

the histidine kinase CheA, and the adaptor protein CheW. The intracellular tips of MCPs

bind CheA and CheW and communicate changes from the external chemical environment

into the cell by modulating CheA kinase activity (Fig. S1A)[6–8]. Activation of CheA initiates an

intracellular phosphorelay that ultimately controls flagellar rotation and cell movement.

CheA functions as a dimer and possesses five domains (P1-P5) with distinct roles in

autophosphorylation and array integration. The P1 domain is the phosphate substrate

domain, P2 interacts with response regulators, P3 is the dimerization domain, P4 binds ATP,

and P5 interacts with CheW. In the model species *Escherichia coli* (*Ec*), CheA P5 and CheW

are paralogs that interact pseudo-symmetrically to form six-subunit rings. In all bacterial and

archaeal species examined thus far, the MCPs are arranged in a trimer-of-dimer oligomeric

state and further organize into a hexagonal lattice (Fig. S1B). In *Ec*, the receptors are

connected by the highly ordered rings of CheA and CheW bound to the cytoplasmic tips of

the receptors[5,9]. These insights have established a widely accepted central model of the

chemotaxis array (Fig. S1B)[4,5,10]. However, emerging research has recently revealed

divergent components and arrangements of the chemotaxis apparatus in non-canonical

organisms. For example, in *Vibrio cholerae* (*Vc*) chemotaxis arrays, CheA and CheW lack

an ordered arrangement in the rings[1,11].

Many of these structural insights have transpired from cryo-electron tomography (cryo-ET)

studies that utilize artificial systems for higher resolution data. Specifically, the advent of so-

called 'mini-cell' bacterial strains produce extremely small cells that are ideal for cryo-ET[1,3],

and lipid-templating methods generate *in vitro* arrays with increased conformational

homogeneity[2,12]. However, these methods generate arrays with non-native curvature and it

is unclear how this may affect array structure and behavior. Here, we examine the *in vivo*

array structure of the pathogenic spirochete *Treponema denticola* (*Td*) by cryo-ET, which

possesses the highest native membrane curvature of any bacteria examined for their

chemotaxis system so far[4,5]. We demonstrate the presence of a novel array architecture with

two-fold (P2) symmetry in *Td*, that likely owes to the high curvature of the cells. Genetics

experiments, bioinformatics analyses, structural investigations, and molecular modeling of

Td chemotaxis proteins reveals adaptations that have likely evolved to accommodate

formation of an extended chemotaxis array in a highly curved membrane. We demonstrate

that a CheR-like fusion domain in a variant CheW is key for maintaining the structural

integrity of the arrays. Furthermore, cryo-ET analysis of *Td* cells lacking the oxygen sensor

ODP reveals substantial changes to the ordering or mobility of CheA[13]. Collectively, these

data demonstrate a greater diversity of the chemotaxis system than previously assumed,

and exemplify the importance of examining biological structures in native *in vivo* conditions.

Results

Conservation of the F2 chemotaxis system

The chemotaxis systems in prokaryotes have been classified into 19 systems based on

phylogenomic markers[14]. These classes include 17 'Flagellar' systems (F), one 'Alternative

Cellular Function' system (ACF), and one 'Type Four Pilus system' (TFP). The spirochete

chemotaxis system belongs to the 'Flagella class 2' (F2) category, which has not been

investigated with structural methods[14]. To explore the characteristics of chemotaxis proteins

in F2 genomes, we analyzed genomes in the Microbial Signal Transduction Database

version 3 (MiST3)[15]. All genomes (306) with at least one CheA of the F2 class (CheA-F2)

are in the Spirochaetota phylum, with a few exceptions of lateral gene transfer to other phyla

(Dataset 1, see Supplementary Methods). Not all Spirochaetota bacteria possess an F2

system however. There are 1096 genomes assigned to the Spirochaetota phylum in GTDB[16]

and 804 of these are present in the MiST3 database (Dataset 2)[15]. With these 804 genomes,

we mapped the different classes of CheA kinases to the Spirochaetota taxonomy tree

(Figure S1). Based on the topology of this tree, it appears that the major chemosensory

systems in the genomes from the Spirochaetota phylum are: F1 (Leptospirae), a transitional

hybrid F1/F2 system (Brachyspirae), and F2 (Spirochaetia) (Fig. S2). Therefore, we

conclude that F2 systems are exclusive to the Spirochaetia class, with a few exceptions of

lateral gene transfer.

The main architectural difference of the F2 system compared to others is the presence of an

unusual scaffold protein that consists of an N-terminal CheW domain and a C-terminal

CheR-like domain, hereafter referred to as CheW-CheR$_{like}$. Typically, CheR is a

methyltransferase that, together with the methylesterase CheB, controls the methylation

state of the receptors and thus provides an adaptation system[17]. Our analyses indicate that

all Spirochaetia genomes in MiST3 contain CheW-CheR$_{like}$. Furthermore, if we limit our

analysis to genomes that are fully sequenced (see Supplementary Methods), CheW-CheR$_{like}$

is only found in F2 chemosensory systems.

To investigate sequence patterns in the CheR protein and the CheR-like domain of CheW-

CheR$_{like}$, we produced a sequence dataset with 83 CheR-F2 and 88 CheW-CheR$_{like}$ proteins

and summarized them in sequence logos (Fig. S3A). Although the key catalytic residues are

conserved in the typical CheR protein, two residues that are essential for CheR to methylate

chemoreceptors, R79 and Y218 in *Td* CheR, are modified in the CheW-CheR$_{like}$ protein

(R79W and Y218F)[17]. Furthermore, the conserved region at the C-terminus of CheR is not

conserved in CheW-CheR$_{like}$. Based on these results we speculate that the CheR$_{like}$ domain

does not possess methyltransferase activity. Collectively, our analyses suggest that CheR

and CheW-CheR$_{like}$ have different biological functions.

F2 systems contain three proteins with a CheW domain: the classical scaffold CheW, CheW-

CheR$_{like}$, and the P5 domain from the histidine kinase CheA. Phylogenetic mapping of

CheW-F2 proteins indicate the location of last common ancestor (Fig. S4, see

Supplementary methods). To investigate sequence patterns of the three CheW domains, we

analyzed non-redundant sequence datasets of CheW, CheW-CheR$_{like}$, and CheA-F2 from all

genomes with at least one CheA-F2. The final alignments for each group contain the CheW

domain portion of 74 CheW proteins, 59 CheW-CheR$_{like}$ proteins, and 73 CheA proteins. The

sequences of each group are summarized in sequence logos and demonstrate the presence

of conserved regions at established interaction interfaces, as well as loop insertions near

these interfaces that could confer altered specificity of binding (Fig. S3B).

The structure of the *Treponema denticola* (*Td*) chemotaxis array in wild-type cells

Cell poles of intact *Td* cells were imaged by cryo-electron tomography (cryo-ET) and used

for three-dimensional reconstructions. Top views (cross-sections through the array) and side

views (visualizing the long axes of the receptors) of membrane-associated arrays were

clearly visible (Fig. 1A, S5A). Sub-volume averaging revealed the conserved receptor trimer-

of-dimers in the typical hexagonal arrangement. Remarkably, several novel features of the

chemotaxis arrays are apparent (Fig. 1B). Specifically, a density of unknown origin is located

in the center of the receptor hexagons and slightly above the plane of the CheA:CheW rings.

This density, which will hereafter be referred to as the middle density, extends from two

subunits in the rings (Fig. 2A). Additionally, there are small but distinct puncta of density in

between some of the trimer-of-dimer modules (Fig. 1B). However, averages of the arrays at

the CheA:CheW layer did not reveal discernible CheA density, indicating either a sparse or

disordered distribution of CheA or a highly mobile kinase (Fig. 1C). Remarkably, the sub-

tomogram averages reveal the axis of the *Td* cells relative to the chemotaxis arrays,

demonstrating that the arrays occupy a preferred orientation with respect to the cell axis

(Fig. S5B).

Arrays in *T. denticola* deletion mutants

The oxygen-binding diiron protein (ODP) functions as an oxygen sensor for chemotaxis in

Td, however it is uknown if ODP is an integral component of the array[13]. This protein is

genetically coupled to an MCP homolog, TDE2496, that lacks both transmembrane and

sensing modules. TDE2496 likely integrates into the cytoplasmic regions of the membrane-

bound arrays based on the observation that no cytoplasmic arrays were observed in the

tomograms. Moreover, the *Td* genome encodes only one CheA homolog, and cytoplasmic

receptors often associate with distinct kinases[9,18]. TDE2496 forms trimers-of-dimers *in vitro*

and can modulate CheA activity (Fig. S6A,B). To determine if the presence of ODP

(TDE2498) and its cognate receptor (TDE2496) impacts array architecture or integrity, we

conducted cryo-ET with two *Td* gene knock-out strains, Δ2498 and Δ2498Δ2496[13].

Importantly, deletion of ODP does not impact transcription of TDE2496[13]. The sub-volume

averages of these strains reveal distinct differences in array densities compared to the wild-

type (WT) strain (Fig. 1B,C, Fig. S5A). Namely, the location of CheA at the CheA:CheW

layer (Layer 2, Fig 1C) is now clearly visible. Interestingly, CheA arranges in well-ordered

linear rows. Placement of CheA necessarily positions the P3 domain in between two of the

trimer-of-dimer modules in each hexagon. This position exclusively corresponds to the

Fig. 1 Cryo-electron tomography of whole *T. denticola* (*Td*) cells reveals the protein arrangement of chemotaxis machinery. (A) Side-views of the membrane-associated chemotaxis apparatus illustrate the location of the receptor layer (Layer 1) and CheA:CheW baseplate (Layer 2). These layers are spaced ~90 Å from one another. (B) Sub-volume averaging of three *Td* strains reveals the universally-conserved receptor trimer-of-dimer arrangement with 12 nm spacing between opposing trimer-of-dimer modules. Notably, density is apparent in the center of the receptor hexagons (blue arrow) and between some receptor trimer-of-dimer modules (green arrow). (C) Sub-volume averages at Layer 2 reveal the organization of CheA. In this arrangement, the density between the trimer-of-dimer modules (green arrow) corresponds to the CheA P3 domain.

location of the puncta between receptor trimer-of-dimer modules observed in the WT array,

indicating that this density represents the P3 domain (Layer 1, Fig. 1B). Like the WT arrays,

the cell axis relative to the chemotaxis arrays is also apparent in these sub-tomogram

averages and matches the preferred array orientation in the WT cells (Fig. S5B).

Analyses of the CheW-CheR$_{like}$ protein in _T. denticola_

To determine the composition of the middle density in the _Td_ arrays, we considered the

unique chemotaxis proteins in spirochete

genomes. As shown above, CheW-CheR$_{like}$ is

a conserved component of the F2 Spirochaetia

chemotaxis system (Fig. S2). In _Td,_ a 28-

residue linker in CheW-CheR$_{like}$ separates the

two domains and is predicted to be largely

helical (JPred), and the gene is co-transcribed

with the only CheA, CheX, and CheY proteins

in the genome (Fig. S6C,D)[18]. Furthermore,

the purified CheW-CheR$_{like}$ protein dimerizes

and interacts with CheA, as evidenced by its

ability to alter autophosphorylation activity (Fig.

2C,D, Fig. S6E,F). Therefore, we postulated

that the middle density may be comprised of

two dimerized CheR$_{like}$ domains that extend

from the CheW units harbored within the

CheA:CheW rings.

A _Td_ strain lacking the CheR$_{like}$ domain

(ΔCheR$_{like}$) reveals a significant decrease in

Fig. 2 Investigations into the identity of the density located in the center of the receptor hexagons, referred to as the middle density. (A) In Layer 1, the middle density extends to two positions in the CheA:CheW rings for all three _Td_ strains (red arrows). (B) Deletion of a CheR-like domain that is covalently attached to the C-terminus of a CheW domain generates sub-tomogram averages that lack the middle density feature. (C) Radioisotope assays that monitor CheA autophosphorylation with [γ-^{32}P]ATP demonstrate that CheW-CheR$_{like}$ influences CheA activity _in vitro_. Bands are from non-adjacent lanes on the same gel. (D) Purification of CheW-CheR$_{like}$ followed by Native PAGE gel analysis reveals the presence of a protein monomer (49 kDa) and protein dimer (98 kDa). Bands are from non-adjacent lanes on the same gel.

the prevalence and size of the arrays (Fig. S5A). Due to the small size of the arrays in

ΔCheR$_{like}$, only 194 particles were available for sub-tomogram averaging, but the resulting

averages clearly demonstrate that the middle density is no longer present (Fig. 2B, S5A).

Like the WT strain, the CheA density below the rings in ΔCheR$_{like}$ is not apparent.

Protein interfaces in the CheA:CheW:CheW-CheR$_{like}$ rings

Bioinformatics analyses demonstrate that all functional Spirochaetota F2 chemotaxis

systems possess a CheW-CheR$_{like}$ homolog and at least one classical CheW protein. As

only two of the CheW subunits in the hexagonal rings extend to the middle density (which

probably arises from the CheW-CheR$_{like}$ protein), and two of the ring positions are occupied

by CheA P5, it is likely that the other two positions are occupied by the classical CheW

protein. (Fig. 2A, S7). Within this arrangement three unique interaction interfaces are

possible, interface 1 occurs between CheA P5 and the classical CheW (as seen in canonical

systems), interface 2 occurs between CheA P5 and the CheW domain of CheW-CheR$_{like}$,

and a third interface (interface 3) occurs between the classical CheW and the CheW domain

of CheW-CheR$_{like}$ (Fig. S7). To explore the binding interfaces within the *Td* rings, we

analyzed homology models of the classical CheW, the CheW domain of CheW-CheR$_{like}$, and

the CheA P5 domain. The CheW models were generated using a crystal structure of

Thermoanaerobacter tengcongensis (*Tt*) CheW as the template (PDB ID: 2QDL), and the

CheA P5 model was generated using a cryo-EM structure of *E. coli* CheA P5 (PDB ID:6S1K)

(Fig. S8A-C)[2,19]. Three of the four regions with lowest sequence conservation among the

three domains are located at interfaces 1-3 (Fig. S9). Alignment of the *Td* CheW and CheA

P5 models to a crystal structure of *Tm* CheW in complex with *Tm* CheA P5 (PDB ID: 3UR1)

further illustrates that these regions are located at the CheW:P5 ring interfaces (Fig S9)[20].

Mapping the variable regions onto the sequence logos of the F2 CheW domains

demonstrates that they evolved different sequence patterns in these regions, with the

exception of the variable region that is not located at the interaction interface (region 2, Fig.

S9, S3B).

CheA arrangement and array curvature in *T. denticola*

Sub-tomogram averaging reveals that *Td* CheA forms a linear arrangement across the

chemotaxis array, linking the CheA:CheW rings into extended 'strands' that are held

together by receptor:CheA/W interactions (Fig. 3A,B). The orientation of the strands are

apparent in the sub-tomogram averages and run relatively perpendicular to the axis of the

cells (Fig. S10A). The array densities at Layer 1 (receptors, P3, and CheR$_{like}$) produce

apparent lines in the cryo-ET reconstructions that, like the sub-tomogram averages, run

parallel to the cell axis (Fig. 3C,D)[21]. This arrangement allows interactions that hold the

strands together to occur with minimal bending (Fig. 3E,F). Indeed, the angle between the

cell axis and these lines in the cells is 10.4 +/- 8.6°, (*n* = 26 cells) and no significant

difference was found among the three *Td* strains measured (Table S1A). As the

CheA:CheW strands are perpendicular to these lines in the sub-tomogram averages, they

will be oriented at an angle of 55.4 +/- 8.6° with respect to the cell axis (Fig. S10A).

Previous studies in *Ec* and *Vc* demonstrate that the chemotaxis arrays can

accommodate vastly different curvatures between lysed and artificially small mini-cell

strains[1,3,22]. However, *Td* cells demonstrate a significantly higher curvature of the cell

membrane than organisms investigated for their chemotaxis arrangement thus far [1,5,22]. As a

measure of comparison, the *Vc* mini-cells used in a previous study have an inner membrane

curvature of 9.15 +/- 4.5 /μm (radius 1092 Å, *n* = 6 cells), and the *Td* cells have an inner

membrane curvature of 35.8 +/- 6.6 /μm (radius: 279 Å, *n* = 10 cells) (Fig. S10B,C, Table

S1B,C). Additionally, the measured curvature of the *Td* CheA:CheW baseplate is 65.6 +/- 19

/μm (radius: 152 Å, *n* = 10 cells) (Fig. S10B, Table S1C).

The CheA:CheW rings present in a crystal structure (PDB ID: 3UR1) are flat and ~95 Å in

diameter (Fig. S11A)[20]. The length across two rings connected by a dimeric CheA is 224 Å

(Fig. 11B). To determine the extent of bending that would need to occur in the two

CheA:CheW rings if they ran perpendicular to the cell axis, the 224 Å rings were modeled as

a chord in a circle with radius 152 Å. Using the equation $h = r - \sqrt{r^2 - L^2}$ (where h is the

height of the circular segment, r is the circle radius, and L is half the chord length (224 Å /2)),

the height of the circular segment is 49.2 Å (Fig. S11B). Therefore, the center of the two

rings (the P3 domain) must bend by an average of 49.2 Å toward the cell membrane to

accommodate the cell curvature. Using the same equation above (where L is 95 Å /2), the

height of the circular segment is 7.6 Å (Fig. S11A). Therefore, in *Td,* the center of a single

ring must bend toward the membrane by an average of 7.6 Å to align to the measured

baseplate curvature.

Fig. 3 The organization of CheA in *Td* differs from all previously reported arrangements. (A) The hexagonal arrangement of CheA in *E. coli* and other canonical chemotaxis systems. (B) In *Td,* CheA is arranged with a strict linear organization, producing 'strands' of CheA:CheW rings linked by the CheA P3 domain. (C) The receptor densities in *Ec* (Layer 1) are apparent and hexagonal. Figure adapted from Briegel A. et al. 2010.[21] (D) The densities at Layer 1 in *Td* (red oval) produce apparent lines of receptors, P3 and the middle density (CheR-like) that follow the axis of *Td* cells (black arrow). (E) *Ec* arrays possess radial P6 symmetry where all interaction interfaces are equally curved. (F) The *Td* array has a two-fold symmetry arrangement that allows receptor:CheA/W interactions that hold the CheA:CheW strands together to occur with minimal bending. Dashed lines represent the apparent lines in the tomograms.

Spirochetes possess an atypical dimerization domain

The cryo-ET results reveal density corresponding to the P3 domain, which has not been previously reported in *in vivo* arrays. Sequence alignments of *Td* CheA with CheA homologs from a variety of model bacteria with previously characterized chemotaxis systems indicate that in *Td* CheA an additional ~50 residues join the canonical dimerization domain (P3) helices (Figure S12A)[4]. CheA homologs from other spirochete genera including Borrelia and Brachyspira also possess additional residues in this region (Fig. S12B)[23]. Analysis of non-redundant P3 domains from all CheA classes reveal general sequence conservation in the canonical helices but highly divergent sequences at these additional residues (Fig. S13A). Furthermore, CheA-F2 proteins contain the most residues in this non-conserved region (Fig. S13B,C). The x-ray crystal structure of the isolated *Td* P3 domain (PDB ID: 6Y1Y, Fig. 4, Table S2) reveals that the additional residues adopt the coil-coiled motif of the classic dimerization domain with the exception of a break in one of the helices, producing a discontinuous coiled-coil (Fig. 4, S14A). Interestingly, aromatic residues (Phe, Tyr) cluster near the helix breakages and unusual core packing of these residues allows for maintenance of a coiled-coil register despite a disruption of helical heptad repetition in the C-terminal helix (Fig. S14A). The net result is a distortion in the alignment of the hairpin tip, the consequence of which is currently unknown. Differing orientations of Tyr83 in the two subunits produces asymmetry in the added tip region (Fig. S14B,C). Fitting the new P3 domain into an all-atom chemotaxis

Fig. 4 The arrangement of chemotaxis proteins in *Td*. The linear organization of CheA produces 'strands' of CheA:CheW rings that are perpendicular to the cell axis and are linked by an atypical P3 domain.

array that was generated for previous molecular dynamics simulations (PDB ID: 3JA6)

shows that these additional helices are within a ~15 Å from receptors (Fig. S14D)[12].

Additionally, the handedness of the helix connection in the *Td* P3 domain differs from that of

Thermotoga maritima CheA and rather matches helix connectivity of sensor kinase DHP

domains[24,25].

Discussion

Here, we reveal the protein arrangement of F2 chemotaxis arrays through cryo-ET of intact

T. denticola (*Td*) cells. *Td* cells are thin and cyndrillical, producing a cell that is polarized in

shape and membrane curvature; the membrane has extremely high curvature perpendicular

to the cell axis and lower curvature parallel to the axis. In accordance with this feature, the

Td arrays are polarized and assume a preferred orientation with respect to the cell axis. In

this system, three proteins comprise the rings at the receptor tips: CheA, CheW, and a

CheW-CheR$_{like}$ protein. Like the *Ec* system, these proteins in *Td* are integrated into the array

with strict organization[1,22]. However, a linear arrangement of CheA is present that generates

'strands' of rings interlinked by the CheA dimerization domain (P3). The strands run

perpendicular to the axis of the *Td* cells, resulting in substantial bending of inter-linked rings.

Presumably, the unique binding interfaces in the CheA:CheW rings generate bent interfaces

with the appropriate curvature. Furthermore, this directional distinction allows for some array

contacts to follow the cell axis and remain undistorted. The deleterious effect on array

integrity with the loss of the CheR$_{like}$ domain strongly suggests that CheR$_{like}$ dimerization

plays a key role in array assembly and stabilization. The strict linear arrangement of CheA

could be facilitated by the composition of the *Td* rings; three unique protein interfaces are

present in the rings and restrict CheA P5 integration (i.e. CheA P5 can only occupy these

two positions in the six-member ring). Furthermore, the *Td* CheA P3 domain is clearly

discernible in the averaged tomograms, which has not been previously observed *in vivo*[5,9,22].

As the CheA:CheW rings have to undergo substantial bending to accommodate the

baseplate curvature, the elongated P3 domain may have evolved to stabilize CheA

dimerization by increasing the interface area and furthermore may encourage interactions

with neighboring receptors[12]. Due to the curvature, the receptor trimer-of-dimer modules are

expected to be further splayed, and the elongated P3 may compensate for the increased

distance between receptors and P3. As *Td* cells have the smallest average diameter (0.1 –

0.4 µm) of all bacteria with determined chemotaxis architectures thus far, and the novel

linear arrangement of CheA produces arrays that can better accommodate its unique

membrane curvature, we surmise that the array architecture in *Td* is an adaptation that

evolved to produce an extended chemoreceptor apparatus in a highly curved and

asymmetric membrane[4,5,9]. Importantly, bioinformatics analyses indicate that the unique

protein features seen in *Td* are exclusive to Spirochaetia F2 systems.

Unexpectedly, the placement of CheA in WT *Td* arrays could not be discerned (with the

exception of the P3 domain), but was clearly visible in two *Td* mutants (Δ2498,

Δ2498Δ2496). As the density corresponding to the P3 domain in the WT strain is clearly

discernible, the sparse density corresponding to all other CheA domains (P1, P2, P4, P5) is

not attributed to low CheA incorporation in these arrays. These results indicate that the

kinase is highly mobile or more disordered in the WT strain, but is more constrained when

ODP (TDE2498) is deleted, suggesting that ODP directly affects array structure. However,

densities in the three strains do not designate an obvious position for ODP, indicating that

ODP may not be an integral component of the array, but rather peripherally interacts with the

chemotaxis machinery or influences its architecture through other means, perhaps related to

its signaling properties.

In summary, we illustrate a novel chemotaxis arrangement that has evolved to compliment

the spirochetes' high membrane curvature and asymmetry. Therefore, it is likely that the

behavior and characteristics of chemoreceptor arrays in general can be influenced by

perturbing the shape of the cell membrane. Recent studies with *Ec* ultra-minicells (the

smallest mini-cells available to date) produce densities corresponding to portions of

chemoreceptors that have not been observed before[3]. However, it is possible that increased

membrane curvature limits movement of the transmembrane receptors, resulting in

increased receptor localization and resolution. Furthermore, the use of lipid-templating has

been extensively used to assemble arrays *in vitro* for cryo-ET and reconstitutes the

chemotaxis apparatus in a perfectly flat formation[2,12]. While cryo-ET experiments of *in vivo*

Ec arrays consistently shows the CheA P3 density to be too sparse to determine its position,

cryo-ET of *in vitro* lipid-templated *Ec* arrays clearly defines the P3 position. If the P3 domain

does engage receptors, this discrepancy may suggest that lipid-templating abrogates this

native behavior. Furthermore, it is unclear how the *Td* chemotaxis arrangements affects

cooperative behavior in the arrays as linear CheA:CheW strands are only connected through

receptor interactions. Investigations into this system may reveal alternative mechanisms for

cooperative behavior than what have been reported. Collectively, our results illustrate the

importance of investigating transmembrane systems *in situ*, and that examining systems in

non-model organisms can lead to new, unexpected advances for understanding the

remarkable signaling system of bacterial chemotaxis.

Acknowledgements

This work is part of the research programme National Roadmap for Large-Scale Research

Infrastructure 2017 – 2018 with project number 184.034.014, which is financed in part by the

Dutch Research Council (NWO). This work was funded by a grant from the National

Institutes of Health: R35GM122535 awarded to BRC, R01AI078958 and R01DE023080 to

CLI, and by the European Union under a Marie-Sklodowska-Curie COFUND LEaDing

fellowship to ARM. We thank the Netherlands Centre for Electron Nanoscopy (NeCEN) for

access to cryo-ET data collection facilities, and NE-CAT at the Advanced Photon Source for

access to X-ray crystallography data collection facilities. NE-CAT is supported by

NIH/NIGMS awards P30 GM124165 and S10 RR029205.

Author contributions. A.R.M., D.R.O, K.K., C.L., B.R.C, and A.B. designed research; A.R.M.,

D.R.O., and K.K. performed research; A.R.M., D.R.O, W.Y., K.K., and A.S. analyzed data; and

A.R.M., D.R.O, and A.B. wrote the paper with input from all authors.

Competing interest. The authors declare no competing interests.

Materials and Correspondence. Correspondence and materials requests should be addressed to A.

Briegel at a.briegel@biology.leidenuniv.nl.

1.   Yang, W., Alvarado, A., Glatter, T., Ringgaard, S. & Briegel, A. Baseplate variability of

Vibrio cholerae chemoreceptor arrays. *Proc. Natl. Acad. Sci.* **115**, 201811931 (2018).

2.   Cassidy, C. K. *et al.* Structure and dynamics of the E. coli chemotaxis core signaling

complex by cryo-electron tomography and molecular simulations. *Commun. Biol.* **3**,

1–10 (2020).

3.   Burt, A. *et al.* Complete structure of the core signalling unit of the E. coli

chemosensory array in an optimised minicell strain. *Nat. Commun.* 1–9 (2019).

doi:10.1038/s41467-020-14350-9

4.   Briegel, A. *et al.* Universal architecture of bacterial chemoreceptor arrays. *Proc. Natl.*

Acad. Sci. U. S. A. **106**, 17181–17186 (2009).

5.   Briegel, A. *et al.* Bacterial chemoreceptor arrays are hexagonally packed trimers of

receptor dimers networked by rings of kinase and coupling proteins. *Proc. Natl. Acad.*

Sci. **109**, 3766–3771 (2012).

6.   Hazelbauer, G. L., Falke, J. J. & Parkinson, J. S. Bacterial chemoreceptors: high-

performance signaling in networked arrays. *Trends Biochem. Sci.* **33**, 9–19 (2008).

7.   Pedetta, A., Parkinson, J. S. & Studdert, C. A. Signalling-Dependent Interactions

Between the Kinase-Coupling Protein CheW and Chemoreceptors in Living Cells. **93**,

1144–1155 (2015).

8.   Muok, A. R., Briegel, A. & Crane, B. R. Regulation of the chemotaxis histidine kinase

CheA: A structural perspective. *Biochim. Biophys. Acta - Biomembr.* **1862**, 183030

(2019).

9.   Briegel, A. *et al.* Structure of bacterial cytoplasmic chemoreceptor arrays and

implications for chemotactic signaling. *Elife* **2014**, 1–16 (2014).

10.   Bhatnagar, J. *et al.* Structure of the ternary complex formed by a chemotaxis receptor

signaling domain, the CheA histidine kinase, and the coupling protein CheW as

determined by pulsed dipolar ESR spectroscopy. *Biochemistry* **49**, 3824–41 (2010).

11.   Zhang, K. *et al.* Two CheW coupling proteins are essential in a chemosensory

pathway of Borrelia burgdorferi. *Mol. Microbiol.* **85**, 782–794 (2012).

12.   Cassidy, C. K. *et al.* CryoEM and computer simulations reveal a novel kinase

conformational switch in bacterial chemotaxis signaling. *Elife* **4**, 1–20 (2015).

13.   Muok, A. R. *et al.* A di-iron protein recruited as an Fe[II] and oxygen sensor for

bacterial chemotaxis functions by stabilizing an iron-peroxy species. *Proc. Natl. Acad.*

Sci. **116**, 14955–14960 (2019).

14.   Wuichet, K. & Zhulin, I. Origins and Diversification of a Complex Signal Transduction

System in Prokaryotes. *Sci Signal* **128**, 1–27 (2010).

15.   Gumerov, V. M., Ortega, D. R., Adebali, O., Ulrich, L. E. & Zhulin, I. B. MiST 3.0: an

updated microbial signal transduction database with an emphasis on chemosensory

systems. *Nucleic Acids Res.* **48**, D459–D464 (2020).

16.   Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996 (2018).

17.   García-Fontana, C. *et al.* High specificity in CheR methyltransferase function: CheR2

of pseudomonas putida is essential for chemotaxis, whereas CheR1 is involved in

biofilm formation. *J. Biol. Chem.* **288**, 18987–18999 (2013).

18.   Greene, S. R. & Stamm, L. V. Molecular characterization of a chemotaxis operon in

the oral spirochete, Treponema denticola. *Gene* **232**, 59–68 (1999).

19.   Yao, W., Shi, L. & Liang, D. C. Crystal structure of scaffolding protein CheW from

thermoanaerobacter tengcongensis. *Biochem. Biophys. Res. Commun.* **361**, 1027–1032 (2007).

20. Li, X. *et al.* The 3.2 Å resolution structure of a receptor:CheA:CheW signaling complex defines overlapping binding sites and key residue interactions within bacterial chemosensory arrays. *Biochemistry* **52**, 3852–3865 (2013).

21. Briegel, A. *et al.* The mobility of two kinase domains in the Escherichia coli chemoreceptor array varies with signaling state . **14**, 384–399 (2010).

22. Yang, W. *et al.* In situ conformational changes of the escherichia coli serine chemoreceptor in different signaling states. *MBio* **10**, 1–14 (2019).

23. Sze, C. W., Zhang, K., Kariu, T., Pal, U. & Li, C. Borrelia burgdorferi needs chemotaxis to establish infection in mammals and to accomplish its enzootic cycle. *Infect. Immun.* **80**, 2485–2492 (2012).

24. Greenswag, A., Muok, A., Li, X. & Crane, B. Conformational Transitions that Enable Histidine Kinase Autophosphorylation and Receptor Array Integration. *J. Mol. Biol.* **427**, 87–92 (2016).

25. Bilwes, A. M., Alex, L. A., Crane, B. R. & Simon, M. I. Structure of CheA, a signal-transducing histidine kinase. *Cell* **96**, 131–141 (1999).

 Supplementary Methods and Materials

 Data availability

 The cryo-ET sub-tomogram averages that support these findings are deposited in the Electron

 Microscopy Data Bank (EMDB) with accession codes EMD-10784, EMD-10785, EMD-10786, EMD-

 10787, EMD-10788, EMD-10790. The protein x-ray crystallography structure that supports these

 findings is deposited in the Protein Data Bank (PDB) with accession code 6Y1Y. These depositions

 will be publicly available with publication.

 Code availability

 All custom-made scripts are available to editors and referees upon request. These scripts will be

 publicly available with publication.

 Bacterial strains, culture conditions, and oligonucleotide primers

 Treponema denticola (*Td*) ATCC 35405 (wild-type) was used in this study. The *Td* deletion

 mutants, Δ2498 and Δ2498Δ2986, were generated in a previous study[2]. Cells were grown in

 tryptone-yeast extract-gelatin-volatile fatty acids-serum (TYGVS) medium at 37°C in an

 anaerobic chamber in presence of 85% nitrogen, 10% carbon dioxide, and 5% hydrogen[1].

 Td mutants were grown with an appropriate antibiotic for selective pressure as needed:

 erythromycin (50 μg/ml) and gentamycin (20 μg/ml). *Escherichia coli* 5α strain (New England

 Biolabs, Ipswich, MA) was used for DNA cloning. The *E. coli* strains were cultivated in

 lysogeny broth (LB) supplemented with appropriate concentrations of antibiotics. The

 oligonucleotide primers for PCR amplifications used in this study are listed in Table S3.

 These primers were synthesized by IDT (Integrated DNA Technologies, Coralville, IA).

 Construction of a CheR truncated mutant (ΔCheR$_{like}$)

 TDE1492::ermB (Fig. S15) was constructed to replace the CheR-like domain (8781-1,308 nt) in

 TDE1492 with a previously documented erythromycin B resistant cassette (*ermB*)[3]. The

 TDE1492::ermB vector was constructed by two-step PCR and DNA cloning. To construct this vector,

 the 5` end of *TDE1492* region and the downstream flanking region were PCR amplified with primers

P_1/P_2 and P_3/P_4, respectively, and then fused together with primers P_1/P_4, generating Fragment 1. The

Fragment 1 was cloned into the pMD19 T-vector (Takara Bio USA, Inc, Mountain View, CA). The

ermB cassette was PCR amplified with primers P_5/P_6, generating Fragment 2. The Fragment 2 was

cloned into the pGEM-T easy vector (Promega, Madison, WI). The Fragment 1 and 2 were digested

using *Not*I and ligated, generating the *TDE1492::ermB* plasmid. The primers used here are listed in

Table S3. To delete *TDE1492*, the plasmid of *TDE1492::ermB* was transformed into *Td* wild-type

competent cells via heat shock, as previously described[4]. Erythromycin-resistance colonies that

appeared on the plates were screened by PCR for the presence of *ermB* and absence of *TDE1492*

(781-1,308 nt) gene**.** The PCR results showed that the *TDE1492 (781-1,308 nt)* gene was replaced by

ermB cassette as expected (Fig. S15). One positive clone (*ΔTDE1492*) was selected for further study.

Bioinformatics software and resources

The datasets used in the bioinformatics analysis were built using data from Microbial Signal

Transduction Database v3 (MiST3) accessed February 2020[5] and the Genome Taxonomy Database

v89 (GTDB) [6]. We built custom scripts using TypeScript-3.7.5 and NodeJS-12.13. To make these

scripts, we also used packages publicly available at the node package manager repository (npm): we

used RegArch-1.0.1[7] to separate CheW-CheR$_{like}$ from other CheWs, gtdb-local-0.0.12

(https://npmjs.com/package/gtdb-local) to use the GTDB taxonomy, Phylogician-TS-0.10.1-4

(https://npmjs.com/package/phylogician-ts) to visualize and manipulate the phylogenetic trees,

BioSeq-TS-0.2.4 (https://npmjs.com/package/bioseq-ts) to handle protein sequences and MiST3-TS-

0.7.6 (https://npmjs.com/package/mist3-ts) to access MiST3 API. Multiple sequence alignments were

produced using L-INS-I algorithm from the MAFFT package. To reduce redundancy in sequence

datasets we used CD-HIT v4.6[8] with unaligned sequences and Jalview[9] with aligned sequences.

RAxML v8.2.10[10] was used to perform phylogenetic reconstructions, and low support branches in the

phylogenetic trees were collapsed with TreeCollapseCL4[11]. Sequence logos were built using Weblogo

3.7[12].

Bioinformatics scripts and pipelines

We collected all information on the proteins classified as CheA (96,434) and CheW (134,165). To

process this dataset we built several scripts and pipelines to produce the tables, figures and datasets

used in this analysis (Fig. S16). The scripts are found in Supplementary File 1.

Chemosensory profile of Spirochaetotas

The "spiro-pipeline" selects all genomes from Spirochaetota phylum using gtdb-local package to

access GTDB v89, then it filters only the genomes that are also present in MiST3 database. It collects

the information on MiST for each genome and appends the complete taxonomy information from

GTDB and signal transduction profiles. Finally, the pipeline builds the table with the information in

markdown (Dataset 2).

Chemosensory profile of genomes with at least one CheA-F2

The "chea-pipeline" starts from the raw dataset taken from MiST3 database with information on

96,434 CheA genes. Based on MiST3 classification it selects the genomes with at least one CheA-F2

sequences and fetches information about these genomes. At this step, it also checks with the list

generated by the "wr-pipeline" of the genomes containing CheW-CheR$_{like}$. It proceeds to append

chemosensory information for each genome and GTDB taxonomy. At this step the pipeline splits into

four pathways, where one parses the information to build the Dataset 1 and the other three build

FASTA formatted files with sequences from: CheA-F2, CheR-F2, CheW-CheR$_{like}$ and all CheWs

belonging to genomes with at least 1 CheA-F2. We noticed that MiST3 currently misclassifies some

CheR proteins, so we used RegArch to filter out false positives. We also used RegArch to separate

CheW and CheW-CheR$_{like}$ sequences as MiST3 classifies both as CheW (scaffold). The RegArch

definitions can be found in the script 'regArchDefinitions.ts' of the source code in Supplementary File

1.

In MiST3[12], 306 genomes contain at least 1 CheA-F2 (Dataset 1) with three exceptions: two of them

belong to Acidobacteriota phylum,and one from Planctomycetota, which suggest that the presence of

an F2 system in these genomes (outside of the Spirochaetota phylum) is the consequence of lateral

gene transfer. Of the 306 genomes with at least one CheA-F2, only the following lack the CheW-

CheR$_{like}$ protein: the three non-Spirochaetota genomes mentioned previously, 40 genomes of the

Brachyspirae class, and three genomes from the order Borreliales (Fig. S2). Interestingly, the

Borreliales genomes do appear to have the CheW-CheR$_{like}$ gene, except there is no gene product

associated with them in the MiST3 database.

Classification of CheW

CheW proteins are not classified in MiST3. In order to make comparisons between the sequences of

CheW domains in F2 systems, we must first select only CheW-F2. We first assign to the F2 class all

the canonical CheW found in genomes with a single CheA-F2. Contrary to the F2 systems where the

canonical CheW is not present in the chemosensory gene clusters, other classes do contain their

canonical CheW within the rest of the gene cluster. CheW found within 5 genes from a classified

CheA were assigned to the same class as CheA. To perform this classification, we selected the 598

full-length CheW sequences generated by the chea-pipeline. We used CD-HIT to remove 405

redundant sequences (-c 1). Next, we ran the "classify-w" pipeline on the remaining sequences (193).

The pipeline reads the identifiers of the sequences and fetches the chemotaxis profile from MiST3 for

each genome. It classifies CheWs as F2 classes if there is only 1 CheA of the class F2 in the profile.

Next, it fetches the gene neighborhood (5 genes up and downstream) of the remaining CheWs and

assigns a matching class if there is a classified CheA within these genes. We also aligned the

sequences (193) with the L-INS-I algorithm of the MAFFT package, produced a phylogeny with 1000

rapid bootstrap using RAxML (-f a -m PROTGAMMAIAUTO -N 1000) and collapsed the phylogeny

using TreeCollapse4 at 50% bootstrap. We mapped the CheW classification to the CheW tree in

Figure S2. We expanded the F2 classification to the 74 sequences within the branch with only CheW-

F2 sequences.

Comparison of the CheW domains in CheA, CheW-CheR$_{like}$ and CheW

We put together the sequences from CheA-F2 and CheW-CheR$_{like}$ (both trimmed by the PFAM model

for the CheW domain, already annotated in MiST3) and the full sequence of the CheW-F2 selected in

the previous step. We then use L-INS-I algorithm to align the sequences and Jalview to manually

inspect and eliminate identical redundant sequences. Finally, we trimmed the whole alignment based

on the boundaries of CheA-F2 and CheW-CheR$_{like}$ and eliminated one incomplete sequence:

GCF_000413015.1-HMPREF1221_RS07250. The final alignment had a total of 206 sequences: 73

CheA, 59 CheW-CheR$_{like}$ and 74 CheW. We separated the alignments into individual files and built

sequence logos to summarize the amino-acid diversity in each position for each group (Fig. S3B).

Comparison of CheR domains

557 First, we added together the trimmed part matching the CheR domain of the CheW-CheR$_{like}$

558 sequences and the 292 sequences of the CheR protein. We then aligned the 550 sequences using L-

559 INS-I. We used Jalview to inspect the alignment and remove identical redundant sequences. The final

560 alignment had the CheR domain of 83 CheR and 83 CheW-CheR$_{like}$ proteins. We separated the

561 alignments into individual files and generated independent sequence logos (Fig. S3A).

Analyses of CheA P3 domains

The "p3-pipeline" pipeline processes the data for the analysis of the length of P3 domains of all CheA

homologs in the MiST3 database. It reads the information for all 96,434 CheAs in the MiST3

database, trims the sequence matching the Pfam mode H-kinase_dim and builds FASTA formatted

datasets for each chemotaxis class. For each dataset, we used CD-HIT with 75% identity cutoff and

aligned them using the L-INS-I algorithm from MAFFT. Using Jalview we manually inspect and edited

the alignment to remove divergent sequences that opens major gaps in the alignment. We removed 6

F1 sequences, 37 F7 sequences, 20 F8 sequences and 5 F9 sequences. Then we merged the

alignment using mafft-profile with each dataset as a seed alignment in a single shot (Fig. S13A). We

selected the non-conserved central region of the alignment and measured the number of amino-acids

in each sequence (Fig. S13B).

Cryo-ET and sub-tomogram averaging of *T. denticola* chemotaxis arrays

Cells were concentrated by centrifugation, and a 1/10 dilution of protein A- treated 10-nm colloidal

gold solution (Cell Microscopy Core, Utrecht University, Utrecht, The Netherlands) was added to the

cells and mixed by pipeting. 3 μL aliquots of the cell suspension were applied to glow-discharged

R2/2 200 mesh copper Quanti-foil grids (Quantifoil Micro Tools, GmbH), the sample was pre-blotted

for 30 seconds, and then blotted for 2 seconds. Grids were pre-blotted and blotted at 20 °C and at

95% humidity. The grids were plunge-frozen in liquid ethane using an automated Leica EM GP

system (Leica Microsystems).

Data collection was achieved on a Titan Krios transmission electron microscope (Thermo Fisher

Scientific) operating at 300 kV. Images for three strains (WT, Δ2498, Δ2498Δ2986) were recorded

with a Gatan K2 Summit direct electron detector with a GIF Quantum energy filter (Gatan) operating

with a slit width of 20 eV. Images were taken at a magnification of 42,000×, which corresponds to a

pixel size of 3.513 Å. Tilt series were collected using SerialEM with a modified bidirectional tilt scheme

(-20° to 60°, followed by -22° to -60°) with a 2° increment. Images for the ΔCheR$_{like}$ strain were

recorded with a Gatan K3 Summit direct electron detector equipped with a GIF Quantum energy filter

(Gatan) operating with a slit width of 20 eV. Images were taken at a magnification of 26,000X, which

corresponds to a pixel size of 3.27 Å. Tilt series were collected using SerialEM with a bidirectional

dose-symmetric tilt scheme (-60° to 60°, starting from 0°) with a 2° increment. For all strains, the

defocus was set to − 6μm and the cumulative exposure per cell was 100 e-/A^2.

Bead tracking-based tilt series alignment and drift correcting were done using IMOD[13]. CTFplotter

was used for contrast transfer function determination and correction[14]. Tomograms were

reconstructed using simultaneous iterative reconstruction with iteration number set to 6. Dynamo was

used for manual particle picking and sub-tomogram averaging[15,16].

Purification of CheA, CheA P3, CheW-CheR$_{like}$, and TDE2496 proteins of *T. denticola*

DNA segments encoding the CheA P3 domain, CheW-CheR$_{like}$ protein, and TDE2496 in *T. denticola*

were PCR amplified from *Td* genomic DNA using a forward oligonucleotide encoding an NdeI

restriction site and a reverse primer encoding an EcoRI restriction site. The CheA protein was

amplified using a forward oligonucleotide encoding an NdeI restriction site and a reverse primer

encoding an BamHI restriction site. The PCR products were treated with the appropriate restriction

enzymes, purified, and ligated into a pet28a plasmid with a poly-Histidine tag and kanamycin

resistance marker. The plasmids were transformed into *Escherichia coli* BL21-DE3 cells and 4-8 L of

cell culture were grown at 37°C until an O.D. of 0.6 was reached. The flasks were cooled to 21°C and

1 mM of IPTG was added to the culture. The cells were harvested after 16 hours of growth. The cells

were lysed via sonication in lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 5 mM Imidazole) while

cooled on ice. The lysate was centrifuged at 20,000 X G for 1 hour at 4°C. The lysate was then run

over a gravity-flow purification column containing 3 ml of Nickel-NTA resin. The resin was washed

with 10 ml wash buffer (50 mM Tris pH 7.5, 150 mM NaCl, 20 mM Imidazole) and the protein was

eluted with 10 ml elution buffer (50 mM Tris pH 7.5, 150 mM NaCl, 200 mM Imidazole) and collected

in 1 ml fractions. The fractions were assessed for protein concentration via Bradford reagent and the

fractions containing protein were run on a size-exclusion s75 and s200 column systems that

monitored absorbance at 280 nm and collected 6 ml fractions. Fractions that contain CheA were

concentrated to ~20 mg/ml via centrifugation in a protein concentrator containing a regenerated

cellulose filter with a 50 kDa molecular-weight cut-off (MWCO). Fractions that contain CheA P3,

CheW-CheR$_{like}$ and TDE2496 were concentrated with a 10 kDa MWCO filter to 32 mg/ml, 11 mg/ml

and 7 mg/ml, respectively. The protein solutions were aliquoted, flash frozen in liquid nitrogen, and

stored at -80°C. For CheW-CheR$_{like}$, CheA, and CheA P3 domain, the purifications were prepared at

ambient temperatures. For TDE2496, the purification was prepared at 4°C.

Radioisotope assays

23 µl samples containing 2 µM CheA alone, or 2 µM CheA and 2 µM CheW-CheR$_{like}$, or 2 µM CheA

and 2 µM CheW-CheR$_{like}$ with TDE2496 (2 µM or 12 µM), were incubated in 50 mM MOPS pH 7.5,

150 mM KCl, 10 mM MgCl$_2$ for 30 minutes at ambient temperatures. Phosphorylation of CheA was

initiated by the addition of 2 µl of a solution containing 1 mM ATP mixed and 2-10 µl radiolabeled γ-

P$_{32}$ ATP (3000 Ci/mmol, 10 µCi/µL; Perkin Elmer) and quenched with 25 µl of 3X LDS buffer

containing 100 mM EDTA pH 8.0 after 1-12 minutes. The samples were run on a native Tris-glycine

gel for 2 hours at 120 volts. The gels were dried, placed in a radiocassette for 24 hours, and then

imaged with a Typhoon phosphor-imager (GE Healthcare). The intensity of the radio-labeled protein

bands were quantified using ImageJ.

Quantification of cell curvature

The cell curvature of *Td* whole cells and *V. cholera* minicells was quantified by analyzing images of

cross-sections of the cells where top views of chemotaxis arrays are present. For *Td* cells, the inner

membrane curvature and CheA:CheW base-plate curvature was quantified. For *Vc* minicells, the

inner membrane curvature was quantified. These images were pre-processed with Fiji by placing

points along the desired area with a distance of 10 nm between each point. The curvature of the inner

membrane was measured with a pre-built plugin for python, called Sabl_mpl, written by Jewett, A.

from the Jensen lab (Pasadena, CA)[23]. The "measure 3-point curvature" function was used to select

three adjacent points along the inner membrane of the cell and calculate the radius of these points.

The radius of the three selected points allowed for the calculation of the local curvature by dividing 1

with the measured radius (1/R = c). This was repeated for all points with a "sliding window" approach,

where the second point of the initial three points would become the first point, until the desired area

was covered.

Quantification of array alignment to the cell axis

The angle between the strands of CheA:CheW rings and the Td cell axis was quantified using Image

J software. 2D images from reconstructions that clearly locate the orientation of the strands and cell

axis were chosen for analysis. First, a straight line was drawn from the cell pole down the axis of the

cells. Then, a second line was drawn through one of the strands in the array and the angle between

the two intersecting lines was quantified. In some cases, the angle was too small (<3°) to be

accurately determine so the angle was annotated as 0°.

Residue conservation and molecular modeling of Td CheW, CheA P5, and the CheW domain of

CheW-CheR$_{like}$

The protein sequences of the two Td CheW domains and Td P5 were aligned (Clustal Omega),

conservation was calculated based on the alignment (JalView[9]), and the highest variable regions

were selected based on conservation (10+ adjacent residues with conservation scores lower than 8).

Homology models of Td CheA P5, CheW and the CheW domain of CheW-CheR$_{like}$ were generated

via the Swiss-Model server using complete residue sequences of each protein as a target[24]. The

CheW protein from *Thermoanaerobacter tengcongensis* (Tt) (PDB ID: 2QDL) had the highest percent

identity to the CheW proteins (37% and 31%) and was therefore used as the structural template. The

resulting homology models for Td CheW and the CheW domain of CheW-CheR$_{like}$ had a QMEAN of -

1.3 and -1.2, respectively. The P5 structure from *Escherichia coli* produced the best homology model

for Td CheA P5, with QMEAN -0.98 (PDB ID: 6S1K). The homology models were then aligned to the

CheW protein and P5 domain in a crystal structure containing *Thermotoga maritima* CheW and CheA

P4P5 in complex using PyMol (PDB ID: 3UR1).

Crystallization and structural determination of the P3 domain of *T. denticola* CheA

The isolated P3 domain was concentrated to 32 mg/ml and crystallized via hanging drop in 0.1 M

Imidazole pH 7.0, 25% PEG 400 using a 1:1 ratio of protein solution to crystallization solution with a

final volume of 3 µl. Crystals were apparent within eight hours but increased in size over three days.

Crystals were manually picked up in loops, flash cooled and shipped in liquid nitrogen to a beamline

(APS, line NE-CAT 24-ID-C, Dectris Pilatus 6M-F Pixel Array detector). The crystals diffracted to ~1.3

Å with a C2 symmetry and data was cut-off at 1.5 Å. The diffraction data was scaled and integrated

using XDS[17], and phased by molecular replacement with Phaser MR using *ab initio* search models

generated through the QUARK server and then ran through the AMPLE pipeline on the CCP4 web

server[18–20]. Model improvement was done by several rounds of manual model improvement in COOT

followed by automated refinement using Phenix Refine software[21,22].

1.   Ohta, K., Makinen, K. K. & Loesche, W. J. Purification and characterization of an enzyme

     produced by Treponema denticola capable of hydrolyzing synthetic trypsin substrates. *Infect.*

     Immun. **53**, 213–20 (1986).

2.   Muok, A. R. *et al.* A di-iron protein recruited as an Fe[II] and oxygen sensor for bacterial

     chemotaxis functions by stabilizing an iron-peroxy species. *Proc. Natl. Acad. Sci.* **116**, 14955–

     14960 (2019).

3.   Goetting-Minesky, M. P. & Fenno, J. C. A simplified erythromycin resistance cassette for

     Treponema denticola mutagenesis. *J. Microbiol. Methods* **83**, 66–68 (2010).

4.   Kurniyati, K. & Li, C. pyrF as a counterselectable marker for unmarked genetic manipulations

     in Treponema denticola. *Appl. Environ. Microbiol.* **82**, 1346–1352 (2016).

5.   Gumerov, V. M., Ortega, D. R., Adebali, O., Ulrich, L. E. & Zhulin, I. B. MiST 3.0: an updated

     microbial signal transduction database with an emphasis on chemosensory systems. *Nucleic*

     Acids Res. **48**, D459–D464 (2020).

6.   Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny

     substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996 (2018).

7.   Ortega, D. R. & Jensen, G. J. Regular Architecture (RegArch): A standard expression

     language for describing protein architectures. *bioRxiv* 679910 (2019). doi:10.1101/679910

8.   Li, W., Jaroszewski, L. & Godzik, A. Clustering of highly homologous sequences to reduce the

     size of large protein databases. *Bioinformatics* **17**, 282–283 (2004).

9.   Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview Version

2-A multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–

1191 (2009).

10.   Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

11.   Hodcroft, E. TreeCollapserCL4: Collapse Trees by Bootstrap. Available at:

http://emmahodcroft.com/TreeCollapseCL3.html.

12.   Crooks, G., Hon, G., Chandonia, J. & Brenner, S. WebLogo: A Sequence Logo Generator.

Genome Res **14**, 1188–1190 (2004).

13.   Mastronarde, D. N. Dual-axis tomography: An approach with alignment methods that preserve

resolution. *J. Struct. Biol.* **120**, 343–352 (1997).

14.   Xiong, Q., Morphew, M. K., Schwartz, C. L., Hoenger, A. H. & Mastronarde, D. N. CTF

determination and correction for low dose tomographic tilt series. *J. Struct. Biol.* **168**, 378–387

(2009).

15.   Castaño-Díez, D., Kudryashev, M., Arheit, M. & Stahlberg, H. Dynamo: A flexible, user-friendly

development tool for subtomogram averaging of cryo-EM data in high-performance computing

environments. *J. Struct. Biol.* **178**, 139–151 (2012).

16.   Castaño-Díez, D., Kudryashev, M. & Stahlberg, H. Dynamo Catalogue: Geometrical tools and

data management for particle picking in subtomogram averaging of cryo-electron tomograms.

J. Struct. Biol. **197**, 135–144 (2017).

17.   Kabsch, W. *et al.* XDS. Acta Crystallogr. Sect. D Biol. Crystallogr. **66**, 125–132 (2010).

18.   Bibby, J., Keegan, R. M., Mayans, O., Winn, M. D. & Rigden, D. J. AMPLE: A cluster-and-

truncate approach to solve the crystal structures of small proteins using rapidly computed ab

initio models. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **68**, 1622–1631 (2012).

19.   Keegan, R. M. *et al.* Exploring the speed and performance of molecular replacement with

AMPLE using QUARK ab initio protein models. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **71**,

338–343 (2015).

20.   Thomas, J. M. H. *et al.* Routine phasing of coiled-coil protein crystal structures with AMPLE.

IUCrJ **2**, 198–206 (2015).

21.   Adams, P. *et al.* The Phenix Software for Automated Determination of Macromolecular

Structures. *Methods* **292**, 342–351 (2009).

22.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*

Crystallogr. Sect. D Biol. Crystallogr. **66**, 486–501 (2010).

23.   Yao, Q. *et al.* Short FtsZ filaments can drive asymmetric cell envelope constriction at the onset

of bacterial cytokinesis. *EMBO J.* **36**, 1577–1589 (2017).

24.   Schwede, T., Kopp, J., Guex, N. & Peitsch, M. C. SWISS-MODEL: An automated protein

homology-modeling server. *Nucleic Acids Res.* **31**, 3381–3385 (2003).

741
742
743
744

Fig. S1 The arrangement of the canonical chemoreceptor array, exemplified in *Escherichia coli* (*Ec*). (A) Transmembrane chemoreceptors form dimers that function as a trimer-of-dimers module to modulate the activity of the histidine kinase CheA (blue) via a coupling protein CheW (green). In cryo-ET experiments, cross-sections at Layer 1 reveals the arrangement of chemoreceptors and Layer 2 reveals the position of the kinase. (B) In all organisms investigated previously, CheA is integrated into the array with hexagonal (pseudo-P6) symmetry.

Fig. S2 The chemosensory class profile of Spirochaetota. Sequences belonging to the four major classes of Spirochaetota are marked: UBA4802 (dark red), Leptospira (purple), Brachyspirae (cyan), UBA6919 (blue), Brevinematia (yellow), GWE2-31-10 (green) and Spirochaetia (light green). Nodes are marked for presence of classes F1 (orange), F2 (red), F5 (blue), F7 (light green), F8 (green), ACF (light blue), and Tfp (purple). Genomes not present in MiST3 database are marked as grey. F2 systems with CheW-CheR$_{like}$ are marked with a black outline around the red circle.

A

CheR-R

CheWR-R

B

CheW

CheA-P5

CheWR-W

Fig. S3 (A) Sequence logo of representative sequences of the two groups of CheR-containing proteins in the F2 system. (B) Sequence logo of representative sequences of the three groups of CheW-containing proteins in the F2 system. Blue boxes denote the location of variable regions identified in the *Td* CheW and P5 homologs. In these locations, all regions possess unique conserved residues with the exception of Region 2, which is not at the CheA:CheW ring interfaces.

Fig. S4 Phylogenetic tree of CheW sequences in genomes with at least one CheA-F2 suggests a last common ancestor of CheW-F2 sequences. We mapped the sequences of CheW from the classes F2 (red), F5(purple), F7 (light green), F8 (green) and ACF (blue). The larger red internal node marks a candidate of the last common ancestor of CheW-F2.

Fig. S5 (A) The number of arrays and particles selected to generate sub-tomogram averages for each *Td* strain. (B) Sub-tomogram averages reveal the orientation of the cell axis with respect to the orientation of the chemotaxis arrays. Intriguingly, the arrays have a preferred orientation in the cells and this orientation is conserved in WT and deletion mutants.

763
764

765
Fig. S6 (A) The soluble receptor TDE2496 deactivates *Td* CheA in a concentration-dependent manner. (B) MALS
experiments demonstrate that TDE2496 forms the expected dimer (52 kDa) and hexamer (156 kDa) oligomeric states.
(C) The CheW-CheR$_{like}$ protein is conserved on the same operon as the only CheA, CheX, and CheY proteins in *Td*. (D)
The CheW-CheR$_{like}$ linker is predicted to forma single alpha helix flanked by unordered regions (Jpred). (E) Radioisotope
assays that monitor CheA autophosphorylation over time in the presence and absence of CheW-CheR$_{like}$ demonstrate a
difference in CheA activity. (F) In the *T. maritima* system, the presence of CheW also activates CheA kinase activity.
     Both samples are at time point 3 minutes.

Fig. S7 The arrangement of chemotaxis CheA:CheW rings in *Td.* Top: The CheA:CheW rings consists of three domains: CheA P5 (blue), a classical CheW protein (green), and the CheW domain of CheW-CheR$_{like}$ (white). Three unique interfaces are produced and the CheR$_{like}$ domain (purple) extends to the center of the rings to interact with one another. Bottom: The location of the CheW-CheR$_{like}$ protein in the rings is evident by examining the location of the CheA domains with respect to the CheW-CheR$_{like}$ linker.

777
778
779
780

Fig. S8 Homology models of *Td* CheW domains and CheA P5. (A) A homology model of the CheW domain of CheW-CheRlike using *Thermoanaerobacter tengcongensis* (*Tt*) CheW (PDB ID: 2QDL) as the template. (B) A homology model of the classical *Td* CheW using *Tt* CheW (PDB ID: 2QDL) as the template. (C) A homology model of CheA P5 using *E. coli* CheA P5 (PDB ID: 6S1K) as the template.

Fig. S9 Homology models of *Td* CheA P5, CheW, and the CheW domain of CheW-CheR$_{like}$ mapped onto a previously determined crystal structure of *Thermotoga maritima* CheA P5 and CheW (PDB ID: 3UR1). Variable regions between the three domains (blue boxes) were determined by sequence alignments of the three domains followed by conservation analysis. These variable regions are located at the rings interface regions. The CheW domains also have variable N-terminal and C-terminal regions that are not complimented in P5 (grey boxes). Variable regions for P5 and the CheW domains are denoted on the homology models by italicized numbers and underlined numbers, respectively.

Fig. S10 (A) Sub-tomogram averaging of *Td* WT and mutant strains reveal that linked CheA:CheW rings (yellow) run perpendicular to the cell axis via a strict linear orientation of CheA (blue). Averages from WT *Td* are illustrated here but apply to all strains. (B) The curvature of the inner membrane of *Td* at chemotaxis arrays in the reconstructions is 35.8 +/- 6.6 /µm (270 Å radius). The curvature of the CheA:CheW baseplate (red arrow) is 65.6 +/- 19 /µm (152 Å radius). (C) *V. cholerae* minicells have an inner membrane curvature of 9.15 +/- 4.5 /µm (radius 1092 Å).

Fig. S11 Modeling of CheA:CheW rings to the curvature of the *Td* baseplate. (A) For single CheA:CheW rings to follow the baseplate curvature, it must buckle by an average of 7.6 Å toward the membrane. (B) In order for two linked CheA:CheW rings to run perpendicular to the cell axis, the center of the rings (P3) must buckle by 49.2 Å toward the cell membrane.

A
```
T.denticola   LRVDANRIDYLLNLVSETVITKASLNQSTIEFAELYDKFQNSSTIYKDKTRRLLDKMPEY  60
V.cholerae    LKVDQEKVDLLMDLVGELIVAKNSMQYLAYRAENEFGVR----------------------  39
C.jejuni      IRVEVKRLDHLMNLIGELVLGKNRLLKIYDDVEERYEGE----------------------  39
E.coli        IRVAVEKVDQLINLVGELVITQSMLAQRSSELDPVN-------------------------  36
S.enterica    IRVAVEKVDQLINLVGELVITQSMLAQRSNELDPVN-------------------------  36
T.maritima    VRVDIEKLGNLMDLMGELVIARSRILETLKK---YN-------------------------  33
B.subtilis    IRVNIDRLDSLMNLFEELVIDRGRLEQIAKE---LE-------------------------  33
              ::*   .::.  *::*.  *  ::  :      :

T.denticola   LEKIQQGYDINSIKQDVLNEYSSLLEVFGDFDSLMKAAVTKFKSSSQNLGRISGELQEGV 120
V.cholerae    ------------------------------KLAQDIKAEQSVISRLAEDLQSVV        63
C.jejuni      ------------------------------KFLEELNQVVSQLSIITTDVQLAV        63
E.coli        -----------------------------HGDLITSMGQLQRNARDLQESV         58
S.enterica    -----------------------------HGDLITSMGQLQRNARDLQESV         58
T.maritima    -----------------------------IKELDESLSHLSRITLDLQNVV         55
B.subtilis    -----------------------------HNELTETVERMTRISGDLQSII         55
                                           .:    :   : ::*  :

T.denticola   MKIRMVP        127
V.cholerae    MQVRMVP        70
C.jejuni      MKTRMQP        70
E.coli        MSIRMMP        65
S.enterica    MSIRMMP        65
T.maritima    MKIRMVP        62
B.subtilis    LNMRMVP        62
              :. ** *
```

Fig. S12 Multiple sequence alignments of the CheA P3 domain from several bacteria demonstrates the presence of additional P3 residues in *Td* and other Spirochetes. (A) CheA P3 alignments of *Td* and P3 from other bacteria with previously characterized chemotaxis proteins. *Td* possesses ~50 residues that are not found in the other homologs and are located in between the traditional dimerization helices. (B) *Td* CheA P3 aligned with other Spirochete P3. The additional residues identified in alignment **A** are highlighted in blue. Figures were made using Clustal Omega.

B
```
B.burgdorferi NLIRIDSKKIDYILNLVSEAVISKSSYNQINSEMITLFYNFNYFYDYQESFQRNFLIDLK  60
B.pilosicoli  SYLRVESDRIDAMMNQVGELVTNKSSYVQYDDDLSSYGKLISNSI----------NDIK  49
T.denticola   --LRVDANRIDYLLNLVSETVITKASLNQSTIEFAELYDKFQNSS---------------  43
T.pallidum    SVLRVDAKRIDYLLNLVSETVIIKASLNQSALEFGEVYTLFQNAN--------------- 45

B.burgdorferi IVFKDAGLTLEDEIESH------------INSLMSFKMEKALKDISELRNSFFRLLQNFK 108
B.pilosicoli  KYYRDTIVQILRKFEDY----AQKKEIKDLRANYINSFNSKLNEISKIEDDFKYTLDRYR 105
T.denticola   TIIYKDKTRRLLDKMPEYLEKIQQGYDINSIKQDVLNEYSSLLEVFGDFDSLMKAAVTKFK 103
T.pallidum    GAYKERLRKFFDRVPAYLEKVKNGQDADAVRKEMIAEAVGVFDIFSSFENGLKQSVTKFR 105
              :::    : ::         :.       :.  :.:  :   :      :  :.:

B.burgdorferi MTSGRLSRIITDLHESVLKTRMLP        132
B.pilosicoli  SSYQLLTRVTNELQETVMKIRMVP        129
T.denticola   SSSQNLGRISGELQEGVMKIRMVP        127
T.pallidum    SSAQNLGRISGELQEGVMKIRMVP        129
              :   * *:  :*:* *:* **:*
```

817
818
819

Fig. S13 Analysis of non-redundant CheA P3 domains with a 75% sequence identity cut-off (1450 sequences). (A)
Conservation scores of the P3 alignment with the 1450 sequences. The traditional P3 helices are largely conserved, but
regions located between the helices are non-conserved. (B) Analyses of CheA from different chemotaxis classes
reveals that CheA F2 homologs possess the most residues in the non-conserved region of P3. (C) The non-conserved
regions are llustrated with the two known CheA P3 structures. Top: *Td* P3 possesses 71 residues that align to the non-
conserved region (PDB ID: 6Y1Y). Bottom: *Tm* P3 possesses seven residues that align to the non-conserved region
(PDB ID: 1B3Q).

832
833
834
Fig. S14 The P3 domain of *Td* CheA. (A) The CheA P3 dimer contains a cluster of Phe and Tyr residues
near the breakages in the helices (black arrows). All Phe and Tyr residues are highlighted in red. (B) The
crystal structure of CheA P3 demonstrates asymmetry in the subunits. (C) Repositioning of Y82 in the
subunits induces alterations of adjacent residues and may account for subunit asymmetry. (D) Alignment
of the P3 structure to a previously determined model of the chemotaxis array in *E. coli* (PDB ID: 3JA6)
indicates that the P3 domain lies within ~15 Å of the receptors (when measuring from peptide back-bone).
         CheA and CheW in this model are shown in grey.

846
847
848
849   **Fig. S15** Diagrams illustrating construction of the *TDE1492::ermB* vector (A) for the targeted
850   mutagenesis of *TDE1492 (781-1,308 nt)* by in-frame replacement of *TDE1492* using *ermB* cassette.
      These constructs were constructed by two-step PCR followed by DNA cloning. Arrows represent the
851   relative positions and orientations of these primers, which are listed in Table S3. *ermB* =
852   erythromycin resistance. (B) Characterization of the *ΔTDE1492* strain by PCR analysis. The top
853   panel illustrates how the PCR analysis is designed; the bottom panel is the PCR results. Arrows
854   represent the relative positions and orientations of these primers; the numbers are predicted sizes of
      PCR products generated by the corresponding primers. The primer P_7 is located at the 5'-end of
855   *TDE1492*, P_6 at the 3'-end of *ermB*, P_5 at the 5'-end of *ermB*, P_8 at the flanking region of *TDE1492*,
856   P_9 at the middle of *TDE1492*, and P_{10} at the 3'- end of *TDE1492*. The sequences of these primers
857   are listed in Table S3.

Fig. S16 Flowchart of the three major pipelines used to produce the bioinformatics datasets. Steps marked in red represents fetching information from MiST3 database, in green are steps requiring RegArch as a filter, and in blue indicate writing data to file.

Td strain	Average angle (°)
WT (n=9)	9.67 +/- 9.21
Δ2498 (n=9)	9.55 +/- 8.75
Δ2498Δ2496 (n=8)	12.18 +/- 7.62
Combined average:	**10.40 +/- 8.59**

Table S1A. Statistics for the average angle between the *Td* cell axis and the 'strands' of CheA:CheW rings.

Vc cell number	Inner membrane curvature (/μm)
1	6.2
2	15.4
3	14.2
4	7.4
5	4.3
6	7.5
Average:	**9.155 +/- 4.53**

Table S1B. Statistics of inner membrane curvature for 6 *Vc* mini-cells.

Td cell number	Inner membrane curvature (/µm)	Baseplate curvature (/µm)
1	35.0	49.7
2	25.4	54.0
3	34.2	63.7
4	43.1	96.1
5	30.8	99.7
6	41.1	55.7
7	39.6	74.6
8	33.1	47.7
9	46.2	47.8
10	29.1	67.1
Average:	**35.8 +/- 6.6**	**65.6 +/- 19**

872 **Table S1C.** Statistics of inner membrane and baseplate curvature for 10 WT Td
873 cells.

	Td P3 domain (PDB: 6Y1Y)
Wavelength	0.979100
Resolution range	75.36 – 1.50 (1.55 – 1.50)
Space group	C 1 2 1
Unit cell	a, b, c: 101.64, 28.77, 82.90 α, β, γ: 90, 114.75, 90
Total reflections	53542
Unique reflections	35178 (3490)
Multiplicity	6.9 (6.2)
Completeness (%)	96.72 (99.31)
Mean I/sigma(I)	14.7 (5.4)
R-merge	0.063(0.290)
R-meas	0.075(0.350)
R-pim	0.040(0.193)
CC1/2	0.99(0.94)
Reflections for refinement	33864 (3361)
Reflections for R-free	1308 (143)
R-work	0.187 (0.205)
R-free	0.211 (0.244)
No. non-H atoms	2344
macromolecules	2028
ligands	0
solvent	316
Protein residues	253
RMS(bonds)	0.006
RMS(angles)	0.93
Ramachandran favored (%)	100
Ramachandran allowed (%)	0.00
Ramachandran outliers (%)	0.00
Rotamer outliers (%)	0.00
Clashscore	6.87
Average B-factor	19.1
macromolecules	17.2
solvent	28.2

905
906 **Table S2.** Data collection and refinement statistics for *Td* CheA P3 domain.
907
908
909
910

Primers	Sequences (5'-3')	Note [a]
P$_1$	CGGGCGTAGGCATCGGAGATAC	5` portion for TDE1492 inactivation; [F]
P$_2$	TTACAATATCACCTGAGCTGGCGGCCGCTTAAGGTGACAAGAAAGATGATA	5` portion for TDE1492 inactivation; [R]
P$_3$	TATCATCTTTCTTGTCACCTTAAGCGGCCGCCAGCTCAGGTGATATTGTAA	3` portion for TDE1492 inactivation; [F]
P$_4$	CCCAGAGCACTTATCATAAC	3` portion for TDE1492 inactivation; [R]
P$_5$	ATGAACAAAAATATAAAATATTCTC	Erythromycin B cassette (ermB); [F]
P$_6$	TTATTTCCTCCCGTTAAATAATAG	Erythromycin B cassette (ermB); [R]
P$_7$	ATGGAAGAAATGAAAGAAC	5` flanking region of TDE1492, ΔTDE1492 PCR analysis; [F]
P$_8$	GATATAGTTCTTGGCTCCAAG	3` flanking region of TDE1492, ΔTDE1492 PCR analysis; [R]
P$_9$	GAGGCCTATATAAATGCC	TDE1492, ΔTDE1492 PCR analysis; [F]
P$_{10}$	CTTGGGCATTGCCTCATTATG	TDE1492, ΔTDE1492 PCR analysis; [R]

Table S3. Oligonucleotide primers used in this study
[a] Underlined sequences are engineered restriction cut sites for DNA cloning; [F] forward; [R] reverse.

911
912
913
914
915