Revealing the polarity of actin filaments in intact cells

Bruno Martins[1], Simona Sorrentino[1], Wen-Lu Chung[1], Meltem Tatli[1], Ohad Medalia[1*], and

Matthias Eibauer[1*]

[1] Department of Biochemistry, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

Keywords: Actin filaments, Focal adhesions, Cryo electron tomography, Subtomogram

averaging, Image processing

*Correspondence should be addressed to O.M. (omedalia@bioc.uzh.ch) and to M.E.

(m.eibauer@bioc.uzh.ch)

Abstract

The actin cytoskeleton plays a fundamental role in numerous cellular processes, such as cell

motility, cytokinesis, and adhesion to the extracellular matrix. Revealing the polarity of

individual actin filaments in intact cells, would foster an unprecedented understanding of

cytoskeletal processes and their associated mechanical forces. Cryo-electron tomography

provides the means for high-resolution structural imaging of cells. However, the low signal-to-

noise ratio of cryo-tomograms obscures the high frequencies and therefore the polarity of actin

filaments cannot be directly measured. Here, we developed the ActinPolarityToolbox that

enables to determine the polarity of each actin filament in cellular cryo-tomograms. We applied

this method to reveal the actin polarity distribution in focal adhesions, and show a linear relation

between actin polarity and distance from the apical boundary of the adhesion site. Our method

facilitates the reconstruction of the actin cytoskeleton at sub-nanometric precision.

Introduction

Actin polymerization drives cell motility and is a central factor in mediating contractile forces in

cells [1-3]. Actin filaments [4, 5] assemble into complex networks [6, 7], which are essential for

their activity in the cell. Reconstructing individual actin filaments at sub-nanometer resolution

inside cells, would provide an unparalleled view on actin networks, and would allow a more

fine-grained modelling of cytoskeletal based mechanical processes [8].

A prominent mechanosensitive process, involving actomyosin contractility, occurs at the

integrin-based interaction sites of cells with the extracellular matrix [9, 10]. These interactions

are mediated by adhesive structures such as focal adhesions (FAs) [11-13]. The organization

of proteins and the role of the actin network in FAs has been intensively studied using

fluorescent and electron microscopy [14, 15], however, the high-resolution 3D architecture of

FAs, including the polarity and position of each individual actin filament at sub-nanometer

precision was not yet shown.

Cryo-electron tomography (cryo-ET) allows to reconstruct 3D density maps of unperturbed

cells at a resolution of 2-3 nm [16-18]. Therefore, single actin filaments can readily be detected

in tomograms of eukaryotic cells [19, 20]. Moreover, using direct electron detectors permit to

recognize the characteristic helical shape of F-actin that is in clear distinction to other cellular

filaments. However, cryo-tomograms suffer from a low signal-to-noise ratio (SNR) [21, 22], and

from missing information due to the limited tilt range during data acquisition, referred to as the

missing wedge [16]. Subtomogram averaging can compensate for these issues [17, 23-25],

but it is computationally more demanding and is lacking the robustness and level of

standardization as the procedures used for data analysis in single particle cryo-

electron microscopy [26, 27].

Here we present the ActinPolarityToolbox (APT), comprising a set of MATLAB scripts that

enable the 3D reconstruction of actin filaments from cryo-tomograms at sub-nanometric

resolution, which we based on transforming subtomogram averaging into a single particle task,

and the determination of their polarity build on a robust statistical analysis of the individual

filaments. Furthermore, APT features tools for spatial and topological analysis of the

reconstructed actin filament networks and their visualization.

Using a correlative fluorescence microscopy and cryo-ET approach [15, 28], we applied APT

to describe the 3D architecture of the actin cytoskeleton at FAs. We show that the actin polarity

distribution correlates with the position along the FA and that regions of mixed polarity are

concentrated at the periphery of the characteristic actin bundles.

Results

Actin polarity determination with APT. The first aim of APT is the reconstruction of an actin

filament with clear polarity features. Therefore, APT requires cryo-tomograms and

segmentations of the actin cytoskeleton as input data (Fig. 1a). Subsequently, the segmented

filaments are subdivided into segments with equidistant spacing. We make use of the

observation that a projection of a subtomogram is invariant of the missing wedge orientation

and can be treated as a single particle image (Supplementary Fig. 1). Thus, fast, stable, and

standardized image processing packages, such as cryoSPARC [29] or RELION [27], can be

used to reconstruct the actin filament from the segments. Interestingly, this is a general

approach, and not only restricted to filamentous structures (Supplementary Fig. 2).

The main goal of APT is to determine the polarity of all the initially segmented filaments.

Therefore, based on the filament average we calculate for each segment the position of its

plus-end in the cryo-tomograms. During the 3D reconstruction step, APT obscures the

correlation between segments and filaments, thus for each filament multiple independent

polarity observations can be statistically evaluated, which greatly reduces the error of polarity

determination, as we show in the validation of the method (see below).

The segmentations can be conducted on contrast enhanced tomograms in order to facilitate a

better detection of actin filaments. We do not assume the segmentations to be free of false

positives, because they will be efficiently sorted out at a later stage. However, we assume that

each filament or detected part is represented by a unique set of voxels, therefore the filaments

should not touch each other in the segmentations. The fact that a filament appears divided into

parts, for instance when it runs through low-contrast, dense or crowded regions, where it

cannot be tracked unambiguously, has no detrimental effect on its polarity determination, but

leads to an underestimation of the actual filament length.

The workflow of APT is visualized in Fig. 1a. A detailed description of the method can be found

in the following subsections.

Step I: Coordinates. The input data is a set of N_T cryo-electron tomograms T_i $(i = 1, \cdots, N_T)$

and for each T_i there is an associated segmentation S_i of the actin filaments. Each S_i define a

set of filaments F_i^j $(j = 1, \cdots, N_i)$, with N_i being the number of detected filaments in T_i. Firstly,

APT utilizes the S_i to construct for each F_i^j a set of 3D coordinates $\left\{x_1, x_2, \cdots, x_{N_i^j}\right\}$, that are evenly

spaced along F_i^j, and will be the sampling points for polarity determination. Here N_i^j indicates

the number of sampling points per filament, which we term segments.

Step II: Subtomograms. Next, APT pools all these $N_C = \sum N_i^j$ segments x_k $(k = 1, \cdots, N_C)$,

stores the index relation $k \leftrightarrow F_i^j$ for polarity determination at a later stage, and extracts N_C

subtomograms centred around x_k from a ctf-corrected version of the dataset T_i.

Step III: Projections. Subsequently, APT masks and then projects the subtomograms in

z-direction. The applied mask diminishes the influence of neighbouring filaments in z-direction.

We term the height of the mask as projection thickness parameter.

Step IV: Prealignment. The purpose of this module is to align the (central) filament density in

the projected subtomograms parallel to the x-axis (Methods). The detected orientations will be

used as priors in the next step.

Step V: 2D Classification. In this module high quality class averages of filamentous actin will

be produced (Methods). It allows to sort out false positive and low-quality segments [30].

Because the filaments are prealigned at this stage, a second mask can be applied in order to

further diminish the influence of neighbouring filaments in y-direction.

Step VI: 3D Reconstruction. In order to obtain an actin filament structure from the projected

subtomograms we use a helical reconstruction 3D refine job in RELION [31] (Methods).

Subsequently, APT combines all transformations, and as a result we have the forward

transformation Ω_k, that describes how filament F_i^j has to be transformed at sampling point x_k

in order to align with the filament average (Supplementary Fig. 3a).

Step VII: Polarity. To resolve actin polarity, it is crucial that the filament average reaches a

resolution better than 20 Å. Once this has been achieved, APT calculates the inverse

transformation Ω_k^{-1}, that describes how filament F_i^j is oriented at sampling point x_k with respect

to the filament average (Supplementary Fig. 3a). Particularly filament polarity is encoded in the

inverted psi angle ψ_k^{-1}, which indicates the position of the plus-end of F_i^j relative to the

y-axis of T_i.

Next, APT uses the index relation $k \leftrightarrow F_i^j$ to extract all inverted psi angles $\psi_k^{-1}(T_i)$, that are

originating from the same T_i. If T_i contains a filament bundle with mixed polarity, then the

histogram of $\psi_k^{-1}(T_i)$ shows two distinct peaks, that are separated by 180°. The height of each

peak indicates how many segments were aligned with their plus-ends in the respective

direction, and allows an estimation of the polarity ratio within the bundle, while the peak width

decreases with increasing parallelism of the filaments. Obviously, if a bundle exhibits a uniform

polarity distribution, then the histogram of $\psi_k^{-1}(T_i)$ shows a single peak.

Subsequently, APT assigns to each segment a direction label. Per default, segments included

in the right peak are labelled with 0, segments included in the left peak are labelled with 1.

Therefore, each filament can be represented as a vector $\xi(F_i^j)$ of labelled segments.

The centrepiece of APT is that the elements of $\xi(F_i^j)$ are computed statistically independent.

Segments, whose directionality was detected incorrectly, do not prohibit the correct polarity

determination of a filament as a whole, as long as the majority of segments building that

filament were determined accurately. Consequently, the final polarity label $P(F_i^j)$ of filament F_i^j

is defined as follows:

146
$$P(F_i^j) = \begin{cases} 0 & if\ \mathrm{mean}\left(\xi(F_i^j)\right) < 0.5 \\ 0.5 & if\ \mathrm{mean}\left(\xi(F_i^j)\right) = 0.5 \\ 1 & if\ \mathrm{mean}\left(\xi(F_i^j)\right) > 0.5 \end{cases}.$$

147   If $P(F_i^j) = 0.5$ the polarity of that filament is maximally uncertain.

149   *Step VIII: Confidence.* We suggest that the closer $\mathrm{mean}\left(\xi(F_i^j)\right)$ is to 0 or to 1, respectively,

150   the higher the probability, that $P(F_i^j)$ was determined correctly. We use this relation to define

151   a confidence score for each filament, termed majority confidence score (mcs), as follows:

152
$$\mathrm{mcs}(F_i^j) = \max(\mathrm{mean}\left(\xi(F_i^j)\right), \mathrm{mean}\left(\sim\xi(F_i^j)\right)),$$

with $\sim\xi(F_i^j)$ the logical complement of $\xi(F_i^j)$. The values of mcs range between 0.5 (maximum

uncertainty) to 1 (minimal uncertainty), and are the fraction of segments per filament that are

pointing in the majority direction. The mcs measures the consistency of the segment's

directionality determination along a filament.

Additionally, we define a second confidence score, termed sensitivity confidence score (scs).

Here the reconstruction of the filament average is repeated with a reversed direction of the

template. Subsequently, APT initializes for each F_i^j a vector $\lambda(F_i^j)$, having the same number

of elements than F_i^j segments. Those segments that reverse their plus-end orientation as well,

are orientational sensitive and APT sets the respective elements of $\lambda(F_i^j)$ to 1. On the other

hand, if a segment shows no logical behaviour upon change in template direction, the

respective element of $\lambda(F_i^j)$ is set to 0. Then we define scs as follows:

164
$$\text{scs}(F_i^j) = \text{mean}\left(\lambda(F_i^j)\right).$$

165 The values of scs range between 0 (no segment of F_i^j is orientational sensitive) and 1 (F_i^j is

166 built exclusively from orientational sensitive segments). Comparable to the mcs, the scs is an

167 alternative to measure the self-consistency of the segment's directionality determination.

168 Finally, we define the combined confidence score (ccs) of filament F_i^j as follows:

169
$$\text{ccs}(F_i^j) = \text{mcs}(F_i^j) \cdot \text{scs}(F_i^j).$$

Step IX: Topology. In this module APT performs a 3D neighbourhood analysis of the actin

filaments in order to characterize their polarity distribution in the observed bundles. Therefore,

centred around each segment x_k, APT constructs a sphere $\mathcal{K}_\epsilon(x_k)$ with radius ϵ, defining the

neighbourhood of x_k. Then, APT extracts number, distance and polarity of all filaments passing

through $\mathcal{K}_\epsilon(x_k)$. For instance, let the polarity of the segment x_k be $P(F_i^j) = 0$, and imagine F_i^j

is surrounded by four filaments within $\mathcal{K}_\epsilon(x_k)$, of which one displays same polarity and three

reversed polarity (Supplementary Fig. 3b). Then, the position of the probed filament is located

in a neighbourhood that exhibits 75% mixed polarity and 25% uniform polarity, respectively.

This calculation is performed by APT for all $\mathcal{K}_\epsilon(x_k)$, and subsequently it sums the degrees of

uniform polarity $P_{\parallel}$ and mixed polarity $P_{\nparallel}$ of all neighbourhoods in a given actin bundle. Finally,

the topology score τ of an actin bundle is defined as follows:

182
$$\tau = \frac{P_{\parallel} - P_{\nparallel}}{P_{\parallel} + P_{\nparallel}}.$$

The topology score τ ranges from -1 to +1. The more the filaments are organized in a uniform

polarity configuration, the value of τ will be closer to +1. If the bundle shows a phase separated

topology, so that filaments with opposite polarities adjoin at phase boundaries mainly, τ will be

reduced accordingly. However, the more the bundle favours close proximity between filaments

with opposite polarities, the more τ will approach -1.

Validation of APT. The missing wedge affects the reconstruction of a tomogram in an

anisotropic manner [16]. As a consequence, filaments running parallel to the tilt-axis are better

resolved than filaments oriented orthogonal to the tilt-axis (Supplementary Fig. 1). It is

fundamental to ensure that our projection approach for subtomogram averaging is not biased

by the missing wedge. Additionally, the impact of SNR and polarity distribution on the output

precision of APT should be verified.

For validation purposes we implemented a ground truth data generator (Methods), that creates

modelled tomograms of actin bundles (volume 353 x 353 x 353 nm^3 with a pixelsize of 3.44 Å),

utilizing EMD-6179 [5] as filament density, with three adjustable parameters: the angle φ of

the bundle with the tilt-axis, the SNR of the tomogram [21, 22], and the polarity ratio ρ_r, namely

the fraction of filaments in the bundle oriented in opposite direction. Each modelled bundle

comprises on average 73 filaments, with a uniform length distribution of 86-282 nm and a

uniform distance distribution of 10-37 nm between neighboring filaments.

In total we created three ground truth datasets with three different SNRs, that are 0.01, 0.001,

and 0.0001. Each dataset consists of ten tomograms of different bundles, that are oriented in

10° steps from $\varphi = 0°$ (filaments $\parallel$ to the tilt-axis) to $\varphi = 90°$ (filaments $\perp$ to the tilt-axis).

Additionally, within each set we varied the polarity ratio ρ_r in triplets, meaning that the first four

tomograms ($\varphi = 0°, 10°, 20°, 30°$) were modelled with $\rho_r = 0.1, 0.3, 0.5, 0.1$, and so forth, except

the last tomogram ($\varphi = 90°$) was constructed with a polarity ratio of 0.5.

A top view on one of the modeled bundles ($\varphi = 30°$, $\rho_r = 0.1$) is shown in Fig. 1b (without

noise, only for visualization purposes). A second example is depicted in Fig. 1c. This bundle

is oriented 70° with respect to the tilt-axis and the polarity ratio is 0.3. Its side view is displayed

in Fig. 1d, with the viewing axis oriented parallel to the filaments. A bandpass filtered slice of

a filament, that was boxed out of one of the ground truth tomograms ($\varphi = 0°$, SNR = 0.001), is

shown and can be compared to a filament parallel to the tilt-axis as well, but modelled with a

SNR of 0.0001 (Fig. 1e and f, respectively).

We applied the APT workflow on each of the three ground truth datasets independently. Since

the segmentation of the filaments is given by the ground truth data, we first established the

segment coordinates (*Step I*), with an equidistant spacing of 11 nm, corresponding to ~1000

segments per modelled bundle. Next, the subtomograms were extracted using a box size of

50 x 50 x 50 nm^3 (*Step II*) and subsequently projected using a projection thickness of 11 nm

(*Step III*). In the next step segments were prealigned (*Step IV*), followed by the 2D classification

module (*Step V*). In Supplementary Fig. 4 the results of prealignment and 2D classification are

shown for the modelled dataset with SNR = 0.0001.

Following this, we executed 3D reconstruction (*Step VI*) of the three actin filament averages

from a similar number of particles (~ 10'000 segments per average), distributed approximately

uniform over the modelled bundles. As expected, the resolutions of the filament structures,

drop with decreasing SNR (Supplementary Fig. 5).

Subsequently, we performed polarity determination (*Step VII*) of all modelled bundles.

All ψ^{-1} histograms showed two distinct peaks (Fig. 1g), separated by 180°, indicating the

opposite plus-end orientations of the segments, and the height of the peaks reproduce the

respective polarity ratios. All restored bundles from all three SNR conditions showed

satisfactory agreement with the ground truth bundles (Fig. 1h-i). Furthermore, we visualized

the $\xi(F_i^j)$ vectors of all the recovered bundles (Fig. 1j). The position of incorrect assigned

segments is randomly distributed and their number increases with decreasing SNR.

The results of the confidence analysis (*Step VIII*) are plotted in Fig. 1k. The ccs values of

bundles with SNR = 0.001 dropped slightly in comparison to bundles with SNR = 0.01, but

show no trend to lower ccs values with increasing φ angle. We conclude, that the missing

wedge induced anisotropic deterioration does not negatively influence the precision of APT for

low SNR values, as found in cellular cryo-tomograms [21, 22]. However, for the markedly

challenging SNR = 0.0001 the ccs values decrease with increasing φ angle.

Finally, we evaluated the total segment error (Fig. 1k), namely the fraction of segments with

incorrect orientation determination compared to the ground truth. For low SNR conditions the

total segment error is ≲10% for all modelled bundles, and increases to ~20% for bundles

modelled with SNR = 0.0001. However, if we evaluate the total filament error, that is the fraction

of filaments with incorrect polarity determination compared to the ground truth, we find that the

polarity of solely 5% of the filaments (34 out of 712 filaments) were determined incorrectly for

the dataset modelled with SNR = 0.0001. This shows that APT is capable to reliably correct

noise induced polarity errors on the filament level.

In-situ actin reconstruction at FAs. The first milestone working with APT is to obtain an actin

filament structure with clear polar features. The calculated transformations will facilitate the

reconstruction of the underlying actin network and polarity distribution.

Here we applied APT on the actin cytoskeleton inside FAs, using a correlative fluorescence

microscopy and cryo-ET approach [15, 28] (Methods). Mouse embryonic fibroblasts (MEFs),

expressing vinculin-venus as a FA marker [32, 33], were cultured on electron microscopy grids

with silicon-oxide support and imaged by fluorescence microscopy (Fig. 2a). Conspicuous FAs

were identified and found again under the electron beam.

Tomograms cannot cover a complete FA (≳3 μm in length), and due to electron dose sensitivity

of the sample, only a single tomogram can be acquired per FA. Therefore, we recorded cryo-

tomograms at a spectrum of positions, namely from proximal regions, where a stress fibre

enters the FA, to distal regions, where they are oriented towards plasma membrane. Seven

cryo-tomograms of FAs (Fig. 2b), covering positions from proximal to distal, were selected in

the first place, and the actin filaments were manually segmented (Fig. 2c).

Following this, we applied the APT workflow on this dataset. Firstly, we extracted a total of

43'400 segments with an equidistant spacing of 11 nm (*Step I*). The box size of the

subtomograms was set to 50 x 50 x 50 nm^3 (*Step II*), and the projection thickness to 11 nm

(*Step III*). After prealignment and 2D classification (*Step IV-V*), we finally selected 20'585

segments (Supplementary Fig. 6a) for 3D reconstruction (*Step VI*).

The obtained in-situ actin filament structure was resolved to 13.8 Å, and allows to

unambiguously detect the position of its plus-end (Supplementary Fig. 6b-d). This shows that

the APT workflow is capable to produce sufficient resolution for subsequent mapping of the

filament's polarity distribution, although the dataset was relatively limited and the filaments

originate from a crowded and dense cellular environment, with multiple possible binding

partners and modulations of their helical symmetry.

In order to efficiently increase the size of the dataset, we used the previous seven manual

segmentations to train a convolutional neural network with EMAN2 [34], capable of detecting

actin filaments (Methods). Using this approach, we additionally created 31 segmentations of

the actin cytoskeleton at FAs.

We then applied the APT workflow on this extended dataset. Here we extracted a total of

247'940 segments (11 nm spacing) with a box size of 36 x 36 x 36 nm^3 and a projection

thickness of 11 nm (*Step I-III*). After prealignment and 2D classification (*Step IV-V*), we finally

selected 72'973 segments (Supplementary Fig. 7, Fig. 2d) for 3D reconstruction (*Step VI*).

Subsequently, we performed a 3D classification with RELION. This approach yielded in-situ

actin filament structures that were resolved to 13.3 Å, 9.4 Å, and 9.1 Å, respectively

(Supplementary Fig. 8, Supplementary Fig. 9). The highest resolved class average is shown

in Fig. 2e, together with a docking of EMD-6179 [5] in Fig. 2f.

Polarity distribution of actin bundles at FAs. Finally, we utilized APT to reconstruct actin

networks in-situ, including the polarity of the filaments.

Therefore, we continued the APT workflow with the polarity determination module (*Step VII*),

based on the in-situ actin filament reconstruction (Supplementary Fig. 6), previously obtained

from seven manual segmentations of actin bundles at FAs. All ψ^{-1} histograms showed two

distinct peaks, separated by 180°. We extracted the $\xi\left(F_i^j\right)$ vectors of the resulting 2893

filaments, and used APT to calculate the confidence scores (*Step VIII*), plotted in Fig. 3a-c.

The minimum reliable ccs was set to 0.6 (Fig. 3c), thereby leaving a total amount of 2146 actin

filaments with determined polarity. The total length of the filaments is ~149 µm, with an average

length of ~70 nm, and a maximum length of ~350 nm (Supplementary Fig. 10a,b).

In Fig. 3d the resulting $\xi\left(F_i^j\right)$ vectors of one of the bundles are displayed. Here we found that

62% of the segments are pointing with their plus-ends towards the cell tip (blue segments),

and 38% in opposite direction (red segments). The actin polarity distribution of this actin bundle

is visualized in Fig. 3e. The blue filaments are pointing with their plus-ends towards the cell tip,

and the red filaments in opposite direction.

The two bundles shown in Fig. 3e-f were acquired at proximal regions of FAs, whereas the two

bundles shown in Fig. 3g-h were acquired at distal regions. We applied APT's topology module

(*Step IX*), and sorted the bundles with increasing topology score (Supplementary Fig. 10c)

from left (τ=0.25) to right (τ=0.87), which reflects their transition from a tendentially mixed

polarity distribution to a predominantly uniform polarity distribution. In Fig. 3e, 65% of the

filaments are directed towards the cell tip, whereas in Fig. 3h this fraction is increased to 96%.

The bundles in Fig. 3e and Fig. 3h are shown from the side in Fig. 3i and Fig. 3j. Furthermore,

the long axis of the bundle in Fig. 3e is shown in Fig. 3k. The data suggests that filaments,

which are directed towards the cell tip, forming the core of the FA actin bundle, and the

reversely directed filaments are organized around this core. The positive topology scores of

bundles with mixed polarity distributions are a further indication for this architectural principle.

If the reversed actin filaments would be diffused into the bundles, the topology score would

approach negative values. In addition, we analysed the polarity distribution in layers parallel to

the support, finding that reversed filaments are enriched on top and bottom of the mixed

polarity bundles (Supplementary Fig. 10d).

The number of filaments within neighbourhood spheres $\mathcal{K}_\epsilon(x_k)$, which were evaluated during

topology analysis (*Step IX*) with a radius of ϵ=40 nm, has its maximum at 6 neighbours

(Supplementary Fig. 10e). This indicates that actin filaments in FA bundles resemble a

hexagonal packing with a mean filament distance of 28 nm (Supplementary Fig. 10f).

Finally, we completed the APT workflow (*Step VII-IX*) for the extended dataset with previously

obtained 31 automatic segmentations (Fig. 2d-f, Supplementary Fig. 7-9). Seven tomograms

were excluded, because they contained ill-defined or multiple bundles with similar orientation,

so that the ψ^{-1} histograms did not exhibit two clearly defined peaks. As before, we set the

minimal reliable ccs to 0.6, thereby leaving a total amount of 6152 actin filaments with

determined polarity. The total length of the analyzed filaments was increased to ~388 μm.

In Fig. 4 we plotted the topology score τ of these 24 bundles versus their normalized position

along a FA (Methods). Strikingly, pronounced uniform polarity distribution (high τ values) can

be found in distal regions of FAs exclusively. For most of the analyzed bundles (Fig. 4, green

dots) the plot suggests a correlation between topology score and normalized FA position. In

other words, within a FA, from proximal to distal regions, the polarity distribution of the actin

filaments transforms smoothly from mixed to uniform actin polarity organization. Interestingly,

this correlation does not hold true, when the FA is not positioned towards a pronounced cell

protrusion (Fig. 4, orange dots).

Discussion

Resolving the polarity of actin filaments inside cells, would provide high-resolution insights of

cytoskeletal processes and their functional organization. Therefore, we developed and

implemented APT, a set of MATLAB scripts and RELION procedures that enables

subnanometric reconstructions and polarity determination of actin filaments in cellular cryo-

tomograms. This general approach can be applied to any cryo-tomograms, in which actin

filaments are observed. Moreover, the usage of APT to reconstruct in-situ actin filament

structures at sub-nanometric precision, opens new avenues to explore complex actin networks

and even structural interactions of actin and actin binding proteins.

The basic concept of APT is to dissect an actin filament into multiple segments, and

subsequently measure their polarity with single particle techniques. Thereby we create a set

of statistically independent estimations of the filament's polarity. Based on majority, the final

polarity of the filament is determined. We validated this approach and found that the total

filament error is below 5%, even for very challenging and low SNR = 0.0001.

A previous approach to decipher actin polarity from tomographic data necessitated chemical

extraction and negative staining of the cytoskeleton [35], in order to increase the SNR of the

tomograms. However, the challenge of resolving actin filament structure and polarity in-situ

requires a tomographic data set from intact cells [16, 17], which involves vitrification of the

sample, and therefore the method needs to successfully handle low SNR [21, 22].

As an example, we applied APT on actin bundles found at FAs. These multi-layered

assemblies, around 200-350 nm in thickness, connect integrin receptors, which anchor the

plasma membrane to the extracellular matrix, via force transduction and actin regulation layers,

with the actin cytoskeleton [14, 15].

Actin bundles at FAs (Fig. 3e-k) resemble a hexagonal packing of the filaments with an average

distance of ~28 nm (Supplementary Fig. 10e,f). The majority of the filaments is directed

towards the cell tip (Fig. 3e-h), and form the core of the bundles, whereas filaments with

reversed direction organize around this core (Fig. 3i-k, Supplementary Fig. 10d).

Ultimately, the polarity distribution depends on the position of the bundle along the FA (Fig. 4).

The more distal the position, the more uniform the polarity distribution, while mixed polarity can

be observed mainly in the proximal regions (Fig. 5). Stress fibers are contractile actin

assemblies [6], enriched with actomyosin interactions, which favor a mixed polarity distribution

of actin filaments. Proximal regions of FAs anchor to stress fibers [10], therefore exhibiting a

mixed polarity actin bundle. However, at distal regions of FAs, more vectoral or even protruding

forces may play a role that are resembled in a more uniform polarity distribution.

Technical developments in sample preparation allowed cryo-ET to provide fundamental

insights into cellular assemblies and processes in-situ [36-38]. However, it also requires the

development of novel tools to analyse the data [25, 34, 39, 40]. APT complements these tools

and allows to analyse actin filaments in physiological relevant processes, which would provide

a better understanding of actin networks, such as the actin cortex and remodelling of actin

during phagocytosis.

Methods

Implementation and code availability. All functionality of APT was implemented as MATLAB

(MathWorks, Natick, USA) scripts and functions. In order to interface with RELION [27] (tested

with version 3.0.4), respective functions to write and read starfiles were programmed.

The software is available at https://www.placeholder.com. It is equipped with a tutorial that

explains APT step-by-step based on modelled tomograms of actin bundles.

Step IV: Prealignment. This is the first of three steps performed in RELION. First, a starfile

was created only passing the name of each projected subtomogram (`rlnImageName`) and its

originating tomogram (`rlnMicrographName`) to RELION. Next, the projected subtomograms

were normalized with `relion_preprocess`. Please note that the index relation $k \leftrightarrow F_i^j$ was

kept invisible for RELION in all the following steps (for example no metadata label

`rlnHelicalTubeID` was provided [31]). Furthermore, CTF correction in RELION jobs was

switched off, because the tomograms were CTF corrected by phase-flipping prior to

subtomogram extraction.

For prealignment of the segments a 2D classification with the prepared starfile as input was

performed. The purpose of this step is to align the (central) actin filament density parallel with

the x-axis, which is the RELION convention for helical reconstruction [31]. Therefore, we

created a set of projection images (the template library) from an actin filament structure, which

was oriented along the x-axis, rotated in 3° increments and subsequently projected. The

template library was used as initial references for prealignment (Supplementary Fig. 4a).

Since the resolution of the actin filament averages was determined by Fourier shell correlation

between the averages and EMD-6179 [5] as an external reference (Supplementary Fig. 5,

Supplementary Fig. 6d, and Supplementary Fig. 9), we were reluctant to use the same

structure for the construction of the template library. For that purpose, we used EMD-10737.

Step V: 2D Classification. Subsequently we performed a second 2D classification in RELION,

passing the psi angles and translations found in the previous step as priors

(`rlnAnglePsiPrior`, `rlnOriginXPrior`, `rlnOriginYPrior`). In contrast to the

prealignment step, it is crucial that this 2D classification is unsupervised, in order to extract the

structural heterogeneity in the data unbiasedly (Supplementary Fig. 4c-f).

The psi search was conducted local around the psi prior. Therefore, it was possible to apply a

second mask (the first mask was applied during the projection step in z-direction), which

allowed to further diminish the influence of neighbouring filaments in y-direction.

This second 2D classification is capable of producing high quality class averages of

filamentous actin from inside intact cells (Supplementary Fig. 6a, Supplementary Fig. 7), and

allows to sort out false positive actin detections and segments of lower quality. Based on this

classification we selected the segments for subsequent 3D reconstruction.

Step VI: 3D Reconstruction. For actin filament averaging a 3D refine job in RELION was

performed with helical reconstruction switched on [31]. As helical rise 27.6 Å and as helical

twist 166.7° was used [5]. In the input starfile, the refined psi angle and translations found in

the previous step were passed as priors plus for each segment a random rot angle as prior

was added and the tilt prior was set to 90° [31] (`rlnAngleRotPrior`,

`rlnAngleTiltPrior`). As initial template the same structure as for construction of the

template library was used, low-pass filtered to 30 Å (Supplementary Fig. 6d).

Ground truth data generator. In the first step of modelling a cryo-tomogram of an actin bundle, filament x-z-coordinates were calculated based on random close packing [41], with a uniform distance distribution in the range between 10-37 nm. As a consequence of the modelled x-z-dimension of the bundles of 282 x 150 nm, on average 73 filaments could be packed in each bundle. Next, the position and length of each filament in y-direction was randomly chosen from a uniform distribution between 86-282 nm.

According to these coordinates, actin filaments based on EMD-6179 [5], were pasted in an empty volume of 353 x 353 x 353 nm^3, however, a random fraction of ρ_r filaments was pasted in the volume after 180° rotation around the x-axis (Fig. 1b-d, red coloured filaments).

After that the modelled bundle was rotated φ degrees around the z-axis, then Gaussian noise was added to adjust the targeted SNR, and finally the missing wedge of a tilt-series with tilt-range from -60° to +60° was applied in Fourier space.

Cell culture. MEFs stably expressing vinculin-venus were cultured in Dulbecco's Modified Eagle's Medium (Sigma-Aldrich, D5671), supplemented with 10% (v/v) fetal bovine serum (Sigma-Aldrich, G7524), 2 mM L-glutamine (Sigma-Aldrich, G7513) and 100 µg/ml penicillin-streptomycin (Sigma-Aldrich, P0781), at 37˚ C and 5% CO_2. The MEFs were applied onto glow-discharged EM grids with a silicon-oxide support film (R1/4, Au mesh; Quantifoil, Jena, Germany). After incubation, the cells were fixed in a 4% paraformaldehyde solution (Sigma-Aldrich, 16005), and washed 3 times with 1x PBS (Fisher Scientific, BP399-1).

Fluorescence microscopy. The EM grids were transferred to a 35x14 mm glass-bottom cell culture dish (MatTek, P35G-0-14-C) for fluorescence microscopy analysis of the FAs. Adherent MEFs were imaged using an automated inverted microscope (DMI4000 B, Leica Microsystems, Wetzlar, Germany) equipped with a fluorescence lamp and a monochromatic digital camera (DFC365 FX, Leica Microsystems). Overview images of the EM grids were acquired using a 10x dry objective (HCX PL Fluotar 10x/0.30, Leica Microsystems). All images

of individual cells (Fig. 2a) were acquired in phase contrast and fluorescence mode using a 63x oil objective (HCX PL APO 63x/1.40-0.6, Leica Microsystems).

Cryo-electron tomography. A drop of 4 µl BSA-coated 10 nm fiducial gold markers (Aurion, Wageningen, Netherlands) was applied on the EM grids before plunging them into liquid nitrogen cooled ethane.

A Titan Krios transmission electron microscope (Thermo Fisher Scientific, Waltham, USA) equipped with a Quantum energy filter and a K2-Summit direct electron detector (Gatan, Pleasanton, USA) was used for cryo-EM data acquisition. The microscope was operated at 300 keV in zero-loss mode with the energy filter slit width set to 20 eV.

The position of the cells and FAs were identified by overlaying the fluorescence signal to low magnification EM images of the grids. At tomography positions image stacks were recorded at each tilt angle in super-resolution mode with an electron flux of ~8 electrons per pixel per second using SerialEM [42]. All tomograms were acquired at a magnification of 42'000x, and a dose-fractionated frame rate of 6 frames per 1.2 s. The tilt-series covered an angular range of -60° to +60°, and were recorded with tilt increments of 2° and a defocus of -4 µm. The accumulated electron dose did not exceed ~75 e$^-$/Å^2.

All image stacks were down-sampled and motion corrected using MotionCorr [43], resulting in a final pixel size of 3.44 Å. Next, CTF correction of the tilt-series was applied [44]. Overview tomograms (Fig. 2b) for actin filament segmentation (Fig. 2c) and subtomograms of actin segments were reconstructed by weighed back-projection using the TOM Toolbox [45].

Normalized bundle position of FA. The tomographic position within a FA was determined by overlaying the light microscopy images of the individual cells with cryo-EM images.

First, phase contrast and fluorescence images were overlaid, hereby enabling the identification of the cell membrane and enclosing grid holes. Second, low magnification EM images (4'800x) of the FA tomography position were overlaid such, that the cell membrane and grid holes coincided with the light microscopy images. Third, the zero-degree tilt-series image (42'000x)

was aligned with the low magnification EM image such, that the cell membrane shape

matched. By increasing the opacity of the EM images, we displayed the FA fluorescence signal

on the tomography position.

We defined the center of the tomogram as the tomography position FA_T. In order to determine

the relative position of FA_T within the entire FA, we defined the proximal end (closest to cell

body) and distal end (closest to cell periphery) fluorescence microscopy signal pixels of the FA

as FA_P and FA_D, respectively. The distance d between FA_T and FA_D and the length l of the FA

(distance between FA_P and FA_D) was measured (the pixel size of the fluorescence microscopy

images was 91 nm). Finally, the normalized bundle position of the FA (plotted on the y-axis in

Fig. 4) was calculated as $1 - d/l$.

Automatic segmentation. We developed a script in MATLAB to automatize the convolutional

neural network (CNN) segmentation with EMAN2 [34, 46].

The method requires three image stacks with fixed x-y-dimension of 64 x 64 pixels as input

files. Firstly, the positive training stack contains 2D images of densities targeted for

segmentation. These images are slices, boxed out of cryo-tomograms. In our case they contain

actin filament densities. Secondly, an image stack that provides for each image in the positive

training stack an associated segmentation mask. Thirdly, the negative training stack, which

contains 2D images of densities not targeted for segmentation, for example gold particles used

as fiducial markers, backprojection rays, tomogram edges, plasma membrane, and regions

without actin filaments in general.

As a starting point we created seven manual segmentations of actin filaments in cryo-

tomograms of FA actin bundles (Fig. 2c) with AMIRA (Thermo Fisher Scientific, Waltham,

USA). The segmentations were skeletonized, then the filaments were dissected in segments,

and around each 3D coordinate of a segment one positive training image was extracted from

the tomogram at respective z-position. For each manual segmentation one positive training

stack was created, limited to 3'000 images.

In order to create the associated segmentation masks, we extended the skeletonized filaments to the diameter of an actin filament. Subsequently, the segmentation masks were extracted from the extended segmentation at the same coordinates as the positive training images. For each manual segmentation one stack with segmentation masks was created, with the same dimension as the associated positive training stack.

Negative training images were extracted at random positions from the tomograms, but the volume of the actin bundle was excluded. For each manual segmentation one negative training stack was created, limited to 3'000 images as well.

Based on these image stacks, we trained one CNN for each manually segmented tomogram, using default EMAN2 data augmentation, network and training parameters. In the next step, the resulting seven CNNs were applied to each of the 31 tomograms in the extended dataset, thereby creating a total amount of 217 segmentations. Finally, the segmentations belonging to the same tomogram were averaged and post-processed in UCSF Chimera [47] with the hide dust command.

Visualization. All isosurface visualizations of actin filament structures and actin bundles were rendered with UCSF Chimera or AMIRA.

For the visualizations of actin bundles in Fig. 3e-k we used EMD-6179 [5] to represent the filaments. Therefore, we used a b-spline registration algorithm to bend the filaments in order to match their 3D shape defined by the segment coordinates [48].

Data availability. Actin filament averages will be deposited in the Electron Microscopy Data Bank under accession codes EMD-abcd$_1$ for the actin filament structure obtained from the manual segmented dataset (Supplementary Fig. 6), and EMD-abcd$_2$, EMD-abcd$_3$, and EMD-abcd$_4$ for the actin filament class averages (Supplementary Fig. 8) obtained from the extended dataset.

Acknowledgements

This work was funded by grants from the Swiss National Science Foundation (SNSF 31003A_179418) and the Mäxi Foundation. We thank Carsten Grashoff for the vinculin labelled MEF cells. We thank the Center for Microscopy and Image Analysis at the University of Zurich.

Author contributions

B. M. prepared samples and recorded data. M. E. developed and implemented the method. B. M., S. S., W. C., and M. T. wrote and tested code. B. M and M. E. analyzed data. O. M. and M. E. conceived the research and wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Holmes, K.C., et al., *Atomic model of the actin filament.* Nature, 1990. **347**(6288): p. 44-9.
2. Pollard, T.D. and G.G. Borisy, *Cellular motility driven by assembly and disassembly of actin filaments.* Cell, 2003. **112**(4): p. 453-65.
3. Merino, F., et al., *Structural transitions of F-actin upon ATP hydrolysis at near-atomic resolution revealed by cryo-EM.* Nat Struct Mol Biol, 2018. **25**(6): p. 528-537.
4. Egelman, E.H., N. Francis, and D.J. DeRosier, *F-actin is a helix with a random variable twist.* Nature, 1982. **298**(5870): p. 131-5.
5. Galkin, V.E., et al., *Near-atomic resolution for one state of F-actin.* Structure, 2015. **23**(1): p. 173-182.
6. Malik-Garbi, M., et al., *Scaling behaviour in steady-state contracting actomyosin networks.* Nat Phys, 2019. **15**(5): p. 509-516.
7. Xu, K., H.P. Babcock, and X. Zhuang, *Dual-objective STORM reveals three-dimensional filament organization in the actin cytoskeleton.* Nat Methods, 2012. **9**(2): p. 185-8.
8. Hervas-Raluy, S., J.M. Garcia-Aznar, and M.J. Gomez-Benito, *Modelling actin polymerization: the effect on confined cell migration.* Biomech Model Mechanobiol, 2019. **18**(4): p. 1177-1187.
9. Geiger, B., J.P. Spatz, and A.D. Bershadsky, *Environmental sensing through focal adhesions.* Nat Rev Mol Cell Biol, 2009. **10**(1): p. 21-33.
10. Burridge, K. and C. Guilluy, *Focal adhesions, stress fibers and mechanical tension.* Exp Cell Res, 2016. **343**(1): p. 14-20.
11. Shemesh, T., et al., *Focal adhesions as mechanosensors: a physical mechanism.* Proc Natl Acad Sci U S A, 2005. **102**(35): p. 12383-8.
12. Zaidel-Bar, R., et al., *Functional atlas of the integrin adhesome.* Nat Cell Biol, 2007. **9**(8): p. 858-67.
13. Legate, K.R., et al., *Integrin adhesion and force coupling are independently regulated by localized PtdIns(4,5)2 synthesis.* EMBO J, 2011. **30**(22): p. 4539-53.
14. Kanchanawong, P., et al., *Nanoscale architecture of integrin-based cell adhesions.* Nature, 2010. **468**(7323): p. 580-4.
15. Patla, I., et al., *Dissecting the molecular architecture of integrin adhesion sites by cryo-electron tomography.* Nat Cell Biol, 2010. **12**(9): p. 909-15.
16. Lucic, V., F. Forster, and W. Baumeister, *Structural studies by electron tomography: from cells to molecules.* Annu Rev Biochem, 2005. **74**: p. 833-65.
17. Beck, M. and W. Baumeister, *Cryo-Electron Tomography: Can it Reveal the Molecular Sociology of Cells in Atomic Detail?* Trends Cell Biol, 2016. **26**(11): p. 825-837.
18. Weber, M.S., M. Wojtynek, and O. Medalia, *Cellular and Structural Studies of Eukaryotic Cells by Cryo-Electron Tomography.* Cells, 2019. **8**(1).
19. Medalia, O., et al., *Macromolecular architecture in eukaryotic cells visualized by cryoelectron tomography.* Science, 2002. **298**(5596): p. 1209-13.
20. Jasnin, M., et al., *The Architecture of Traveling Actin Waves Revealed by Cryo-Electron Tomography.* Structure, 2019. **27**(8): p. 1211-1223 e5.
21. Pei, L., et al., *Simulating cryo electron tomograms of crowded cell cytoplasm for assessment of automated particle picking.* BMC Bioinformatics, 2016. **17**(1): p. 405.

22. Forster, F., et al., *Classification of cryo-electron sub-tomograms using constrained correlation.* J Struct Biol, 2008. **161**(3): p. 276-86.

23. Forster, F., et al., *Retrovirus envelope protein complex structure in situ studied by cryo-electron tomography.* Proc Natl Acad Sci U S A, 2005. **102**(13): p. 4729-34.

24. Schur, F.K., et al., *An atomic model of HIV-1 capsid-SP1 reveals structures regulating assembly and maturation.* Science, 2016. **353**(6298): p. 506-8.

25. Himes, B.A. and P. Zhang, *emClarity: software for high-resolution cryo-electron tomography and subtomogram averaging.* Nat Methods, 2018. **15**(11): p. 955-961.

26. Nogales, E., *The development of cryo-EM into a mainstream structural biology technique.* Nat Methods, 2016. **13**(1): p. 24-7.

27. Scheres, S.H., *RELION: implementation of a Bayesian approach to cryo-EM structure determination.* J Struct Biol, 2012. **180**(3): p. 519-30.

28. Sartori, A., et al., *Correlative microscopy: bridging the gap between fluorescence light microscopy and cryo-electron tomography.* J Struct Biol, 2007. **160**(2): p. 135-45.

29. Punjani, A., et al., *cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination.* Nat Methods, 2017. **14**(3): p. 290-296.

30. Bharat, T.A.M. and S.H.W. Scheres, *Resolving macromolecular structures from electron cryo-tomography data using subtomogram averaging in RELION.* Nature Protocols, 2016. **11**(11): p. 9-20.

31. He, S. and S.H.W. Scheres, *Helical reconstruction in RELION.* J Struct Biol, 2017. **198**(3): p. 163-176.

32. Grashoff, C., et al., *Measuring mechanical tension across vinculin reveals regulation of focal adhesion dynamics.* Nature, 2010. **466**(7303): p. 263-6.

33. Ringer, P., et al., *Multiplexing molecular tension sensors reveals piconewton force gradient across talin-1.* Nat Methods, 2017. **14**(11): p. 1090-1096.

34. Chen, M., et al., *Convolutional neural networks for automated annotation of cellular cryo-electron tomograms.* Nat Methods, 2017. **14**(10): p. 983-985.

35. Narita, A., et al., *Direct determination of actin polarity in the cell.* J Mol Biol, 2012. **419**(5): p. 359-68.

36. Marko, M., et al., *Focused-ion-beam thinning of frozen-hydrated biological specimens for cryo-electron microscopy.* Nat Methods, 2007. **4**(3): p. 215-7.

37. Schaffer, M., et al., *A cryo-FIB lift-out technique enables molecular-resolution cryo-ET within native Caenorhabditis elegans tissue.* Nat Methods, 2019. **16**(8): p. 757-762.

38. Mahamid, J., et al., *Visualizing the molecular sociology at the HeLa cell nuclear periphery.* Science, 2016. **351**(6276): p. 969-972.

39. Song, K., et al., *In situ structure determination at nanometer resolution using TYGRESS.* Nat Methods, 2019.

40. Martinez-Sanchez, A., et al., *Template-free detection and classification of membrane-bound complexes in cryo-electron tomograms.* Nat Methods, 2020.

41. Desmond, K.W. and E.R. Weeks, *Random close packing of disks and spheres in confined geometries.* Physical Review E, 2009. **80**(5).

42. Mastronarde, D.N., *Automated electron microscope tomography using robust prediction of specimen movements.* Journal of Structural Biology, 2005. **152**(1): p. 36-51.

43. Li, X.M., et al., *Electron counting and beam-induced motion correction enable near-atomic-resolution single-particle cryo-EM.* Nature Methods, 2013. **10**(6): p. 584-+.

643   44.   Eibauer, M., et al., *Unraveling the structure of membrane proteins in situ by transfer
644   function corrected cryo-electron tomography.* Journal of Structural Biology, 2012.
645   **180**(3): p. 488-496.
646   45.   Nickell, S., et al., *TOM software toolbox: acquisition and analysis for electron
647   tomography.* J Struct Biol, 2005. **149**(3): p. 227-34.
648   46.   Chen, M., et al., *A complete data processing workflow for cryo-ET and subtomogram
649   averaging.* Nat Methods, 2019. **16**(11): p. 1161-1168.
650   47.   Pettersen, E.F., et al., *UCSF chimera - A visualization system for exploratory research
651   and analysis.* Journal of Computational Chemistry, 2004. **25**(13): p. 1605-1612.
652   48.   Rueckert, D., et al., *Nonrigid registration using free-form deformations: Application
653   to breast MR images.* Ieee Transactions on Medical Imaging, 1999. **18**(8): p. 712-
654   721.
655

Figures

Figure 1 | Polarity determination of modelled actin bundles. (**a**) The workflow of APT consists of consecutive modules. The orange modules symbolize the required input data. The blue modules were implemented in MATLAB, and the green modules were executed in RELION. (**b**) Top view of one of the modelled actin bundles, which were used for the validation of APT. The bundles were rendered without noise for visualization purposes. Here the angle φ between bundle and tilt-axis is 30°, and the fraction ρ_r of actin filaments, that are oriented in opposite direction (red colored filaments) was set to 0.1. (**c**) Top view of an additional bundle with φ increased to 70°, and ρ_r to 0.3. (**d**) The same bundle seen from the side with the viewing axis adjusted parallel to the filaments. The distance between the filaments is 10-37 nm. Scale bar 100 nm. (**e**) In order to quantify the impact of noise on the precision of APT, the bundles were modelled with defined SNRs. The depicted bandpass filtered slice of a filament was extracted from a bundle with SNR = 0.001. For comparison, the filament shown in (**f**) originates from a bundle with SNR = 0.0001. Scale bar 10 nm.

(**g**) The modelled bundles were processed with APT. The plot shows the resulting ψ^{-1}

histogram of the bundle displayed in (**c**). Segments in the blue peak mainly originate from blue

filaments, and the opposite for the red peak. The filaments point with their plus-ends in opposite

directions, therefore the peaks are separated by 180°. Since ρ_r was set to 0.3 in this bundle,

the number of segments in the red peak is ≈30% of the total number of segments. (**h**) The line

plot shows the filament positions and polarities of the bundle, shown in (**c**), as recovered

by APT. It was affected with a SNR = 0.001 in this case, however, the match between original

and recovered bundle (the side view is shown in (**i**), and can be compared to (**d**)) is almost

error free. (**j**) Here the $\xi\left(F_i^j\right)$ vectors of the recovered bundle are depicted. Segments, that are

originating from the same filament are plotted as columns of circles. The colour scheme reflects

to which peak the segments were assigned in the ψ^{-1} histogram, shown in (**g**), and the $\xi\left(F_i^j\right)$

vectors are sorted according to their fraction of segments linked to the blue peak. In this

representation of the filaments, segments with incorrect orientation determination appear as

distortions of otherwise uniformly colored columns. (**k**) Three datasets of ten modelled actin

bundles each, with SNRs of 0.01, 0.001, and 0.0001, were produced, subsequently processed

with APT, and finally the ccs values and errors in the polarity determination were analyzed.

Each cross marks the average of all ccs values in a bundle. The SNR can be identified by the

color scheme: black, SNR = 0.01, red, SNR = 0.001, and blue, SNR=0.0001, respectively.

A regression line was fitted for each SNR condition, and show that there is no trend to lower

ccs values with increasing φ for SNRs ≳ 0.001 (black and red lines). Thus, the missing wedge

induced anisotropy, which is proportional to φ, can be neglected for those SNR levels.

However, for SNRs ≲ 0.0001 a correlation can be detected (blue line). Next, we calculated the

average total segment error for each bundle, plotted as diamond symbols. Respective

regression lines are delineated as dashed lines. Even for the markedly challenging

SNR = 0.0001 the total segment error rarely exceeds 20%. Because APT uses a majority

criterion for final polarity determination of a whole filament, segment errors <50% can

be compensated.

Figure 2 | Cryo-tomography of FAs and sub-nanometer actin filament structure inside cells. Correlated microscopy combining fluorescence microscopy (**a**, **b**, inset) and cryo-ET (**b**) was used to identify FA sites. Scale bar in (**a**) 10 μm. (**b**) A 1.4 nm thick slice through a cryo-tomogram of a FA (**a**, **b**, white circles), shows individual actin filaments and plasma membrane. Scale bar 200 nm. (**c**) Surface rendering view of the FA site. Actin is depicted in yellow, membranes in blue, macromolecules in red and grey, and receptor densities in green. The segmentation of the actin filaments served as input for APT. (**d**) Class averages obtained by 2D classification of the extended dataset. Scale bar 18 nm. (**e**) Further processing yielded a sub-nanometer resolution structure of actin filaments at FAs (Supplementary Fig. 9), shown as grey isosurface. (**f**) The in-vitro structure EMD-6179 [5] (green isosurface) was docked into the in-situ structure. Scale bar 5 nm.

Figure 3 | Architecture of actin bundles at FAs. (**a**) Histogram of the mcs values from 2893

reconstructed actin filaments, extracted from seven manually segmented cryo-tomograms

acquired at FAs. (**b**) Histogram of the respective scs values. (**c**) Plot of the respective ccs

values. Each red cross marks the ccs of one filament, and the scores are plotted as a function

of the bundle orientation φ. The blue crosses indicate the average of all ccs values that are

originating from the same actin bundle, and a regression line was fitted to these values (blue

line). Only filaments reaching a ccs≥0.6 (dashed line) were accepted, therefore 2146

reconstructed actin filaments were used for subsequent bundle visualization and topology

analysis. (**d**) Visualization of the resulting $\xi\left(F_i^j\right)$ vectors of a bundle that was recorded at a

proximal FA region. Segments, that are originating from the same filament are plotted as

columns of circles. Blue segments are oriented with their plus-ends towards the cell tip, and

red segments point in the direction of the cell body. In this bundle 495 filaments with an average

length of 7.1 segments were analysed. Approximately 2/3 of the segments are directed

towards the cell tip. (**e-h**) Top views on the architecture and polarity distribution of four actin

bundles, recorded at proximal (**e,f**) and distal (**g,h**) FA regions. Blue actin filaments are

oriented with their plus-ends towards the cell tip, and red filaments point in the direction of the

cell body. Proximal FA bundles are characterized by a tendentially mixed polarity distribution,

and distal FA bundles converge to a predominantly uniform polarity distribution. (**i,j**) Side views

on the bundles shown in (**e,h**), respectively. (**i**) Filaments, which are directed towards the cell

body (red filaments), are found with a higher probability at outer regions of proximal FA actin

bundles. This relation is displayed in (**k**) as well, showing the long axis of bundle (**e**). In all

visualizations (**e-k**), actin filaments were rendered from EMD-6179 [5], the dashed rectangles

are bounding boxes surrounding the actin bundles. Scale bars 100 nm. Additionally, positions

of plasma membrane are suggested by light grey lines, and support planes by dark grey lines.

Figure 4 | The distribution of actin polarity at FAs. Plot of topology score versus normalized

bundle position along a FA. The green datapoints (fitted by the red dashed regression line)

suggest a correlation between these two parameters. The polarity distribution along FAs

transitions smoothly from mixed to uniform. However, in case that the distal region of a FA

does not coincide with a pronounced cell protrusion, the polarity distribution exhibits a

tendentially mixed character (orange datapoints). Some datapoints in the plot are

accompanied by their fluorescence microscopy signal. The plasma membrane is outlined by

white dashed lines, and the tomogram position is indicated by a red circle. Scale bar 5 µm.

Figure 5 | A model for actin polarity at FAs. Actin bundles within FAs are characterized by

a mixed polarity distribution at proximal regions. However, there is a smooth transition towards

a uniform polarity distribution at distal regions. Actin filaments, which are directed with their

plus-ends towards the cell body (red filaments), are arranged around a central core of filaments

pointing towards the cell periphery (blue filaments).