Aberrant axonal pathfinding and exuberant myelination in an inducible model of neocortical

heterotopia

Authors: Alice M. Li[1,3], Robert A. Hill[2,3*], and Jaime Grutzendler[1,3,4*]

[1] Interdepartmental Neuroscience Program, Yale School of Medicine, New Haven, CT, 06520,

USA.

[2] Department of Biological Sciences, Dartmouth College, Hanover, NH, 03755, USA.

[3] Department of Neurology, Yale School of Medicine, New Haven, CT, 06520, USA.

[4] Department of Neuroscience, Yale School of Medicine, New Haven, CT, 06520, USA.

*Correspondence: robert.hill@dartmouth.edu; jaime.grutzendler@yale.edu

ABSTRACT

Neocortical heterotopia consist of ectopic neuronal clusters that are frequently found in

individuals with cognitive disability and epilepsy. However, their pathogenesis remains poorly

understood due in part to a lack of tractable animal models. We have developed an inducible

model of focal heterotopia that enables their precise spatiotemporal control and high-resolution

optical imaging in live mice. Here we report that heterotopia are associated with striking patterns

of hypermyelinated and circumferentially projecting axons around neuronal clusters. Despite their

aberrant axonal patterns, *in vivo* calcium imaging revealed that heterotopic neurons remain

functionally connected to other brain regions, highlighting their potential to influence global neural

networks. These aberrant patterns only form when heterotopia are induced during a critical

embryonic temporal window, but not in early postnatal development. Our model provides a new

way to investigate heterotopia formation *in vivo* and revealed features suggesting the existence

of developmentally-modulated, neuron-derived axon guidance and myelination factors.

INTRODUCTION

Up to one third of routine postmortem examinations reveal the presence of neocortical

heterotopia[1–4], a heterogeneous group of focal cortical lamination defects characterized by

abnormally positioned clusters of neurons[5,6]. Heterotopia have been linked to many neurological

conditions including epilepsy, intellectual disability, and dyslexia[3,7–12]. However, comprehensive

exploration of their pathogenesis and pathophysiology has been limited by a lack of tools for their

investigation in the live animal. Several genetic, traumatic, chemotoxic and other neocortical

heterotopia models have been described[13–21]. However, these previous models have not been

utilized for cellular intravital optical imaging analyses largely due to the lack of control over the

position and timing of heterotopia induction and/or limited means for targeted cell labelling.

Here, we developed a methodology enabling the specific induction and visualization of

focal heterotopia in the live mammalian neocortex. Our model uses *in utero* electroporation[22–24]

combined with *in vivo* optical imaging, to generate and track layer I cortical heterotopia. When

visualized using label free myelin imaging[25–27], we identify striking patterns of aberrantly projecting

and hypermyelinated axons surrounding the heterotopic neurons. These distinct patterns emerge

only when the heterotopia are induced during a critical embryonic period, suggesting the presence

of locally-derived, developmentally-modulated signals that initiate the abnormal structural

organization and myelination of the heterotopic axons. Identification and characterization of the

neuronal and glial subtypes within the induced heterotopia revealed many consistent features that

are used to define spontaneously occurring heterotopia in both humans[28,29] and rodent

models[18,30–33]. Finally, using genetically encoded calcium biosensors, we reveal that heterotopic

neurons display similar calcium transient frequencies as neighboring layer II/III neurons and

respond to sensorimotor stimulation in behaving mice, opening new possibilities for the

exploration of their influence on cortical function.

RESULTS

Spatiotemporally precise layer I heterotopia generation and visualization in the live mouse

brain

In utero electroporation (IUE) is a powerful technique that permits labeling and genetic

manipulation of targeted cortical neurons[22–24]. As part of the IUE procedure, a thin glass

microcapillary needle is used to deliver genetic material into the lateral ventricle of embryonic

stage animals for the electroporation of neuronal progenitors (Fig. 1a). While implementing IUE

to label and image cortical axons *in vivo*, we unexpectedly found that cortical sites that had been

injected during the procedure developed marked accumulations of labelled neuronal cell bodies

in layer I of adult mice (Fig. 1c, top row). Electroporated pyramidal cells are normally found in

layers II/III of cortical regions, thus their distinct presence at injected layer I sites stood out in

contrast to non-injected surrounding cortical areas (Fig. 1c, top row). *In vivo* labeling of neurons

with a fluorescent dye called NeuO[34] further revealed that the IUE-labelled neuronal cell bodies

constituted only a small fraction of the total neuronal cell bodies ectopically positioned (Fig.1b, c;

for detailed statistics see Supplementary File 1; mean diameter of NeuO$^+$ ectopic cell clusters 474

± 174 μm s.d. in n = 23 mice). Abnormal superficial clustering of neuronal cell bodies is a defining

feature of layer I heterotopia[1,4,28], a subtype of neocortical heterotopia that has been linked to

learning impairments in humans[7,35,36]. Their presence demonstrates that direct injections into the

cortex performed as part the IUE procedure can induce molecularly tractable neocortical

heterotopia *in vivo.*

Aberrant axonal and myelin patterns characterize layer I heterotopia formed during a

critical embryonic time window.

Using this methodology, we set out to examine the cellular composition of the layer I

heterotopia. Interestingly, we observed markedly aberrant patterns of myelinated axons

exclusively associated with the ectopic neuronal clusters (Fig. 2a and Fig. 3a), as visualized *in*

vivo by spectral confocal reflectance microscopy (SCoRe), a technique that enables high

resolution label-free imaging of myelinated axons[25,26]. This was further corroborated by fixed

tissue immunohistochemistry of myelin markers, showing increased oligodendrocyte and myelin

densities at the sites of injection (Fig. 2b-d, Fig. 3e-g, and Figure 3–figure supplement 1; for

detailed statistics see Supplementary File 1). On closer examination *in vivo*, we observed both

myelinated and unmyelinated axons swirling in a nest-like fashion around heterotopic neurons

(Fig. 2a, Fig. 3a-d, and Figure 3–figure supplement 1), often forming thick concentric borders that

were most prominent towards the pial surface (Fig. 3e-g and Video 1). Radially-oriented,

myelinated fiber bundles were also observed projecting out from underneath the heterotopia (Fig.

2b-c and Figure 2–figure supplement 1), similar to previous descriptions of spontaneously

occurring layer I heterotopia[28–31]. Thus, by applying our methodology in combination with SCoRe

microscopy and immunohistochemical analyses, we discovered that distinct axon guidance and

myelin abnormalities occur in embryonically-induced layer I heterotopia.

To investigate whether the marked axonal and myelin changes were dependent on

induction of heterotopia at specific developmental ages, we performed IUE at various time points

during cortical development between embryonic day 14 (E14) to E17, and postnatal day 0 (P0).

We found that whereas embryonic injections always resulted in similar nest-like patterns of

densely myelinated fiber bundles around heterotopic neurons (Fig. 4, Fig. 5b), P0 injections did

not result in robust axon guidance or myelination abnormalities, despite the presence of ectopic

neural clusters (Fig. 5a). Although postnatally-induced heterotopia tended to be smaller and more

variable in size (data not shown), the axonal and myelin abnormalities were never observed even

around larger P0-induced clusters (Fig. 5a). Consistent with this, we found no significant

difference in the myelin or oligodendrocyte cell body densities (Fig. 5c; for detailed statistics see

Supplementary File 1) between heterotopia and contralateral control regions of P0-injected mice.

Likewise, immunolabelling for neurofilament heavy chain (NF-H) did not reveal aberrant

accumulations of axonal fibers (Fig. 5a). Together, our data indicate previously unrecognized

differences in the cellular composition and organization of neocortical heterotopia based on the

timing of their induction.

Astrocyte and microglia cell densities are not altered in induced heterotopia

Given the striking abnormalities in oligodendrocyte production, myelination and axon

pathfinding observed in the embryonically-induced heterotopia, we next wondered whether other

glial cell types such as astrocytes and microglia also exhibited altered morphology or density.

Immunolabelled cortical brain sections against Aldh1L1 revealed no significant difference in the

astrocyte cell density between layer I heterotopia and corresponding contralateral control regions

(Fig. 6a-c; for detailed statistics see Supplementary File 1). Similarly, using Iba1 immunolabelling,

we found no regional differences in microglia density or morphology (Fig. 6d-f; for detailed

statistics see Supplementary File 1). Moreover, there was no evidence of pronounced astrocytic

or microglial accumulation or altered cellular morphology at the borders of the heterotopia

suggesting that a glial scar had not formed due to the embryonic injection. These data suggest

that while heterotopia induce marked oligodendrocyte generation, astrocytes and microglia are

not significantly influenced by potential local factors derived from ectopic neuronal clusters.

Heterotopia contain a variety of excitatory and inhibitory neurons born at different

embryonic ages

To characterize the neural composition of the induced heterotopia, we examined the

expression of neuronal subtype markers Cux1, Tle4, and GAD-67 (Fig. 7). Cux1 is a transcription

factor predominantly expressed in callosal projection neurons in layers II-IV[37], whereas Tle4 is

primarily restricted to deeper corticothalamic projection neurons of layers V and VI[38,39]. We found

both Cux1$^+$ and Tle4$^+$ neurons present to varying degrees in all layer I heterotopia (Fig. 7b, c and

Figure 7–figure supplement 1), consistent with a mixed population of excitatory projection neurons

from different cortical layers. Further analyses revealed the presence of GAD67[+] interneurons,

which occurred at similar densities within heterotopia as in corresponding layer I control regions

(Fig. 7d-f; for detailed statistics see Supplementary File 1). In addition, layer I heterotopia always

contained GAD67[+] puncta, consistent with inhibitory synapses (Fig. 7d, e). Together, our findings

suggest that induced layer I heterotopic neural clusters comprise a diverse cohort of glutamatergic

and GABAergic neurons, in line with the heterogeneous population of neuronal subtypes

previously described in those occurring spontaneously[31,33,40,41].

We next used birthdating techniques to determine whether neurons within the induced

heterotopia originate at similar or different time points during development. 5-Ethynyl-2´-

deoxyuridine (EdU) pulse labelling at E11.5 revealed small but distinct populations of EdU[+]NeuN[+]

cells in all embryonically-generated layer I heterotopia (Figure 7–figure supplement 2), suggesting

that heterotopia contain neurons born during early corticogenesis[42,43]. We also observed cells

"birthdate-labelled" via IUE[22] at E15 (~90% heterotopia in $n > 30$ mice; Fig. 1c) in heterotopia,

suggesting the additional presence of later-born neurons. These data indicate that induced

heterotopia contain neurons born at various stages of cortical development.

Heterotopic neurons exhibit similar calcium dynamics as neighboring normotopic neurons

Despite their linkage to a wide spectrum of neurological disorders, the cellular dynamics

and mechanisms by which individual heterotopic cells contribute to neural circuit dysfunction are

poorly understood. We examined the calcium activity of single cortical neurons in induced

heterotopia and in surrounding, non-injected cortical layer II/III regions of awake head-fixed mice

using the genetically encoded calcium indicator, GCaMP6f. Surprisingly, although we found highly

variable calcium spike patterns of individual cells within heterotopia (Fig. 8a-c and Video 2), there

were no significant differences in their overall calcium spike event frequency, variance, or

synchrony compared to surrounding non-heterotopic layer II/III regions (Fig. 8d; for detailed

statistics see Supplementary File 1). Furthermore, we did not identify any epileptiform activity in

our imaging sessions, although we cannot rule out that continuous recordings could have

revealed sporadic aberrant activity (Fig. 8a-c and Video 2). Interestingly, we found that mice that

had been startled with brief whisker stimulation consistently responded via neuronal calcium

spikes within layer I heterotopia (Fig. 8e, f and Video 3). This result indicates that heterotopic

neurons are connected to other brain areas and could thus influence network function.

DISCUSSION

Here we describe an inducible model that allows the visualization of focal heterotopia at

cellular resolution in the live animal. By mechanically puncturing the cortex as part of the IUE

procedure we induce a targeted layer I heterotopion that can be repeatedly imaged *in vivo*. Pairing

this methodology with intravital fluorescence and label-free (SCoRe) microscopy of neuronal cell

bodies, axons and myelin, as well as time lapse axonal and calcium imaging[22,23,25,34,44], enables

detailed studies of heterotopia in the intact brain.

Many animal models of focal heterotopia have been described however most involve

subcortical or intrahippocampal heterotopia[13,14,17,19,45], which are not easily amenable to intravital

optical imaging due to their distance from the cortical surface. Moreover, other animal models of

more superficially-occurring heterotopia[15,16,18,21,46] have not been imaged successfully at high

resolution *in vivo* mainly as a result of their temporally and spatially unpredictable nature and/or

lack of visible cell transfection. Our methodology improves upon these limitations by employing

minimally invasive microinjections to both induce layer I heterotopia and deliver plasmid DNA or

viral vectors for cellular genetic manipulation in the live mouse.

Using our model, we revealed several striking characteristics of neocortical heterotopia.

First, we discovered that axonal projections within layer I heterotopia formed swirled and

concentric morphologies specifically around heterotopic neuronal clusters. Interestingly there was

no evidence of glial scarring or mechanical tissue barriers surrounding the heterotopia, suggesting

that the directional changes in axon pathfinding were instead likely mediated by disruptions in the

precise balance of repulsive and attractive gradients of local axon guidance cues[47–51]. These axon

pathfinding abnormalities led us to investigate whether the developmental timing of heterotopia

induction could modulate their formation. We found that whereas embryonically-induced

heterotopia always displayed distinct axon guidance abnormalities, similar defects were not

observed in postnatally-induced heterotopia. This surprising finding suggested the presence of a

critical gestational time period during which cortical axon pathfinding is uniquely sensitive to local

environmental disturbances. The distinct responsiveness of axons during this critical period could

be mediated by the enhanced expression of axon growth cone receptors that increase the ability

of axons to extend toward guidance cues[52,53]. However, developmentally modulated changes in

the signaling gradients of guidance cues themselves could also play a role[54].

In addition to aberrant axonal organization, we used label-free SCoRe *in vivo* imaging to

discover restricted hypermyelination specifically within the heterotopia. This focal

hypermyelination was due to an increased production of myelinating oligodendrocytes that

deposited myelin sheaths primarily along the aberrantly projecting and concentric axons.

Interestingly, the hypermyelination was only observed in embryonically-induced but never in

postnatally-induced heterotopia. Similar to the critical period for disrupted axonal patterning, these

data suggest that the myelination of embryonically-induced heterotopia is instructed by

developmentally regulated molecules or biophysical cues, which could accelerate the local

production of myelin by stimulating oligodendrocyte precursor cell recruitment and/or

differentiation. Myelin formation is influenced by axon caliber[55] and genetic manipulations

increasing axon caliber can trigger oligodendrocyte precursor cell proliferation and the

ensheathment of classically unmyelinated fibers[56]. Thus, axon caliber could serve as one

mechanism that differs between the aberrantly projecting axons found in embryonically-induced

heterotopia compared to adjacent non-myelinated cortical axons. Other potential causes for the

localized hypermyelination of heterotopic axons could include cell surface markers or molecular

identity. Specific subpopulations of excitatory[57] and inhibitory[58] neurons exhibit variable

myelination patterns, seemingly not directly linked to axon caliber, but instead potentially due to

their differential expression of adhesion molecules or altered patterns of neuronal activity. Our

model provides a novel means to test these possibilities in future studies.

Despite the striking axon and myelin abnormalities associated with the induced

heterotopia, we found no evidence of spontaneous epileptiform activity occurring in heterotopic

regions or nearby non-heterotopic cortices of awake animals, consistent with previous

studies[40,59,60]. Moreover, spontaneous calcium fluctuations and relative synchrony of firing

between neurons in layer I heterotopia and neighboring layer II/III cortex were similar. This lack

of observable difference in activity may reflect the similar neural composition of both brain areas,

which could enable a similar balance of excitatory and inhibitory inputs to these regions. Indeed,

we identified Cux1[+] cells, GABAergic neurons and GABAergic puncta in layer I heterotopia that

are also prevalent in layers II/III[37,61–63]. A second, non-mutually exclusive possibility is that more

metabolically demanding conditions are required to trigger aberrant activity in mice with induced

heterotopia. Consistent with this hypothesis, the application of normally subthreshold doses of

convulsant drugs can provoke epileptiform activity in mice with heterotopia *in vitro*[64] and *in*

vivo[59,64,65]. Other animal models of heterotopia, such as the *tish* mutant[13] and Ihara's genetically

epileptic rat[19] have documented spontaneous epileptiform activity *in vivo*. Reasons for this

difference from our findings remain unclear, however they may be related to disparities in the

number, location or size of heterotopia, or unknown off-target effects of the inherited mutations

themselves in the genetic models.

There is little known about the functional dynamics of individual heterotopic neurons during

behavioral stimulation in the live animal. Using time-lapse calcium imaging, we revealed that

heterotopic neurons display robust responses following brief whisker stimulation in awake mice,

consistent with the functional connectivity implicated by previous anatomical, electrophysiological,

and behavioral studies[13,40,59,66–69]. This functional connectivity suggests that, even though we did

not detect aberrant spontaneous activity within heterotopia, these neurons are integrated into the

local neural network and are potentially capable of altering neural network function at baseline or

under metabolically demanding conditions.

We describe an inducible model of heterotopia amenable to intravital visualization that

closely mimics the neural, glial, axonal, and myelin profiles of those occurring spontaneously in

layer I[28,33]. Building upon previous models, this system employs *in utero* microinjections to directly

provoke the disruptions in pial basement membrane and radial glial scaffolding thought to

culminate in the neural migration defects leading to their development[70]. Importantly, this model

allows imaging of the downstream effects of this disruption using multiple optical, functional, and

molecular probes for different cell subtypes[22,23,25,34,44], adding a powerful new tool for the study of

cortical malformations in the live mammalian brain.

MATERIALS AND METHODS

Animals

All experimental approaches and procedures were conducted in accordance with Yale

University Institutional Animal Care and Use Committee regulations. Timed pregnant outbred

CD1 mice were purchased from Charles River Laboratories, Inc, with the first 24 hours of

postnatal life designated as P0. We included both male and female mice, aged to P30 – P60, for

this study. For some birth dating experiments as described in the text, pregnant dams were given

a single intraperitoneal (i.p.) injection of EdU (30 µg g^{-1} body weight) at E11.5, prior to IUE surgery

at E15. Litters were kept in individual ventilated cages until weaning age (P21), after which mice

were housed in single-sex groups with 2 – 5 animals per unit. Cages were maintained in

temperature-controlled facilities with 12-hour light/12-hour dark cycles.

In utero intracranial injection, electroporation and viral infection

Embryonic cortical injections were performed as part of the IUE procedure during needle

insertion into the lateral ventricle[23,71]. Embryos were injected once unilaterally

between embryonic day 14 – 17 (E14 - E17) for all electroporations, as specified in the

text. Intracranial injections were not performed at gestational ages below E14 or over E17 due to

the technical challenges associated with maintaining embryo viability. To

keep consistent experimental design parameters, however, all quantifications for *in vivo* and

fixed tissue analyses used only E15-injected embryos.

All *in utero* injections were targeted toward prospective somatosensory cortices.

Injection solutions included the following plasmid and viral components for neuronal

labelling: pCAG-tdTomato (based on Addgene plasmid 11150) and rAAV8-hSyn-eGFP (UNC

Vector Core, Lot AV5075D; titer 3.9e12 GC ml^{-1}). Injection solutions contained either pCAG-

tdTomato (1.5 µg µL^{-1} final concentration) only, or both pCAG-tdTomato (1.5 ug µl^{-1} final

concentration) and AAV8-hSyn-eGFP (final titer 3.9e11 GC ml^{-1}) in saline solution. All injection

solutions contained Fast Green FCF dye (TCI; 2 mg ml^{-1}) to facilitate their visual tracking during

the injection procedure. Procedures for *in utero* electroporation have been previously

described[23,71]. Briefly, timed pregnant CD1 dams were anesthetized with a saline solution

containing both ketamine (100 - 120 mg kg^{-1}) and xylazine (10 - 12 mg kg^{-1}), delivered i.p.

Following induction of deep surgical anesthesia, midline incisions (1¼ inch) were made into the

abdominal skin and muscle wall to access the underlying uterus. Pulled glass capillary needles

(10 µL Drummond Scientific Glass Capillaries, Cat# 3-000-210-G; pulled to ~ 50 µm diameter at

tip and ~ 125 µm diameter at 1mm above tip) were then used to puncture the uterine wall and

deliver (Picospritzer II, General Valve) 0.5 µl of injection solution into the lateral ventricle of

individual embryos. Each embryo was injected only once and subsequently electroporated using

BTX tweezertrodes aimed at the somatosensory cortex of the injected hemisphere.

All electroporations were conducted using four 50 ms, 50 V electrical pulses, delivered at 1

second intervals (BTX Harvard Apparatus 8300 pulse generator).

Postnatal intracranial viral injection

Intracranial injections were performed on P0 pups from timed pregnant CD1 dams that

were naïve to the *in utero* electroporation procedure. Injection solutions included one of the

following viruses to mark injected regions: rAAV8-hSyn-eGFP (UNC Vector Core, Lot AV5075D;

titer 3.9e12 GC ml^{-1}) and rAAV2-CaMKIIa-mCherry (UNC Vector Core, Lot AV4377d, 3.8e12 titer

GC ml^{-1}). P0 intracranial injections were performed essentially as previously described[72]. Briefly,

neonates were cryoanesthetized[73] within 24 hours following birth. After confirming loss of

voluntary movement, neonates were placed in the prone position on a polymer cooling block. A

pulled glass capillary needle (10 µl Drummond Scientific Glass Capillaries, Cat# 3-000-210-G;

pulled to ~ 75 µm diameter at tip and ~ 325 µm diameter at 4 mm above tip), advanced 4mm past

the dorsal scalp at a 90° angle, was then used to deliver into the brain parenchyma 1 µl of virus

solution (diluted 1:10 in PBS and 2 mg ml^{-1} Fast Green FCF, TCI from stock). Each neonate

received only one unilateral intracranial injection targeted over somatosensory cortices. Injected

neonates were then rewarmed on a heating pad and placed back with their biological mother. All

P0-injected mice were sacrificed for analysis at P30, with quantifications and analyses performed

in fixed tissue to circumvent the obstructive meningeal scarring and parenchymal adhesions

associated with P0-injections on *in vivo* imaging.

Postnatal subarachnoid viral injection

Injection solutions were prepared using AAV9-Syn-GCaMP6f-WPRE-SV40 (Penn Vector

Core, Lot CS1001; titer 7.648e13 GC ml^{-1}) virus diluted 1:100 in PBS and Fast Green FCF (TCI;

2 mg ml^{-1}). P21- P30 mice that had previously been electroporated *in utero* were anesthetized by

intraperitoneal injections of ketamine (100 mg kg^{-1}) and xylazine (10 mg kg^{-1}). The scalp was

shaved, cleaned, and then incised to expose the underlying bone. A high speed drill was next

employed to introduce a small burr hole (~ 0.75 mm diameter) over the transfected hemisphere,

taking care to avoid cortical regions suspected to have been directly punctured as part of the *in*

utero electroporation procedure. The underlying dura was gently detached, and a 12 µL volume

of injection solution (prepared as described above) was infused into the subarachnoid space to

achieve viral transfection of both layer I heterotopic and layer II/III neurons via topical cortical

application. The scalp incision was then closed with sutures. Cranial windows for *in vivo* imaging

were prepared over injected cortical hemispheres 3 – 4 weeks following the subarachnoid AAV

infusion.

Cranial window surgery and in vivo imaging

All *in vivo* imaging was performed using cranial windows[26]. Briefly, mice were

anesthetized using ketamine (100 mg kg^{-1}) and xylazine (10 mg kg^{-1}), delivered via i.p. injection.

The dorsal skull surgical field was shaved and cleaned, and a ~4 mm diameter circular region of

skull and dura mater was excised from the injected hemisphere. A #0 transparent glass coverslip

was then gently implanted on top of uncovered pial surface to serve as the cranial window. Glue

and dental cement were applied to secure the window to surrounding skull bone.

To detect neuronal cell bodies *in vivo* as described in the text, the fluorescent membrane-

permeable probe NeuO (NeuroFluor, Stemcell Technologies Cat# 01801, diluted 1:25 in PBS)

was applied to exposed cortex for 20 min followed by a 1-2 min PBS rinse, before the placement

of the #0 glass coverslip during the cranial window surgery. In some cases, 100 µL Evans blue

(TCI; 1 mg ml^{-1}) was injected intravenously after the cranial window surgery to label the cortical

vasculature.

Except for GCaMP6f calcium imaging experiments, all *in vivo* imaging studies used mice

that were anesthetized via i.p. ketamine and xylazine injection. *In vivo* imaging of anesthetized

mice was performed at P30 immediately after cranial window surgery. All

intravital GCaMP6f calcium imaging was carried out in P50 - P60 awake head-fixed mice starting

four hours after arousal from cranial window surgical anesthesia.

Confocal *in vivo* images of previously injected cortical areas with or without the needle

tract sites were acquired using a 20X water immersion objective (Leica, 1.0 NA) on a Leica SP5

upright laser scanning microscope. Spectral confocal reflectance (SCoRe) imaging to detect

myelinated axon segments was performed as previously described[25,26] by capturing the

simultaneously reflected light signals from 488 nm, 561 nm, and 633 nm multi-wavelength laser

excitation outputs. Single-photon laser outputs were tuned to the following excitation wavelengths

for fluorescence imaging: 488 nm for GFP and NeuO; 561nm for tdTomato and mCherry; 633nm

for Evans Blue. Sequential imaging was employed to minimize overlap between SCoRe reflection

and individual fluorescence emission signals for all *in vivo* confocal imaging experiments.

Needle tracts from injections performed *in utero* were identified by the abrupt changes in

orientations of labelled dendritic and axonal processes around breaks in the cortical surface *in*

vivo. Since needle tracts identified in this manner always displayed ectopic neural clusters of

similar expanse in layer I, the outer boundaries of 'heterotopia' *in vivo* were defined as the layer I

needle tract borders for all quantifications. Ipsilateral, non-injected cortical areas were defined as

layer I regions at least 150 μm away from a discernable injection site border. Heterotopia and

ipsilateral control regions were imaged using identical laser output and image acquisition

configurations that were determined for each experimental data set. Confocal z-stacks were

acquired at 1024 × 1024 pixel resolution starting from the pia through depths of up to 120 μm

below the cortical pial surface. In some cases, time lapse imaging of *in utero*-induced

heterotopia was also performed at 512 X 512 pixel resolution.

For some GCaMP6f calcium imaging experiments that involved visualization of deeper

cortical regions in layers II/III as indicated in the text, time-lapse fluorescence images were

acquired using a 20X water immersion objective (Zeiss, 1.0 NA) on a Prairie Technologies two-

photon microscope fitted with a mode-locked, tunable Spectra Physics Mai-Tai laser. In these

experiments, the two-photon laser was tuned to 920nm for excitation of both GCaMP6f and

TdTomato fluorophores. All two-photon time-lapse imaging was performed at 512 X 512

pixel resolution.

In vivo image processing and quantification

Except for GCaMP6f calcium imaging experiments which were conducted using P50 - P60

mice, all intravital imaging studies were performed using P30 mice, with quantifications carried

out on those that had previously been injected at E15. All *in vivo* images were processed and

quantified using ImageJ/FIJI.

For SCoRe density quantifications, we analyzed single z-sections located 10 μm deep to

the cortical pial surface from both heterotopic and ipsilateral control regions. SCoRe density

values were assessed using a custom-built, automated thresholding and binarization macro in

Image/FIJI, with Robust Automatic Threshold parameters set to noise = 25, lambda = 3, min =

31. Randomly selected, equally sized regions of interest (ROIs) were used to

determine SCoRe density values within the centers and edges of the heterotopia versus

control regions. A 'heterotopion center' was defined as the circular region with a radius extending

from the needle tract center to 1/3 of the radius of the needle tract. A 'heterotopion edge'

was defined as the needle tract concentric circular region just outside the 'heterotopion center',

extending from the 'heterotopion center' outer edge to the outermost border of the needle tract.

Average SCoRe densities were determined for the heterotopion center, heterotopion edge and

control area for each mouse (n = 8 mice). Statistical analyses were carried out using Wilcoxon

matched pairs, signed-rank non-parametric tests.

For NeuO dye-labelled cell body density quantifications, data were analyzed from the first

75 μm of cortex deep to the pial surface of both heterotopic and control regions in layer I. Equal-

sized volumes (50 μm^3) were randomly selected from the superficial cortical z-stacks captured

from both heterotopic and control regions. The number of NeuO$^+$ cell bodies was manually

counted in each volume. Average NeuO$^+$ cell body densities were then determined for

the heterotopic and control region of each mouse (n = 6 mice). Statistical analyses were

performed using Wilcoxon matched pairs, signed-rank non-parametric tests.

For quantification of neuronal calcium dynamics in heterotopia vs. adjacent non-injected

cortical areas, we used the two-photon microscope to capture 150 X 150 μm field of view

(FOV) time lapse images from both layer I heterotopia and non-injected ipsilateral layer

II/III regions. Time lapse imaging (512 X 512 pixel resolution; 2Hz) was performed in awake,

head-fixed mice during a 2-hour time window that started 4 hours after their arousal from cranial

window surgical anesthesia. For heterotopic regions, FOVs were randomly selected from within

the first 75 μm below the cortical surface in layer I. For layer II/III regions, FOVs were randomly

selected from between 120 to 175 μm below the cortical surface. Six to eight separate FOVs

(exactly half from layer I heterotopia and half from ipsilateral layer II/III regions) were imaged in

each mouse, with each FOV recorded for 120s per trial for three trials. The order in which FOVs

were acquired from heterotopia and layer II/III regions was alternated between mice. Time series

analyses were performed using ImageJ/FIJI. Prior to quantifications, TurboReg plugin in

ImageJ/FIJI was used to align all time series images in the XY plane.

To quantify GCaMP6f fluorescence changes in individual cells, ROIs were manually

selected to encapsulate neuronal cell bodies. GCaMP6f$^+$ cell bodies that exhibited tdTomato

labelling from the IUE procedure were excluded from all analyses. Approximately 10 - 25

GCaMP6f$^+$ cells were analyzed in each FOV. Baseline GCaMP6f fluorescence intensity values

(F) for each cell were defined in each trial as the average of the lowest 20% of the recorded values

per trial, with fluctuations from this baseline denoted as ΔF/F. A spike event was defined in each

trial as a ΔF/F > 0.5. Spike event frequencies were averaged across 3 trials for each cell in both

layer I heterotopia and layer II/III regions for n = 7 mice. Statistical analyses were performed using

Wilcoxon matched pairs, signed-rank non-parametric tests.

For quantifications of the global synchronization index reflecting the relative coordination

of GCaMP6f activity, analyses were performed using Fluorescence Single Neuron and Network

Analysis Package (FluoroSNNAP)[74]. Briefly, this semi-automated software implements a

correlation matrix-based algorithm[74–76] to compute the normalized global synchronization indices

(ranging in value from 0 to 1) for time series calcium imaging data of neural populations. The

highest value, 1, signifies entirely synchronized firing throughout an identified cluster of neurons,

whereas 0 indicates the total absence of synchrony. Synchronization cluster analyses were

performed using template-based calcium event detection parameters set to detection threshold =

0.85, minimum size of synchronization clusters = 2, and surrogate resampling = 20. A cluster of

neurons was defined as the GCaMP6f$^+$ cell population within one FOV. Global synchronization

indices were averaged across three trials for each FOV in both layer I heterotopia and layer II/III

regions for n = 7 mice. Statistical analyses were performed using Wilcoxon matched pairs,

signed-rank non-parametric tests.

Tissue processing and immunohistochemistry

At P30, mice were deeply anesthetized using ketamine and xylazine, and then perfused

transcardially with 4% paraformaldehyde in phosphate buffered saline (PBS). Harvested brains

were post-fixed overnight using the same solution at 4 °C, and then vibratome sectioned (coronal;

75 µm thickness) for fixed tissue analysis.

To identify brain regions that had been injected as part of the IUE procedure, a

fluorescence microscope was next used to examine the sections for visibly disrupted layer I

regions showing transfected neurons and/or neuronal processes, similar to what has been

previously described[46]. The disrupted regions often revealed cellular clusters

containing eGFP$^+$ neurons that protruded past the pial surface, which were never observed in

intact cortical regions or in non-injected mice. Sections containing visibly disrupted layer I cortical

cytoarchitecture indicative of injection-associated trauma were then selected for further

processing. Immunohistochemistry, performed as described below, was used to confirm the

presence of layer I heterotopia at all identified IUE injection sites. Coronal sections from P0-

injected mice were screened in a similar manner for injected areas, which were identified by virally

transfected axonal fibers and neuronal cell bodies concentrated around visible needle tracts (data

not shown).

Prior to immunostaining, all free-floating sections were heated to 95 °C for 30 minutes in

50 mM sodium citrate buffer (0.05% Tween-20, pH 6.0) for antigen retrieval and elimination of

endogenous eGFP expression. After rinsing the sections in PBS at room temperature, EdU

labelling was next carried out in some experiments as specified by the Click-iT EdU Alexa Fluor-

647 Imaging Kit protocol (Cat# C10340). Before proceeding to antibody staining, all tissues

were pre-incubated for 1-2 hr in 0.1% Triton X-100 and 5% Normal Goat Serum (NGS;

Jackson Immunoresearch, Cat# 005-000-121) in PBS at room temperature. Slices were then

incubated with primary antibody for 1.5 hrs - 2 days as needed in 0.1% Triton X-100 and 5% NGS

in PBS at 4 °C. The primary antibodies used were: rabbit anti-NeuN (Abcam, Cat# ab177487,

1:3000), mouse anti-NeuN (Abcam, Cat# ab104224, 1:1000), mouse anti- Myelin CNPase (clone

SMI 91, Biolegend, Cat# 836404, 1:1000), rabbit anti-Cux1 (Novus Biologicals, Cat# NBP2-

13883, 1:100), mouse anti-Tle4 (E-10) Alexa Fluor 647 (Santa Cruz Biotechnology, Cat# sc-

365406 AF647, 1:100), mouse anti- GAD-67 (EMD Millipore, Cat# MAB5406, 1:1000), rabbit anti-

Iba1 (Wako, Cat# 019-19741, 1:600), mouse anti-Aldh1l1 (clone N103/39, NeuroMab, Cat# 75-

140, 1:500), chicken anti-Neurofilament NF-H (EnCor, Cat# CPCA-NF-H, 1:500), chicken anti-

GFP (Abcam, Cat# 13970, 1:500), and rabbit anti-Myelin Basic Protein (MBP; Abcam, Cat#

40390, 1:1000). Following primary antibody incubation, slices were washed in PBS and then

incubated with Alexa Fluor dye-conjugated secondary antibodies of the appropriate host species

at 1:600 dilution in 0.1% Triton X-100 and 5% NGS in PBS for 1 - 2 days at 4 °C. After secondary

antibody incubation, sections were washed again in PBS and incubated with 2.5 µg mL^{-1} 4′,6-

diamidino-2-phenylindole (DAPI) in PBS for 15 min at room temperature to counterstain cell

nuclei. Following an additional subsequent wash, stained sections were mounted with 25%

mounting media solution (Dako Ultramount, Cat# S1964; diluted 1:4 in PBS) onto glass slides for

imaging.

Fixed tissue imaging and quantification

Images of stained sections were collected using a Leica SP5 upright confocal laser

scanning microscope. Lower magnification views of analyzed sections for presentation were

acquired using 10X and 20X Leica objectives. All fixed tissue images for quantification were

captured through a 40X Leica water immersion objective. For immunohistochemical analyses of

in utero-injected mice, 'heterotopia' were defined as the layer I cortical regions demarcated by

ectopic NeuN$^+$ cell body clusters identified at IUE needle tract sites. For P0-injected mice,

because the heterotopia associated with needle tracts were more variable in size, ranging from

several cells to 350 µm in diameter (data not shown), data for these injected areas were obtained

from 300 µm X 90 µm FOVs centered on visible needle tracts that were within the first 100 µm

below the pial surface. Control, non-injected regions were specified on corresponding

contralateral cortices at similar mediolateral distances from the midline. Identical laser output and

image acquisition configurations were used to capture images of both heterotopia and

contralateral control regions throughout each immunostaining set.

All data for Aldh1l1, Iba1, CNPase, and GAD-67 cell density quantifications were acquired

from the most superficial 100 μm of cortical tissue of both heterotopic and contralateral

homologous control regions in layer I. The number of cell bodies confirmed by DAPI labelling that

were positive for each marker was manually counted in randomly selected volumes within z-

stacks acquired from both regions. The volumes were kept at identical dimensions between

heterotopia and contralateral control regions for each experiment. For GAD-67 immunostaining,

the presence of neuronal cell bodies was also confirmed by NeuN labelling. Two coronal sections

containing heterotopia were analyzed per animal, with each immunostaining set comprising $n = 6$

animals. Statistical analyses were performed using Wilcoxon matched pairs, signed-rank non-

parametric tests.

Data for CNPase density quantifications were acquired from the most superficial 100 μm

of cortical tissue in both heterotopia and contralateral control regions. CNPase intensity was

assessed using an automated thresholding and binarization plugin in Image/FIJI (Robust

Automatic Threshold parameters set to noise = 1, lambda = 2, min = 208) for equally-sized ROIs

that were randomly selected from within heterotopic and contralateral control z-projections (15

μm thickness). CNPase density values for each section were determined by subtracting the

automated measurements made in additional background regions from those made in assessed

heterotopic and control regions. Two coronal sections containing heterotopia were analyzed per

animal ($n = 6$ animals). Statistical analyses were performed using Wilcoxon matched pairs,

signed-rank non-parametric tests.

To examine the neuronal composition of the embryonically-induced heterotopia, coronal

sections were stained for NeuN, DAPI and one of the other following stains: Cux1, Tle4,

or EdU as indicated in the text. Data were acquired from within the most superficial 100 μm of the

501 neocortex for layer I heterotopia, and across the entire cortical wall for contralateral homologous

502 control regions. For quantification of layer I heterotopia, equally sized VOIs were randomly

503 selected from within the first 100 μm below the pial surface. For quantification of control regions

504 on the contralateral hemisphere, the cortical wall was partitioned into 11 equal volume bins

505 extending from the pial surface to white matter, with Bin1 encompassing layer 1 and Bin 11

506 adjoining white matter at its lower boundary. For each heterotopic VOI and control bin, the 1)

507 number of NeuN$^+$ cells and 2) proportion of NeuN$^+$ cells that were also Cux1, Tle4,

508 or EdU positive was manually quantified in ImageJ. Cells were identified as EdU positive if >

509 50% of their nuclear volume, defined by DAPI, was occupied by EdU. Two sections were

510 analyzed per mouse, with sample sizes for each group denoted in the text and figure legends.

511  The sites of injection analyzed in this study were positioned in somatosensory cortices

512 and immediately adjacent areas in both male and female mice. Given the minimal variation in the

513 morphological appearances of needle tracts between male and female mice and across different

514 cortical areas, we pooled together these variables in our analyses. No animal subjects or

515 experimental data points were excluded from analysis, and the nature of our study did not require

516 subject randomization or experimenter blinding. No statistical methods were used to predetermine

517 sample sizes, although our sample sizes are comparable to those published and generally

518 accepted in the field. GraphPad Prism 7 was utilized for all statistical analyses, and Wilcoxon

519 matched pairs signed-rank non-parametric tests were used to determine statistical significance

520 (declared for p-values below 0.05; two-tailed) because a normal distribution of differences

521 between the paired data could not be assumed. Data are reported as mean ± s.e.m., unless

522 otherwise noted.

REFERENCES

1. Kaufmann WE, Galaburda AM. Cerebrocortical microdysgenesis in neurologically normal

subjects: a histopathologic study. *Neurology*. 1989;39(2 Pt 1):238-244.

doi:10.1212/WNL.39.2.238

2. Kasper BS, Stefan H, Buchfelder M, Paulus W. Temporal Lobe Microdysgenesis in

Epilepsy Versus Control Brains. *J Neuropathol Exp Neurol*. 1999;58(1):22-28.

doi:10.1097/00005072-199901000-00003

3. Meencke HJ, Veith G. Migration disturbances in epilepsy. *Epilepsy Res Suppl*. 1992;9:31-

39; discussion 39-40. http://www.ncbi.nlm.nih.gov/pubmed/1285910.

4. Schulze KD, Braak H. Hirnwarzen. *Zeitschrift fur Mikroskopisch-Anatomische Forsch -*

Abteilung 2. 1978;92(4):609-623.

5. Sisodiya SM. *Malformations of Cortical Development: Burdens and Insights from*

Important Causes of Human Epilepsy. Vol 3. Lancet Publishing Group; 2004:29-38.

doi:10.1016/S1474-4422(03)00620-3

6. Barkovich AJ, Gressens P, Evrard P. Formation, maturation, and disorders of brain

neocortex. *Am J Neuroradiol*. 1992;13(2):423-446.

http://www.ncbi.nlm.nih.gov/pubmed/1566709.

7. Galaburda AM, Kemper TL. Cytoarchitectonic abnormalities in developmental dyslexia: a

case study. *Ann Neurol*. 1979;6(2):94-100. doi:10.1002/ana.410060203

8. Hardiman O, Burke T, Phillips J, et al. Microdysgenesis in resected temporal neocortex:

incidence and clinical significance in focal epilepsy. *Neurology*. 1988;38(7):1041-1047.

doi:10.1212/wnl.38.7.1041

9.   Palmini A, Andermann F, Olivier A, Tampieri D, Robitaille Y. Focal neuronal migration

disorders and intractable partial epilepsy: Results of surgical treatment. *Ann Neurol*.

1991;30(6):750-757. doi:10.1002/ana.410300603

10.   Kakita A, Hayashi S, Moro F, et al. Bilateral periventricular nodular heterotopia due to

filamin 1 gene mutation: Widespread glomeruloid microvascular anomaly and dysplastic

cytoarchitecture in the cerebral cortex. *Acta Neuropathol*. 2002. doi:10.1007/s00401-002-

0594-9

11.   Farrell MA, DeRosa MJ, Curran JG, et al. Neuropathologic findings in cortical resections

(including hemispherectomies) performed for the treatment of intractable childhood

epilepsy. *Acta Neuropathol*. 1992;83(3):246-259.

http://www.ncbi.nlm.nih.gov/pubmed/1557956.

12.   Nordborg C, Eriksson S, Rydenhag B, Uvebrant P, Malmgren K. Microdysgenesis in

surgical specimens from patients with epilepsy: occurrence and clinical correlations. *J

Neurol Neurosurg Psychiatry*. 1999;67(4):521-524.

http://www.ncbi.nlm.nih.gov/pubmed/10486403.

13.   Lee KS, Schottler F, Collins JL, et al. A genetic animal model of human neocortical

heterotopia associated with seizures. *J Neurosci*. 1997;17(16):6236-6242.

doi:10.1523/jneurosci.17-16-06236.1997

14.   Bai J, Ramos RL, Ackman JB, Thomas AM, Lee R V., LoTurco JJ. RNAi reveals

doublecortin is required for radial migration in rat neocortex. *Nat Neurosci*.

2003;6(12):1277-1283. doi:10.1038/nn1153

15.   Rosen GD, Sherman GF, Richman JM, Stone L V., Galaburda AM. Induction of molecular

layer ectopias by puncture wounds in newborn rats and mice. *Dev Brain Res*.

1992;67(2):285-291. doi:10.1016/0165-3806(92)90229-P

16. Brunstrom JE, Gray-Swain MRR, Osborne PA, Pearlman AL. Neuronal Heterotopias in

the Developing Cerebral Cortex Produced by Neurotrophin-4. *Neuron*. 1997;18(3):505-

517. doi:10.1016/S0896-6273(00)81250-7

17. Riggs HE, McGrath JJ, Schwarz HP. Malformation of the adult brain (albino rat) resulting

from prenatal irradiation. *J Neuropathol Exp Neurol*. 1956;15(4):432-447.

doi:10.1097/00005072-195610000-00006

18. Sherman GF, Galaburda AM, Behan PO, Rosen GD. Neuroanatomical anomalies in

autoimmune mice. *Acta Neuropathol*. 1987;74(3):239-242. doi:10.1007/BF00688187

19. Amano S, Ihara N, Uemura S, et al. Development of a Novel Rat Mutant with

Spontaneous Limbic-Like Seizures. *Am J Pathol*. 1996;149(1):329–336.

20. Dvořàk K, Feit J, Juránková Z. Experimentally induced focal microgyria and status

verrucosus deformis in rats - Pathogenesis and interrelation histological and

autoradiographical study. *Acta Neuropathol*. 1978;44(2):121-129.

doi:10.1007/BF00691477

21. Ferrer I, Alcántara S, Catalá I, Zújar MJ. Experimentally induced laminar necrosis, status

verrucosus, focal cortical dysplasia reminiscent of microgyria, and porencephaly in the

rat. *Exp brain Res*. 1993;94(2):261-269. http://www.ncbi.nlm.nih.gov/pubmed/8359242.

Accessed July 14, 2017.

22. Lo Turco J, Manent JB, Sidiqi F. New and improved tools for in utero electroporation

studies of developing cerebral cortex. *Cereb Cortex*. 2009;19(Suppl. 1):i120-5.

doi:10.1093/cercor/bhp033

23. Saito T, Nakatsuji N. Efficient Gene Transfer into the Embryonic Mouse Brain Using in

Vivo Electroporation. *Dev Biol*. 2001;240(1):237-246. doi:10.1006/dbio.2001.0439

24.   Tabata H, Nakajima K. Efficient in utero gene transfer system to the developing mouse

594   brain using electroporation: Visualization of neuronal migration in the developing cortex.

595   *Neuroscience*. 2001. doi:10.1016/S0306-4522(01)00016-1

596   25.   Schain AJ, Hill RA, Grutzendler J. Label-free in vivo imaging of myelinated axons in

597   health and disease with spectral confocal reflectance microscopy. *Nat Med*.

598   2014;20(4):443-449. doi:10.1038/nm.3495

599   26.   Hill RA, Li AM, Grutzendler J. Lifelong cortical myelin plasticity and age-related

600   degeneration in the live mammalian brain. *Nat Neurosci*. 2018;21(May):1-13.

601   doi:10.1038/s41593-018-0120-6

602   27.   Hill RA, Grutzendler J. In vivo imaging of oligodendrocytes with sulforhodamine 101. *Nat

603   Methods*. 2014;11(11):1081-1082. doi:10.1038/nmeth.3140

604   28.   Jacob H. Die feinere Oberflächengestaltung der Hirnwindungen, die Hirnwarzenbildung

605   und die Mikropolygyrie. *Zeitschrift für die gesamte Neurol und Psychiatr*. 1940;170:68-84.

606   doi:10.1007/bf02869355

607   29.   Morel F, Wildi E. Dysgénésie nodulaire disséminée de l'écorce frontale. *Rev Neurol

608   (Paris)*. 1952;87(3):251-270.

609   30.   Sherman GF, Stone JS, Press DM, Rosen GD, Galaburda AM. Abnormal architecture

610   and connections disclosed by neurofilament staining in the cerebral cortex of autoimmune

611   mice. *Brain Res*. 1990;529(1-2):202-207. doi:10.1016/0006-8993(90)90828-Y

612   31.   Ramos RL, Smith PT, DeCola C, Tam D, Corzo O, Brumberg JC. Cytoarchitecture and

613   Transcriptional Profiles of Neocortical Malformations in Inbred Mice. *Cereb Cortex*.

614   2008;18(11):2614-2628. doi:10.1093/cercor/bhn019

32.   Sherman GF, Galaburda AM, Geschwind N. Cortical anomalies in brains of New Zealand

mice: a neuropathologic model of dyslexia? *Proc Natl Acad Sci U S A*. 1985;82(23):8072-

8074. doi:10.1073/pnas.82.23.8072

33. Ramos RL, Siu NY, Brunken WJ, et al. Cellular and axonal constituents of neocortical

molecular layer heterotopia. *Dev Neurosci*. 2014;36(6):477-489. doi:10.1159/000365100

34. Er JC, Leong C, Teoh CL, et al. NeuO: a Fluorescent Chemical Probe for Live Neuron

Labeling. *Angew Chemie Int Ed*. 2015;54(8):2442-2446. doi:10.1002/anie.201408614

35. Humphreys P, Kaufmann WE, Galaburda AM. Developmental dyslexia in women:

neuropathological findings in three patients. *Ann Neurol*. 1990;28(6):727-738.

doi:10.1002/ana.410280602

36. Galaburda AM, Sherman GF, Rosen GD, Aboitiz F, Geschwind N. Developmental

dyslexia: Four consecutive patients with cortical anomalies. *Ann Neurol*. 1985;18(2):222-

233. doi:10.1002/ana.410180210

37. Molyneaux BJ, Arlotta P, Fame RM, MacDonald JL, MacQuarrie KL, Macklis JD. Novel

subtype-specific genes identify distinct subpopulations of callosal projection neurons. *J

Neurosci*. 2009;29(39):12343-12354. doi:10.1523/jneurosci.6108-08.2009

38. Molyneaux BJ, Goff LA, Brettler AC, et al. DeCoN: Genome-wide Analysis of In Vivo

Transcriptional Dynamics during Pyramidal Neuron Fate Selection in Neocortex. *Neuron*.

2015;85(2):275-288. doi:https://doi.org/10.1016/j.neuron.2014.12.024

39. Sorensen SA, Bernard A, Menon V, et al. Correlated Gene Expression and Target

Specificity Demonstrate Excitatory Projection Neuron Diversity. *Cereb Cortex*.

2015;25(2):433-449. doi:10.1093/cercor/bht243

40. Gabel LA, LoTurco JJ. Electrophysiological and morphological characterization of

neurons within neocortical ectopias. *J Neurophysiol*. 2001;85(2):495-505.

doi:10.1152/jn.2001.85.2.495

41.   Gabel LA. Layer I neocortical ectopia: Cellular organization and local cortical circuitry.

Brain Res. 2011;1381:148-158. doi:10.1016/J.BRAINRES.2011.01.040

42.   Angevine JB, Sidman RL. Autoradiographic Study of Cell Migration during Histogenesis

of Cerebral Cortex in the Mouse. *Nature*. 1961;192(4804):766-768.

doi:10.1038/192766b0

43.   Raedler E, Raedler A. Autoradiographic study of early neurogenesis in rat neocortex.

Anat Embryol (Berl). 1978;154(3):267-284. doi:10.1007/bf00345657

44.   Hill RA, Damisah EC, Chen F, Kwan AC, Grutzendler J. Targeted two-photon chemical

apoptotic ablation of defined cell types in vivo. *Nat Commun*. 2017;8(May):1-15.

doi:10.1038/ncomms15837

45.   Singh SC. Ectopic neurones in the hippocampus of the postnatal rat exposed to

methylazoxymethanol during foetal development. *Acta Neuropathol*. 1977;40(2):111-116.

doi:10.1007/bf00688698

46.   Rosen GD, Bai J, Wang Y, et al. Disruption of Neuronal Migration by RNAi of Dyx1c1

Results in Neocortical and Hippocampal Malformations. *Cereb Cortex*. 2007;17(11):2562-

2572. doi:10.1093/cercor/bhl162

47.   Bagnard D, Lohrum M, Uziel D, Püschel AW, Bolz J. Semaphorins act as attractive and

repulsive guidance signals during the development of cortical projections. *Development*.

1998;125(24):5043-5053. http://www.ncbi.nlm.nih.gov/pubmed/9811588.

48.   Richards LJ, Koester SE, Tuttle R, O'Leary DD. Directed growth of early cortical axons is

influenced by a chemoattractant released from an intermediate target. *J Neurosci*.

1997;17(7):2445-2458. http://www.ncbi.nlm.nih.gov/pubmed/9065505.

49. Métin C, Deléglise D, Serafini T, Kennedy TE, Tessier-Lavigne M. A role for netrin-1 in
the guidance of cortical efferents. *Development*. 1997;124(24):5063-5074.
https://www.scopus.com/inward/record.uri?eid=2-s2.0-
0031441084&partnerID=40&md5=16c8fc81c15621923b5da3fd9e86e059. Accessed
September 23, 2019.

50. Polleux F, Giger RJ, Ginty DD, Kolodkin AL, Ghosh A. Patterning of cortical efferent
projections by semaphorin-neuropilin interactions. *Science (80-)*. 1998;282(5395):1904-
1906. doi:10.1126/science.282.5395.1904

51. Gao PP, Yue Y, Zhang JH, Cerretti DP, Levitt P, Zhou R. Regulation of thalamic neurite
outgrowth by the Eph ligand ephrin-A5: implications in the development of thalamocortical
projections. *Proc Natl Acad Sci U S A*. 1998;95(9):5329-5334.
doi:10.1073/pnas.95.9.5329

52. Baudet M-L, Zivraj KH, Abreu-Goodger C, et al. miR-124 acts through CoREST to control
onset of Sema3A sensitivity in navigating retinal growth cones. *Nat Neurosci*.
2012;15(1):29-38. doi:10.1038/nn.2979

53. Shewan D, Dwivedy A, Anderson R, Holt CE. Age-related changes underlie switch in
netrin-1 responsiveness as growth cones advance along visual pathway. *Nat Neurosci*.
2002;5(10):955-962. doi:10.1038/nn919

54. Skaliora I, Singer W, Betz H, Püschel AW. Differential patterns of semaphorin expression
in the developing rat brain. *Eur J Neurosci*. 1998;10(4):1215-1229. doi:10.1046/j.1460-
9568.1998.00128.x

55. Sturrock RR. Myelination of the mouse corpus callosum. *Neuropathol Appl Neurobiol*.
1980;6(6):415-420. doi:10.1111/j.1365-2990.1980.tb00219.x

56.   Goebbels S, Wieser GL, Pieper A, et al. A neuronal PI(3,4,5)P3-dependent program of

oligodendrocyte precursor recruitment and myelination. *Nat Neurosci*. 20(1):10-15.

doi:10.1038/nn.4425

57.   Tomassy GS, Berger DR, Chen H-HH, et al. Distinct Profiles of Myelin Distribution Along

Single Axons of Pyramidal Neurons in the Neocortex. *Science (80-)*.

2014;344(6181):319-324. doi:10.1126/science.1249766

58.   Micheva KD, Wolman D, Mensh BD, et al. A large fraction of neocortical myelin

ensheathes axons of local inhibitory neurons. *Elife*. 2016;5. doi:10.7554/eLife.15784

59.   Ishii K, Kubo KI, Endo T, et al. Neuronal heterotopias affect the activities of distant brain

areas and lead to behavioral deficits. *J Neurosci*. 2015. doi:10.1523/JNEUROSCI.3648-

14.2015

60.   Roper SN, Gilmore RL, Houser CR. Experimentally induced disorders of neuronal

migration produce an increased propensity for electrographic seizures in rats. *Epilepsy*

Res. 1995;21(3):205-219. doi:10.1016/0920-1211(95)00027-8

61.   Nieto M, Monuki ES, Tang H, et al. Expression of Cux-1 and Cux-2 in the subventricular

zone and upper layers II-IV of the cerebral cortex. *J Comp Neurol*. 2004;479(2):168-180.

doi:10.1002/cne.20322

62.   Chmielowska J, Stewart MG, Bourne RC. gamma-Aminobutyric acid (GABA)

immunoreactivity in mouse and rat first somatosensory (SI) cortex: description and

comparison. *Brain Res*. 1988;439(1-2):155-168. doi:10.1016/0006-8993(88)91472-2

63.   Meyer HS, Schwarz D, Wimmer VC, et al. Inhibitory interneurons in a cortical column

form hot zones of inhibition in layers 2 and 5A. *Proc Natl Acad Sci*. 2011;108(40):16807-

16812. doi:10.1073/pnas.1113648108

64.   Gabel LA, LoTurco JJ. Layer I Ectopias and Increased Excitability in Murine Neocortex. *J*
Neurophysiol. 2002;87(5):2471-2479. doi:10.1152/jn.2002.87.5.2471

65.   Manent JB, Wang Y, Chang Y, Paramasivam M, LoTurco JJ. Dcx reexpression reduces
subcortical band heterotopia and seizure threshold in an animal model of neuronal
migration disorder. *Nat Med*. 2009;15(1):84-90. doi:10.1038/nm.1897

66.   Chevassus-Au-Louis N, Rafiki A, Jorquera I, Ben-Ari Y, Represa A. Neocortex in the
hippocampus: An anatomical and functional study of CA1 heterotopias after prenatal
treatment with methylazoxymethanol in rats. *J Comp Neurol*. 1998;394(4):520-536.
doi:10.1002/(SICI)1096-9861(19980518)394:4<520::AID-CNE9>3.0.CO;2-3

67.   Jenner AR, Galaburda AM, Sherman GF. Connectivity of ectopic neurons in the
molecular layer of the somatosensory cortex in autoimmune mice. 2000;10(10):1005-
1013.

68.   Chevassus-Au-Louis N, Congar P, Represa A, Ben-Ari Y, Gaïarsa JL. Neuronal migration
disorders: heterotopic neocortical neurons in CA1 provide a bridge between the
hippocampus and the neocortex. *Proc Natl Acad Sci U S A*. 1998;95(17):10263-10268.
doi:10.1073/PNAS.95.17.10263

69.   Schottler F, Fabiato H, Leland JM, et al. Normotopic and heterotopic cortical
representations of mystacial vibrissae in rats with subcortical band heterotopia.
Neuroscience. 2001. doi:10.1016/S0306-4522(01)00395-5

70.   Romero DM, Bahi-Buisson N, Francis F. Genetics and mechanisms leading to human
cortical malformations. *Semin Cell Dev Biol*. 2018;76:33-75.
doi:10.1016/J.SEMCDB.2017.09.031

71.   Saito T. In vivo electroporation in the embryonic mouse central nervous system. *Nat*

Protoc. 2006;1(3):1552-1558. doi:10.1038/nprot.2006.276

72.   Kim J-Y, Ash RT, Ceballos-Diaz C, et al. Viral transduction of the neonatal brain delivers

controllable genetic mosaicism for visualising and manipulating neuronal circuits in vivo.

Eur J Neurosci. 2013;37(8):1203-1220. doi:10.1111/ejn.12126

73.   Phifer CB, Terry LM. Use of hypothermia for general anesthesia in preweanling rodents.

Physiol Behav. 1986;38(6):887-890.

74.   Patel TP, Man K, Firestein BL, Meaney DF. Automated quantification of neuronal

networks and single-cell calcium dynamics using calcium imaging. *J Neurosci Methods*.

2015;243:26-38. doi:10.1016/j.jneumeth.2015.01.020

75.   Li X, Ouyang G, Usami A, Ikegaya Y, Sik A. Scale-Free Topology of the CA3

Hippocampal Network: A Novel Method to Analyze Functional Neuronal Assemblies.

Biophys J. 2010;98(9):1733-1741. doi:10.1016/J.BPJ.2010.01.013

76.   Li X, Cui D, Jiruska P, Fox JE, Yao X, Jefferys JGR. Synchronization Measurement of

Multiple Neuronal Populations. *J Neurophysiol*. 2007;98(6):3341-3348.

doi:10.1152/jn.00977.2007

ACKNOWLEDGEMENTS

We thank F. Chen for his guidance *in utero* electroporation techniques. This study was funded by grants from the U.S. National Institutes of Health (R01NS089734, R21NS088411 and R21NS087511 to J.G.; R00NS099469 and P20GM113132 to R.A.H.; T32NS007224 and T32GM007205 to A.M.L.), Donors Cure Foundation (New Vision Award to R.A.H.), and National Multiple Sclerosis Society (#RR-1602-07686 to J.G.).

AUTHOR CONTRIBUTIONS

A.M.L. and J.G. conceived the initial project. A.M.L., R.A.H, and J.G. designed the experiments. A.M.L. performed all experiments, analyzed the data, and prepared the figures. A.M.L., R.A.H. and J.G. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

MATERIALS & CORRESPONDENCE

Correspondence and requests for materials should be addressed to J.G. (email: jaime.grutzendler@yale.edu) and R.A.H. (email: robert.hill@dartmouth.edu).

Fig. 1. Targeted induction and high-resolution imaging of layer I heterotopia in the live

mouse. a, Diagram showing a single neocortical heterotopion induced at the needle tract

during *in utero* electroporation (IUE) at embryonic day (E15). The site is relocated postnatally for

detailed investigation of the resulting layer I heterotopion by intravital imaging. **b**, Quantification

showing significantly increased neuron density (n = 6 mice) within heterotopia compared to

ipsilateral neighboring layer I control regions (Wilcoxon non-parametric, matched-pairs signed

rank test *P < 0.05). Each point represents the heterotopion or control region from one animal.

Horizontal lines and error bars denote mean and s.e.m, respectively. Descriptive statistics are

indicated in Supplementary File 1. **c**, *In vivo* fluorescence images of a representative layer

I heterotopion (right) and neighboring ipsilateral, non-injected control region in a P30 mouse (left).

Notice the tightly packed cluster of fluorescently labelled neuronal cell bodies (NeuO; green).

Neurons within the heterotopion can also be readily labeled during the electroporation procedure

and visualized *in vivo* (tdTomato; red). In contrast, neurons in more sparsely populated control

layer I regions never demonstrate IUE-mediated labelling (left column). HTP, heterotopia. Scale

bars, 50 µm (**c**).

Fig. 2. Heterotopia contain an abundance of hypermyelinated axons. a, *In vivo* combined

fluorescence and label-free SCoRe myelin images of a representative induced heterotopion (right;

same as in Figure 1c) and neighboring ipsilateral, non-injected control region (left) in a P30

mouse. The tightly packed heterotopic neuronal cell bodies (NeuO) are surrounded by a

circumscribed abundance of aberrantly projecting, myelinated axon segments (SCoRe). **b,c** Low

magnification (**b**) and high magnification (**c**) immunostainings of an induced heterotopion and its

corresponding contralateral control region taken from a P30 mouse, showing increased

oligodendrocyte CNPase expression associated with the heterotopion, as defined by the

ectopically positioned NeuN$^+$ neuronal cell bodies in layer I (**b**, white arrowheads). Notice the

more horizontal orientation of myelin segments located closer to the pial surface of the

heterotopion (**c**, blue arrowhead). Deeper myelin segments, in contrast, are organized in a more

radial fashion (**c**, blue arrow) and fasciculate into densely myelinated fiber bundles that project

through lower cortical layers (**b**, white arrow). **d**, Quantifications of SCoRe *in vivo* (n = 8 mice;

top) and CNPase in fixed tissue (n = 6 mice; middle and bottom), showing increased myelination

and oligodendrocyte cell body densities in induced layer I heterotopia compared to non-injected

contralateral control regions. Each point represents the layer I heterotopion or control region from

one animal. Horizontal lines and error bars denote mean and s.e.m, respectively (Wilcoxon

matched-pairs signed rank non-parametric test; *P < 0.05, **P < 0.01). HTP, Heterotopia.

Descriptive statistics are given in Supplementary File 1. Scale bars, 50 µm (**a**), 200 µm in (**b**), and

50 µm (**c**).

Fig. 3. Myelinated and unmyelinated axons follow aberrantly looping and concentric paths.

a-d, Intravital imaging of a nest-like heterotopion at P30 in (**a**) using SCoRe and confocal

fluorescence microscopy showing **b**, labelled neuronal cell bodies (tdTomato; green) dispersed

throughout the aberrantly oriented myelinated fibers (SCoRe; magenta) and **c**, winding axonal

projections that occasionally fasciculate into bundles at the edges of the heterotopion. **d**,

Myelinated (arrowheads) and unmyelinated axon segments (arrows) are observed within the

heterotopion. **e-g**, Low magnification (**e**) and high magnification (**f**, **g**) immunostainings of an

embryonically-induced heterotopion, confirming the presence of both myelinated (**g**, arrowhead)

and unmyelinated (**f**, arrow) axon segments (**f** and **g** show high magnification images of areas

indicated by arrow and arrowhead in **e**, respectively). NF-H, neurofilament heavy chain. Images

are representative of experiments performed in at least six animals. Scale bars, 100 μm (**a**), 15

μm (**b-d**), 100 μm (**e**), and 25 μm (**f**, **g**).

Fig. 4. Heterotopia can be induced at various embryonic stages of corticogenesis. a, *In vivo*

NeuO dye neuron staining (green) and SCoRe myelin imaging (magenta) of P30 mouse cortices

that were previously electroporated at E14 (left column), E15 (middle column), and E17 (right

column), all showing similar aberrantly projecting axons and hypermyelination encircling ectopic

neuron clusters. Images are representative of observations made in at least three animals per

IUE-injected age group. Scale bars, 100 μm.

Fig. 5. Postnatally-induced heterotopia do not develop aberrant axons or myelination. a,

Confocal images of an immunolabelled coronal section from P30 mouse cortex, showing no

changes in myelination (CNPase) or axonal pathfinding (NF-H) associated with a P0-induced

heterotopion (NeuN; arrowheads), compared to its contralateral control region (left). **b**, In contrast,

neuronal heterotopia induced *in utero* at E15 (NeuN; arrowheads) show dramatic changes in local

layer I myelin (CNPase) expression and patterning. **c**, Quantifications of CNPase expression,

showing no significant differences in the myelin or oligodendrocyte cell body densities ($n = 6$

animals) between postnatally-induced heterotopia and control regions. Each point represents the

P0-induced layer I heterotopion or control region from one animal, with horizontal lines and error

bars denoting mean and s.e.m, respectively. Wilcoxon matched-pairs signed rank non-parametric

test was used to determine significance; NS, no significance. NF-H, neurofilament heavy chain.

Descriptive statistics are given in Supplementary File 1. Scale bars, 100 µm (**a**, **b**).

Fig. 6. Astrocytes and microglia retain a normal density despite aberrant axonal and myelin

distribution. a-b, Low magnification (**a**) and high magnification (**b**) images of Aldh1l1-positive

astrocytes in P30 mouse cortex, showing similar cell densities in layer I heterotopia (right images,

arrowheads) compared to corresponding contralateral control regions (left images). **c**,

Quantifications using P30 mouse forebrain tissue showing no significant difference in astrocyte

cell density (Wilcoxon non-parametric matched-pairs, signed rank test, n = 6 animals; ns, no

significance). **d-e** Low magnification (**d**) and high magnification (**e**) images of Iba1-

immunolabelled P30 mouse forebrain, showing no difference in microglia cell densities between

heterotopia (right images, arrowheads) and contralateral control regions (left images). **f**,

Quantifications using P30 mouse forebrain tissue demonstrating no significant difference in

microglia cell density between layer I heterotopia and corresponding contralateral control regions

(Wilcoxon matched-pairs, signed rank non-parametric test, n = 6 animals; ns, no significance).

For graphs in (**c**) and (**f**), each point represents an embryonically-induced layer I heterotopion or

control region from one animal, with horizontal lines and accompanying error bars denoting mean

and s.e.m., respectively. Descriptive statistics are given in Supplementary File 1. HTP,

heterotopia. Scale bars, 100 µm (**a**), 20 µm (**b**), 100 µm (**d**), and 20 µm (**e**).

Fig. 7. Heterotopia contain a mixed population of excitatory and inhibitory neurons. a,

Approach for quantifying layer-specific neuronal marker expression (symbolized by blue and

green dots) in layer I heterotopia and across corresponding contralateral control cortices in P30

mice. Contralateral control cortices are subdivided into 11 equally-sized bins that together span

the entire thickness of the cortex. **b-c** Both Cux1$^+$ (**b**, cyan) and Tle4$^+$ (**c**, green) neuronal cell

bodies occur within layer I heterotopia (right images, arrowheads). In contralateral control regions

(left images), Cux1 is concentrated in more superficial cortical areas corresponding to layers II/III,

whereas deeper areas corresponding to layers V/VI encapsulate the majority of Tle4 labelling.

Quantifications (far right) show the percentages of NeuN$^+$ cell bodies that also express Cux1 (top)

or Tle4 (bottom) within heterotopia and across corresponding contralateral control bins. Each dot

corresponds to the layer I heterotopion or control bin of a single animal, with all dots of the same

color belonging to the same animal (*n* = 4 animals for Cux1; *n* = 5 animals for Tle4). The black

line denotes the mean. WM, white matter. **d,e** Low (**d**) and high (**e**) magnification z-projections of

GAD-67-immunostained P30 mouse forebrain, showing the presence of GAD-67-labelled cells

and puncta in both layer I heterotopia (right images, arrowheads) and corresponding contralateral

control regions (left images). **f**, Quantifications from P30 mouse tissue showing no significant

difference in GAD-67$^+$ neuronal cell density between layer I heterotopia and contralateral control

regions (Wilcoxon matched-pairs, signed rank non-parametric test, *n* = 6 animals; ns, no

significance). Each point represents the layer I heterotopion or contralateral control region from

one animal, with horizontal lines and accompanying error bars denoting mean and s.e.m.,

respectively. HTP, heterotopia. Descriptive statistics are given in Supplementary File 1. Scale

bars 200 µm (**b**, **c**), 100 µm (**d**), and 10 µm (**e**).

Fig. 8. Heterotopic neurons display calcium dynamics similar to adjacent layer II/III

neurons and respond to sensorimotor input. a-c *In vivo* time lapse imaging of layer I

heterotopic neurons expressing the calcium sensor GCaMP6f in an awake, head-fixed mouse.

Dotted lines in (**a**) indicate an analyzed heterotopic region that is displayed in (**b**). **c**, Example

traces of spontaneous neuronal calcium transients from layer I heterotopic (right) and ipsilateral

layer II/III non-heterotopic (left) regions displayed in (**b**). **d**, Quantifications showing no significant

difference in calcium spike event frequency (Hz), variance (s.d.) or global synchronization index

between layer I heterotopia and adjacent layer II/III non-heterotopic regions. Each dot

corresponds to the average value for each measure in a layer I heterotopion or non-heterotopic

layer II/III region from a single animal. Data represent 493 layer I heterotopic neurons and 594

layer II/III neurons from *n* = 7 mice. Wilcoxon matched-pairs, signed rank non-parametric test was

used for all three quantifications; ns, no significance; HTP, heterotopia. Descriptive statistics are

given in Supplementary File 1. **e**, *In vivo* z-projection of the same GCaMP6f-transfected

heterotopion displayed in (**a**), captured at a different depth, showing neurons that were imaged

during brief applications of whisker stimulation. **f**, Baseline calcium dynamics (left column) and

calcium fluctuations (middle and right columns) of individual heterotopic neurons in response to

stimuli (denoted by solid blue line in two trials) obtained from the awake, head-fixed mouse

imaged in (**e**). Experimental observations in (**e**, **f**) were replicated in three mice. All data and

images from (**a**-**f**) were obtained from P50-60 mice with heterotopia induced at E15, and virally

transfected with AAV-GCaMP6f at P21-P30. Some neuronal cell body and axonal labelling by

pCAG-TdTomato (red) is the result of E15 IUE-mediated transfection. Scale bars, 50 μm (**a**), 25

μm (**b**), and 50 μm (**e**).

Figure 3–figure supplement 1. Myelinated axons project aberrantly around heterotopic

cells. a, High resolution immunostaining showing the winding trajectories followed by CNPase[+]

fibers (bottom, blue arrow) within a layer I heterotopion (NeuN; white arrowheads) in a P30 mouse

coronal section. The aberrantly-projecting myelinated fiber bundles are absent from immediately

adjacent non-heterotopic cortex and corresponding contralateral control regions (top row).

Images are representative of experiments performed in at least six animals. Scale bars, 100 μm.

Figure 2–figure supplement 1. Disrupted myelin patterning and axonal pathfinding occurs

in deeper cortical layers. a, Myelinated fibers (MBP, cyan) accumulate in thick radially oriented

cords (white arrow) beneath a layer I heterotopion (NeuN; arrowheads) in an immunostained P30

mouse forebrain coronal section. **b-c,** Confocal images of a different coronal section from the

same heterotopion in (**a**), revealing similar aberrant myelin and axon accumulations as identified

by oligodendrocyte CNPase (green) and axonal NF-H (red) immunostaining, respectively. A

higher magnification view of the densely packed myelinated axons fasciculating underneath

the heterotopion (**b**, blue arrow) is shown in (**c**). Images in (**b**) and (**c**) are of same section shown

in Figure 3e. Images are representative of fixed tissue experiments performed in at least three

mice. Scale bars, 200 µm (**a**, **b**), and 20 µm (**c**).

Figure 7–figure supplement 1. Heterotopic neurons express both deep and superficial

cortical layer markers. a-b, Additional fixed tissue images captured from the heterotopia

displayed in Figures 7b and c, confirming the presence of $Cux1^+$ (**a**, cyan) and $Tle4^+$ (**b**, green)

neuronal cell bodies within the representative heterotopia (bottom images, arrowheads). In

corresponding contralateral control regions (top images), Cux1 (left) largely concentrates in

superficial cortical layers whereas Tle4 (right) tends to localize toward deeper cortical layers. **c-**

d, Quantifications showing the densities of $Cux1^+$ (**c**, $n = 4$ animals), $Tle4^+$ (**d**, $n = 5$ animals), and

$NeuN^+$ (**c** and **d**, right) cells in heterotopia and across corresponding contralateral control regions

of P30 mice. Each dot corresponds to the layer I heterotopion or control bin of a single animal,

with all dots of the same color belonging to the same animal. The black line denotes the mean.

Images are representative of experiments performed in at least four animals. WM, white matter;

HTP, heterotopia. Scale bars, 200 µm.

Figure 7–figure supplement 2. Birth-dating of heterotopic neurons. a, Diagram of the pulse

labelling strategy for birthdating heterotopic cortical neurons. Dividing cells are labelled by a single

EdU injection at E11.5, with heterotopia induced at E15. Mice are sacrificed at P30 for analysis.

b, Low magnification images showing the EdU$^+$ neurons incorporated in a layer I heterotopion

(right, arrowheads) of a P30 mouse. In corresponding contralateral control cortex (left column),

EdU expression is predominantly concentrated in deeper cortical layers. **c**, Quantifications

showing the percentages of NeuN$^+$ cells that also express EdU (top), EdU$^+$NeuN$^+$ cell body

densities (middle), and NeuN$^+$ cell body densities (bottom) in layer I heterotopia and across

corresponding contralateral control cortices of P30 mice. Each dot corresponds to the layer I

heterotopion or control bin of a single animal, with all dots of the same color belonging to the

same animal (n = 4 animals total). Images are representative of the observations made in each

mouse brain. The black line denotes the mean. WM, white matter; HTP, heterotopia. Scale bars,

200 µm.

Video 1. *In vivo* imaging of a layer I heterotopion and adjacent control area. Confocal z-

stacks of a heterotopion (right) and ipsilateral non-injected layer I control region (left) captured

from a living P30 mouse using label-free SCoRe (magenta) and fluorescence microscopy,

showing the aberrantly projecting myelinated axons and ectopic neurons located inside a layer I

heterotopion induced at E15. Neuronal cell bodies and cerebral blood vessels are visualized using

fluorescent neuronal dye NeuO (green) and Evans Blue (white), respectively. The density of the

horizontally crisscrossing, aberrantly projecting fibers diminishes with increasing cortical depth

(indicated at bottom left).

Video 2. Layer I heterotopia and adjacent layer II/III regions display highly variable

spontaneous neuronal calcium transients. Representative *in vivo* two-photon time-lapse

recordings of GCaMP6f-labelled neurons (green) within a layer I heterotopion (right) and an

adjacent layer II/III cortical region (left) in an awake, head-fixed P55 mouse. Diverse patterns of

spontaneous calcium transients are observed in neuronal cell bodies throughout both imaged

regions. Note the absence of obvious epileptiform activity. Images were acquired at 2 Hz from an

E15-induced layer I heterotopion. Some neuronal cell body and axonal labelling by pCAG-

TdTomato (red) is the result of E15 IUE-mediated transfection.

Video 3. Heterotopic neurons respond to sensory stimuli. Two *in vivo* confocal time-lapse

recording examples of GCaMP6f-labelled neurons within a layer I heterotopion before, during,

and after the application of a brief whisker stimulus, administered one minute into each imaging

session. The timing of the stimulus and accompanying movements are indicated by the white

circle (positioned at the top left of the embedded video). Neuronal calcium changes are observed

upon the application of each stimulus, indicating the responsiveness of the heterotopic neurons

to sensorimotor input. Both recordings were obtained from P55 awake, head-fixed mice with layer

I heterotopia that were induced at E15 and virally transfected using AAV-GCaMP6f at P21-P30.

Supplementary Table 1. Details for all statistical analyses, including sample sizes and p-values.

Figure 1

a

E15 IN UTERO ELECTROPORATION

Step 1
Insertion of needle into cortex

Step 2
Plasmid DNA delivery into lateral ventricle

Step 3
Electroporation

P30 IN VIVO IMAGING

Live imaging of electroporated cortex

Induced HTP at needle tract

b

c

CONTROL　　　　**HETEROTOPIA**

tdTomato NeuO

Figure 2

a CONTROL HETEROTOPIA NeuO SCoRe Evans Blue

b NeuN CNPase DAPI
CONTROL HETEROTOPIA

c CONTROL HETEROTOPIA

d

SCoRe density (a.u.)
Control HTP Edge HTP Center
** **

CNPase density (a.u.)
Control HTP
*

CNPase⁺ cells (10³/mm³)
Control HTP
*

Figure 2 - figure supplement 1

NeuN MBP DAPI   NeuN CNPase NF-H DAPI

Figure 3

Figure 3 - figure supplement 1

a

CONTROL   NeuN CNPase DAPI

HETEROTOPIA

a

SCoRe NeuO

E14 IUE → P30   E15 IUE → P30   E17 IUE → P30

Figure 5

NeuN CNPase NF-H DAPI

a Control | P0 injection | P30

b E15 injection | P30

c
CNPase density (a.u.)
ns
Control | P0 injection

CNpase+ cells (10³/mm³)
ns
Control | P0 injection

Figure 6

Figure 7

Figure 7–figure supplement 1

Figure 8

Supplementary Table 1					
Figure Number	**Statistical Test**	**Sample Size**	**Sample Definition**	**P Value**	**Degrees of Freedom and F/T/z/R/ETC value**
1b	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.0313 for NeuO, HTP vs. Control	n/a
2d	Wilcoxon non-parametric matched-pairs signed rank test	8	number of mice (paired samples)	SCoRe density, p=0.0078 for HTP edge vs. Control; p=0.0078 for HTP center vs. Control	n/a
	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.0313 for CNPase density, HTP vs. Control.	n/a
	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.0313 for CNPase$^+$ cell bodies, HTP vs. Control.	n/a
5c	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p > 0.9999 for CNPase$^+$ cell bodies, HTP vs. Control.	n/a
	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.0625 for CNPase density, HTP vs. Control.	n/a
6c	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.8438 for Aldh1L1$^+$ cell density, HTP vs. Control.	n/a
6f	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.9375 for Iba$^+$ cell density, HTP vs. Control	n/a
7f	Wilcoxon non-parametric matched-pairs signed rank test	6	number of mice (paired samples)	p = 0.5313 for GAD67$^+$ cell density, HTP vs Control	n/a
8d	Wilcoxon non-parametric matched-pairs signed rank test	7	number of mice (paired samples)	p = 0.6875 for spike event frequency, HTP vs Layers II/III	n/a
	Wilcoxon non-parametric matched-pairs signed rank test	7	number of mice (paired samples)	p = 0.9375 for variance (s.d.), HTP vs. Layers II/III	n/a
	Wilcoxon non-parametric matched-pairs signed rank test	7	number of mice (paired samples)	p = 0.2969 for global synchronization index, HTP vs. Layers II/III	n/a