From work stress to disease: A computational model

Remco Benthem de Grave, Fred Hasselman, & Erik Bijleveld

Radboud University

Remco Benthem de Grave, Behavioural Science Institute, Radboud University, The Netherlands; Fred Hasselman, Behavioural Science Institute, Radboud University, The Netherlands; Erik Bijleveld, Behavioural Science Institute, Radboud University, The Netherlands.

Remco Benthem de Grave is now at the Institute of Neuroscience, Newcastle University.

Erik Bijleveld was supported by grant 016-165-100 from the Netherlands Organisation for Scientific Research.

Correspondence concerning this article should be addressed to Remco Benthem de Grave, Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, NE2 4HH, United Kingdom. E-mail: r.benthemdegrave2@ncl.ac.uk

Abstract

In modern society, work stress is highly prevalent. Problematically, work stress can cause disease. To help understand the causal relationship between work stress and disease, we present a computational model of this relationship. That is, drawing from allostatic load theory, we captured the link between work stress and disease in a set of mathematical formulas. With simulation studies, we then examined our model's ability to reproduce key findings from previous empirical research. Specifically, results from Study 1 suggested that our model could accurately reproduce established findings on daily fluctuations in cortisol levels (both on the group level and the individual level). Results from Study 2 suggested that our model could accurately reproduce established findings on the relationship between work stress and cardiovascular disease. Finally, results from Study 3 yielded new predictions about the relationship between workweek configurations (i.e., how working hours are distributed over days) and the subsequent development of disease. Together, our studies suggest a new, computational approach to studying the causal link between work stress and disease. We suggest that this approach is fruitful, as it aids the development of falsifiable theory, and as it opens up new ways of generating predictions about why and when work stress is (un)healthy.

Keywords: work stress, occupational health, allostatic load theory, computational modeling, cortisol

From work stress to disease: A computational model

Work stress has been estimated to cost between $221 to $187 billion annually (Hassard et al., 2018). Considering these high societal costs, it is not surprising that scientists have thoroughly studied the origins, the nature, and the consequences of work stress. Broadly, work psychologists have examined how and when work stressors shape well-being and performance; biological psychologists have examined the nature of the physiological stress response; and, epidemiologists have examined how and when work stress may cause disease on the long run. Yet, despite the maturity of the science of work stress, this important area has a clear shortcoming: it does not yet have computational models that explain how work stress may cause disease.

Computational models can advance scientific knowledge in various ways. For example, they contribute to the transparency and falsifiability of theory, they facilitate the understanding of potential causal mechanisms, and they help generate new predictions (Epstein, 2008; Weinhardt, & Vancouver, 2012; Yarkoni, & Westfall, 2017). In what follows, drawing from *allostatic load theory* (McEwen, 1998), we propose a computational model of the putative causal effect of work stress on disease. In turn, we test whether our model can explain core findings from several previous empirical studies. Finally, we use our model to generate new predictions about the relationship between workweek configurations (i.e., how work hours are distributed over the working week) and the development of disease. Together, our research uses computational modelling to address a central problem in the domain of occupational health: how does work stress affect people's health?

A brief introduction to computational models

This section is intended for readers who are new to computational models. Readers who are familiar with computational modeling may skip this section.

We will start explaining what computational models are, by comparing them to *verbal theories* (with *theory* synonymous for *model*). By *verbal theories*, we refer to theories not described through mathematical equations (nor through some other structure of formal logic). Most theories in the social sciences are verbal theories. An example of a well-known verbal theory is *cognitive dissonance theory* (Festinger, 1962). Cognitive dissonance theory predicts that attitude change takes place when people's previously-held attitudes are inconsistent with their behavior. Another well-known verbal theory in social sciences is *social facilitation theory* (Allport, 1924; Zajonc, 1965), which predicts that, when in presence of others, people perform better on well-learned and simple tasks, but worse on new and complex tasks (Strauss, 2002). What these theories have in common is that they are provided in words, not in mathematical equations. As a result, they lack specificity. Little is clear about, say, the dynamics of how attitude change develops over time. Is the speed of change constant? Or does it increase first and then plateau? Or, how about the relationship between the presence of other people and performance? Is this relationship linear? What conditions are required for this relationship to hold? In order to perform any quantitative test of a verbal theory, e.g., using statistical analyses, researchers always need to make an interpretation of the theory first, leaving room for flexibility. Tests of verbal theory thus do not strictly test the theory—rather, they test the researcher's *interpretation* of the theory.

In physics and engineering, most (if not all) theories are computational. An example of a computational theory is Newton's theory of universal gravitation. This theory explains the force that works between two masses with the equation $= G \cdot \frac{m_1 \cdot m_2}{r^2}$, in which m_1 and m_2 are the two masses, r the distance between the two masses, G the universal gravitation constant, and F the resulting force. Each combination of values that are inserted for the parameters in the right side of the equation result in a single, exact, resulting force. More generally, for

researchers to test a computational theory, they do not need to make any additional interpretations.

Benefits of computational modeling. First, more so than verbal models, computational models are falsifiable. As mentioned, the limited specificity of verbal theory leaves room for various ways in which to define relationships between parameters. This flexibility makes it difficult to falsify verbal theories (Edwards, & Berry, 2010; Meehl, 1978), thus limiting what can be learned from a study that attempts to empirically validate a verbal theory. In contrast, computational models are fully specific. The model provides transparency about the included parameters and the assumed relationships between them, thus limiting interpretational freedom, facilitating falsifiability (Frankenhuis & Tiokhin, 2018; Smaldino, 2017; Weinhardt, & Vancouver, 2012).

Second, computational models can help the understanding of underlying causal mechanisms of the relationship that is being studied. Rather than attempting to create a model that exactly mimics reality, the goal of modeling is to provide a model that can satisfyingly approximate empirical observations, while maintaining simplicity as much as possible (Kokko, 2007; Epstein, 2008). As such, modeling can give insight in the dynamics that govern the process under investigation. Computational models may also increase understanding through analogies. In particular, processes that seem unrelated can have models that are computationally the same (e.g., the same model may explain the behavior of both arteries in the body, and the behavior of rubber bands that hold objects together). This way, researchers can borrow principles from well-developed theories from other scientific areas, accelerating scientific progress (Epstein, 2008).

Third, computational models can be used to perform simulations, which has several important advantages. Through simulations, researchers can explore mechanisms that can explain empirical findings (e.g., Nettle, Frankenhuis, & Rickard, 2013; Vancouver, Tamanini,

& Yoder, 2010); clarify inconsistencies among previous findings (e.g., Niv, Daw, Joel, &

Dayan, 2007; Vancouver, Li, Weinhardt, Steel, & Purl, 2016; Vancouver & Purl, 2017);

examine a model's robustness, i.e., examine the parameter ranges under which a model

explains existing empirical data (Epstein, 2008); and scrutinize the logic of intuitive reasoning

behind a theory, which is sometimes flawed (Farrell, & Lewandowsky, 2010). In other words,

simulations can examine whether a theory is viable to begin with. Furthermore, simulations of

computational models can also generate novel predictions of behavior that may be observed in

real populations (Epstein, 2008). In some cases, intuitive interpretation of a theory does not

lead to clear predictions. To provide an example from the present research: Does taking free

days on Wednesdays and Sundays—instead of Saturdays and Sundays—influence the risk of

developing disease in the long run? While it is not easy to use verbal models to formulate

hypotheses about particular situations such as these, simulations of computational models can

be used for this purpose.

The present research

With this research, we aim to meet three goals. Our first goal is to develop a first

computational model of the *work stress–disease* relationship. Rather than providing a detailed

model, including many possible parameters involved in the process of becoming diseased, we

aim to create a simple and compact model that focuses on the most important candidate

mechanisms. We chose to prioritize parsimony as we believe that it will benefit

interpretability of the model, and as it will limit the number of arbitrary assumptions that we

will need to make. Rather than formulating a new theory from scratch, we will be using

knowledge from leading, existing verbal theory (i.e., allostatic load theory) to develop our

model.

Our second goal is to investigate our model's ability to reproduce previously-reported

data, by using Monte Carlo simulations (i.e. simulations in which the value of a sample is

determined by its previous value and random sampling from a distribution). For these simulations, we will simulate individuals for whom we will verify that their collective behavior is comparable to the behavior that is described in large-scale empirical studies. This investigation will provide us with feedback about the credibility of the model that has been developed .

Our third goal is to formulate new predictions, again by using MCMC simulations. Specifically, we aim to formulate predictions about the impact of how working hours are distributed over the working week. When future research tests our novel predictions, this would provide a transparent means to verify or falsify our model—and, subsequently, to improve—our model.

In the next sections, we will lay out our research. In Study 1, we will perform simulations based on our model, to verify that our model can produce cortisol time courses similar to those reported in the literature. In Study 2, we will describe the relationship between cortisol levels and disease, completing our computational model. In both Studies 1 and 2, through simulations, we will examine if our simulated people exhibit the same relationship between stressors and disease, as was reported in previous large-scale studies. In Study 3, we will use the model to make novel predictions.

Study 1: From stress to cortisol

Allostatic Load Theory. We base our computational model, summarized in Figure 1, on allostatic load theory (McEwen, 1998). This influential theory describes the physiological processes that mediate the causal relationship between stress and disease. Allostatic load theory starts out from the assumption that stressful situations disrupt the stable resting state (*homeostasis*) of physiological systems, causing an alternative equilibrium (*allostasis*) in which physiological systems adaptively respond to deal with the stressful situation. Generally, allostasis involves increased activity of the sympathetic nervous system and the

hypothalamus–pituitary–adrenal axis (HPA axis). These activations generally cause an increase in heart rate, a release of nutrients into the bloodstream, a suppression of digestion, as well as a number of other changes (Sapolsky, Romero, & Munck, 2000; for an accessible introduction, see Sapolsky, 2004).

Although the ability to transition into allostasis is usually considered to be a healthy adaptation—i.e., it helps people to effectively deal with stressors—allostasis does put a burden on physiological systems. In many cases, this impact is fully reversible; it causes no permanent damage. For example, an artery under high pressure may readily return to its normal, resting state. However, when allostasis occurs too frequently (Type I), when allostasis is maintained for too long (Type II), or when physiological adaptation to allostasis is inadequate (Type III), lasting damage may occur. In this paper, we refer to allostasis' reversible burden on physiological systems as *allostatic strain*, and allostasis' lasting damage as *allostatic load*. As allostatic load accumulates, *disease* may emerge.

To develop a parsimonious computational model based on allostatic load theory, we have applied a simplification: we will consider only allostatic load due to allostasis occurring too frequently (i.e., Type I). To be able to test our computational model against previous empirical data, we needed to make an additional assumption: we assume that the concentration of the hormone cortisol—which is, indeed, frequently used as a direct measure of the human stress response (Dickerson, & Kemeny, 2004; Hellhammer, Wüst, & Kudielka, 2009)—provides an index of the current physiological response that is recruited to achieve allostasis.

Daily fluctuations in cortisol levels. The *HPA axis* regulates a chain of bodily events that leads to the production of cortisol (Sapolsky et al., 2000). The HPA axis consists of three

Figure 1. Schematic illustration of our computational model. In essence, the model explains how activation of the HPA axis, due to work stress and circadian inputs, elevates circulating cortisol levels. In turn, the cortisol response burdens the physiological system, in a fully reversible way (*allostatic strain*). Sustained allostatic strain, in turn, causes permanent damage (*allostatic load*), which ultimately causes *disease*. This chain of events is described in detail in the main text. HPA axis = Hypothalamus Pituitary Adrenal-axis.

hormonal glands: the *hypothalamus* and the *pituitary gland* (both located deep in the brain) and the *adrenal gland*s (located on the kidneys). These glands produce *corticotropin-releasing hormone* (CTH), *adrenocorticotropic hormone* (ACTH) and *cortisol*, respectively. These glands operate in a cascading manner. That is, neural impulses into the hypothalamus (e.g., due to stressors) cause the hypothalamus to release CTH, which causes the pituitary gland to release ACTH, which in turn causes the adrenal glands to release cortisol. However, all three glands have receptors for their own and each other's hormonal output. Thus, they together form an intricate dynamic system, which also involves the liver, which removes cortisol from the bloodstream, and then decomposes it (Hosseinichimeh, Rahmandad & Wittenborn, 2015).

Bloodstream cortisol concentrations follow a characteristic circadian pattern. Generally, blood cortisol steeply rises during the last few hours of the night, peaks about half an hour after awakening (we will refer to this peak as the *morning peak*), and then gradually decreases during the day (Wilhelm, Born, Kudielka, Schlotz, & Wüst, 2007). Importantly,

though, individual cortisol levels are very heterogeneous (Carroll et al., 2007). One key

source of variation is the strength of the response to neural impulses into the HPA axis, which

differs from person to person (Dickenson & Kemeny, 2004). Also, people substantially differ

in the rate with which the liver decomposes cortisol: cortisol half-life varies between 60 and

90 minutes (Carroll et al. 2007; Kovacs & Ojeda, 2011). Bloodstream cortisol concentrations

also respond to acute stressors. After people encounter a stressor, the cortisol level peaks after

about 30 minutes, after which cortisol levels decrease again (Dickenson & Kemeny, 2004;

Goodman, Janson, & Wolf, 2017).

It is likely that the amplitude of the morning peak and strength of cortisol responses to

acute stressors (during the day), are related. In particular, the morning peak may reflect the

anticipation of physiological and psychological demands for that day (Fries, Dettenborn, &

Kirschbaum, 2009; see also Brosschot, Pieper, & Thayer, 2005). In support of this idea,

research shows that people show higher morning peaks on weekdays, as compared to

weekend days (Schlotz, Hellhammer, Schulz, & Stone, 2004; Thorn, Hucklebridge, Evans, &

Clow, 2006). Similarly, a study among competitive dancers showed that dancers had a larger

morning peak on competition days, as compared to training days (Rohleder, Beulen, Chen,

Wolf, & Kirschbaum, 2007). So, it is plausible that the morning peak reflects the anticipation

of upcoming events, at least in part, causing a correlation between (a) the amplitude of the

morning peak and (b) the strength of cortisol responses to acute stressors. In our simulations,

we will consider this possibility. That is, we will run separate simulations assuming vs. not

assuming this correlation.

In a computational model. In this part of the model we describe the relationship

between neural impulses to the HPA axis, I, and the change in cortisol concentration, dC/dt.

We assume that all activity of the HPA axis is the result of neural impulses to the HPA axis,

which may either be the result of a circadian process taking place during the night (we will

further refer to these impulses as *night impulses*), or the result of work stressors (for consistency, further referred to as *work impulses*). We assume that each impulse has a binary intensity: impulses are either happening or not happening. Due to this assumption, impulses to the HPA axis, at any specific time point, can be described as countable number, $\sum I_t$. Thus, a higher number of impulses at the same time can be interpreted as a stronger impulse.

We simplify the dynamics of the HPA axis by assuming that it can be described by the following equation:

$$\frac{dC}{dt} = -\rho_C\, C_t + \kappa_{HPA} \sum(I_t - \tau) \tag{1}$$

In this equation, a rise in the cortisol concentration that is proportional, by $\kappa_{HPA,}$ to the neural impulses, $\sum I_t$, delayed by τ. The decay of cortisol is described by a linear relation between the current cortisol concentration, C_t, and a decay constant, ρ_C.

Simulations. To examine the ability of the model to reproduce the cortisol patterns that were previously reported (specifically in Born, Hansen, Marshall, Molle, & Fehm, 1999; Hosseinichimeh et al., 2015; Miller et al., 2016), we performed MCMC simulations with the model, with parameters as defined in Table 1 (code available as [online supplementary material](#)). As input to the model, a population of 10,000 people was simulated. Each simulated person was characterized by an individual cortisol half-life value. Specifically, for each person, a random draw from a normal distribution of cortisol half-life values determined this person's cortisol half-life, with a mean and standard deviation chosen such to reflect the variation of half-life values reported in previous literature (Carroll et al. 2007; Kovacs & Ojeda, 2011).

To mirror the time course of a cortisol response to an acute stressor as described in literature (Dickenson & Kemeny, 2004; Goodman, et al. 2017), impulses always lasted 30 minutes. Each simulated person was given a unique average number of daily *night impulses* and *work impulses*.

For night impulses, this unique *average number* of daily impulses was randomly drawn from a Gamma distribution (i.e. a continuous probability distribution for values that can only be positive, such as frequency values; Hazewinkel, 2002; see online supplementary material). In turn, this average number was used to randomly assign a number of impulses to each day. In particular, the number of night impulses that a person received on a particular day, was determined from a Poisson distribution (i.e., the discrete frequency distribution) with as expected frequency λ, the average number of daily night impulses for that person. For *work impulses*, we followed the same procedure.

In our simulation, we assumed work that impulses could occur at any moment within the working hours. Thus, work impulses were determined through random draws from a uniform distribution, ranging from work start to work end. Weekends were free of work impulses. We further assumed that night impulses should become exponentially more likely towards the end of the night, with the maximum probability at awakening. Thus, night impulses were randomly drawn from a negative exponential distribution (see online supplemental material for a plot of this exponential distribution).

To illustrate the simulation procedures laid out above, Figures 2 and 3 describe the outcomes of the procedures for four representative, simulated people. Figure 2 shows frequency distributions of night impulses and work impulses. Figure 3 shows how impulses were distributed over time during a single day.

Table 1

Parameter settings for simulating cortisol time courses in Study 1

Parameter	Value
Sampling frequency	2
People simulated	10'000
Days simulated per person	200
Night impulses, quantity	$X \sim \Gamma(k = 3, \theta = 14)$[a]
Night impulses, moment	$X \sim T_{wake} - \text{EXP}(\lambda = 1)$[a]
Work impulses, quantity	$X \sim \Gamma(k = 3, \theta = 6)$[a]
Work impulses, moment	$X \sim \mathcal{U}(a = T_{ws}, b = T_{we})$
Cortisol decay constant (ρ_C)	$X \sim \mathcal{N}(\mu = .52, \sigma^2 = .05^2)$
HPA axis scaling constant (κ_{HPA})	2.20
Cortisol response delay, τ	30 min
Wake time, T_{wake}	7AM
Work start, T_{ws}	8:30AM
Work end, T_{we}	4:30PM
Working days	Mon-Fri

a = Parameter values were determined through visual calibration on cortisol time courses published in Miller et al. (2016; see Figure 5a). Specifically, we adjusted these parameter values until simulations yielded a cortisol time course similar to the one reported in Miller et al. (2016).

As introduced above, we ran separate simulations in which we assumed no correlation between night impulses and work impulses (Scenario I), and in which we assumed a correlation between night impulses and work impulses (Scenario II; see Fries et al., 2009). In Scenario I, we drew average night impulses and average work impulses independently from a Gamma distribution. In Scenario II, we first drew the average number of work impulses from a Gamma distribution. Then, to define the average number of night impulses, we summed the average number of work impulses with a random value from a gamma distribution with shape parameter $k = 1$ and scale parameter $\theta = 8$. This procedure ensured (a) that the distribution for the average number of night impulses was visually similar across Scenarios I and II, and that (b) there was a correlation between average night impulses and average work impulses only in Scenario II (illustrated in Figure 4).

Simulation results. Figures 5–7 show comparisons of cortisol time courses from the simulations with cortisol time courses from findings from the published literature. For each person, only the first day of the simulation was used to create these plots. Figure 5 and 6 only show results only for Scenario I, as results for Scenario II were extremely similar (see online supplemental material).

First, we examined our model's ability to reproduce group-level cortisol time courses during waking time. To this end, we used data from a meta-dataset from Miller et al. (2016) as a starting point. This meta-dataset combined 15 previous field studies (total N ≈ 19,000), in order to obtain reference ranges for salivary cortisol levels in humans. Data from this previous study are replotted in Figure 5a; results from our simulations are plotted in Figure 5b.

Figure 2. The count of days (y-axis) that a simulated person received a specific number of impulses (x-axis).

Four random, representative, simulated people are shown. Note: the total amount of simulated days is 200.

Figure 3. Distribution of all impulses over two consecutive days (48 hours). Two representative days are shown

for each of four random, representative, simulated people. The plots illustrate that the number of impulses varies

from day to day. This is because the number of impulses in a given day results from a random draw from a

Poisson distribution, based on the person's overall mean (see main text).

Figure 4. Correlations between average daily work impulses and average daily night impulses in Scenario I,

uncorrelated and Scenario II, correlated.

Figure 5. a) Empirical data. Cortisol time courses based on data from ~19,000 people, replotted with permission

from Miller et al. (2016). b) Simulation results. Note: in both panels, data are shown until 16 hours after

awakening.

Figure 6. a) Empirical data. Night-time cortisol time courses (N = 15; error bars reflect 95% confidence interval

around the mean), replotted with permission from Clow et al. (2009). b) Simulation results. Note: in (b), the 95%

confidence interval around the average is too small to be discernable.

Figure 7. a) Empirical data. 24-Hour cortisol time courses of five random, representative healthy individuals,

replotted with permission from Hosseinichimeh et al. (2015). b) Simulation results, showing five random,

representative individuals based on Scenario I. c) Simulation results, showing five random, representative

individuals based on Scenario II.

Second, we examined our model's ability to reproduce group-level cortisol time courses in the hours around waking. To this end, we used data from a dataset from Born, Hansen, Marshall, Molle, and Fehm (1999; data presented in Clow, Hucklebridge, Stalder, Evans, & Thorn, 2010), who took blood samples from 15 participants every 15 minutes during the night and the morning, which they then assayed for cortisol. Data from this previous study are replotted in Figure 6a; results from our simulations are plotted in Figure 6b.

Third, we examined our model's ability to simulate individual-level cortisol time courses that are similar to empirical observations. To this end, we used data from a dataset from Carroll et al. (2007; shared for public use by Hosseinichimeh et al., 2015). Carroll et al. (2007) took blood samples from people with depression (N = 12) and healthy controls (N = 17) every 10 minutes for 24 hours, starting at midnight. Data from five random, representative healthy control participants from this previous study are plotted in Figure 7a; representative results from our simulations are plotted in Figure 7b and 7c.

Discussion. In sum, the model simulations suggest that our model is able to reproduce three key sets of empirical findings (Miller et al., 2016; Clow et al., 2009; Hosseinichimeh et al., 2015). In particular, by applying our model, we could successfully reproduce empirically-observed cortisol concentrations, both during the night and during the day, both on the group and the individual level. Thus, at this stage, our model passes the bar to move on to the next step—that is, we conclude that we can build on this model to examine if we can predict the occurrence of disease from work stress.

Nevertheless, we found two slight differences between the empirical data and the simulation results. First, whereas empirical data suggest that cortisol levels never truly approach zero, this does happen in the simulation results (see Figures 5a vs. 5b; 6a vs. 6b; 7a vs. 7b and 7c). Perhaps, this discrepancy stems from the way we simulated work impulses. In

particular, for *all* simulated people, work always ended 9.5 hours after awakening; we modelled no work impulses after that. By contrast, in life, many people remain active well beyond 9.5 hours after awakening, which may expose them to stressors (which may or may not be related to work). Such evening- and night-time activity can explain why measured cortisol levels, at least on the group level, stay above zero throughout the night. Alternatively, it may be the case that some homeostatic mechanism blocks the degradation of cortisol, after cortisol concentrations fall below some threshold. We are not aware of the existence of such a homeostatic mechanism—but if it exists, it may explain the slight discrepancy between the data and the model simulations.

Second, when we examine the group-level cortisol time courses in the hours around waking (Clow et al., 2009), it seems that our simulation results (Figure 5b) suggest a somewhat steeper rise and fall than the empirical data (Figure 5a). However, the simulated time course does seem to fall within the 95% confidence interval of the empirical data (Figure 5a). On a macroscopic level, the two plots are similar.

Study 2: From cortisol to disease

Study 2 explains how cortisol dynamics can lead to disease. Like in Study 1, we compare our model simulations to previously-reported empirical data. Finally, we test the robustness of the predictions for variation in parameter values.

Background. High cortisol concentrations can cause disease via several routes. That is, each organ system is affected in its own way, and even within each organ system, there are often multiple ways in which cortisol can cause damage. A detailed review of all of these routes is beyond the scope of this article; however, in what follows, we will discuss three well-established core pathways. Then, we capture the essence of these pathways in our computational model.

First, cortisol can cause disease through its actions on the cardiovascular system. Cortisol triggers a rise in blood pressure, heart rate, and cardiac output (Grunfeld & Eloy, 1987; Sambhi, Weil, & Udhoji, 1965; Sapolsky et al., 2000). These cardiovascular changes help people deal with stressors and they are fully reversible (*allostatic strain*). Yet, when they are prolonged, these cardiovascular changes increase the risk of vascular lesions, which in turn promote the buildup of arterial plaque at the lesion sites (*allostatic load*; Singh et al., 2002). These plaques harden and clog the arteries, which leads to a disease state called *atherosclerosis* (Dekker et al., 2008; Fantidis, 2010; Singh et al, 2002). Atherosclerosis causes symptoms depending on the location of the artery that is affected. For example, potential symptoms include shortness of breath, trouble speaking, dizziness, pain, and nausea (Mayo Clinic, 2018). Moreover, plaques can get loose, after which they can get stuck in narrower arteries, where they block blood supply to the distal tissue (Falk, 2006). This process is potentially lethal, especially when it takes place in the heart (*myocardial infarction*) or the brain (*stroke*).

Second, cortisol can cause disease through its actions on the immune system. As a part of the acute stress response, the body's capacity to initiate inflammation and fever rapidly increases (e.g., through the cytokine *interleukin 6*; Zhou, Kusnecov, Shurin, DePaoli, & Rabin, 1993). Under normal circumstances, cortisol plays a role in suppressing this rapid inflammatory response, preventing it from overshooting (Sapolsky et al., 2000). This suppressive effect of cortisol is adaptive and fully reversible (*allostatic strain*). Yet, when cortisol levels are high for a longer period of time, the immune system becomes less sensitive to cortisol; that is, prolonged high cortisol causes *glucocorticoid receptor resistance* (Cohen et al., 2012; Cole, 2008; Miller, Cohen, & Ritchley, 2002). People who have glucocorticoid receptor resistance, in turn, are more vulnerable to *nonresolving inflammation*, a condition

that may progress into many disease states, including arthritis, asthma, and cancer (Nathan &

Ding, 2010; Scrivo, Vasile, Bartosiewicz, & Valesini, 2011).

Third, cortisol can cause disease through its actions on the metabolic system. Cortisol

increases circulating glucose, which helps people to sustain their ongoing attempts to deal

with stressors (Sapolsky et al., 2000). This process is adaptive and fully reversible (*allostatic

strain*). However, when cortisol levels stay high, a cascade of physiological processes takes

off, and these processes can together cause damage. Perhaps most notably, although the

interaction between cortisol and insulin is complex, it is fair to say that cortisol has the

potential to disrupt insulin functioning (Joseph & Golden, 2017). Under normal

circumstances, insulin prompts cells to take up glucose, so that the glucose can be used for

glycolysis, the process that energizes cells. Yet, when cortisol levels continue to be high,

maintaining the high levels of circulating glucose, more and more insulin is required to

achieve the same results as before; that is, high cortisol can cause *insulin resistance* (Rizza,

Mandarino, & Gerich, 1982). Moreover, sustained high levels of cortisol, facilitate the

accumulation of abdominal fat (Björntorp, 2001; Rosmond, Dallman, & Björntorp, 1998).

These two processes—the development of insulin resistance and the accumulation of

abdominal fat (*allostatic load*)—can together progress into *diabetes type 2* (Joseph & Golden,

2017). This is a disease state characterized by symptoms such as thirst, hunger (also after

eating), unexplained weight loss, fatigue, and headaches.

As we aimed to develop a parsimonious model, we attempted to capture only the

essence of these cortisol-to-disease mechanisms in our model's formulas. In particular, we

modelled the link between cortisol and disease as follows: (a) the cortisol response puts a

burden on the body that is, in principle, fully reversible (*allostatic strain*); (b) however, when

this burden is sustained, lasting tissue damage occurs (*allostatic load*); (c) such lasting tissue

damage can lead to *disease* (see Figure 1).

To further illustrate how we modelled the cortisol-to-disease pathway, we note that our approach is similar to the so-called *rubber band analogy* (e.g., Logan & Barksdale, 2008; Taylor & Gerrard, 1977a, 1977b). The rubber band analogy holds that the pathway from stress to disease is analogous to how rubber bands can get damaged when they are stretched. Also in this domain, three stages can be distinguished: (a) the rubber stretches, but it can regain its original shape (this is called *elastic deformation*); (b) the rubber stretches further, and it can no longer return to its original shape (this is called *plastic deformation*); (c) the rubber tears (this is called *failure*). These three stages are akin to *allostatic strain*, *allostatic load*, and *disease*, respectively. That is, in our model, the pathway from stress to disease is analogous to how a rubber band, when stretched, subsequently undergoes elastic deformation, plastic deformation, and failure.

We will evaluate our model by examining its ability to reproduce the relationship between work stressors and disease, which is well-supported by previous empirical work. Specifically, the highest level of evidence is available for the relationship between *job strain* (the combination of high job demand and low job control) and cardiovascular disease (Kivimäki & Kawachi, 2015; Nyberg et al., 2013; Steptoe & Kivimäki, 2013; Theorell et al., 2016). That is, a meta-analysis (Kivimäki and Kawachi, 2015), which included over 600,000 participants from 27 cohort studies, revealed a positive relationship between work strain and cardiovascular disease with a cumulative effect size of 1.33 (95% confidence interval: 1.19, 1.49). As *job strain* can be seen as a close proxy for the prevalence of *work stressors*, we will use this previously-observed, positive relationship as a benchmark for our model predictions.

In a computational model. We refer to the current burden on physiological systems as *allostatic strain*, S. Like rubber bands, physiological systems usually return to their original state. In line with this principle, we describe change in *allostatic strain*, dS/dt, as follows:

$$\frac{dS}{dt} = -\rho_R\, S_t + C_t \tag{2}$$

In Equation 2, S_t represents *allostatic strain* at a given time point; ρ_R is the *recovery coefficient*, which determines the speed of recovery; and, C_t is the current force exerted on the physiological system (represented by cortisol in our model).

When allostatic strain exceeds beyond a threshold, we assume that lasting damage occurs. We refer to lasting damage as *allostatic load, L*. The *allostatic threshold*, ε_L, is the point where allostatic load starts to form. We can write:

$$\frac{dL}{dt} = \begin{cases} 0, & S_t < \varepsilon_L \\ S_t - \varepsilon_L, & S_t \geq \varepsilon_L \end{cases} \tag{3}$$

Equation 3 states that allostatic load does not change if the current allostatic strain, S_t, is lower than ε_L. However, when the current allostatic strain exceeds the allostatic threshold, allostatic load increases by the excess, $S_t - \varepsilon_L$.

When physiological systems are repeatedly stretched beyond the allostatic threshold, this will cause allostatic load to accumulate. We assume that when allostatic load passes the *disease threshold*, ε_D, people become diseased. We refer to people's *disease state* as D. We can write:

$$D_t = \begin{cases} False, & L_t < \varepsilon_D \\ True, & L_t \geq \varepsilon_D \end{cases} \tag{4}$$

Equation 4 states that the *disease state*, D_t, is *False* (i.e., the person is not diseased) if allostatic load, L_t, is below the disease threshold ε_D. Conversely, D_t is *True* (i.e., the person is diseased) if L_t exceeds ε_D.

Together, equations 2–4 describe how cortisol can cause disease. Figure 8 illustrates what variation of allostatic strain, allostatic load and disease can be expected in our model

Figure 8. Sample of seven days showing variation on all model variables from five random simulated people.

simulations. The reader should note that because choices of ρ_R, ε_L and ε_D are arbitrary, the

absolute time until disease in our model is meaningless. Rather, the relative time until disease

between simulated individuals is what is of interest in our simulations.

Simulations. We performed simulations based on our model, as defined in Equations

1–4, using several values for the parameters ρ_R, ε_L and ε_D (code available in the online

supplemental material). Specifically, we used three values for each parameter, yielding 27

combinations of parameter values. As the model is not scaled to resemble real-life

observations (e.g., there is no clear-cut measure for *allostatic strain* and *allostatic load*), the

selected parameter values were arbitrary. We selected these parameter values after

exploration, such that, in each simulation, at least some people would become diseased *and* at

least some people would not. Apart from this restriction, we selected parameter values over

the full possible range.

As model input, cortisol time courses were created exactly as described in Study 1

(Table 1), apart from two minor variations. First, we aimed to increase the precision of

detecting differences that exist in the simulated population, without needing to increase the

number of simulated people. Thus, instead of simulating a population with a variation in work

impulses that is described by a Gamma distribution, we simulated individual variation in work

impulses from a uniform distribution over the same range of work impulses. In other words,

instead of creating a population with a variation in work impulses that may resemble real-life

populations, we now simulated a population with work impulses that are evenly distributed.

Apart from increasing the precision of detecting effects, this variation does not otherwise

influence the outcome of the simulations. Second, to save computational time, we simulated

5,000 instead of 10,000 people per parameter combination. Like before, for each parameter

combination, we simulated both a scenario where night impulses and work impulses were

uncorrelated (Scenario I), as well as a scenario where they were correlated (Scenario II).

Simulation results. Figure 9 shows the results of the simulations. The figure shows the simulated relative risk of people becoming diseased, as a function of the number of work impulses. The online supplemental material includes a plot of accumulated allostatic load against the average amount of daily work stressors.

Results indicate that, in case of Scenario II, where work and night impulses are correlated, the model was robust to variation of the parameters ρ_R, ε_L and ε_D. Irrespective of the variation in the parameter values, we found a positive relationship between the frequency of work impulses and the simulated risk of becoming diseased (note that, because choices of ρ_R, ε_L and ε_D are arbitrary, the absolute effect sizes are meaningless). Also, we observed little variation in the predicted effect sizes, regardless of the specific parameter values (Figure 9b). These results are in line with the well-established empirical relationship between work stressors and disease (Kivimäki & Kawachi, 2015), which we aimed to reproduce.

We did not observe this same robustness when we examined Scenario I. Although we did observe a positive relationship between work impulses and the simulated risk of disease for a number of parameter combinations, this relationship did not emerge in all cases (Figure 9a). Also, further exploratory analyses revealed that the cases where a positive relationship was observed for Scenario I, an unrealistically high proportion of simulated people had become diseased (see online supplemental material).

Discussion. At least when we assume that work and night impulses are correlated (Scenario II), which we would expect if people anticipate the events of the following workday (Fries et al., 2005; Rohleder et al., 2007), the model successfully produced a positive correlation between work impulses and disease. This relationship, which is in line with the relationship between work stress and disease that is often observed in empirical literature (Kivimäki & Kawachi, 2015), holds for a wide range of values of the parameters ρ_R, ε_L and ε_D, supporting the robustness of the model. However, we did not find a robust relationship

Figure 9. Simulation results of relative risk of becoming diseased at various time points in the simulation for

Scenario I (a; no correlation between work and night impulses) Scenario II (b; correlation between work and

night impulses). The odds ratios are calculated from the standardized averages of work impulses (i.e., the effect

sizes represent one SD difference in the average number work impulses). Odd ratios greater than 100 are

presented as 100.

between work impulses and disease if work and night impulses were uncorrelated (Scenario

I).

With caution, this set of findings raises the possibility that acute responses to work

stressors may play only a limited role in the development of disease. Rather, it appears that, in

comparison to the response to acute work stressors (the work impulses in our model), the

contribution of the night impulses is so pronounced, that the cortisol spikes from the acute

stressors during the working day contribute relatively little to the accumulation of allostatic

load. In line with this observation, exploration of individual time courses of allostatic load

show that the increase in allostatic load is mainly seen at the time point of the morning peak

(for an illustration, see Figure 8).

Study 3: New predictions for variations in workweek configurations

To recap, so far, we developed a computational model of work stress and disease. We

compared the predictions from model simulations against empirical data. We found that, if we

assume that night impulses (i.e., activity of the HPA axis during the night) and work impulses

work impulses (i.e., activity of the HPA axis due to work stressors) are correlated, our model

can reproduce results from empirical research. Based on this finding, we conclude that our

model is an appropriate model of the relationship between work stress and disease.

We will now proceed with generating new predictions, based on our model. In

particular, we examine the effect of how working hours are distributed over the week. We

examine the consequences of several workweek configurations on the development of

disease.

Simulations. In all simulations, we used a *standard working week* (Monday to Friday,

8 hours per day), like we used in all previous sections (Table 1), as a benchmark. Table 2

presents all workweek configurations that we examined with separate simulations; Table 3

describes the parameter values that were identical across all simulations. Note that these

values are identical to the values that we used previously, except that we used only one

combination of the parameters ρ_R, ε_L and ε_D (simulations for other parameter combinations

are reported in online supplemental material). Also, based on our previous findings (see Study

2), in all simulations that follow, we assumed a correlation between night and work impulses.

Table 2

Workweek configurations that we explored with simulations

Configuration	Working days	Days worked (per week)	Hours worked (per day)		
			30h/week	40h/week	50h/week
#1	Mon–Fri	5	6h	8h*	10h
#2	Mon–Tue and Thu–Sat	5	6h	8h	10h
#3	Mon–Tue and Thu–Fri	4	7h30m	10h	12h30m
#4	Mon–Tue and Thu	3	10h	13h20m	16h40m
#5	Mon–Sat	6	5h	6h40m	8h20m
#6	Mon–Sun	7	4h17m	5h43m	7h09m

Note: * = Benchmark for all other configurations.

Table 3

Parameter settings that were held constant between the simulations of varying worktime configurations.

Parameter	Value
Sampling frequency	2
People simulated	5'000
Days simulated per person	200
Night impulses, quantity	$X \sim \Gamma(k = 3, \theta = 14)$
Night impulses, moment	$X \sim T_{wake} - \text{EXP}(\lambda = 1)$
Work impulses, quantity	$X \sim \mathcal{U}(a = 1, b = 50)$
Work impulses, moment	$X \sim \mathcal{U}(a = T_{ws}, b = T_{we})$
Cortisol decay constant (ρ_C)	$X \sim \mathcal{N}(\mu = .52, \sigma^2 = .05^2)$
HPA Axis scaling constant (κ_{HPA})	2.20
Cortisol response delay, τ	30 min
Wake time, T_{wake}	7AM
Work start, T_{ws}	8:30AM
Elasticity constant (ρ_R)	0.6
Allostatic threshold (ε_L)	20
Disease threshold (ε_D)	400

Simulation results and discussion. Figure 10 provides an overview of all simulation results, showing the predicted effects of all workweek configurations in Table 2 on the development of disease. The results show a clear pattern. Specifically, we found an increase in predicted disease risk for configurations #3 and #4, in which working hours are concentrated in a limited number of days (a so-called *compressed working week;* Baltes, Briggs, Huff, Wright, & Neuman, 1999). In contrast, the results show a decrease in disease risk in configurations #5 and #6, in which working hours are spread out over the week. We found no effect of configuration #2, in which work-free days are distributed throughout the week, not chunked together in a weekend. These effects were independent of the total number of hours (30, 40, or 50) that simulated people worked in a week. In sum, our model predicts that spreading out working hours over more days, rather than concentrating working hours in less days, helps prevent the development of disease.

We note another, related regularity from these simulation results (Figure 10). That is, independently of the number of working hours per week, there is a clear, positive relation between the number of working hours per day and the predicted disease risk. In line with this regularity, our simulations suggest that 50-hour working weeks can be relatively healthy—i.e., healthier than the standard 40-hour/5-day working week—as long as these 50 hours are distributed over 7 short days (7.09 hours per day). Similarly, our simulations suggest that 30-hour working weeks can be relatively unhealthy, when these 30 hours are cramped into 3 long days (10 hours per day). In sum, although the total number of weekly working hours does matter (on average, disease risk is higher when people work more hours per week), our predictions suggests that the total number of daily working hours plays a more dominant role.

Figure 10. New predictions based on our model. The plot represents the relative risk of developing disease, as a

function of different workweek configurations. All predictions are relative to a standard working week (i.e.,

configuration #1: 5 working days of 8 hours, Monday to Friday). See Table 2 for an explanation of all workweek

configurations that we examined.

These simulation results speak to a recent set of trials conducted in Sweden, where

employees were allowed to work 30-hour/5-day working weeks (instead of 40-hour/5-day

working weeks; no pay cut applied). Although results of these trials were reported

informally—to our knowledge, mainly in media publications (e.g., Alderman, 2016;

Oltermann, 2017)—our simulation results can be used to make new predictions about whether

such trials are likely to support employees' long-term health. Specifically, we predict that the

Swedish 30-hour configuration (i.e., configuration #1; 30-hour/5-day) is indeed healthier than

its 40-hour counterpart (i.e., configuration #1; 40-hour/5-day), but that there may be even

healthier ways to distribute 30 hours over a working week (i.e., configuration #5, 30-hour/6-

day; and configuration #6, 30-hour/7-day).

General discussion

In this research we aimed to explain how work stress can cause disease, by developing a new computational model. In Study 1, we found that a model that defines cortisol dynamics in two linear relationships (i.e., release of cortisol is proportional to the number of stressors; decay of cortisol is proportional to the current cortisol level), can reproduce the characteristic shape of the cortisol day curve very well (Born et al., 1999; Carroll et al. 2007; Miller et al., 2016). In Study 2, we found that we could reproduce the previously-reported relationship between work stress and disease (Kivimäki & Kawachi, 2015), by formalizing the central predictions from *allostatic load theory*. In Study 3, based on our model, we formulated several novel predictions about the relationship between work stress and disease.

Implications

Our research makes several contributions to the literature on the link between work stress and disease. First, we provide a first formalization of the core theoretical ideas in this domain, which previously existed only in the form of verbal theory. Thus, our computational model *contributes to the falsifiability* of these existing core ideas. Indeed, in our model, there are no implicit assumptions and there is no flexibility in interpretation of how parameters interact. As a result, going beyond previous theories, we can be explicit as to what observations would be needed to falsify our model: any pattern of data that is inconsistent with our predictions (Figure 10), would falsify our model. In that case, our model would need to be improved or replaced by an alternative model.

Second, by developing our model, we gained *new insights about the mechanisms* that together form the causal chain between work stress and disease. For example, we learned that the key logic of allostatic load theory holds when its predictions are scrutinized using a computational model, suggesting that allostatic load theory provides, in principle, a set of premises that is useful to understand how work stress can cause disease. Perhaps more

importantly, we found that to show a robust link between work stress and disease, we need to assume that night impulses into the HPA axis are correlated with work impulses to the HPA axis. This finding is consistent with the idea that the morning peak, at least in part, stems from the *anticipation* of upcoming stressors (Fries et al., 2009; Rohleder et al., 2007). Our findings suggest that the acute cortisol response to work stressors by themselves, does not strongly contribute to the development of disease. Instead, work stressors may cause disease as their anticipation (e.g., during the night before) augments the morning peak. Interestingly, in line with the latter idea, several studies show that cardiovascular incidents are indeed most frequent in the first hours after awakening (e.g., Manfredini et al., 2005; Muller et al., 1985).

Third, through simulations, we made several *new predictions* about the relationship between working hours and the development of disease. In essence, our model predicts that spreading out working hours over more days, rather than concentrating working hours in less days, prevents the development of disease. Relatedly, our model predicts that the number of working hours per day (more so than the number of working hours per week) strongly predicts the development of disease. In the following paragraphs, we will discuss these predictions against the background of existing literature.

The *compressed working week.* Our predictions are related to previous research on the concept of the *compressed working week.* The compressed working week is defined as a working week in which the weekly number of working hours is completed in fewer than five working days (Baltes et al., 1999). Generally, researchers hypothesized that the compressed working week should benefit employees' well-being via the restorative effects of having longer weekends. In particular, longer weekends should enable employees to pursue more leisure activities; to invest more time in social relationships (Baltes et al., 1999; Bambra, Whitehead, Sowden, Akers, & Petticrew, 2007; Dall'ora, Ball, Recio-Saucedo, & Griffiths, 2016); and to detach and recover more strongly from work (Geurts, Beckers, & Tucker,

2014). However, findings are mixed: some studies show that compressed working weeks

increase well-being (or job satisfaction; Bambra et al., 2007; Baltes et al., 1999), but other

studies show negative effects (Dall'ora et al., 2016).

How to reconcile these mixed findings with our prediction that compressed working

weeks can cause disease on the long run? We should point out that it is questionable whether

these previous studies are relevant to our prediction at all. After all, they do not directly speak

to employees' long-term physical health. Nevertheless, if we sidestep this important issue,

these mixed findings are inconsistent with our predictions. This apparent inconsistency may

be explained by one specific assumption in our model simulations, i.e., the assumption that

the number of daily work stressors is linearly related to the number of working hours. In other

words, we assumed to longer work days cause proportionally more impulses into the HPA

axis. It is possible, though, that employees experience longer working days (in compressed

working weeks) as an expression of *control* over their working hours, mitigating the negative

effect from work stress. Indeed, the perception of control over working hours is a powerful

mechanism to increase the impact of stressors (Beckers, Kompier, Kecklund, & Härmä,

2012). In future research, our model (and our simulation approach) can be used to further

explore this issue.

Independent of this assumption regarding longer work days, we should note that our

simulations still do not suggest a special health benefit from having longer weekends. In

particular, in our simulations, increases in *allostatic load* happened on each individual

workday, rather than that increases in allostatic load gradually developed throughout the

working week. So, our simulations suggest that it does not matter whether two (or more)

work-free days are chunked together in a weekend. If research proves otherwise—i.e., if data

would convincingly show that, all else being equal, longer weekends protect long-term

health—this would falsify our model, and an improvement to our model would be necessary (e.g., we would need to add a parameter).

Chronic stress and the cortisol *morning peak*. Our simulations suggest a relationship between the cortisol *morning peak* and the development of disease. A larger morning peak, so we predict, can cause disease in the long run. We assume that the morning peak stems from night impulses into the HPA axis, which are in part due to the anticipation of the upcoming day.

Interestingly, previous research suggests that the amplitude of the morning peak changes when people chronically experience stress (Miller, Chen, & Zhou, 2007). In particular, some studies report that chronic stress leads to *reduced* responsiveness of the HPA axis, causing *hypocortisolism*, of which a lower morning peak is a key symptom (Fries, Hesse, Hellhammer, & Hellhammer, 2005). Hypocortisolism has been reported in people who have previously experienced sustained or intense stress, such as people with post-traumatic stress disorder (Rohleder, Joksimovic, Wolf, & Kirschbaum, 2004) and perhaps people with severe burnout (Lennartson, Sjörs, Währborg, Ljung, & Jonsdottir, 2015). Speculatively, stress-induced hypocortisolism may function to protect people from further physical damage (Fries et al., 2005). After all, lower cortisol levels lead to less sustained burden on the cardiovascular, metabolic, and immune systems, potentially preventing disease.

In sum, (a) morning peaks may contribute to disease, but (b) the central nervous system may suppress morning peaks after exposure to sustained stress, and (c) such suppression, which happens in people with burnout, may protect people from further damage. Connecting these three ideas, it is possible to formulate a novel perspective on the nature of burnout. In particular, burnout is traditionally conceptualized as a highly aversive syndrome that emerges from known work-related conditions (Demerouti, Bakker, Nachreiner, & Schaufeli, 2001), that can be best treated in a way that is tailored to the client (e.g., Farber,

2000), making use of established treatments for depression (Schonfeld & Bianchi, 2016). We suggest that burnout can also be seen as an adaptive state that shields people from developing potentially lethal conditions, such as myocardial infarctions. This alternative conceptualization (burnout as a protective mode of functioning) is not necessarily better than the original (burnout as a work-related syndrome related to depression), but we suggest that it is potentially productive to consider both conceptualizations together in future research.

Limitations and future directions

As we strived to develop a minimal model, with only few parameters, we have simplified reality in several ways. Each of these simplifications constitutes a potential limitation of our model, as they may cause our model to have one or more blind spots.

First, perhaps reflecting the most rigorous simplification of reality in our study (for a discussion, see Kagan, 2016), we used cortisol as the only indicator of allostasis. However, allostasis is thought to involve a host of regulatory systems, including not just the HPA axis, but also the sympathetic branch of the autonomic nervous system (McEwen, 1998; Rohleder et al. 2007). We chose to focus only on cortisol, because the role of cortisol in stress is very well-studied, which allowed us to calibrate our model to large samples (Miller et al., 2016). Moreover, cortisol affects several regulatory systems, including the sympathetic nervous system. Still, our reductionistic approach constitutes a departure from previous, broader attempts to operationalize allostasis and allostatic load (Seeman, Epel, Gruenewald, Karlamangla, & McEwen, 2010), and future research is needed to examine whether this simplification can be justified.

Second, our model only considers variation in the *frequency* of work stressors, not their *duration*. In fact, we assumed that all impulses into the HPA axis have the same duration. Importantly, despite this limitation, our model was well able to reproduce core characteristics of the cortisol day curve (Figures 5–7). So, on first sight, it seems that adding

variation in the duration of stressors would not lead to a big improvement in our model. Nevertheless, real-life stressors vary in their duration, and stressors may differ in the duration in the physiological response that they trigger (e.g., stressors may have a sustained impact if they lead to perseverative cognition, Brosschot et al., 2005). More broadly, at least in its current form, our model cannot be used to make predictions about the impact of between-stressor differences (e.g., whether stressors stem from work vs. from other sources; whether stressors are short vs. long). We acknowledge that these differences exist, and that they may matter.

Third, in our simulations, we assumed the same values for all simulated individuals for the parameters ρ_R (recovery coefficient), ε_L (allostatic load threshold) and ε_D (disease threshold). As some of our parameters were already unscaled (i.e., they were not directly linked to quantities measured in real life, e.g., this was the case for S, allostatic strain, and L, allostatic load), we chose not to vary values for ρ_R, ε_L and ε_D in order to avoid adding additional arbitrary assumptions of inter-individual variability. However, it unlikely that this simplification is realistic; in fact, it is plausible that inter-individual variation in health and healthy lifestyle are represented in these parameters. Future research is needed to examine the impact of such variation.

Fourth, we should note that the fact that we included unscaled parameters prevented us from estimating precise effect sizes and time-to-disease values. However, we should note that our model does allow for making relative comparisons, to determine which scenario has a higher risk of leading to disease. If researchers succeed in determining meaningful values for some of the parameters (e.g., S and L), it will be possible to estimate effect sizes of different workweek configurations and to estimate time-to-disease values, which we think would present a valuable and meaningful addition to the current model. Although this was not the aim of this study, we do feel this constitutes an interesting direction for future research.

Fifth, other mechanisms than the one modeled in our computational model, can potentially explain the relation between work stress and disease. Perhaps most importantly, our model does not consider the possibility that stress may lead to disease through poor health decisions (e.g., unhealthy eating, Araiza & Lobel, 2018; smoking, Kouvonen, Kivimäki, Virtanen, Pentti, & Vahtera, 2005; alcohol use, Frone, 2016). Moreover, some research suggests that gradual changes in the circadian cortisol profile after a long period of chronic stress (specifically, a flattened profile with blunted morning cortisol peaks and elevated cortisol levels during the rest of the day) contributes to stress related disease (e.g. Heim, Ehlert, & Hellhammer, 2000). Our computational model models neither of these mechanisms. However, as we mentioned in the introduction, rather than to provide a comprehensive model, we aimed to create and test a parsimonious and straightforward model of the stress–disease relationship. This being said, we do envision that modelling these alternative mechanisms will lead to interesting insights. In particular, a model that includes the health behavior route may lead to interesting, counter-intuitive predictions, as such a model is likely to involve complex dynamics.

Taken together, we see several avenues for future research. Most importantly, the first next step should be to collect empirical data (or to re-use existing empirical data) to critically test our model's predictions. This would help examine whether our model's development is on the right track, or whether large changes are needed. However, even in its current form, our model can already be used to generate predictions about specific configurations of working weeks, beyond those covered in this paper. Finally, our model can be adapted or refined, e.g., by adding and/or removing parameters, to make new predictions about the role of job characteristics (e.g., worktime control, job control, social support, emotional and physical demands of work, task variety), and perhaps, to examine the long-term health

implications of other aspects of working life (e.g., whether people can recover well at home; whether people encounter stressors for non-work-related reasons).

Conclusion

The consequences of work stress are costly to individuals, organizations, and society at large. By formalizing the central assumptions of allostatic load theory, we were able to derive new hypotheses (about the effects of workweek configurations), which would have been difficult to derive through intuitive reasoning. At the same time, we contributed to the science of work stress by providing a model that is directly falsifiable. Reflecting on these results, we strongly feel that computational modeling is a fruitful approach not just for science, technology, and engineering—but that this approach can also help to make progress in the domain of occupational health. We hope this research may serve as a blueprint for making further progress in this important area.

References

Alderman, L. (May 20, 2016). In Sweden, experiment turns shorter workdays into bigger gains. *The New York Times.* Retrieved from http://www.nytimes.com

Allport, F. H. (1924). *Social psychology.* Boston: Houghton Mifflin. Retrieved from https://brocku.ca/MeadProject/Allport/1924/1924_toc.html

Araiza, A. M., & Lobel, M. (2018). Stress and eating: Definitions, findings, explanations, and implications. *Social and Personality Psychology Compass, 12*, e12378. https://doi.org/10.1111/spc3.12378

Baltes, B. B., Briggs, T. E., Huff, J. W., Wright, J. A., & Neuman, G. A. (1999). Flexible and compressed workweek schedules: A meta-analysis of their effects on work-related criteria. *Journal of Applied Psychology*, *84*, 496–513. https://doi.org/10.1037/0021-9010.84.4.496

Bambra, C., Whitehead, M., Sowden, A., Akers, J., & Petticrew, M. (2008). "A hard day's night?" The effects of compressed working week interventions on the health and work-life balance of shift workers: A systematic review. *Journal of Epidemiology and Community Health*, *62*, 764-777. https://doi.org/10.1136/jech.2007.067249

Beckers, D. G. J., Kompier, M. A. J., Kecklund, G., & Härmä, M. (2012). Worktime control: Theoretical conceptualization, current empirical knowledge, and research agenda. *Scandinavian Journal of Work, Environment and Health*, *38*, 291–297. https://doi.org/10.5271/sjweh.3308

Björntorp, P. (2001). Do stress reactions cause abdominal obesity and comorbidities? *Obesity Reviews, 2*, 73-86. https://doi.org/10.1046/j.1467-789x.2001.00027.x

Born, J., Hansen, K., Marshall, L., Molle, M., & Fehm, H. L. (1999). Timing the end of

nocturnal sleep. *Nature, 397,* 29–30. https://doi.org/10.1038/16166

Brosschot, J. F., Pieper, S., & Thayer, J. F. (2005). Expanding stress theory: Prolonged

activation and perseverative cognition. *Psychoneuroendocrinology,* 30, 1043-1049.

https://doi.org/10.1016/j.psyneuen.2005.04.008

Carroll, B. J., Cassidy, F., Naftolowitz, D., Tatham, N. E., Wilson, W. H., Iranmanesh, …

Veldhuis, J. D. (2007). Pathophysiology of hypercortisolism in depression. *Acta*

Psychiatrica Scandinavica, *115*, 90–103. https://doi.org/10.1111/j.1600-

0447.2007.00967.x

Cohen, S., Janicki-Deverts, D., Doyle, W. J., Miller, G. E., Frank, E., Rabin, B. S., & Turner,

R. B. (2012). Chronic stress, glucocorticoid receptor resistance, inflammation, and

disease risk. *Proceedings of the National Academy of Sciences, 109*, 5995-5999.

https://doi.org/10.1073/pnas.1118355109

Cole, S. W. (2008). Social regulation of leukocyte homeostasis: the role of glucocorticoid

sensitivity. *Brain, Behavior, and Immunity, 22*, 1049-1055.

https://doi.org/10.1016/j.bbi.2008.02.006

Clow, A., Hucklebridge, F., Stalder, T., Evans, P., & Thorn, L. (2010). The cortisol

awakening response: More than a measure of HPA axis function. *Neuroscience and*

Biobehavioral Reviews, 35, 97-103. https://doi.org/10.1016/j.neubiorev.2009.12.011

Dall 'ora, C., Ball, J., Recio-Saucedo, A., & Griffiths, P. (2016). Characteristics of shift work

and their impact on employee performance and wellbeing: A literature review.

International Journal of Nursing Studies, *57*, 12–27.

https://doi.org/10.1016/j.ijnurstu.2016.01.007

Das, S., & O'Keefe, J. H. (2008). Behavioral cardiology: Recognizing and addressing the

profound impact of psychosocial stress on cardiovascular health. *Current Hypertension*

Reports, *10*, 374–381. https://doi.org/10.1007/s11906-008-0070-6

Dekker, M. J. H. J., Koper, J. W., Van Aken, M. O., Pols, H. A. P., Hofman, A., De Jong, F.

H., ... & Tiemeier, H. (2008). Salivary cortisol is related to atherosclerosis of carotid

arteries. *Journal of Clinical Endocrinology & Metabolism, 93*, 3741-3747.

https://doi.org/10.1210/jc.2008-0496

Demerouti, E., Bakker, A. B., Nachreiner, F., & Schaufeli, W. B. (2001). The job demands-

resources model of burnout. *Journal of Applied Psychology, 86*, 499-512.

https://doi.org/10.1037/0021-9010.86.3.499

Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: a

theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, *130*,

355–391. https://doi.org/10.1037/0033-2909.130.3.355

Downey, A. B. (2017). *Modeling and Simulation in Python*. Needham, Massachusetts: Green

Tea Press. Retrieved from http://greenteapress.com/ModSimPy/ModSimPy.pdf

Edwards, J. R., & Berry, J. W. (2010). The presence of something or the absence of nothing:

Increasing theoretical precision in management research. *Organizational Research*

Methods, *13*, 668–689. https://doi.org/10.1177/1094428110380467

Epstein, J. (2008). Why model? *Journal of Artificial Societies and Social Simulation, 11*, 12.

https://doi.org/10.1080/01969720490426803

Esch, T., Stefano, G. B., Fricchione, G. L., & Benson, H. (2002). Stress in cardiovascular

   diseases. *Medical Science Monitor*, *8*, 93–102. Retrieved from

   https://www.medscimonit.com/download/index/idArt/420828

Falk, E. (2006). Pathogenesis of atherosclerosis. *Journal of the American College of*

   Cardiology, 47, C7-C12. https://doi.org/10.1016/j.jacc.2005.09.068

Fantidis, P. (2010). The role of the stress-related anti-inflammatory hormones ACTH and

   cortisol in atherosclerosis. *Current Vascular Pharmacology, 8*, 517-525.

   https://doi.org/10.2174/157016110791330889

Farber, B. A. (2000). Treatment strategies for different types of teacher burnout. *Journal of*

   Clinical Psychology, 56, 675-689. https://doi.org/10.1002/(SICI)1097-

   4679(200005)56:5<675::AID-JCLP8>3.0.CO;2-D

Farrell, S., & Lewandowsky, S. (2010). Computational models as aids to better reasoning in

   psychology. *Current Directions in Psychological Science*, *19*, 329–335.

   https://doi.org/10.1177/0963721410386677

Frankenhuis, W. E., & Tiokhin, L. (2018). Bridging evolutionary biology and developmental

   psychology: Toward an enduring theoretical infrastructure. *Child Development*, *89*,

   2303-2306. https://doi.org/10.1111/cdev.13021

Fries, E., Dettenborn, L., & Kirschbaum, C. (2009). The cortisol awakening response (CAR):

   Facts and future directions. *International Journal of Psychophysiology*, *72*, 67–73.

   https://doi.org/10.1016/j.ijpsycho.2008.03.014

Fries, E., Hesse, J., Hellhammer, J., & Hellhammer, D. H. (2005). A new view on

    hypocortisolism. *Psychoneuroendocrinology, 30*, 1010-1016.

    https://doi.org/10.1016/j.psyneuen.2005.04.006

Frone, M. R. (2016). Work stress and alcohol use: Developing and testing a biphasic self-

    medication model. *Work & Stress, 30*, 374-394.

    https://doi.org/10.1080/02678373.2016.1252971

Geurts, S. A. E., Beckers, D. G. J., & Tucker, P. (2014). Recovery from demanding work

    hours. In M. C. W. Peeters, J. de Jonge, & T. W. Taris (Eds.), *An introduction to*

    contemporary work psychology (1st Ed., pp. 196–219). Hoboken, NJ: Wiley. Retrieved

    from https://cronfa.swan.ac.uk/Record/cronfa18761

Geurts, S. A. E., & Sonnentag, S. (2006). Recovery as an explanatory mechanism in the

    relation between acute stress reactions and chronic health impairment. *Scandinavian*

    Journal of Work, Environment and Health, 32, 482-492.

    https://doi.org/10.5271/sjweh.1053

Goodman, W. K., Janson, J., & Wolf, J. M. (2017). Meta-analytical assessment of the effects

    of protocol variations on cortisol responses to the Trier Social Stress Test.

    Psychoneuroendocrinology, 80, 26-35. https://doi.org/10.1016/j.psyneuen.2017.02.030

Grünfeld, J. P., & Eloy, L. (1987). Glucocorticoids modulate vascular reactivity in the rat.

    Hypertension, 10, 608-618. https://doi.org/10.1161/01.HYP.10.6.608

Juster, R. P., McEwen, B. S., & Lupien, S. J. (2010). Allostatic load biomarkers of chronic

    stress and impact on health and cognition. *Neuroscience and Biobehavioral Reviews*.

    https://doi.org/10.1016/j.neubiorev.2009.10.002

Hassard, J., Teoh, K.R.H., Visockaite, G., Dewe, P., & Cox, T. (2018). The Cost of Work-Related Stress to Society: A Systematic Review. *Journal of Occupational Health Psychology, 23,* 1-17. https://dx.doi.org/10.1037/ocp0000069

Hazewinkel, M. (2002). Gamma-distribution. In *Encyclopedia of Mathematics*. New York: Springer. Retrieved from http://www.encyclopediaofmath.org/index.php?title=Gamma-distribution&oldid=18532

Heim, C., Ehlert, U., & Hellhammer, D. H. (2000). The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology*, *25*, 1–35.

Hellhammer, D. H., Wüst, S., & Kudielka, B. M. (2009). Salivary cortisol as a biomarker in stress research. *Psychoneuroendocrinology*, *34*, 163–171. https://doi.org/10.1016/j.psyneuen.2008.10.026

Hosseinichimeh, N., Rahmandad, H., & Wittenborn, A. K. (2015). Modeling the hypothalamus-pituitary-adrenal axis: A review and extension. *Mathematical Biosciences*, *268*, 52–65. https://doi.org/10.1016/j.mbs.2015.08.004

Joseph, J. J., & Golden, S. H. (2017). Cortisol dysregulation: the bidirectional link between stress, depression, and type 2 diabetes mellitus. *Annals of the New York Academy of Sciences, 1391*, 20-34. https://doi.org/10.1111/nyas.13217

Lennartsson, A. K., Sjörs, A., Währborg, P., Ljung, T., & Jonsdottir, I. H. (2015). Burnout and hypocortisolism–a matter of severity? A study on ACTH and cortisol responses to acute psychosocial stress. *Frontiers in Psychiatry, 6*, 8. https://doi.org/10.3389/fpsyt.2015.00008

Logan, J. G., & Barksdale, D. J. (2008). Allostasis and allostatic load: expanding the discourse on stress and cardiovascular disease. *Journal of Clinical Nursing, 17*, 201-208. https://doi.org/10.1111/j.1365-2702.2008.02347.x

Kagan, J. (2016). An overly permissive extension. *Perspectives on Psychological Science, 11*, 442-450. https://doi.org/10.1177/1745691616635593

Kaplan, & Glass. (1995). Finite-difference equations. In *Understanding Nonlinear Dynamics* (pp. 1–33). New York: Springer. https://doi.org/10.1007/978-1-4612-0823-5

Kivimäki, M., & Kawachi, I. (2015). Work stress as a risk factor for cardiovascular disease. *Current Cardiology Reports*, 17-74. https://doi.org/10.1007/s11886-015-0630-8

Kokko, H. (2007). *Modelling for field biologists and other interesting people*. New York: Cambridge University Press. Retrieved from www.cambridge.org/9780521831321

Kouvonen, A., Kivimäki, M., Virtanen, M., Pentti, J., & Vahtera, J. (2005). Work stress, smoking status, and smoking intensity: an observational study of 46 190 employees. Journal of Epidemiology & Community Health, 59, 63-69. http://dx.doi.org/10.1136/jech.2004.019752

Kovacs, W. J., & Ojeda, S. R. (2011). *Textbook of Endocrine Physiology* (6th Ed.). New York: Oxford University Press. Retrieved from https://global.oup.com/academic/product/textbook-of-endocrine-physiology-9780199744121?cc=nl&lang=en&

Manfredini, R., Boari, B., Smolensky, M. H., Salmi, R., la Cecilia, O., Maria Malagoni, A., ... & Manfredini, F. (2005). Circadian variation in stroke onset: identical temporal pattern

in ischemic and hemorrhagic events. *Chronobiology International, 22*, 417-453.

https://doi.org/10.1081/CBI-200062927

Maslach, C., Schaufeli, W. B., & Leiter, M. P. (2001). Job burnout. *Annual Review of*

Psychology, *52*, 397–422. https://doi.org/10.1146/annurev.psych.52.1.397

Mayo Clinic (April 24, 2018). *Arteriosclerosis / atherosclerosis.* Retrieved from

https://www.mayoclinic.org/diseases-conditions/arteriosclerosis-

atherosclerosis/symptoms-causes/syc-20350569

McEwen, B. S. (1998). Stress, adaptation, and disease. Allostasis and allostatic load. *Annals*

of the New York Academy of Sciences, *840*, 33–44. https://doi.org/10.1111/j.1749-

6632.1998.tb09546.x

Meehl, P. E. (1978). Theoretical risks and tabular asterisks: Sir Karl, Sir Ronald, and the slow

progress of soft psychology. *Journal of Consulting and Clinical Psychology, 46*, 806.

https://doi.org/10.1016/j.appsy.2004.02.002

Miller, G. E., Chen, E., & Zhou, E. S. (2007). If it goes up, must it come down? Chronic

stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological*

Bulletin, *133*(1), 25–45. https://doi.org/10.1037/0033-2909.133.1.25

Miller, G. E., Cohen, S., & Ritchey, A. K. (2002). Chronic psychological stress and the

regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health*

Psychology, 21, 531-541. https://doi.org/10.1037/0278-6133.21.6.531

Miller, R., Stalder, T., Jarczok, M., Almeida, D. M., Badrick, E., Bartels, M., … Kirschbaum,

C. (2016). The CIRCORT database: Reference ranges and seasonal changes in diurnal

salivary cortisol derived from a meta-dataset comprised of 15 field studies.

Psychoneuroendocrinology, *73*, 16–23. https://doi.org/10.1016/j.psyneuen.2016.07.201

Muller, J. E., Stone, P. H., Turi, Z. G., Rutherford, J. D., Czeisler, C. A., Parker, C., ... &

Sobel, B. E. (1985). Circadian variation in the frequency of onset of acute myocardial

infarction. *New England Journal of Medicine, 313,* 1315-1322.

https://doi.org/10.1056/NEJM198511213132103

Nathan, C., & Ding, A. (2010). Nonresolving inflammation. *Cell, 140,* 871-882.

https://doi.org/10.1016/j.cell.2010.02.029

Nettle, D., Frankenhuis, W. E., & Rickard, I. J. (2013). The evolution of predictive adaptive

responses in human life history. *Proceedings of the Royal Society*, *280*, 20131343.

https://doi.org/10.1098/rspb.2013.1343

Niv, Y., Daw, N. D., Joel, D., & Dayan, P. (2007). Tonic dopamine: Opportunity costs and

the control of response vigor. *Psychopharmacology*, *191*, 507–520.

https://doi.org/10.1007/s00213-006-0502-4

Nyberg, S. T., Fransson, E. I., Heikkilä, K., Alfredsson, L., Casini, A., Clays, E., …

Kivimäki, M. (2013). Job strain and cardiovascular disease risk factors: Meta-analysis of

individual-participant data from 47,000 men and women. *PLoS ONE*, *8*, e67323.

https://doi.org/10.1371/journal.pone.0067323

Oltermann, P. (2017, January 4). Sweden sees benefits of six-hour working day in trial for

care workers. *The Guardian*. Retrieved from https://www.theguardian.com

Oosterholt, B. G., Maes, J. H. R., Van der Linden, D., Verbraak, M. J. P. M., & Kompier, M.

A. J. (2015). Burnout and cortisol: Evidence for a lower cortisol awakening response in

both clinical and non-clinical burnout. *Journal of Psychosomatic Research*, *78*, 445–451. https://doi.org/10.1016/j.jpsychores.2014.11.003

Oosterholt, B. G., Maes, J. H. R., Van Der Linden, D., Verbraak, M. J. P. M., & Kompier, M. A. J. (2016). Getting better, but not well: A 1.5 year follow-up of cognitive performance and cortisol levels in clinical and non-Clinical burnout. *Biological Psychology*, *117*, 89–99. https://doi.org/10.1016/j.biopsycho.2016.02.009

Rizza, R. A., Mandarino, L. J., & Gerich, J. E. (1982). Cortisol-induced insulin resistance in man: impaired suppression of glucose production and stimulation of glucose utilization due to a postreceptor defect of insulin action. *Journal of Clinical Endocrinology & Metabolism, 54*, 131-138. https://doi.org/10.1210/jcem-54-1-131

Rohleder, N., Beulen, S. E., Chen, E., Wolf, J. M., & Kirschbaum, C. (2007). Stress on the dance floor: The cortisol stress response to social-evaluative threat in competitive ballroom dancers. *Personality and Social Psychology Bulletin*, *33*, 69–84. https://doi.org/10.1177/0146167206293986

Rohleder, N., Joksimovic, L., Wolf, J. M., & Kirschbaum, C. (2004). Hypocortisolism and increased glucocorticoid sensitivity of pro-inflammatory cytokine production in Bosnian war refugees with posttraumatic stress disorder. *Biological Psychiatry, 55*, 745-751. https://doi.org/10.1016/j.biopsych.2003.11.018

Rosmond, R., Dallman, M. F., & Björntorp, P. (1998). Stress-related cortisol secretion in men: relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities. *Journal of Clinical Endocrinology & Metabolism, 83,* 1853-1859. https://doi.org/10.1210/jcem.83.6.4843

Sambhi, M. P., Weil, M. H., & Udhoji, V. N. (1965). Acute pharmacodynamic effects of
glucocorticoids: Cardiac output and related hemodynamic changes in normal subjects
and patients in shock. *Circulation, 31*, 523-530. https://doi.org/10.1161/01.CIR.31.4.523

Sapolsky, R. M. (2004). *Why zebras don't get ulcers* (3rd Ed.)*.* New York: Holt.

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence
stress responses? Integrating permissive, suppressive, stimulatory, and preparative
actions. *Endocrine Reviews, 21*, 55–89. https://doi.org/10.1210/edrv.21.1.0389

Schlotz, W., Hellhammer, J., Schulz, P., & Stone, A. (2004). Perceived work overload and
chronic worrying predict weekend-weekday differences in the cortisol awakening
response. *Psychosomatic Medicine*, *66*, 207–214.
https://doi.org/10.1097/01.psy.0000116715.78238.56

Schonfeld, I. S., & Bianchi, R. (2016). Burnout and depression: two entities or one? *Journal
of Clinical Psychology, 72*, 22-37. https://doi.org/10.1002/jclp.22229

Scrivo, R., Vasile, M., Bartosiewicz, I., & Valesini, G. (2011). Inflammation as "common
soil" of the multifactorial diseases. *Autoimmunity Reviews, 10,* 369-374.
https://doi.org/10.1016/j.autrev.2010.12.006

Seeman, T., Epel, E., Gruenewald, T., Karlamangla, A., & McEwen, B. S. (2010). Socio-
economic differentials in peripheral biology: Cumulative allostatic load. *Annals of the
New York Academy of Sciences, 1186*, 223-239. https://doi.org/10.1111/j.1749-
6632.2009.05341.x.

Silverman, M. N., & Sternberg, E. M. (2012). Glucocorticoid regulation of inflammation and
its functional correlates: From HPA axis to glucocorticoid receptor dysfunction. *Annals*

of the New York Academy of Sciences, *1261*, 55–63. https://doi.org/10.1111/j.1749-6632.2012.06633.x

Singh, R. B., Mengi, S. A., Xu, Y.-J., Arneja, A. S., Dhalla, N. S., Naranjan, D., & Dhalla, S. (2002). Pathogenesis of atherosclerosis: A multifactorial process. *Experimental & Clinical Cardiology, 7*, 40–53. Retrieved from https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2716189/pdf/ecc07040.pdf

Smaldino, P. E. (2017). Models are stupid, and we need more of them. In *Computational Social Psychology* (pp. 311–331). https://doi.org/10.4324/9781315173726

Steptoe, A., & Kivimäki, M. (2013). Stress and cardiovascular disease: An update on current knowledge. *Annual Review of Public Health*, *34*, 337–354. https://doi.org/10.1146/annurev-publhealth-031912-114452

Strauss, B. (2002). Social facilitation in motor tasks: a review of research and theory. *Psychology of Sport and Exercise, 3,* 237-256. https://doi.org/10.1016/S1469-0292(01)00019-X

Taylor, L. A., & Gerrard, J. H. (1977a). Pressure-radius relationships for elastic tubes and their application to arteries: Part 1-Theoretical relationships. *Medical & Biological Engineering & Computing*, *15*, 11–17. https://doi.org/10.1007/BF02441569

Taylor, L. A., & Gerrard, J. H. (1977b). Pressure-radius relationships for elastic tubes and their applications to arteries: Part 2-A comparison of theory and experiment for a rubber tube. *Medical & Biological Engineering & Computing*, *15*, 18–21. https://doi.org/10.1007/BF02441570

Theorell, T., Jood, K., Järvholm, L. S., Vingård, E., Perk, J., Östergren, P. O., & Hall, C. (2016). A systematic review of studies in the contributions of the work environment to ischaemic heart disease development. *The European Journal of Public Health*, *26*, 470–477. https://doi.org/10.1093/eurpub/ckw025

Thorn, L., Hucklebridge, F., Evans, P., & Clow, A. (2006). Suspected non-adherence and weekend versus week day differences in the awakening cortisol response. *Psychoneuroendocrinology*, *31*, 1009–1018. https://doi.org/10.1016/j.psyneuen.2006.05.012

Thosar, S. S., Butler, M. P., & Shea, S. A. (2018). Role of the circadian system in cardiovascular disease. *Journal of Clinical Investigation, 128*, 2157–2167. https://doi.org/10.1172/JCI80590

U.S. Department of Labor. (2008). *Fact sheet #22: Hours worked under the fair labor standards*. Retrieved from https://www.dol.gov/whd/regs/compliance/whdfs22.pdf

Vancouver, J. B., Li, X., Weinhardt, J. M., Steel, P., & Purl, J. D. (2016). Using a Computational Model to Understand Possible Sources of Skews in Distributions of Job Performance. *Personnel Psychology*, *69*, 931–974. https://doi.org/10.1111/peps.12141

Vancouver, J. B., & Purl, J. D. (2017). A computational model of self-efficacy's various effects on performance: Moving the debate forward. *Journal of Applied Psychology*, *102*, 599–616. https://doi.org/10.1037/apl0000177

Vancouver, J. B., Tamanini, K. B., & Yoder, R. J. (2010). Using dynamic computational models to reconnect theory and research: Socialization by the proactive newcomer as

example. *Journal of Management*, *36*(3), 764–793.

https://doi.org/10.1177/0149206308321550

Vroom, V. H. (1964). *Work and Motivation (1st Ed.).* San Francisco: Wiley.

Weinhardt, J. M., & Vancouver, J. B. (2012). Computational models and organizational

psychology: Opportunities abound. *Organizational Psychology Review*, *2*, 267–292.

https://doi.org/10.1177/2041386612450455

Wilhelm, I., Born, J., Kudielka, B. M., Schlotz, W., & Wüst, S. (2007). Is the cortisol

awakening rise a response to awakening? *Psychoneuroendocrinology*, *32*, 358–366.

https://doi.org/10.1016/j.psyneuen.2007.01.008

Whitworth, J. A., Williamson, P. M., Mangos, G., & Kelly, J. J. (2005). Cardiovascular

consequences of cortisol excess. *Vascular Health and Risk Management*. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/17315601

Yarkoni, T., & Westfall, J. (2017). Choosing prediction over explanation in psychology:

Lessons from machine learning. *Perspectives on Psychological Science*, *12*, 1100–1122.

https://doi.org/10.1177/1745691617693393

Zajonc, R. B. (1965). Social facilitation. *Science, 149*, 269–274.

https://doi.org/10.1126/science.149.3681.269

Zhou, D., Kusnecov, A. W., Shurin, M. R., DePaoli, M., & Rabin, B. S. (1993). Exposure to

physical and psychological stressors elevates plasma interleukin 6: relationship to the

activation of hypothalamic-pituitary-adrenal axis. *Endocrinology, 133,* 2523-2530.

https://doi.org/10.1210/en.133.6.2523