TITLE

Serial sarcomere number is substantially decreased within the paretic biceps brachii in

chronic hemiparetic stroke

AUTHORS & AFFILIATIONS:

[1,4,5]Amy N. Adkins, [1,2,3]Julius P.A. Dewald, [1,2,6]Lindsay Garmirian, [1,2,7]Christa M. Nelson,

[1,2,3,4,5]Wendy M. Murray*

[1]Department of Biomedical Engineering, Northwestern University, Evanston, IL, USA

[2]Department of Physical Therapy and Human Movement Sciences, Northwestern University

Feinberg School of Medicine, Chicago, IL, USA

[3]Department of Physical Medicine & Rehabilitation, Northwestern University Feinberg School of

Medicine, Chicago, IL, USA

[4]Shirley Ryan AbilityLab (formerly the RIC), Chicago, IL, USA

[5]Edward Hines, Jr. VA Hospital, Hines, IL, USA

[6]Department of Health, Human Function & Rehabilitation Sciences, George Washington

University, Washington, DC, USA (Current Address)

[7]Department of Physical Therapy and Rehabilitation Science, University of Maryland School of

Medicine, Baltimore, MD, USA (Current Address)

ABSTRACT

Stroke-induced neural deficits substantially alter input to and usage of individual

muscles. We combined novel *in vivo* imaging methods across three scales of muscle to

measure sarcomere length, fascicle length, and volume of the biceps brachii in both limbs of

individuals with chronic hemiparetic stroke and unimpaired controls. We calculated serial

sarcomere number and physiological cross-sectional area. The paretic biceps had fewer serial

sarcomeres compared to the contralateral limb; no interlimb differences were observed in

controls, suggesting the observed differences were muscle adaptations associated with stroke

rather than natural interlimb variability. We provide the first direct evidence of the loss of serial

sarcomeres in human muscle, observed in a population with neural impairments that lead to

disuse and chronically place the affected muscle at a shortened position. This adaptation is

consistent with functional consequences (increased passive resistance to elbow extension) that

would amplify already problematic, neurally driven motor impairments.

INTRODUCTION

Three-fourths of the nearly 7 million stroke survivors currently living in the United States

report substantial motor impairments that limit their independence in tasks of everyday life[1,2].

Damage to the corticofugal motor pathways and the resulting reliance on indirect

corticoreticulospinal pathways following stroke alter neuronal input to contralesional muscles[3-6].

Such changes include decreased voluntary neural drive (weakness or paresis)[7,8] and increased

involuntary neural drive (hypertonicity), causing the inability to fully activate and fully relax the

muscle, respectively. Further, abnormal muscle coactivation patterns[9] also arise from the

altered neuronal drive, commonly leading to abnormal limb synergies[10] or a loss of independent

joint control[9,11]. For example, in the upper limb, the flexion synergy presents when the individual

attempts to abduct their shoulder but involuntarily and concomitantly flexes all of their distal

joints (elbow, wrist, and fingers)[9,12]. The flexion synergy limits the ability to combine shoulder

abduction and extension of the limb, as needed to use the arm and hand to reach and acquire

an object at a distance from the body. Thus, although stroke is primarily an injury of the brain,

stroke-induced neural deficits substantially alter input to and use of the contralesional upper

limb.

Skeletal muscle is a plastic tissue; changes to both the stimulus it receives and how it is

used can alter its functional capacity (e.g.[13-19]). A muscle's function, its ability to contract and

produce force, can be delineated via its architecture[20-23]. Specifically, optimal fascicle length

(OFL) is the fascicle length at which the muscle generates its maximum force. OFL is a

measure of the number of sarcomeres in series in the muscle, so it also provides a measure of

the absolute range of lengths over which the muscle can actively generate force[21,24].

Physiological cross-sectional area (PCSA) is a correlate of the maximum isometric force a

muscle can produce given maximum activation[25]. These two critical muscle architectural

features are calculated from more primary measures of a muscle's structural anatomy that occur

across different scales, including muscle volume, fascicle length, pennation angle (in pennate

muscle), and sarcomere length (see Methods, Eqs. 4 and 5)[24,26]. Quantifying the adaptation of

these two parameters to changes in use and stimulus can provide insight into the functional

implications of such changes. For example, a series of classic studies in animal models

demonstrate that when limbs were immobilized for an extended time, muscles immobilized at a

joint angle that shortened muscle-tendon lengths (i.e., with origin-to-insertion distances that

were decreased compared to resting length) lost serial sarcomeres (i.e., adapted such that OFL

was shorter)[18,27-29]. Conversely, when immobilized at a joint posture that lengthened muscles,

sarcomeres were added in series. These fundamental studies suggest that when *in vivo* length

is chronically altered, a muscle's architecture changes in a way that maximizes its function at

the new length. Specifically, in the immobilization studies, the muscle's length "re-optimized" so

OFL (the length at which the muscle generates its maximum force) occurred at the joint angle of

immobilization.

Despite these important studies, it is unclear the extent to which this mechanism is

expressed, *in vivo*, in human muscle. For example, at this point, only a single published study

has calculated serial sarcomere number (i.e., measured both sarcomere length and fascicle

length *in vivo*) under conditions in which a human muscle has chronically been placed at a

shortened position[32]. In this case, the calf muscles in children with cerebral palsy with equinus

contractures (functionally shortened muscle-tendon units that are more resistant to stretch[30,31])

severe enough to require surgical intervention did not "re-optimize" to the shorter muscle

lengths imposed by the participants' chronically plantarflexed joint posture. Rather, the soleus

muscles in these children had sarcomeres substantially longer (4.07μm[32]) than optimal length

(2.70μm[33]). Unfortunately, identical measurements could not be collected in a control population

because sarcomere length could not be measured in children who were not undergoing surgery.

Sarcomere length measurements in living human subjects have traditionally been limited to

biopsy or intraoperative studies, (i.e. during distraction surgeries for limb discrepancy[13], tendon

transfer surgeries[34,35], or muscle lengthening in children with cerebral palsy[30]) and only recently

have become measurable via minimally or non-invasive techniques[36,37]. As a result, the authors

estimated serial sarcomere number for control participants by combining fascicle length

measures obtained from typically developing children via ultrasound with sarcomere lengths

measured from adult cadavers via dissection. Based on these methods, the authors concluded

the soleus in the children with CP had fewer sarcomeres in series than typically developing

muscle[32]. This pediatric study suggested that, instead of re-optimizing so that OFL occurred at

the chronically plantarflexed ankle angle, the shortened muscle shifted to much longer

sarcomere lengths, at which active force-generating capabilities are weaker and passive forces

are greater.

Stroke-induced neural deficits lead to decreased use and a chronically more flexed

resting posture of the paretic upper limb. Given the muscle adaptation observed following

immobilization in animal models, as well as a general assumption that these type of muscle

adaptations do occur in humans with severe neural impairments and may result in muscles that

are both structurally weaker and stiffer, there is concern about functional consequences of

muscle adaptation following stroke. Thus, several studies have sought to directly quantify *in vivo*

muscle structure in this cohort. In general, *in vivo* medical imaging has provided strong evidence

that many muscle anatomical structural parameters are different in the paretic limb compared to

the contralateral limb. Specifically, shorter fascicles[38], as well as smaller pennation angles[39],

muscle masses[40], volumes[41,42], and anatomical cross-sectional areas (the area of muscle

perpendicular to the line of action of the external tendons[43]) have been demonstrated in

numerous thigh and shank muscles[41,44]. Similarly, in the upper limb, shorter fascicles[45,46] and

smaller muscle volumes[47] have been reported in various paretic muscles as compared to the

contralateral side.

The primary weakness of existing studies that demonstrate differences in anatomical

parameters in paretic muscles in chronic stroke is that none also simultaneously measured

sarcomere length. Neither OFL nor PCSA is calculable in any existing study that describes

muscle structure following chronic stroke because they do not report the corresponding

measures of both a muscle's sarcomere and fascicle lengths. Thus, insight into how the

observed structural differences relate to muscle function in the paretic limb is limited. For

example, shorter fascicles in the paretic limb relative to the contralateral limb have been widely

demonstrated[38,45,46]. One explanation of this observation in the elbow flexors could be that they

experience conditions similar to muscle immobilization at shortened muscle-tendon lengths; the

paretic upper limb is generally used much less than the non-paretic limb and it rests in a more

flexed elbow posture[48]. Thus, similar to immobilization studies in animal models (i.e.[18,27,29]),

shorter fascicles in the paretic elbow flexors could result from a loss of sarcomeres in series.

Functionally, this adaptation would indicate a decrease in the absolute range of lengths over

which the muscle can generate active force[21]. In addition, the animal studies demonstrated that

muscles that lost sarcomeres in series exhibited a shift in the onset of passive force

development to shorter lengths; the muscles that lost sarcomeres in series were also described

as less "extensible"[29]. However, without a concomitant measure of the length of the muscle's

sarcomeres, the possibility that fascicle lengths measured in paretic muscle are shorter because

there are fewer sarcomeres in series cannot be distinguished from the possibility that the paretic

muscle has the same number of sarcomeres in series as in the contralateral limb, but its

sarcomeres have adapted by shifting to shorter lengths. In this case, the paretic muscle would

be capable of generating active force over the same range of lengths, but the muscle would

operate over shorter sarcomere lengths throughout the joint's range of motion. Based on basic

muscle physiology, we would expect a muscle operating at shorter sarcomere lengths to

generate smaller passive forces throughout the joint's range of motion[43]. A single study has

reported *in vivo* measures of biceps brachii sarcomere lengths following chronic hemiparetic

stroke (fascicle lengths were not measured); 2 of 4 stroke participants had longer sarcomeres in

the paretic biceps compared to the non-paretic side, and 2 had shorter sarcomeres[49]. Thus, the

implications for OFL, the range of lengths over which the muscle can generate active force, and

the passive forces the muscle produces over the joint's range of motion, remain unclear. These

distinct possibilities for sarcomere adaptation would also have different effects on maximum

isometric force capacity. While a decrease in muscle volume is regularly reported in paretic

limbs[41,47], PCSA is the architectural correlate of force-generating capacity, and it is calculated

using the ratio between muscle volume and OFL. Critically, stroke alters both neural input[5-7] to

and use[8,9,11,12] of the paretic limb. Thus, it is also possible that paretic muscle adapts in an

unexpected manner post-stroke; animal models of immobilization limited the use of the limb and

altered muscle length but did not involve neural injury.

In this study, we quantify multiscale muscle parameter differences, *in vivo*, in the biceps

brachii of the paretic and contralateral limbs of individuals with chronic hemiparetic stroke and in

both limbs of a group of age-matched individuals with no history of musculoskeletal or

neurological diseases or injuries to the upper limb. Specifically, sarcomere length, fascicle

length, and muscle volume are measured from images obtained using second harmonic

generation microendoscopy, extended field-of-view ultrasound, and fat suppression magnetic

resonance imaging, respectively. Our most prominent finding was that the paretic biceps of

individuals with chronic hemiparetic stroke has fewer sarcomeres in series (i.e., shorter OFL)

compared to the contralateral muscle. In the limb posture we evaluated, this result was

manifested by systematically shorter fascicles in the paretic muscle without significantly different

sarcomere lengths from the non-paretic, contralateral biceps. Our data provide the first direct

evidence of muscle adaptation involving the loss of serial sarcomeres in living human subjects

and is observed in a population with neural impairments that chronically place the affected

muscle at a shortened position. This muscle architectural difference post chronic hemiparetic

stroke is consistent with functional consequences that would amplify the already problematic

neural driven motor impairments (i.e. the ability to reach away from the body to grab an object).

RESULTS

SSN, Fascicle length, and Sarcomere Length

Relative to the contralateral, non-paretic limb, the paretic biceps brachii of individuals

with chronic hemiparetic stroke had fewer sarcomeres in series; this inter-limb difference was

not observed in participants without stroke. Specifically, there was a statistically significant

difference in overall SSN between paretic and non-paretic limbs (**p = 0.0044**) with an estimated

mean SSN in the paretic muscle of 30,152 ± 2,163 (95% CI, 25,038 to 35,267) compared to

38,697 ± 2,162 (95% CI, 33,584 to 43,810) serial sarcomeres in the non-paretic muscle (Fig 1).

We also observed significantly shorter biceps fascicles (**p < 0.0001**) in these participants'

paretic limbs; mean fascicle lengths on the paretic side were shorter (10.66 ± 0.61 cm; 95% CI,

9.46 to 11.85 cm) than non-paretic fascicle lengths (13.59 ± 0.61 cm; 95% CI, 12.40 to 14.79

cm) in the single joint posture we tested (Fig. 2A). There was no significant difference in

sarcomere length in this posture (p = 0.6787; paretic 3.58 ± 0.08 µm 95% CI, 3.42 to 3.73 µm;

non-paretic 3.54 ± 0.08 μm 95% CI, 3.39 to 3.70 μm) (Fig. 2B). For the individuals who did not

have a stroke, there were no observed interlimb differences in SSN (p = 0.2463; dominant

40,102 ± 1,451 95% CI, 37,253 to 42,951; non-dominant 39,545 ± 1,453 95% CI, 36,691 to

42,398), fascicle length (p = 0.0790; dominant 14.32 ± 0.27 cm 95% CI, 13.78 to 14.86 cm;

non-dominant 14.11 ± 0.27 cm 95% CI, 13.57 to 14.65 cm), or sarcomere length (p = 0.9423;

dominant 3.58 ± 0.07 μm 95% CI, 3.44 to 3.72 μm; non-dominant 3.59 ± 0.07 μm 95% CI, 3.45

to 3.73 μm) (Fig 2-3). The relationship between Fugl-Meyer Assessment (FMA) and percent

difference in fascicle length yielded a modest linear relationship (R^2 = 0.5953) which was

significant (**p = 0.0249**) (Fig. 3A); the relationship between FMA and percent difference in SSN

was weaker (R^2 = 0.4676; p = 0.0615) (Fig. 3B).

PCSA and Muscle Volume

There was a substantial difference in muscle volume and a modest difference in PCSA

in the paretic biceps brachii in participants with chronic hemiparetic stroke as compared to the

contralateral limb. Muscle volume was significantly smaller on the paretic side (**p = 0.0127**;

paretic 105 ± 21 cm³ 95% CI, 55 to 154 cm³; non-paretic 148 ± 21 cm³ 95% CI, 99 to 197 cm³);

however, one stroke participant had a larger muscle volume on the paretic side compared to the

non-paretic side (Fig. 4A). This individual also had the smallest volume among all participants'

non-paretic or dominant limbs. On average, post-stroke participants had larger PCSAs in their

non-paretic limb (p = 0.2075; paretic 12.59 ± 1.71 cm² 95% CI, 8.55 to 16.64 cm²; non-paretic

13.72 ± 1.71 cm² 95% CI, 9.66 to 17.76 cm²) (Fig 4B). We did not observe interlimb differences

in either volume (p = 0.7743; dominant 125 ± 40 cm³ 95% CI, -3 to 253 cm³; non-dominant 124

± 40 cm³ 95% CI, -3 to 253 cm³) or PCSA across limbs (p = 0.5561; dominant 11.38 ± 3.45 cm²

95% CI, 4.60 to 18.16 cm²; non-dominant 11.52 ± 3.45 cm² 95% CI, 4.74 to 18.29 cm²) in the

participants who had not had a stroke (Fig 4). In the seven stroke participants with smaller

biceps volumes in the paretic limb, normalized interlimb differences in PCSA were smaller than

those in volume due to fewer sarcomeres in series (Fig 5).

DISCUSSION

This study aimed to quantify *in vivo* differences in muscle architecture parameters

between the paretic and non-paretic biceps brachii of individuals with chronic hemiparetic

stroke. Most notably, this comprehensive, multi-scale study found fewer sarcomeres in series in

the paretic muscle compared to the contralateral side. In 7 of 8 stroke participants, we observed

strikingly smaller muscle volumes on the paretic side. However, the corresponding deficit in

PCSA of the paretic biceps, the architectural parameter that predicts the maximum isometric

force the muscle can generate with full activation[25], was more modest. In these seven stroke

participants, the fact that each had fewer serial sarcomeres partially explains the smaller paretic

muscle volume. We quantified the same architectural parameters in age-range matched

individuals who had not undergone a stroke; we found no substantial or significant interlimb

differences in any muscle architecture parameter, suggesting the interlimb differences we

observed were adaptations associated with chronic stroke rather than natural interlimb

variability. Thus, the architectural parameters suggest a functional re-organization of the

muscle. Specifically, shorter optimal fascicle lengths are generally understood to indicate a

proportional decrease in the width of the muscle's isometric force-length curve[21,24,43], or – more

explicitly – a proportional decrease in the absolute range of lengths over which the muscle can

generate active force. Given this adaptation in length, the loss of muscle volume we observed in

7 of the 8 paretic limbs was not a direct measure of the loss in the muscle's force-generating

capacity (Fig. 5). The substantial decrease in serial sarcomere number that occurs when a

muscle is held at a joint posture which places it at a shortened length was first reported in

classic limb immobilization studies in animal models[18,27,29]; it is widely assumed to be a

fundamental muscle adaptation process. We now provide the first direct evidence of this

phenomenon in living human subjects.

The most complete demonstration of in vivo muscle adaptation that accompanies

chronic length changes has been via animal models involving limb immobilization[18,27-29]. The

main difference between the adaptations observed in our study compared to these studies of

limb immobilization is the magnitude of the observed adaptation. The decrease in SSN we

observed in the paretic biceps post stroke was less substantial than the loss of serial

sarcomeres observed following immobilization at a shortened muscle-tendon length in animal

models (~22% vs ~40%). The prime differences between individuals post-stroke and animal

immobilization studies, which may explain this difference in magnitude, is that post-stroke

individuals tend to disuse their paretic limb, but still have active and passive joint motion

whereas animal immobilization studies eliminated movement entirely. In our stroke participants,

the studied muscle also receives altered neural inputs; the animal studies did not involve a

neural injury.

In human subjects, only a single previous study has ever measured both sarcomere

lengths and fascicle lengths in the same muscle under conditions that are generally assumed to

result in a loss of serial sarcomeres (i.e., in a muscle that has been chronically placed in a

shortened position)[32]. This previous work evaluated the soleus muscle in children with cerebral

palsy who were undergoing surgery to address equinus contractures. However, because of the

invasive methods required to measure sarcomere lengths before the development of second

harmonic generation microendoscopy, measures of sarcomere length were obtained

intraoperatively. Reasonably, these data could not also be collected from typically developing

children. Thus, despite the valuable data obtained from the clinical population, serial sarcomere

number in the chronically shortened soleus muscle could not be compared to direct measures of

the same parameters obtained in a control population. Despite this limitation, the intraoperative

data provided evidence that the sarcomere lengths in the chronically shortened soleus were

extremely long compared to optimal sarcomere length, even in an extremely plantarflexed limb

posture. This finding was surprising because the results from the immobilization studies in

animal models suggest that muscle "re-optimizes" such that optimal length occurs in the limb

position of immobilization. Importantly, this result in the soleus replicated a previous finding in

children with CP undergoing surgery for wrist contracture. In this case, sarcomere lengths for

the wrist flexor in a flexed wrist position were shown to be much longer compared to typically

developing children undergoing surgery to treat radial nerve palsy (3.48μm vs 2.41μm)[50]. Unlike

the results in children with CP, we did not observe systematically longer sarcomeres in the

paretic muscles of individuals with chronic hemiparetic stroke (Fig. 2B). We expect that a

critically important difference between muscle adaptation following stroke and CP is that stroke

occurs in a fully developed system. Similar to the biceps brachii in our population, the affected

muscles in the CP studies were chronically in a shortened positioned due to the primary neural

impairments. However, it has been posited that while chronically shortened CP muscle loses

sarcomeres in series as would be expected from the classic immobilization studies, a

malfunctioning sensing system within the muscles prevents the addition of serial sarcomeres

during bone growth, resulting in the abnormally long sarcomeres[30]. Thus, the additional

physiological process of bone growth in children with CP confounds direct comparison with

muscle adaptation that occurs in adults with stroke.

In general, our work provides the most direct confirmation in humans to date that chronic

impairments that lead to disuse and place a muscle in a shortened position are associated with

the loss of serial sarcomeres. In the context of the literature discussed above, we note that this

adaptation process also seems to be moderated by the presence of confounding factors (i.e.

bone growth (CP), altered neural input, disuse resulting in a reduced use of available range of

motion, etc.). Such confounds are common *in vivo* following disease, neural injury, and clinical

interventions such as surgery or immobilization. With the comprehensive multiscale imaging

techniques utilized in this study, adaptation of muscle structure in the context of confounding

factors can now be better explored and more fully understood in humans, which is necessary for

the development of more targeted interventions that seek to improve outcomes.

The substantial decrease in serial sarcomere number found in the paretic biceps brachii

likely amplifies motor impairments which stem from stroke-induced neural impairment.

Decreased voluntary neural drive (weakness or paresis)[7,8], increased involuntary neural drive

(hypertonicity), and abnormal muscle coactivation patterns[9] occur following damage of the

corticofugal motor pathways due to a stroke. In addition, it is clear that the stroke induced neural

deficits result in altered use of the contralesional or paretic limbs. Particularly notable is the

difficulty (and in severe cases impossibility) of coordinated extension of the upper limb to reach

and grab an object which is some distance from the body[12,51]. In this study we find that the

muscle structure itself has been altered via a loss of serial sarcomeres in the paretic biceps

brachii muscle. Functionally, for the individuals in our study who had survived a stroke, this

means that the paretic muscle has a narrower range of lengths and joint angles over which it

can produce active force. Reduction in serial sarcomeres resulted in an increase in the muscle's

passive resistance to stretch in animal models of limb immobilization[29]. For the biceps, this

would exacerbate neurally-driven motor impairments that diminish the ability to extend the

elbow, in the presence of significant elbow extensor weakness, to reach for an object away from

their body.

Results from our study are reasonable in the context of previously reported in vivo

studies which independently measure either fascicle length, muscle volume, or sarcomere

length. Specifically, muscle fascicle lengths in the paretic biceps brachii in this study were on

average 20.6% shorter than the non-paretic side. Previous studies performed in elbow flexors

report similar, substantial decreases in fascicle length in extended joint postures (18.6%

decrease in biceps brachii at 25⁰ elbow flexion [46],15% decrease in brachialis at 10⁰ elbow

flexion[45]). Studies of muscle volume differences between paretic and non-paretic limbs are

variable. On average, our muscle volume differences (29%) are in the same direction (paretic

muscle is smaller) and of slightly greater magnitude than other upper limb studies (no difference

to 25% difference[40,47]) and lie within the range of lower limb difference (no difference to

33%[35,49]). The exclusion of intramuscular fat in our measure of muscle volume is relatively novel

and may explain our slightly higher percent differences. Measurements of biceps brachii

sarcomere lengths have only been obtained in vivo in a single study that enrolled 4 individuals

with chronic hemi-paretic stroke. Although there was not a significant difference in the mean

sarcomere length between limbs among the stroke participants we studied, similar to this

previous work, we found that some individuals had shorter sarcomere lengths in their paretic

limb while others had longer[49]. Notably, independently measured muscle anatomical parameters

(particularly sarcomere lengths and non-normalized fascicle lengths) are difficult to directly

compare between studies as they are sensitive to the limb posture chosen for testing[52]; muscle

fiber and sarcomere lengths change with joint position.

There are various limitations to this study. While it provides novel *in vivo* evidence of

comprehensive muscle architecture changes following chronic hemiparetic stroke, more

participants, a larger number of muscles studied in more joint positions, and inclusion of factors

that quantify the extent participants use their upper limbs (i.e. passive and active range of

motion, elbow joint resting posture, amount of voluntary use of limb, etc.) would broaden our

knowledge of muscle adaptation post brain injury. We would expect that altered tendon

properties or increased tendon length[53] could accompany the paretic muscle adaptation we

observed, but we did not include these measures in this study. This study is limited in that we

did not directly measure stiffness or changes in muscle or tendon properties which may further

elucidate muscle-tendon adaptation post-stroke.

Beyond the addition of the first comprehensive measurements of in vivo muscle

architecture for the investigation of muscle plasticity to stroke-induced neural deficits, this study

demonstrates the need for such comprehensive in vivo studies and provides insight for the

design of therapeutic interventions for stroke survivors. Without the combination of fascicle and

sarcomere lengths measured at the same joint angle with quantification of muscle volume,

ambiguity would remain in the functional impact of these individual muscle parameters. Many

prior studies in stroke and other populations demonstrate interlimb differences in fascicle length

without normalizing to sarcomere length. With the addition of sarcomere length, we were able to

explicitly demonstrate that the paretic biceps brachii muscle has fewer serial sarcomeres. Our

finding leads us to conclude that stroke-induced neural deficits, which lead to altered input and

disuse of the contralesional limb, ultimately change the basic biceps brachii muscle architectural

parameters in a way which may amplify functional impairments.

METHODS

Participants

Measurements of sarcomere length, fascicle length, and muscle volume of the biceps

brachii were obtained using in vivo imaging techniques in both arms of twelve participants; eight

participants with chronic hemiparetic stroke (3 female/5 male, 60 ± 9 yrs, Fugl-Meyer 26 ± 9, 13

± 10 yrs post-stroke) and four participants with no history of musculoskeletal or neurological

diseases or injuries to the upper limb (2 female/2 male, 62 ± 6 yrs). Fugl-Meyer Assessment

scores reported in this study were performed by a licensed physical therapist prior to

experimentation. All the individuals who participated in this study provided informed consent

prior to experimentation; Northwestern University's Institutional Review Board approved this

study's procedures.

Sarcomere length

Sarcomere lengths of both arms were acquired with the participants seated in a comfortable

chair. The arm being imaged was secured at 85⁰ shoulder adduction, 10⁰ horizontal shoulder

flexion, 25⁰ of elbow flexion, mid pro-supination, and 0⁰ wrist and finger flexion (Fig. 6). Joint

posture was verified by goniometric measurement. A microendoscopic system (Zebrascope,

Zebra Medical Technologies, Palo Alto, CA) which consists of a laser (class IV, output power >

500mW, center wavelength 1030 nm), microscope, and microendoscopic probe, was used to

image sarcomeres in vivo. This system utilizes the second-harmonic generation optical

technique to capture the intrinsic striation pattern of sarcomeres[36,49].

A sterile microendoscopic probe was inserted into the long head of the biceps brachii. The

probe consisted of two 1.8cm long, 20-gauge needles with beveled tips (Fig. 6B); one needle

with a transmitting lens used to excite the muscle tissue and one needle housing a receiving

lens to capture the reflection of the signal after it has interacted with tissue. Ultrasound and

palpation techniques were used, prior to insertion, to verify placement of the probe at mid-belly

of the muscle with the probe's optical lenses aligned parallel to the fascicle direction. A spring-

loaded injector was used to rapidly insert the probe, minimizing pain and improving precision of

probe placement. The microscope was attached to the microendoscopic needle for imaging.

Images with a field-of-view of 82μm by 82μm were collected at 1.9Hz for approximately 2-5

minutes. The image produced from the microendoscopic system captures the A-bands (myosin

protein) of the sarcomeres and enables direct measurement of sarcomeres from the resulting

striation pattern[49]. Surface EMG (Bagnoli-16 system, Delsys Inc., Boston, MA) of the biceps

brachii was obtained simultaneously using a custom written MATLAB script. Baseline EMG

activity was collected for 10 seconds with the needle inserted and participant relaxed. Muscle

activation was visually monitored during data collection and analysis was performed offline.

Fascicle length

Fascicle length measurements of the long head of the biceps brachii in both arms of all

participants were obtained using extended field-of-view ultrasound (EFOV-US) under the same

conditions (same joint posture, passive muscle) as sarcomere length measures. The extended

field-of-view technique involves sweeping the ultrasound probe along the length of the muscle

as sequential B-mode ultrasound images are acquired and stitched together to form a single

composite image with a field-of-view longer than the ultrasound probe's aperture (±60 cm)[54].

This method has been demonstrated to be accurate and reliable for measurement of fascicle

length in different individuals and muscles[52,55]. Approximately 10 qualitatively good images were

captured per arm. EFOV-US images (Acuson S2000, linear array transducer 18L6, SieScape,

Siemens Medical Solutions USA, Inc., Mountain View, CA) and surface EMG (Bagnoli-16

system, Delsys Inc., Boston, MA) of the biceps brachii were simultaneously recorded (Spike,

Power1401, and Micro1401-3, Cambridge Electronic Design Limited, Cambridge, England).

Muscle Volume

To determine the volume of the biceps brachii muscle, excluding intramuscular fat, the

Dixon method, a fat suppression MRI sequence, was implemented on both upper limbs of all

participants (3D GRE, TR = 7ms, flip angle = 12°, matrix size = 256 x 304, slice thickness =

3mm, TE of 2.45ms and 3.68ms)[47]. As increases in intramuscular fat within muscle has been

demonstrated in the lower limb of stroke participants[41] and patients with other pathologies (i.e

whiplash[56], spinal cord injury[57]) correcting muscle volumes for the amount of intramuscular fat is

necessary to avoid potential overestimation of volume of muscle. The participants were lying

supine in a 3T MRI (Area, Siemens Medical Solutions USA, Inc., Mountain View, CA) scanner

with their arm as close to the center of the scanner as possible. To minimize participant

movement during scanning, the lower arm was splinted using an orthosis.

Data analysis

Sarcomere length

Image sequences obtained from the microendoscopic system were analyzed to

determine mean sarcomere length offline using a custom MATLAB script. The raw image

sequence was combined into a multipage Tiff prior to processing. A Fast Fourier Transform

(FFT) was used and the edges and vertical center which contained only noise were cleared.

The FFT was then symmetrically, low pass Gaussian filtered. Frequency bounds were set to

filter out all frequencies which would yield unphysiologic sarcomere length values, specifically,

values smaller than 2μm or larger than 5μm. Images without sarcomeres (without frequencies in

the 2-5 μm range) were removed from the multipage Tiff sequence. White noise in the FFT was

calculated and removed. To determine mean sarcomere length in the image, the peak

frequency of a least squares fit of a Gaussian was calculated for each image. Mean sarcomere

length and standard deviation were calculated across all analyzed images for each arm and

each participant. Any images collected when the biceps EMG signal was 3 standard deviations

above the resting baseline EMG were excluded.

Fascicle Length

EFOV-US images were exported as DICOM images and measurements of fascicle

length were made offline using the segmented line tool in ImageJ (ImageJ with Fiji, version

1.51h, Wayne Rasband, National Institutes of Health, Bethesda, MD[58]). Measurements were

made on 3 images per arm per participant. The experimenter selected the 3 images which best

captured the entire muscle and had visible fascicles which extended from central tendon to

aponeurosis. Four fascicles were measured per image. Mean fascicle length was calculated

across the 3 images (3 images X 4 fascicles per image = 12 fascicles)[46,52].

Muscle Volume

Manual segmentation of the biceps brachii (long and short head) muscle was performed

using Analyze12.0 (AnalyzeDirect, Overland Park, KS). To calculate the volume of muscle

without intramuscular fat (V_{m-f}), the following equations were implemented:

$$(1) \% Fat = \frac{intensity\ of\ fat}{intensity\ of\ water\ +\ intensity\ of\ fat}$$

$$(2) V_{intramuscular \atop fat} = V_{total} * \% Fat$$

$$(3) V_{m-f} = V_{total} - V_{intramuscular \atop fat}$$

Where V_{total} is the total volume segmented from the MRI images. Measurements of biceps

muscle volume without fat were made by two different raters (rater 1 n = 7, rater 2 n=5).

Segmentation and calculation of muscle volume without intramuscular fat (V_{m-f}) has been

shown to be reliable within and across raters[47].

Calculation of Functional Parameters

With the quantification of sarcomere length, fascicle length, and muscle volume, optimal

fascicle length (OFL), serial sarcomere number (SSN), and physiological cross-sectional area

(PCSA), were calculated for both arms of all participants using the following equations.

$$(4) SSN = \frac{l^F}{l^S}$$

$$(5) OFL = SSN * l_o^S$$

$$(6) \ PCSA = \frac{V_{m-f}}{OFL}$$

Where $l_s^o \ is \ 2.7 \mu m$ or optimal sarcomere length[33], and l^F (fascicle length), l^S (sarcomere length),

and V_{m-f} (volume of muscle without fat infiltration) were measured in this study. For statistical

analysis involving equations (4-6), average fascicle length and all measures of sarcomere

length were utilized.

Statistical Analysis

To determine if there were significant interlimb differences in any of the muscle

architecture parameters or calculations, generalized linear mixed-effects models were

implemented (SAS 9.4, SAS Institute Inc., Cary, NC). Each model had one of the parameters or

calculations as the outcome variable. Whether or not the participant had a stroke was a fixed

effect in the model. Within-subject correlation and the correlation between paretic and non-

paretic (or dominant non-dominant) arms were modeled as random effects. A significant

difference between limbs was present if the p-value was less than 0.05 for all models. To

determine if there is a linear relationship between the percent difference in SSN or OFL in the

participants studied with stroke and their clinical function score (Fugl-Meyer assessment), a

linear regression was performed.

An a priori power analysis was conducted to determine sufficient sample size to test the

hypothesis that participants with stroke have interlimb differences in optimal fascicle length

which are not present in individuals who have not had a stroke. The analysis indicated that with

8 participants with stroke and 4 participants without stroke and an effect size greater of 1.85, a

power greater than 0.8 would be achieved. This effect size was established from interlimb

differences in fascicle and sarcomere length obtained from two previous studies performed on

different sets of individuals[46,49]. The effect size from our data was verified to exceed the effect

size from the a priori analysis.

<u>REFERENCES</u>

1   Benjamin, E. J., Muntner, P. & Bittencourt, M. S. Heart disease and stroke statistics-

2019 update: a report from the American Heart Association. *Circulation* **139**, e56-e528

(2019).

2   Faria-Fortini, I., Michaelsen, S. M., Cassiano, J. G. & Teixeira-Salmela, L. F. Upper

extremity function in stroke subjects: relationships between the international

classification of functioning, disability, and health domains. *Journal of Hand Therapy* **24**,

257-265 (2011).

3   Ellis, M. D., Drogos, J., Carmona, C., Keller, T. & Dewald, J. P. Neck rotation modulates

flexion synergy torques, indicating an ipsilateral reticulospinal source for impairment in

stroke. *Journal of neurophysiology* **108**, 3096-3104 (2012).

4   Zaaimi, B., Edgley, S. A., Soteropoulos, D. S. & Baker, S. N. Changes in descending

motor pathway connectivity after corticospinal tract lesion in macaque monkey. *Brain*

135, 2277-2289 (2012).

5   Karbasforoushan, H., Cohen-Adad, J. & Dewald, J. P. A. Brainstem and spinal cord MRI

identifies altered sensorimotor pathways post-stroke. *Nature Communications* **10**, 3524,

doi:10.1038/s41467-019-11244-3 (2019).

6   McPherson, J. G. *et al.* Progressive recruitment of contralesional cortico-reticulospinal

pathways drives motor impairment post stroke. *The Journal of Physiology* **596**, 1211-

1225, doi:10.1113/jp274968 (2018).

7   Bourbonnais, D. & Noven, S. V. Weakness in patients with hemiparesis. *American*

Journal of Occupational Therapy **43**, 313-319 (1989).

8   Carin-Levy, G. *et al.* Longitudinal changes in muscle strength and mass after acute

stroke. *Cerebrovascular Diseases* **21**, 201-207 (2006).

9   Dewald, J. P., Pope, P. S., Given, J. D., Buchanan, T. S. & Rymer, W. Z. Abnormal

muscle coactivation patterns during isometric torque generation at the elbow and

shoulder in hemiparetic subjects. *Brain* **118**, 495-510 (1995).

10   Brunnstrom, S. *Movement Therapy in Hemiplegia: A Neurophysiological Approach.* 1st

edn, (Harper & Row, 1970).

11   Miller, L. C. & Dewald, J. P. A. Involuntary paretic wrist/finger flexion forces and EMG

increase with shoulder abduction load in individuals with chronic stroke. *Clinical*

Neurophysiology **123**, 1216-1225, doi:https://doi.org/10.1016/j.clinph.2012.01.009

(2012).

12   Sukal, T. M., Ellis, M. D. & Dewald, J. P. Shoulder abduction-induced reductions in

reaching work area following hemiparetic stroke: neuroscientific implications.

Experimental brain research **183**, 215-223 (2007).

13 Boakes, J. L., Foran, J., Ward, S. R. & Lieber, R. L. CASE REPORT: Muscle Adaptation

by Serial Sarcomere Addition 1 Year after Femoral Lengthening. *Clinical orthopaedics*

and related research **456**, 250-253 (2007).

14 Csapo, R., Maganaris, C., Seynnes, O. & Narici, M. On muscle, tendon and high heels.

Journal of Experimental Biology **213**, 2582-2588 (2010).

15 Eisenberg, B. R., Brown, J. M. & Salmons, S. Restoration of fast muscle characteristics

following cessation of chronic stimulation. *Cell and tissue research* **238**, 221-230 (1984).

16 Kernell, D., Donselaar, Y. & Eerbeek, O. Effects of physiological amounts of high-and

low-rate chronic stimulation on fast-twitch muscle of the cat hindlimb. II. Endurance-

related properties. *Journal of neurophysiology* **58**, 614-627 (1987).

17 Kernell, D., Eerbeek, O., Verhey, B. & Donselaar, Y. Effects of physiological amounts of

high-and low-rate chronic stimulation on fast-twitch muscle of the cat hindlimb. I. Speed-

and force-related properties. *Journal of neurophysiology* **58**, 598-613 (1987).

18 Williams, P. E. & Goldspink, G. The effect of immobilization on the longitudinal growth of

striated muscle fibres. *Journal Of Anatomy* **116**, 45 (1973).

19 Thomason, D., Herrick, R., Surdyka, D. & Baldwin, K. Time course of soleus muscle

myosin expression during hindlimb suspension and recovery. *Journal of applied*

physiology **63**, 130-137 (1987).

20 Gans, C. & Bock, W. J. The functional significance of muscle architecture: a theoretical

analysis. . *Adv Anat Embryol Cell Biol* **38**, 115-142 (1965).

21 Lieber, R. L. & Fridén, J. Functional and clinical significance of skeletal muscle

architecture. *Muscle & Nerve: Official Journal of the American Association of*

Electrodiagnostic Medicine **23**, 1647-1666 (2000).

22 Gans, C. Fiber architecture and muscle function. *Exercise and Sports Sciences Reviews*

10, 160-207 (1982).

23   Sacks, R. D. & Roy, R. R. Architecture of the hind limb muscles of cats: Functional

significance. *Journal of Morphology* **173**, 185-195, doi:10.1002/jmor.1051730206 (1982).

24   Zajac, F. E. Muscle and tendon: properties, models, scaling, and application to

biomechanics and motor control. *Critical reviews in biomedical engineering* **17**, 359-411

(1989).

25   Powell, P. L., Roy, R. R., Kanim, P., Bello, M. A. & Edgerton, V. R. Predictability of

skeletal muscle tension from architectural determinations in guinea pig hindlimbs.

Journal of Applied Physiology **57**, 1715-1721 (1984).

26   Murray, W. M., Buchanan, T. S. & Delp, S. L. The isometric functional capacity of

muscles that cross the elbow. *J. Biomech.* **33**, 943-952 (2000).

27   Williams, P. E. & Goldspink, G. Changes in sarcomere length and physiological

properties in immobilized muscle. *Journal of Anatomy* **127**, 459-468 (1978).

28   Williams, P. E. & Goldspink, G. Longitudinal growth of striated muscle fibres. *Journal of

cell science* **9**, 751-767 (1971).

29   Tabary, J., Tabary, C., Tardieu, C., Tardieu, G. & Goldspink, G. Physiological and

structural changes in the cat's soleus muscle due to immobilization at different lengths

by plaster casts. *The Journal of physiology* **224**, 231 (1972).

30   Lieber, R. L. & Fridén, J. Muscle contracture and passive mechanics in cerebral palsy.

Journal of Applied Physiology **126**, 1492-1501, doi:10.1152/japplphysiol.00278.2018

(2019).

31   Mathewson, M. A. & Lieber, R. L. Pathophysiology of muscle contractures in cerebral

palsy. *Physical Medicine and Rehabilitation Clinics* **26**, 57-67 (2015).

32   Mathewson, M. A., Ward, S. R., Chambers, H. G. & Lieber, R. L. High resolution muscle

measurements provide insights into equinus contractures in patients with cerebral palsy.

Journal of Orthopaedic Research **33**, 33-39, doi:10.1002/jor.22728 (2015).

33 Gordon, A., Huxley, A. F. & Julian, F. The variation in isometric tension with sarcomere
length in vertebrate muscle fibres. *The Journal of physiology* **184**, 170-192 (1966).

34 Lieber, R. L., Murray, W. M., Clark, D. L., Hentz, V. R. & Fridén, J. Biomechanical
properties of the brachioradialis muscle: Implications for surgical tendon transfer. *The*
Journal of Hand Surgery **30**, 273-282, doi:http://dx.doi.org/10.1016/j.jhsa.2004.10.003
(2005).

35 Murray, W. M., Hentz, V. R., Fridén, J. & Lieber, R. L. Variability in Surgical Technique
for Brachioradialis Tendon Transfer: Evidence and Implications. *JBJS* **88**, 2009-2016,
doi:10.2106/jbjs.e.00973 (2006).

36 Llewellyn, M. E., Barretto, R. P. J., Delp, S. L. & Schnitzer, M. J. Minimally invasive high-
speed imaging of sarcomere contractile dynamics in mice and humans. *Nature* **454**, 784-
788, doi:10.1038/nature07104 (2008).

37 Sanchez, Gabriel N. *et al.* In Vivo Imaging of Human Sarcomere Twitch Dynamics in
Individual Motor Units. *Neuron* **88**, 1109-1120,
doi:http://dx.doi.org/10.1016/j.neuron.2015.11.022 (2015).

38 Gao, F. & Zhang, L.-Q. Altered contractile properties of the gastrocnemius muscle
poststroke. *Journal of Applied Physiology* **105**, 1802-1808,
doi:10.1152/japplphysiol.90930.2008 (2008).

39 Ramsay, J. W., Buchanan, T. S. & Higginson, J. S. Differences in Plantar Flexor
Fascicle Length and Pennation Angle between Healthy and Poststroke Individuals and
Implications for Poststroke Plantar Flexor Force Contributions. *Stroke Res Treat* **2014**,
919486-919486, doi:10.1155/2014/919486 (2014).

40 English, C., McLennan, H., Thoirs, K., Coates, A. & Bernhardt, J. Loss of Skeletal
Muscle Mass after Stroke: a Systematic Review. *International Journal of Stroke* **5**, 395-
402, doi:10.1111/j.1747-4949.2010.00467.x (2010).

41   Ramsay, J. W., Barrance, P. J., Buchanan, T. S. & Higginson, J. S. Paretic muscle

atrophy and non-contractile tissue content in individual muscles of the post-stroke lower

extremity.   *Journal   of   biomechanics*   **44**,   2741-2746,

doi:https://doi.org/10.1016/j.jbiomech.2011.09.001 (2011).

42   Metoki, N., Sato, Y., Satoh, K., Okumura, K. & Iwamoto, J. Muscular atrophy in the

hemiplegic thigh in patients after stroke. *American journal of physical medicine &*

rehabilitation **82**, 862-865 (2003).

43   Zajac, F. E. How musculotendon architecture and joint geometry affect the capacity of

muscles to move and exert force on objects: a review with application to arm and

forearm tendon transfer design. *The Journal of hand surgery* **17**, 799-804 (1992).

44   Ryan, A. S., Dobrovolny, C. L., Smith, G. V., Silver, K. H. & Macko, R. F. Hemiparetic

muscle atrophy and increased intramuscular fat in stroke patients. *Archives of physical*

medicine and rehabilitation **83**, 1703-1707 (2002).

45   Li, L., Tong, K. Y. & Hu, X. The Effect of Poststroke Impairments on Brachialis Muscle

Architecture as Measured by Ultrasound. *Arch Phys Med Rehabil* **88**, 243-250 (2007).

46   Nelson, C. M., Murray, W. M. & Dewald, J. P. A. Motor Impairment–Related Alterations

in Biceps and Triceps Brachii Fascicle Lengths in Chronic Hemiparetic Stroke.

Neurorehabilitation and Neural Repair **32**, 799-809, doi:10.1177/1545968318792618

(2018).

47   Garmirian, L. R. P., Acosta, A. M., Schmid, R. & Dewald, J. P. A. Volume and

intramuscular fat content of upper extremity muscles in individuals with chronic

hemiparetic stroke. *bioRxiv*, 687699, doi:10.1101/687699 (2019).

48   Bhadane, M. Y., Gao, F., Francisco, G. E., Zhou, P. & Li, S. Correlation of resting elbow

angle with spasticity in chronic stroke survivors. *Frontiers in neurology* **6**, 183 (2015).

49   Sanchez, Gabriel N. *et al.* In Vivo Imaging of Human Sarcomere Twitch Dynamics in

Individual Motor Units. *Neuron* **88**, 1109-1120, doi:10.1016/j.neuron.2015.11.022 (2015).

50   Lieber, R. L. & Fridén, J. Spasticity causes a fundamental rearrangement of muscle–joint
609   interaction. *Muscle & Nerve* **25**, 265-270, doi:10.1002/mus.10036 (2002).

610   51   Lan, Y., Yao, J. & Dewald, J. P. A. The Impact of Shoulder Abduction Loading on
611   Volitional Hand Opening and Grasping in Chronic Hemiparetic Stroke.
612   *Neurorehabilitation and neural repair* **31**, 521-529, doi:10.1177/1545968317697033
613   (2017).

614   52   Nelson, C. M., Dewald, J. P. A. & Murray, W. M. In vivo measurements of biceps brachii
615   and triceps brachii fascicle lengths using extended field-of-view ultrasound. *Journal of*
616   *biomechanics* **49**, 1948-1952, doi:doi:10.1016/j.jbiomech.2016.03.040 (2016).

617   53   Takahashi, M., Ward, S. R., Marchuk, L. L., Frank, C. B. & Lieber, R. L. Asynchronous
618   muscle and tendon adaptation after surgical tensioning procedures. *J Bone Joint Surg*
619   *Am* **92**, 664-674 (2010).

620   54   Weng, L. *et al.* US extended-field-of-view imaging technology. *Radiology* **203**, 877-880
621   (1997).

622   55   Adkins, A. N., Franks, P. W. & Murray, W. M. Demonstration of extended field-of-view
623   ultrasound's potential to increase the pool of muscles for which in vivo fascicle length is
624   measurable.   *Journal   of   biomechanics*   **63**,   179-185,
625   doi:https://doi.org/10.1016/j.jbiomech.2017.08.012 (2017).

626   56   Elliott, J. M. *et al.* Magnetic Resonance Imaging Findings of Fatty Infiltrate in the Cervical
627   Flexors in Chronic Whiplash. *Spine* **35**, 948-954, doi:10.1097/BRS.0b013e3181bb0e55
628   (2010).

629   57   Elder, C. P., Apple, D. F., Bickel, C. S., Meyer, R. A. & Dudley, G. A. Intramuscular fat
630   and glucose tolerance after spinal cord injury – a cross-sectional study. *Spinal Cord* **42**,
631   711-716, doi:10.1038/sj.sc.3101652 (2004).

632   58   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
633   *methods* **9**, 676-682 (2012).

AKNOWLDGEMENTS

We would like to thank the study participants, Vikram Darbhe for assistance in data collection,

and Dr. Masha Kocherginsky and Liqi Chen for aid with statistical analysis. We would also like

to thank Sabeen Adamani for assistance in equipment set up and troubleshooting and Zebra

Medical Technologies (now Enspectra Health) for their support with data collection and image

processing.

AUTHOR CONTRIBUTIONS

ANA, WMM, and JPAD contributed to the conception, design, and interpretation of the data and

have drafted or substantially revised the work. ANA was responsible for the acquisition and

analysis of the sarcomere length data and contributed to the acquisition and analysis of all other

data obtained in this study. CMN contributed to the design and analysis of the fascicle length

data obtained via extended field-of-view ultrasound. LG contributed to the design, acquisition,

analysis and interpretation of the MRI data. All authors have approved the submission of this

work.

COMPETING INTEREST STATMENT

The authors have no competing interests to report.

FUNDING

This work is supported by the National Science Foundation Graduate Research Fellowship

Program under Grant No. DGE-1324585, as well as NIH R01 D084009. Any opinions, findings,

and conclusions or recommendations expressed in this material are those of the authors and do

not necessarily reflect the views of the NSF or NIH.

FIGURE LEGENDS (limit 8 fig + tables)

Figure 1: Serial Sarcomere Number/ Optimal Fascicle Length. Data showing interlimb

differences in serial sarcomere number and, proportionally, optimal fascicle length for all

participants who had a stroke and the age-range matched controls (no stroke). Each participant

is represented by a different shape/color and each individual's limbs are connected by a line.

Black circular points and error bars which are offset from individual participant data, represent

mean and standard deviations estimated from the generalized mixed effects model. The star (*)

indicates a significant interlimb difference ($p<0.05$).

Figure 2: Fascicle and Sarcomere Length. Graphs displaying mean sarcomere length (A) and

fascicle length (B) measurements from both limbs and in all participants in the stroke and

control (no-stroke) group. Each participant is represented by a different shape and or color.

Each individual's limbs are connected by a solid line with the exception of one healthy individual

whom had the same average sarcomere length on both limbs as another healthy individual

(dashed line to enable visualization of both healthy individuals). Black circular points and error

bars which are offset from individual participant data, represent mean and standard deviations

estimated from the generalized mixed effects model. The star (*) indicates a significant interlimb

difference ($p<0.05$).

Figure 3: Relationship between change in muscle parameters and clinical assessment.

Graphs showing the relationship between percent difference in serial sarcomere number (A)

and percent difference in fascicle length (B) versus the Fugl-Meyer Assessment clinical

impairment score. There is a trend toward a greater difference in OFL as the impairment level

increases (Fugl-Meyer becomes smaller number).

Figure 4: **Muscle Volume and PCSA.** Data for muscle volume with intramuscular fat removed (A) and calculated PCSA (B) for both biceps braciii of all participants who had a stroke and the age matched controls (no stroke). Each participant is represented by a different shape and or color and each individual's limbs are connected by a solid line. Black circular points and error bars which are offset from individual participant data, represent mean and standard deviations estimated from the generalized mixed effects model. The star (*) indicates a significant interlimb difference (p<0.05).

Figure 5: Interlimb Muscle Differences Post-Stroke. Bar graph showing the percent difference in muscle architectural (sarcomere length "SL" and muscle volume "MV") and functional parameters (serial sarcomere number "SSN" and physiological cross sectional area "PCSA") estimated from the general linear mixed-effects model for all stroke participants who had smaller muscle volume on the paretic biceps. A positive percent difference indicates that the paretic parameter is smaller than the non-paretic side. Error bars represent one standard deviation from the mean of the percent difference across the subjects.

Figure 6: Illustration of experimental set up for muscle architecture measures. **A.** Arm posture used for measurement of fascicle length using ultrasound (right) and for sarcomere length (see B). The white dashed line overlaid on the ultrasound image demonstrates a muscle fascicle. **B.** The illustration shows the Zebrascope which utilizes second harmonic generation microendoscopy to capture the natural striation pattern of sarcomeres in vivo. The Biodex chair and arm fixture (middle) used for measurement of sarcomere and fascicle lengths (see A). The microscope is blown up on the left (Blue dotted box) to show the flow of laser light through the microscope (orange arrows). Below the microscope is a graphic of the probe which is inserted into the bicep brachii muscle. Between the two needles of the probe at 1.5cm the laser light interacts with myofibrils and the striation pattern of the sarcomeres can be captured. To the

right, a raw image which would be seen during image collection is shown. After post-processing,

sarcomere length is measured from the processed image which is showing the length of 10

sarcomeres in series (white line). **C.** Participant in the supine positioning (left) in the MRI bore.

Orange inset shows the splinting of the arm to reduce artifacts due to hand and arm movement.

(Right) 3D rendering of the biceps brachii muscle and humerus bone with a single MRI slice

superimposed.

Table 1: Demographic information of individuals with stroke (1-8) and healthy age-range

matched controls (9-12) who participated in this study. Age and year post-stroke are reported as

of the time at which the experimental data collection for that participant was completed.

FIGURES AND TABLES

Figure 1:

Figure 2

Figure 3

734   Figure 4

736   Figure 5

738   Figure 6

740 Table 1

	Gender	Age	Paretic	Years Post Stroke	FMA
1	F	65	R	32	25
2	F	65	R	11	15
3	F	63	L	15	24
4	M	48	L	8	18
5	M	60	R	7	15
6	M	72	R	22	38
7	M	62	L	5	47
8	M	44	R	5	39
	Gender	Age	Dominant		
9	F	53	R		
10	F	64	R		
11	M	62	R		
12	M	67	R		