Lipogenesis and innate immunity in hepatocellular carcinoma cells reprogrammed by an isoenzyme switch of hexokinases

Laure Perrin-Cocon[1],†, Pierre-Olivier Vidalain[1],†, Clémence Jacquemin[1], Anne Aublin-Gex[1], Keedrian Olmstead[2], Baptiste Panthu[1,3], Gilles J. P. Rautureau[4], Patrice André[1], Piotr Nyczka[5], Marc-Thorsten Hütt[5], Nivea Amoedo[6], Rodrigue Rossignol[6,7], Fabian Volker Filipp[2,8], Vincent Lotteau[1],*‡ Olivier Diaz[1],*‡

1- CIRI, Centre International de Recherche en Infectiologie, Univ Lyon, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, 21 Avenue Tony Garnier, F-69007, Lyon, France.
2- Cancer Systems Biology, Institute of Computational Biology, Helmholtz Zentrum München, Ingolstädter Landstraße 1, D-85764, München, Germany.
3- Actual address: Univ Lyon, CarMeN Laboratory, Inserm, INRA, INSA Lyon, Université Claude Bernard Lyon 1, F-69921, Oullins Cedex, France.
4- Université de Lyon, CNRS, Université Claude Bernard Lyon 1, ENS de Lyon, Centre de RMN à Très Hauts Champs (CRMN), FRE 2034, F-69100 Villeurbanne, France.
5- Department of Life Sciences and Chemistry, Jacobs University, Bremen, Germany.
6- CELLOMET, Centre de Génomique Fonctionnelle de Bordeaux, 146 Rue Léo Saignat, F-33000 Bordeaux, France.
7- Univ. Bordeaux, Inserm, MRGM, U1211, F-33076, Bordeaux, France.
8- School of Life Sciences Weihenstephan, Technical University München, Maximus-von-Imhof-Forum 3, D-85354, Freising, Germany.

† These authors contributed equally to this work
‡These authors jointly supervised this work

* Corresponding authors:
vincent.lotteau@inserm.fr (ORCID: 0000-0003-0997-3282)
olivier.diaz@inserm.fr (ORCID: 0000-0002-3451-9914)

Keywords
Hepatocellular carcinoma, hexokinase, lipogenesis, innate immunity, lipoproteins.

This PDF file includes:
Main Text
Figures 1 to 7

Abstract

During the cancerous transformation of normal hepatocytes into hepatocellular carcinoma (HCC),

the enzyme catalyzing the first rate-limiting step of glycolysis, namely the glucokinase (GCK), is

replaced by the higher affinity isoenzyme, hexokinase 2 (HK2). The transcriptomic analysis of HCC

tumors shows that highest expression level of *HK2* in tumor lesions is inversely correlated to *GCK*

expression, and is associated to poor prognosis for patient survival. To further explore functional

consequences of the GCK-to-HK2 isoenzyme switch occurring during carcinogenesis, *HK2* was

knocked-out in the HCC cell line Huh7 and replaced by *GCK*, to generate the Huh7-*GCK$^+$/HK2$^-$* cell

line. HK2 knockdown and GCK expression rewired central carbon metabolism, stimulated

mitochondrial respiration and restored essential metabolic functions of normal hepatocytes such

as lipogenesis, VLDL secretion, glycogen storage. It also reactivated innate immune responses and

sensitivity to natural killer cells, showing that consequences of the HK switch extend beyond

metabolic reprogramming.

Main Text

Introduction

Hepatocellular carcinoma (HCC) is the most common liver cancer and the fourth leading cause of

cancer-related death (1). HCC is closely linked to chronic liver inflammation, chronic viral hepatitis,

exposure to toxins, and metabolic dysfunction such as non-alcoholic steatohepatitis (NASH). HCC

is of poor prognosis, and treatments are essentially based on surgical resection, liver

transplantation or aggressive chemo and/or radiotherapy. In patients with advanced HCC, broad-

spectrum kinase inhibitors are approved (2) but with limited benefit (3). Effective personalized

therapies are needed but their development is impeded by our poor understanding of molecular

mechanisms underlying HCC onset and progression. Efforts to characterize the disease on the

basis of etiology and outcomes revealed metabolic deregulation as a hallmark of HCC progression

(4). Indeed, metabolic remodeling is critically required for tumor growth, since bioenergetic

requirements and anabolic demands drastically increase (5–7). For instance, HCC cells have lost

their ability to secrete very low-density lipoproteins (VLDL), a highly specialized function of

hepatocyte and can only secrete low-density lipoproteins (LDL)-like lipoproteins, indicating a

defective lipogenesis and/or lipoprotein assembly (8).

Metabolic reprogramming in cancer cells involves the modulation of several enzymes by

oncogenic drivers (6). Targeting these enzymes is now considered as a therapeutic strategy for

several types of cancers (6). Among these enzymes, hexokinase 2 (HK2) stands out because of its

elevated or induced expression in numerous cancers, including HCC (9). Hexokinases control the

first rate-limiting step of glucose catabolism by phosphorylating glucose to glucose-6-phosphate

(G6P), fueling glycolysis as well as glycogen, pentose phosphate and triglyceride synthesis. The

human genome contains four genes encoding distinct hexokinase isoenzymes, named HK1 to HK4

(HK4 is also known as glucokinase or GCK), with distinct enzymatic kinetics and tissue

distributions. A fifth putative hexokinase enzyme was recently discovered but has not been fully

characterized yet (10). A switch from GCK to HK2 isoenzymes is occurring during the transition

from primary to tumor hepatocytes so that HCC cell lines express HK2 but no longer GCK. HK2

expression level has been correlated with disease progression and dedifferentiation of HCC cells

(11). When HK2 is artificially knocked-down in HCC cell lines, glycolysis is repressed, and

tumorigenesis is inhibited while cell death increases (9). In addition, hexokinase function extends

beyond metabolism towards autophagy, cell migration, and immunity, suggesting that the GCK-

to-HK2 isoenzyme switch has broader consequences than initially suspected (12–15). Here, we

analyzed transcriptomic data of HCC biopsies and correlated hexokinase isoenzyme expression

level with patient survival. This led us to generate a new human HCC model expressing *GCK*

instead of *HK2*. A comparative analysis of *GCK$^+$ vs HK2$^+$* HCC cell lines provided a unique

opportunity to look into HK isoenzyme-dependent metabolic features, lipoprotein production and

resistance to immune signals of liver cancer cells.

Results

Relative expression level of GCK and HK2 in HCC patients

Although an isoenzyme switch from GCK to HK2 has been observed during the carcinogenesis

process (16), whether hexokinase isoenzymes expression is predictive of patient survival is

unclear. We first analyzed the transcriptomes (RNAseq data) of 365 HCC biopsies from The Cancer

Genome Atlas (TCGA) database (17, 18)(Supplementary Table 1). For each HK, the individual gene

expression level was used to stratify patients into two subgroups according to Uhlen et al. (18)

and overall survival in the two subgroups was determined using a Kaplan-Meier's estimator.

Although *HK1* or *HK3* expression level were not associated to patient survival rate (Fig. 1a), highest

expression levels of *HK2* as previously described (19) and lowest expression levels of *GCK* in the

tumors were associated with a lower survival rate. We thus stratified patients based on the

GCK/HK2 expression ratio to combine these two markers (Fig. 1b). When patients were stratified

on the basis of *HK2* or *GCK* expression levels, the median survival between the corresponding

subgroups differed by 33.8 and 36.5 months, respectively (Fig. 1a). This difference reached 42.8

months when the stratification of patients was based on the *GCK/HK2* ratio (Fig. 1b). This

demonstrated that the *GCK/HK2* ratio outperforms *HK2* or *GCK* expression alone as predictor of

patient survival. Finally, correlation coefficients between patient survival in months and *HK2* or

GCK expression level were determined. For this, we only considered the subset of 130 patients

for whom the period between diagnosis and death is precisely known (uncensored data), and

performed a Spearman's rank correlation test (Fig. 1c). Patient survival was positively correlated

to *GCK* expression but inversely correlated to *HK2* expression in line with the Kaplan-Meier

analysis. In addition, *GCK* and *HK2* expression tends to be inversely correlated in tumor samples

(Fig. 1c). Therefore, there is a trend for mutual exclusion of *GCK* and *HK2* expression in HCC

tumors, and this profile is associated to clinical outcome.

Engineering a cellular model of the hexokinase isoenzyme switch

To decipher functional consequences of GCK or HK2 expression in a HCC model, we restored GCK

expression by lentiviral transduction in the reference HCC cell line Huh7, and knocked-out the

endogenous *HK2* gene by CRISPR/Cas9. The exclusive expression of HK2 and *GCK* in Huh7 and

Huh7-*GCK⁺/HK2⁻* cell lines, respectively, was validated, while HK1 and HK3 were not expressed

(Fig. 2a). The hexokinase activity in the presence of increasing concentration of glucose was

determined in protein lysates from the two respective cell lines. Hexokinase activity in Huh7 lysate

reached its maximum at low glucose concentration, presenting a saturation curve according to

Michaelis–Menten kinetics (Fig. 2b). In contrast, the hexokinase activity in Huh7-*GCK⁺/HK2⁻*

lysates followed a pseudo-allosteric response to glucose (20, 21). Thus, the expected HK2 and GCK

activities were observed in the Huh7 and Huh7-*GCK⁺/HK2⁻* cells respectively. The cell proliferation

capacity remained identical between the two cell lines (Supplementary Fig. 1). We then compared

the genome edited Huh7-*GCK⁺/HK2⁻* and the parental Huh7 cell lines (*i.e.* Huh7-*GCK⁻/HK2⁺*) at a

transcriptomic, metabolic and immunological level.

Transcriptomic data revealed extended modifications of metabolic connections in Huh7-

GCK+/HK2-

Transcriptomic profiles of Huh7 and Huh7-*GCK⁺/HK2⁻* cells were determined by next generation

sequencing (Supplementary Table 2). Overall, 4.2% of the gene transcripts were reduced and 6.9%

were induced in Huh7-*GCK+/HK2-* compared to Huh7 (Fig. 2c; |fold-change (FC)|>2 and *p*

value<0.05). We first determined the metabolic consequences of the HK isoenzyme switch by

mapping the differentially expressed genes onto the a well-established bipartite metabolic

network Recon2, connecting gene products and metabolites (Supplementary Fig. 2-4) (22, 23).

After trimming highest-degree metabolites as currency metabolites, clusters of genes that are

both differentially expressed and connected by common metabolites emerged. Interestingly, we

found that across a wide range of analysis parameters, including varying rates of currency

metabolites and gene expression fold-change, the differentially expressed metabolic genes are

substantially better connected than expected by chance (Fig. 2d). This highlights the specificity of

the transcriptomic changes with respect to metabolic pathways. The spanned network presented

in Figure 2e corresponds to a stringent fold-change threshold for transcriptomic data

($\log_2(|FC|)>3$) while removing 2 percent of highest-degree currency metabolites. This network

shows connected components within glycolysis, but also across distant modules including the

gamma-aminobutyric acid (GABA) shunt (ALDH5A1), urea cycle (CPS1, OTC), glycogen metabolism

(GYS1, GYS2, AGL) and lipid synthesis (GPAM, AGPAT4, DGKG, CDS1, A4GALT) or degradation

(ACADL, HSD17B4, AMACR). This analysis highlights the global impact of the HK isoenzyme switch

that spreads beyond glycolysis across distant connected metabolic modules.

Enrichment of molecular and cellular functions in differentially expressed genes was also analyzed

using Ingenuity Pathway Analysis (IPA). This revealed that cellular movement and lipid

metabolism were the most affected functions (Fig. 3a and Supplementary Table 3). A closer look

at these annotations pointed to differences in the migratory capacities (Fig. 3b) as well as lipid

concentration and synthesis (Fig. 3c). The migratory capacities of Huh7 and Huh7-*GCK+/HK2-* were

compared using transwell-migration cell assays (Fig. 3d-e). Results showed a higher migratory

capacity of Huh7-*GCK+/HK2-* cells, in line with Kishore M. et al showing that GCK expression

induced by pro-migratory signals controls the trafficking of CD4+/CD25+/FOXP3+ regulatory T (T$_{reg}$)

cells (15). To validate the differences in lipid metabolism highlighted by the transcriptomic

analysis, intracellular content in neutral lipids was first determined with lipophilic dyes. As

assessed by Oil-Red-O or BODIPY staining, an accumulation of intracellular neutral lipids was

observed in Huh7-*GCK+/HK2-* in comparison to Huh7 (Fig. 3f and 3g). The accumulation of neutral

lipids in Huh7 expressing both *HK2* and *GCK* indicates that the phenotype is driven by *GCK*

expression and not by *HK2* knockdown (Supplementary Fig. 5). Lipid accumulation upon GCK

expression was also observed in Huh6 hepatoblastoma cells but not in epithelial kidney Vero cells,

indicating that this phenomenon occurs in metabolically relevant cells (Supplementary Fig. 5).

Differential lipid metabolism in Huh7 and Huh7-*GCK+/HK2-*

The intracellular lipid content of the two cell lines was further analyzed. In Huh7-*GCK+/HK2-*, an

enrichment in phosphatidylcholine, cholesterol, triglycerides (TG) and free fatty acids was

observed compared to Huh7 (Fig. 4a). One major function of hepatocytes is to secrete triglyceride-

rich VLDL and this function is altered in HCC cells that secrete smaller lipoproteins with the density

of LDL (24, 25). The secretion of lipids and lipoproteins by both cell lines was analyzed after a 24h-

culture in the absence of FCS to exclude any participation of exogenous lipids in the production

of lipoproteins. Huh7-*GCK+/HK2-* secreted more free fatty acids than Huh7 while secretion of

cholesterol and triglycerides (TG) remained unchanged (Fig. 4b). However, under the same

conditions, the secretion of apolipoprotein B (ApoB) by Huh7-*GCK+/HK2-* was reduced compared

to Huh7. Since ApoB is a non-exchangeable protein with only one copy in VLDL and LDL particles,

an elevated TG/ApoB ratio indicates that ApoB+-lipoproteins secreted by Huh7-*GCK+/HK2-* cells

are enriched in TG compared to those secreted by Huh7 (Fig. 4c). This was confirmed by the ApoB

distribution in density gradient fractions. As expected, lipoproteins secreted by Huh7 sediment at

the density of LDL, while those secreted by Huh7-*GCK+/HK2-* (Fig. 4d) match the density of VLDL

found in human plasma or secreted by primary human hepatocytes in culture (26, 27). This

indicates that GCK expression is essential for the VLDL assembly/secretion pathway and could

explain the loss of this crucial metabolic pathway in hepatoma cells expressing HK2 instead of GCK

(28).

Differential activity of the tricarboxylic acid cycle (TCA) in Huh7 and Huh7-*GCK+/HK2-*

We observed that GCK expression increased the intracellular content in lipids, resulting in

accumulation of lipid droplets and secretion of VLDL. A rewiring of cellular metabolism towards

energy storage in Huh7-*GCK+/HK2-* was thus suspected and confirmed by the accumulation of

glycogen, creatinine and creatinine-P (Fig. 5a and b), a feature of functional hepatocytes. To

further determine the consequences of replacing HK2 by GCK, we quantified prominent

intracellular metabolites via GC-MS. Figure 5c shows relative intracellular quantities of

metabolites that are significantly different between Huh7 and Huh7-*GCK+/HK2-*. Among

differentially represented metabolites, higher levels of glucose, glycerol-3-phosphate and lactic

acid were detected in Huh7-*GCK+/HK2- cells*. Several intermediates of the TCA cycle (succinic acid,

fumaric acid, alpha-ketoglutaric acid), and metabolites directly connected to it (GABA, glutamic

acid, glutamine, aspartic acid) were also differentially present between the two cell lines. This

supports a significant modulation of central carbon metabolism at both the level of glycolysis and

TCA cycle. This led to investigate glucose catabolism in further details. Glucose consumption and

stable isotope incorporation from [U-^{13}C]-glucose into pyruvate were both increased in Huh7-

*GCK+/HK2-*cells compared to Huh7 cells (Fig. 5d and f). This increased glycolytic flux together with

a reduced lactate secretion (Fig. 5e) is likely to account for the elevation of lactate levels and

suggest that the increased pyruvate production essentially fuels mitochondrial TCA cycle in Huh7-

$GCK^+/HK2^-$ cells.

Pyruvate entering the mitochondria downstream of glycolysis can be either oxidized by pyruvate

dehydrogenase (PDH), producing acetyl-CoA, or converted into oxaloacetate (OAA) by pyruvate

carboxylase (PC). Acetyl-CoA and OAA are then combined in the TCA cycle to form citrate. *De novo*

lipogenesis requires citrate egress from the TCA cycle to serve as a precursor of cytosolic acetyl-

CoA for further synthesis of fatty acids. In Huh7-$GCK^+/HK2^-$ cells, we observed both an increased

activity of PC (Fig. 5g) without changes in protein expression (Fig. 5h) and an increased

phosphorylation of pyruvate dehydrogenase (PDH), which is indicative of a reduced activity of this

enzyme (Fig. 5i). This is consistent with the increased expression of the PDH kinase PDK2 and the

decreased expression of the PDH phosphatase PDP2 in Huh7-$GCK^+/HK2^-$ cells that regulate the

PDH phosphorylation state (Fig. 5j). A rebalanced usage of pyruvate in Huh7-$GCK^+/HK2^-$ cells

maintains a functional TCA cycle and supports lipogenesis. In Huh7-$GCK^+/HK2^-$ cells, we also

observed an increased phosphorylation of ATP citrate lyase (ACLY), the first enzyme of the fatty

acid synthesis pathway, indicating an enhanced activity of this enzyme (Fig. 5k). This reaction also

regenerates OAA in the cytosolic compartment. Interestingly, transcriptomic data show that PCK1

which converts OAA to phosphoenolpyruvate (PEP), is overexpressed in Huh7-$GCK^+/HK2^-$ cells

compared to Huh7 (FC > 32).

A shift from pyruvate oxidation to carboxylation is observed in cancer cells where succinate

dehydrogenase (SDH) is inactivated by mutation and OAA can only be generated through PC

activity (29). SDH inhibition leads to succinate accumulation, especially in activated immune cells

(30). Interestingly, higher levels of succinate and a reduced activity of SDH were measured in

Huh7-$GCK^+/HK2^-$ cells compared to Huh7 (Fig. 5l and m). Even though SDH is also part of the

complex II of the mitochondrial respiratory chain, we observed that the overall oxygen

consumption was increased in Huh7-$GCK^+/HK2^-$ (Fig. 5n) with increased basal and maximal

respiration, ATP production and spare respiration capacity (Supplementary Fig. 6). Functional

analysis of the respiratory chain showed that oxygen consumption in Huh7 and Huh7-$GCK^+/HK2^-$

cells was mainly dependent on complex I which is fueling complex III (Fig. 5n and o). Thereby, the

HK isoenzyme switch rewired the TCA cycle promoting carboxylation of pyruvate into OAA in the

presence of a reduced SDH activity and increased respiration through complex I.

Restored innate immune sensitivity in Huh7-$GCK^+/HK2^-$

Lipid accumulation in hepatocytes is incriminated in hepatic inflammation (31) and TCA cycle

rewiring is associated with innate immunity activation (32). These two events were observed

when replacing HK2 by GCK in Huh7 cells, questioning the immune status of these cells and their

sensitivity to antitumor immunity. The functional analysis of gene ontology (GO) terms associated

to differentially expressed transcripts revealed an enrichment in terms related to the regulation

of innate immunity. The gene signature associated with type-I interferon (IFN) signaling pathway

scored among the top enriched terms of upregulated genes in Huh7-$GCK^+/HK2^-$ cells

(Supplementary Fig. 7). Within the 91 gene members of this GO term, 20 transcripts of Type I-IFN

signaling were significantly up-regulated in Huh7-$GCK^+/HK2^-$ compared to Huh7 (Fig. 6a). This

includes interferon regulatory factors (IRF1, IRF7 and IRF9), IFN-stimulated genes (ISGs) such as

ISG15, MX1, OAS1, OAS3, RNaseL and signaling intermediates such as IKBKE coding for IKKε (Fig.

6b). The chaperon HSP90AB1, which is involved in the phosphorylation and activation of STAT1,

was also induced (33). In contrast, two genes were significantly down-regulated in Huh7-

$GCK^+/HK2^-$ compared to Huh7, the RNaseL inhibitor ABCE1, and TRIM6, an E3 ubiquitin-protein

ligase regulating IKKε. Among the members of the RIG-I-like receptor (RLR) family involved in

immune sensing and antitumor defense (34), the expression of IFIH1 (also known as MDA5) was

slightly increased in Huh7-$GCK^+/HK2^-$, while RIG-I itself remained unchanged. Such identified gene

sets suggest that Huh7-$GCK^+/HK2^-$ cells may be better equipped than Huh7 cells to respond to

innate immune signals. Thus, we compared the innate immune response of the two cell lines

when stimulated by RIG-I or MDA5 ligands, known to sense different double-stranded RNA

(dsRNA) in viral immune recognition. In order to quantify cellular activation by dsRNA, immuno-

stimulatory challenge assays were conducted with triphosphate-hairpin RNA (3p-hpRNA) or

polyinosinic-polycytidylic acid (poly(I:C)). Interferon-sensitive response element (ISRE)-

dependent transcription was efficiently induced by these RLR ligands in Huh7-$GCK^+/HK2^-$ cells

whereas Huh7 response was very limited even at high doses of immuno-stimulatory ligands (Fig.

6c).

To investigate whether this differential sensitivity to RLR ligands is linked to GCK expression or

HK2 knockout, we used Huh7 cells transduced for GCK expression so that they express both HK2

and GCK (Huh7-$GCK^+/HK2^+$). In these cells, the response to RIG-I ligation did not differ from that

of Huh7 cells suggesting that GCK expression alone is not sufficient to restore immune sensitivity

(Fig. 6d). When HK2 expression was repressed in these cells with a shRNA (Huh7-$GCK^+/HK2^-$Sh)

(see Supplementary Fig. 8 showing 95% extinction of HK2 protein), ISRE response to RIG-I signaling

was restored to a level similar to that observed in Huh7-$GCK^+/HK2^-$ cells (Fig. 6d). This is pointing

towards HK2 as a negative regulator of RLR signaling in HCC cells and suggests that the *GCK*-to-

HK2 isoenzyme switch during malignant transformation of hepatocytes is accompanied by a

reduced sensitivity to innate immune signals. The higher sensitivity to RLR ligands of Huh7-

$GCK^+/HK2^-$ cells also resulted in increased secretion of inflammatory interleukins (IL-6 and IL-8),

antiviral cytokines (IFN-λ_1, IFN-$\lambda_{2/3}$, IFN-β), and IP-10 (Fig. 6e), indicating that both NF-κB- and

IRF3-dependent signaling pathways were induced. IL-1β, TNFα, IL-12p70, GM-CSF, IL-10 and IFNγ

were not detected in the supernatants of none of the cell lines, whether they were stimulated or

not.

As natural killer (NK) cell-mediated lysis of tumor cells is crucial for the anti-cancer immune

defense, we compared the susceptibility of the two cell lines to NK cells cytotoxicity. Figure 6f

shows that Huh7 cells are resistant to NK cell-mediated lysis in contrast to Huh7-$GCK^+/HK2^-$. Thus,

replacing HK2 by GCK restored Huh7 sensitivity to NK cell-mediated lysis. Similar results were

obtained when NK cells were pre-activated with IL-2 (Supplementary Fig. 9). Altogether, these

results demonstrate that HCC cells expressing HK2 instead of GCK exhibit an impaired response

to immune signals and also a strong resistance to NK cells. Significantly, these two observations

are in line with clinical data showing that elevated GCK expression is associated with prolonged

survival, while elevated HK2 expression coinciding with GCK reduction correlates with shorter

overall survival (Fig. 1).

Discussion

Metabolic network rewiring is a hallmark of cancer although for most tumors, mechanisms at the

origin of this metabolic reprogramming have not been elucidated. While GCK, but not HK2, is

expressed in normal hepatocytes, the expression of HK2 occurs during cirrhosis and increases as

the disease progresses to carcinoma. Several signaling pathways such as hypoxia inducible factors

(HIF), peroxisome proliferator-activated receptors (PPAR) and phosphatidylinositol-4,5-

bisphosphate 3-kinase (PI3K) might contribute to HK2 induction in fatty liver disease and its

evolution towards cirrhosis and carcinogenesis (35–37). Consequently HK2 induction has been

proposed as a risk marker of HCC development (16). Analyzing TCGA data from human HCC

tumors, we observed that not only high levels of *HK2* but also low levels of *GCK* are of poor

prognosis. In contrast, neither *HK1* nor *HK3* expression levels were correlated with survival of HCC

patients. *GCK* expression is very low or not detected in biopsies from a majority of patients (65.8%

of patients show RSEM values <10), whereas *HK2* is widely expressed (16) (only 5.8% of patients

show RSEM values <10). This probably explains that HK2 expression is a better prognostic marker

than GCK for HCC. However, when GCK and HK2 expression were combined into a single ratio,

this prognostic marker outperformed HK2 or GCK expression alone. This suggests that both HK2

induction and GCK loss play a role in HCC progression. As HK2 and GCK expression tend to be

mutually exclusive, both HK2 induction and GCK downregulation might have consequences on the

metabolic reprogramming during malignant transformation of hepatocytes. To compare the

functional consequences of the HK isoenzyme switch in HCC, we therefore expressed GCK in the

reference HCC cell line Huh7 and knocked-down HK2 expression. Our comparative transcriptomic,

metabolic and functional studies demonstrate that the replacement of HK2 by GCK not only

restored some essential metabolic functions of normal hepatocytes such as lipogenesis, VLDL

secretion and glycogen storage but also reactivated innate immune responses and sensitivity to

NK-mediated cell lysis.

HCC cell lines predominantly secrete LDL-like particles, unlike normal hepatocytes, which secrete

VLDL. Lipid loading of Huh7 cells with oleic acid can boost the secretion of ApoB$^+$ particles but

does not induce a shift from LDL to VLDL density, indicating that intracellular fatty acid

accumulation of exogenous origin cannot rescue VLDL production (28). Here we show that

replacing HK2 by GCK in Huh7 cells restored *de novo* fatty acid synthesis, allowing VLDL

assembly/secretion in the absence of exogenous lipids. To our knowledge Huh7-*GCK$^+$/HK2$^-$* is the

first human cell model with a functional VLDL secretion pathway. Such a tool will strongly benefit

the field of cardiovascular diseases and hepatic steatosis.

De novo fatty acid synthesis from carbohydrates requires an adequate supply in metabolic

substrates, especially citrate that is produced by the TCA cycle from incoming pyruvate. The

glycolytic entry point into the TCA cycle is controlled by PDH and PC that convert pyruvate into

acetyl-CoA or OAA, respectively. Our data revealed that in addition to the increased production

of pyruvate from glucose, PC activity is increased whereas PDH is inhibited. This suggests that

pyruvate metabolism is rebalanced in favor of OAA in Huh7-*GCK$^+$/HK2$^-$* cells, as described in

healthy liver. Such a mechanism of anaplerosis is known to replenish TCA cycle intermediates and

compensate citrate export out of the mitochondria for lipogenesis fueling. Increased PC activity

is observed in both normal and pathological situations, mainly as a result of an increased

transcription of the PC gene. In our model, mRNA and protein levels were not affected, indicating

that PC activity can be regulated by alternative mechanisms depending on HK isoenzyme

expression. This may relate to lower levels of oxalate, a known inhibitor of PC activity, in Huh7-

GCK$^+$/HK2$^-$ cells (Fig. 5c and Fig. 7 discussed below).

A rebalanced pyruvate usage in favor of OAA is also described for instance in SDH-deficient

neuroendocrine tumor cells, where succinate accumulates and PC activity is increased to maintain

OAA production, replenish the oxidative TCA cycle and support aspartate synthesis (29).

Interestingly, in comparison to Huh7 cells, succinate and aspartate levels are elevated in Huh7-

GCK$^+$/HK2$^-$ where SDH activity is reduced, suggesting a direct link between PC and SDH activity in

hepatocytes. Several mechanisms inhibiting SDH have been described (38). Modification of the

expression of SDH subunits is unlikely as no variation was observed at the transcriptomic level*.

Itaconate is a weak inhibitor of SDH produced by IRG1 from aconitate but this metabolite was not

detected and IRG1 mRNA was absent from the transcriptome of both cell lines. Whether fumarate

or other metabolites are responsible for the reduced SDH activity in GCK-expressing cells remains

to be investigated. Finally, SDH-deficient cells and LPS-stimulated macrophages have been shown

to elicit a hypoxic-like phenotype through accumulation of large amounts of succinate and

stabilization of HIF-1α (39, 40). Despite an elevated succinate steady-state level in Huh7-

GCK[+]/*HK2*[-] compared to Huh7 cells, we observed no difference in HIF-1α stabilization neither at

basal level nor upon induction (Supplementary Fig. 10). This suggested that the reduction of SDH

activity in Huh7-*GCK*[+]/*HK2*[-] cells was not strong enough to induce such a pseudo-hypoxic

phenotype.

Our gene-centric metabolic analysis of transcriptomic data revealed a wide spreading of

metabolic modifications resulting from HK isoenzyme switch. Illustrating these modifications,

Figure 7 is an attempt to integrate the observed changes in central carbon metabolism and closely

connected metabolic pathways. In particular, decreased level of alanine and increased aspartate

concentration in Huh7-*GCK*[+]/*HK2*[-] cells could be an indirect effect of PC activation that uses

pyruvate for the synthesis of OAA. As a consequence, hepatic transaminases may balance

intracellular pools of OAA, aspartate, alanine and pyruvate. Glutamate and GABA levels were also

modified, thus supporting anaplerosis of the TCA cycle through glutamine consumption and the

GABA shunt pathway, respectively. We also observed lower levels of oxalate, an end-product of

glyoxylate degradation. In Huh7-*GCK*[+]/*HK2*[-] *cells,* increased levels of the enzyme alanine-

glyoxylate and serine-pyruvate aminotransferase (AGXT) could account for this phenotype as it

converts alanine and glyoxylate into pyruvate and glycine, which is also increased. Interestingly,

high level of AGXT is a good prognostic marker for HCC (41). Consistently, it was found that oxalate

inhibits liver PC, resulting in reduced gluconeogenesis and lipogenesis (42, 43). Thus, a higher PC

activity could be explained by lower levels of oxalate in Huh7-*GCK*[+]/*HK2*[-] cells. We also observed

that isoleucine and valine levels increased while BCAT1 (branched chain amino acid transaminase

1) predominant transcripts decreased. This suggests a reduced catabolism of branched chain

amino acids in Huh7-*GCK⁺/HK2⁻* cells. Again, low levels of BCAT1 is a good prognostic marker for

HCC and oral supplementation with branched chain amino acids has been shown to reduce the

risk of liver cancer in cirrhotic patients (44, 45). If some metabolic modifications seem to advocate

for the restoration of a normal hepatocyte phenotype following the replacement of HK2 by GCK,

it cannot be a general statement. Indeed, the urea cycle was also impacted in Huh7-*GCK⁺/HK2⁻*

cells with lower levels of CPS1 and OTC, which are also observed in aggressive HCC tumors (46).

Altogether, our results demonstrate the broad impact of replacing HK2 by GCK in HCC cells, and

the key role played by the HK isoenzyme switch in HCC tumor metabolism.

We discovered that HK isoenzyme expression not only controls hepatic metabolic functions but

also interferes with intrinsic innate immunity of hepatocytes and antitumor immune surveillance.

Several reports have recently established functional links between glucose metabolism and

signaling pathways downstream of innate immunity sensors of the RLR family, RIG-I and MDA5

(47–49), which are usually associated with antiviral responses. However, RIG-I expression is

downregulated in hepatic cancer tissues and low RIG-I expression is correlated with poor survival

of patients, whereas RIG-I expression in HCC cell lines enhanced IFN response and cancer cell

apoptosis (34). This suggests an unexpected role of this receptor family in the antitumor response.

Here we show that Huh7 cells expressing GCK instead of HK2 exhibit a higher sensitivity to RIG-I

and MDA5 ligands, and produce higher levels of type I/III IFNs and inflammatory cytokines. This

immune phenotype occurs in a context of reduced SDH activity and increased intracellular content

in succinate (Fig. 5l-m). A pro-inflammatory function of immune cells such as macrophages was

previously linked to TCA rewiring, with reduced SDH activity resulting in succinate accumulation

(39, 50, 51). Succinate can also be secreted from LPS-activated macrophages and activate its

cognate receptor, succinate receptor 1 (SUCNR1, previously known as GPR91) in an autocrine and

paracrine manner to further enhance production of IL-1β (52). Interestingly, glucose metabolism

promotes RIG-I and MDA5 signaling through the O-GlcNAcylation of the mitochondrial adaptor

MAVS (47). Thus, an intriguing hypothesis is that GCK expression could facilitate MAVS signaling

by increasing UDP-GlcNAc through upregulation of the hexosamine biosynthetic pathway. HK2

binding at the surface of mitochondria may also compete with pyruvate carboxylase, metabolites

or mitochondria factors known to control MAVS signaling (47–49). Here we show that HK2

knockdown promotes RIG-I-induced ISRE-dependent transcription (Fig. 6d). This is consistent with

the results obtained by Zhang W. et al. (48), indicating that HK2 interaction with MAVS restrains

RIG-I-induced IFN-β secretion. Further investigations are now required to decipher the molecular

links between metabolism and immune responses.

Beyond the inhibition of RLR signaling, other mechanisms might contribute to tumor escape from

immune surveillance in HCC patients. In advanced-stage HCC patients, NK cells often exhibit

reduced infiltration and impaired functional activities (53). We thus compared the sensitivity of

Huh7-$GCK^+/HK2^-$ cells to Huh7, and found that sensitivity to NK cell lysis is restored when HK2 is

replaced by GCK. When analyzing cell surface expression of the NK cells inhibitors, HLA class I and

MICA/B, no significant changes were observed between cell lines (Supplementary Fig. 9). In

contrast, an increased transcription and surface expression of ICAM1 (FC=2.6; Supplementary Fig.

9) was observed in Huh7-$GCK^+/HK2^-$. Since ICAM-1 binding to active LFA-1 at the surface of NK

cells is essential for granule polarization and efficient killing of the target cells (54), its enhanced

exposition at the surface of Huh7-$GCK^+/HK2^-$ may contribute to their higher sensitivity to NK cell-

mediated killing. These results suggest that HK2 expression at the expense of GCK in HCC tumors

decreases immune responsiveness and sensitivity to NK cytotoxicity, thus favoring immune

escape.

Taken together, our data demonstrate that beyond glycolysis, the hexokinase isoenzyme switch

in an HCC model rewires central carbon metabolism, promotes lipogenesis, enhances innate

immune functions, and restores sensitivity to natural killer cells.

Materials and Methods
422
Materials. Unless otherwise specified, chemicals were from Merck Sigma-Aldrich. The RIG-I

specific ligand 3p-hpRNA and the MDA5/TLR3 ligand poly(I:C) HMW (High Molecular Weight) were

from Invivogen.

Cell cultures. Huh7 cells and derivatives were grown as previously described (55) in DMEM, 10%

fetal calf serum (FCS), penicillin/streptomycin, 1 mM pyruvate, 2 mM L-glutamine. Culture

medium and additives were from Gibco except FCS (Dominique Dutcher).

Cell lines. 15×10^4 Huh7 cells were transduced for GCK expression at different multiplicities of

infection (lentiviral transduction using the pLEX-GCK construct). The Huh7-$GCK^+/HK2^+$ cells were

then cultured for 7 days with puromycin (1 µg/mL) before amplification. HK2 knock-out was

achieved using the CRISPR/Cas9 system as previously described (56) to obtain Huh7-$GCK^+/HK2^-$

cells. A single guide RNA (sgRNA) pair was designed for double nicking using the CRISPR Design

Tool (http://tools.genome-engineering.org). The guide sequence oligos (sgRNA$_1$(*HK2*): 5'-

CACCGTGACCACATTGCCGAATGCC-3' and sgRNA$_2$(*HK2*): 5'-CACCGTTACCTCGTCTAGTTTAGTC-3')

were cloned into a plasmid containing sequences for Cas9 expression and the sgRNA scaffold

(pSpCas9(BB)-2A-GFP, Addgene plasmid #48138). 48 h post-transfection, cells were sorted by

FACS based on the transient expression of GFP and cloned by limiting dilution. Effective deletion

of HK2 was assessed by qPCR.

For *HK2* knock-down, Huh7-*GCK+/HK2+* cells were transduced with lentiviral vectors expressing

HK2-targeting shRNAs, and antibiotic selection was applied (hygromycin; 100 μg/ml). The *HK2*-

targeting sequence 5'-

CCGGCCAGAAGACATTAGAGCATCTCTCGAGAGATGCTCTAATGTCTTCTGGTTTTTT-3' was cloned in

the pLKO.1 hygro vector (a gift from Bob Weinberg; Addgene plasmid #24150). HK2 expression in

Huh7-*GCK+/HK2+* and Huh7-*GCK+/HK2-Sh* was analyzed on cell lysates by western blotting

(Supplementary Fig. 8).

Enzymatic activity assays. Cells were trypsinized, washed twice, and cell pellets were stored at -

80°C. Protein extractions and assays were performed as previously described for hexokinase (57–

59) and pyruvate carboxylase (60). Specific enzymatic assays are described in Supplementary

Methods.

Metabolomics profiling. Cells were seeded at $13x10^5$ cells in $75cm^2$ dishes. After 24 h, culture

medium was replaced and cells were further incubated during 24 h. Culture supernatants were

collected and stored at -80°C. Cells were harvested, washed twice with ice-cold PBS, and cell

pellets were stored at -80°C. To analyze catabolic glucose flux, cells were incubated with [U-^{13}C]-

glucose (Sigma-Aldrich; 389374-2G) at 12.5 mM and unlabeled glucose at 12.5 mM for 24h.

Extraction, separation and analysis of the metabolites by GC-MS are described in Supplementary

Methods.

Transcriptome profiling of Huh7 and Huh7-*GCK+/HK2-* cell lines. Transcriptome profiling was

performed by next generation sequencing (Profilexpert, Lyon). To identify differentially expressed

genes, raw data were processed using the DESeq2 analysis pipeline (61). See Supporting

Information and Gene Expression Omnibus database with the accession number GSE144214 for

entire raw data.

Pathway analysis. The list of transcripts differentially expressed in Huh7 and Huh7-*GCK+/HK2-* cell

lines was analyzed by gene set enrichment analysis (IPA, Build version: 486617M, Qiagen)

weighted by their corresponding fold-change and *p* value. The fold-change cut-off of mean

expression for each transcript was set at 2 with an adjusted *p* value<0.05. The list of genes

associated with "Type I-IFN signaling pathway" was defined in the AmiGO 2 database. Expression

data of these genes were retrieved from the transcriptomes of Huh7-*GCK+/HK2-* and Huh7, and

correspond for each gene to the most differentially expressed transcript.

Cell migration assay. $2x10^4$ cells were plated in the upper chamber of transwells (Sarstedt, PET

8.0-μm, TL - 833932800) with DMEM without FCS to allow migration for 24 h at 37°C. DMEM with

10% FCS was distributed in each well, below the chamber. Chambers were gently picked up before

a brief PBS rinse and 0.05% crystal violet coloration. The migrated cells were analyzed a Leica M50

microscope using a magnification factor of 20x. The number of cells that have migrated through

the membrane and attached on the underside of the membrane were counted using the software

Image J.

Respiration Assay. Respiration was measured using an Extracellular Flux Analyzer (Seahorse

Bioscience). The assay was performed according to the Agilent Seahorse XF Cell Mito Stress Test

as detailed in Supplementary Methods. The number of cells was determined at the end of the run

using the SRB assay as previously described (62) or in plate cell counting by Cytation 1 imaging

reader (Biotek).

Intracellular lipid staining. For microscopic observations, cells were fixed 48 h post-seeding, and

intracellular lipids stained with Oil-Red-O as described in Supplementary Methods. For the

quantification of intracellular lipid droplets by FACS, cells were stained with the 4,4-difluoro-

1,3,5,7,8-Pentamethy-4-Bora-3a,4a-Diaza-s-indacene (BODIPY) 493/503 dye (Tocris Bio-Techne).

Protein, ApoB, and Lipid Quantification. Protein concentration was determined using the DC

Protein Assay (Bio-Rad). ApoB concentration in medium and gradients fractions was determined

by ELISA as previously described (63). Total concentrations of cholesterol, phospholipids, and

triglycerides (TG) were determined using specific assays from Millipore Sigma-Aldrich (ref.

MAK043, MAK122 and MAK266 respectively). Free Fatty Acids were quantified using a specific

assay kit from Abcam (ref. ab65341).

Iodixanol density gradients. Iodixanol gradients were prepared as previously described (64). One

ml of culture supernatant was applied to the top of 6 to 56% iodixanol gradients and centrifuged

for 10h at 41,000rpm and 4°C in a SW41 rotor. The gradient was harvested by tube puncture from

the bottom and collected into 22 fractions (0.5 ml each). The density of each fraction was

determined by weighing.

Metabolic network coherence computational analysis. In order to measure the consistency of

differentially expressed genes with a metabolic network, we employed the metabolic network

coherence measure introduced by Sonnenschein *et al.* (65) (see Supplementary Methods for

details). This approach was previously applied to various disease-related transcriptome profiles

(66, 67) and for extracting information on the genetic control of metabolic organization (68).

Recently, detailed theoretical analysis of the extended version of the method has been performed

by Nyczka and Hütt (69)

Western-Blot Analysis. Protein expression was determined by standard western-blot analysis

(See Supplementary Methods for details).

RLR stimulation

Cells were seeded in 96-well or 24-well plates. After 24 h, cells were co-transfected with indicated

doses of the RIG-I ligand 3p-hpRNA or the MDA5/TLR3 ligand poly(I:C) HMW together with the

pISRE-luc (1.25 µg/ml) and pRL-SV40 (0.125 µg/ml) reporter plasmids using the JetPEI-Hepatocyte

reagent (Polyplus Transfection). Manufacturer's instructions were followed. After 48 h,

supernatants were collected for cytokine quantification. Firefly and Renilla luciferase expressions

within cells were determined using the Dual-Glo luciferase Assay system (Promega) and an Infinite

M200 microplate reader (TECAN).

Cytokine assays

Clarified culture supernatants were collected and stored at -20°C. IL-8 was quantified using the

Cytometric Bead Array for human IL-8 (BD Biosciences). Other cytokines were assayed using the

LEGENDPlex multiplex assay (Human Anti-Virus Response Panel, BioLegend). Fluorescence was

analyzed using a FACS Canto II (BD Biosciences).

NK cell cytotoxicity test

NK cells were isolated as described in Supplementary Methods from human buffy coats of healthy

donors obtained from the Etablissement Français du Sang. Informed consent was obtained from

donors and experimental procedures were approved by the local institutional review committee.

Huh7 or Huh7-$GCK^+/HK2^-$ were seeded at $1x10^5$ cells per well in a 24-well plate in RPMI-1640

(Gibco) with 10% FCS and 40 µg/ml gentamycin. After 24 h, $3x10^5$ or $3x10^6$ NK cells were added

to the culture wells. The cytotoxicity assay was performed for 4 h at 37°C, under 5% CO_2. Target

hepatoma cells were harvested after trypsination, labelled with propidium iodide (PI) and

analyzed by FACS. Cell death was monitored after morphological gating on hepatocytes.

Statistics and reproducibility

All the statistical analyses were performed with GraphPad Prism or Analyse-it softwares. Details

of statistical analyses can be found in figure legends.

Acknowledgments

We acknowledge the contribution of the Genomics and Microgenomics platform ProfileXpert

(University Lyon 1, SFR santé LYON-EST, UCBL-Inserm US 7-CNRS UMS3453) and SFR Biosciences

(UMS3444/CNRS, US8/Inserm, ENS de Lyon, UCBL) facilities: AniRA-Cytometry, AniRA-ImmOs

metabolic phenotyping and LYMIC-PLATIM microscopy. We gratefully thank Laurence Canaple for

technical assistance. This work was supported by the Fondation pour la Recherche Médicale

(FRM), grant number DEQ20160334893 to VL. F.V.F. is grateful for the support by grants

CA154887, GM115293, CRN-17-427258, NSF GRFP, and the Science Alliance on Precision

Medicine and Cancer Prevention by the German Federal Foreign Office, implemented by the

Goethe-Institute, Washington, DC, USA, and supported by the Federation of German Industries

(BDI), Berlin, Germany.

Author Contributions

L.P-C., P-O.V., O.D., V.L. designed the experiments with critical advices from G.J.P.R., P.A., R.R. and

F.V.F.; C.J., A.A-G., K.O., B.P., G.J.P.R., N.A, R.R. and F.V.F. performed experiments and analyzed

the data; P.N. and M-T.H. performed metabolic network computational analysis; L.P-C., P-O.V.,

P.A., F.V.F, V.L. and O.D. analyzed the data, prepared figures and wrote the manuscript.

References

1.   A. Villanueva, Hepatocellular Carcinoma. *N. Engl. J. Med.* **380**, 1450–1462 (2019).

2.   M. Kudo, *et al.*, Lenvatinib versus sorafenib in first-line treatment of patients with
unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *The*
Lancet **391**, 1163–1173 (2018).

3.   J. M. Llovet, R. Montal, D. Sia, R. S. Finn, Molecular therapies and precision medicine for
hepatocellular carcinoma. *Nat Rev Clin Oncol* **15**, 599–616 (2018).

4. S. De Matteis, *et al.*, Aberrant Metabolism in Hepatocellular Carcinoma Provides
Diagnostic and Therapeutic Opportunities. *Oxidative Medicine and Cellular Longevity* **2018**,
1–13 (2018).

5. F. V. Filipp, *et al.*, Glutamine-fueled mitochondrial metabolism is decoupled from glycolysis
in melanoma. *Pigment Cell & Melanoma Research* **25**, 732–739 (2012).

6. N. Hay, Reprogramming glucose metabolism in cancer: can it be exploited for cancer
therapy? *Nat Rev Cancer* **16**, 635–649 (2016).

7. M. G. Vander Heiden, L. C. Cantley, C. B. Thompson, Understanding the Warburg effect:
the metabolic requirements of cell proliferation. *Science* **324**, 1029–1033 (2009).

8. J. Jiang, P. Nilsson-Ehle, N. Xu, Influence of liver cancer on lipid and lipoprotein
metabolism. *Lipids Health Dis* **5**, 4 (2006).

9. D. DeWaal, *et al.*, Hexokinase-2 depletion inhibits glycolysis and induces oxidative
phosphorylation in hepatocellular carcinoma and sensitizes to metformin. *Nature
Communications* **9**, 446 (2018).

10. C. Guo, *et al.*, Coordinated regulatory variation associated with gestational hyperglycaemia
regulates expression of the novel hexokinase HKDC1. *Nat Commun* **6**, 6069 (2015).

11. G. Guzman, *et al.*, Evidence for Heightened Hexokinase II Immunoexpression in
Hepatocyte Dysplasia and Hepatocellular Carcinoma. *Dig Dis Sci* **60**, 420–426 (2015).

12. A. J. Wolf, *et al.*, Hexokinase Is an Innate Immune Receptor for the Detection of Bacterial
Peptidoglycan. *Cell* **166**, 624–636 (2016).

13. V. P. Tan, S. Miyamoto, HK2/hexokinase-II integrates glycolysis and autophagy to confer
cellular protection. *Autophagy* **11**, 963–964 (2015).

14. D. J. Roberts, V. P. Tan-Sah, E. Y. Ding, J. M. Smith, S. Miyamoto, Hexokinase-II Positively
Regulates Glucose Starvation-Induced Autophagy through TORC1 Inhibition. *Mol. Cell* **53**,
521–533 (2014).

15. M. Kishore, *et al.*, Regulatory T Cell Migration Is Dependent on Glucokinase-Mediated
Glycolysis. *Immunity* **47**, 875-889.e10 (2017).

16. N. C. W. Lee, M. A. Carella, S. Papa, C. Bubici, High Expression of Glycolytic Genes in
Cirrhosis Correlates With the Risk of Developing Liver Cancer. *Front Cell Dev Biol* **6**, 138
(2018).

17. M. Uhlén, *et al.*, Tissue-based map of the human proteome. *Science* **347** (2015).

18. M. Uhlen, *et al.*, A pathology atlas of the human cancer transcriptome. *Science* **357** (2017).

19. J.-J. Yoo, *et al.*, Hexokinase-II Inhibition Synergistically Augments the Anti-tumor Efficacy of
Sorafenib in Hepatocellular Carcinoma. *Int J Mol Sci* **20** (2019).

20. C. M. Porter, B. G. Miller, Cooperativity in monomeric enzymes with single ligand-binding
sites. *Bioorganic Chemistry* **43**, 44–50 (2012).

21. M. Larion, R. K. Salinas, L. Bruschweiler-Li, B. G. Miller, R. Brüschweiler, Order–Disorder
Transitions Govern Kinetic Cooperativity and Allostery of Monomeric Human Glucokinase.
PLoS Biology **10**, e1001452 (2012).

22. I. Thiele, *et al.*, A community-driven global reconstruction of human metabolism. *Nat.*
Biotechnol. **31**, 419–425 (2013).

23. D. C. Zielinski, *et al.*, Pharmacogenomic and clinical data link non-pharmacokinetic
metabolic dysregulation to drug side effect pathogenesis. *Nature Communications* **6**, 7101
(2015).

24. G. van den Berghe, The role of the liver in metabolic homeostasis: Implications for inborn
errors of metabolism. *Journal of Inherited Metabolic Disease* **14**, 407–420 (1991).

25. J. M. Berg, J. L. Tymoczko, L. Stryer, Each Organ Has a Unique Metabolic Profile.
Biochemistry. 5th edition (2002) (January 13, 2020).

26. F. Beilstein, *et al.*, Lysophosphatidylcholine acyltransferase 1 is downregulated by hepatitis
C virus: impact on production of lipo-viro-particles. *Gut* **66**, 2160–2169 (2017).

27. H. N. Ginsberg, LIPOPROTEIN PHYSIOLOGY. *Endocrinology and Metabolism Clinics of North*
America **27**, 503–519 (1998).

28. S. J. R. Meex, U. Andreo, J. D. Sparks, E. A. Fisher, Huh-7 or HepG2 cells: which is the better
model for studying human apolipoprotein-B100 assembly and secretion? *J. Lipid Res.* **52**,
152–158 (2011).

29. C. Lussey-Lepoutre, *et al.*, Loss of succinate dehydrogenase activity results in dependency
on pyruvate carboxylation for cellular anabolism. *Nat Commun* **6**, 8784 (2015).

30. V. Lampropoulou, *et al.*, Itaconate Links Inhibition of Succinate Dehydrogenase with
Macrophage Metabolic Remodeling and Regulation of Inflammation. *Cell Metab.* **24**, 158–
166 (2016).

31. A. Mansouri, C.-H. Gattolliat, T. Asselah, Mitochondrial Dysfunction and Signaling in
Chronic Liver Diseases. *Gastroenterology* **155**, 629–647 (2018).

32. E. L. Mills, B. Kelly, L. A. J. O'Neill, Mitochondria are the powerhouses of immunity. *Nature*
Immunology **18**, 488 (2017).

33. M. Cheng, *et al.*, Stat1 mediates an auto-regulation of hsp90β gene in heat shock
response. *Cellular Signalling* **22**, 1206–1213 (2010).

34. J. Hou, *et al.*, Hepatic RIG-I predicts survival and interferon-α therapeutic response in
hepatocellular carcinoma. *Cancer Cell* **25**, 49–63 (2014).

35. G. Panasyuk, *et al.*, PPARγ contributes to PKM2 and HK2 expression in fatty liver. *Nat*
Commun **3**, 672 (2012).

36. O. A. Mesarwi, *et al.*, Hepatocyte Hypoxia Inducible Factor-1 Mediates the Development of
Liver Fibrosis in a Mouse Model of Nonalcoholic Fatty Liver Disease. *PLoS One* **11** (2016).

37. D. J. Roberts, S. Miyamoto, Hexokinase II integrates energy metabolism and cellular
protection: Akting on mitochondria and TORCing to autophagy. *Cell Death and*
Differentiation **22**, 248–257 (2015).

38. J. Van den Bossche, L. A. O'Neill, D. Menon, Macrophage Immunometabolism: Where Are
We (Going)? *Trends in Immunology* **38**, 395–406 (2017).

39. G. M. Tannahill, *et al.*, Succinate is an inflammatory signal that induces IL-1β through HIF-
1α. *Nature* **496**, 238–242 (2013).

40. K. Kluckova, D. A. Tennant, Metabolic implications of hypoxia and pseudohypoxia in
pheochromocytoma and paraganglioma. *Cell Tissue Res.* **372**, 367–378 (2018).

41. Y. Sun, *et al.*, Loss of alanine-glyoxylate and serine-pyruvate aminotransferase expression
accelerated the progression of hepatocellular carcinoma and predicted poor prognosis. *J*
Transl Med **17**, 390 (2019).

42. E. A. Yount, R. A. Harris, Studies on the inhibition of gluconeogenesis by oxalate. *Biochim.*
Biophys. Acta **633**, 122–133 (1980).

43. I. E. O'Neill, D. W. Bannister, The effects of oxalate and glucose on lipogenesis by isolated
hepatocytes from normal and biotin-deficient chicks (Gallus domesticus). *Int. J. Biochem.*
16, 517–521 (1984).

44. Y. Muto, *et al.*, Overweight and obesity increase the risk for liver cancer in patients with
liver cirrhosis and long-term oral supplementation with branched-chain amino acid
granules inhibits liver carcinogenesis in heavier patients with liver cirrhosis. *Hepatol. Res.*
35, 204–214 (2006).

45. T. Kawaguchi, *et al.*, Branched-chain amino acids prevent hepatocarcinogenesis and
prolong survival of patients with cirrhosis. *Clin. Gastroenterol. Hepatol.* **12**, 1012-1018.e1
(2014).

46. A. Ally, *et al.*, Comprehensive and Integrative Genomic Characterization of Hepatocellular
Carcinoma. *Cell* **169**, 1327-1341.e23 (2017).

47. T. Li, *et al.*, O-GlcNAc Transferase Links Glucose Metabolism to MAVS-Mediated Antiviral
Innate Immunity. *Cell Host Microbe* **24**, 791-803.e6 (2018).

48.   W. Zhang, *et al.*, Lactate Is a Natural Suppressor of RLR Signaling by Targeting MAVS. *Cell*
178, 176-189.e15 (2019).

49.   Z. Cao, *et al.*, Pyruvate Carboxylase Activates the RIG-I-like Receptor-Mediated Antiviral
Immune Response by Targeting the MAVS signalosome. *Sci Rep* **6**, 22002 (2016).

50.   L. A. J. O'Neill, A Broken Krebs Cycle in Macrophages. *Immunity* **42**, 393–394 (2015).

51.   E. L. Mills, *et al.*, Succinate Dehydrogenase Supports Metabolic Repurposing of
Mitochondria to Drive Inflammatory Macrophages. *Cell* **167**, 457-470.e13 (2016).

52.   A. Littlewood-Evans, *et al.*, GPR91 senses extracellular succinate released from
inflammatory macrophages and exacerbates rheumatoid arthritis. *J. Exp. Med.* **213**, 1655–
1662 (2016).

53.   L. Cai, *et al.*, Functional impairment in circulating and intrahepatic NK cells and relative
mechanism in hepatocellular carcinoma patients. *Clin. Immunol.* **129**, 428–437 (2008).

54.   Y. T. Bryceson, M. E. March, D. F. Barber, H.-G. Ljunggren, E. O. Long, Cytolytic granule
polarization and degranulation controlled by different receptors in resting NK cells. *J Exp*
Med **202**, 1001–1012 (2005).

55.   C. Scholtes, *et al.*, Enhancement of genotype 1 hepatitis C virus replication by bile acids
through FXR. *J. Hepatol.* **48**, 192–199 (2008).

56.   F. A. Ran, *et al.*, Genome engineering using the CRISPR-Cas9 system. *Nature Protocols* **8**,
2281–2308 (2013).

57.   Y. Kuang, S. J. Schomisch, V. Chandramouli, Z. Lee, Hexokinase and glucose-6-phosphatase
activity in woodchuck model of hepatitis virus-induced hepatocellular carcinoma. *Comp.*
Biochem. Physiol. C Toxicol. Pharmacol. **143**, 225–231 (2006).

58.   C. Ramière, *et al.*, Activity of hexokinase is increased by its interaction with hepatitis C
virus protein NS5A. *J. Virol.* **88**, 3246–3254 (2014).

59.   L. Perrin-Cocon, *et al.*, Toll-like Receptor 4-Induced Glycolytic Burst in Human Monocyte-
Derived Dendritic Cells Results from p38-Dependent Stabilization of HIF-1α and Increased
Hexokinase II Expression. *J. Immunol.* **201**, 1510–1521 (2018).

60.   J. Payne, J. G. Morris, Pyruvate carboxylase in Rhodopseudomonas spheroides. *J. Gen.*
Microbiol. **59**, 97–101 (1969).

61.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

62.   P. Skehan, *et al.*, New colorimetric cytotoxicity assay for anticancer-drug screening. *J. Natl.*
Cancer Inst. **82**, 1107–1112 (1990).

63.   C. Scholtes, *et al.*, High plasma level of nucleocapsid-free envelope glycoprotein-positive
691   lipoproteins in hepatitis C patients. *Hepatology* **56**, 39–48 (2012).

692   64.   V. Icard, *et al.*, Secretion of hepatitis C virus envelope glycoproteins depends on assembly
693   of apolipoprotein B positive lipoproteins. *PLoS ONE* **4**, e4233 (2009).

694   65.   N. Sonnenschein, M. Geertz, G. Muskhelishvili, M.-T. Hütt, Analog regulation of metabolic
695   demand. *BMC Syst Biol* **5**, 40 (2011).

696   66.   N. Sonnenschein, *et al.*, A network perspective on metabolic inconsistency. *BMC Syst Biol*
697   **6**, 41 (2012).

698   67.   C. Knecht, C. Fretter, P. Rosenstiel, M. Krawczak, M.-T. Hütt, Distinct metabolic network
699   states manifest in the gene expression profiles of pediatric inflammatory bowel disease
700   patients and controls. *Sci Rep* **6**, 32584 (2016).

701   68.   K. Schlicht, *et al.*, The metabolic network coherence of human transcriptomes is associated
702   with genetic variation at the cadherin 18 locus. *Hum. Genet.* **138**, 375–388 (2019).

703   69.   P. Nyczka, M.-T. Hütt, Generative network model of transcriptome patterns in disease
704   cohorts with tunable signal strength. *Phys. Rev. Research* **2**, 033130 (2020).

705   70.   J. Gao, *et al.*, Integrative Analysis of Complex Cancer Genomics and Clinical Profiles Using
706   the cBioPortal. *Sci Signal* **6**, pl1 (2013).

707   71.   E. Cerami, *et al.*, The cBio Cancer Genomics Portal: An Open Platform for Exploring
708   Multidimensional Cancer Genomics Data. *Cancer Discov* **2**, 401–404 (2012).

Figure 1. Correlation between hexokinase expression levels in HCC tumors and patient survival.
a Kaplan–Meier estimates of the survival of HCC patients depending on the expression of HK1,
HK2, HK3 and GCK (HK4) genes in tumor biopsies (n=365; TCGA expression data retrieved from
cBioPortal)(70, 71). Duplicate analyses from the same patient were removed as well as patients
who died when biopsied (overall survival=0 months). Optimal stratification based on highest and
lowest gene expression values was determined using Protein Atlas database (18). **b** Same as above
but patients were stratified based on the GCK/HK2 gene expression ratio. The stratification
showing the lowest *p* value when comparing subgroups of patients with the highest to the lowest

GCK/HK2 expression ratio is displayed. Patient TCGA-DD-AAE9 exhibiting undetectable levels of
GCK and HK2 was removed from this analysis as the GCK/HK2 ratio could not be calculated. **c**
Correlations between patient survival, GCK expression and HK2 expression. Spearman's rank
correlation test on the subset 130 patients for whom the period between diagnosis and death is
precisely known (uncensored data).

Figure 2. Hexokinase isoenzyme switch in Huh7 cells induces extended modifications of
metabolic connections. a Western-blot analysis of HK1, HK2, HK3 and GCK expression in Huh7
and Huh7-*GCK⁺/HK2⁻*. **b** Hexokinase activity in homogenates of Huh7 and Huh7-*GCK⁺/HK2⁻* cells. **c**
Number of genes changing their expression pattern in Huh7 and Huh7-*GCK⁺/HK2⁻* cells (see
Supplementary Table 2 for details). **d** Heatmap showing clustering enrichment scores of the
networks obtained when mapping differentially expressed genes to the human metabolic model
Recon2. Clustering enrichment scores from the highest in red to the lowest in blue were
calculated for different gene expression thresholds ($Log_2|FC|$) and percentages of retained
currency metabolites. **e** Gene network corresponding to the maximal clustering enrichment score
($Log_2|FC|>3$; removed currency metabolites = 2%). The transcription of nodes in green was
upregulated and those in red downregulated in Huh7-*GCK⁺/HK2⁻* compared to Huh7 cells. Plain

a

Molecular and Cellular Functions		
Name	*p*-value range	Number of genes involved
Cellular Movement	$7.68 \times 10^{-6} - 4.66 \times 10^{-25}$	701
Lipid Metabolism	$2.00 \times 10^{-6} - 3.12 \times 10^{-14}$	414
Molecular Transport	$5.63 \times 10^{-6} - 3.12 \times 10^{-14}$	361
Small Molecule Biochemistry	$5.63 \times 10^{-6} - 3.12 \times 10^{-14}$	462
Protein Synthesis	$1.87 \times 10^{-6} - 3.42 \times 10^{-14}$	199

b

Cellular Movement Functional Annotations		
Name	*p*-value	Number of genes involved
Migration of cells	4.66×10^{-25}	585
Cell movement	3.48×10^{-24}	642
Cell movement of blood cells	1.52×10^{-18}	276
Leucocyte migration	1.59×10^{-18}	274
Invasion of cells	5.33×10^{-17}	306

c

Lipid Metabolism Functional Annotations		
Name	*p*-value	Number of genes involved
Concentration of lipid	3.12×10^{-14}	256
Synthesis of lipid	2.37×10^{-10}	233
Fatty acid metabolism	4.31×10^{-10}	166
Quantity of steroid	4.48×10^{-09}	138
Concentration of cholesterol	6.58×10^{-08}	94

743
744
Figure 3. Huh7-*GCK$^+$/HK2$^-$* cells have a higher migration capacity and lipid droplets content. a
Analysis of differentially expressed genes in the two cell lines using gene set enrichment analysis
(|FC|>2 with a *p* value <0.05). Top five enriched molecular and cellular functions are presented.
b and **c** Ranked IPA-annotations associated with 'cellular movement' and 'lipid metabolism'. **d, e**

749 Results of transwell-migration tests. **d** Representative images and **e** count of migrating cells. **f** Oil
750 Red-O staining of lipid droplets (red) with nucleus counterstaining (blue). **g** Quantification of
751 intracellular lipids by FACS after BODIPY staining. E and G correspond to means ± SEM with n≥3
752 (*p* value determined by Student's t-test).
753

Figure 4. Lipogenesis and very-low-density lipoproteins (VLDL) secretion are restored in Huh7-
***GCK$^+$/HK2$^-$* cells. a** Quantification of intracellular lipids in total cell extracts of Huh7 and Huh7-
GCK$^+$HK2$^-$ cells. **b** Lipids and ApoB secretions in supernatants of cells cultured 24h without FCS. **c**
TG/ApoB molar ratio calculated from quantifications determined in b. **d** Supernatants of Huh7
and Huh7-*GCK$^+$/HK2$^-$* were analyzed by ultracentrifugation on iodixanol density gradients. ApoB
was quantified in each fraction by ELISA. Presented data correspond to means ± SEM (n≥3, *p* value
determined by Student's t-test) except for d) that shows one representative experiment.
762

Figure 5. TCA rewiring after hexokinase isoenzyme switch in Huh7 cells. **a** Glycogen quantification. **b** Creatinine and creatinine-P quantification. **c** This bubble chart compares intracellular metabolomes of Huh7 and Huh7-*GCK⁺/HK2⁻* cells. Metabolite pool sizes larger in Huh7 are indicated in blue, whereas the one larger in Huh7-*GCK⁺/HK2⁻* are shown in red. The size of bubbles inversely scales with p values between 5.10^{-2} and 1.10^{-17} of differential metabolomics responses. **d, e** Metabolic fluxes for overall glucose consumption (d) and lactate secretion (e) by Huh7 and Huh7-*GCK⁺/HK2⁻* cells. Indicated values correspond to differences in glucose or lactate concentrations in extracellular culture medium before and after 24h of culture. **f** Mass isotopomer distribution vector of pyruvate in cells cultured with [U-^{13}C]-glucose. **g** Pyruvate carboxylase (PC) activity determined in cell homogenates. **h** Western-blot analysis of PC expression in Huh7 and Huh7-*GCK⁺/HK2⁻* cells. **i**) Western-blot analysis of pyruvate dehydrogenase (PDH) E1-alpha subunit phosphorylation at Ser293. **j** RNAseq quantification of pyruvate dehydrogenase kinase 2 (PDK2) and pyruvate dehydrogenase phosphatase 2 (PDP2)(BH adjusted p value<0.05 from transcriptomic data). **k** Western-blot analysis of ATP-citrate Lyase (ACLY) phosphorylation at Ser455. **l** Succinate quantification in cell homogenates. **m** Succinate dehydrogenase (SDH) activity determined in cell homogenates. **n** Oxygen consumption rate (OCR) in Huh7 and Huh7-*GCK⁺/HK2⁻* cells was determined with a Seahorse analyzer before and after the addition of oligomycin (Complex V inhibitor), FCCP (uncoupling agent), rotenone (Complex I inhibitor) and antimycin A (Complex III inhibitor). **o** Non-mitochondrial, complex I-dependent and complex III-dependent maximal OCR were calculated from n. Data correspond to means ± SEM (n≥3).

Gene ID	Huh7 (mean normalized counts)	Huh7-GCK+/HK2- (mean normalized counts)	FC (calculated by DESeq2 analysis)
EGR1	178,2	537,5	3,0
GBP2	243,0	816,5	3,3
HLA-A	247,2	608,1	2,5
HLA-B	13,4	29,8	2,2
HLA-H	177,8	579,0	3,3
HSP90AB1	0,0	622,9	10000,0
IFI35	50,2	252,5	5,0
IFI6	21,7	142,8	6,6
IFIH1	34,5	106,1	3,1
IFITM3	180,9	1081,4	6,0
IKBKE	0,3	419,4	1284,3
IRF1	212,9	505,2	2,4
IRF3	38,5	84,4	2,2
IRF9	109,3	388,8	3,6
ISG15	89,9	197,6	2,2
MX1	0,0	28,6	10000,0
OAS1	0,8	157,5	173,5
OAS3	195,1	1095,0	5,6
PSMB8	0,0	13,4	80,0
RNASEL	0,0	4,9	10000,0
ABCE1	6559,7	3037,7	-2,2
TRIM6	663,1	263,9	-2,5

Figure 6. Innate immune response is enhanced in Huh7-GCK$^+$/HK2$^-$ cells. a Sector chart from the transcriptomic study showing genes included in the GO-term "Type I-IFN signaling pathway". **b** List of genes significantly up-regulated in purple or down-regulated in green (|FC|>2, BH adjusted *p* value<0.05) in Huh7-*GCK$^+$/HK2$^-$* compared to Huh7 cells. **c-e** Cells were stimulated or not for 48h with 3p-hpRNA (RIG-I ligand) or poly(I:C) (IFIH1/MDA5 ligand). ISRE-luciferase expression was monitored and normalized to Renilla luciferase (c, d). Cell supernatants were assayed for cytokine concentration by multiplex assays (e). **f** NK cell mediated lysis of Huh7 or Huh7-*GCK$^+$/HK2$^-$*. Hepatoma cells were seeded 24h before NK cells addition for 4h at effector to target (E:T) ratio of 0, 3 or 30. After harvesting, cell lysis was determined by the percentage of PI+ cells on gated hepatocytes. *p* values were obtained from 2-way ANOVA analyses comparing matched cell means with Sidak's correction for multiple comparison, with α=0.05.

Figure 7. Simplified scheme of central carbon metabolism and connected pathways showing differences between Huh7-*GCK+/HK2-* vs Huh7. Highlighted metabolites, enzymatic activities, and metabolism-associated genes were selected from transcriptomic (Fig. 2e), metabolomic (Fig. 5c) and enzymatic analyses (Fig. 5g-m).