Different Proteins Mediate Step-wise Chromosome Architectures in *Thermoplasma acidophilum* and *Pyrobaculum calidifontis*

Hugo Maruyama[1][†][*], Eloise I. Prieto[2][†], Takayuki Nambu[1], Chiho Mashimo[1], Kosuke Kashiwagi[3], Toshinori Okinaga[1], Haruyuki Atomi[4], Kunio Takeyasu[5]

[1] Department of Bacteriology, Osaka Dental University, Hirakata, Japan

[2] National Institute of Molecular Biology and Biotechnology, University of the Philippines Diliman, Quezon City, Philippines

[3] Department of Fixed Prosthodontics, Osaka Dental University, Hirakata, Japan

[4] Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Kyoto, Japan

[5] Graduate School of Biostudies, Kyoto University, Kyoto, Japan

[†] These authors have contributed equally to this work

[*] **Correspondence:**

Hugo Maruyama

maruyama@cc.osaka-dent.ac.jp; maruyama.hugo@gmail.com

Keywords: archaea, higher-order chromosome structure, nucleoid, chromatin, HTa, histone, transcriptional regulator, horizontal gene transfer

Running Title: Step-wise chromosome architecture in Archaea

Manuscript length: 5565 words

Number of Figures: 6

Number of Tables: 3

Abstract

Archaeal species encode a variety of distinct lineage-specific chromosomal proteins. We have
previously shown that in *Thermococcus kodakarensis*, histone, Alba, and TrmBL2 play distinct
roles in chromosome organization. Although our understanding of individual archaeal
chromosomal proteins has been advancing, how archaeal chromosomes are folded into higher-
order structures and how they are regulated are largely unknown. Here, we investigated the
primary and higher-order structures of archaeal chromosomes from different archaeal lineages.
Atomic force microscopy of chromosome spreads out of *Thermoplasma acidophilum* and
Pyrobaculum calidifontis cells revealed 10-nm fibers and 30–40-nm globular structures,
suggesting the occurrence of higher-order chromosomal folding. Our results also indicated that
chromosome compaction occurs toward the stationary phase. Micrococcal nuclease digestion
indicated that fundamental structural units of the chromosome exist in *T. acidophilum* and *T.*
kodakarensis but not in *P. calidifontis* or *Sulfolobus solfataricus*. In vitro reconstitution showed
that, in *T. acidophilum*, the bacterial HU protein homolog HTa formed a 6-nm fiber by wrapping
DNA, and that Alba was responsible for the formation of the 10-nm fiber by binding along the
DNA without wrapping. Remarkably, Alba could form different higher-order complexes with
histone or HTa on DNA in vitro. Mass spectrometry detected HTa in the *T. acidophilum*
chromosome but not in other species. A putative transcriptional regulator of the AsnC/Lrp family
(Pcal_1183) was detected on the *P. calidifontis* chromosome, but not on that of other species
studied. Putative membrane-associated proteins were detected in the chromosomes of the three
archaeal species studied, including *T. acidophilum*, *P. calidifontis*, and *T. kodakarensis*.
Collectively, our data show that Archaea use different combinations of proteins to achieve
chromosomal architecture and functional regulation.

1.   Introduction

Genomic DNA needs to be folded properly in the cell to simultaneously accomplish compaction of the genetic material and accessibility to the transcription and replication machinery in all three domains of life. In eukaryotic cells, genomic DNA folding is primarily achieved through the nucleosome, a structure composed of histone octamers with approximately 150 bp of DNA wrapped around them (Luger et al., 1997; Davey et al., 2002). Additional proteins, including linker histone H1 and DNA topoisomerases, have been proposed to contribute to the higher-order folding of the eukaryotic genome (Thomas, 1984; Woodcock et al., 2006; Hizume et al., 2007; Fyodorov et al., 2018). Chromosome structural regulation is also pivotal for cell cycle progression. For example, condensin, a structural maintenance of chromosomes (SMC)-superfamily protein, is required for mitotic chromosome condensation in eukaryotes (Hirano et al., 1997; Ono et al., 2003). Epigenetic modifications also reportedly regulate gene expression (Grewal and Moazed, 2003; Li and Reinberg, 2011). Moreover, recent technological advances in chromosome conformation capture (3C) have shown that the eukaryotic genome is hierarchically folded into chromosomal domains, such as topologically associating domains (TADs) (Gibcus and Dekker, 2013; Barutcu et al., 2017). In bacteria, genome compaction is achieved through the functions of nucleoid-associated proteins (NAPs) including HU, Fis, H-NS, and IHF (Skoko et al., 2006; Sarkar et al., 2009; Wang et al., 2013; Badrinarayanan et al., 2015; Hammel et al., 2016; Japaridze et al., 2017). Non-protein factors, such as macromolecular crowding, also contribute to the higher-order folding and regulation of the bacterial nucleoid (De Vries, 2010). It has also become evident that the bacterial nucleoid is segmented into many highly self-interacting regions called chromosomal interaction domains (CIDs), which are equivalent to the eukaryotic TADs, and the CIDs are further organized into large spatially distinct domains called macrodomains (Dame and Tark-Dame, 2016; Brocken et al., 2018; Verma et al., 2019).

Archaea constitute one of the three domains of life, along with Eukarya and Bacteria. Although Archaea are considered extremophiles that live in extreme environments, including high temperature, high salinity, or low pH (Forterre, 2016), they have also been found in moderate environments, including the marine (Lloyd et al., 2013), soil (Bates et al., 2011), and even the human body (Probst et al., 2013; Koskinen et al., 2017; Nkamga et al., 2017; Pausan et al., 2019). Moreover, Archaea have been associated with human diseases, including periodontal disease (Lepp et al., 2004) and inflammatory bowel disease (Blais Lecours et al., 2014). Thus, understanding how Archaea respond to and influence their environment has become crucial. Archaea encode transcription factors (TFs) resembling bacterial TFs, whereas their transcription and replication machineries are more similar to eukaryotic machineries than bacterial ones (Peeters et al., 2015; Lemmens et al., 2019).

Archaeal chromosomal proteins are diverse (Sandman and Reeve, 2005; Luijsterburg et al., 2008; Driessen and Dame, 2011). For example, most species in Euryarchaeota, one of the major archaeal phyla, encode proteins homologous to eukaryotic histone (Nishida and Oshima, 2017; Henneman et al., 2018) Species in newly proposed phyla such as Nanoarchaeota and Thaumarchaeota also encode histone (Henneman et al., 2018). Exceptions in Euryarchaeota are the members of the order *Thermoplasmatales*, which lack histone and instead encode HTa (**H**istone-like protein of *Thermoplasma acidophilum*), a protein homologous to bacterial HU (DeLange et al., 1981 ; Hocher et al., 2019). Whether HTa has a role similar to that of bacterial

HU or an Archaea-specific function remains elusive. Proteins such as Cren7, CC1, and Sul7d are
specific to species in Crenarchaeota, another major archaeal phylum (Driessen and Dame, 2011).
Alba, a 10-kDa DNA/RNA-binding protein, is found in both Euryarchaeota and Crenarchaeota
(Guo et al., 2003; Laurens et al., 2012), as well as in newly proposed phyla including
Nanoarchaeota, Korarchaeota, and Thaumarchaeota (Sanders et al., 2019). Alba has been shown
to undergo post-translational modification (acetylation), which was proposed to be involved in
transcriptional regulation (Bell et al., 2002). Alba has also been shown to have several different
modes of interaction with DNA, such as DNA stiffening or bridging (Laurens et al., 2012).
However, how Alba cooperates with other chromosomal proteins in higher-order chromosome
folding and whether it plays different roles in Euryarchaeota and Crenarchaeota remain to be
elucidated. Besides these DNA-binding proteins that are usually smaller than 10 kDa, some larger
transcription factor-like proteins also behave as chromosomal proteins in Archaea. TrmB-like 2
(TrmBL2), a protein homologous to the sugar-responsive transcriptional regulator TrmB, is an
abundant chromosomal protein that is completely conserved among *Thermococcales* (Gindner et
al., 2014; Kim et al., 2016). Lines of experimental evidence suggest that TrmBL2 is a
chromosome-associated protein resembling bacterial H-NS, with functions that include filament
formation on DNA and the global suppression of genes (Maruyama et al., 2011; Efremov et al.,
2015).

In the present study, we analyzed the protein composition and structure of chromosomes
from diverse archaeal lineages, aiming to understand the general principles of genome folding.
Species used in this study belong to either Euryarchaeota or Crenarchaeota, two major phyla in
Archaea, and encode different combinations of chromosomal proteins. *Thermococcus*
kodakarensis is a euryarchaeon that encodes histone, Alba, and TrmBL2 as major chromosomal
proteins (Maruyama et al., 2011). *Thermoplasma acidophilum* is a euryarchaeon that is an
exception, as it lacks histones, but it encodes a protein HTa homologous to bacterial HU, together
with Alba (Ruepp et al., 2000). Since it is unclear whether HTa plays roles similar to that of
bacterial HU, we investigated how HTa contributes to genome folding. *Pyrobaculum calidifontis*
is a hyperthermophilic crenarchaeon that encodes Alba, Cren7, CC1, and Sso10a (Amo et al.,
2002; Peeters et al., 2015). *Sulfolobus solfataricus* is a crenarchaeon that encodes Alba, Cren7,
Sul7, and Sso10a (Peeters et al., 2015). Both the primary and higher-order chromosome structures
were analyzed with atomic force microscopy (AFM). Chromosomal proteins in each species were
identified through a combination of chromosome isolation and mass spectrometry. The minimal
structural unit of each chromosome was investigated with micrococcal nuclease (MNase)
digestion analysis and the in vitro reconstitution of protein-DNA complexes.

2.   Results

2.1.   The archaeal chromosome is commonly composed of 10-nm fibers and 30–40-nm
globular structures

We previously showed that, in *T. kodakarensis*, the chromosome is more compact in the
stationary phase than in the log-phase (Maruyama et al., 2011). Here, the extent of chromosome
compaction in two archaeal species from different phyla, *T. acidophilum* and *P. calidifontis*, were
analyzed by using the on-substrate lysis method (see Materials and Methods). Cells were grown

and applied on coverslips, then mildly lysed with detergent. Staining of the lysed cells with 4',6-
diamidino-2-phenylindole (DAPI) showed a higher degree of genomic material released from log-
phase cells than from stationary phase cells for both archaeal species (Figure 1A). In contrast,
compact chromosome structures were dominant after the lysis of stationary phase cells based on
both DAPI staining and AFM (Figure 1B). These results indicate that chromosome compaction
toward the stationary phase commonly occurs in two major archaeal phyla, Euryarchaeaota and
Crenarchaeota.

Detailed AFM analysis of chromosomes released from the lysed cells identified two distinct
chromosomal structures common to both species, 10-nm fiber and 30–40-nm globular structures.
The 10-nm fiber structure was prevalent on the well-spread chromosome in the log-phase in both
species (Figure 2A). The diameter of the fibers was 10.1 ± 4.2 nm (mean $\pm$ SD, n = 119) in *T.*
acidophilum and 11.5 ± 4.2 nm (n = 110) in *P. calidifontis* (Figure 2B). Larger globular structures
were found where chromosomes were less spread in both species in the stationary phase, wherein
most cells showed limited chromosome release (Figure 2C). The diameters of the globular
structures differed between the two species and were specifically 38.6 ± 10.1 nm (n = 217) nm in
T. acidophilum and 29.1 ± 5.1 nm (n = 152) nm in *P. calidifontis* (Figure 2D), suggesting
differences in the fundamental structure or the higher-order architecture of the chromosome.
Taken together, these results show that higher-order folding exists in the archaeal chromosome
and that the extent of chromosome folding is dynamically regulated throughout the growth phase
of the cells.

2.2. Minimal unit of the archaeal chromosome revealed with MNase digestion

On-substrate lysis results revealed that the size of the primary structure of the archaeal
chromosome is approximately 10 nm in diameter (Figure 2A and 2B). We next investigated the
minimal structural unit through MNase digestion of purified chromosomes from four archaeal
species: *T. kodakarensis* and *T. acidophilum* from Euryarchaeota, and *P. calidifontis* and *S.*
solfataricus from Crenarchaeota. The analysis revealed a variation in minimal structural units in
each species (Figure 3). As we have previously shown, MNase digestion of the *T. kodakarensis*
chromosome resulted in a 30-bp laddered pattern with a minimum of 60 bp of DNA (Figure 3A).
The 30-bp intervals detected are typical of some histone-containing Archaea and reflect the
flexible histone multimeric structure, which was shown previously (Maruyama et al., 2013;
Mattiroli et al., 2017; Henneman et al., 2018). Interestingly, MNase digestion of the *T.*
acidophilum chromosome resulted in the accumulation of ~40 bp of DNA, suggesting the size of
DNA involved in the primary unit (Figure 3B). The larger DNA fragment of 80~90 bp identified
with a lower MNase concentration indicated that this unit is positioned at a regular interval. In
contrast, the MNase digestion of the chromosomes of the two crenarchaeal species, *P. calidifontis*
and *S. solfataricus*, did not yield the accumulation of DNA fragments of a particular size (Figure
3C and 3D). These results demonstrate that in crenarchaeal species, which encode neither histone
nor HTa, the chromosomes do not have a defined-size structural unit that protects the genomic
DNA from MNase digestion. Whereas on-substrate cell lysis revealed similar fibrous
chromosome structures among *T. acidophilum* and *P. calidifontis* (Figure 2), as well as in *T.*
kodakarensis, as previously reported (Maruyama et al., 2011), the MNase digestion patterns

indicated that the fine details in the organization of these structures vary, probably due to
differences in the chromosomal proteins involved.

2.3. Mass spectrometry identifies chromosome-associated proteins in archaeal species

Protein components of isolated chromosomes of *T. acidophilum*, *P. calidifontis* and *T.*
kodakarensis were analyzed using mass spectrometry. Tables 1 to 3 provide an overview of the
distribution of chromosomal proteins in each archaeal species. Proteins predicted to be involved
in replication, transcription, and translation were detected in the three species. These include the
DNA primase DnaG detected in *T. acidophilum* and *P. calidifontis* and DNA-directed RNA
polymerase subunits detected in *T. acidophilum* and *T. kodakarensis*. Ribosome subunits were
commonly found in all species and constituted a large portion of the identified proteins, which is
reasonable considering that transcription and translation are coupled in Archaea (French et al.,
2007).

Apart from these general information processing proteins, DNA-binding proteins that might
contribute to chromosome folding and regulation were also detected. Alba was detected in all
three species studied (Tables 1–3). In the *T. acidophilum* chromosome, HTa was detected together
with Alba (Table 1). In the *P. calidifontis* chromosome, a protein annotated as 'Transcriptional
regulator, AsnC family' (tremble ID: A3MVE1, gene ID: Pcal_1183) was detected in addition to
Alba (Table 2). It has been suggested that some AsnC/Lrp family transcriptional regulators have
global gene regulatory roles as well as an architectural function, instead of being a specific
transcriptional regulator (Peeters et al., 2015). The identification of Pcal_1183 in *P. calidifontis*,
which has not been reported as a chromosomal protein, supports the view that TF-like proteins act
as chromosome proteins in Archaea, as is the case for TrmBL2 in *Thermococcales* (Maruyama et
al., 2011). Cren7, although encoded in the *P. calidifontis* genome, was not detected in the present
mass spectrometry analysis, suggesting that the amount of Cren7 on the *P. calidifontis*
chromosome is lower than the detection limit of our method. In *T. kodakarensis*, histones, Alba,
and TrmBL2 were detected as previously reported (Table 3) (Maruyama et al., 2011).
Interestingly, in addition to proteins with apparent DNA-binding ability, a few putative
membrane-associated proteins were detected in the chromosome. For example, V-type ATPases
were found in all three species (Tables 1–3), suggesting a general role of these proteins in
archaeal chromosome folding and regulation.

2.4. Distinct fundamental structures formed with Alba and HTa

Alba was common among the three species studied (Tables 1–3). When recombinant Alba
proteins from *T. acidophilum* or *P. calidifontis* genes expressed in *E. coli* were mixed with linear
3-kb DNA in vitro, a 10-nm fiber structure was formed at sub-saturating concentrations (Figure
4A and 4B). The diameter of the fiber was 10.0 ± 2.8 nm (n = 80) in *T. acidophilum* and $12.1 \pm$
2.2 nm (n = 119) in *P. calidifontis* (Figure 4B). These correlate well with the structures visualized
in the on-substrate lysis experiment (Figure 2A and 2B), suggesting that Alba is responsible for
the formation and maintenance of the 10-nm fiber structures. Closer inspection of the structures

reconstituted with Alba showed that both Alba from *T. acidophilum* and *P. calidifontis* could join two DNA strands (Figure 4A). At higher concentrations, Alba formed filaments without changing the DNA contour length (Figure 4A and 4C), indicating that DNA does not wrap around Alba molecules. These filaments were capable of folding onto themselves and forming more complex structures. The results strongly support a role of Alba protein contributing to primary order folding of the DNA into 10-nm fibers, as well as to more complex, higher-order structures.

Whereas most Euryarchaeota commonly encode histones, species belonging to *Thermoplasmatales* lack histones and encode HTa, a homolog of bacterial HU. It has been proposed that *Thermoplasmatales* acquired the HU gene from bacteria through horizontal gene transfer (HGT) and lost the histone gene and that HTa plays a role similar to that of archaeal histone (Hocher et al., 2019). We investigated how HTa interacts with DNA at the primary level. The reconstitution using recombinant histidine-tagged HTa from *T. acidophilum* showed a beaded structure (Figure 4D), similar to the beads-on-a-string structure formed with archaeal histone (Maruyama et al., 2011; Maruyama et al., 2013). The diameter of the HTa beads was 6.3 ± 1.5 nm ($n = 77$) (Figure 4E), which is thinner than the 10-nm Alba filaments (Figure 4A and 4B). At higher concentrations, HTa formed a filament of 5.5 ± 1.4 nm in width ($n = 221$) on the DNA (Figures 4D and 4F). The contour DNA length of reconstituted nucleoprotein structure was shorter than that of naked DNA, with an average of 60 bp per bead (Figure 4G), indicating the length of DNA wrapped around an HTa particle. These results support the notion that archaeal HTa is functionally more closely related to histones than bacterial HU (Hocher et al., 2019). The ~6-nm particle might be a fundamental unit of the *Thermococcales* chromosome, together with the 10-nm fiber formed with Alba (Figure 4A and 4B).

2.5. Relationship between Alba and other chromosomal proteins in forming structural complexes with DNA

Alba can form various structures with DNA, depending on its amount (Figure 4A). This flexibility might facilitate its interaction with other structural proteins, forming DNA-based complexes. Alba coexists with other DNA-binding proteins in the cell. For example, *T. kodakarensis* encodes histone and Alba, whereas *T. acidophilum* encodes HTa and Alba. We therefore combined Alba with the chromosomal proteins, histone and HTa, to determine how they cooperate in compacting DNA. In vitro reconstitution using histone alone at a 1:1 histone:DNA weight ratio resulted in compact beaded fiber structures displaying decreased contour length of the DNA-protein complex (Figure 5A). Reconstitution using both histone and Alba from *T. kodakarensis* at a 1:3 or 1:7 histone:Alba ratio resulted in the formation of structures somewhat similar to those formed with DNA and histone alone (Figure 5A). However, the resulting complexes had a larger proportion of DNA molecules with a longer contour length (> 500 nm) than that with histone alone. As Alba does not wrap DNA (Figure 4A and 4B), this result suggests that Alba binding decreases the extent of histone-mediated DNA folding or that Alba competes with histone for DNA-binding, leading to less compact structures.

Reconstitution using HTa and Alba from *T. acidophilum* at 2:6 and 2:15 weight ratios (Figure 5B) resulted in structures similar to those formed by Alba alone at sub-saturating amounts (Figure 4A). Structures with DNA strands joined together were dominant, and DNA molecules

were considerably folded (Figure 5B). Moreover, some complexes were able to fold upon
themselves, providing additional compaction (Figure 5B). These data indicate that unlike with
histone (Figure 5A), Alba with HTa facilitates DNA compaction through DNA strand bridging
(Figure 5B). Thus, Alba encoded in each species might play different roles depending on which
proteins coexist on the chromosome in each archaeal lineage. Binding of HTa might increase the
efficacy of Alba-mediated DNA bridging, leading to complexes that are more compact than those
with either HTa or Alba alone. Whether this compaction is achieved through direct protein-
protein interaction of HTa and Alba needs to be elucidated.

3.   Discussion

In the present study, we investigated the fundamental and higher-order structures of archaeal
chromosomes from different lineages. We show that archaeal chromosomes comprise step-wise
structures that consist of primary fibrous structures of 10 nm in diameter and higher-order
globular structures of 30–40 nm. The minimal unit of the chromosome varies between species.
Archaeal chromosomes undergo compaction toward the stationary phase (Figure 6A). We also
show that archaeal HTa can wrap DNA and form a 6-nm beaded fiber, which might be folded into
the higher-order structures in association with Alba in *T. acidophilum*. Interestingly, TF-like
proteins were found in *T. kodakarensis* and *P. calidifontis*. Membrane-associated proteins were
detected in the chromosomes of all three species. Finally, we demonstrate that Alba-mediated
chromosomal structures vary depending on the presence and identity of additional architectural
proteins.

3.1 Primary and higher-order folding of archaeal chromosomes

Higher-order chromosome architecture and its regulation have been extensively studied in
bacteria and eukaryotes. In bacteria, the coordination of various NAPs is required for proper
folding and regulation of the nucleoid (Verma et al., 2019). For example, the *Escherichia coli*
nucleoid is composed of 10-, 40-, and 80-nm fibers and undergoes compaction in the stationary
phase, in which Dps is highly expressed (Ali Azam et al., 1999; Kim et al., 2004). Oxidative
stress also causes Dps-dependent nucleoid compaction in *Staphylococcus aureus* (Morikawa et
al., 2019). In eukaryotes, it has been viewed that beads-on-a-string chromatin is folded into a 30-
nm fiber with the linker histone H1 (Hizume et al., 2005; Prieto and Maeshima, 2019). Our results
indicate that archaeal chromosomes also comprise step-wise structures, sharing features with both
bacterial and eukaryotic chromosomes. A hierarchical mode of archaeal genome packing can be
inferred wherein the 6- or 10-nm fiber structures form the 30–40-nm globular structures. The 6-
or 10-nm fibers might represent the first order of folding in the archaeal chromosome, wherein
DNA is bound and constrained by one or more abundant chromosomal protein(s). The abundant
chromosomal proteins present in each species and their interactions with each other might
contribute to species-specific primary and higher-order chromosome structures.

We show that the archaeal chromosome undergoes compaction toward the stationary phase,
similarly to what happens in the bacterial nucleoid. However, how compaction is achieved
remains elusive. SMC-superfamily proteins are known to be involved in higher-order architecture
and the regulation of chromosomes in eukaryotes and bacteria (Uhlmann, 2016). The 3C
experiments on *Sulfolobus* species showed that an SMC-family protein contributes to the
formation of distinct chromosome domains and the higher-order interactions of crenarchaea

(Takemata et al., 2019). Although SMC homologs are present in archaeal genomes based on
genomic sequences, they were not detected using mass spectrometry. Low amounts of SMC
proteins might be sufficient for their function in archaeal cells, probably cooperating with other
chromosomal proteins to achieve higher-order folding of the chromosome (Figure 6). Post-
translational modifications of archaeal DNA-binding proteins such as Alba and Cren7 have been
reported (Bell et al., 2002; Niu et al., 2013). Detailed analyses of the influence of such post-
translational modifications on protein-protein and protein-DNA interactions might also provide
insights into mechanisms of chromosome folding and regulation.

3.2 Role of HTa in *T. acidophilum*

HTa is an abundant protein in members of *Thermococcales*, which lack histones. Our present
study shows that *T. acidophilum* HTa can form a beads-on-a-string structure by wrapping DNA
and that the wrapping protects DNA from MNase digestion (Figure 3B and 4D). Taken together
with a recently reported analysis of HTa (Hocher et al., 2019), these results indicate that HTa has
characteristics more similar to those of archaeal histone than to those of bacterial HU. Moreover,
HTa was recently reported to form oligomers (Hocher et al., 2019) resembling the flexible
"hypernucleosome" formed by multiple archaeal histone dimers stacked onto each other
(Maruyama et al., 2013; Mattiroli et al., 2017; Henneman et al., 2018). Thus, it seems that for
species in Euryarchaeota, a protein that fulfills the DNA wrapping function (i.e., histone or HTa)
seems to be indispensable for the proper folding and regulation of the chromosome. Based on the
results of the MNase assay (Figure 3C and 3D), we assume that most crenarchaeal species do not
utilize DNA wrapping as the primary level of DNA folding (Figure 6B), although a minority of
crenarchaea encode histone that could wrap DNA (Henneman et al., 2018). Why most
crenarchaeal chromosomes do not require DNA wrapping is an interesting question that needs to
be answered in the future.

The phylogenetic reconstitution of the HTa/HU gene family has suggested that HTa was
acquired through the horizontal transfer of bacterial HU at the root of the clade which includes
Thermoplasmatales (Hocher et al., 2019). From an evolutionary aspect, an intriguing question is
how histone could be replaced by a protein with apparently different characteristics; for example,
archaeal histone wraps DNA, whereas bacterial HU bends it (Henneman et al., 2018; Verma et
al., 2019). Although some species have been shown to tolerate the complete deletion of histone
gene(s) from their genome (Weidenbach et al., 2008), complete deletion of histone genes seems
to be impossible in most species (Čuboňová et al., 2012). Besides, since all species in
Euryarchaeota (except for *Thermoplasmatales)* encode histone, loss of histone does not seem to
be favorable in the living environments. Thus, it seems unlikely that the loss of histone occurred
prior to the acquisition of HU. Interestingly, the expression of archaeal histone in *E. coli* does not
cause a severe growth defect (Rojec et al., 2019), indicating that histone and HU can coexist in a
single organism. These facts are consistent with the idea that the horizontal transfer of bacterial
HU to Archaea and its adaptation might have occurred before the loss of the histone gene.

3.3 Role of transcription factor-like proteins in *P. calidifontis* and *T. kodakarensis*

Our present results showed the abundance of a putative TF-like protein in the *P. calidifontis*
chromosome. Several archaeal TFs and TF-like proteins have been found to combine a global
gene regulatory role with an architectural role (Maruyama et al., 2011; Peeters et al., 2015). For
example, TrmBL2 (TK0471) in *T. kodakarensis* has a helix-turn-helix motif with non-specific
DNA-binding ability (Maruyama et al., 2011; Ahmad et al., 2015). Single-molecule studies have
shown that TrmBL2 forms a stiff nucleoprotein filament and that histone competes with TrmBL2,
modulating the stiffness of the filament (Efremov et al., 2015). The functions of TrmBL2
resemble those of bacterial H-NS, a NAP that suppresses the transcription of horizontally
acquired genetic elements by forming a nucleoprotein filament on genomic DNA (Navarre et al.,
2006; Oshima et al., 2006; Higashi et al., 2016). Considering the functional similarity between H-
NS and TrmBL2 and the fact that HGT occurs between bacteria and Archaea (López-García et
al., 2015; Wagner et al., 2017), TrmBL2 might function in suppressing horizontally transferred
genes (HTGs) in Archaea (Maruyama et al., 2019). Proteins homologous to TrmBL2 have been
found in Archaea other than members of the *Thermococcales*, as well as in bacteria (Maruyama et
al., 2011). The discovery of Pcal_1183, a TF that belongs to the AsnC/Lrp family, as an abundant
chromosomal protein in *P. calidifontis*, adds another candidate HTG suppressor to Archaea.
Searching for abundant chromosomal proteins with the method used in the present study would
facilitate the identification of novel TF-like nucleoid proteins possibly involved in chromosomal
folding or the suppression of HTGs both in bacteria and Archaea in the future.

3.4 Conclusions and perspectives

In conclusion, although the chromosomes of Euryarchaeota and Crenarchaeota are commonly
folded into higher-order structures, their structural units are fundamentally different. Specifically,
DNA wrappers such as histones or HTa are necessary for Euryarchaeota, whereas DNA wrapping
as a primary DNA folding is missing in Crenarchaeota (Figure 6A and 6B). Both Euryarchaeota
and Crenarchaeota use Alba, which basically functions as a DNA bridger. The function of Alba
might be different depending on what other proteins are present in the cell. TF-like proteins are
found in both Euryarchaeota and Crenarchaeota. Possible functions of such TF-like proteins
might include architectural roles as well as the suppression of horizontally transferred genetic
elements (Maruyama et al., 2019). Further, the interplay between the chromosomal proteins, as
well as differential expression and post-translational modifications, might contribute to higher-
order folding and regulation of the archaeal chromosomes (Figure 6A).

We also propose that membrane-associated proteins might play roles in chromosome folding
and regulation. In the present study, membrane-associated proteins, together with proteins
involved in translation, transcription, and replication, were detected in archaeal chromosomes.
The association between membrane-bound proteins and chromosomes has been proposed in both
eukaryotes and bacteria (Maruyama et al., 2014; Chang et al., 2015; Monterroso et al., 2019). In
eukaryotes, the linker of nucleoskeleton and cytoskeleton (LINC) complex has been shown to
interconnect the chromosome, nuclear membrane, and cytoskeletal filaments, performing diverse
functions, including mechanotransduction and meiotic chromosome movements in mice, yeast,
and nematodes (Hiraoka and Dernburg, 2009; Sato et al., 2009; Chang et al., 2015). In *E. coli*, the
actin-like protein MreB, a member of a membrane-bound complex (MreBCD), is associated with
the nucleoid, playing a role in bacterial chromosome segregation (Thanbichler et al., 2005). The

present findings suggest that membrane-associated proteins might be involved in genome
architecture and dynamics in all three domains of life.

4.   Materials and methods

4.1.   Strain and growth conditions

Thermoplasma acidophilum strain DSM 1728 was grown at 58 °C under aerobic conditions in
modified Allan's basal salt medium (in 1 L: 0.2 g $(NH_4)_2SO_4$, 3.0 g KH_2PO_4, 0.5 g $MgSO_4 \cdot 7H_2O$,
0.25 g $CaCl_2 \cdot 2 H_2O$) adjusted to pH 1.5 with H_2SO_4 and supplemented with 0.1% yeast extract
and 1% glucose (Searcy, 1975). *Pyrobaculum calidifontis* strain VA1 was cultivated in the TY
medium containing 0.3% sodium thiosulfate pentahydrate at 90 °C (Pelve et al., 2012).
Thermococcus kodakarensis strain KOD1 was cultured as previously described (Maruyama et al.,
2011). *Sulfolobus solfataricus* strain P1 was cultured at 80 °C in *Sulfolobus solfataricus* medium
171 specified by the Japan Collection of Microbes.

4.2.   Nuclear staining

Live-cell staining of DNA was performed according to a previous study (Poplawski and
Bernander, 1997). Briefly, exponentially growing (log) and stationary phase cells were incubated
with 1–2 µg/ml of DAPI for 1 hour. These were then harvested, mounted on slides with agarose
pads, and visualized using epi-fluorescence microscopy.

4.3.   On-substrate lysis

On-substrate lysis of archaeal cells was performed as previously described (Maruyama et al.,
2011; Ohniwa et al., 2018) with slight modifications. *T. acidophilum* and *P. calidifontis* cells at
the log and stationary phases were harvested, washed, and resuspended to a final density of
approximately 0.5 OD_{600}. The cell suspension was then applied to a round coverslip (15-mm
round; Matsunami Glass Ind., Kishiwada, Japan) and dried with nitrogen gas. The cells were
lysed with the addition of lysis buffer (10 mM tris pH 6.8, 10 mM EDTA, 0.5% Triton X-100)
onto the coverslip and then dried again with nitrogen gas. The sample was imaged with AFM. To
assess the extent of chromosome spreading, samples were stained with DAPI.

4.4.   Atomic force microscopy

Samples were imaged in the air with the tapping mode of Nanoscope IIIa or IV (Digital
Instruments, Santa Barbara, CA) using an OMCL AC160TS probe (Olympus, Tokyo, Japan).
Images were captured in a 512×512-pixel format, plane-fitted, and flattened using the computer
program supplied in the imaging module. Measurements were made, and correction for the "tip-
effect" was performed accordingly (Maruyama et al., 2011). Dimensions of the chromosomal

structures were calculated based on the measured bottom-to-bottom distance of particles
(apparent measured distance) using the previously described equation $S = 0.75W - 16.4$, where S
is the corrected width of the sample and W is the apparent measured distance (Ohniwa et al.,
2007).

4.5. Isolation of archaeal chromosome

Chromosomes from *T. acidophilum*, *P. calidifontis*, and *T. kodakarensis* were obtained as
described previously with some modifications (Maruyama et al., 2011). Cells were harvested at
log or stationary phase and lysed for 10 minutes at 4 °C with extraction buffer (25 mM HEPES,
pH 7.0, 15 mM $MgCl_2$, 100 mM NaCl, 0.4 M sorbitol, 0.5% Triton X-100). The lysate was then
layered on top of a pre-cooled linear sucrose gradient (15–45% sucrose) in 10 mM tris pH 6.8 and
3 mM $Mg(CH_3COO)_2$. The samples were centrifuged in a swinging-bucket MLS-50 rotor
(Beckman) at $10,000 \times g$ and 4 °C for 20 minutes. The DNA-enriched material, visible as an
opalescent band, was collected, washed with MNase buffer (20 mM Tris pH 8.0, 5 mM NaCl, 2.5
mM $CaCl_2$), and stored at −20 °C until further use.

4.6. MNase digestion of archaeal chromosome

T. acidophilum, *P. calidifontis*, and *T. kodakarensis* genomic DNA was incubated with MNase
(Worthington Biochemical, Lakewood, NJ) at various concentrations (0.3, 1, 3, 10 and 30 U
MNase/100 µl) for 20 minutes at 37 °C. The digestion reaction was stopped by adding stop
solution (100 mM EDTA, 100 mM EGTA, 2% SDS). Chromosomal fragments were then purified
with phenol/chloroform extraction and ethanol precipitation. The DNA pellet was dissolved in TE
buffer (10 mM tris pH 7.5, 1 mM EDTA), resolved with a 2.5% agarose gel, and visualized with
ethidium bromide staining.

4.7. Mass spectrometry

Chromosomes were isolated as described previously herein, digested with MNase (Worthington
Biochemical Corp., Lakewood, NJ), and run on a 15% SDS-PAGE gel to separate proteins
associated with archaeal chromosomes. After staining with Coomassie Brilliant Blue, visible
protein bands were excised from the gel and processed for liquid chromatography coupled with
tandem mass spectrometry (LC-MS/MS) as described previously (Maruyama et al., 2011). Top
hits of each band are summarized in Tables 1–3.

4.8. Expression and purification of recombinant proteins

455 Genes encoding the protein Alba from *T. acidophilum* and *P. calidifontis* and HTa from *T.*
456 *acidophilum* were PCR-amplified from their respective genomic DNA template and inserted into
457 a pET16b vector containing 10x histidine tag at the N-terminus (Novagen, Madison, WI). The

following primers were used: for *T. acidophilum* Alba, forward 5'-ATCGCCATGGCAGAGGAGAACATAATCTTTG-3' and reverse 5'-ATCGGGATCCTCAACGTGACAGC-3'; for *P. calidifontis* Alba, forward 5'-ATCGCCATGGCGACAGAACAGACAATAC-3' and reverse 5'-ATCGGATCCTTAGGCCATTTCGAGCACT-3'; for *T. acidophilum* HTa, forward 5'-ATCGCATATGTAGGAATCAGTGAGCTAT-3' and reverse 5'-ATCGGGATCCTTACTGCTGGTATTTTATCTTGC -3'. The Alba gene was inserted using the NcoI and BamHI restriction sites, resulting in the removal of the his-tag. HTa gene was inserted via the NdeI and BamHI sites and thus tagged with histidine. Competent *E. coli* BL21-CodonPlus (DE3)-RIL was transformed with the resulting constructs. Protein expression was induced via the addition of IPTG. Recombinant proteins were then extracted and purified. Alba proteins were run on an ion-exchange column (HiTrap SP column, 1 ml, GE Healthcare UK Ltd. Buckinghamshire, England) following the manufacturer's protocol and as described previously (Maruyama et al., 2011). The His-tagged HTa was purified with Ni-NTA agarose beads (Qiagen) following the manufacturer's protocol. Histones from *T. kodakarensis* were expressed and purified as previously described (Maruyama et al., 2011).

4.9. In vitro reconstitution

The plasmid pBlueScriptII (Stratagene) was linearized through restriction enzyme digestion with Hind III. Each recombinant protein was mixed with the 3-kb linearized DNA at various DNA-to-protein weight ratios (1:0, 1:2, 1:6, 1:15) in 10 mM tris pH 6.8 and 200 mM NaCl. The mixture was heated at appropriate temperatures, 58 °C for *T. acidophilum* and 90 °C for *P. calidifontis*, and then processed for AFM imaging. For AFM observation, the samples were diluted 1:20 with AFM fixation buffer (10 mM tris pH 8.0, 5 mM NaCl, 0.3% glutaraldehyde) and incubated for 30 minutes at 25 °C. The protein-DNA complexes were then deposited onto a freshly cleaved mica surface pretreated with 10 mM spermidine. After 10 minutes, the mica was washed with 1 ml of pure water, dried with nitrogen gas, and observed with AFM. Reconstitution experiments using a combination of different chromosomal proteins were also performed by mixing the 3-kb linearized DNA with the protein pair of interest at specified amounts. These were then processed for AFM imaging.

5. Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6. Funding

This work was supported by a JSPS KAKENHI Grant-in-Aid for Young Scientist (B), Grant numbers 26850057 and 17K15254, to HM.

7. Acknowledgments

We would like to thank Mr. Yuzo Watanabe (Proteomics Facility, Graduate School of Biostudies,
Kyoto University) and Dr. Ryosuke Ohniwa (Faculty of Medicine, University of Tsukuba) for
mass spectrometry assistance and Editage (www.editage.com) for English editing.

8.   Author contributions

HM and KT conceived and designed the study. EP performed experiments, analyzed data, and
wrote the first draft of the manuscript. TN, CM, and TO wrote sections of and gave critical
comments on the manuscript. HM, EP, HA, and KT revised the manuscript.

 Figure legends

Figure 1. Genomic material released from *Thermoplasma acidophilum* and *Pyrobaculum*
calidifontis cells. DAPI staining was performed after on-substrate lysis of the cells during (A) log
and (B) stationary phases. (A) Chromosomes were well-spread in the log-phase for both *T.*
acidophilum (upper panel) and *P. calidifontis* (lower panel). (B) The stationary phase
chromosome was more compact than the log-phase chromosome (arrow). Scale bars: 2 µm.

Figure 2. Structural details of genomic materials from *Thermoplasma acidophilum* and
Pyrobaculum calidifontis after on-substrate lysis visualized with atomic force microscopy (AFM).
(A-B) Extensively released chromosome fibers of log-phase cells. (A) AFM of the released
genomic materials. (B) Histograms indicate the diameters of the released structures. (C-D)
Compact chromosome of stationary phase cells. (C) AFM of the lysed cells. Boxed areas in the
upper panels were investigated in detail in the lower panels. Globular structures were observed
(arrows) (D) Histograms indicate the diameters of the globular structures. Scale bars: 200 nm.

Figure 3. Digestion pattern of archaeal chromosomes with micrococcal nuclease (MNase).
Purified chromosomes of (A) *Thermococcus kodakarensis*, (B) *Thermoplasma acidophilum*, (C)
Pyrobaculum calidifontis, and (D) *Sulfolobus solfataricus* were digested with increasing
concentrations of MNase and separated with 2.5% agarose gels in 1X TBE. The accumulation of
DNA of particular sizes (arrows) was observed with *T. kodakarensis* and *T. acidophilum* but not
with *P. calidifontis* and *S. solfataricus* chromosomes. MNase concentration was 0.3, 1, 3, 10 U
MNase in 100 µl reaction (A) or 0, 0.3, 1, 3, 10 and 30 U MNase in 100 µl reaction (B-D).

Figure 4. In vitro reconstitution of chromosome structures with recombinant Alba and HTa. (A-C)
Reconstitution on a linear 3-kb plasmid using recombinant Alba from *Thermoplasma*
acidophilum and *Pyrobaculum calidifontis* at different Alba to DNA weight ratios. (A) Atomic
force microscopy (AFM) showing various structures formed with Alba and DNA. Alba:DNA
weight ratios are indicated. (B) Histograms show the diameters of the fibrous structures formed,
as well as (C) Respective contour lengths of DNA at a 15:1 Alba:DNA ratio. The theoretical
length of a linear 3-kb DNA in B-form is ~1000 nm (Maruyama et al., 2011). (D-G)
Reconstitution with histidine-tagged HTa from *T. acidophilum* at varying protein to DNA ratio.
(D) AFM showing beaded (arrows) and filamentous (triangles) structures. (E) Diameter of the
beads formed at a relatively lower HTa concentration. (F) Width of the filaments formed at
relatively higher HTa concentration. (G) Histogram shows the contour DNA lengths of the
structures formed at a 15:1 HTa:DNA ratio. Scale bars: 100 nm.

Figure 5. In vitro reconstitution using a combination of archaeal proteins. Chromosomal
structures were reconstituted on a linear 3-kb plasmid using recombinant proteins at varying Alba
concentrations. (A) Atomic force microscopy (AFM) showing reconstitution using histone and
Alba from *Thermococcus kodakarensis*. Histograms below the AFM images show the contour
DNA length in each condition. (B) AFM showing reconstitution using HTa and Alba from
Thermoplasma acidophilum. Scale bars: 100 nm.

Figure 6. Model of archaeal chromosome folding. (A) In Euryarchaeota, histone or HTa wraps
genomic DNA, whereas there is no DNA wrapper in Crenarchaeota. Alba can bridge distant
genomic DNA. Chromosome compaction occurs toward the stationary phase, where 30~40-nm
globular structures are formed. (B) Schematic phylogenetic tree of selected Archaea (Sandman
and Reeve, 2005) together with the distributions of histone and HTa. Species in Euryarchaeota
utilize histone or HTa (*Thermoplasmatales*) to wrap DNA at the primary level. Members of
Nanoarchaeota and Thaumarchaeota also encode histone that might wrap DNA. Species in
Crenarchaeota neither encode histone nor wrap DNA. The branch lengths do not correspond to
evolutionary distances.

Table 1. Proteins identified by mass spectrometry from isolated chromosomes of *Thermoplasma*
acidophilum. Protein and gene information based on the UniProt database (The UniProt
Consortium, 2019).

Protein ID	Protein description	Mass (kDa)	Gene	Gene locus
Q03587	DNA-directed RNA polymerase subunit B	134.7	rpoB	Ta0390
Q9HLR8	DNA double-strand break repair Rad50 ATPase	103.4	rad50	Ta0157
Q9P997	V-type ATP synthase alpha chain	84.8	atpA	Ta0004
O05209	VCP-like ATPase	83.1	vat	Ta0840
P26752	Elongation factor 2	81.2	fusA	Ta0446
Q9HL99	Threonine--tRNA ligase	76.1	thrS	Ta0330
P50023	Chaperone protein DnaK	66.4	dnaK	Ta1087
Q9HJ60	Probable translation initiation factor IF-2	65.4	infB	Ta1112
Q9HJM5	Glutamate--tRNA ligase	63.5	gltX	Ta0942
P48424	Thermosome subunit alpha	58.3	thsA	Ta0980
P57693	Phenylalanine--tRNA ligase alpha subunit	57.5	pheS	Ta0639
Q03586	DNA-directed RNA polymerase subunit A''	56.6	rpoA2	Ta0392
Q9HM64	V-type ATP synthase beta chain	51	atpB	Ta0005
Q9HLX7	DNA primase DnaG	50	dnaG	Ta0097
P19486	Elongation factor 1-alpha	47	tuf	Ta0444
Q9HM12	S-adenosylmethionine synthase	44.4	mat	Ta0059
Q9HM67	V-type ATP synthase subunit C	41.4	atpC	Ta0002
Q9HIQ9	50S ribosomal protein L3	37.6	rpl3	Ta1271
Q9HKS5	Pyridoxal 5'-phosphate synthase subunit PdxS	36.5	pdxS	Ta0522
P57692	50S ribosomal protein L10	34.6	rpl10	Ta0359
Q9HJA6	Riboflavin kinase	25	ribK	Ta1064
Q9HIR5	30S ribosomal protein S3	24.7	rps3	Ta1265
Q9HL71	50S ribosomal protein L1	23.9	rpl1	Ta0360
P25156	Proteasome subunit alpha	25.8	psmA	Ta1288
Q9HM16	Translation initiation factor 6	23.5	eif6	Ta0055
Q9HJD7	30S ribosomal protein S4	22.9	rps4	Ta1032

P28061	Proteasome subunit beta	23.1	psmB	Ta0612
P57712	30S ribosomal protein S2	22.5	rps2	Ta1190
Q9HJ11	PyrE-like protein	21.9	Ta1164	Ta1164
Q9HLY1	30S ribosomal protein S7	20.5	rps7	Ta0092
Q9HIS3	50S ribosomal protein L6	19.6	rpl6	Ta1255
Q9HJB3	50S ribosomal protein L10e	19.4	rpl10e	Ta1057
P60746	50S ribosomal protein L24	16	rpl24	Ta1260.1
Q9HIS8	50S ribosomal protein L30	18.7	rpl30	Ta1250
Q9HL70	50S ribosomal protein L11	16.4	rpl11	Ta0361
Q9HIR4	50S ribosomal protein L22	16.9	rpl22	Ta1266
Q03589	Probable transcription termination protein NusA	16.8	nusA	Ta0393
Q9HIS9	50S ribosomal protein L15	15.8	rpl15	Ta1249
P02345	DNA-binding protein HTa	10.1	Ta0093	Ta0093
Q9HJQ5	DNA/RNA-binding protein Alba	9.9	albA	Ta0912

555   Table 2. Proteins identified by mass spectrometry from isolated chromosomes of *Pyrobaculum*
556   *calidifontis*

Protein ID	Protein description	Mass (kDa)	Gene name	Gene locus
A3MWL6	Pullulanase / alpha-amylase	111.5	Pcal_1616	Pcal_1616
A3MUT3	Oligopeptide binding protein, putative	95.6	Pcal_0975	Pcal_0975
A3MTK9	V-type ATP synthase subunit I	85.8	Pcal_0549	Pcal_0549
A3MY13	AAA family ATPase, CDC48 subfamily	81.8	Pcal_2115	Pcal_2115
A3MY25	Succinate dehydrogenase subunit A	64.5	Pcal_2127	Pcal_2127
A3MV10	Thermosome subunit	60.1	Pcal_1052	Pcal_1052
A3MX21	Thermosome subunit	60.2	Pcal_1771	Pcal_1771
A3MV59	ATPase, PilT family	58	Pcal_1101	Pcal_1101
A3MV69	Elongation factor 1-alpha	48.9	tuf	Pcal_1111
A3MV64	rRNA biogenesis protein Nop56/Nop58	46.8	Pcal_1106	Pcal_1106
A3MV08	DNA primase DnaG	45	dnaG	Pcal_1050
A3MY01	S-adenosylmethionine synthase	44.4	mat	Pcal_2103
A3MX84	50S ribosomal protein L10	37.5	rpl10	Pcal_1834
A3MUU8	DNA polymerase sliding clamp	29.4	pcn	Pcal_0990
A3MY29	DNA polymerase sliding clamp	28.5	pcn	Pcal_2131
A3MWF6	Thiamine thiazole synthase	27.7	thi4	Pcal_1555
A3MX83	50S ribosomal protein L1	25.1	rpl1	Pcal_1833
A3MU88	30S ribosomal protein S5	22.1	rps5	Pcal_0779
A3MXZ7	50S ribosomal protein L13	21.7	rpl13	Pcal_2099
A3MTM7	DJ-1_PfpI domain-containing protein	20.9	Pcal_0567	Pcal_0567
A3MXA2	Uncharacterized protein	20	Pcal_1852	Pcal_1852
A3MVE1	Transcriptional regulator, AsnC family	18.7	Pcal_1183	Pcal_1183
A3MUQ5	DNA/RNA-binding protein Alba	9.9	albA	Pcal_0947

559 Table 3. Proteins identified by mass spectrometry from isolated chromosomes of *Thermococcus*
560 *kodakarensis*

Protein ID	Protein description	Mass (kDa)	Gene	Gene locus
Q5JH54	Predicted endonuclease-methyltransferase fusion protein	150.7	TK1460	TK1460
Q5JE98	CDC48/VCP homolog, AAA superfamily	88.9	TK1157	TK1157
Q5JE33	DNA-directed RNA polymerase subunit	102.9	TK1082	TK1082
Q5JH85	Putative 5-methylcytosine restriction system, GTPase subunit	79.7	TK0795	TK0795
Q5JHC8	Acylamino acid-releasing enzyme	72.1	TK0752	TK0752
Q5JH81	Type 2 DNA topoisomerase 6 subunit B	64.2	top6B	TK0799
Q5JCX3	AMP phosphorylase	53.7	deoA	TK0352
Q5JIR2	V-type ATP synthase beta chain	52.4	atpB	TK1603
Q5JFZ4	Elongation factor 1-alpha	47.5	tuf	TK0308
Q5JE34	DNA-directed RNA polymerase subunit A''	43.7	rpoA2	TK1081
Q5JGA2	Uncharacterized protein	40.8	TK0462	TK0462
Q5JDJ0	50S ribosomal protein L3	39	rpl3	TK1542
Q5JHG6	Uncharacterized protein	36.7	TK2157	TK2157
Q5JHT2	Transcription regulator, SpoVT/AbrB family, containing PhoU domain	35.3	TK2245	TK2245
Q5JD67	HTH-type transcriptional regulator TrmBL2	30.8	trmBL2	TK0471
Q5JGL5	Uncharacterized protein	29.3	TK1263	TK1263
Q5JFN1	Fibrillarin-like rRNA/tRNA 2'-O-methyltransferase	25.3	flpA	TK0183
Q5JF30	Peroxiredoxin	24.6	TK0537	TK0537
Q5JDD5	Transcription factor E	22.1	tfe	TK2024
Q5JG73	Iron-molybdenum cofactor-binding protein	21.5	TK0724	TK0724
Q5JJH0	50S ribosomal protein L15	16.5	rpl15	TK1519
Q5JH90	UPF0179 protein TK0790	15.9	TK0790	TK0790
Q5JJF8	50S ribosomal protein L14	15.2	rpl14	TK1531
Q5JF39	DNA/RNA-binding protein Alba	10.1	albA	TK0560
Q9Y8I1	Archaeal histone A	7.4	hpkA	TK1413
Q9Y8I2	Archaeal histone B	7.2	hpkB	TK2289

References

Ahmad, M.U.D., Waege, I., Hausner, W., Thomm, M., Boos, W., Diederichs, K., et al. (2015).
Structural Insights into Nonspecific Binding of DNA by TrmBL2, an Archaeal Chromatin
Protein. *J Mol Biol* 427(20)**,** 3216-3229. doi: 10.1016/j.jmb.2015.08.012.
Ali Azam, T., Iwata, A., Nishimura, A., Ueda, S., and Ishihama, A. (1999). Growth phase-
dependent variation in protein composition of the Escherichia coli nucleoid. *J Bacteriol*
181(20)**,** 6361-6370.
Amo, T., Paje, M.L.F., Inagaki, A., Ezaki, S., Atomi, H., and Imanaka, T. (2002). Pyrobaculum
calidifontis sp. nov., a novel hyperthermophilic archaeon that grows in atmospheric air.
Archaea 1(2)**,** 113-121.
Badrinarayanan, A., Le, T.B., and Laub, M.T. (2015). Bacterial chromosome organization and
segregation. *Annual review of cell and developmental biology* 31**,** 171-199.
Barutcu, A.R., Lian, J.B., Stein, J.L., Stein, G.S., and Imbalzano, A.N. (2017). The connection
between BRG1, CTCF and topoisomerases at TAD boundaries. *Nucleus* 8(2)**,** 150-155.
doi: 10.1080/19491034.2016.1276145.
Bates, S.T., Berg-Lyons, D., Caporaso, J.G., Walters, W.A., Knight, R., and Fierer, N. (2011).
Examining the global distribution of dominant archaeal populations in soil. *The ISME*
Journal 5(5)**,** 908-917. doi: 10.1038/ismej.2010.171.
Bell, S.D., Botting, C.H., Wardleworth, B.N., Jackson, S.P., and White, M.F. (2002). The
interaction of Alba, a conserved archaeal chromatin protein, with Sir2 and its regulation
by acetylation. *Science* 296(5565)**,** 148-151.
Blais Lecours, P., Marsolais, D., Cormier, Y., Berberi, M., Haché, C., Bourdages, R., et al.
(2014). Increased Prevalence of Methanosphaera stadtmanae in Inflammatory Bowel
Diseases. *PLOS ONE* 9(2)**,** e87734. doi: 10.1371/journal.pone.0087734.
Brocken, D.J.W., Tark-Dame, M., and Dame, R.T. (2018). The organization of bacterial
genomes: Towards understanding the interplay between structure and function. *Current*
Opinion in Systems Biology 8**,** 137-143. doi: https://doi.org/10.1016/j.coisb.2018.02.007.
Chang, W., Worman, H.J., and Gundersen, G.G. (2015). Accessorizing and anchoring the LINC
complex for multifunctionality. *J Cell Biol* 208(1)**,** 11-22. doi: 10.1083/jcb.201409047.
Čuboňová, L., Katano, M., Kanai, T., Atomi, H., Reeve, J.N., and Santangelo, T.J. (2012). An
archaeal histone is required for transformation of Thermococcus kodakarensis. *Journal of*
bacteriology 194(24)**,** 6864-6874.
Dame, R.T., and Tark-Dame, M. (2016). Bacterial chromatin: converging views at different
scales. *Current Opinion in Cell Biology* 40**,** 60-65. doi:
https://doi.org/10.1016/j.ceb.2016.02.015.
Davey, C.A., Sargent, D.F., Luger, K., Maeder, A.W., and Richmond, T.J. (2002). Solvent
mediated interactions in the structure of the nucleosome core particle at 1.9 Å resolution.
Journal of molecular biology 319(5)**,** 1097-1113.
De Vries, R. (2010). DNA condensation in bacteria: Interplay between macromolecular crowding
and nucleoid proteins. *Biochimie* 92(12)**,** 1715-1721.
DeLange, R.J., Green, G., and Searcy, D. (1981). A histone-like protein (HTa) from
Thermoplasma acidophilum. I. Purification and properties. *Journal of Biological*
Chemistry 256(2)**,** 900-904.
Driessen, Rosalie P.C., and Dame, Remus T. (2011). Nucleoid-associated proteins in
Crenarchaea. *Biochemical Society Transactions* 39(1)**,** 116-121. doi: 10.1042/bst0390116.
Efremov, A.K., Qu, Y., Maruyama, H., Lim, C.J., Takeyasu, K., and Yan, J. (2015).
Transcriptional Repressor TrmBL2 from Thermococcus kodakarensis Forms Filamentous
Nucleoprotein Structures and Competes with Histones for DNA Binding in a Salt- and
DNA Supercoiling-dependent Manner. *J Biol Chem* 290(25)**,** 15770-15784. doi:
10.1074/jbc.M114.626705.

Forterre, P. (2016). *Microbes from Hell.* University of Chicago Press.

French, S.L., Santangelo, T.J., Beyer, A.L., and Reeve, J.N. (2007). Transcription and translation are coupled in Archaea. *Molecular biology and evolution* 24(4)**,** 893-895.

Fyodorov, D.V., Zhou, B.R., Skoultchi, A.I., and Bai, Y. (2018). Emerging roles of linker histones in regulating chromatin structure and function. *Nat Rev Mol Cell Biol* 19(3)**,** 192-206. doi: 10.1038/nrm.2017.94.

Gibcus, Johan H., and Dekker, J. (2013). The Hierarchy of the 3D Genome. *Molecular Cell* 49(5)**,** 773-782. doi: https://doi.org/10.1016/j.molcel.2013.02.011.

Gindner, A., Hausner, W., and Thomm, M. (2014). The TrmB family: a versatile group of transcriptional regulators in Archaea. *Extremophiles* 18(5)**,** 925-936. doi: 10.1007/s00792-014-0677-2.

Grewal, S.I., and Moazed, D. (2003). Heterochromatin and epigenetic control of gene expression. *science* 301(5634)**,** 798-802.

Guo, R., Xue, H., and Huang, L. (2003). Ssh10b, a conserved thermophilic archaeal protein, binds RNA in vivo. *Mol Microbiol* 50(5)**,** 1605-1615.

Hammel, M., Amlanjyoti, D., Reyes, F.E., Chen, J.-H., Parpana, R., Tang, H.Y., et al. (2016). HU multimerization shift controls nucleoid compaction. *Science advances* 2(7), e1600650.

Henneman, B., van Emmerik, C., van Ingen, H., and Dame, R.T. (2018). Structure and function of archaeal histones. *PLoS Genet* 14(9)**,** e1007582. doi: 10.1371/journal.pgen.1007582.

Higashi, K., Tobe, T., Kanai, A., Uyar, E., Ishikawa, S., Suzuki, Y., et al. (2016). H-NS Facilitates Sequence Diversification of Horizontally Transferred DNAs during Their Integration in Host Chromosomes. *PLoS Genet* 12(1)**,** e1005796. doi: 10.1371/journal.pgen.1005796.

Hirano, T., Kobayashi, R., and Hirano, M. (1997). Condensins, chromosome condensation protein complexes containing XCAP-C, XCAP-E and a Xenopus homolog of the Drosophila Barren protein. *Cell* 89(4)**,** 511-521.

Hiraoka, Y., and Dernburg, A.F. (2009). The SUN Rises on Meiotic Chromosome Dynamics. *Developmental Cell* 17(5)**,** 598-605. doi: https://doi.org/10.1016/j.devcel.2009.10.014.

Hizume, K., Araki, S., Yoshikawa, K., and Takeyasu, K. (2007). Topoisomerase II, scaffold component, promotes chromatin compaction in vitro in a linker-histone H1-dependent manner. *Nucleic Acids Res* 35(8)**,** 2787-2799. doi: 10.1093/nar/gkm116.

Hizume, K., Yoshimura, S.H., and Takeyasu, K. (2005). Linker histone H1 per se can induce three-dimensional folding of chromatin fiber. *Biochemistry* 44(39)**,** 12978-12989. doi: 10.1021/bi050623v.

Hocher, A., Rojec, M., Swadling, J.B., Esin, A., and Warnecke, T. (2019). The DNA-binding protein HTa from Thermoplasma acidophilum is an archaeal histone analog. *Elife* 8. doi: 10.7554/eLife.52542.

Japaridze, A., Renevey, S., Sobetzko, P., Stoliar, L., Nasser, W., Dietler, G., et al. (2017). Spatial organization of DNA sequences directs the assembly of bacterial chromatin by a nucleoid-associated protein. *Journal of Biological Chemistry* 292(18)**,** 7607-7618.

Kim, J., Yoshimura, S.H., Hizume, K., Ohniwa, R.L., Ishihama, A., and Takeyasu, K. (2004). Fundamental structural units of the Escherichia coli nucleoid revealed by atomic force microscopy. *Nucleic Acids Res* 32(6)**,** 1982-1992.

Kim, M., Park, S., and Lee, S.-J. (2016). Global transcriptional regulator TrmB family members in prokaryotes. *Journal of Microbiology* 54(10)**,** 639-645.

Koskinen, K., Pausan, M.R., Perras, A.K., Beck, M., Bang, C., Mora, M., et al. (2017). First Insights into the Diverse Human Archaeome: Specific Detection of Archaea in the Gastrointestinal Tract, Lung, and Nose and on Skin. *MBio* 8(6). doi: 10.1128/mBio.00824-17.

Laurens, N., Driessen, R.P., Heller, I., Vorselen, D., Noom, M.C., Hol, F.J., et al. (2012). Alba shapes the archaeal genome using a delicate balance of bridging and stiffening the DNA. *Nature communications* 3**,** 1328.

Lemmens, L., Maklad, H.R., Bervoets, I., and Peeters, E. (2019). Transcription Regulators in Archaea: Homologies and Differences with Bacterial Regulators. *J Mol Biol*. doi: 10.1016/j.jmb.2019.05.045.

Lepp, P.W., Brinig, M.M., Ouverney, C.C., Palm, K., Armitage, G.C., and Relman, D.A. (2004). Methanogenic Archaea and human periodontal disease. *Proc Natl Acad Sci U S A* 101(16), 6176-6181. doi: 10.1073/pnas.0308766101.

Li, G., and Reinberg, D. (2011). Chromatin higher-order structures and gene regulation. *Current opinion in genetics & development* 21(2), 175-186.

Lloyd, K.G., Schreiber, L., Petersen, D.G., Kjeldsen, K.U., Lever, M.A., Steen, A.D., et al. (2013). Predominant archaea in marine sediments degrade detrital proteins. *Nature* 496(7444), 215-218. doi: 10.1038/nature12033.

López-García, P., Zivanovic, Y., Deschamps, P., and Moreira, D. (2015). Bacterial gene import and mesophilic adaptation in archaea. *Nature Reviews Microbiology* 13(7), 447-456.

Luger, K., Mader, A.W., Richmond, R.K., Sargent, D.F., and Richmond, T.J. (1997). Crystal structure of the nucleosome core particle at 2.8 A resolution. *Nature* 389(6648), 251-260.

Luijsterburg, M.S., White, M.F., van Driel, R., and Dame, R.T. (2008). The major architects of chromatin: architectural proteins in bacteria, archaea and eukaryotes. *Crit Rev Biochem Mol Biol* 43(6), 393-418.

Maruyama, H., Harwood, J.C., Moore, K.M., Paszkiewicz, K., Durley, S.C., Fukushima, H., et al. (2013). An alternative beads-on-a-string chromatin architecture in Thermococcus kodakarensis. *EMBO Rep* 14(8), 711-717. doi: 10.1038/embor.2013.94.

Maruyama, H., Kent, N.A., Nishida, H., and Oshima, T. (2019). "Functions of Archaeal Nucleoid Proteins: Archaeal Silencers are Still Missing," in *DNA Traffic in the Environment*. Springer), 29-45.

Maruyama, H., Ohniwa, R.L., Prieto, E., Hejna, J., and Takeyasu, K. (2014). "Genome-folding mechanisms in the three domains of life revealed by AFM," in *Atomic Force Microscopy in Nanobaiology*. Pan Stanford Publishing), 275-310.

Maruyama, H., Shin, M., Oda, T., Matsumi, R., Ohniwa, R.L., Itoh, T., et al. (2011). Histone and TK0471/TrmBL2 form a novel heterogeneous genome architecture in the hyperthermophilic archaeon Thermococcus kodakarensis. *Mol Biol Cell* 22(3), 386-398. doi: 10.1091/mbc.E10-08-0668.

Mattiroli, F., Bhattacharyya, S., Dyer, P.N., White, A.E., Sandman, K., Burkhart, B.W., et al. (2017). Structure of histone-based chromatin in Archaea. *Science* 357(6351), 609-612. doi: 10.1126/science.aaj1849.

Monterroso, B., Zorrilla, S., Sobrinos-Sanguino, M., Robles-Ramos, M.Á., Alfonso, C., Söderström, B., et al. (2019). The bacterial DNA binding protein MatP involved in linking the nucleoid terminal domain to the divisome at midcell interacts with lipid membranes. *mBio* 10(3), e00376-00319.

Morikawa, K., Ushijima, Y., Ohniwa, R.L., Miyakoshi, M., and Takeyasu, K. (2019). What Happens in the Staphylococcal Nucleoid under Oxidative Stress? *Microorganisms* 7(12), 631.

Navarre, W.W., Porwollik, S., Wang, Y., McClelland, M., Rosen, H., Libby, S.J., et al. (2006). Selective silencing of foreign DNA with low GC content by the H-NS protein in Salmonella. *Science* 313(5784), 236-238.

Nishida, H., and Oshima, T. (2017). Archaeal histone distribution is associated with archaeal genome base composition. *J Gen Appl Microbiol* 63(1), 28-35. doi: 10.2323/jgam.2016.07.003.

Niu, Y., Xia, Y., Wang, S., Li, J., Niu, C., Li, X., et al. (2013). A prototypic lysine methyltransferase 4 from archaea with degenerate sequence specificity methylates chromatin proteins Sul7d and Cren7 in different patterns. *Journal of Biological Chemistry* 288(19), 13728-13740.

Nkamga, V.D., Henrissat, B., and Drancourt, M. (2017). Archaea: Essential inhabitants of the
human digestive microbiota. *Human Microbiome Journal* 3**,** 1-8. doi:
https://doi.org/10.1016/j.humic.2016.11.005.
Ohniwa, R.L., Maruyama, H., Morikawa, K., and Takeyasu, K. (2018). "Atomic Force
Microscopy Imaging and Analysis of Prokaryotic Genome Organization," in *Bacterial*
Chromatin. Humana Press, New York, NY), 147-160.
Ohniwa, R.L., Morikawa, K., Takeshita, S.L., Kim, J., Ohta, T., Wada, C., et al. (2007).
Transcription-coupled nucleoid architecture in bacteria. *Genes Cells* 12(10)**,** 1141-1152.
Ono, T., Losada, A., Hirano, M., Myers, M.P., Neuwald, A.F., and Hirano, T. (2003). Differential
contributions of condensin I and condensin II to mitotic chromosome architecture in
vertebrate cells. *Cell* 115(1)**,** 109-121.
Oshima, T., Ishikawa, S., Kurokawa, K., Aiba, H., and Ogasawara, N. (2006). Escherichia coli
histone-like protein H-NS preferentially binds to horizontally acquired DNA in
association with RNA polymerase. *DNA Res* 13(4)**,** 141-153.
Pausan, M.R., Csorba, C., Singer, G., Till, H., Schöpf, V., Santigli, E., et al. (2019). Exploring the
Archaeome: Detection of Archaeal Signatures in the Human Body. *Frontiers in*
Microbiology 10(2796). doi: 10.3389/fmicb.2019.02796.
Peeters, E., Driessen, R.P., Werner, F., and Dame, R.T. (2015). The interplay between nucleoid
organization and transcription in archaeal genomes. *Nat Rev Microbiol* 13(6)**,** 333-341.
doi: 10.1038/nrmicro3467.
Pelve, E.A., Lindås, A.C., Knöppel, A., Mira, A., and Bernander, R. (2012). Four chromosome
replication origins in the archaeon Pyrobaculum calidifontis. *Molecular microbiology*
85(5)**,** 986-995.
Poplawski, A., and Bernander, R. (1997). Nucleoid structure and distribution in thermophilic
Archaea. *J Bacteriol* 179(24)**,** 7625-7630. doi: 10.1128/jb.179.24.7625-7630.1997.
Prieto, Eloise I., and Maeshima, K. (2019). Dynamic chromatin organization in the cell. *Essays in*
Biochemistry 63(1)**,** 133-145. doi: 10.1042/ebc20180054.
Probst, A.J., Auerbach, A.K., and Moissl-Eichinger, C. (2013). Archaea on human skin. *PLoS*
One 8(6)**,** e65388. doi: 10.1371/journal.pone.0065388.
Rojec, M., Hocher, A., Stevens, K.M., Merkenschlager, M., and Warnecke, T. (2019).
Chromatinization of Escherichia coli with archaeal histones. *Elife* 8. doi:
10.7554/eLife.49038.
Ruepp, A., Graml, W., Santos-Martinez, M.-L., Koretke, K.K., Volker, C., Mewes, H.W., et al.
(2000). The genome sequence of the thermoacidophilic scavenger Thermoplasma
acidophilum. *Nature* 407(6803)**,** 508-513. doi: 10.1038/35035069.
Sanders, T.J., Marshall, C.J., and Santangelo, T.J. (2019). The Role of Archaeal Chromatin in
Transcription. *J Mol Biol* 431(20)**,** 4103-4115. doi: 10.1016/j.jmb.2019.05.006.
Sandman, K., and Reeve, J.N. (2005). Archaeal chromatin proteins: different structures but
common function? *Curr Opin Microbiol* 8(6)**,** 656-661.
Sarkar, T., Petrov, A.S., Vitko, J.R., Santai, C.T., Harvey, S.C., Mukerji, I., et al. (2009).
Integration host factor (IHF) dictates the structure of polyamine-DNA condensates:
implications for the role of IHF in the compaction of bacterial chromatin. *Biochemistry*
48(4)**,** 667-675.
Sato, A., Isaac, B., Phillips, C.M., Rillo, R., Carlton, P.M., Wynne, D.J., et al. (2009).
Cytoskeletal forces span the nuclear envelope to coordinate meiotic chromosome pairing
and synapsis. *Cell* 139(5)**,** 907-919. doi: 10.1016/j.cell.2009.10.039.
Searcy, D.G. (1975). Histone-like protein in the prokaryote Thermoplasma acidophilum.
Biochimica et Biophysica Acta (BBA)-Nucleic Acids and Protein Synthesis 395(4)**,** 535-
547.
Skoko, D., Yoo, D., Bai, H., Schnurr, B., Yan, J., McLeod, S.M., et al. (2006). Mechanism of
chromosome compaction and looping by the Escherichia coli nucleoid protein Fis. *J Mol*
Biol 364(4)**,** 777-798.

Takemata, N., Samson, R.Y., and Bell, S.D. (2019). Physical and Functional
Compartmentalization of Archaeal Chromosomes. *Cell* 179(1)**,** 165-179 e118. doi:
10.1016/j.cell.2019.08.036.
Thanbichler, M., Wang, S.C., and Shapiro, L. (2005). The bacterial nucleoid: A highly organized
and dynamic structure. *Journal of Cellular Biochemistry* 96(3)**,** 506-521. doi:
10.1002/jcb.20519.
The UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*
Res 47(D1)**,** D506-D515. doi: 10.1093/nar/gky1049.
Thomas, J.O. (1984). The higher order structure of chromatin and histone H1. *J Cell Sci Suppl*
1(Supplement 1)**,** 1-20. doi: 10.1242/jcs.1984.supplement_1.1.
Uhlmann, F. (2016). SMC complexes: from DNA to chromosomes. *Nature Reviews Molecular*
Cell Biology 17(7)**,** 399-412. doi: 10.1038/nrm.2016.30.
Verma, S., Qian, Z., and Adhya, S. (2019). Architecture of the Escherichia coli nucleoid. *PLoS*
genetics 15(12)**,** e1008456.
Wagner, A., Whitaker, R.J., Krause, D.J., Heilers, J.-H., Van Wolferen, M., Van Der Does, C., et
al. (2017). Mechanisms of gene flow in archaea. *Nature Reviews Microbiology* 15(8)**,** 492.
Wang, X., Llopis, P.M., and Rudner, D.Z. (2013). Organization and segregation of bacterial
chromosomes. *Nature Reviews Genetics* 14(3)**,** 191.
Weidenbach, K., Gloer, J., Ehlers, C., Sandman, K., Reeve, J.N., and Schmitz, R.A. (2008).
Deletion of the archaeal histone in Methanosarcina mazei Go1 results in reduced growth
and genomic transcription. *Mol Microbiol* 67(3)**,** 662-671.
Woodcock, C.L., Skoultchi, A.I., and Fan, Y. (2006). Role of linker histone in chromatin
structure and function: H1 stoichiometry and nucleosome repeat length. *Chromosome*
Research 14(1)**,** 17-25.

Figure 1. Genomic material released from *Thermoplasma acidophilum* and *Pyrobaculum calidifontis* cells. DAPI staining was performed after on-substrate lysis of the cells during (A) log and (B) stationary phases. (A) Chromosomes were well-spread in the log-phase for both T. acidophilum (upper panel) and P. calidifontis (lower panel). (B) The stationary phase chromosome was more compact than the log-phase chromosome (arrow). Scale bars: 2 μm.

Figure 2. Structural details of genomic materials from *Thermoplasma acidophilum* and *Pyrobaculum calidifontis* after on-substrate lysis visualized with atomic force microscopy (AFM). (A-B) Extensively released chromosome fibers of log-phase cells. (A) AFM of the released genomic materials. (B) Histograms indicate the diameters of the released structures. (C-D) Compact chromosome of stationary phase cells. (C) AFM of the lysed cells. Boxed areas in the upper panels were investigated in detail in the lower panels. Globular structures were observed (arrows) (D) Histograms indicate the diameters of the globular structures. Scale bars: 200 nm.

Figure 3. Digestion pattern of archaeal chromosomes with micrococcal nuclease (MNase). Purified chromosomes of (A) *Thermococcus kodakarensis*, (B) *Thermoplasma acidophilum*, (C) *Pyrobaculum calidifontis*, and (D) *Sulfolobus solfataricus* were digested with increasing concentrations of MNase and separated with 2.5% agarose gels in 1X TBE. The accumulation of DNA of particular sizes (arrows) was observed with *T. kodakarensis* and *T. acidophilum* but not with *P. calidifontis* and *S. solfataricus* chromosomes. MNase concentration was 0.3, 1, 3, 10 U MNase in 100 μl reaction (A) or 0, 0.3, 1, 3, 10 and 30 U MNase in 100 μl reaction (B-D).

Figure 4. In vitro reconstitution of chromosome structures with recombinant Alba and HTa. (A-C) Reconstitution on a linear 3-kb plasmid using recombinant Alba from *Thermoplasma acidophilum* and *Pyrobaculum calidifontis* at different Alba to DNA weight ratios. (A) Atomic force microscopy (AFM) showing various structures formed with Alba and DNA. Alba:DNA weight ratios are indicated. (B) Histograms show the diameters of the fibrous structures formed, as well as (C) Respective contour lengths of DNA at a 15:1 Alba:DNA ratio. The theoretical length of a linear 3-kb DNA in B-form is ~1000 nm (Maruyama et al., 2011). (D-G) Reconstitution with histidine-tagged HTa from *T. acidophilum* at varying protein to DNA ratio. (D) AFM showing beaded (arrows) and filamentous (triangles) structures. (E) Diameter of the beads formed at a relatively lower HTa concentration. (F) Width of the filaments formed at relatively higher HTa concentration. (G) Histogram shows the contour DNA lengths of the structures formed at a 15:1 HTa:DNA ratio. Scale bars: 100 nm.

Figure 5. In vitro reconstitution using a combination of archaeal proteins. Chromosomal structures were reconstituted on a linear 3-kb plasmid using recombinant proteins at varying Alba concentrations. (A) Atomic force microscopy (AFM) showing reconstitution using histone and Alba from *Thermococcus kodakarensis*. Histograms below the AFM images show the corresponding contour DNA length with each condition. (B) AFM showing reconstitution using HTa and Alba from *Thermoplasma acidophilum*. Scale bars: 100 nm.

Figure 6. Model of archaeal chromosome folding. (A) In Euryarchaeota, histone or HTa wraps genomic DNA, whereas there is no DNA wrapper in Crenarchaeota. Alba can bridge distant genomic DNA. Chromosome compaction occurs toward the stationary phase, where 30~40-nm globular structures are formed. (B) Schematic phylogenetic tree of selected Archaea (Sandman and Reeve, 2005) together with the distributions of histone and HTa. Species in Euryarchaeota utilize histone or HTa (*Thermoplasmatales*) to wrap DNA at the primary level. Members of Nanoarchaeota and Thaumarchaeota also encode histone that might wrap DNA. Species in Crenarchaeota neither encode histone nor wrap DNA. The branch lengths do not correspond to evolutionary distances.