Human and macaque pairs employ different coordination strategies in a transparent decision game

Sebastian Möller[1,2,3], Anton M. Unakafov[1,2,3,4,5], Julia Fischer[3,6,7], Alexander Gail[1,2,3,8,*], Stefan Treue[1,2,3,8,*], Igor Kagan[1,3,✉,*]

[1]Cognitive Neurosciences Laboratory, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077, Goettingen, Germany

[2]Georg-Elias-Müller-Institute of Psychology, University of Goettingen, Gosslerstrasse 14, 37073, Germany

[3]Leibniz ScienceCampus Primate Cognition, Kellnerweg 4, 37077, Goettingen, Germany

[4]Max Planck Institute for Dynamics and Self-Organization, Am Fassberg 17, 37077, Goettingen, Germany

[5]Campus Institute for Dynamics of Biological Networks, Hermann-Rein-Straße 3 37075, Goettingen, Germany

[6]Cognitive Ethology Laboratory, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077, Goettingen, Germany

[7]Department of Primate Cognition, Johann-Friedrich-Blumenbach Institute for Zoology and Anthropology, University of Goettingen, Germany

[8]Bernstein Center for Computational Neuroscience, Am Fassberg 17, 37077, Goettingen, Germany

✉ Corresponding author: ikagan@dpz.eu

* Shared senior authorship

Short title: Decision coordination in humans and macaques

1 Abstract

Many real-world decisions in social contexts are made while observing a partner's actions. To
study dynamic interactions during such decisions, we developed a setup where two agents seated
face-to-face engage in game-theoretical tasks on a shared transparent touchscreen display
("transparent games"). We compared human and macaque pairs in a transparent version of the
coordination game "Bach-or-Stravinsky", which entails a conflict about which of two individually-
preferred opposing options to choose to achieve coordination. Most human pairs developed
coordinated behavior and adopted dynamic turn-taking to equalize the payoffs. All macaque pairs
converged on simpler, static coordination. Remarkably, two animals learned to coordinate
dynamically after training with a human confederate. This pair selected the faster agent's
preferred option, exhibiting turn-taking behavior that was captured by modeling the visibility of the
partner's action before one's own movement. Such competitive turn-taking was unlike the
prosocial turn-taking in humans, who equally often initiated switches to and from their preferred
option. Thus, the dynamic coordination is not restricted to humans, but can occur on the
background of different social attitudes and cognitive capacities in rhesus monkeys. Overall, our
results illustrate how action visibility promotes emergence and maintenance of coordination when
agents can observe and time their mutual actions.

Introduction

The majority of primate species lives in complex social groups, in which interactions range from

intense competition to cooperation (Byrne and Bates, 2010; Chang et al., 2013; Jaeggi et al.,

2010; Joly et al., 2017; Platt et al., 2016). To adjust behavior optimally, individuals need to assess

their own actions and goals, and those of other group members, while taking into account the

history of interactions with and between these individuals (Frith and Frith, 2010; Platt et al., 2016;

Waal and Waal, 2007). Coordination is essential for maintaining cohesion between group

members, avoiding conflicts, and achieving individual and joint action goals. Here we focus on

understanding how such coordination can be achieved and maintained in a dyadic "transparent"

setting in which agents can observe each other's ongoing actions and social cues (Unakafov et

al., 2020, 2019), like in many natural situations (Conty et al., 2012; Fruteau et al., 2013; Rizzolatti

et al., 2001; Silk, 2007).

Game theory, developed to study strategic interactions in rational decision-makers (von Neumann

and Morgenstern, 1944), offers a powerful framework to investigate dyadic social interactions

(Rilling and Sanfey, 2011; Sanfey, 2007; Tremblay et al., 2017) and the evolution of coordination

(Brosnan, 2018; Santos and Rosati, 2015; Smith, 1979). In 2×2 games (Rapoport et al., 1976;

Rapoport, 1973) each of two agents chooses one of two actions and the outcome depends on

the combination of their choices. Non-zero-sum 2×2 games that have both cooperative and

competitive elements with aligned and opposing interests are particularly interesting for

understanding many realistic scenarios. Here, the agents need to coordinate their choices to

maximize individual and/or joint reward (Rapoport, 1967). For example, in the Stag Hunt (or

Assurance) game, agents choose between maximizing the individual as well as joint reward and

minimizing the individual risk (Skyrms, 2003). Humans *(Homo sapiens)*, chimpanzees *(Pan*

troglodytes), rhesus macaques *(Macaca mulatta)* and capuchin monkeys *(Cebus apella)*

coordinate at above chance level on mutually beneficial high-reward / high-risk choices in the

iterated Stag Hunt (Brosnan et al., 2012, 2011; Bullinger et al., 2011; Duguid et al., 2014). Similar

to Stag Hunt, in Prisoner's Dilemma (PD) two agents maximize their *joint* reward if they both

choose to cooperate. Yet, an individual agent obtains the highest reward when defecting while

the partner cooperates (and receives the lowest reward), creating a dilemma between the selfish

temptation to defect and a riskier cooperation that requires trust. Humans often show a bias

towards cooperative behavior (Camerer, 2003; Rilling et al., 2002; Wedekind and Milinski, 1996).

Remarkably, it was demonstrated that macaques, while mostly defecting, chose mutual

cooperation significantly more and mutual defection significantly less than expected by chance,

especially after preceding cooperation, in an iterated PD game (Haroush and Williams, 2015).

This finding suggests that macaques can at least partially overcome the selfish motives and

reciprocate cooperation.

Another type of conflict that contrasts selfish and prosocial preferences is implemented in the

Conflict game (also known as Hawk-Dove, or game of Chicken) (Brosnan et al., 2017; Smith,

1979). These anti-coordination games model a competition over a shared resource that can be

monopolized by one agent, while an actual clash is highly detrimental to both. Humans, capuchins

and macaques often converge on anti-coordination (Brosnan et al., 2017), but importantly, only

humans alternate between the two individually-optimal anti-coordinated choices (Brosnan et al.,

2017; Grueneisen and Tomasello, 2017; Helbing et al., 2005). Such alternating behavior results

in the maximal and equal payoff for both agents and is called "cooperative turn-taking" (Bornstein

et al., 1997; Colman and Browning, 2009; Noë, 2006).

In the above games, reward-optimizing rational agents should coordinate on one option (Stag

Hunt and Prisoner's Dilemma) or anti-coordinate on opposite options (Conflict game). A third, less

studied type of games that emphasizes the coordination on *either of two options* is known as

Battle of the Sexes, or Bach-or-Stravinsky game (BoS) (Kilgour and Fraser, 1988). Each agent

has an individually preferred option, but coordinating on either one of these options adds the same

bonus to both agents. This renders any coordinated choice better than no coordination for both

agents, but one coordinated choice is better for the first agent and the other better for the second

agent. The rational choice is to coordinate, but unlike the conceptually simpler optimal

convergence on one option in Stag Hunt and PD, BoS includes an inherent conflict about who

profits the most. This combination of cooperation and conflict offers an interesting opportunity for

studying social interactions in nonhuman primates. As in the Conflict game, games of the BoS

type revealed that humans often take turns, switching between the two coordinated options to

ensure fairness (Ioannou and Romero, 2014; Sonsino and Sirota, 2003). Interestingly, in a rope-

pulling task that requires cooperation but rewards only one agent at a time, trial-by-trial turn-taking

frequently took place in 5-year-old children, but neither in 3-year-old children nor in chimpanzees

(Melis et al., 2016). This result begs the question whether turn-taking requires special social or

cognitive abilities that are unique to humans. We therefore compared the behavior of (adult)

humans and rhesus macaques in the BoS economic game.

Traditionally, economic games are played either *"simultaneously"* (neither agent knows the choice

of the other before making its own decision) or *sequentially* in a predetermined order. Yet, real

dyadic interactions often play out with the partner's actions in direct sight (Dugatkin et al., 1992;
van Doorn et al., 2014). Thus, the timing of one's own and the other's actions becomes part of
the strategy space (McDonald et al., 2019). Such a "transparent" continuous time setting can
change choice strategies as compared to the classic simultaneous and sequential settings (Noë,
2006).

To account for changes in strategies during action visibility compared to "opaque" (simultaneous)
choices, we recently developed the concept of "transparent games" that extends classical
evolutionary game-theoretic analysis to real-time interactions in which the visibility of partners'
actions depends on their relative action times (Unakafov et al., 2020, 2019). Here, we use the
theoretical insights from that work and analyze short- and long-term dynamics of choices, mutual
information and action times in humans and rhesus monkeys playing a transparent version of an
iterated movement-based BoS game. To this end, we designed a novel dyadic interaction platform
where two human or monkey agents face each other and act on the same visual objects on a
vertical touch-sensitive transparent display between them. This configuration provides action
visibility, i.e. it allows agents to monitor and react to each other's actions in real-time, emulating
naturalistic interactions under well-controlled laboratory conditions.

To facilitate inter-species comparability, we paired human or macaque agents without explicit task
instructions. We expected that the transparency and instantaneous coaction (van Doorn et al.,
2014) would facilitate efficient coordination in both species. Based on their cognitive abilities
(Proto et al., 2019), propensity for fairness (Peysakhovich et al., 2014) and perspective taking
(Devaine et al., 2017; Herrmann et al., 2007), we predicted humans to engage in pro-social turn-
taking, where the reward maximization is alternated between partners. We predicted that
macaques would also utilize the information about the partner's actions, but would be less likely
to grant an advantage to the other player. Instead, we expected that the monkeys would
coordinate to increase individual (and joint) rewards, but without turn-taking to balance rewards
between players.

Results

Two human or macaque agents sat face-to-face and performed a task on a dyadic interaction
platform using a shared transparent vertical workspace (**Figure 1A**). Agents could simultaneously
see the stimuli and each other's actions. In our variant of the BoS paradigm, each agent could
choose one of two simultaneously presented color targets (red and blue, **Figure 1B**). The left/right

position of each color was randomized across trials (50% left, 50% right). The rewards followed

the payoff matrix shown in **Figure 1C** and depended on the choices of both agents. Prior to the

first dyadic session, each human or macaque agent in a pair was individually trained to associate

one of the two color targets to a larger reward, so that the two agents preferred different colors.

In the dyadic trials, selecting the same target resulted in an additional reward on top of the

individually trained values - such that choosing the same target was always better than choosing

different targets. But in such coordinated trials, the agent whose individually preferred ("own")

color was selected received a larger reward. This paradigm probes the ability to coordinate in

order to obtain higher rewards, and the capacity to deal with the unfairness introduced in each

coordinated trial.

Human pairs mainly converge on fair coordination

We recruited 38 naïve human subjects to assess performance in the transparent BoS paradigm

(**Table S1**). **Figure 1E,F** illustrates an exemplary human pair that developed such alternating turn-

taking behavior, showing the *fraction of choosing own* color (FCO) - the likelihood for each agent

to select the own preferred color, and the *fraction of choosing left* side (FCL) - the likelihood for

each agent to select the target on the left side. These measures provide direct insight into each

agent's target and side biases. Here the FCO curves show that this pair mostly jointly selected

A's or B's color. After an initial period of long blocks of trials, the pair switched to shorter blocks.

The FCL curves show that the agents did not have a side bias, as was also the case for the

majority of pairs (**Figure S1B**).

Human dyads showed mainly one of three different choice patterns. **Figure 2A** plots the larger of

the two average fraction of choosing own values in a pair versus the smaller of the two, for all 19

pairs. The FCOs of 53% (10 of 19 pairs, teal underlay) had balanced values close to 0.5 for both

agents, indicating alternating between colors, using either trial-by-trial or block-wise turn-taking.

The remaining 47% (9 of 19 pairs) had FCO of at least one agent close to one or to zero (both

cases indicate a fixed color selection). Five pairs coordinated on a fixed color favoring one agent

(blue underlay). Of the four pairs that did not coordinate successfully, in two pairs both agents

selected only the own preferred color (red underlay). In remaining two pairs one agent selected

the non-preferred color while the other switched between the two colors or selected the one side.

Most pairs achieved an average joint reward above the chance level (2.5 units), and about half

the pairs close to the optimum for the dyad (3.5) (**Figure 2C**).

Figure 1. Experimental setup. (A) Photos of the dyadic interaction platform showing humans and macaques reaching to targets in a shared transparent workspace. **(B)** A sketch and **(C)** the payoff matrix describing reward outcomes for each player for four possible combinations of individual choices (numbers show the monetary/liquid reward units for agent A in red and agent B in blue, 1.5 euro cents per "unit" for humans and ~0.1 ml of liquid for monkeys). The coordinated selection of the same target (green bold diagonal) resulted in a "bonus" of 2 reward units. **(D)** Schematic top-view of two macaque agents in different stages of the task, and the timing of one trial. **(E,F)** Session timecourse of an example human pair (pair 2). **(E)** The fraction of choosing own (FCO) for both agents (A: red; B: blue) as running average over 8 trials. In coordinated trials one agent always selects the own target (FCO = 1), while the other selects the non-preferred target (FCO = 0). The dashed lines show the session average FCO for each agent. The color bar below shows for each trial the selected combination (red: A and B selected A's preferred target; blue: A and B selected B's preferred target; magenta: both agents selected their individually preferred target; green: both agents selected their non-preferred target). Here both human agents coordinated by selecting the same target in blocks. **(F)** The fraction of choosing left (FCL, from A's viewpoint) as running average over 8 trials. The dashed lines show the session average FCL for each agent. Here side selection fluctuated around random level 0.5 for both players.

To characterize the dynamics of coordinated behavior, we derived a measure of the *dynamic coordination reward* (DCR). The DCR describes the average reward a pair earned *above* (or *below*) the reward expected for independent random choices, given the observed color choice frequencies per agent. Zero DCR indicates coordination by chance, values significantly different from zero indicate above chance levels of dynamic coordination. The DCR values (**Figure 2D**) and the fairly balanced average rewards (**Figure S2B**) suggest that 10 of 19 human pairs converged on a form of turn-taking, or reciprocal coordination. These are the same 10 pairs that showed roughly balanced FCO values around 0.5 (**Figure 2A**).

Figure 2. Performance of human and macaque pairs. (A) Human pairs. The larger of the two average FCO values in each pair versus the smaller of the two. The diagonal line indicates equal frequency of own choices for both agents. Humans converged mainly on three different strategies: balanced turn taking (teal underlay), static selection of the same color (blue underlay), or static selection of different colors (red underlay), two out of 19 pairs did not converge on any of these strategies. **(B)** Macaque pairs: same as (A) for a late session for each pair. Macaque pairs converged on jointly selecting either one of the two color targets (blue underlay) or the same side (teal underlay). Note that the clusters within the teal underlay in (A) and (B) have different underlying choice behavior: turn-taking for humans and fixed side choices for monkeys which results in automatic alternation of the color targets; to disambiguate these two strategies see **Figure 3**. **(C)** Violin plots of the average joint reward for each pair; the colored dots show the individual pairs; the white dot shows the median of each group. A reward of 2 indicates uncoordinated choice of the respective preferred target, a reward of 3.5 indicates perfect coordination, and a reward of 2.5 indicates random independent choices. All but 5 human pairs and all monkey pairs achieved an average joint reward > 3. Note that the average joint reward cannot disambiguate between balanced and unbalanced reward for each agent, see **Figure S2B, G** for individual rewards. **(D)** The dynamic coordination reward (DCR) for human and monkey pairs. Ten human pairs showed a significant DCR with amplitude >0.2 (teal dots, corresponding to the teal underlay in A, see also **Figure S2C**) while no monkey pair showed high DCR (only one pair showed a small (<0.2) but significant DCR, see also **Figure S2H**).

Macaque pairs converge to simpler coordination strategies

Like humans, macaque monkeys were individually trained to internalize the reward values of the two color targets. All six macaques developed a strong preference for the target associated with the large reward (range 78-100%, mean 95% large reward color selection in the last solo session), before being paired with a conspecific with the opposite target color preference. We then collected the data from 9 macaque pairs for multiple sessions to assess on which strategy they converged and how their behavior evolved over time. In the late dyadic sessions, all pairs converged on coordination, in one of two distinct ways: either by converging on the same color target (**Figure 2B**, blue underlay, 5 pairs) or on the same side (teal underlay, 4 pairs). This side coordination can be seen in high FCL values (**Figure S1F**, see **Figure S3** for example of developing such coordination in one pair). All pairs achieved above chance joint reward (**Figure 2C**).

Across all pairs, the average joint reward increased from early to late sessions (median early 2.75 versus late 3.29; p < 0.004, two-sided exact Wilcoxon signed rank test), and in 8 of 9 pairs, the proportion of coordinated trials increased significantly (p < 0.05 Bonferroni corrected, Fisher's exact test on the number of coordinated versus uncoordinated trials in the early and the late session, calculated separately for each pair; **Figure S1E**).

Comparison of behavior between species

To better understand what drives high dynamic coordination in humans (**Figure 2D**), we calculated the *mutual information* (MI) between the sequences of target color choice (MI target) and the side choice (MI side) of both agents. These measures indicate how well one agent's choice of color or side can be predicted from the other's choice. Comparing the DCR and the MI measures revealed that high DCR values coincided with non-zero values of *both* MI side and MI color, indicating that dynamic coordination corresponds to both correlated side and correlated color choices between the agents (**Figure S2D,E**).

In contrast to humans, monkeys showed either high values of MI color or MI side indicating that the resulting coordination was not achieved by dynamic turn-taking (**Figure 2D**). Instead, the four pairs in the middle of **Figure 2B** (teal underlay) that had roughly equal FCO and appear to be similar to the human turn-takers achieved the coordination by consistently selecting the same side regardless of the target color, which was randomized by the task (Materials and Methods).

To directly compare the strategies of human and macaque pairs, we plotted the mutual information values for side and target color choices of each pair (**Figure 3**). The distance of each

pair's location to the origin denotes the strength of (anti)coordination. Locations close to the x-
axis denote static side strategies; locations close to the y-axis denote static target color strategies.
Ten of 19 human pairs were close to the main diagonal, corresponding to similar values of MI
side and MI color. These locations denote trial-by-trial or block-wise turn-taking – a signature of
dynamic coordination. In contrast to humans, nine macaque pairs showed simple static side-
based (4 pairs) or single target color-based (5 pairs) coordination, with different levels of
coordination strength, but did not employ dynamic coordination.

Figure 3. Comparison of coordination strategies in humans and macaques. Each symbol shows the mutual
information between side choices (MI side) and target color choices (MI color) calculated over the last 200 trials of a
session, for human and macaque pairs. Since mutual information for a non-varying sequence of choices is zero,
sessions in which a pair statically converged on a fixed color have MI color around zero and a non-zero MI side, and
sessions in which a pair statically converged on a fixed side have a non-zero MI color and around zero MI side.
Pairs/sessions near the y-axis hence indicate static selection of one of the two target colors (i.e., either coordinating on
A's or B's preferred option, or only selecting the own color; dark gray underlays), while pairs/sessions near the x-axis
indicate static coordination on a side. Pairs/sessions with balanced side and color choices fall onto the main diagonal,
indicating dynamic coordination or turn-taking (light gray underlay). The distance between each symbol and the (0,0)
origin represents the (anti)coordination strength. Ten human pairs showed dynamic coordination, using trial-by-trial or
block-wise turn-taking. Eight human pairs converged on color-based strategies: five pairs largely selected one color,
i.e., one of the two coordinated options (5, 14, 16, 10, 18), two pairs did not coordinate (4, 19), and in one pair (11) one
agent selected other's color but the other randomly switched between the colors. In one pair, both MI side and MI color
did not significantly differ from zero (p > 0.01; pair 15, in this pair one agent mostly selected one side and the other -
one color). In late sessions, macaque pairs either converged on selecting the same color target (pairs FC, LE, TC, TF,
CL) or converged on the same side (TE, CE, MC, MF).

Macaques paired with a human confederate learn to follow

The absence of trial-by-trial coordination in macaque pairs suggests that they did not monitor their partners in order to take their choices into account on a trial-by-trial basis. This might be a consequence of the animals' training history (starting with extended individual solo training to associate one color with higher reward) or a general disinclination to consider the partner's choice when making immediate decisions, at least when a simpler strategy is available. To test the latter, we trained two monkeys in the dyadic condition with a human confederate who followed a strict pattern of alternating between the two colors in blocks of approximately 20 trials ("confederate training"). If the rhesus monkeys were insensitive to partner's choices in our dyadic task, the pattern of confederate's choices should not affect monkey's behavior.

Both monkeys started with a strong bias for selecting their own preferred color but over time changed to reliable coordination (cf. **Figure 4A** with **4B,** and **S4A** with **S4B** for another monkey). Their action times (**Figure 4** and **S4**, right columns) were initially faster than those of the confederate, and did not vary strongly depending on the confederate's actions. In later sessions the animals reacted considerably slower in those blocks in which the confederate was selecting the animals' non-preferred color, as if waiting for the confederate to commit to an action. We therefore labelled this as "following" behavior.

The aggregate measures of choice behavior and coordination development over multiple confederate-paired sessions showed similarities in monkeys' learning (**Figure S5**). Both animals reached close to 0.8 DCR units, similar to the values reached by humans employing turn-taking (cf. **Figure 2C**). These results indicate that macaques can take information from their partner into consideration when making decisions in the BoS paradigm. Action time dynamics suggested that the macaques started to actively monitor the confederate's actions and only selected their non-preferred color if the partner started to reach there first.

Due to the repeating block strategy employed by the human confederate it is not immediately clear whether the monkeys made use of the action visibility and based their decision on the action observed within a trial, or rather on the reward history. We therefore performed a control experiment in which we placed an opaque barrier on the confederate's side of the display for roughly the middle third of a session. The barrier blocked the monkey's view of the confederate hand, while keeping the face visible. In both monkeys, the "following" behavior ceased and they reverted to selecting their own preferred colors irrespective of the confederate's choices, who – by design – continued the block strategy (**Figure 4C**, **S4C**). In line with this observation, there

was a significant difference in monkey's action times when selecting monkey's preferred (blue)

and the non-preferred (red) color in the transparent condition, but no difference in the opaque

condition. In other words, monkeys failed to coordinate when they could not see the confederate's

actions. Note that reverting to mostly own choices in combination with a block-switching

confederate resulted in a reward alternating between 2 and 4 units, clearly different from the

constant reward of 2 units in solo trials. Hence, the experienced reward schedule and the visible

presence of a confederate render the opaque condition distinct from the solo trials. These results

imply that the immediate visual information about partner's actions and not merely monitoring

reward history drove and maintained the macaques' "following" behavior in transparent settings.

Figure 4. Macaques can be trained to take turns when paired with a human confederate. A human partner
alternated between the two colors in pre-defined blocks. **(A)** In an early confederate training session, the monkey F
(agent B) mostly selected his own target, independent of the human's choices. The lower sub-panel shows the running
average (8 trials) of the average joint reward; the cyclic modulation shows that the macaque did not coordinate with the
confederate. The action time (AT) difference histograms on the right (bin size 50 ms) show that in all four choice
combinations the macaque acted faster than the confederate. **(B)** A late confederate training session. After several
sessions (**Figure S5 A-F**) the macaque had learned to coordinate (note anti-correlation of the two FCO curves and the
average joint reward approaching maximum 3.5 value). The AT difference histograms show that in blocks when the
pair selected the macaque's non-preferred target (red curve), the monkey acted slower than the confederate, waiting
to see the human's choice before committing to its own choice (t-test: coordination on red (mean: -259 ms, SD: 113,
N: 193) vs. blue (mean: -7 ms, SD: 153, N: 217), t(395.5): -19.1, p<0.001). **(C)** Macaque turn-taking with human

confederate depended on visibility of the confederate's actions. The gray background denotes the period in which an
opaque barrier was placed over the region of the transparent display encompassing all touch targets; hence, the agents
could not see each other's reach movements but could still see the face. The macaque switched from turn-taking to
mainly selecting own preferred target during the opaque trials. Upward histograms show the transparent condition (solid
curves) and downward histograms show the opaque condition (dashed curves). In the transparent condition the
behavior was similar to (B): the monkey waited for the human to act first in blocks of the monkey's non-preferred target
(t-test: coordination on red (mean -248 ms, SD 101, N 123) vs. blue (mean -39 ms, SD 124, N 166); t(284.5) -15.9, p <
0.001). In the opaque condition the behavior was similar to (A): the monkey mostly selected his preferred target and
there was no AT difference dependence on human's choices (note the overlap between dashed blue and magenta
curves). See **Figure S4** and **S5G-L** for monkey C.

Confederate-trained macaque pair shows dynamic coordination

After two macaques had learned to monitor and follow a partner's actions, we tested how these

animals would perform when paired with each other again. At the end of the initial naïve sessions

of this pair, and before the training with a human confederate, monkey F (agent A) mostly selected

his own preferred color while monkey C (agent B) either selected the left side or monkey F's

preferred color. At the end of the confederate training, both monkeys followed the confederate.

The question was: would they now express turn-taking behavior or would they fall back to their

initial static color coordination behavior?

The pair's coordination behaviour changed noticeably after training with a confederate. In the first

session, after some initial back and forth, the pair converged on mostly selecting B's color, with B

acting faster (**Figure 5A**). The most interesting phase was during the second session, in which

the animals repeatedly alternated between the two colors in long blocks (**Figure 5B**). In the third

session, the pattern was opposite to the first session, with the pair converging mostly on A's color

who acted faster (**Figure 5C**). This time, B tried occasionally to switch to his preferred color, and

A followed for three short periods. The mutual information measures as well as the DCR values

indicate that after confederate training, the coordination strength increased and dynamic

coordination resulted in turn-taking behavior in two sessions (**Figure 5D**, **Figure S6**).

If action visibility is driving this form of dynamic coordination, then the question of who successfully

"insists" on own preference and who "accommodates" the other's preference should depend on

the relative action times of the two agents. Indeed, in the session in which the animals exhibited

apparent turn-taking behavior (**Figure 5B**), the faster animal (Leader) selected the preferred color,

while the slower animal (Follower) had to accommodate. Thus, the dynamic coordination arose

within a competitive interaction.

Figure 5. Confederate-trained macaques show competitive turn-taking behavior. (A) In the first paired session after confederate training (FC-1) indications of turn-taking followed by a convergence on B's target can be seen in the FCO in the top subpanel; note how agent A occasionally tried selecting his preferred target. The action time difference histograms show that agent B acted faster when both agents selected his preferred target (t-test: coordination on red (mean: -211 ms, SD: 112, N: 27) vs. blue (mean: 105 ms, SD: 98, N: 286), t(29.9): -14.1, p<0.001). **(B)** In the next session (FC-2) the pair developed turn-taking; the AT difference histograms show that the pair converged on the faster monkey's target (t-test: coordination on red (mean: -117 ms, SD: 96, N: 238) vs. blue (mean: 90 ms, SD: 104, N: 201), t(411.7): -21.4, p<0.001). **(C)** In the following session (FC-3) the pair mostly converged on A's preferred target, with A now acting faster (t-test: coordination on red (mean: -188 ms, SD: 115, N: 553) vs. blue (mean: -31 ms, SD: 192, N: 33), t(33.4): -4.6, p<0.001); note that as in the first session the agent B occasionally tried his own preferred target, and two of these attempts were reciprocated by A. **(D)** Change in coordination in pair FC from a late session (red circle), to six sessions after confederate training. The "coordination strength" (distance from the origin) increased, and in two sessions there was a dynamic coordination (FC-2, FC-5).

Comparison of dynamic coordination between species

At first glance, the turn-taking in the confederate-trained macaques resembled the long turn-taking observed in some human pairs (cf. **Figure 5D** with **Figure 3**). Yet, the analysis of action time distributions suggests that macaques employed a competitive version of turn-taking in which the faster agent selects its own color (insists) while the slower agent follows (accommodates). Such behavior would be in line with theoretical predictions derived from our evolutionary simulation (Unakafov et al., 2020), showing that competitive turn-taking provides the most effective strategy for a BoS type game when players have high probability to observe partner's choices. To further test the hypothesis that this macaque pair developed a competitive Leader-Follower strategy, we performed a correlation analysis between the modeled visibility of the faster agent's action by the slower agent and the observed likelihood of following to the faster agent's color (the faster animal nearly always selected his own color). **Figure 6A** shows macaque agent A's probability to select his non-preferred color, the "fraction of choosing other's" (the inverted fraction of choosing own) and the modeled probability for A to see B's choice before A made own choice, and the same for

macaque agent B (see Materials and Methods for details on modeling). In both monkeys, there

was a strong correlation between the model and the observation, ranging from 0.92 for smoothed

data to 0.64 for unsmoothed data, all significant at p < 0.001. This analysis further supports the

notion that the confederate-trained macaque pair had developed a competitive turn-taking

(alternating Leader-Follower) strategy. Strong positive correlations between action visibility and

"following" behavior were present in all six post-confederate-trained sessions, indicating bouts of

dynamic coordination (**Table S2**).

Figure 6. Confederate-trained macaque turn-taking is driven by action visibility. The inset on the right shows the
logistic mapping function from action time difference to the probability to see the partner's choice before acting. **(A)** The
2nd session of confederate-trained macaque pair FC (also shown in **Figure 5B**). This panel shows the correlation
between probability of seeing the other move first (p_{see}) and following to the other's preferred target (p_{other}). The "fraction
of choosing other's" is the inverse of the fraction of choosing own (FCO). All curves show running averages of the
respective measure over 8 trials. For agent A the correlation coefficient r(545) was 0.92 and for agent B 0.91 for
smoothed data, and 0.64 and 0.65 for non-smoothed (trial-by-trial) data (all p<0.001). **(B)** The human pair 12 with the
highest positive correlation showed a different pattern: a slow gradual shift from B following A to B leading A over the
course of the session. For A the correlation coefficient r(298) was 0.52 and for B 0.46 for smoothed data, and 0.27 and
0.33 for non-smoothed data (all p<0.001). The same human pair (only one out of ten that exhibited dynamic
coordination) showed a significant AT difference for coordinated choices favoring A vs. B, with each agent being faster
when selecting the own preferred color (coordination on A (mean: -88, SD: 129, N: 96), coordination on B (mean: 12,
SD: 109, N: 180), t(168.8): -6.5, p<0.001, t-test). See **Figure S7** for data on other human pairs.

In contrast, human participants did not show a strong trial-by-trial relationship between relative

action time differences and choices (**Table S3**). Even in the pair with the strongest positive

correlations (pair 12, **Figure 6B**), the "following" behavior was weak and resulted from a gradual

transition from B initially following A to A following B later in the session, rather than from

alternating competitive turn-taking. The lack of such a correlation does not mean that action times

and action visibility played no role in human choices. Indeed, in 47% (9 of 19) human pairs the

relative difference in action times between the agents significantly differed in coordinated trials

compared to uncoordinated trials (**Table S1**). Hence, similarly to macaques, seeing the faster

agent's choice can help the slower agent to coordinate. But unlike the macaque pair, dynamic

human coordination was not driven by the faster agent striving to select the own preferred color

(except human pair 12 shown in **Figure 6B**).

The correlation analysis shows that for the confederate-trained macaques the visibility of the

other's action before making one's own decision correlated with the probability to follow the choice

of the other, with the faster monkey "selfishly" selecting its own preference. Turn-taking in

humans, however, did not rely on temporal competition. To highlight this difference further, we

asked how the two species maintained the coordination. Macaques transitioned between periods

of coordination using short periods of non-coordination, while turn-taking humans tended to switch

seamlessly between the two coordination options without verbal communication (**Figure S8A,B**).

Turn-taking humans showed a high and roughly equal amount of benevolent and selfish seamless

switches by a faster player (**Figure 7A**, teal underlay). The four naïve macaque pairs showing the

highest number of seamless switches used a side-based strategy, which due to the color-to-side

randomization trivially generated seamless color switches. Confederate-trained macaques and

non-turn-taking humans showed only very few seamless switches.

Alternative to seamless switches, agents can end the current coordination block by switching to

non-coordination. These trials can be used to initiate or "signal" a switch of coordination. When

turn-taking humans showed uncoordinated trials they lay along the diagonal, indicating a

benevolent intention to initiate the switch to the other's preferred color in half of these trials (**Figure

7B**). All macaque pairs were below the unity diagonal, indicating selfish preference for own color.

In the confederate-trained macaques the coordination epochs were separated by epochs of non-

coordination that looked like each agent "challenged" the other to accommodate. Indeed, during

challenge initiations and resolutions, the faster macaques tended to act selfishly, while turn-taking

humans showed balanced switching behavior (**Figure S8C,D**). The analysis of action times in

confederate-trained macaques showed that the faster agent slowed down and the slower agent

sped up when transiting to non-coordination, and conversely, the insisting agent sped up to initiate

the coordination on the own color and the accommodating agent slowed down (**Figure S9**).

In sum, there were no indications of competitive turn-taking in humans, as the faster agents

showed balanced selection of both coordination options. Furthermore, in four out of the ten turn-

taking pairs, one agent was significantly faster than the other in coordinated trials as compared

to uncoordinated trials, regardless if it was coordination on its own or the other's color (**Table S1**).

This means that such an agent initiated switches to and from the own color; while being able to

see the faster agent's action likely helped the slower agent to maintain coordination and to

accomplish seamless switches. In contrast, the confederate-trained macaques showed no

seamless switches, but displayed temporal competition in which the faster agent led to his own

color.

Figure 7. Humans and macaques show different switching behavior. A quantification of different transition aspects
as scatter plots of benevolent choices on the y-axis and selfish choices on the x-axis, in the two species. **(A)** The
behavior of a faster agent for all seamless switches between the two coordination modes (red-to-blue and blue-to-red).
Here and in B, the x-axis shows the number of own target selections (selfish), the y-axis the number of non-preferred
target selections (benevolent). Turn-taking humans as well as naïve monkey pairs that coordinated on one side showed
high and balanced numbers of seamless switches, while the confederate-trained macaque pair and the non turn-taking
humans showed only few seamless switches. The dashed contours show the 95% confidence interval around the
diagonal, the teal underlay shows the area with balanced selfish and benevolent choices, the green and the magenta
underlays show the area with significantly more benevolent and selfish choices, respectively. **(B)** The number of
uncoordinated trials in which both subjects selected their respective preferred color (own-own) on the x-axis, and the
number of trials in which both subjects selected their non-preferred color (other-other) on the y-axis. Turn-taking human
pairs mainly lay along the diagonal while macaque pairs cluster in the lower right region (mostly significantly selfish).

Discussion

We studied macaque and human pairs in a coordination game, which offered higher rewards for

selecting the same option, but entailed an inherent conflict about which of the two coordinated

options to select. Both species largely converged on coordinated behavior but in a markedly

different fashion. Half of the human pairs achieved nearly-optimal coordination and fair

(cooperative) turn-taking that equalized the rewards of the two partners, without explicit

communication. Macaques, instead, exploited simpler strategies that maximized reward without

the need to track the actions of the partner in real-time. That is, they used static instead of dynamic
coordination. After two macaques were trained to observe and attend to the partner's choice with
a human confederate, their behavior shifted to competitive turn-taking. In post-confederate-
training sessions, the choice behavior was strongly correlated with trial-by-trial differences in the
reach action times: the faster monkey chose his preferred option and the slower one followed.
Our results show that both humans and macaques make use of action visibility by taking
information about the other's action into account before making their own decisions. There was,
however, a fundamental difference: when coordinating dynamically, monkeys showed competitive
turn-taking while in humans the faster agent often "offered" switching to the partner's preferred
option, exhibiting a form of benevolent and fair turn-taking.

Coordination in humans and macaques

Ten of the 19 human pairs converged on dynamic coordination that balanced the reward,
developing fair, cooperative turn-taking, while another five pairs coordinated on the same fixed
color throughout the session. Similar to the approach of Brosnan and colleagues (Brosnan et al.,
2017, 2011), the human subjects in the present study had to infer the underlying payoff rules while
playing, closely matching the procedure in macaques. Even under such conditions, the majority
of humans arrived at near-optimal behavior within a single session, in our study as well as in other
coordination and anti-coordination games (Brosnan et al., 2017, 2012, 2011). The observed turn-
taking frequency was close to other variants of BoS games with explicit instructions of the payoff
matrix, where approximately 50-60% of human subjects developed turn-taking (Arifovic and
Ledyard, 2018; Ioannou and Romero, 2014; Melis et al., 2016; Sonsino and Sirota, 2003).

In contrast to humans, macaque pairs converged on simpler strategies, either selecting the same
color (56%) or selecting the same side (44%). These two strategies require less cognitive
resources and trial-by-trial coordination effort than turn-taking, because they allow each agent to
make decisions without paying attention to the trial-specific actions of the partner. However, all
but one animal coordinated both via same color or same side selection depending on the partner,
and no animal persisted on the own color with all tested partners. For instance, monkey F insisted
on his preferred color when playing with monkey C, accommodated to the other's color when
playing with monkey T and converged on his right side when playing with monkey M. This result
indicates that macaques took the partner's actions and reward outcomes into account, albeit not
in the strict trial-by-trial or within-trial fashion, resembling the formation of stable color selection
conventions in Guinea baboons in a simple sequential coordination task (Formaux et al., 2022).

For instance, after few sessions of selecting mainly own color, one agent changed either to

selecting their non-preferred color (increasing his reward to 3 and the other's to 4 drops) or to

selecting one side (increasing his reward to an average of 2.5 and the other's to 3 drops). If the

other monkey also started favoring the same side, this increased both agents' reward to 3.5. This

sequence of steps illustrates how macaque pairs can develop their strategy over the course of

several sessions.

While our study is, to the best of our knowledge, the first implementation of a BoS paradigm in

macaques, other 2×2 games have been investigated in several primate species (humans,

chimpanzees, macaques and capuchins). So far, the data suggest that monkeys, unlike humans,

do not spontaneously adopt turn-taking, despite coordinated reciprocity being a successful

evolutionary strategy in simulated non-communicating agents (Colman and Browning, 2009). For

instance, in the anti-coordination Conflict game (Brosnan et al., 2017), capuchins and macaques

converged on only one of the two possible asymmetric equilibria, much like four of our nine

macaque pairs (while many human pairs balanced payoffs by alternating between the two

equilibria). One can argue that turn-taking is not expected in macaques, since it has been

suggested that even apes don't exhibit it. In an exceptionally stringent variant of a BoS-type game,

in which coordinated rope pulls were required to deliver a reward only to one of the two agents

(payoff matrix zero except 0/1 or 1/0 at the coordinated diagonal), chimpanzees, unlike human

children aged 5, did not converge on trial-by-trial alternation/turn-taking (Melis et al., 2016). This

difference was interpreted as a species difference in strategic capability to maintain longer term

cooperation (e.g., by motivating the other to reciprocate). Nevertheless, apes cooperated in 64%

of trials, and in the examples of ape dyad data, one can see some switching between the agents'

choices, albeit not trial-by-trial but in longer blocks. Given that even humans often balance the

fairness over longer time frames, as we observed in several human dyads (e.g., **Figure 1E**), it

was important to test if also macaques might exhibit a similar block-wise turn-taking under

conditions of more trials than ape experiments and less conflict (i.e., when both agents get

increased reward for coordination).

Action visibility and dynamic coordination

The lack of spontaneous dynamic coordination such as turn-taking in naïve macaques in our

experiment cannot be attributed to their inability to infer the upcoming choices of the partner,

because they could observe the partner's actions through the transparent display. Especially for

nonhuman species, coordinating based on mutual choice history might be more demanding than

coordinating using the immediately observable behavior of others. Coordination (or anti-
coordination) typically is improved with increasing information about the other's choice by shifting
from the strictly simultaneous to the sequential mode (Brocas et al., 2018; Rapoport, 1997). The
transparent game approach we adopted here differs from those two classical modes in that each
subject can decide independently when to act (within a certain time window). Under such
conditions, the coordination in chimpanzees is facilitated if one of the agents consistently acts
faster than the partner (Bullinger et al., 2011). Compared to an opaque simultaneous setting,
visual feedback about the partner's choices improved macaques' and capuchins' coordination in
a transparent (called asynchronous) versions of a Stag Hunt game (Brosnan et al., 2012).
Similarly, there were substantial differences in capuchins' and rhesus' behavior in the Conflict /
Chicken game when they had access to the current choice of a partner (Brosnan et al., 2017;
Ong et al., 2021).

In the studies of Brosnan and colleagues, subjects looked at the same split monitor while sitting
next to each other, and could observe each other's decisions as cursor movements. In our setup,
agents sat opposite to each other and saw actual eye and hand movements, combining face-to-
face and action visibility. In humans, face-to-face visibility in a simultaneous iterated Prisoner's
Dilemma (iPD) and in the Ultimatum game significantly improves mutual cooperation, even
without action visibility (Jahng et al., 2017; Tang et al., 2016). These findings support the idea
that non-verbal social signals and, more generally, the observable presence of others influence
decision-making. The ongoing action visibility per se also had a profound effect on mutual
cooperation in iPD (Friedman and Oprea, 2012) and in a web interface-based anti-coordination
game conceptually similar to BoS (Hawkins and Goldstone, 2016). In a competitive reaching task,
a face-to-face transparency allowed human subjects to glean useful information from observing
the relevant hand effector and from seeing the face and the full body of an opponent (Vaziri-
Pashkam et al., 2017). Seeing actual movements rather than relying on abstract representations
of others such as cursor motions might be especially crucial in nonhuman experiments.

The importance of action visibility is also supported by computational modeling. In the continuous-
time cooperation games such as iPD and Stag Hunt where agents could observe and respond to
each other's actions in real-time, cooperation by coaction is more easily obtained and stabilized
against exploitation than the cooperation that relies on delayed reciprocity (van Doorn et al.,
2014). For the iPD and BoS transparent games, we showed that different coordination strategies
are preferable for different probabilities of seeing partner's choice (Unakafov et al., 2020, 2019).
When these probabilities are low (or when agents do not utilize the action visibility), simple

strategies like "win-stay, lose-shift" are most effective. For higher probabilities of seeing partner's

choice more complex strategies emerge, such as coordinated turn-taking and temporal Leader-

Follower, where the faster agent determines the choice of the slower agent. For the selfish agents

this means that the faster one insists on its individually-preferred own target and the slower

accommodates (Unakafov et al., 2020).

Given that many human pairs exhibited spontaneous dynamic coordination, while naïve

macaques did not, we can ask if and how these two species utilized the action visibility. Both

species exploited action visibility to achieve and/or maintain the turn-taking, emphasizing the

importance of the "transparency" of interactive behavior as an important determinant of emerging

strategies. Humans relied on action visibility for seamless switches from one color to another, and

for maintaining coordination within a block of trials. But besides one pair, there was no indication

that humans employed the competitive variant of a Leader-Follower strategy. Instead, faster

agents were as likely to switch from their own to the other's preferred color, indicating benevolent

turn-taking. These results add to the body of literature indicating that humans in a social setting

might base their decisions not only on pure reward maximization (Clavien and Chapuisat, 2016).

Such normative behavior can also reflect prospective planning to avoid conflict and maintain

stable individually-beneficial cooperation in subsequent trials.

Very different dynamics transpired in additional experiments with macaques. Pairing them with a

human confederate who alternated between colors in short blocks demonstrated that macaques

are capable to closely follow partner's actions (Nougaret et al., 2019), and adopt the imposed

"turn-taking". Moreover, temporarily blocking the view of the confederate's hands abruptly

changed macaque behavior, disrupting the coordination. The macaques' coordination was thus

driven by action visibility and not by strategies like "win-stay, lose-shift" or trial counting. In line

with the Leader-Follower strategy, macaques were faster than the confederate when selecting

own color, but waited for the confederate when selecting the confederate's color. Intriguingly,

pairing the two macaques that completed the training with a human confederate resulted in

behavior different from both naïve macaque behavior and from the "following behavior" with the

confederate. These macaques established competitive dynamics, with the faster agent selecting

his preferred color and the slower agent following, as predicted by our evolutionary simulations

(Unakafov et al., 2020). Such competition resulted in either sustained coordination on one fixed

color or in turn-taking. Comparing the temporal signatures of the action time differences in trials

around a switch to or from coordinating on a specific color further indicated that macaque turn-

taking was competitive and dynamic in nature. The break of coordination was triggered by the

faster agent slowing down and the slower one speeding up and selecting his preferred color;
conversely, the transition to coordination was associated with the speeding up agent selecting his
preferred color and the slowing down agent accommodating.

Species differences in general cooperative behavior may explain both the reluctance of naïve
macaques to coordinate dynamically, as well as the competitive nature of dynamic interactions
when they occur. The transparent BoS game requires the agents to cooperate in close proximity
in the pursuit of an immediate reward which necessitates some degree of social tolerance by both
partners. If we can take natural food sharing behavior as a proxy for social tolerance, it is not
equally prevalent among primate species. Humans, apes, some baboon species (Goffe and
Fischer, 2016), and some New World monkey species including capuchins share food between
adults, while rhesus macaques do not even share food with their offspring (Jaeggi and Gurven,
2013). In line with these patterns, mutually beneficial alternating task performance and turn-taking
has been observed in humans (Brosnan et al., 2017; Grueneisen and Tomasello, 2017; Helbing
et al., 2005; Rapoport et al., 1976), apes (de Waal, 1989; Martin et al., 2017), and capuchins
(Parrish et al., 2015), but not in more despotic, less tolerant rhesus macaques (Joly et al., 2017).

Beyond social attitudes, the lack of the cooperative turn-taking in nonhuman primates might reflect
cognitive limitations in long-term planning and perspective taking (Stevens and Hauser, 2004). In
humans performance/level of coordination in BoS and other iterative games was shown to depend
to a larger degree on intelligence (to form good strategies) than on personality or social
predispositions (Proto et al., 2019). While the action transparency decreases the reliance of
coordination on mental representations, the differences between strategies in macaques
compared to humans might still be driven by more limited cognitive capacities.

Limitations and future directions

There are several limitations in our study that have to be considered and possibly addressed in
future experiments. Firstly, we only tested 6 macaques (and 9 pairs), all males, and only one pair
was housed together (others were from neighboring but separate enclosures). Therefore, we
cannot say much about the influence of social rank. Anecdotally, however, it did not seem that
the more dominant agent always prevailed. For instance, one animal (monkey T), the smallest
and likely to be subordinate to all other partners, successfully "insisted" on his own color in 2 out
of 3 cases (cf. (Massen et al., 2010)). Furthermore, we only tested two confederate-trained
macaques so far. While the results are exciting and show that macaques can engage in dynamic
competitive turn-taking, future experiments will need to test how generalizable this pattern is.

Secondly, due to the task target color/side randomization, selecting targets on the same side of

the display resulted in low effort, efficient and fair coordination. We cannot say if fairness was a

contributing factor in convergence of the four naïve macaque pairs to this strategy, although we

deem it highly unlikely. A new experiment where the probability of a specific color to appear on

the same side is parametrically modulated is needed to evaluate the conditions under which this

strategy emerges (or what level of unfairness is accepted). Furthermore, we prompted the human

participants to pay attention to the partner's choices, to facilitate emergence of coordination within

a single session. It would be interesting to test human pairs in several sessions and without any

instructions, similar to macaques, to see if humans might also converge on the fixed side strategy

as means of fair and seamless coordination. It requires preventing the human participants from

any verbal contact after and between the sessions however. Finally, it would be important to vary

the ratio and the range of reward magnitude ("stakes"), as it has been shown that stakes may

strongly affect the individual (Schultz, 2015) and social (Hawkins and Goldstone, 2016) decisions.

Inherently, the nature of reward differed between the species: humans received monetary

compensation while macaques worked for a liquid reward. Note however that in each session

macaques were offered an opportunity to work and obtain more liquid in a solo regime after the

dyadic experiment, so the participation in a dyadic interaction was not forced.

In summary, we show that both, humans and nonhuman primates, dynamically coordinate in real-

time transparent interactions, but macaques compete while humans strive for fairness by

prosocial turn-taking. Enabling action visibility with our transparent setup provides novel

comparative insights into social decision processes as they unfold in real-time during transparent

interactions, and offers a new route to behavioral and neural investigations of dynamic decision-

making in cooperative and competitive contexts (Ferrari-Toniolo et al., 2019; Isoda et al., 2018;

Yoo et al., 2021).

Materials and Methods

Participants

Humans. 38 right-handed subjects (23 females, mean age: 26.1 ± 4.1 SD, range 20 to 41 years)

participated in the study as paid volunteers. Subjects were tested as 19 unique pairs, i.e. each

participant contributed only once. Instructions given to the subjects before the experiment are

provided in Supplementary Materials. In short, subjects were instructed how to operate the setup

and to interpret the auditory feedback as an indicator of the earned reward. They were not given

an explicit description of the task's payoff structure beyond "Your reward will depend on your own
and your partner's choice". Prior to the experiment, subjects were individually familiarized with the
setup and practiced a single-player ("solo") version of the task. All subjects gave written informed
consent for participation after the procedures had been explained to them and before taking part
in the experiment. Experiments were performed in accordance with institutional guidelines for
experiments with humans and adhered to the principles of the Declaration of Helsinki. The
experimental protocol was approved by the ethics committee of the Georg-Elias-Mueller-Institute
for Psychology, University of Goettingen (GEMI 17-06-06 171). We excluded 2 additional pairs
from the analyses due to experimental differences (one pair did not perform the initial individual
training; one pair ended the experiment prematurely).

Macaques. Research with nonhuman primates represents a small but indispensable component
of neuroscience research. The scientists in this study are committed to the responsibility they
have in ensuring the best possible science with the least possible harm to the animals (Roelfsema
and Treue, 2014). The experimental procedures were approved of by the responsible regional
government office (Niedersaechsisches Landesamt fuer Verbraucherschutz und
Lebensmittelsicherheit (LAVES), permits 3392-42502-04-13/1100 and 3319-42502-04-18/2823),
and were conducted in accordance with the European Directive 2010/63/EU, the corresponding
German law governing animal welfare, and German Primate Center institutional guidelines.

Six adult male rhesus monkeys (designated by initials C, E, F, L, M, T) participated in the study,
yielding 9 pairs (each monkey participated in 2 or more pairs). Animals were extensively trained
with positive reinforcement to climb into and stay seated in a primate chair. The animals were
pair- or group-housed in facilities of the German Primate Center (DPZ) in accordance with all
applicable German and European regulations. The facility provides the animals with an enriched
environment, including a multitude of toys and wooden structures (Berger et al., 2018; Calapai et
al., 2017), natural as well as artificial light and access to outdoor space, exceeding the size
requirements of European regulations, and a rich diet including primate biscuits, fruit and
vegetables. During the study the animals had unrestricted access to food and fluid, except on the
days where data were collected or the animal was trained on the behavioral paradigm. On these
days, the animals were allowed access to fluid through their performance in the behavioral
paradigm. The DPZ veterinarians, the animal facility staff, and the lab scientists carefully
monitored the welfare of the animals.

Experimental setup

For maximal comparability of the human and the monkey behavior, we developed a novel Dyadic
Interaction platform in which two human or nonhuman primate subjects co-act in a shared
workspace while sitting face-to-face (**Figure 1A**). Joint dyadic tasks have been previously
implemented in a side-by-side or 90° angle setting with a shared or separate workspaces for each
subject (Brosnan et al., 2017; Chang et al., 2013; Dal Monte et al., 2020; Falcone et al., 2017;
Ferrari-Toniolo et al., 2019; Fujii et al., 2007; Haroush and Williams, 2015), using back-to-back
nontransparent monitors (Jahng et al., 2017; Tang et al., 2016), or in a table-like setting with
opposing subjects acting in a horizontal workspace that is not in the line of sight between the
subjects' faces (Báez-Mendoza et al., 2016; Ong et al., 2021; Yoshida et al., 2011). Only few
studies utilized transparent face-to-face arrangement of the subjects, using a video projector and
two semitransparent mirrors virtually placing the stimuli into the shared plane (Ballesta and
Duhamel, 2015), or incorporating physical targets into transparent Plexiglas screen (Vaziri-
Pashkam et al., 2017). Similarly, with our design we aimed for maximal availability of mutual social
signaling in a face-to-face setting, a shared vertical workspace with computer-controlled stimuli in
the direct line of sight, without risk of injuries due to physical contact, and suitability across
species. We achieved this using a novel transparent display (1920 x 1080 pixels, 121 x 68 cm,
60 Hz, EYE-TOLED-5500, Eyevis, Reutlingen, Germany), amended for dual-side touch sensitivity
(PQLabs, G5S, Freemont, CA, USA) with a custom-built "sandwich" construction. Sitting on either
side of the display, subjects saw each other and the same stimuli. Two proximity sensors (Carlo
Gavazzi CA18-CA30CAF16NA, Lainate, Italy) per subject for the left and right hand respectively,
mounted below the display ("home" buttons) and the two touch panels mounted on either side of
the display registered hand positions of both agents, at 240 Hz temporal resolution. Experimental
control and stimulus presentation were implemented using the EventIDE software package
(Okazolab, Delft, The Netherlands). Liquid reward for monkeys was delivered via computer-
controlled peristaltic fluid pumps, for every correctly performed trial.

Dyadic decision-making task

Bach-or-Stravinsky game. Since we are interested in the effect of mutual action visibility on
coordination behavior, we implemented a transparent version of a Bach-or-Stravinsky (BoS)
game, in which each player's time-continuous visuomotor behavior can be seen by the other
player. Conceptually, in the BoS game two agents are choosing between going to Bach or
Stravinsky concerts. Agent A prefers Bach, Agent B prefers Stravinsky; yet, both prefer going to

the concert together (Osborne and Rubinstein, 1994). Thus, agents wish to coordinate their

behavior but have conflicting interests. This is a classic 2×2 non-zero sum game (also known as

the Battle of the Sexes) with two pure strategy Nash equilibria (Bach-Bach, or Stravinsky-

Stravinsky), and one less efficient mixed strategy Nash equilibrium, where the agents go to their

preferred event more often than the other (Kilgour and Fraser, 1988; Rapoport, 1967).

In our implementation, the agents were choosing between two options represented by two

differently colored targets placed in the left and the right side of the display. Each target color was

associated with a higher reward for one of the agents resulting in individually preferred targets.

This reward schedule, 2 reward units for one color and 1 unit for the other color was trained in

"solo" trials to establish an "own" color preference in each subject. In dyadic trials, an agent

selecting own preferred target was assured to get at least 2 reward units, while selecting the

other's target (individually non-preferred target) yielded at least 1 unit. Additionally, when both

agents selected the same target, a bonus of 2 reward units was added to the payoff of each agent.

Thus, the maximum average joint reward (3.5 units) was obtained on coordinated trials when one

agent selected own preferred target (getting 4 units) while the other chose non-preferred target

(getting 3 units). Note that on any given coordinated trial, i.e. when agents selected the same

target for overall higher reward, this payoff matrix (**Figure 1C**) resulted in an unequal (i.e. unfair)

reward distribution. Hence, the BoS paradigm probes both the ability to realize that coordinated

target-selection results in higher rewards than uncoordinated target-selection, as well as the

ability to perceive and counteract the unfairness/conflict situation.

The task mechanics. On every trial, each subject chose between the individually-preferred and

non-preferred target on the touchscreen. Targets were light blue circles of 35 mm diameter with

either a red (individually-preferred target for Agent A) or a yellow (individually-preferred target for

Agent B) rings (for better visualization, we replaced yellow and red rings with blue and red solid

targets in the figures and the text).

Subjects had to place both hands on the two home buttons for 500 ms to start a trial after the

inter-trial interval (ITI). This allowed us to control which effector (acting hand) was used (**Figure

1D**). Then an initial fixation target without a colored ring appeared on the display (10 cm below

eye level). Subjects had 1500 ms to touch this initial fixation target with the instructed hand. After

both subjects touched the fixation target they had to hold it for 500 ms. Then two choice targets

appeared at one of three different pairs of positions (140 mm to the left and right of the central

fixation target and either at the same height as the central fixation target or 35 mm below or 35

mm above it). We randomized the "red" location equally over all six positions, balanced in sets of

18 trials with each of the six "red" positions appearing three times, and vertically mirrored the

opposite location. Simultaneously with targets' appearance the initial fixation target disappeared,

serving as a go signal. Subjects had 1500 ms to make their choice and touch one of the targets.

After both subjects acquired their chosen target, selected target(s) brightened, and subjects

needed to hold the hand on the target for another 500 ms. At that point, the choices were

evaluated and rewards were dispensed according to the payoff matrix. The amount of reward

earned by each subject was signaled by two sequential series of auditory pulses (always first for

the subject on side B then on side A), with different pitch for each subject. Each pulse was

constructed as a harmonic series with 12 overtones and a fundamental frequency of 443 Hz for

side A and 733 Hz for side B to provide distinct sounds for each agent. For the monkeys, we

delivered liquid as reward (approximately 0.14 ml per pulse) concurrently with the auditory pulses

for the respective monkey's side; humans were instructed to expect "a few" cents per pulse, the

accumulated earnings were paid out as a lump sum after the experiment. After the reward period

subjects had to wait for an inter-trial-interval of 1500 ms before they could initiate the next trial.

The side of the targets was randomized on each trial, i.e., in ~50% the red target was on the right,

the blue on the left, and vice versa. This design can lead to the three following coordination

patterns (or any mix of the three):

i)   Coordinating statically by both agents repeatedly selecting the same fixed color of the
747   target, irrespective of unfair distribution of the rewards.
748   ii)   Coordinating statically by both agents repeatedly selecting the same fixed side. Due to
749   color/side randomization, this pattern "automatically" ensures a fair reward distribution.
750   iii)   Coordinating dynamically by both agents selecting the same target, while picking from
751   both colors and both sides trial-by-trial (e.g. trial 1 red right, trial 2 red left, trial 3 blue
752   left, … etc.). This could result in a fair or unfair reward distribution depending on the
753   ratio of red to blue color selection.

Subjects were not informed about the randomization procedure, so recognizing the "automatic"

fairness of pattern (ii) would have required keeping a count of color occurrences per side from

both subjects.

Human procedure. Human subjects were recruited via a university job website and pairs were

selected based on matching schedules. Subjects were given a brief introduction to the experiment

and a set of verbal instructions on the task mechanics (see Supplementary Materials). This

material included the information "Your reward will depend on your own and your partner's choice"
and a very basic description of the task ("You will have to choose one of the two circles presented
to you. […] You will have to decide and respond quickly."), but did not include details of the payoff
matrix. Participants had to infer the task rules by exploration, similar to the macaques (Brosnan
et al., 2012). After the joint introduction, each subject alone performed 100 individual (solo) trials,
to learn how to operate the touchscreen and to develop a preference for one of the two color
targets (in 5 pairs, we only used 50 solo trials to reduce the experiment's duration for logistical
reasons, **Table S1**). All but one subject who were trained on 100 solo trials and converged to
>75% own color selection did so already in the trials 26-50. Subjects were positioned ~50 cm
from the display with the height of the chairs adjusted such that both subjects' eyes were ~121
cm above the ground. Subjects were instructed to use the right hand. After the solo training, both
subjects entered the setup for the main dyadic task which lasted for 300 or 400 trials. Subjects
were not informed about the number of trials in a session. Each session lasted approximately 1.5
hours. After the experiment we conducted an individual debriefing and paid the earned reward
separately to each subject (10.5 – 26 Euro, mean 19 Euro, **Table S1**).

Macaque procedure. Macaques were brought to the setup in their individual primate chairs. The
chairs were positioned such that the eyes were 30 cm from the display. The monkeys had
previously been trained to perform the basic task structure (hands on proximity sensors, reach to
the initial fixation target with the instructed hand, and select one of the presented choice targets
by reaching to it). Dyads containing monkeys (monkey-monkey, and monkey-human confederate)
were instructed to use the left hand. The animals performed the solo version of the task with
differential rewards to develop a preference for one of the color targets and were only paired with
a conspecific after selecting a higher rewarded target in ≥ 75% of the trials. Thereafter, pairs of
macaques worked together in the dyadic version of the task, for 11 ± 7 sessions (range 4-25).
After the dyadic task we offered monkeys the opportunity to obtain additional liquid in the solo
version of the task for as long as they continued to work, essentially making participation in the
dyadic trials optional.

787   Data analysis

We computed the following six *aggregate measures of choice behavior and coordination,*
explained below*:* fraction of choosing own color, fraction of choosing (objective) left side, mutual
information for color choice, mutual information for side choice, average reward, and dynamic
coordination reward. Unless noted otherwise, we report these measures for the last 200 trials of

each session in order to assess the "steady-state" behavior after allowing for an initial period of
exploration.

Fraction of choosing own / left. The fraction of choosing own color (FCO) is the fraction of trials
(within a certain window) where an agent has selected the individually preferred color (red for
agent A and blue for agent B). Similarly, the fraction of choosing left (FCL) is a fraction of trials
where an agent has selected the target on the "objective" left side of the display (which is the left
side for agent A and right side for agent B). Fractions range from 0 to 1. For figures showing
FCO/FCL over the course of a session we also calculated both measures for the session as a
whole, and in running windows of $w = 8$ trials. In the latter case, FCO and FCL can take the values
of 0, 1/8, 2/8, …, 1. For instance, FCO = 0 means that an agent has selected individually non-
preferred target for 8 trials in a row.

Mutual information. Mutual information (MI) represents the reduction of uncertainty regarding
the values of one time series provided by knowing the values of the other time series. Here we
consider mutual information for the color of the target (MI color) and side (MI side) choices of the
two agents, showing how much information the target/side choices of one agent provide about
the respective choices of the other. Mutual information is measured in bits. Since both target and
side choices are binary (in each trial an agent selects either the preferred or non-preferred target
and either the left or right side), both MI color and MI side range from 0 bit (the choices of one
agent provide no information about the choices of the other) to 1 bit (the choices of one agent can
be inferred precisely from the choices of the other agent).

For instance, if both agents select the objective left target in every odd trial and the objective right
target in every even trial, both MI color = MI side = 1, since for every trial the target and the side
selected by one agent entirely describes the target/side selected by the other. At the same time,
the choices of each agent individually are highly uncertain: both sides and both targets are
selected with the same probability of 0.5. If both agents constantly select the objective left side,
this would result in MI color = 1 due to side-randomization of target color, but MI side = 0, since
there is no uncertainty regarding the side selection and thus no additional knowledge about the
other's choice can reduce the uncertainty.

Formally, mutual information of time series $X = (X_t)$ and $Y = (Y_t)$ is given by

821
$$\mathrm{MI}(X,Y) = \sum_{x,y} p(X_t = x, Y_t = y)\log_2 \frac{p(X_t=x,Y_t=y)}{p(X_t=x)p(Y_t=y)} \qquad (1)$$

where $p(X_t = x)$ is the probability of the value *x* in time series X, $p(Y_t = y)$ is the probability of the

value *y* in time series Y, and $p(X_t = x, Y_t = y)$ is the joint probability to simultaneously have values

x and y in time series *X* and *Y*, respectively, and *x* and *y* can be either 0 or 1, so that the sum is

over all four combinations. Since in our case time series X and Y have finite length, we simply

replace probabilities by relative frequencies. This is known as a naïve estimation of mutual

information, but it is sufficiently precise for binary time series (Grassberger, 1988; Steuer et al.,

2002).

To test whether the MI values were significantly different from zero, we generate Whittle

surrogates for the given choice time series and estimate from them the threshold for the given

significance level (p=0.01 in our case) (Pethel and Hahs, 2014).

Average reward. Average reward (AR) is computed as the average of an agent's payoff across

the session. Note that the reward of individual agent can be in the range of 1 to 4 units, while the

joint reward of a pair ([reward A + reward B]/2) cannot exceed 3.5 (since when one agent gets

payoff of 4, the other agent gets 3). The average reward for completely independent choices of

two agents with 50% probability for either target is 2.5 (but note that an achieved reward of 2.5 is

not a positive proof of independent choices).

Dynamic coordination reward. Dynamic coordination reward (DCR) is the surplus reward of the

two agents compared to the reward they would get by playing randomly. By *playing randomly* we

mean that choices of the agents in each round are independent of the history and of the current

choices of the partner. The "random reward" is computed by selecting the two color targets with

the same probabilities as actually observed in the two agents, but randomly permuting the choices

over trials. For our payoff matrix, the range of DCR is [-1,1], with -1 corresponding to very inefficient

playing (alternating selection of the two anti-coordination options), while 1 corresponds to very

efficient playing with explicit coordination (for instance, turn-taking). DCR is hence a measure of

dynamic (reciprocal) coordination. For instance, if both agents would coordinate statically by

constantly selecting one and the same side, this would result in $\mathrm{DCR} = 0$, even though this

coordination pattern still yields the maximum average reward of 3.5.

Formally, DCR is defined as the actual average reward of a pair (R_{actual}) minus the reward the

agents would get if they were playing randomly (R_{PR}). $\mathrm{DCR} = R_{\mathrm{actual}} - R_{\mathrm{PR}}$. The reward for playing

randomly (R_{PR}) depends only on eight probabilities, four for each agent. Below index i indicates

the agent and stands for either A or B:

$P_{i,1,\text{left}}$ - probability to select non-preferred objective left target (which is the left side for agent A

and right side for agent B),

$P_{i,1,\text{right}}$ - probability to select non-preferred objective right target,

$P_{i,2,\text{left}}$ - probability to select preferred objective left target,

$P_{i,2,\text{right}}$ - probability to select preferred objective right target.

Note that these probabilities are not independent on each other. First, for $i = A, B$, it holds

859
$$P_{i,1,\text{left}} + P_{i,2,\text{left}} + P_{i,1,\text{right}} + P_{i,2,\text{right}} = 1 \qquad (2)$$

860 Second, it holds

861
$$P_{A,2,\text{left}} + P_{A,1,\text{right}} = P_{B,1,\text{left}} + P_{B,2,\text{right}} = Q_{\text{left}} \qquad (3)$$

862 and

863
$$P_{A,1,\text{left}} + P_{A,2,\text{right}} = P_{B,2,\text{left}} + P_{B,1,\text{right}} = Q_{\text{right}} \qquad (4)$$

864 where Q_{left} and Q_{right} are the probability of agent A's preferred target to appear on the left and on

865 the right, respectively (and of B's preferred target appear on the right and on the left). Given the

866 independent random selection from trial to trial described above, Q-values should approximate

867 0.5 for larger N of trials.

868 Average reward of two agents for playing randomly is computed as follows:

869
$$R_{\text{PR}} = 3.5(p_{4,3} + p_{3,4}) + 2p_{2,2} + p_{1,1} \qquad (5)$$

870 where $p_{a,b}$ is the probability that by random playing agent A gets reward a and agent B reward b.

871 These probabilities are given by the following equations:

872
$$p_{1,1} = \frac{P_{A,1,\text{right}} \cdot P_{B,1,\text{left}}}{Q_{\text{left}}} + \frac{P_{A,1,\text{left}} \cdot P_{B,1,\text{right}}}{Q_{\text{right}}} \qquad (6)$$

873
$$p_{2,2} = \frac{P_{A,2,\text{left}} \cdot P_{B,2,\text{right}}}{Q_{\text{left}}} + \frac{P_{A,2,\text{right}} \cdot P_{B,2,\text{left}}}{Q_{\text{right}}} \qquad (7)$$

874
$$p_{4,3} = \frac{P_{A,2,\text{left}} \cdot P_{B,1,\text{left}}}{Q_{\text{left}}} + \frac{P_{A,2,\text{right}} \cdot P_{B,1,\text{right}}}{Q_{\text{right}}} \qquad (8)$$

875
$$p_{3,4} = \frac{P_{A,1,\text{right}} \cdot P_{B,2,\text{right}}}{Q_{\text{left}}} + \frac{P_{A,1,\text{left}} \cdot P_{B,2,\text{left}}}{Q_{\text{right}}} \qquad (9)$$

To see why this is the case, consider, for instance, $p_{1,1}$. Both agents get reward of 1 when they

both select the other's preferred target, either when target of agent A appears on the left side

(probability of this is encoded by the first term) or on the right side (second term).

To compute confidence intervals, we use the fact that FCL_i and FCO_i for an agent *i* can be

considered as binomially distributed, thus radiuses of their confidence intervals ΔFCL_i and

ΔFCO_i can be estimated by the classic method of approximating the distribution of error around

binomially-distributed observation with a normal distribution. To obtain the confidence interval for

DCR, it is sufficient to compute maximal and minimal possible DCR given that FCO and SCL of the

two agents are within the respective confidence intervals. Note that $\text{FCL}_i = P_{i,1,\text{left}} + P_{i,2,\text{left}}$,

$\text{FCO}_i = P_{i,2,\text{left}} + P_{i,2,\text{right}}$, thus all the probabilities necessary for calculating DCR can be computed

from FCL_i and FCO_i given that Q_{left} and Q_{right} are fixed to 0.5 in the reported experiments. Simple

analysis reveals that minimum and maximum DCR should be at the edges of the 4-D confidence

interval formed by FCL_i and FCO_i of the two agents, which reduces the problem to testing 16 DCR

values computed for $\text{FCL}_i \pm \Delta\text{FCL}_i$ and $\text{FCO}_i \pm \Delta\text{FCO}_i$.

For the main analysis presented in the Results, we used last 200 trials to compute DCR values.

Using the last 150 or 250 trials for DCR analysis resulted in the exact same 10 human pairs with

significant results as using the last 200.

Action time measurements. We measured the reaction time from the onset of the choice targets

to the release time of the initial fixation target (t_{release}) and to the acquisition time of the selected

target, $t_{\text{acquisition}}$ individually for each subject. We then calculated the movement time t_{movement}

and action time t_{action} as follows:

897
$$t_{\text{movement}} = t_{\text{acquisition}} - t_{\text{release}} \qquad (10)$$

898
$$t_{\text{action}} = t_{\text{release}} + \frac{t_{\text{movement}}}{2} \qquad (11)$$

We use t_{action}, the half-way point between t_{release} and $t_{\text{acquisition}}$, i.e. the duration from target

stimulus onset to the half-time of the reach movement, as a proxy for the estimated time at which

the trajectory of each subject's reach movement should be evident for the other agent. Typical

movement time values (mean ± SD across trials) in our experiment ranged from 314 ± 104 ms for

humans, 171 ± 77 ms for macaques, and 180 ± 64 ms for the confederate trained macaque pair.

Estimated probability to see partner's choice. When an agent is acting slower than the partner,

there is a chance for this agent to see the partner's choice and use this information to make the

own choice. We therefore modeled the probability to see the partner's choice as a logistic function

of the difference of the agents' action times, in each trial. The logistic function $p_{\text{see}}(t)$ had an

inflection point at 50 ms ($p_{see}(50) = 0.5$) and reached its plateau phase at 150 ms ($p_{see}(150) =$

0.98). Formally, the function was given by the following equation

910
$$p_{see}(t) = \frac{1}{1 + e^{-k(t - \Delta T_0)}} \qquad (12)$$

with $k = 0.04$ (steepness of the slope) and $\Delta T_0 = 50$ ms (inflection point). The values of p_{see} were

used for the analysis of correlation with probability of selecting the other's color (cf. **Figure 6**). We

also tested a wide range of these parameters (k fixed at 0.04, $\Delta T_0 = \{12.5, 25, 50, 75, 100, 200\}$ ms

and ΔT_0 fixed at 50 ms, $k = \{0.01, 0.02, 0.04, 0.08, 0.16\}$), and confirmed that the values of

correlation were robust in respect to these parameters; only the longest ΔT_0 = 200 ms resulted in

a noticeable drop of resultant correlations.

References

Arifovic J, Ledyard J. 2018. Learning to alternate. *Exp Econ* **21**:692–721. doi:10.1007/s10683-018-9568-1

Báez-Mendoza R, Coeverden CR van, Schultz W. 2016. A neuronal reward inequity signal in primate striatum. *Journal of Neurophysiology* **115**:68–79. doi:10.1152/jn.00321.2015

Ballesta S, Duhamel J-R. 2015. Rudimentary empathy in macaques' social decision-making. *Proc Natl Acad Sci USA* **112**:15516–15521. doi:10.1073/pnas.1504454112

Berger M, Calapai A, Stephan V, Niessing M, Burchardt L, Gail A, Treue S. 2018. Standardized automated training of rhesus monkeys for neuroscience research in their housing environment. *J Neurophysiol* **119**:796–807. doi:10.1152/jn.00614.2017

Bornstein G, Budescu D, Zamir S. 1997. Cooperation in intergroup, N-person, and two-person games of chicken. *Journal of Conflict Resolution* **41**:384–406.

Brocas I, Carrillo JD, Sachdeva A. 2018. The path to equilibrium in sequential and simultaneous games: A mousetracking study. *Journal of Economic Theory* **178**:246–274. doi:10.1016/j.jet.2018.09.011

Brosnan SF. 2018. Insights into human cooperation from comparative economics. *Nature Human Behaviour* **2**:432–434. doi:10.1038/s41562-018-0383-7

Brosnan SF, Parrish A, Beran MJ, Flemming T, Heimbauer L, Talbot CF, Lambeth SP, Schapiro SJ, Wilson BJ. 2011. Responses to the Assurance game in monkeys, apes, and humans using equivalent procedures. *PNAS* **108**:3442–3447. doi:10.1073/pnas.1016269108

Brosnan SF, Price SA, Leverett K, Prétôt L, Beran M, Wilson BJ. 2017. Human and monkey responses in a symmetric game of conflict with asymmetric equilibria. *Journal of Economic Behavior & Organization* **142**:293–306. doi:10.1016/j.jebo.2017.07.037

Brosnan SF, Wilson BJ, Beran MJ. 2012. Old World monkeys are more similar to humans than New World monkeys when playing a coordination game. *Proceedings of the Royal Society of London B: Biological Sciences* **279**:1522–1530. doi:10.1098/rspb.2011.1781

Bullinger AF, Wyman E, Melis AP, Tomasello M. 2011. Coordination of chimpanzees (Pan troglodytes) in a stag hunt game. *International Journal of Primatology* **32**:1296–1310.

Byrne RW, Bates LA. 2010. Primate Social Cognition: Uniquely Primate, Uniquely Social, or Just Unique? *Neuron* **65**:815–830. doi:10.1016/j.neuron.2010.03.010

Calapai A, Berger M, Niessing M, Heisig K, Brockhausen R, Treue S, Gail A. 2017. A cage-based training, cognitive testing and enrichment system optimized for rhesus macaques in neuroscience research. *Behavior research methods* **49**:35–45.

Camerer CF. 2003. Behavioral Game Theory: Experiments in Strategic Interaction, First Edition edition. ed. New York, N.Y. : Princeton, N.J: Princeton University Press.

Chang SWC, Brent LJN, Adams GK, Klein JT, Pearson JM, Watson KK, Platt ML. 2013. Neuroethology of primate social behavior. *Proc Natl Acad Sci USA* **110 Suppl 2**:10387–10394. doi:10.1073/pnas.1301213110

Clavien C, Chapuisat M. 2016. The evolution of utility functions and psychological altruism. *Studies in History and Philosophy of Science Part C: Studies in History and Philosophy of Biological and Biomedical Sciences* **56**:24–31. doi:10.1016/j.shpsc.2015.10.008

Colman AM, Browning L. 2009. Evolution of cooperative turn-taking. *Evolutionary Ecology Research* 949–963.

Conty L, Dezecache G, Hugueville L, Grèzes J. 2012. Early Binding of Gaze, Gesture, and Emotion: Neural Time Course and Correlates. *J Neurosci* **32**:4531–4539. doi:10.1523/JNEUROSCI.5636-11.2012

Dal Monte O, Chu CCJ, Fagan NA, Chang SWC. 2020. Specialized medial prefrontal–amygdala coordination in other-regarding decision preference. *Nat Neurosci* **23**:565–574. doi:10.1038/s41593-020-0593-y

de Waal FB. 1989. Food sharing and reciprocal obligations among chimpanzees. *Journal of Human Evolution* **18**:433–459.

Devaine M, San-Galli A, Trapanese C, Bardino G, Hano C, Jalme MS, Bouret S, Masi S, Daunizeau J. 2017. Reading wild minds: A computational assay of Theory of Mind sophistication across seven primate species. *PLOS Computational Biology* **13**:e1005833. doi:10.1371/journal.pcbi.1005833

Dugatkin LA, Mesterton-Gibbonsand M, Houston AI. 1992. Beyond the prisoner's dilemma: Toward models to discriminate among mechanisms of cooperation in nature. *Trends in ecology & evolution* **7**:202–205.

Duguid S, Wyman E, Bullinger AF, Herfurth-Majstorovic K, Tomasello M. 2014. Coordination strategies of chimpanzees and human children in a Stag Hunt game. *Proceedings of the Royal Society of London B: Biological Sciences* **281**:20141973.

Falcone R, Cirillo R, Ferraina S, Genovesio A. 2017. Neural activity in macaque medial frontal cortex represents others' choices. *Scientific Reports* **7**. doi:10.1038/s41598-017-12822-5

Ferrari-Toniolo S, Visco-Comandini F, Battaglia-Mayer A. 2019. Two Brains in Action: Joint-Action Coding in the Primate Frontal Cortex. *J Neurosci* **39**:3514–3528. doi:10.1523/JNEUROSCI.1512-18.2019

Formaux A, Paleressompoulle D, Fagot J, Claidière N. 2022. The experimental emergence of convention in a non-human primate. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**:20200310. doi:10.1098/rstb.2020.0310

Friedman D, Oprea R. 2012. A Continuous Dilemma. *American Economic Review* **102**:337–363. doi:10.1257/aer.102.1.337

Frith U, Frith C. 2010. The social brain: allowing humans to boldly go where no other species has been. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**:165–176. doi:10.1098/rstb.2009.0160

Fruteau C, van Damme E, Noë R. 2013. Vervet Monkeys Solve a Multiplayer "Forbidden Circle Game" by Queuing to Learn Restraint. *Current Biology* **23**:665–670. doi:10.1016/j.cub.2013.02.039

Fujii N, Hihara S, Iriki A. 2007. Dynamic Social Adaptation of Motion-Related Neurons in Primate Parietal Cortex. *PLoS ONE* **2**:e397. doi:10.1371/journal.pone.0000397

Goffe AS, Fischer J. 2016. Meat sharing between male and female Guinea baboons (*Papio papio*). *Primate Biology* **3**:1–8. doi:https://doi.org/10.5194/pb-3-1-2016

Grassberger P. 1988. Finite sample corrections to entropy and dimension estimates. *Physics Letters A* **128**:369–373.

Grueneisen S, Tomasello M. 2017. Children coordinate in a recurrent social dilemma by taking turns and along dominance asymmetries. *Developmental psychology* **53**:265.

Haroush K, Williams ZM. 2015. Neuronal prediction of opponent's behavior during cooperative social interchange in primates. *Cell* **160**:1233–1245. doi:10.1016/j.cell.2015.01.045

Hawkins RXD, Goldstone RL. 2016. The Formation of Social Conventions in Real-Time Environments. *PLoS One* **11**. doi:10.1371/journal.pone.0151670

Helbing D, Schönhof M, Stark H-U, Ho\lyst JA. 2005. How individuals learn to take turns: Emergence of alternating cooperation in a congestion game and the prisoner's dilemma. *Advances in Complex Systems* **8**:87–116.

Herrmann E, Call J, Hernàndez-Lloreda MV, Hare B, Tomasello M. 2007. Humans Have Evolved Specialized Skills of Social Cognition: The Cultural Intelligence Hypothesis. *Science* **317**:1360–1366. doi:10.1126/science.1146282

Ioannou CA, Romero J. 2014. A generalized approach to belief learning in repeated games. *Games and Economic Behavior* **87**:178–203. doi:10.1016/j.geb.2014.05.007

Isoda M, Noritake A, Ninomiya T. 2018. Development of social systems neuroscience using macaques. *Proceedings of the Japan Academy, Series B* **94**:305–323. doi:10.2183/pjab.94.020

Jaeggi AV, Burkart JM, Van Schaik CP. 2010. On the psychology of cooperation in humans and other primates: combining the natural history and experimental evidence of prosociality. *Philos Trans R Soc Lond B Biol Sci* **365**:2723–2735. doi:10.1098/rstb.2010.0118

Jaeggi AV, Gurven M. 2013. Natural cooperators: food sharing in humans and other primates. *Evol Anthropol* **22**:186–195. doi:10.1002/evan.21364

Jahng J, Kralik JD, Hwang D-U, Jeong J. 2017. Neural dynamics of two players when using nonverbal cues
to gauge intentions to cooperate during the Prisoner's Dilemma Game. *NeuroImage* **157**:263–274.
doi:10.1016/j.neuroimage.2017.06.024

Joly M, Micheletta J, De Marco A, Langermans JA, Sterck EHM, Waller BM. 2017. Comparing physical and
social cognitive skills in macaque species with different degrees of social tolerance. *Proc Biol Sci*
284. doi:10.1098/rspb.2016.2738

Kilgour DM, Fraser NM. 1988. A taxonomy of all ordinal 2 × 2 games. *Theor Decis* **24**:99–117.
doi:10.1007/BF00132457

Martin CF, Biro D, Matsuzawa T. 2017. Chimpanzees spontaneously take turns in a shared serial ordering
task. *Scientific Reports* **7**:14307. doi:10.1038/s41598-017-14393-x

Massen JJM, van den Berg LM, Spruijt BM, Sterck EHM. 2010. Generous Leaders and Selfish Underdogs:
Pro-Sociality in Despotic Macaques. *PLOS ONE* **5**:1–5. doi:10.1371/journal.pone.0009734

McDonald KR, Broderick WF, Huettel SA, Pearson JM. 2019. Bayesian nonparametric models characterize
instantaneous strategies in a competitive dynamic game. *Nature Communications* **10**:1–12.
doi:10.1038/s41467-019-09789-4

Melis AP, Grocke P, Kalbitz J, Tomasello M. 2016. One for you, one for me: Humans' unique turn-taking
skills. *Psychological science* **27**:987–996.

Noë R. 2006. Cooperation experiments: coordination through communication versus acting apart together.
Animal Behaviour **71**:1–18. doi:10.1016/j.anbehav.2005.03.037

Nougaret S, Ferrucci L, Genovesio A. 2019. Role of the social actor during social interaction and learning
in human-monkey paradigms. *Neuroscience & Biobehavioral Reviews* **102**:242–250.
doi:10.1016/j.neubiorev.2019.05.004

Ong WS, Madlon-Kay S, Platt ML. 2021. Neuronal correlates of strategic cooperation in monkeys. *Nature
Neuroscience* **24**:116–128. doi:10.1038/s41593-020-00746-9

Osborne MJ, Rubinstein A. 1994. A Course in Game Theory, First Edition, Twelf edition. ed. Cambridge,
Mass: The MIT Press.

Parrish A, Brosnan S, Beran M. 2015. Capuchin monkeys alternate play and reward in a dual computerized
task. *Animal Behavior and Cognition* **2**:334–347.

Pethel S, Hahs D. 2014. Exact test of independence using mutual information. *Entropy* **16**:2839–2849.

Peysakhovich A, Nowak MA, Rand DG. 2014. Humans display a 'cooperative phenotype' that is domain
general and temporally stable. *Nature Communications* **5**:4939. doi:10.1038/ncomms5939

Platt ML, Seyfarth RM, Cheney DL. 2016. Adaptations for social cognition in the primate brain. *Philos Trans
R Soc Lond, B, Biol Sci* **371**:20150096. doi:10.1098/rstb.2015.0096

Proto E, Rustichini A, Sofianos A. 2019. Intelligence, Personality, and Gains from Cooperation in Repeated
Interactions. *Journal of Political Economy* **127**:1351–1390. doi:10.1086/701355

Rapoport A. 1997. Order of Play in Strategically Equivalent Games in Extensive Form. *Int J Game Theory*
26:113–136. doi:10.1007/s001820050021

Rapoport A. 1973. Two-Person Game Theory. University of Michigan Press.

Rapoport A. 1967. Exploiter, leader, hero, and martyr: The four archetypes of the 2 × 2 game. *Syst Res*
12:81–84. doi:10.1002/bs.3830120202

Rapoport A, Guyer MJ, Gordon DG. 1976. The 2x2 Game. University of Michigan Press.

Rilling JK, Gutman DA, Zeh TR, Pagnoni G, Berns GS, Kilts CD. 2002. A Neural Basis for Social
Cooperation. *Neuron* **35**:395–405. doi:10.1016/S0896-6273(02)00755-9

Rilling JK, Sanfey AG. 2011. The neuroscience of social decision-making. *Annu Rev Psychol* **62**:23–48.
doi:10.1146/annurev.psych.121208.131647

Rizzolatti G, Fogassi L, Gallese V. 2001. Neurophysiological mechanisms underlying the understanding
and imitation of action. *Nature reviews Neuroscience* **2**:661–70. doi:10.1038/35090060

Roelfsema PR, Treue S. 2014. Basic Neuroscience Research with Nonhuman Primates: A Small but
Indispensable Component of Biomedical Research. *Neuron* **82**:1200–1204.
doi:10.1016/j.neuron.2014.06.003

Sanfey AG. 2007. Social Decision-Making: Insights from Game Theory and Neuroscience. *Science*
318:598–602. doi:10.1126/science.1142996

Santos LR, Rosati AG. 2015. The Evolutionary Roots of Human Decision Making. *Annu Rev Psychol*
66:321–347. doi:10.1146/annurev-psych-010814-015310

Schultz W. 2015. Neuronal Reward and Decision Signals: From Theories to Data. *Physiological Reviews*
95:853–951. doi:10.1152/physrev.00023.2014

Silk JB. 2007. The strategic dynamics of cooperation in primate groups. *Advances in the Study of Behavior*
37:1–41.

Skyrms B. 2003. The Stag Hunt and the Evolution of Social Structure. Cambridge University Press.

Smith JM. 1979. Game theory and the evolution of behaviour. *Proc R Soc Lond, B, Biol Sci* **205**:475–488.
doi:10.1098/rspb.1979.0080

Sonsino D, Sirota J. 2003. Strategic pattern recognition—experimental evidence. *Games and Economic*
Behavior **44**:390–411.

Steuer R, Kurths J, Daub CO, Weise J, Selbig J. 2002. The mutual information: detecting and evaluating
dependencies between variables. *Bioinformatics* **18**:S231–S240.

Stevens JR, Hauser MD. 2004. Why be nice? Psychological constraints on the evolution of cooperation.
Trends Cogn Sci (Regul Ed) **8**:60–65. doi:10.1016/j.tics.2003.12.003

Tang H, Mai X, Wang S, Zhu C, Krueger F, Liu C. 2016. Interpersonal brain synchronization in the right
temporo-parietal junction during face-to-face economic exchange. *Soc Cogn Affect Neurosci* **11**:23–
32. doi:10.1093/scan/nsv092

Tremblay S, Sharika KM, Platt ML. 2017. Social Decision-Making and the Brain: A Comparative
Perspective. *Trends in Cognitive Sciences* **21**:265–276. doi:10.1016/j.tics.2017.01.007

Unakafov AM, Schultze T, Gail A, Moeller S, Kagan I, Eule S, Wolf F. 2020. Emergence and suppression
of cooperation by action visibility in transparent games. *PLOS Computational Biology* **16**:e1007588.
doi:10.1371/journal.pcbi.1007588

Unakafov AM, Schultze T, Kagan I, Moeller S, Gail A, Treue S, Eule S, Wolf F. 2019. Evolutionary
successful strategies in a transparent iterated Prisoner's Dilemma. *EvoApplications 2019, Lecture*
Notes in Computer Science **11454**:204–219. doi:10.1101/524348

van Doorn GS, Riebli T, Taborsky M. 2014. Coaction versus reciprocity in continuous-time models of
cooperation. *Journal of Theoretical Biology* **356**:1–10. doi:10.1016/j.jtbi.2014.03.019

Vaziri-Pashkam M, Cormiea S, Nakayama K. 2017. Predicting actions from subtle preparatory movements.
Cognition **168**:65–75. doi:10.1016/j.cognition.2017.06.014

von Neumann J, Morgenstern O. 1944. Theory of Games and Economic Behavior. Princeton University
Press.

Waal F de, Waal FBM. 2007. Chimpanzee Politics: Power and Sex Among Apes. JHU Press.

Wedekind C, Milinski M. 1996. Human cooperation in the simultaneous and the alternating Prisoner's
Dilemma: Pavlov versus Generous Tit-for-Tat. *Proceedings of the National Academy of Sciences of*
the United States of America **93**:2686–2689.

Yoo SBM, Hayden BY, Pearson JM. 2021. Continuous decisions. *Phil Trans R Soc B* **376**:20190664.
doi:10.1098/rstb.2019.0664

Yoshida K, Saito N, Iriki A, Isoda M. 2011. Representation of Others' Action by Neurons in Monkey Medial
Frontal Cortex. *Current Biology* **21**:249–253. doi:10.1016/j.cub.2011.01.004

1107

1108

Acknowledgements

We thank Elisheba Crecca, Tarana Nigam and Roberta Nocerino for their help with the data collection, and Daniela Lazzarini, Janine Kuntze, Sina Plümer and Klaus Heisig for technical support.

Funding

This work was supported by a funding from the Ministry for Science and Education of Lower Saxony ("Top Level Research in Lower Saxony") and the Volkswagen Foundation through the "Niedersächsisches Vorab", https://www.volkswagenstiftung.de/en/funding/niedersaechsisches-vorab (JF, AG, ST, IK). Additional support was provided by the Leibniz Association through funding for the Leibniz ScienceCampus Primate Cognition, https://www.primate-cognition.eu (JF, AG, ST, IK). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

JF, ST and IK wrote the key grants to support his study. SM, AMU, AG, ST and IK conceptualized and designed the task. SM implemented the task and conducted the experiments. SM and AMU analyzed the data. SM, AMU and IK prepared figures. SM, AMU and IK wrote the initial version of the manuscript. All authors discussed and interpreted the findings, and revised the manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability

The code and datasets generated and analyzed in the current study are available from the corresponding author on reasonable request, and will be uploaded to Open Science Framework data repository (https://osf.io/), pending scientific review.

Supplementary Materials for

Human and macaque pairs employ different coordination strategies in a dyadic decision game with face-to-face action visibility

Sebastian Möller, Anton M. Unakafov, Julia Fischer, Alexander Gail, Stefan Treue, Igor Kagan

This PDF file includes:

Supplementary Materials and Methods

Written instructions read by human subjects

Instructions read to human subjects by experimenter

Abbreviations

Figures S1 to S9

Tables S1 to S3

Supplementary Materials and Methods

Written instructions read by human subjects

Description of the study

This study is the investigation of the behavioral correlates of social decision making while playing a game with a partner. Every day we have to make decisions that depend not only on our own needs and goals but also on the needs and goals of others: for instance, while working on a project with colleagues, planning vacation with friends and family, getting on the bus or shopping groceries, etc. With the help of the task presented in this study we will investigate how people make such decisions.

The course of the study

You will complete one session of a decision making task on the computer together with your partner. You will have to choose one of the two circles presented to you. Your partner will have to perform the same task on his/her side of the touchscreen. You will have to decide and respond quickly. If either you or your partner are too slow, the trial will be aborted without any reward. Your reward will depend on your own and your partner's choice. After the decision is made, you both will receive different auditory feedbacks, denoting your reward and the reward of your partner. Please do not talk to your partner during the session. After the session, we will ask you several questions about the experiment.

Instructions read to human subjects by experimenter

1) You rest both hands on the gray board at the two round objects (touch sensors).
2) A central touch target will appear.
3) Move your right hand to the target and hold. Use the right hand during the entire session.
4) While the target brightens up keep holding your finger on the target.
5) Then two colored choice targets appear and the central target disappears.
6) Make your choice and touch the chosen target within 1.5 seconds.
7) All touched targets will brighten up, keep holding until the targets disappear.
8) Please note that both selected targets will brighten up:
   In case both players selected the same target only that single target will brighten;
   In case both players selected different targets, both targets will brighten.
9) Now two streams of auditory beeps will signify the earned reward for each player: each beep corresponds to a few cents. Please try to learn the sound related to your reward during the training trials.
10) While the audio plays move the hand back to the two touch sensors.
11) Go to 1.

Abbreviations

AT   Action time

BoS   Bach-or-Stravinsky game (also known as the Battle of the Sexes)

DCR   Dynamic coordination reward

FCL   Fraction of choosing left

FCO   Fraction of choosing own

MI   Mutual information

MIS   Mutual information between side choices

MIT   Mutual information between target color choices

(i)PD   (iterated) Prisoner's dilemma

SD   Standard deviation

Supplementary Figures

Supplementary Figure S1, related to Figure 2. Performance improvement in human and monkey pairs. (A) Scatter plot of the average joint reward earned by all 19 human pairs for the first 100 versus the last 200 trials. Note that 1 is the minimal possible average joint reward and 3.5 is the maximum. Filled circles denote pairs with significant

change of the ratio of uncoordinated to coordinated choices between the early (first 100) and late (last 200) trials ($p < 0.05$, Bonferroni-corrected Fisher's exact test of the proportion of coordinated versus uncoordinated trials early versus late); in 9 of 19 pairs the increase in coordination was significant. Over all pairs there was a significant increase in median reward from early to late trials (Wilcoxon signed rank test: N: 19; first 100 median 2.83, last 200 median 3.39, signed rank 12, $p < 0.0011$, median change 0.56). **(B)** Average fraction of choosing left for all human pairs for early versus late trials. Note that human pairs cluster around the main diagonal indicating balanced side choices. **(C-F)** Dyadic task increased average joint reward in monkey pairs: **(C)** Average reward increased in early versus late trials in the initial early session of monkey pairs (Wilcoxon signed rank test: N: 9, first 100 median 2.53, last 200 median 2.75, signed rank 0, $p < 0.004$, median change 0.23; 4 of 9 pairs individually significant). **(D)** The increase was not significant in late sessions (Wilcoxon signed rank test: N: 9, first 100 median 3.22, last 200 median 3.29, signed rank 7, $p < 0.08$, median change: 0.08; 3 of 9 pairs individually significant). **(E)** Comparing the average reward between the last 200 trials of early and late session for each monkey pair also showed a significant increase in coordination (Wilcoxon signed rank test: N: 9, early session median 2.75, late session median 3.29, signed rank 0, $p < 0.004$, median change: 0.54; 8 of 9 pairs individually significant). Monkeys showed some mild intra-session increase in coordination (as seen in the increased median average reward) but they only reached the human level of increase (~0.5 units) in median average reward/coordination when comparing late with early session data (cf. A with E). **(F)** Average fraction of choosing left for monkey early versus late sessions; note the two groups, those that converged on selecting the same side in the upper right corner (same pairs clustered around an FCO of 0.5, see **Figure 2B**), and those that converged on selecting the same target color, around the center of the plot (same pairs clustered at the top left of **Figure 2B**).

Supplementary Figure S2, related to Figure 2. Performance of naïve human and macaque pairs. (A) Human pairs: aggregate measures of choice behavior and coordination for 19 naïve human pairs. Panels C to E are sorted by the average joint reward; the example in **Figure 1E,F** is from pair 2. Individual measures for A are shown in red, for B in blue, measures for the pair are in purple. **(A)** The larger of the two average FCO values in each pair versus the smaller of the two. The diagonal line indicates equal frequency of own choices for both agents. **(B)** The average reward for each agent. The dotted line at 2.5 shows the reward value for random choices, the line at 3.5 shows the maximal attainable average joint reward. **(C)** The dynamic coordination reward (DCR), the error bars show 95% confidence intervals; 53% of pairs (3, 1, 13, 2, 12, 7, 9, 17, 8, 6) showed significant DCR. **(D)** Mutual information for target choices. **(E)** Mutual information for side choices. **(F-J) Macaque pairs late session:** same measures as A-E for a late session for each pair. Macaque pairs converged on jointly selecting either the same side or on jointly selecting one of the two color targets. **(K-O) Macaque pairs early session:** same measures as A-E for an early session for each pair. Macaques tended to insist on their respective higher-reward color, with only pair MF converging mostly on side coordination.

Supplementary Figure S3, related to Figure 2. Exemplary development of choice behavior in macaque pair TE over 6 sessions. **(A, C)** The running average of the fraction of choosing own from an early **(A)** and a late session **(C)** of pair TE. **(B, D)** The running average of the fraction of choosing (objective) left from an early **(B)** and a late session **(D)** for the same pair. Conventions are the same as in **Figure 1E,F**. The pair shifted from each selecting their individually preferred target to mostly jointly selecting the objective left side. **(E-J)** The change in aggregate measures of choice behavior and coordination over the course of 6 sessions for the last 200 trials of each session. There was a clear development from selecting own targets to coordination by virtue of converging on mostly selecting the target on the same side of the transparent display. This is also reflected in the gradual increase of the target MI and the gradual decrease of the side MI. The zero DCR values indicate no significant dynamic coordination. In a typical macaque pair, both agents initially retained a strong preference for their own preferred colors, showed no strong side selection bias and little coordination (**Figure S3A,B**; see **Figure S2K-O** for data from all pairs). In the late session the behavior had changed, with both agents exhibiting less own color selection and a stronger convergence on the left side of the display (**Figure S3C,D**). This shift in strategy was gradual (**Figure S3E,F**) and resulted in a better outcome for both agents with nearly equalized reward (**Figure S3G**).

Supplementary Figure S4, related to Figure 4. Macaques can be trained to take turns when paired with a human confederate. Same as the **Figure 4**, for monkey C. **(A)** Early confederate-training session, **(B)** late confederate-training session. The reaction time difference histograms of the late session show that in blocks when the pair selected the macaque's non-preferred target (red curve), the monkey acted slower than the confederate, waiting to see the human's choice before committing to its own choice (t-test: coordination on red (mean -142 ms, SD 138, N 86) vs. blue (mean 0.5 ms, SD 119, N 108); t(168.4) -7.6, p < 2*10^{-12}). **(C)** Monkey C also ceased to take turns when an opaque barrier was placed over the region of the transparent display containing touch targets (gray background in C) and mainly selected his own target. The action time difference histograms show that in the transparent condition the behavior was similar to (B): the monkey waited for the human to act first in blocks of the monkey's non-preferred target (t-test: coordination on red (mean -192 ms, SD 180, N 98) vs. blue (mean 27 ms, SD 130, N 169); t(156.3) -10.6, p < 5*10^{-20}). In the opaque condition the behavior was similar to (A): the monkey mostly selected his preferred target and there was no AT difference dependence on human's choices (note the overlap between dashed blue and magenta curves).

Supplementary Figure S5, related to Figure 4. Monkeys gradually learn to follow a human confederate's turn-taking behavior. (A-F) The development of aggregate measures of choice behavior and coordination for monkey F (same conventions as **Figure S3E-J**). Note the gradual increase in average rewards, and the overall slow changes in the first 9 sessions as compared to the steeper increases in the MIs and the dynamic coordination reward (DCR) in the last 6 sessions. **(G-L)** Corresponding data for monkey C.

Supplementary Figure S6, related to Figure 5. Confederate trained macaques. (A-F) The development of aggregate measures of choice behavior and coordination for monkey pair FC after confederate training (same conventions as **Figure S3E-J**). Note sessions FC-2 and FC-5, which show a more equal FCO, and average reward between the two monkeys, as well as both non-zero mutual information for side ad target, and especially significant DCR values.

Supplementary Figure S7, related to Figure 6. Correlations between the probability to see the partner's choice p_{see} and the likelihood to select the partner's target p_{other}. The left column shows the correlations for different sessions of the same pair of macaques (FC) before (FC naïve), during and after the confederate training, as well as different human pairs. The middle and right columns show p_{see} and p_{other} for one exemplary session for each group. Note that training with a confederate increased magnitude of the correlation (Confederate training) and this carried into the post-confederate training sessions of the two macaques (FC trained). **Tables S2** and **S3** contain the statistical information for each individual correlation.

Supplementary Figure S8, related to Figure 7. Humans and macaques show different switching behavior. (A) A magnification of the middle part of the macaque FC-2 session shown in **Figure 5B** (trials 211 - 495); color bars represent the choice combinations; the magenta arrows point to uncoordinated switches that resulted in a change of the coordination color. **(B)** Four subsequent trials (n-2, n-1, n, n+1) illustrating 8 possible switch types occurring in trial *n*, classified as "selfish" (an agent initiates a transition to own preferred target) and "benevolent" (an agent initiates a transition to other's preferred target). Seamless switches involve uninterrupted change from one to another coordinated option; uncoordinated switches involve one or more uncoordinated trials. **(C-D)** A quantification of different transition aspects as scatter plots of benevolent choices on the y-axis and selfish choices on the x-axis, in the two species. **(C)** The switches from coordination to non-coordination, or "challenge initiations", for the faster agent. Monkey pairs fall below the diagonal (with the two competitive turn-taking sessions in pair FC even below the 95% confidence intervals around the diagonal). Turn-taking human pairs again cluster around the diagonal. **(D)** The behavior of the faster agent for all switches from non-coordination to coordination, or "challenge resolutions". Turn-taking humans cluster along the diagonal, while the trained monkey pair clustered below the diagonal (with the two competitive turn-taking sessions below 95% confidence interval around the diagonal).

Supplementary Figure S9, related to Figure 7. Action time changes around choice combination switches. The graphs show the action time transitions around 6 specific choice switches, for agent A (monkey F) in red and for agent B (monkey C) in blue. Each curve shows the mean action time for 3 trials before, at the actual switch trial and 3 trials after the switch, the shaded areas show 95% confidence intervals around the mean. R – red, B – blue, M – magenta. The choice transitions are R→M: from both agents selecting red to each selecting own color; B→M: from both selecting blue to each selecting own color; R→B and B→R: from both selecting one color to both selecting another color (seamless transition, rare in macaques, see **Figure 8A**); as well as the respective reverse transitions. The choices of agent B are indicated on the top, the choices of agent A on the bottom. Note that both macaques speed up when transitioning from coordinating on their non-preferred color to challenging (R→M for agent B, and B→M for agent A), and also when transitioning from challenging to coordinating on their preferred color (M→B for agent B, and M→R for agent A). Similarly, both macaques slow down when transitioning from coordination on their preferred color to challenging (B→M for agent B, and R→M for agent A) and when transitioning from challenging to coordination on their non-preferred color (M→R for agent B, and M→B for agent A). In rare seamless switches, monkeys speed up on coordination on the preferred color and slow down for transitions to the non-preferred color. Comparing action times in 3 trials before the transitions from coordinated choices on an agent's preferred color to uncoordinated selection of individually preferred colors (RRR→M for agent A, BBB→M for agent B) versus the action times in 3 trials after the respective reverse transitions from uncoordinated own color selection to coordination on an agent's preferred color (M→RRR for agent A, M→BBB for agent B) showed no significant differences (agent A: RRR→M: mean 526.7, SD 56.9; M→RRR: mean 527.8, SD 55.4, t(533.6) -0.23, p < 0.83; Agent B: BBB→M: mean 590.8, SD 128.1, M→BBB transitions: mean 573.7, SD 116.6, t(298.3) 1.23 p < 0.22). Note we excluded the actual transition trial. The absence of action time differences in the beginning and the end of coordination bouts indicates that transitions were not "invited" by a faster agent slowing down prior to the transition.

Supplementary Tables

Pair ID	N solo trials	Solo trials: fraction of choosing own, agent A [%]			Solo trials: fraction of choosing own, agent B [%]			Choice preference (own vs. other's) last 200 dyadic trials		AT A – AT B, coordinated trials			AT A – AT B, uncoordinated trials			Test on coordinated vs uncoordinated AT differences			Total reward [EUR]	
		all trials	26-50	last 25	all trials	26-50	last 25	agent A	agent B	mean	SD	N trials	mean	SD	N trials	df (corr.)	t	p	agent A	agent B
1	100	43	44	48	54	52	60	bal.	bal.	-110.4	90.8	306	-89.7	76.2	94	181.3	-2.19	**2.95E-02**	20	19.2
2	100	93	**100**	**100**	96	**100**	**100**	bal.	bal.	72.4	85.1	383	64.6	63.7	17	18.6	0.48	6.33E-01	22.8	24
3	100	86	**88**	**100**	55	60	52	bal.	bal.	130.5	116.1	263	60.2	114.6	90	156.0	5.01	**1.48E-06**	19.5	19
4	100	92	**100**	**100**	99	**100**	**100**	own	own	n/a	n/a	0	-21.3	52.3	400	n/a	n/a	n/a	14.9	15
5	100	98	**100**	**100**	95	**100**	**100**	own	bal.	-7.0	57.4	121	-30.5	52.3	279	210.3	3.87	**1.47E-04**	18.6	16.7
6	100	92	**100**	**100**	87	**92**	**92**	bal.	bal.	-7.3	77.9	396	104.3	70.0	4	3.1	-3.17	**4.88E-02**	23.8	23.7
7	100	26	32	4	50	64	48	bal.	bal.	-27.2	72.4	367	-32.8	96.3	33	35.3	0.33	7.46E-01	21.6	22.4
8	100	54	56	52	99	**100**	**100**	bal.	bal.	-177.6	80.3	377	-56.4	114.6	23	23.3	-5.00	**4.52E-05**	23.1	23
9	100	97	**100**	**100**	93	**100**	**100**	bal.	bal.	5.9	82.9	320	-5.1	77.3	80	128.3	1.13	2.61E-01	22	22
10	50	86	**96**	**96**	24	16	16	own	other's	-36.4	73.7	300	156.2	n/a	1	n/a	n/a	n/a	19.4	14.7
11	50	46	48	48	2	0	0	bal.	other's	-43.2	66.4	176	-33.2	55.5	124	289.4	-1.42	1.58E-01	13.6	10.5
12	50	34	64	64	61	64	64	bal.	bal.	-22.5	125.9	276	-23.2	97.0	24	30.2	0.03	9.77E-02	15.7	17.2
13	50	89	**92**	**92**	60	44	44	bal.	bal.	106.2	109.2	239	75.1	84.6	61	116.5	2.41	**1.77E-02**	15.8	15.1
14	50	94	**100**	**100**	50	40	40	own	other's	-40.3	53.8	240	-14.2	61.3	60	83.1	-3.03	**3.28E-03**	18.7	13.7
15	100	1	0	0	53	44	40	other's	bal.	55.5	70.2	166	15.3	67.0	235	345.0	5.76	**1.84E-08**	13.1	16.3
16	100	58	44	**80**	20	40	0	own	other's	0.2	66.5	388	-31.0	142.8	12	11.2	0.75	4.67E-01	26	19.6
17	100	94	**100**	**100**	94	**96**	**100**	bal.	bal.	114.5	85.4	337	153.5	87.1	63	85.8	-3.27	**1.53E-03**	22.4	22.5
18	100	96	**96**	**96**	31	48	24	other's	own	-96.0	108.8	398	9.5	65.6	2	1.0	-2.26	2.60E-01	21	25.9
19	100	93	**100**	**100**	1	4	0	own	own	n/a	n/a	0	22.2	90.8	407	n/a	n/a	n/a	15.1	13.7

Table S1. Human pairs. Columns 2-10: No relationship between own target preference in solo trials and dyadic behavior. 19 of 38 human subjects preferred their own target at the end of the solo training trials (own color selection >75% in 26-50 and last 25 trials is marked by bold font). Percentage of choice for the own preferred target for both agents of each human pair, for all solo trials, trials 26-50, and for the last 25 solo trials. Note that out of the 28 subjects trained with 100 solo trials only one (16A) did not reach 75% threshold in trials 26-50 trials but eventually reached it in trials 76-100. Pairs displaying dynamic coordination (turn-taking behavior) are designated by a gray background. Choice preference in dyadic trials: other's (FCO<25%), balanced (25%≤FCO≤75%), own (FCO>75%). Out of 10 pairs (20 subjects) that showed balanced turn-taking, 11 subjects showed a strong preference for own targets already in solo trials, and 9 did not. In the remaining 9 pairs (18 subjects), 7 subjects who had own preference in solo trials also showed it in dyadic trials, 4 subject who had other's preference in solo trials also showed it in dyadic trials, and 5 out remaining 7 subjects switched the preference from solo to dyadic trials, e.g. other's to own. Thus, the behavior in the solo trials is not predicting the behavior in the dyadic trials. This indicates that human participants did not simply carry their established preferences into the dyadic setting, but continued to explore and modify their strategies. **Columns 11-19. Action time differences and coordination. Nine of 19 human pairs showed significant differences between coordinated and uncoordinated trials' action time difference, indicating that seeing the other's action can help maintain coordination.** Comparison of trial-by-trial action time differences between the two agents in the pair (AT A – AT B), for coordinated and uncoordinated trials for all human pairs by t-test using Satterthwaite's approximation to allow for unequal variance. Note that pairs 4, 10, and 19 had too few trials for one of the trial types to calculate the test statistic (n/a). Pairs displaying turn-taking behavior are designated by a gray background. Bold font denotes correlations with a p<0.05. Green font shows pairs in which absolute AT differences increased with coordination, purple – pair in which absolute AT differences decreased with coordination. **Columns 20-21. The total reward earned by each agent in solo and dyadic trials**. Note that pairs 10-14 were offered only 350 (50 solo and 300 dyadic) trials instead of 500 (100 solo and 400 dyadic) trials, explaining the lower totals.

Session	df	r(A)	r(B)	p(A)	p(B)	raw r(A)	raw r(B)	raw p(A)	raw p(B)
Prior to confederate training (naïve pair)									
FC-1	304	-0.544	0.169	**5.12E-25**	**0.002952**	-0.068	0.043	0.23521	0.45327
FC-2	365	0.236	-0.536	**4.62E-06**	**1.17E-28**	0.388	-0.114	**1.27E-14**	**0.028859**
FC-3	183	-0.351	0.200	**9.42E-07**	**0.006239**	-0.124	-0.062	0.092404	0.40478
FC-4	207	-0.376	-0.200	**2.00E-08**	**0.003778**	-0.128	0.218	0.064765	**0.001527**
FC-5	171	0.194	0.161	**0.010527**	**0.034044**	0.111	-0.014	0.14464	0.85860
FC-6	397	n/a	-0.096	n/a	0.054399	n/a	-0.039	n/a	0.44066
FC-7	314	-0.044	0.560	0.433	**1.68E-27**	0.024	0.144	0.67187	**0.010417**
FC-8	447	0.447	0.228	**1.67E-23**	**9.91E-07**	0.171	0.027	**0.00027874**	0.5692
During confederate training									
F-1	431		0.063		0.19125		0.079		0.10121
F-2	714		0.125		**0.000804**		0.108		**0.003798**
F-3	578		-0.053		0.20033		0.140		**0.000699**
F-4	80		0.040		0.72184		0.248		**0.024749**
F-5	547		0.044		0.30683		0.173		**4.56E-05**
F-6	317		0.280		**3.54E-07**		0.205		**0.000225**
F-7	540		0.141		**0.001023**		0.170		**6.82E-05**
F-8	582		0.077		0.063123		0.076		0.068242
F-9	547		0.260		**5.64E-10**		0.298		**9.42E-13**
F-10	479		0.544		**2.19E-38**		0.390		**6.99E-19**
F-11	473		0.820		**7.89E-117**		0.586		**3.44E-45**
F-12	525		0.829		**1.15E-134**		0.579		**1.40E-48**
F-13	594		0.815		**3.75E-143**		0.589		**7.59E-57**
F-14	554		0.867		**2.42E-169**		0.557		**1.10E-46**
F-15	446		0.868		**1.40E-137**		0.642		**2.38E-53**
C-1	652		0.161		**3.51E-05**		0.188		**1.37E-06**
C-2	544		0.311		**9.50E-14**		0.172		**5.18E-05**
C-3	320		0.482		**3.56E-20**		0.307		**1.78E-08**
C-4	455		0.145		**0.001896**		0.188		**5.03E-05**
C-5	412		0.588		**7.67E-40**		0.285		**3.63E-09**
C-6	476		0.697		**7.88E-71**		0.408		**1.46E-20**
C-7	581		0.670		**3.11E-77**		0.388		**2.42E-22**
C-8	228		0.715		**2.36E-37**		0.471		**4.42E-14**
After confederate training (trained pair)									
FC-1	371	0.806	0.834	**2.45E-86**	**1.24E-97**	0.516	0.603	**8.13E-27**	**3.13E-38**
FC-2	545	0.916	0.909	**3.57E-218**	**1.28E-209**	0.635	0.655	**4.15E-63**	**2.93E-68**
FC-3	625	0.753	0.518	**6.88E-116**	**2.65E-44**	0.423	0.321	**1.19E-28**	**1.76E-16**
FC-4	471	0.856	0.517	**9.73E-137**	**9.49E-34**	0.420	0.305	**1.20E-21**	**1.22E-11**
FC-5	683	0.707	0.654	**1.16E-104**	**9.32E-85**	0.562	0.430	**2.71E-58**	**3.53E-32**
FC-6	370	0.901	0.822	**3.33E-136**	**1.29E-92**	0.601	0.580	**7.55E-38**	**7.34E-35**

Table S2. Action time correlations in macaque pair FC. Correlation between the probability to see other's action and the selection of the other's target. Prior to confederate training (8 naïve sessions), monkeys F (agent A) and C (agent B) showed only relatively weak and inconsistent (both positive and negative) correlations between probability to see other's action (r(A) and r(B)) and the selection of the other's target. Correlation degree of freedom (df), correlation coefficients (r) and correlation p-values (p) for the running average over 8 trials as displayed in **Figure S7**, and for the raw non-smoothed data are shown. Bold font denotes correlations with p<0.05. n/a – not applicable: the correlation is undefined because at least one variable was constant (e.g. the fraction of choosing other's 0). The few high correlations on smoothed data were mainly driven by the slow changes in the timecourse of the session. During confederate training both monkeys (F, C) developed strong and highly significant positive correlations between being able to see the confederate's actions and following the confederate: 15 sessions of monkey F and 8 sessions of monkey C during confederate training. Note the human confederate worked on position A, and is excluded here. Bold font denotes correlations with a p<0.05. After confederate training this pair continued to show strong correlations between seeing the other's choice and following: 6 sessions of monkeys F and C after confederate training. Note that even in the sessions without significant DCR in the last 200 trials (i.e. FC-1, FC-3, FC-4, FC-6), where monkeys largely converged on one color, the high positive correlations reflect brief bouts of competitive turn-taking or challenging.

Pair ID	df	r(A)	r(B)	p(A)	p(B)	raw r(A)	raw r(B)	raw p(A)	raw p(B)
4	398	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
19	405	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
15	399	-0.123	-0.110	**0.014003**	**0.026945**	0.067	0.176	0.1802	**0.000399**
11	298	-0.097	n/a	0.09230	n/a	0.040	n/a	0.48759	n/a
5	398	-0.009	-0.427	0.85640	**3.93E-19**	-0.028	-0.171	0.58233	**0.000598**
3	351	0.164	0.006	**0.001941**	0.91763	-0.012	0.054	0.82342	0.30902
1	398	-0.009	0.025	0.85616	0.62476	0.027	0.026	0.59697	0.60243
14	298	0.060	0.079	0.29914	0.17498	0.010	0.154	0.86265	**0.007371**
13	298	0.126	0.108	**0.029495**	0.061976	0.026	-0.037	0.65319	0.52015
2	398	0.243	0.394	**9.12E-07**	**2.54E-16**	0.095	0.124	0.05769	**0.012791**
12	298	0.523	0.456	**1.98E-22**	**8.08E-17**	0.268	0.330	**2.41E-06**	**4.90E-09**
16	398	0.046	-0.569	0.36308	**1.18E-35**	0.142	-0.214	**0.004470**	**1.64E-05**
7	398	0.053	0.287	0.29108	**4.94E-09**	0.001	0.066	0.97791	0.18541
9	398	0.126	-0.271	**0.011469**	**3.59E-08**	0.009	-0.025	0.85404	0.61258
17	398	-0.487	0.187	**3.10E-25**	**0.000171**	-0.054	0.038	0.28453	0.44647
8	398	-0.114	0.075	**0.022805**	0.13386	-0.036	0.055	0.47255	0.2706
6	398	-0.370	0.071	**2.04E-14**	0.15713	0.040	0.017	0.41953	0.73857
10	299	0.251	n/a	**1.07E-05**	n/a	0.221	n/a	**0.000109**	n/a
18	398	0.048	0.176	0.34150	**0.000399**	-0.000	0.094	0.99829	0.06041

Table S3. Action time correlations in humans. Correlation between the probability to see other's action and the selection of the other's target in 19 human pairs. Human pairs showed weak and inconsistent (both positive and negative) correlations between probability of seeing the other's choice and selecting the other's target. Correlation degree of freedom, correlation coefficients and correlation p-values for the running average over 8 trials, as displayed in **Figure S7**, and for the raw non-smoothed data are shown. Letters (A) and (B) denote the positions of the agents. Pairs are sorted by the average joint reward in ascending order (cf. **Figure S2B**). Pairs displaying dynamic turn-taking behavior are designated by a gray background. Bold font denotes correlations with a p<0.05. n/a – not applicable: the correlation is undefined because at least one variable was constant (e.g. the fraction of choosing other's 0).