Human and macaque pairs employ different coordination strategies in a dyadic decision game with face-to-face action visibility

Sebastian Möller[1,2,3], Anton M. Unakafov[1,2,3,4,5], Julia Fischer[3,6,7], Alexander Gail[1,2,3,8,†], Stefan Treue[1,2,3,8,†], Igor Kagan[1,3, *,†]

[1]Cognitive Neurosciences Laboratory, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077, Goettingen, Germany

[2]Georg-Elias-Müller-Institute of Psychology, University of Goettingen, Gosslerstrasse 14, 37073, Germany

[3]Leibniz ScienceCampus Primate Cognition, Kellnerweg 4, 37077, Goettingen, Germany

[4]Max Planck Institute for Dynamics and Self-Organization, Am Fassberg 17, 37077, Goettingen, Germany

[5]Campus Institute for Dynamics of Biological Networks, Hermann-Rein-Straße 3 37075, Goettingen, Germany

[6]Cognitive Ethology Laboratory, German Primate Center – Leibniz Institute for Primate Research, Kellnerweg 4, 37077, Goettingen, Germany

[7]Department of Primate Cognition, Johann-Friedrich-Blumenbach Institute for Zoology and Anthropology, University of Goettingen, Germany

[8]Bernstein Center for Computational Neuroscience, Am Fassberg 17, 37077, Goettingen, Germany

[*]Corresponding author: ikagan@dpz.eu

[†] Shared senior authorship

Short title: Decision coordination in humans and monkeys

Abstract

Interactions of group-living primates range from cooperation to competition. Many corresponding real-world decisions critically depend on observing partner's actions. In classical game theory studies, however, agents act without knowing the partner's choices. To overcome these limitations, we developed a setup where two agents engage face-to-face in cooperative or competitive tasks on a shared transparent touchscreen display ("transparent games"), ensuring mutual action visibility. We compared human and macaque pairs in a transparent version of the coordination game "Bach-or-Stravinsky", which entails the conflict about which of the two individually-preferred coordinated options to choose. Most human pairs developed coordinated behavior and adopted dynamic turn-taking to equalize the payoffs. All macaque pairs also converged on coordination, but in simpler, static ways. Remarkably, two animals that underwent training with a human confederate learned to coordinate dynamically, exhibiting turn-taking driven by temporal competition. The observed choices were captured by modeling the probability to see the other's action before one's own movement. Such competitive turn-taking is unlike the prosocial turn-taking in humans, who equally often initiated switches to and from their preferred option. Our results show that dynamic coordination is not restricted to humans, but it can be subserved by different social attitudes, and emphasize the importance of action visibility in studies of the emergence and maintenance of coordinated behavior.

Teaser

In real-time transparent interactions, both species show dynamic coordination, but macaques compete while humans strive for fairness by taking turns.

Introduction

The majority of primate species lives in complex social groups, in which interactions range from intense competition to cooperation (*1–5*). To adjust behavior optimally, individuals need to assess their own actions and goals, and those of other group members, while taking into account the history of interactions with and between these individuals (*4*, *6*, *7*). Coordination is essential for maintaining cohesion between group members, avoiding conflicts, and achieving individual and joint action goals. Here we focus on understanding how such coordination can be achieved and maintained in a dyadic "transparent" setting in which both agents can observe each other's ongoing actions and social cues (*8*, *9*), like in many natural situations.

Game theory, developed to study strategic interactions in rational decision-makers (*10*), offers a powerful framework to investigate dyadic social interactions (*11–13*) and the evolution of coordination (*14–16*). One fundamental class of games are 2×2 games (*17*, *18*) in which each agent chooses one of two actions and the outcome depends on the combination of their choices, as defined by the payoff matrices describing the gain or loss the two agents earn for the four possible combinations. Such choices can be presented as one-shot games, or as iterated games with repeated interactions that encourage tracking the interaction history, as well as the formulation of predictions regarding the other agent's decisions (*19*).

For understanding many realistic scenarios, non-zero-sum games are a particularly interesting class of 2×2 games. In non-zero-sum games, the agents need to coordinate their choices to maximize individual and/or joint reward (*20*). As opposed to the strictly competitive zero-sum games, non-zero-sum games have both cooperative and competitive elements, with aligned and opposing interests. For example, in the Stag Hunt (or Assurance) game, agents choose between maximizing the individual as well as joint reward and minimizing the individual risk (*21*). Humans *(Homo sapiens)*, chimpanzees *(Pan troglodytes)*, rhesus macaques *(Macaca mulatta)* and capuchin monkeys *(Cebus apella)* coordinate at above chance level on mutually beneficial high-reward / high-risk choices in the iterated Stag Hunt (*22–25*). Similar to Stag Hunt, in Prisoner's Dilemma (PD) two agents maximize their *joint* reward if they both choose to cooperate. Yet, an individual agent obtains the highest reward when defecting while the partner cooperates (and receives the lowest reward), creating a dilemma between the self-interested temptation to defect and a riskier cooperation. For cooperation to emerge in PD, agents need to "trust" the partner, i.e., they need to expect the partner to cooperate for the mutual benefit. Humans often show a bias towards cooperative behavior (*26–28*). Remarkably, it was

demonstrated that macaques, while mostly selecting defection, chose mutual cooperation significantly more and mutual defection significantly less than expected by chance, especially after preceding cooperation, in an iterated PD game (*29*). This suggests that macaques can at least partially overcome the selfish motives and reciprocate cooperation.

Another type of conflict that contrasts selfish and prosocial preferences is implemented in the Conflict game (also known as Hawk-Dove, or game of Chicken) (*16, 30*). These anti-coordination games model a competition over a shared resource that can be monopolized by one agent, while an actual clash is highly detrimental to both. When both agents exhibit cooperative behavior (e.g., yield), their joint reward is less or equal to the joint reward of the two anti-coordinated choices (e.g., fight-yield and yield-fight) maximizing the individual reward of one agent at the expense of the other. Humans, capuchins and macaques often converged on anti-coordination in this game (*30*), but importantly, only humans alternated between the two individually-optimal anti-coordinated choices (*30–32*). Such alternating behavior results in the maximal and equal payoff for both agents and is called "cooperative turn-taking" (*33–35*).

In the above games, reward-optimizing rational agents should coordinate on one option (Stag Hunt and Prisoner's Dilemma) or anti-coordinate on opposite options (Conflict game). A third, less studied type of games that emphasizes the coordination on *either of two options* is known as Battle of the Sexes, or Bach-or-Stravinsky game (BoS) (*36*). Each agent has an individually preferred option, but coordinating on either one of these options adds the same bonus to both agents. This renders any coordinated choice better than no coordination for both agents, but one coordinated choice better for the first agent and the other better for the second agent. The rational choice is to coordinate, but unlike the conceptually simpler optimal convergence on one option in Stag Hunt and PD, BoS includes an inherent conflict about who profits the most. This combination of cooperation and conflict offers an interesting opportunity for studying social interactions that has not been sufficiently explored in nonhuman primates. As in the Conflict game, games of the BoS type revealed that humans often take turns, switching between the two coordinated options to ensure fairness (*37–39*). Interestingly, cooperative turn-taking frequently took place in 5-year-old children, but neither in 3-year-old children nor in chimpanzees (*38*). This begs the question whether turn-taking requires special social or cognitive abilities that are unique to humans. We therefore compared the behavior of (adult) humans and rhesus macaques in the BoS economic game.

Traditionally, economic games are played either *"simultaneously"* (neither agent knows the choice of the other before making its own decision) or *sequentially* in a predetermined order; yet real dyadic interactions often play out in a real-time with the partner's actions in direct sight (*40*, *41*). Thus, the timing of one's own and other's actions becomes part of the strategy space (*42*). Such a "transparent" continuous time setting can change choice strategies as compared to the classic simultaneous and sequential settings (*33*). Especially for nonhuman species, coordinating based on mutual choice history might be more demanding than coordinating based on the immediately observable behavior of others. For example, visual feedback about the partner's choices improves coordination in the iterated Stag Hunt game in humans, capuchins and rhesus macaques (*25*), and such coordination in chimpanzees is facilitated if one of the agents consistently acts faster than the partner (*23*). Similarly, there were substantial differences in capuchins' and rhesus' behavior in the Conflict / Chicken game when they had access to the current choice of a partner (*30*, *43*). In humans, a real-time anti-coordination game revealed that action visibility (and the possibility to change an already initiated action) increased efficiency and fairness (*44*). Besides those few pioneering studies, the experimental and theoretical effects of action visibility on (anti)coordination have not been systematically addressed.

To account for changes in strategies during action visibility compared to simultaneous ("opaque") discrete choices, we recently developed the concept of "transparent games" that extends classical evolutionary game-theoretic analysis to real-time interactions in which the visibility of partners' actions depends on their relative reaction times (*8*, *9*). Here, we use the theoretical insights from that work and analyze short- and long-term dynamics of choices, mutual information and reaction times in humans and rhesus monkeys playing a transparent version of the iterated movement-based BoS game. Since many direct interactions between primates occur face-to-face (*45–49*), we designed a novel dyadic interaction platform that allows two human or monkey agents to observe each other and to act on the same visual objects on a vertical touch-sensitive transparent display between them. The advantage of this configuration is that agents can monitor and react to each other's actions in real-time, emulating naturalistic interactions while still maintaining well-controlled laboratory conditions. Extending beyond static choice proportions, our approach captures dynamics of such interactions, affording a mechanistic model of trial-to-trial choice behavior driven by within-trial temporal patterns.

To facilitate inter-species comparability, we paired human or macaque agents without explicit task instructions. We expected that the transparency and instantaneous coaction (*41*) would

facilitate efficient coordination. Based on the human propensity for fairness (*50*) and perspective taking (*51, 52*), we predicted that humans would engage in pro-social turn-taking where the reward maximization is alternated between partners. We predicted that macaques would also utilize the information about the partner's actions, but would be less likely to grant an advantage to the other player. Instead, we expected that the monkeys would coordinate to increase individual (and joint) rewards, but without turn-taking to balance rewards between players.

Results

In our experiments, two human or macaque agents were sitting face-to-face and shared a transparent vertical workspace between them (**Figure 1A,B**). Agents could simultaneously see the stimuli and each other's actions. In the transparent variant of the BoS paradigm we used, each agent could choose one of two simultaneously presented color targets (red and blue). The left/right position of each color was randomized across trials (50% left, 50% right). The resulting rewards followed the payoff matrix shown in **Figure 1C** and depended on the choices of both agents. Prior to the first dyadic session, each agent in a pair was individually trained to associate one of the two color targets to a larger reward, so that the agents preferred different colors. In the dyadic trials, selecting the same target resulted in an additional reward on top of the individually trained values - hence choosing the same target was always better than choosing different targets. But in such coordinated trials, the agent whose individually preferred ("own") color was selected received a larger reward. This paradigm probes the ability to coordinate in order to obtain higher rewards, and the capacity to deal with the unfairness introduced in each coordinated trial.

Human pairs mainly converge on fair coordination

We recruited 38 naïve human subjects to assess performance in the transparent BoS paradigm. Each subject was trained individually for 50-100 solo trials to operate the touch panel and to associate a higher auditory pulse count and the corresponding monetary reward with one of the two color targets (**Table S1**). We then paired pre-trained agents and let them explore the full payoff matrix. Subjects were instructed not to talk during the experiments (see Supplementary Materials for instructions given to the subjects). For the aggregate measures of performance, we analyzed the last 200 dyadic trials of each session, to allow for convergence on a strategy (Materials and Methods).

Figure 1. Experimental setup. (A) A sketch and **(B)** a photo of the dyadic interaction platform showing two macaques reaching to targets in a shared workspace. **(C)** The payoff matrix describing the reward outcome for each player for four possible combinations of individual choices (numbers show the monetary/liquid reward units for agent A in red and agent B in blue). The coordinated selection of the same target resulted in a "bonus" of 2 reward units. **(D)** Schematic top-view of two macaque agents in different stages of the task, and the timing of one trial. **(E,F)** Performance of a naïve human pair. **(E)** Example session showing for both agents (A: red; B: blue) the share of own choices (SOC) as running average over 8 trials. In coordinated choices one agent always selects the own target (SOC = 1), while the other selects the non-preferred target (SOC = 0). The dashed lines show the session average SOC for each agent. The color bar below shows for each trial the selected combination (red: A and B selected A's preferred target; blue: A and B selected B's preferred target; magenta: both agents selected their individually preferred target; green: both agent selected a non-preferred target). Here both human agents coordinated by selecting the same target in blocks. **(F)** The share of objective left choices (SLC, from A's viewpoint) as running average over 8 trials. The dashed lines show the session average SLC for each agent. Here side selection fluctuated around random level 0.5 for both players.

Human strategies ranged from strict adherence to the pre-trained higher value colors to joint selection of the same target with alternating between the two colors trial-by-trial or in blocks. **Figure 1E,F** illustrates an exemplary human pair that developed such alternating turn-taking behavior, showing the *share of own choices* (SOC) - the likelihood for each agent to select the own preferred color, and the *share of left choices* (SLC) - the likelihood for each agent to select

the target on the left side. These measures provide direct insight into each agent's target and side biases. Here the SOC curves show that this pair mostly jointly selected A's or B's color. After an initial period of long blocks of trials, the pair switched to shorter blocks. The SLC curves show that the agents did not display a side bias.

The SOC of all 19 pairs (**Figure 2A**) shows 53% (10 of 19) with balanced values close to 0.5 for both agents (indicating alternating between colors), and the remaining 47% (9 of 19) with at least one agent close to one or close to zero (indicating fixed color selection). **Figure 2B** shows the average reward (AR) for each agent, which in most pairs was above chance level (2.5 units), and in about half the pairs close to the optimum for the dyad (3.5).

From the sequence of choices and obtained rewards we derived three measures of coordination (Materials and Methods). The *dynamic coordination reward* (DCR) describes the average reward a pair earned *above (or below)* the reward expected for independent choices, given the observed choice frequencies per agent. Zero DCR indicates coordination by chance, values significantly different from zero indicate above chance levels of coordination. The DCR values (**Figure 2C**) and the fairly balanced average rewards (**Figure 2B**) suggest that 10 of 19 human pairs converged on a form of turn-taking, or reciprocal coordination. These are the same 10 pairs that showed roughly balanced SOC values around 0.5 (**Figure 2A**). To better understand what drives high dynamic coordination seen in **Figure 2C**, we calculated the *mutual information* (MI) between the sequences of target color choice (MI target, **Figure 2D**) and the side choice (MI side, **Figure 2E**) of both agents. These measures indicate how well one agent's choice of color or side can be predicted from the other's choice (Materials and Methods). Comparing the DCR and the MI measures revealed that high DCR values coincided with non-zero values of both MI side and MI target, indicating that dynamic coordination corresponds to both correlated side and color choices between the agents.

The nine human pairs who did not show significant dynamic coordination used strategies other than turn-taking. Five pairs converged on a fixed color favoring one agent (5, 14, 16, 10, 18), indicating that at least one of the two agents understood the value of coordination. Four pairs did not coordinate successfully – in two pairs both agents selected only the own preferred color (4, 19); and in two pairs one agent selected the non-preferred color while the other switched between the two colors (11) or selected the right side (15). To summarize, about half of the human pairs exhibited dynamic coordination and aimed for fairness.

Figure 2. Performance of naïve human (A-E) and macaque pairs (F-J). Human pairs: Aggregate measures of choice behavior and coordination for 19 naïve human pairs. panels B and C are sorted by the average joint reward (not shown); the example in Figure 1E,F is from pair 2. Individual measures for A are shown in red, for B in blue, measures for the pair are in purple. **(A)** The maximal of the two average SOC values in each pair, versus the minimal of the two. The diagonal line indicates equal frequency of own choices for both agents. **(B)** The average reward for each agent. The dotted line at 2.5 shows the reward value for random choices, the line at 3.5 shows the maximal attainable average joint reward. **(C)** The dynamic coordination reward (DCR), the error bars show 95% confidence intervals; 53% of pairs (3, 1, 13, 2, 12, 7, 9, 17, 8, and 6) showed significant DCR. **(D)** Mutual information for target choices. **(E)** Mutual information for side choices. **Macaque pairs:** Panels **(F-J)** show the same measures as A-E for a late session for each pair. Macaque pairs converged on jointly selecting either the same side or on jointly selecting one of the two color targets.

Macaque pairs converge to simpler coordination strategies

Six macaque monkeys were individually trained to internalize the preferred reward values of the two color targets. All six macaques developed a strong preference for the target associated with the large reward (range 78-100%, mean 95% large reward color selection in the last solo session), before being paired with a conspecific with the opposite target color preference. Generally, macaques changed their choice behavior more slowly than human participants, who often exhibited clear changes within a single session. We therefore collected the data from macaque pairs for multiple sessions to assess how their behavior evolved over time and on which strategy they converged. In a typical pair, both agents initially retained a strong preference for their own preferred colors (reward size 2 drops), and no strong side selection bias (**Figure S1A,B**). In the later session the behavior had changed, with both agents exhibiting less own color selection and a stronger convergence on the left side of the display (**Figure**

S1C,D). This shift in strategy was gradual (**Figure S1E,F**) and resulted in a better outcome for both agents with nearly equalized reward (**Figure S1G**).

A similar picture emerged in all nine pairs we tested (**Figure 2F-J** and **S2**). Typically, one agent changed either to selecting the non-preferred color (increasing his reward to 3 and the other's to 4 drops) or to selecting one side (increasing his reward to an average of 2.5 and the other's to 3 drops). If the other monkey also started favoring the same side, this increased both agents' reward to 3.5. This sequence of steps illustrates how macaque pairs developed their strategy over the course of several sessions. Comparing the average rewards in early and late sessions shows that most pairs reached better coordination over time (**Figure S2G**; median early 2.75 versus late 3.29; $p < 0.0039$, two-sided exact Wilcoxon signed rank test). In 8 of 9 pairs, the proportion of coordinated trials increased significantly ($p < 0.05$, Fisher's exact test on the number of coordinated versus uncoordinated trials in the early and the late session, calculated separately for each pair). This increase in coordination was achieved either by converging on the same color target or by converging on the same side, as evident in high values of *either* target *or* side MI (**Figure 2I,J**). Importantly, only high values of either MI target or MI side, and low and with one exception non-significant DCR values (**Figure 2H**) indicate that the resulting coordination was not achieved by dynamic turn-taking. Instead, the four pairs in the middle of **Figure 2F** that had roughly equal SOC and appear to be similar to the human turn-takers achieved the coordination by consistently selecting the same side regardless of the target color, which was randomized by the task (Materials and Methods). This side coordination can be seen from their high MI target and low MI side values, and high SLC values (**Figure S2F**).

Comparison of behavior between species

To directly compare the strategies of naïve human and macaque pairs, we plotted the mutual information values for side and target color choices of each pair, for the last 200 trials (**Figure 3**). The distance of each pair's location to the origin denotes the strength of (anti)coordination. Locations close to the x-axis denote static side strategies; locations close to the y-axis denote static target color strategies. Ten of 19 human pairs were close to the main diagonal, corresponding to similar values of side and target MI. These locations denote trial-by-trial or block-wise turn-taking – a signature of dynamic coordination. In contrast to humans, nine macaque pairs showed simple static side-based (4 pairs) or single target color-based (5 pairs) coordination, with different levels of coordination strength, but did not employ dynamic coordination.

Figure 3. Comparison of coordination strategies in humans and macaques. Each symbol shows the mutual information between side choices (MI side) and target color choices (MI target) calculated over the last 200 trials of a session, for human and macaque pairs. Since mutual information for a non-varying sequence of choices is zero, sessions in which a pair statically converged on a fixed color have MI target around zero and a non-zero MI side, and sessions in which a pair statically converged on a fixed side have a non-zero MI target and around zero MI side. Pairs/sessions near the y-axis hence indicate static selection of one of the two target colors (i.e., either coordinating on A's or B's preferred option, or only selecting the own color), while pairs/sessions near the x-axis indicate static coordination on a side. Pairs/sessions with balanced side and color choices fall onto the main diagonal, indicating dynamic coordination or turn-taking. The distance between each symbol and the (0,0) origin represents the (anti)coordination strength. Ten human pairs showed dynamic coordination, using trial-by-trial or block-wise turn-taking. Eight human pairs converged on color-based strategies: five pairs largely selected one color, i.e., one of the two coordinated options (5, 14, 16, 10, 18), two pairs did not coordinate (4, 19), and in one pair (11) one agent selected other's color but the other randomly switched between the colors. In one pair, both MI side and MI target did not significantly differ from zero (p > 0.01; pair 15, in this pair one agent selected one side and the other - one color). In late sessions, naïve macaque pairs either converged on selecting the same color target (pairs FC, LE, TC, TF, CL) or converged on the same side (TE, CE, MC, MF). In addition to naïve humans and macaque pairs, diamonds show 6 sessions of the confederate-trained macaque pair (F-C), see subsequent sections.

Macaques paired with a human confederate learn to follow

The absence of trial-by-trial coordination in macaque pairs suggests that they did not monitor their partners in order to take their choices into account on a trial-by-trial basis. This might be a consequence of the animals' training history (starting with extended individual solo training to associate one color with higher reward) or a general disinclination to consider the partner's

choice when making immediate decisions, at least when a simpler strategy is available. To test the latter, we trained two monkeys in the dyadic condition with a human confederate who followed a strict pattern of alternating between the two colors in blocks of approximately 20 trials ("confederate training"). If the rhesus monkeys were insensitive to partner's choices in our dyadic task, the pattern of confederate's choices should not affect monkey's behavior.

Both monkeys started with a strong bias for selecting their own preferred color but over time changed to a reliable coordination (cf. **Figure 4A** with **4B** and **S3A** with **S3B**). Their reaction times (**Figure 4** and **S3**, right columns) were initially faster than those of the confederate, and did not vary strongly depending on the confederate's actions. In later sessions the animals reacted considerably slower in those blocks in which the confederate was selecting the animals' non-preferred color, waiting for the confederate to commit to an action. We therefore labelled this as "following" behavior.

The aggregate measures of choice behavior and coordination development over multiple confederate-paired sessions showed similarities in monkeys' learning (**Figure S4**). Both animals reached close to 0.8 DCR units, similar to the values reached by humans employing turn-taking (cf. **Figure 2C**). These results indicate that macaques can take information from their partner into consideration when making decisions in the BoS paradigm. Reaction time dynamics suggested that the macaques started to actively monitor the confederate's actions and only selected their non-preferred color if the partner started to reach there first.

Due to the repeating block strategy employed by the human confederate it is not immediately clear whether the monkeys made use of the action visibility and based their decision on the action observed within a trial, or rather on the reward history. We therefore performed a control experiment in which we placed an opaque barrier on the confederate's side of the display for roughly the middle third of a session. The barrier blocked the monkey's view of the confederate hand, while keeping the face visible. In both monkeys, the "following" behavior ceased and they reverted to selecting their own preferred colors irrespective of the confederate's choices, which – by design – continued to follow the block strategy (**Figure 4C, S3C**). In line with this observation, there was a significant difference in reaction time when selecting monkey's preferred (blue) and the non-preferred (red) color in the transparent condition, but no difference in the opaque condition. In other words, monkeys failed to coordinate when they could not see the confederate's actions. These results imply that the immediate visual information about

partner's actions and not merely monitoring reward history drove and maintained the macaques' "following" behavior in transparent settings.

Figure 4. Macaques can be trained to take turns if paired with a human confederate. Pairing a macaque with a human partner who alternated between the two colors in pre-defined blocks leads to turn-taking. **(A)** In an early confederate training session, the monkey F (agent B) mostly selected his own target, independent of the human's choices. The lower sub-panel shows the running average (8 trials) of the average joint reward over both agents; the cyclic modulation shows that the macaque did not coordinate with the confederate. The reaction time (RT) difference histograms on the right (bin size 50 ms) show that in all four combinations of choices the macaque acted faster than the confederate. **(B)** A late confederate training session. After several sessions (Figure S4) the macaque had learned to coordinate (note anti-correlation of the two SOC curves and the average joint reward approaching maximum 3.5 value). The RT difference histograms show that in blocks when the pair selected the macaque's non-preferred target (red curve), the monkey acted slower than the confederate, waiting to see the human's choice before committing to its own choice (t-test: coordination on red (mean: -259 ms, SD: 113, N: 193) vs. blue (mean: -7 ms, SD: 153, N: 217), t(395.5): -19.1, p<0.001). **(C)** Macaque turn-taking with human confederate depended on direct visibility of the confederate's actions. The gray background denotes the period in which an opaque barrier was placed over the region of the transparent display encompassing all touch targets; hence, the agents could not see each other's reach movements but could still see the other's face. Note how the macaque F switched from turn-taking to mainly selecting own preferred target. Upward histograms show the transparent condition (solid curves) and downward histograms show the opaque condition (dashed curves). In the transparent condition the behavior was similar to (B): the monkey waited for the human to act first in blocks of the monkey's non-preferred target (t-test: coordination on red (mean: -248 ms, SD: 101, N: 123) vs. blue (mean: -39 ms, SD: 124, N: 166), t(284.5): -15.9, p<0.001). In the opaque condition the behavior was similar to (A): the monkey mostly selected his preferred target and there was no RT difference dependence on human's choices (note the overlap between dashed blue and magenta curves).

Confederate-trained macaque pair shows dynamic coordination

After two macaques had learned to monitor and follow a partner's actions, we tested how these animals would perform when paired with each other again. At the end of the initial naïve sessions of this pair, and before the training with a human confederate, monkey F (agent A) mostly selected his own preferred color while monkey C (agent B) either selected the left side or monkey F's preferred color. At the end of the confederate training, both monkeys followed the confederate. The question was: would they now express turn-taking behavior or would they fall back to their initial static color-coordination behavior?

In the first session, after some initial back and forth, the pair converged on mostly selecting B's color, with B acting faster (**Figure 5A**). The most interesting phase was during the second session, in which the animals repeatedly alternated between the two colors in long blocks (**Figure 5B**). In the third session, the pattern was opposite to the first session, with the pair converging mostly on A's color who acted faster (**Figure 5C**). This time, B tried occasionally to switch to his preferred color, and A followed for three short periods.

If action visibility is driving this form of dynamic coordination, then the question of who successfully "insists" on own preference and who "accommodates" the other's preference should depend on the relative reaction times of the two agents. Indeed, in the session in which the animals exhibited apparent turn-taking behavior (**Figure 5B**), the faster animal (Leader) selected the preferred color, while the slower animal (Follower) had to accommodate. Thus, the dynamic coordination arose within a competitive interaction. The mutual information measures as well as the DCR values indicate that after confederate training, this pair exhibited robust dynamic coordination resulting in turn-taking behavior in two sessions (**Figure S5**).

Figure 5. Confederate-trained macaques show competitive turn-taking behavior. (A) In the first paired session after confederate training indications of turn-taking followed by a convergence on B's target can be seen in the SOC in the top subpanel; note how agent A occasionally tried selecting his preferred target. The reaction time difference histograms show that agent B acted faster when both agents selected his preferred target (t-test: coordination on red (mean: -211 ms, SD: 112, N: 27) vs. blue (mean: 105 ms, SD: 98, N: 286), t(29.9): -14.1, p<0.001). **(B)** In the next session the pair developed turn-taking; the RT difference histograms show that the pair generally converged on the faster monkey's target (t-test: coordination on red (mean: -117 ms, SD: 96, N: 238) vs. blue (mean: 90 ms, SD: 104, N: 201), t(411.7): -21.4, p<0.001). In the following session **(C)** the pair mostly converged on A's preferred target, with A now acting faster (t-test: coordination on red (mean: -188 ms, SD: 115, N: 553) vs. blue (mean: -31 ms, SD: 192, N: 33), t(33.4): -4.6, p<0.001); note that as in the first session the agent B occasionally tried his own preferred target, and two of these attempts were reciprocated by A. **(D,E)** Confederate-trained macaque turn-taking is driven by action visibility. **(D)** The 2nd session of confederate-trained macaque pair FC (also shown in B). The inset (top right) shows

the logistic mapping function from reaction time difference to the probability to see the partner's choice before acting. This panel shows the correlation between probability of seeing the other move first (p_{see}) and following to the other's preferred target (p_{other}). The "share of other's choices" is the inverse of the share of own choices (SOC). All curves show running averages of the respective measure over 8 trials. For agent A the correlation coefficient r(545) was 0.92 and for agent B 0.91 for smoothed data, and 0.64 and 0.65 for non-smoothed (trial-by-trial) data (all p<0.001). (**E**) The human pair 12 with the highest positive correlation showed a different pattern: a slow gradual shift from B following A to B leading A over the course of the session. For A the correlation coefficient r(298) was 0.52 and for B 0.46 for smoothed data, and 0.27 and 0.33 for non-smoothed data (all p<0.001). The same human pair (only one out of ten that exhibited dynamic coordination) showed a significant RT difference for coordinated choices favoring A vs. B, with each agent being faster when selecting the own preferred color (coordination on A (mean: -88, SD: 129, N: 96), coordination on B (mean: 12, SD: 109, N: 180), t(168.8): -6.5, p<0.001, t-test).

Comparison of dynamic coordination between species

At the first glance, the observed turn-taking in the confederate-trained macaques resembled the long turn-taking observed in some human pairs (cf. **Figure 3**, FC-2,5). Yet, the analysis of reaction time distributions suggests that macaques employed a competitive version of turn-taking in which the faster agent selects its own color (insists) while the slower agent follows (accommodates). Such behavior would be in line with theoretical predictions derived from our evolutionary simulation (*8*), showing that competitive turn-taking provides the most effective strategy for a BoS type game when players have high probability to observe partner's choices. To further test the hypothesis that this macaque pair developed a competitive Leader-Follower strategy, we performed a correlation analysis between the modeled visibility of the faster agent's action by the slower agent and the observed likelihood of following to the faster agent's color (the faster animal nearly always opted to select his own color). **Figure 5D** shows macaque agent A's probability to select his non-preferred color, the "share of other's choices" (the inverted share of own choices) and the modeled probability for A to see B's choice before A made own choice, and the same for macaque agent B (see Materials and Methods for details on modeling). In both monkeys, there was a strong correlation between the model and the observation, ranging from 0.92 for smoothed data to 0.64 for unsmoothed data, all significant at p < 0.001. This analysis further supports the notion that the confederate-trained macaque pair had developed a competitive turn-taking (alternating Leader-Follower) strategy. Strong positive correlations between action visibility and "following" behavior were present in all six post-confederate-trained sessions, indicating bouts of dynamic coordination (**Table S2**).

In contrast, human participants did not show a strong trial-by-trial relationship between relative reaction time differences and choices (**Table S3**). Even in the pair with the strongest positive correlations (pair 12, **Figure 5E**), the "following" behavior was weak and resulted from a gradual transition from B initially following A to A following B later in the session, rather than from alternating competitive turn-taking. The lack of such a correlation does not mean that reaction

times and action visibility played no role in human choices. Indeed, in 47% (9 of 19) human pairs the relative difference in reaction times between the agents significantly differed in coordinated trials compared to non-coordinated trials (**Table S1**). Hence, similarly to macaques, seeing the faster agent's choice can help the slower agent to coordinate. But unlike the macaque pair, dynamic human coordination was not driven by the faster agent striving to select the own preferred color (except human pair 12 shown in **Figure 5E**).

The correlation analysis shows that for the confederate-trained macaques the visibility of the other's action before making one's own decision correlated with the probability to follow the choice of the other in a trial-by-trial fashion, with the faster monkey "selfishly" selecting its own preference. Turn-taking in humans, however, did not rely on temporal competition to systematically steer the pair towards the faster agent's preferred color. To highlight this difference further, we asked how the two species established and maintained the dynamic coordination (when it was present). The trial-by-trial sequences of choices suggested that confederate-trained macaques transitioned between periods of coordination using short periods of non-coordination (**Figure 6A**, magenta arrows), while turn-taking humans tended to switch seamlessly between the two coordination options (despite verbal communication being prohibited). **Figure 6B** illustrates all possible uncoordinated and seamless switch patterns, sorting them to "selfish" (an agent initiates a transition to own preferred target) and "benevolent" (an agent initiates a transition to other's preferred target). **Figure 6C-F** provides a quantification of different aspects of transitions as scatter plots of benevolent choices on the y-axis and selfish choices on the x-axis, in the two species. Specifically, **Figure 6C** compares the frequency of benevolent and selfish choices after the faster agent switched seamlessly from one coordination combination to the other. Turn-taking humans showed a high and roughly equal amount of benevolent and selfish seamless switches, indicating overall balanced switching behavior. The four naïve macaque pairs showing the highest number of seamless switches all used a side-based strategy, which due to the color-to-side randomization trivially generated seamless color switches. Confederate-trained macaques and non-turn-taking humans showed only very few seamless switches.

Alternative to seamless switches, agents can end the current coordination block by switching to non-coordination (cf. **Figure 6B**). **Figure 6D** shows the total numbers of own-own vs. other-other non-coordinated trials. These trials can be used to initiate a switch of coordination (for example refer to **Figure 1E**, from ~trial 275 to the end: in this human pair three coordination switches were initiated by "selfish" own-own trials and four by "benevolent" other-other trials). All

macaque pairs/sessions are below the unity-diagonal, indicating selfish preference to select own color. Turn-taking humans tend to show fewer non-coordinated trials and lay along the diagonal, indicating that in a half of trials there was a benevolent intention to initiate the switch to the other's preferred color.

The above analysis and the inspection of choice sequences (cf. **Figure 6A**) shows that in the confederate-trained macaques coordination epochs were separated by epochs of non-coordination that looked like each agent "challenged" the other to accommodate. This poses the question whether there was a temporal component in how a pair initiated the challenging epochs and how they resolved those challenges. Of particular interest are the confederate-trained macaques because they showed a strong choice dependence on the relative reaction time difference. Focusing on the transitions from coordinated to non-coordinated epochs and vice versa, **Figure S7** shows that the faster agent slowed down and the slower agent sped up during the transition to non-coordination. Conversely, the agents sped up to initiate the coordination on the own color, and the accommodating agents slowed down.

Figure 6E compares, for all coordination-to-non-coordination transitions, the number of own vs. other's color selections by the faster agent. Here all macaque pairs/sessions are located below the diagonal indicating that the faster agent tended to act selfish, while turn-taking humans mostly lay along the diagonal, again indicating balanced switching behavior. Finally, **Figure 6F** compares the number of times the faster agent selected own or other's color in trials when the pair when they resolved a challenge by switching from non-coordination to coordination. Confederate-trained macaque sessions and 6 out of 9 naïve macaque pairs fall below the diagonal indicating selfish choices of the faster agent. Turn-taking humans, however, lay along the diagonal indicating balanced challenge resolution.

Taken together this shows that in turn-taking human pairs there are no indications of competitive turn-taking, as the faster agents showed balanced selection of both coordination options. Furthermore, in four out of the ten turn-taking pairs, one agent was significantly faster than the other in coordinated trials as compared to non-coordinated trials, regardless if it was coordination on its own or the other's color (**Table S1**). This means that such an agent initiated switches to and from the own color; while being able to see the faster agent's action likely helped the slower agent to maintain coordination and to accomplish the observed high rate of seamless switches. In contrast, the confederate-trained macaques showed no seamless switches, but displayed temporal competition in which the faster agent led to his own color.

Figure 6. Humans and macaques show different switching behavior. (A) A magnification of the middle part of the macaque F-C session shown in Figure 5B (trials 211 - 495); color bars represent the choice combinations; the magenta arrows point to uncoordinated switches that resulted in a change of the coordination color. **(B)** 8 possible switch types, classified as selfish or benevolent based on which agent acted faster. Seamless switches involve uninterrupted change from one to another coordinated option; uncoordinated switches involve one or more uncoordinated trials. **(C)** The behavior of a faster agent for all seamless switches between the two coordination

modes (red-to-blue and blue-to-red). Here and in E-F, the x-axis shows the number of own target selections (selfish), the y-axis the number of non-preferred target selections (benevolent). Turn-taking humans, as well as naïve monkey pairs that coordinated on one side, showed relatively high and balanced numbers of seamless switches, while the confederate-trained macaque pair and the non turn-taking humans showed only few seamless switches. Here and in D-F, the diagonal dashed line shows the line of equality, the black dashed contours show the 95% confidence interval around the diagonal. **(D)** The number of uncoordinated trials in which both subjects selected their respective preferred target (own-own) on the x-axis, and the number of trials in which both subjects selected their non-preferred target (other-other) on the y-axis. Turn-taking human pairs mainly cluster along the diagonal while naïve and trained macaque pairs cluster in the lower right half (mostly significantly below the diagonal); non turn-taking human pairs spread along the main axes. Thus in macaques uncoordinated trials that interrupt blocks of coordinated trials show selfish behavior while turn-taking humans show fair behavior with similar amounts of selfish and benevolent uncoordinated trials. **(E)** The switches from coordination to non-coordination, or "challenge initiations", for the faster agent. Monkey pairs fall below the diagonal (with the two competitive turn-taking sessions in pair FC even below the 95% confidence intervals around the diagonal). Turn-taking human pairs again cluster around the diagonal. **(F)** The behavior of the faster agent for all switches from non-coordination to coordination, or "challenge resolutions". Turn-taking humans cluster along the diagonal, while the trained monkey pair clustered below the diagonal (with the two competitive turn-taking sessions below 95% confidence interval around the diagonal).

Discussion

We studied macaque and human pairs in a coordination game, which offered higher rewards for selecting the same option, but entailed an inherent conflict about which of the two coordinated options to select. Both species largely converged on coordinated behavior but in a markedly different fashion. Many human pairs achieved nearly-optimal coordination and fair (cooperative) turn-taking that equalized the rewards of the two partners, without communicating explicitly. Macaques, instead, tended to exploit simpler solutions that allowed them to maximize their reward without the need to track the actions of the partner in real-time. That is, they used static instead of dynamic coordination. The macaques' behavior significantly shifted from such simple strategies to competitive turn-taking after they were trained to observe and attend to the other's choice with a human confederate. In post-confederate-training sessions, the choice behavior was strongly correlated with trial-by-trial differences in the reach reaction times: the faster monkey chose his preferred option and the slower one followed. Our results show that both humans and macaques can take available information about the other's action into account before making their own decisions. There was, however, a fundamental difference: when coordinating dynamically, monkeys showed competitive turn-taking while in humans the faster agent often "offered" switching to the partner's preferred color, exhibiting a form of benevolent and fair turn-taking.

Coordination in naïve humans and macaques

Half of the human pairs (10 of 19) converged on dynamic coordination that balanced the reward, developing fair, cooperative turn-taking, while another five pairs coordinated on the same fixed color throughout the session. Similar to the approach of Brosnan and colleagues (*22*, *30*), the human subjects in the present study had to infer the underlying payoff rules while playing, to closely match the procedure in macaques. Even under such conditions, the majority of humans arrived at optimal behavior within a single session, in our study as well as in other coordination and anti-coordination games (*22*, *25*, *30*). The observed turn-taking frequency was close to other variants of BoS games with explicit instruction of the payoff matrix where approximately 50-60% of human subjects developed turn-taking (*37–39*, *53*).

In contrast to humans, macaque pairs converged on simpler strategies, either selecting the same color (56%) or selecting the same side (44%). These two strategies require less trial-by-trial coordination effort than the turn-taking, because it allows each agent to make decisions without paying attention to the trial-specific actions of the partner. However, all but one macaque coordinated both via same color or same side selection depending on the partner, and no macaque persisted on the own color with all tested partners. For instance, monkey F insisted on his preferred color when playing with monkey C, accommodated to the other's color when playing with monkey T and converged on his right side when playing with monkey M. This indicates that macaques took the partner's actions into account, albeit not in the strict trial-by-trial or within trial fashion.

While our study is, to the best of our knowledge, the first implementation of a BoS paradigm in macaques, other 2×2 games have been investigated in several primate species (humans, chimpanzees, macaques and capuchins). Human and nonhuman species show qualitatively comparable capabilities to coordinate choices and to maximize payoffs in simple coordination games such as Stag Hunt (*22*, *25*), and in the mixed-motive Prisoner's Dilemma, where macaques showed mutual cooperation significantly above chance and mutual defection below chance (*29*), similarly to humans (*26*). Conversely, the behavior of humans and nonhuman primates differs when the conflict can be resolved by turn-taking. In the anti-coordination Conflict game where maximum joint reward is obtained if each agent selects a different icon (*30*), capuchins and macaques converged on only one of the two possible asymmetric equilibria, much like four of our nine naïve macaque pairs (while 11 out of 27 human pairs balanced payoffs by alternating between the two equilibria, similar to the 10 turn-takers out of

our 19 pairs). Hence, the data suggest that monkeys do not spontaneously adopt turn-taking, despite coordinated reciprocity being a successful evolutionary strategy in simulated non-communicating agents (*34*).

Action visibility and dynamic coordination

The lack of spontaneous dynamic coordination such as turn-taking in naïve macaques in our experiment cannot be attributed to their inability to infer the upcoming choices of the partner, because they could observe partner's actions through the transparent display. This setup is different from traditional economic games, which are either played simultaneously without mutual action visibility or sequentially. It has been demonstrated that coordination (or anti-coordination) typically is improved with increasing information about the other's choice by shifting from the strictly simultaneous to the sequential mode (*54*, *55*). The transparent game approach we adopted here differs from those two classical modes in that each subject can decide independently when to act (within a certain time window). Brosnan and colleagues used a similar approach to show that compared to an opaque simultaneous setting, macaques and capuchins improved their anti-coordination in a transparent (called asynchronous) version of a Conflict game. In this study, subjects looked at the same monitor while sitting next to each other and could observe each other's decisions as cursor movements (*30*).

In our setup, agents sat opposite to each other and saw actual eye and hand movements, combining face-to-face and action visibility. In humans, face-to-face visibility in a simultaneous iterated Prisoner's Dilemma (iPD) and in the Ultimatum game significantly improves mutual cooperation, even without action visibility (*56*, *57*). These findings support the idea that non-verbal social signals and, more generally, the observable presence of others influence decision-making. The ongoing action visibility per se also had a profound effect on mutual cooperation in iPD (*58*) and in a web interface-based anti-coordination game conceptually similar to BoS (*44*). In a competitive reaching task, a face-to-face transparency allowed human subjects to glean useful information from observing the relevant hand effector and from seeing the face and the full body of an opponent (*59*). Seeing actual movements rather than relying on abstract representations of others such as cursor motions might be especially crucial in nonhuman experiments.

The importance of action visibility is also supported by computational modeling. In the continuous-time cooperation games such as iPD and Stag Hunt where agents could observe and respond to each other's actions in real-time, cooperation by coaction is more easily

obtained and stabilized against exploitation than the cooperation that relies on delayed reciprocity (*41*). For the iPD and BoS transparent games, we showed that different coordination strategies are preferable for different probabilities of seeing partner's choice (*8*, *9*). When these probabilities are low (or when agents do not utilize the action visibility), simple strategies like "win-stay, lose-shift" are most effective. For higher probabilities of seeing partner's choice more complex strategies emerge, such as coordinated turn-taking and temporal Leader-Follower, where the faster agent determines the choice of the slower agent. For the selfish agents this means that the faster one insists on its individually-preferred own target and the slower accommodates (*8*).

Given that many human pairs exhibited spontaneous dynamic coordination, while naïve macaques did not, we can ask if and how these two species utilized the action visibility. Our findings suggest that humans relied on action visibility for seamless switches from one color to another, and for maintaining the coordination within a block of trials. But besides one pair, there was no indication that humans employed the competitive variant of the Leader-Follower strategy. Instead, faster agents were as likely to switch from their own to the other's color, indicating benevolent turn-taking.

Very different dynamics transpired in additional experiments with macaques. While naïve macaques did not seem to monitor each other's choices in each trial, pairing them with a human confederate playing a color alternation strategy in short blocks demonstrated that macaques are capable to closely follow the other's actions, and adopt the imposed "turn-taking". Moreover, temporarily blocking the view of the confederate's hands abruptly changed macaque behavior, disrupting the coordination. The macaques' coordination was thus driven by action visibility and not by strategies like "win-stay, lose-shift" or trial counting. In line with the Leader-Follower strategy, macaques were faster than the confederate when selecting own color, but waited for the confederate when selecting the confederate's color. Intriguingly, pairing the two macaques that completed the training with a human confederate resulted in behavior different from both naïve macaque behavior and from the "following behavior" with the confederate. The confederate-trained macaques competed, with the faster agent selecting the own color and the slower agent following. This resulted in either sustained coordination on one fixed color or in turn-taking. Comparing the temporal signatures of the reaction time differences in trials around a switch to or from coordinating on a specific color further indicated that macaque turn-taking was competitive and dynamic in nature. The break of coordination was triggered by the faster agent slowing down and the slower one speeding up and selecting the own color; conversely, the

transition to coordination was associated with the speeding up agent selecting the own color and the slowing down agent accommodating.

In summary, both species exploited action visibility to achieve and/or maintain the turn-taking, emphasizing the importance of the "transparency" of interactive behavior as an important determinant of emerging strategies. Human agents initiated switches to and from their preferred colors to "fairly" balance the payoffs, similarly to the Conflict game (*30*), while macaques established competitive dynamics, as predicted by our evolutionary simulations (*8*). These results add to the body of literature indicating that humans in a social setting might base their decisions not only on pure reward maximization (*60*). It is also plausible that such normative behavior can reflect prospective planning to ensure a stable persistence of the individually-beneficial cooperation. The lack of the cooperative turn-taking in nonhuman primates, as compared to the humans' propensity to engage in it, might reflect cognitive limitations in long-term planning and perspective taking (*61*). Species differences in general cooperative behavior may explain both the reluctance of naïve macaques to coordinate dynamically, as well as the competitive nature of dynamic interactions. The transparent BoS game requires the agents to cooperate in close proximity in the pursuit of an immediate reward which necessitates some degree of social tolerance by both partners. If we can take natural food sharing behavior as a proxy for social tolerance, it is not equally prevalent among primate species. Humans, apes, some baboon species (*62*), and some New World monkey species including capuchins share food even between adults, while rhesus macaques do not even share food with their offspring (*63*). In line with these patterns, mutually beneficial alternating task performance and turn-taking has been observed in humans (*17*, *30–32*), apes (*64*, *65*), and capuchins (*66*), but not in more despotic, less tolerant rhesus macaques (*5*).

Limitations and future directions

There are several limitations in our study that have to be considered and possibly addressed in future experiments. Firstly, we only tested 6 macaques (and 9 pairs), all males, and only one pair was housed together (others were from neighboring but separate enclosures). Therefore, we cannot say much about the influence of social rank. Anecdotally, however, it did not seem that the more dominant agent always prevailed. For instance, one animal (monkey T), the smallest and likely to be subordinate to all other partners, successfully "insisted" on his own color in 2 out of 3 cases (cf. (*67*)). Furthermore, we only tested two confederate-trained macaques so far. While the results are exciting and show that macaques can engage in

dynamic competitive turn-taking, future experiments will need to test how generalizable this pattern is.

Secondly, due to the task target color/side randomization, selecting targets on the same side of the display resulted in low effort, efficient and fair coordination. We cannot say if fairness was a contributing factor in convergence of the four naïve macaque pairs to this strategy, although we deem it highly unlikely. A new experiment where the probability of a specific color to appear on the same side is parametrically modulated is needed to evaluate the conditions under which this strategy emerges (or what level of unfairness is accepted). Furthermore, it would be interesting to test human pairs in several sessions, similar to macaques, to see if humans might also converge on the fixed side strategy as means of fair and seamless coordination. It requires preventing the human participants from any verbal contact after and between the sessions however. Finally, it would be important to vary the ratio and the range of reward magnitude ("stakes"), as it has been shown that stakes may strongly affect the individual (*68*) and social (*44*) decisions.

In summary, we show that both, humans and nonhuman primates, dynamically coordinate in real-time transparent interactions, but also, that macaques compete while humans strive for fairness by prosocial turn-taking. Ensuring real-time action visibility with our shared transparent display setup provides novel comparative insights into social decision processes as they unfold in real-time during transparent interactions, and offers a new route to behavioral and neural investigations of dynamic decision-making in cooperative and competitive contexts (*69*, *70*).

Materials and Methods

Participants

Humans. 38 right-handed subjects (23 females, mean age: 26.1 $\pm$ 4.1 SD, range 20 to 41 years) participated in the study as paid volunteers. Subjects were tested as 19 unique pairs, i.e. each participant contributed only once. Instructions given to the subjects before the experiment are provided in Supplementary Materials. In short, subjects were instructed how to operate the setup and to interpret the auditory feedback as an indicator of the earned reward. They were not given an explicit description of the task's payoff structure beyond "Your reward will depend on your own and your partner's choice". Prior to the experiment, subjects were individually familiarized with the setup and practiced a single-player ("solo") version of the task. All subjects gave written informed consent for participation after the procedures had been explained to them

and before taking part in the experiment. Experiments were performed in accordance with institutional guidelines for experiments with humans and adhered to the principles of the Declaration of Helsinki. The experimental protocol was approved by the ethics committee of the Georg-Elias-Mueller-Institute for Psychology, University of Goettingen (GEMI 17-06-06 171). We excluded 2 additional pairs from the analyses due to experimental differences (one pair did not perform the initial individual training; one pair ended the experiment prematurely).

Macaques. Research with nonhuman primates represents a small but indispensable component of neuroscience research. The scientists in this study are committed to the responsibility they have in ensuring the best possible science with the least possible harm to the animals (*71*).

Six adult male rhesus monkeys (designated by initials C, E, F, L, M, T) participated in the study, yielding 9 pairs (each monkey participated in 2 or more pairs). Animals were extensively trained with positive reinforcement to climb into and stay seated in a primate chair. The experimental procedures were approved of by the responsible regional government office (Niedersaechsisches Landesamt fuer Verbraucherschutz und Lebensmittelsicherheit (LAVES), permits 3392-42502-04-13/1100 and 3319-42502-04-18/2823). The animals were pair- or group-housed in facilities of the German Primate Center (DPZ) in accordance with all applicable German and European regulations. The facility provides the animals with an enriched environment, including a multitude of toys and wooden structures (*72, 73*), natural as well as artificial light and access to outdoor space, exceeding the size requirements of European regulations, and a rich diet including primate biscuits, fruit and vegetables. During the study the animals had unrestricted access to food and fluid, except on the days where data were collected or the animal was trained on the behavioral paradigm. On these days, the animals were allowed access to fluid through their performance in the behavioral paradigm. The DPZ's veterinarians, the animal facility staff and the lab's scientists carefully monitor the animals' welfare.

Experimental setup

For maximal comparability of the human and the monkey behavior, we developed a novel Dyadic Interaction platform in which two human or nonhuman primate subjects co-act in a shared workspace while sitting face-to-face (**Figure 1A, B**). Joint dyadic tasks have been previously implemented in a side-by-side or 90° angle setting with a shared or separate workspaces for each subject (*3, 29, 30, 70, 74–76*), using back-to-back nontransparent monitors (*56, 57*), or in a table-like setting with opposing subjects acting in a horizontal workspace that is

not in the line of sight between the subjects' faces (*43*, *77*, *78*). Only few studies utilized transparent face-to-face arrangement of the subjects, using a video projector and two semitransparent mirrors virtually placing the stimuli into the shared plane (*79*), or incorporating physical targets into transparent Plexiglas screen (*59*). Similarly, with our design we aimed for maximal availability of mutual social signaling in a face-to-face setting, a shared vertical workspace with computer-controlled stimuli in the direct line of sight, without risk of injuries due to physical contact, and suitability across species. We achieved this using a novel transparent display (1920 x 1080 pixels, 121 cm x 68 cm, 60 Hz, EYE-TOLED-5500, Eyevis, Reutlingen, Germany), amended for dual-side touch sensitivity (PQLabs, G5S, Freemont, CA, USA) with a custom-built "sandwich" construction. Sitting on either side of the display, both subjects saw each other and the same stimuli. Two proximity sensors (Carlo Gavazzi CA18-CA30CAF16NA, Lainate, Italy) per subject for the left and right hand respectively, mounted below the display ("home" buttons) and the two touch panels mounted on either side of the display registered hand positions of both agents, at 240 Hz temporal resolution.

Experimental control and stimulus presentation were implemented using the EventIDE software package (Okazolab, Delft, The Netherlands). Liquid reward for monkeys was delivered via computer-controlled peristaltic fluid pumps, for every correctly performed trial.

Dyadic decision-making task

Bach-or-Stravinsky game. Since we are interested in the effect of mutual action visibility on coordination behavior, we implemented a transparent version of a Bach-or-Stravinsky (BoS) game, in which each player's time-continuous visuomotor behavior can be seen by the other player. Conceptually, in the BoS game two agents are choosing between going to Bach or Stravinsky concerts. Agent A prefers Bach, Agent B prefers Stravinsky; yet, both prefer going to the concert together (*80*). Thus, agents wish to coordinate their behavior but have conflicting interests. This is a classic 2×2 non-zero sum game (also known as the Battle of the Sexes) with two pure strategy Nash equilibria (Bach-Bach, or Stravinsky-Stravinsky), and one less efficient mixed strategy Nash equilibrium, where the agents go to their preferred event more often than the other (*20*, *36*).

In our implementation, the agents were choosing between two options represented by two differently colored targets placed in the left and the right side of the display. Each target color was associated with a higher reward for one of the agents resulting in individually preferred targets. An agent selecting own preferred target was assured to get at least 2 reward units,

while selecting the other's target (individually non-preferred target) yielded at least 1 reward unit. Additionally, when both agents selected the same target, a bonus of 2 reward units was added to the payoff of each agent. Thus, the maximum average joint reward (3.5 units) was obtained on coordinated trials when one agent selected own preferred target (getting 4 reward units) while the other chose non-preferred target (getting 3 reward units). Note that on any given coordinated trial, i.e. when agents selected the same target for overall higher reward, this payoff matrix (**Figure 1C**) resulted in an unequal (i.e. unfair) reward distribution. Hence, the BoS paradigm probes both the ability to realize that coordinated target-selection results in higher rewards than non-coordinated target-selection, as well as the ability to perceive and counteract the unfairness/conflict situation.

The task mechanics. On every trial, each subject chose between the individually-preferred and non-preferred target on the touchscreen. Targets were light blue circles of 35 mm diameter with either a red (individually-preferred target for Agent A) or a yellow (individually-preferred target for Agent B) rings (for better visualization, we replaced yellow and red rings with blue and red solid targets in the figures and the text).

Subjects had to place both hands on the two home buttons for 500 ms to start a trial after the inter-trial interval (ITI). This allowed us to control which effector (acting hand) was used. (**Figure 1D**). Then an initial fixation target without a colored ring appeared on the display (10 cm below eye level). Subjects had 1500 ms to touch this initial fixation target with the instructed hand. After both subjects touched the fixation target they had to hold it for 500 ms. Then two choice targets appeared at one of three different pairs of positions (140 mm to the left and right of the central fixation target and either at the same height as the central fixation target or 35 mm below or 35 mm above it). We randomized the "red" location equally over all six positions, balanced in sets of 18 trials with each of the six "red" positions appearing three times, and vertically mirrored the opposite location. Simultaneously with targets' appearance the initial fixation target disappeared, serving as a go signal. Subjects had 1500 ms to make their choice and touch one of the targets. After both subjects acquired their chosen target, selected target(s) brightened, and subjects needed to hold the hand on the target for another 500 ms. At that point, the choices were evaluated and rewards were dispensed according to the payoff matrix. The amount of reward earned by each subject was signaled by two sequential series of auditory pulses, with different pitch for each subject. Each pulse was constructed as a harmonic series with 12 overtones and a fundamental frequency of 443 Hz for side A and 733 Hz for side B to provide distinct sounds for each agent. For the monkeys, we delivered water as reward

(approximately 0.14 ml per pulse) concurrently with the auditory pulses for the respective monkey's side; humans were instructed to expect "a few" cents per pulse, the accumulated earnings were paid out as a lump sum after the experiment. After the reward period subjects had to wait for an inter-trial-interval of 1500 ms before they could initiate the next trial.

The side of the targets was randomized on each trial, i.e., in ~50% the red target was on the right, the blue on the left, and vice versa. This design can lead to the three following coordination patterns (or any mix of the three):

i) Coordinating statically by each agent repeatedly selecting the same fixed color of the target, irrespective of unfair distribution of the rewards.

ii) Coordinating statically by each agent repeatedly selecting the same fixed side. Due to color/side randomization, this pattern ensures a fair reward distribution.

iii) Coordinating dynamically by both agents selecting the same target, while picking from both colors and both sides trial-by-trial (e.g. trial 1 red right, trial 2 red left, trial 3 blue left, … etc.). This could result in a fair or unfair reward distribution depending on the ratio of red to blue color selection.

Human procedure. Human subjects were recruited via a university job website and pairs were selected based on matching schedules. Subjects were given a brief introduction to the experiment (see Supplementary Materials); this material included the information "Your reward will depend on your own and your partner's choice" and a very basic description of the task ("You will have to choose one of the two circles presented to you. […] You will have to decide and respond quickly."), but did not include details of the payoff matrix. After the joint introduction, each subject alone performed 50 or 100 individual (solo) trials, to learn how to operate the touchscreen and to develop a preference for one of the two color targets (see SI for the exact verbal instructions given prior to solo training). Subjects were positioned ~50 cm from the display with the height of the chairs adjusted such that both subjects' eyes were ~121 cm above the ground. After the solo training, both subjects entered the setup for the main dyadic task which lasted for 300 or 400 trials. Participants had to infer the task rules by exploration, similar to the macaques (*25*). Each session lasted approximately 1.5 hours. After the experiment we conducted an individual debriefing and paid the earned reward separately for each subject.

Macaque procedure. Macaques were brought to the setup in their individual primate chairs. The chairs were positioned such that the eyes were 30 cm from the display. The monkeys had

previously been trained to perform the basic task structure (hands on proximity sensors, reach to the initial fixation target with the instructed hand, and select one of the presented choice targets by reaching to it). The animals performed the solo version of the task with differential rewards to develop a preference for one of the color targets and were only paired with a conspecific after selecting a higher rewarded target in ≥ 75% of the trials. Thereafter, pairs of macaques worked together in the dyadic version of the task, for 11 ± 7 sessions (range 4-25).

Data analysis

We computed the following six *aggregate measures of choice behavior and coordination,* explained below*: s*hares of "own" choices, shares of objective left choices, mutual information for target choice, mutual information for side choice, average reward, and dynamic coordination reward. We computed each of these measures for the last 200 trials of each session in order to assess the "steady-state" behavior after allowing for an initial period of exploration.

Shares of own and objective left choices. The share of own choices (SOC) is the fraction of trials where an agent has selected the individually preferred target. Similarly, the share of left choices (SLC) is a fraction of trials where an agent has selected the target on the "objective" left side of the display (which is the left side for Agent A and right side for Agent B). Fractions range from 0 to 1. For figures showing SOC/SLC over the course of a session we also calculated both measures for the session as a whole, and in running windows of w = 8 trials. In the latter case, SOC and SLC can take the values of 0, 1/8, 2/8, …, 1. For instance, SOC = 0 means that an agent has selected individually non-preferred target for 8 trials in a row.

Mutual information. Mutual information (MI) represents the reduction of uncertainty regarding the values of one time series provided by knowing the values of the other time series. Here we consider mutual information for the color of the target (MIT) and side (MIS) choices of the two agents, showing how much information the target/side choices of one agent provide about the respective choices of the other. Mutual information is measured in bits. Since both target and side choices are binary (in each trial an agent selects either the preferred or non-preferred target and either the left or right side), both MIT and MIS range from 0 bit (the choices of one agent provide no information about the choices of the other) to 1 bit (the choices of one agent can be inferred precisely from the choices of the other agent).

For instance, if both agents select the objective left target in every odd trial and the objective right target in every even trial, both MIT = MIS = 1, since for every trial the target and the side

selected by one agent entirely describes the target/side selected by the other. At the same time, the choices of each agent individually are highly uncertain: both sides and both targets are selected with the same probability of 0.5. If both agents constantly select the objective left side, this would result in MIT = 1 due to side-randomization of target color, but MIS = 0, since there is no uncertainty regarding the side selection and thus no additional knowledge about the other's choice can reduce the uncertainty.

Formally, mutual information of time series $X = (X_t)$ and $Y = (Y_t)$ is given by

$$\mathrm{MI}(X, Y) = \sum_{x,y} p(X_t = x, Y_t = y) \log_2 \frac{p(X_t = x, Y_t = y)}{p(X_t = x) p(Y_t = y)} \qquad (1)$$

where $p(X_t = x)$ is the probability of the value x in time series X, $p(Y_t = y)$ is the probability of the value y in time series Y, and $p(X_t = x, Y_t = y)$ is the joint probability to simultaneously have values x and y in time series X and Y, respectively, and x and y can be either 0 or 1, so that the sum is over all four combinations.

Since in our case time series X and Y have finite length, we simply replace probabilities by relative frequencies. This is known as a naïve estimation of mutual information, but it is sufficiently precise for binary time series (*81*, *82*).

To test whether the MI values were significantly different from zero, we generate Whittle surrogates for the given choice time series and estimate from them the threshold for the given significance level (p=0.01 in our case) (*83*).

Average reward. Average reward (AR) is computed as the average of an agent's payoff across the session. Note that the average reward of each individual agent can be in the range of 1 to 4 points, while the average joint reward of a pair cannot exceed 3.5 (since when one agent gets payoff of 4, the other agent gets only 3). The average reward for completely independent choices of two agents with 50% probability for either target is 2.5 (but note that an achieved reward of 2.5 is not a positive proof of independent choices).

Dynamic coordination reward. Dynamic coordination reward (DCR) is the surplus reward of the two agents compared to the reward they would get by playing randomly. By *playing randomly* we mean that choices of the agents in each round are independent of the history and of the current choices of the partner. The "random reward" is computed by selecting the two color targets with the same probabilities as actually observed in the two agents, but randomly permuting the choices over trials. For our payoff matrix, the range of DCR is [-1,1], with -1

corresponding to very inefficient playing (alternating selection of the two anti-coordination options), while 1 corresponds to very efficient playing with explicit coordination (for instance, turn-taking). DCR is hence a measure of dynamic (reciprocal) coordination. For instance, if both agents would coordinate statically by constantly selecting one and the same side, this would result in $\mathrm{DCR} = 0$, even though this coordination pattern still yields the maximum average reward of 3.5.

Formally, DCR is defined as the actual average reward of a pair (R_{actual}) minus the reward the agents would get if they were playing randomly (R_{PR}). $\mathrm{DCR} = R_{\mathrm{actual}} - R_{\mathrm{PR}}$. The reward for playing randomly (R_{PR}) depends only on eight probabilities, four for each agent. Below index i indicates the agent and stands for either A or B:

$P_{i,1,\mathrm{left}}$ - probability to select non-preferred objective left target (which is the left side for agent A and right side for agent B),

$P_{i,1,\mathrm{right}}$ - probability to select non-preferred objective right target,

$P_{i,2,\mathrm{left}}$ - probability to select preferred objective left target,

$P_{i,2,\mathrm{right}}$ - probability to select preferred objective right target.

Note that these probabilities are not independent on each other. First, for $i = \mathrm{A}, \mathrm{B}$, it holds

$$P_{i,1,\mathrm{left}} + P_{i,2,\mathrm{left}} + P_{i,1,\mathrm{right}} + P_{i,2,\mathrm{right}} = 1 \qquad (2)$$

Second, it holds

$$P_{\mathrm{A},2,\mathrm{left}} + P_{\mathrm{A},1,\mathrm{right}} = P_{\mathrm{B},1,\mathrm{left}} + P_{\mathrm{B},2,\mathrm{right}} = Q_{\mathrm{left}} \qquad (3)$$

and

$$P_{\mathrm{A},1,\mathrm{left}} + P_{\mathrm{A},2,\mathrm{right}} = P_{\mathrm{B},2,\mathrm{left}} + P_{\mathrm{B},1,\mathrm{right}} = Q_{\mathrm{right}} \qquad (4)$$

where Q_{left} and Q_{right} are the probability of agent A's preferred target to appear on the left and on the right, respectively (and of B's preferred target too appear on the right and on the left). Given the independent random selection from trial to trial described above, Q-values should approximate 0.5 for larger N of trials.

Average reward of two agents for playing randomly is computed as follows:

$$R_{\text{PR}} = 3.5\big(p_{4,3} + p_{3,4}\big) + 2p_{2,2} + p_{1,1} \qquad (5)$$

where $p_{a,b}$ is the probability that by random playing agent A gets reward a and agent B reward b. These probabilities are given by the following equations:

$$p_{1,1} = \frac{P_{\text{A,1,right}} \cdot P_{\text{B,1,left}}}{Q_{\text{left}}} + \frac{P_{\text{A,1,left}} \cdot P_{\text{B,1,right}}}{Q_{\text{right}}} \qquad (6)$$

$$p_{2,2} = \frac{P_{\text{A,2,left}} \cdot P_{\text{B,2,right}}}{Q_{\text{left}}} + \frac{P_{\text{A,2,right}} \cdot P_{\text{B,2,left}}}{Q_{\text{right}}} \qquad (7)$$

$$p_{4,3} = \frac{P_{\text{A,2,left}} \cdot P_{\text{B,1,left}}}{Q_{\text{left}}} + \frac{P_{\text{A,2,right}} \cdot P_{\text{B,1,right}}}{Q_{\text{right}}} \qquad (8)$$

$$p_{3,4} = \frac{P_{\text{A,1,right}} \cdot P_{\text{B,2,right}}}{Q_{\text{left}}} + \frac{P_{\text{A,1,left}} \cdot P_{\text{B,2,left}}}{Q_{\text{right}}} \qquad (9)$$

To see why this is the case, consider, for instance, $p_{1,1}$. Both agents get reward of 1 when they both select the other's preferred target, either when target of agent A appears on the left side (probability of this is encoded by the first term) or on the right side (second term).

To compute confidence intervals, we use the fact that SLC_i and SOC_i for an agent i can be considered as binomially distributed, thus radiuses of their confidence intervals ΔSLC_i and ΔSOC_i can be estimated by the classic method of approximating the distribution of error around binomially-distributed observation with a normal distribution. To obtain the confidence interval for DCR, it is sufficient to compute maximal and minimal possible DCR given that SOC and SCL of the two agents are within the respective confidence intervals. Note that $\text{SLC}_i = P_{i,1,\text{left}} + P_{i,2,\text{left}}$, $\text{SOC}_i = P_{i,2,\text{left}} + P_{i,2,\text{right}}$, thus all the probabilities necessary for calculating DCR can be computed from SLC_i and SOC_i given that Q_{left} and Q_{right} are fixed to 0.5 in the reported experiments. Simple analysis reveals that minimum and maximum DCR should be at the edges of the 4-D confidence interval formed by SLC_i and SOC_i of the two agents, which reduces the problem to testing 16 DCR values computed for $\text{SLC}_i \pm \Delta\text{SLC}_i$ and $\text{SOC}_i \pm \Delta\text{SOC}_i$.

For the main analysis presented in the Results, we used last 200 trials to compute DCR values. Using the last 150 or 250 trials for DCR analysis resulted in the exact same 10 human pairs with significant results as using the last 200.

Reaction and movement time measurements. We measured the time from the onset of the choice targets to the release time of the initial fixation target (t_{release}) and to the acquisition time

of the selected target, $t_{\text{acquisition}}$ individually for each subject. We then calculated the movement time t_{movement} and reaction time t_{reaction} as follows:

$$t_{\text{movement}} = t_{\text{acquisition}} - t_{\text{release}} \qquad (10)$$

$$t_{\text{reaction}} = t_{\text{release}} + \frac{t_{\text{movement}}}{2} \qquad (11)$$

We use t_{reaction}, the half-way point between t_{release} and $t_{\text{acquisition}}$, i.e. the duration from target stimulus onset to the half-time of the reach movement, as a proxy for the estimated time at which the trajectory of each subject's reach movement should be evident for the other agent. Typical movement time values (mean ± SD across trials) in our experiment ranged from 314 ± 104 ms for humans, 171 ± 77 ms for macaques, and 180 ± 64 ms for the confederate trained macaque pair.

Estimated probability to see partner's choice. When an agent is acting slower than the partner, there is a chance for this agent to see the partner's choice and use this information for making the own choice. We therefore modeled the probability to see the partner's choice as a logistic function of the difference of the agents' reaction times, in each trial. The logistic function $p_{\text{see}}(t)$ had an inflection point at 50 ms ($p_{\text{see}}(50) = 0.5$) and reached its plateau phase at 150 ms ($p_{\text{see}}(150) = 0.98$). Formally, the function was given by the following equation

$$p_{\text{see}}(t) = \frac{1}{1 + e^{-k(t - \Delta T_0)}} \qquad (12)$$

with $k = 0.04$ (steepness of the slope) and $\Delta T_0 = 50$ ms (inflection point). The values of p_{see} were used for the analysis of correlation with probability of selecting the other's color (cf. **Figure 5**). We also tested a wide range of these parameters (k fixed at 0.04, $\Delta T_0 = \{12.5, 25, 50, 75, 100, 200\}$ ms and ΔT_0 fixed at 50 ms, $k = \{0.01, 0.02, 0.04, 0.08, 0.16\}$), and confirmed that the values of correlation were robust in respect to these parameters; only the longest $\Delta T_0 = 200$ ms resulted in a noticeable drop of resultant correlations.

References

1. R. W. Byrne, L. A. Bates, Primate Social Cognition: Uniquely Primate, Uniquely Social, or Just Unique? *Neuron*. **65**, 815–830 (2010).

2. A. V. Jaeggi, J. M. Burkart, C. P. Van Schaik, On the psychology of cooperation in humans and other primates: combining the natural history and experimental evidence of prosociality. *Philos Trans R Soc Lond B Biol Sci*. **365**, 2723–2735 (2010).

3. S. W. C. Chang, L. J. N. Brent, G. K. Adams, J. T. Klein, J. M. Pearson, K. K. Watson, M. L. Platt, Neuroethology of primate social behavior. *Proc. Natl. Acad. Sci. U.S.A.* **110 Suppl 2**, 10387–10394 (2013).

4. M. L. Platt, R. M. Seyfarth, D. L. Cheney, Adaptations for social cognition in the primate brain. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **371**, 20150096 (2016).

5. M. Joly, J. Micheletta, A. De Marco, J. A. Langermans, E. H. M. Sterck, B. M. Waller, Comparing physical and social cognitive skills in macaque species with different degrees of social tolerance. *Proc Biol Sci*. **284** (2017), doi:10.1098/rspb.2016.2738.

6. F. de Waal, F. B. M. Waal, *Chimpanzee Politics: Power and Sex Among Apes* (JHU Press, 2007).

7. U. Frith, C. Frith, The social brain: allowing humans to boldly go where no other species has been. *Philosophical Transactions of the Royal Society B: Biological Sciences*. **365**, 165–176 (2010).

8. A. M. Unakafov, T. Schultze, A. Gail, S. Moeller, I. Kagan, S. Eule, F. Wolf, Emergence and suppression of cooperation by action visibility in transparent games. *PLOS Computational Biology*. **16**, e1007588 (2020).

9. A. M. Unakafov, T. Schultze, I. Kagan, S. Moeller, A. Gail, S. Treue, S. Eule, F. Wolf, Evolutionary successful strategies in a transparent iterated Prisoner's Dilemma. *EvoApplications 2019, Lecture Notes in Computer Science*. **11454**, 204–219 (2019).

10. J. von Neumann, O. Morgenstern, *Theory of Games and Economic Behavior* (Princeton University Press, 1944).

11. J. K. Rilling, A. G. Sanfey, The neuroscience of social decision-making. *Annu Rev Psychol*. **62**, 23–48 (2011).

12. A. G. Sanfey, Social Decision-Making: Insights from Game Theory and Neuroscience. *Science*. **318**, 598–602 (2007).

13. S. Tremblay, K. M. Sharika, M. L. Platt, Social Decision-Making and the Brain: A Comparative Perspective. *Trends in Cognitive Sciences*. **21**, 265–276 (2017).

14. L. R. Santos, A. G. Rosati, The Evolutionary Roots of Human Decision Making. *Annu. Rev. Psychol.* **66**, 321–347 (2015).

15. S. F. Brosnan, Insights into human cooperation from comparative economics. *Nature Human Behaviour*. **2**, 432–434 (2018).

16. J. M. Smith, Game theory and the evolution of behaviour. *Proc. R. Soc. Lond., B, Biol. Sci.* **205**, 475–488 (1979).

17. A. Rapoport, M. J. Guyer, D. G. Gordon, *The 2x2 Game* (University of Michigan Press, 1976).

18. A. Rapoport, *Two-Person Game Theory* (University of Michigan Press, 1973).

19. M. Isoda, A. Noritake, T. Ninomiya, Development of social systems neuroscience using macaques. *Proceedings of the Japan Academy, Series B*. **94**, 305–323 (2018).

20. A. Rapoport, Exploiter, leader, hero, and martyr: The four archetypes of the 2 × 2 game. *Syst. Res.* **12**, 81–84 (1967).

21. B. Skyrms, *The Stag Hunt and the Evolution of Social Structure* (Cambridge University Press, 2003).

22. S. F. Brosnan, A. Parrish, M. J. Beran, T. Flemming, L. Heimbauer, C. F. Talbot, S. P. Lambeth, S. J. Schapiro, B. J. Wilson, Responses to the Assurance game in monkeys, apes, and humans using equivalent procedures. *PNAS*. **108**, 3442–3447 (2011).

23. A. F. Bullinger, E. Wyman, A. P. Melis, M. Tomasello, Coordination of chimpanzees (Pan troglodytes) in a stag hunt game. *International Journal of Primatology*. **32**, 1296–1310 (2011).

24. S. Duguid, E. Wyman, A. F. Bullinger, K. Herfurth-Majstorovic, M. Tomasello, Coordination strategies of chimpanzees and human children in a Stag Hunt game. *Proceedings of the Royal Society of London B: Biological Sciences*. **281**, 20141973 (2014).

25. S. F. Brosnan, B. J. Wilson, M. J. Beran, Old World monkeys are more similar to humans than New World monkeys when playing a coordination game. *Proceedings of the Royal Society of London B: Biological Sciences*. **279**, 1522–1530 (2012).

26. C. Wedekind, M. Milinski, Human cooperation in the simultaneous and the alternating Prisoner's Dilemma: Pavlov versus Generous Tit-for-Tat. *Proceedings of the National Academy of Sciences of the United States of America*. **93**, 2686–2689 (1996).

27. J. K. Rilling, D. A. Gutman, T. R. Zeh, G. Pagnoni, G. S. Berns, C. D. Kilts, A Neural Basis for Social Cooperation. *Neuron*. **35**, 395–405 (2002).

28. C. F. Camerer, *Behavioral Game Theory: Experiments in Strategic Interaction* (Princeton University Press, New York, N.Y. : Princeton, N.J, First Edition edition., 2003).

29. K. Haroush, Z. M. Williams, Neuronal prediction of opponent's behavior during cooperative social interchange in primates. *Cell*. **160**, 1233–1245 (2015).

30. S. F. Brosnan, S. A. Price, K. Leverett, L. Prétôt, M. Beran, B. J. Wilson, Human and monkey responses in a symmetric game of conflict with asymmetric equilibria. *Journal of Economic Behavior & Organization*. **142**, 293–306 (2017).

31. S. Grueneisen, M. Tomasello, Children coordinate in a recurrent social dilemma by taking turns and along dominance asymmetries. *Developmental psychology*. **53**, 265 (2017).

32. D. Helbing, M. Schönhof, H.-U. Stark, J. A. Ho\lyst, How individuals learn to take turns: Emergence of alternating cooperation in a congestion game and the prisoner's dilemma. *Advances in Complex Systems*. **8**, 87–116 (2005).

33. R. Noë, Cooperation experiments: coordination through communication versus acting apart together. *Animal Behaviour*. **71**, 1–18 (2006).

34. A. M. Colman, L. Browning, Evolution of cooperative turn-taking. *Evolutionary Ecology Research*, 949–963 (2009).

35. G. Bornstein, D. Budescu, S. Zamir, Cooperation in intergroup, N-person, and two-person games of chicken. *Journal of Conflict Resolution*. **41**, 384–406 (1997).

36. D. M. Kilgour, N. M. Fraser, A taxonomy of all ordinal 2 × 2 games. *Theor Decis*. **24**, 99–117 (1988).

37. D. Sonsino, J. Sirota, Strategic pattern recognition—experimental evidence. *Games and Economic Behavior*. **44**, 390–411 (2003).

38. A. P. Melis, P. Grocke, J. Kalbitz, M. Tomasello, One for you, one for me: Humans' unique turn-taking skills. *Psychological science*. **27**, 987–996 (2016).

39. C. A. Ioannou, J. Romero, A generalized approach to belief learning in repeated games. *Games and Economic Behavior*. **87**, 178–203 (2014).

40. L. A. Dugatkin, M. Mesterton-Gibbonsand, A. I. Houston, Beyond the prisoner's dilemma: Toward models to discriminate among mechanisms of cooperation in nature. *Trends in ecology & evolution*. **7**, 202–205 (1992).

41. G. S. van Doorn, T. Riebli, M. Taborsky, Coaction versus reciprocity in continuous-time models of cooperation. *Journal of Theoretical Biology*. **356**, 1–10 (2014).

42. K. R. McDonald, W. F. Broderick, S. A. Huettel, J. M. Pearson, Bayesian nonparametric models characterize instantaneous strategies in a competitive dynamic game. *Nature Communications*. **10**, 1–12 (2019).

43. W. S. Ong, S. Madlon-Kay, M. L. Platt, Neuronal correlates of strategic cooperation in monkeys. *Nature Neuroscience*. **24**, 116–128 (2021).

44. R. X. D. Hawkins, R. L. Goldstone, The Formation of Social Conventions in Real-Time Environments. *PLoS One*. **11** (2016), doi:10.1371/journal.pone.0151670.

45. G. Rizzolatti, L. Fogassi, V. Gallese, Neurophysiological mechanisms underlying the understanding and imitation of action. *Nature reviews. Neuroscience*. **2**, 661–70 (2001).

46. J. B. Silk, The strategic dynamics of cooperation in primate groups. *Advances in the Study of Behavior*. **37**, 1–41 (2007).

47. C. Sueur, O. Petit, Signals use by leaders in Macaca tonkeana and Macaca mulatta: group-mate recruitment and behaviour monitoring. *Animal cognition*. **13**, 239–248 (2010).

48. L. Conty, G. Dezecache, L. Hugueville, J. Grèzes, Early Binding of Gaze, Gesture, and Emotion: Neural Time Course and Correlates. *J. Neurosci.* **32**, 4531–4539 (2012).

49. C. Fruteau, E. van Damme, R. Noë, Vervet Monkeys Solve a Multiplayer "Forbidden Circle Game" by Queuing to Learn Restraint. *Current Biology*. **23**, 665–670 (2013).

50. A. Peysakhovich, M. A. Nowak, D. G. Rand, Humans display a 'cooperative phenotype' that is domain general and temporally stable. *Nature Communications*. **5**, 4939 (2014).

51. M. Devaine, A. San-Galli, C. Trapanese, G. Bardino, C. Hano, M. S. Jalme, S. Bouret, S. Masi, J. Daunizeau, Reading wild minds: A computational assay of Theory of Mind sophistication across seven primate species. *PLOS Computational Biology*. **13**, e1005833 (2017).

52. E. Herrmann, J. Call, M. V. Hernàndez-Lloreda, B. Hare, M. Tomasello, Humans Have Evolved Specialized Skills of Social Cognition: The Cultural Intelligence Hypothesis. *Science*. **317**, 1360–1366 (2007).

53. J. Arifovic, J. Ledyard, Learning to alternate. *Exp Econ*. **21**, 692–721 (2018).

54. A. Rapoport, Order of Play in Strategically Equivalent Games in Extensive Form. *Int. J. Game Theory*. **26**, 113–136 (1997).

55. I. Brocas, J. D. Carrillo, A. Sachdeva, The path to equilibrium in sequential and simultaneous games: A mousetracking study. *Journal of Economic Theory*. **178**, 246–274 (2018).

56. H. Tang, X. Mai, S. Wang, C. Zhu, F. Krueger, C. Liu, Interpersonal brain synchronization in the right temporo-parietal junction during face-to-face economic exchange. *Soc Cogn Affect Neurosci*. **11**, 23–32 (2016).

57. J. Jahng, J. D. Kralik, D.-U. Hwang, J. Jeong, Neural dynamics of two players when using nonverbal cues to gauge intentions to cooperate during the Prisoner's Dilemma Game. *NeuroImage*. **157**, 263–274 (2017).

58. D. Friedman, R. Oprea, A Continuous Dilemma. *American Economic Review*. **102**, 337–363 (2012).

59. M. Vaziri-Pashkam, S. Cormiea, K. Nakayama, Predicting actions from subtle preparatory movements. *Cognition*. **168**, 65–75 (2017).

60. C. Clavien, M. Chapuisat, The evolution of utility functions and psychological altruism. *Studies in History and Philosophy of Science Part C: Studies in History and Philosophy of Biological and Biomedical Sciences*. **56**, 24–31 (2016).

61. J. R. Stevens, M. D. Hauser, Why be nice? Psychological constraints on the evolution of cooperation. *Trends Cogn. Sci. (Regul. Ed.)*. **8**, 60–65 (2004).

62. A. S. Goffe, J. Fischer, Meat sharing between male and female Guinea baboons (*Papio papio*). *Primate Biology*. **3**, 1–8 (2016).

63. A. V. Jaeggi, M. Gurven, Natural cooperators: food sharing in humans and other primates. *Evol. Anthropol.* **22**, 186–195 (2013).

64. F. B. de Waal, Food sharing and reciprocal obligations among chimpanzees. *Journal of Human Evolution*. **18**, 433–459 (1989).

65. C. F. Martin, D. Biro, T. Matsuzawa, Chimpanzees spontaneously take turns in a shared serial ordering task. *Scientific Reports*. **7**, 14307 (2017).

66. A. Parrish, S. Brosnan, M. Beran, Capuchin monkeys alternate play and reward in a dual computerized task. *Animal Behavior and Cognition*. **2**, 334–347 (2015).

67. J. J. M. Massen, L. M. van den Berg, B. M. Spruijt, E. H. M. Sterck, Generous Leaders and Selfish Underdogs: Pro-Sociality in Despotic Macaques. *PLOS ONE*. **5**, 1–5 (2010).

68. W. Schultz, Neuronal Reward and Decision Signals: From Theories to Data. *Physiological Reviews*. **95**, 853–951 (2015).

69. S. B. M. Yoo, B. Y. Hayden, J. M. Pearson, Continuous decisions. *Phil. Trans. R. Soc. B.* **376**, 20190664 (2021).

70. S. Ferrari-Toniolo, F. Visco-Comandini, A. Battaglia-Mayer, Two Brains in Action: Joint-Action Coding in the Primate Frontal Cortex. *J. Neurosci.* **39**, 3514–3528 (2019).

71. P. R. Roelfsema, S. Treue, Basic Neuroscience Research with Nonhuman Primates: A Small but Indispensable Component of Biomedical Research. *Neuron*. **82**, 1200–1204 (2014).

72. A. Calapai, M. Berger, M. Niessing, K. Heisig, R. Brockhausen, S. Treue, A. Gail, A cage-based training, cognitive testing and enrichment system optimized for rhesus macaques in neuroscience research. *Behavior research methods*. **49**, 35–45 (2017).

73. M. Berger, A. Calapai, V. Stephan, M. Niessing, L. Burchardt, A. Gail, S. Treue, Standardized automated training of rhesus monkeys for neuroscience research in their housing environment. *J. Neurophysiol.* **119**, 796–807 (2018).

74. N. Fujii, S. Hihara, A. Iriki, Dynamic Social Adaptation of Motion-Related Neurons in Primate Parietal Cortex. *PLoS ONE*. **2**, e397 (2007).

75. R. Falcone, R. Cirillo, S. Ferraina, A. Genovesio, Neural activity in macaque medial frontal cortex represents others' choices. *Scientific Reports*. **7** (2017), doi:10.1038/s41598-017-12822-5.

76. O. Dal Monte, C. C. J. Chu, N. A. Fagan, S. W. C. Chang, Specialized medial prefrontal–amygdala coordination in other-regarding decision preference. *Nat Neurosci*. **23**, 565–574 (2020).

77. K. Yoshida, N. Saito, A. Iriki, M. Isoda, Representation of Others' Action by Neurons in Monkey Medial Frontal Cortex. *Current Biology*. **21**, 249–253 (2011).

78. R. Báez-Mendoza, C. R. van Coeverden, W. Schultz, A neuronal reward inequity signal in primate striatum. *Journal of Neurophysiology*. **115**, 68–79 (2016).

79. S. Ballesta, J.-R. Duhamel, Rudimentary empathy in macaques' social decision-making. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 15516–15521 (2015).

80. M. J. Osborne, A. Rubinstein, *A Course in Game Theory* (The MIT Press, Cambridge, Mass, First Edition, Twelf edition., 1994).

81. P. Grassberger, Finite sample corrections to entropy and dimension estimates. *Physics Letters A*. **128**, 369–373 (1988).

82. R. Steuer, J. Kurths, C. O. Daub, J. Weise, J. Selbig, The mutual information: detecting and evaluating dependencies between variables. *Bioinformatics*. **18**, S231–S240 (2002).

83. S. Pethel, D. Hahs, Exact test of independence using mutual information. *Entropy*. **16**, 2839–2849 (2014).

Acknowledgements

We thank Elisheba Crecca, Tarana Nigam and Roberta Nocerino for their help with the data collection, and Daniela Lazzarini, Janine Kuntze, Sina Plümer and Klaus Heisig for technical support.

Funding

This work was supported by a funding from the Ministry for Science and Education of Lower Saxony ("Top Level Research in Lower Saxony") and the Volkswagen Foundation through the "Niedersächsisches Vorab", https://www.volkswagenstiftung.de/en/funding/niedersaechsisches-vorab (JF, AG, ST, IK). Additional support was provided by the Leibniz Association through funding for the Leibniz ScienceCampus Primate Cognition, https://www.primate-cognition.eu (JF, AG, ST, IK). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

JF, ST and IK wrote the key grants to support his study. SM, AMU, AG, ST and IK conceptualized and designed the task. SM implemented the task and conducted the experiments. SM and AMU analyzed the data. SM, AMU and IK prepared figures. SM, AMU and IK wrote the initial version of the manuscript. All authors discussed and interpreted the findings, and revised the manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability

The code and datasets generated and analyzed in the current study are available from the corresponding author on reasonable request, and will be uploaded to Open Science Framework data repository (https://osf.io/), pending scientific review.

Supplementary Materials for

Human and macaque pairs employ different coordination strategies in a dyadic decision game with face-to-face action visibility

Sebastian Möller, Anton M. Unakafov, Julia Fischer, Alexander Gail, Stefan Treue, Igor Kagan

This PDF file includes:

Supplementary Materials and Methods

Written instructions read by human subjects

Instructions read to human subjects by experimenter

Abbreviations

Figs. S1 to S7

Tables S1 to S3

Supplementary Materials and Methods

Written instructions read by human subjects

Description of the study

This study is the investigation of the behavioral correlates of social decision making while playing a game with a partner. Every day we have to make decisions that depend not only on our own needs and goals but also on the needs and goals of others: for instance, while working on a project with colleagues, planning vacation with friends and family, getting on the bus or shopping groceries, etc. With the help of the task presented in this study we will investigate how people make such decisions.

The course of the study

You will complete one session of a decision making task on the computer together with your partner. You will have to choose one of the two circles presented to you. Your partner will have to perform the same task on his/her side of the touchscreen. You will have to decide and respond quickly. If either you or your partner are too slow, the trial will be aborted without any reward. Your reward will depend on your own and your partner's choice. After the decision is made, you both will receive different auditory feedbacks, denoting your reward and the reward of your partner. Please do not talk to your partner during the session. After the session, we will ask you several questions about the experiment.

Instructions read to human subjects by experimenter

1) You rest both hands on the gray board at the two round objects (touch sensors).

2) A central touch target will appear.

3) Move your right hand to the target and hold. Use the right hand during the entire session.

4) While the target brightens up keep holding your finger on the target.

5) Then two colored choice targets appear and the central target disappears.

6) Make your choice and touch the chosen target within 1.5 seconds.

7) All touched targets will brighten up, keep holding until the targets disappear.

8) Please note that both selected targets will brighten up:

   In case both players selected the same target only that single target will brighten;

   In case both players selected different targets, both targets will brighten.

9) Now two streams of auditory beeps will signify the earned reward for each player: each beep corresponds to a few cents. Please try to learn the sound related to your reward during the training trials.

10) While the audio plays move the hand back to the two touch sensors.

11) Go to 1.

Abbreviations

AR   Average reward

BoS   Bach-or-Stravinsky game (also known as the Battle of the Sexes)

DCR   Dynamic coordination reward

MI   Mutual information

MIS   Mutual information between side choices

MIT   Mutual information between target color choices

(i)PD   (iterated) Prisoner's dilemma

RT   Reaction time

SOC   Share of own choices

SLC   Share of left choices

Supplementary Figures

Supplementary Figure S1. Exemplary development of choice behavior in macaque pair TE over 6 sessions.
(A, C) The running average of the share of own choices from an early **(A)** and a late session **(C)** of pair TE. **(B, D)** The running average of the share of objective left choices from an early **(B)** and a late session **(D)** for the same pair. Conventions are the same as in Figure 1E,F. The pair shifted from each selecting their individually preferred target to mostly jointly selecting the objective left side. **(E-J)** The change in aggregate measures of choice behavior and coordination over the course of 6 sessions for the last 200 trials of each session. There was a clear development from selecting own targets to coordination by virtue of converging on mostly selecting the target on the same side of the transparent display. This is also reflected in the gradual increase of the target MI and the gradual decrease of the side MI. The zero DCR values indicate no significant dynamic coordination.

Supplementary Figure S2. Performance of naïve monkey pairs. (A-E) Data sorted as in Figure 2 F-J, in early sessions. **(F)** Scatter plot of the average share of left choices for all pairs between early and late sessions; note two groups, those that converged on selecting the same side in the upper right (same pairs clustered around an SOC of 0.5, see Fifure 2F), and those that converged on selecting the same target around the center of the plot (same pairs clustered at the top left of Figure 2F). **(G)** Paired training increases average joint reward in naïve monkey pairs. Scatter plot of the average reward for each macaque pair for an early versus a late session. Wilcoxon signed rank test showed that across all 9 pairs the average reward increased from early (median 2.75) to late sessions (median 3.29), Z: -2.67, p < 0.0077, r = -0.89. Note that 1 is the minimal possible average joint reward and 3.5 is the maximum. In all but one pair (TC, open circle), training increased the efficiency of coordination significantly (p < 0.05, Fisher's exact test of the proportion of coordinated versus uncoordinated trials in the early and the late session).

Supplementary Figure S3. Macaques can be trained to take turns when paired with a human confederate. Same as the Figure 4, for monkey C. **(A)** Early confederate-training session, **(B)** late confederate-training session. The reaction time difference histograms of the late session show that in blocks when the pair selected the macaque's non-preferred target (red curve), the monkey acted slower than the confederate, waiting to see the human's choice before committing to its own choice (t-test: coordination on red (mean: -142 ms, SD: 138, N: 86) vs. blue (mean: 0.5 ms, SD: 119, N: 108), t(168.4): -7.6, p: 2*10[-12]). **(C)** Monkey C also ceased to take turns when an opaque barrier was placed over the region of the transparent display containing touch targets (gray background in C) and mainly selected his own target. The reaction time difference histograms show that in the transparent condition the behavior was similar to (B): the monkey waited for the human to act first in blocks of the monkey's non-preferred target (t-test: coordination on red (mean: -192 ms, SD: 180, N: 98) vs. blue (mean: 27 ms, SD: 130, N: 169), t(156.3): -10.6, p: 5*10[-20]). In the opaque condition the behavior was similar to (A): the monkey mostly selected his preferred target and there was no RT difference dependence on human's choices (note the overlap between dashed blue and magenta curves).

Supplementary Figure S4. Monkeys gradually learn to follow a human confederate's turn-taking behavior. (A-F) The development of aggregate measures of choice behavior and coordination for monkey F (same conventions as Figure S1 E-J). Note the gradual increase in average rewards, and the overall slow changes in the first 9 sessions as compared to the steeper increases in the MIs and the dynamic coordination reward (DCR) in the last 6 sessions. **(G-L)** Corresponding data for monkey C.

Supplementary Figure S5. Confederate trained macaques. (A-F) The development of aggregate measures of choice behavior and coordination for monkey pair FC after confederate training (same conventions as Figure S1 E-J). Note sessions FC-2 and FC-5, which show a more equal SOC, and average reward between the two monkeys, as well as both non zero mutual information for side ad target, and especially significant DCR values.

Supplementary Figure S6. Correlations between the probability to see the partner's choice p_see and the likelihood to select the partner's target p_other. The left column shows the correlations for different sessions of the same pair of macaques (FC) before (FC naïve), during and after the confederate training, as well as different human pairs. The middle and right columns show p_see versus p_other for one exemplary session for each group. Note that training with a confederate increased magnitude of the correlation (Confederate training) and this carried into the post-confederate training sessions of the two macaques (FC trained). Tables S2 and S3 contain the statistical information for each individual correlation.

Supplementary Figure S7. Reaction time changes around choice combination switches. The graphs show the reaction time transitions around 6 specific choice switches, for agent A (monkey F) in red and for agent B (monkey C) in blue. Each curve shows the mean reaction time for 3 trials before, at the actual switch trial and 3 trials after the switch, the shaded areas show 95% confidence intervals around the mean. R – red, B – blue, M – magenta. The choice transitions are R→M: from both agents selecting red to each selecting own color; B→M: from both selecting blue to each selecting own color; R→B and B→R: from both selecting one color to both selecting another color (seamless transition, rare in macaques, see Figure 8C); as well as the respective reverse transitions. The choices of agent B are indicated on the top, the choices of agent A on the bottom. Note that both macaques speed up when transitioning from coordinating on their non-preferred color to challenging (R→M for agent B, and B→M for agent A), and also when transitioning from challenging to coordinating on their preferred color (M→B for agent B, and M→R for agent A). Similarly, both macaques slow down when transitioning from coordination on their preferred color to challenging (B→M for agent B, and R→M for agent A) and when transitioning from challenging to coordination on their non-preferred color (M→R for agent B, and M→B for agent A). In rare seamless switches, monkeys speed up on coordination on the preferred color and slow down for transitions to the non-preferred color.

Supplementary Tables

Pair ID	N solo trials	Solo trials: share of own choices A [%]		Solo trials: share of own choices B [%]		Choice preference (own vs. other's) in last 200 dyadic trials		RT A – RT B, coordinated trials			RT A – RT B, non-coordinated trials			Test on coordinated vs non-coordinated RT differences		
		all trials	last 25	all trials	last 25	A	B	mean	SD	N trials	mean	SD	N trials	df(corr.)	t	p
1	100	43	48	54	60	balanced	balanced	-110.4	90.8	306	-89.7	76.2	94	181.25	-2.19	**2.95E-02**
2	100	93	**100**	96	**100**	balanced	balanced	72.4	85.1	383	64.6	63.7	17	18.62	0.48	6.33E-01
3	100	86	**100**	55	52	balanced	balanced	130.5	116.1	263	60.2	114.6	90	155.99	5.01	**1.48E-06**
4	100	92	**100**	99	**100**	own	own	n/a	n/a	0	-21.3	52.3	400	n/a	n/a	n/a
5	100	98	**100**	95	**100**	own	balanced	-7.0	57.4	121	-30.5	52.3	279	210.32	3.87	**1.47E-04**
6	100	92	**100**	87	**92**	balanced	balanced	-7.3	77.9	396	104.3	70.0	4	3.08	-3.17	**4.88E-02**
7	100	26	4	50	48	balanced	balanced	-27.2	72.4	367	-32.8	96.3	33	35.33	0.33	7.46E-01
8	100	54	52	99	**100**	balanced	balanced	-177.6	80.3	377	-56.4	114.6	23	23.34	-5.00	**4.52E-05**
9	100	97	**100**	93	**100**	balanced	balanced	5.9	82.9	320	-5.1	77.3	80	128.34	1.13	2.61E-01
10	50	86	**96**	24	16	own	other's	-36.4	73.7	300	156.2	n/a	1	n/a	n/a	n/a
11	50	46	48	2	0	balanced	other's	-43.2	66.4	176	-33.2	55.5	124	289.36	-1.42	1.58E-01
12	50	34	64	61	64	balanced	balanced	-22.5	125.9	276	-23.2	97.0	24	30.18	0.03	9.77E-01
13	50	89	**92**	60	44	balanced	balanced	106.2	109.2	239	75.1	84.6	61	116.53	2.41	**1.77E-02**
14	50	94	**100**	50	40	own	other's	-40.3	53.8	240	-14.2	61.3	60	83.09	-3.03	**3.28E-03**
15	100	1	0	53	40	other's	balanced	55.5	70.2	166	15.3	67.0	235	345.03	5.76	**1.84E-08**
16	100	58	**80**	20	0	own	other's	0.2	66.5	388	-31.0	142.8	12	11.15	0.75	4.67E-01
17	100	94	**100**	94	**100**	balanced	balanced	114.5	85.4	337	153.5	87.1	63	85.75	-3.27	**1.53E-03**
18	100	96	**96**	31	24	other's	own	-96.0	108.8	398	9.5	65.6	2	1.03	-2.26	2.60E-01
19	100	93	**100**	1	0	own	own	n/a	n/a	0	22.2	90.8	407	n/a	n/a	n/a

Table S1. Human pairs. Columns 2-8: No relationship between own target preference in solo trials and dyadic behavior. Twenty of 38 human subjects preferred their own target at the end of the solo training trials (last 25 trials own color selection >75%, marked by bold font). Percentage of choice for the own preferred target for both agents of each human pair, for all solo trials and for the last 25 solo trials. Pairs displaying dynamic coordination (turn-taking behavior) are designated by a gray background. Choice preference in dyadic trials: other's (SOC<25%), balanced (25%≤SOC≤75%), own (SOC>75%). Out of 10 pairs (20 subjects) that showed balanced turn-taking, 11 subjects showed a strong preference for own targets already in solo trials, and 9 did not. In the remaining 9 pairs (18 subjects), 7 subjects who had own preference in solo trials also showed it in dyadic trials, 4 subject who had other's preference in solo trials also showed it in dyadic trials, and 5 out remaining 7 subjects switched the preference from solo to dyadic trials, e.g. other's to own. Thus, the behavior in the solo trials is not predicting the behavior in the dyadic trials. This indicates that human participants did not simply carry their established preferences into the dyadic setting, but continued to explore and modify their strategies. **Columns 9-17. Reaction time differences and coordination. Nine of 19 human pairs showed significant differences between coordinated and non-coordinated trials' reaction time difference, indicating that seeing the other's action can help maintain coordination.** Comparison of trial-by-trial reaction time differences between the two agents in the pair (RT A – RT B), for coordinated and non-coordinated trials for all human pairs by t-test using Satterthwaite's approximation to allow for unequal variance. Note that pairs 4, 10, and 19 had too few trials for one of the trial types to calculate the test statistic (n/a). Pairs displaying turn-taking behavior are designated by a gray background. Bold font denotes correlations with a p<0.05. Green font shows pairs where absolute RT differences increased with coordination, purple – where absolute RT differences decreased with coordination.

Session	df	r(A)	r(B)	p(A)	p(B)	raw r(A)	raw r(B)	raw p(A)	raw p(B)
Prior to confederate training (naïve pair)									
FC-1	304	-0.544	0.169	**5.12E-25**	**0.002952**	-0.068	0.043	0.23521	0.45327
FC-2	365	0.236	-0.536	**4.62E-06**	**1.17E-28**	0.388	-0.114	**1.27E-14**	**0.028859**
FC-3	183	-0.351	0.200	**9.42E-07**	**0.006239**	-0.124	-0.062	0.092404	0.40478
FC-4	207	-0.376	-0.200	**2.00E-08**	**0.003778**	-0.128	0.218	0.064765	**0.001527**
FC-5	171	0.194	0.161	**0.010527**	**0.034044**	0.111	-0.014	0.14464	0.85860
FC-6	397	n/a	-0.096	n/a	0.054399	n/a	-0.039	n/a	0.44066
FC-7	314	-0.044	0.560	0.433	**1.68E-27**	0.024	0.144	0.67187	**0.010417**
FC-8	447	0.447	0.228	**1.67E-23**	**9.91E-07**	0.171	0.027	**0.00027874**	0.5692
During confederate training									
F-1	431		0.063		0.19125		0.079		0.10121
F-2	714		0.125		**0.000804**		0.108		**0.003798**
F-3	578		-0.053		0.20033		0.140		**0.000699**
F-4	80		0.040		0.72184		0.248		**0.024749**
F-5	547		0.044		0.30683		0.173		**4.56E-05**
F-6	317		0.280		**3.54E-07**		0.205		**0.000225**
F-7	540		0.141		**0.001023**		0.170		**6.82E-05**
F-8	582		0.077		0.063123		0.076		0.068242
F-9	547		0.260		**5.64E-10**		0.298		**9.42E-13**
F-10	479		0.544		**2.19E-38**		0.390		**6.99E-19**
F-11	473		0.820		**7.89E-117**		0.586		**3.44E-45**
F-12	525		0.829		**1.15E-134**		0.579		**1.40E-48**
F-13	594		0.815		**3.75E-143**		0.589		**7.59E-57**
F-14	554		0.867		**2.42E-169**		0.557		**1.10E-46**
F-15	446		0.868		**1.40E-137**		0.642		**2.38E-53**
C-1	652		0.161		**3.51E-05**		0.188		**1.37E-06**
C-2	544		0.311		**9.50E-14**		0.172		**5.18E-05**
C-3	320		0.482		**3.56E-20**		0.307		**1.78E-08**
C-4	455		0.145		**0.001896**		0.188		**5.03E-05**
C-5	412		0.588		**7.67E-40**		0.285		**3.63E-09**
C-6	476		0.697		**7.88E-71**		0.408		**1.46E-20**
C-7	581		0.670		**3.11E-77**		0.388		**2.42E-22**
C-8	228		0.715		**2.36E-37**		0.471		**4.42E-14**
After confederate training (trained pair)									
FC-1	371	0.806	0.834	**2.45E-86**	**1.24E-97**	0.516	0.603	**8.13E-27**	**3.13E-38**
FC-2	545	0.916	0.909	**3.57E-218**	**1.28E-209**	0.635	0.655	**4.15E-63**	**2.93E-68**
FC-3	625	0.753	0.518	**6.88E-116**	**2.65E-44**	0.423	0.321	**1.19E-28**	**1.76E-16**
FC-4	471	0.856	0.517	**9.73E-137**	**9.49E-34**	0.420	0.305	**1.20E-21**	**1.22E-11**
FC-5	683	0.707	0.654	**1.16E-104**	**9.32E-85**	0.562	0.430	**2.71E-58**	**3.53E-32**
FC-6	370	0.901	0.822	**3.33E-136**	**1.29E-92**	0.601	0.580	**7.55E-38**	**7.34E-35**

Table S2. Reaction time correlations in macaque pair FC. Correlation between the probability to see other's action and the selection of the other's target. Prior to confederate training (8 naïve sessions), monkeys F (agent A) and C (agent B) showed only relatively weak and inconsistent (both positive and negative) correlations between probability to see other's action (r(A) and r(B)) and the selection of the other's target. Correlation degree of freedom (df), correlation coefficients (r) and correlation p-values (p) for the running average over 8 trials as displayed in Figure S6, and for the raw non-smoothed data are shown. Bold font denotes correlations with p<0.05. n/a – not applicable: the correlation is undefined because at least one variable was constant (e.g. the share of other's choice 0). The few high correlations on smoothed data were mainly driven by the slow changes in the timecourse of the session. During confederate training both monkeys (F, C) developed strong and highly significant positive correlations between being able to see the confederate's actions and following the confederate: 15 sessions of monkey F and 8 sessions of monkey C during confederate training. Note the human confederate worked on position A, and is excluded here. Bold font denotes correlations with a p<0.05. After confederate training this pair continued to show strong correlations between seeing the other's choice and following: 6 sessions of monkeys F and C after confederate training. Note that even in the sessions without significant DCR in the last 200 trials (i.e. FC-1, FC-3, FC-4, FC-6), where monkeys largely converged on one color, the high positive correlations reflect brief bouts of competitive turn-taking or challenging.

Pair ID	df	r(A)	r(B)	p(A)	p(B)	raw r(A)	raw r(B)	raw p(A)	raw p(B)
4	398	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
19	405	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
15	399	-0.123	-0.110	**0.014003**	**0.026945**	0.067	0.176	0.1802	**0.000399**
11	298	-0.097	n/a	0.09230	n/a	0.040	n/a	0.48759	n/a
5	398	-0.009	-0.427	0.85640	**3.93E-19**	-0.028	-0.171	0.58233	**0.000598**
3	351	0.164	0.006	**0.001941**	0.91763	-0.012	0.054	0.82342	0.30902
1	398	-0.009	0.025	0.85616	0.62476	0.027	0.026	0.59697	0.60243
14	298	0.060	0.079	0.29914	0.17498	0.010	0.154	0.86265	**0.007371**
13	298	0.126	0.108	**0.029495**	**0.061976**	0.026	-0.037	0.65319	0.52015
2	398	0.243	0.394	**9.12E-07**	**2.54E-16**	0.095	0.124	0.05769	**0.012791**
12	298	0.523	0.456	**1.98E-22**	**8.08E-17**	0.268	0.330	**2.41E-06**	**4.90E-09**
16	398	0.046	-0.569	0.36308	**1.18E-35**	0.142	-0.214	**0.004470**	**1.64E-05**
7	398	0.053	0.287	0.29108	**4.94E-09**	0.001	0.066	0.97791	0.18541
9	398	0.126	-0.271	**0.011469**	**3.59E-08**	0.009	-0.025	0.85404	0.61258
17	398	-0.487	0.187	**3.10E-25**	**0.000171**	-0.054	0.038	0.28453	0.44647
8	398	-0.114	0.075	**0.022805**	0.13386	-0.036	0.055	0.47255	0.2706
6	398	-0.370	0.071	**2.04E-14**	0.15713	0.040	0.017	0.41953	0.73857
10	299	0.251	n/a	**1.07E-05**	n/a	0.221	n/a	**0.000109**	n/a
18	398	0.048	0.176	0.34150	**0.000399**	-0.000	0.094	0.99829	0.06041

Table S3. Reaction time correlations in humans. Correlation between the probability to see other's action and the selection of the other's target in 19 human pairs. Human pairs showed weak and inconsistent (both positive and negative) correlations between probability of seeing the other's choice and selecting the other's target. Correlation degree of freedom, correlation coefficients and correlation p-values for the running average over 8 trials, as displayed in Figure S6, and for the raw non-smoothed data are shown. (A) and (B) denote the positions of the agents. Pairs are sorted by the average joint reward in ascending order (cf. Figure 2). Pairs displaying dynamic turn-taking behavior are designated by a gray background. Bold font denotes correlations with a p<0.05. n/a – not applicable: the correlation is undefined because at least one variable was constant (e.g. the share of other's choice 0).