Cytoplasmic incompatibility between Old and New World populations of a tramp ant

Çiğdem Ün[1]°, Eva Schultner[1]°, Alejandro Manzano-Marín[2], Laura V. Flórez[3], Bernhard
Seifert[4], Jürgen Heinze[1], Jan Oettler[1]*

[1]Zoology/Evolutionary Biology, University of Regensburg, Universitätsstrasse 31, 93053
Regensburg, Germany

[2]Centre for Microbiology and Environmental Systems Science, University of Vienna,
Althanstraße 14, 1090 Vienna, Austria

[3]Institute of Organismic and Molecular Evolution, Evolutionary Ecology Department,
Johannes Gutenberg University Mainz, Hanns-Dieter-Hüsch-Weg-15, 55128 Mainz,
Germany

[4] Senckenberg Museum of Natural History Görlitz, Am Museum 1, 02826 Görlitz, Germany

°contributed equally

*corresponding author:
Jan Oettler
joettler@gmail.com

Author ORCIDs:
Çiğdem Ün: 0000-0002-0817-3630
Eva Schultner: 0000-0002-5069-9732
Alejandro Manzano-Marín: 0000-0002-0707-9052
Laura V. Flórez: 0000-0002-0761-3729
Jürgen Heinze: 0000-0002-9403-5946
Jan Oettler: 0000-0002-8539-6029

Classification: Biological Sciences: Evolution

Key words: *Wolbachia*, social insects, speciation, endosymbiont, antibiotics

Contributions
CÜ performed the crosses and qPCR
ES analyzed and visualized the data
AMM assembled *Wolbachia* genomes, performed comparative genomic analyses,
calculated the phylogeny and annotated CI genes
LVF conducted FISH analyses
BS revised the taxonomic status of the populations
JH acquired funding and reviewed and edited the manuscript
JO conceived and supervised the study
JO, ES and CÜ wrote the original draft of the manuscript
All authors revised and approved the final version of the manuscript

Abstract
As we enter the Anthropocene, the evolutionary dynamics of species will change
drastically, and as yet unpredictably, due to human activity. Already today, increases in
global human traffic have resulted in the rapid spread of species to new areas, leading to
the formation of geographically isolated populations. These go on to evolve in allopatry,
which can lead to reproductive isolation, and potentially, the formation of new species.
Surprisingly, little is known about such eco-evolutionary processes in ants, even though
numerous invasive ant species are globally distributed in geographically isolated
populations. Here, we describe the first case of cytoplasmic incompatibility (CI) between
populations of a cosmotropic distributed tramp ant with Asian roots, *Cardiocondyla*
obscurior, which has acquired a novel *Wolbachia* strain in the New World. Our study
uncovers the first symbiont-induced mechanism of reproductive isolation in ants,
providing a novel perspective on the biology of globally distributed ants.

61
Introduction
Ants are the most abundant group of insects on earth, and numerous ant species are
classified as highly invasive on a global scale. Their distribution is, on the one hand,
strongly facilitated by humans. This is evidenced, for instance, by the rapid worldwide
spread of the Argentine ant and the fire ant from their origin in South America [1,2]. On the
other hand, the particular biology of ants, characterized by reproductive division of labor
between highly fecund queens and sterile workers, allows the establishment of large
populations from just one founding propagule. Introduced populations thus go through a
genetic bottleneck, reducing diversity within the population while at the same time
increasing diversity between populations [3]. Such populations then evolve in allopatry, and
differences caused by genetic drift may be further amplified over time by selection,
eventually leading to reproductive isolation. Reproductive isolation, in turn, together with
the evolution of mating systems, underlies the formation of new species and, hence, the
vast diversity of life. By allowing ants (and other organisms) to establish allopatric
populations around the world, humans have inadvertently created a valuable scenario for
studying species evolution in real time.
78
Reproductive isolation between populations that live in allopatry can evolve in the form of
post-mating mechanisms of reproductive isolation (PoMRI). Importantly, PoMRI can result
from Bateson–Dobzhansky–Muller (BDM) incompatibilities, in which two parental loci that
have diverged cause hybrid sterility or inviability. Extending such negative epistatic
interactions to the host's microbiome adds three mechanisms that can reduce hybrid
fitness: hybrid susceptibility (i.e. inferior immune systems and higher pathogen loads in
hybrids), hybrid autoimmunity (i.e. upregulation of immune functions in hybrids because
of negative epistasis), and cytoplasmic incompatibility (CI, i.e. hybrid embryo death
caused by incompatibilities between members of the parental microbiomes) [4]. PoMRI
mechanisms can operate at any developmental stage and can directly affect the fitness of
both hybrid offspring and mated parental females. The best-known cases of PoMRI caused
by bacteria involve *Wolbachia*, the enigmatic evolutionary omnia rotundiora comprising
endosymbionts and endoparasites [5], which infects at least 20% of the world's insect
species, with estimates reaching up to 66% [6]. *Wolbachia* is famous for manipulating
maternal transmission routes, most likely driven by an ancestral genomic association with
a prophage [7]. *Wolbachia* strains capable of reproductive manipulation can sweep through

a population by inducing CI, or by favoring vertical maternal transmission via interference
with the host's development, resulting in parthenogenesis, feminization, or male death [8].
97
Although numerous ant species are highly invasive and form allopatric populations on a
global scale, nothing is known about PoMRI mechanisms involving endosymbiotic bacteria
in these ecologically dominant insects. In particular, the functional role of *Wolbachia* in ant
biology has thus far remained elusive, in spite of its omnipresence [9]. The globally
distributed tramp ant *Cardiocondyla obscurior* has spread from its putative origin in
Southeast Asia and today occurs in fruit tree plantations, city parks and other disturbed
habitats in the tropics and sub tropics. In addition to *Wolbachia* populations of *C.*
obscurior carry a main bacterial endosymbiont, *Cand.* Westeberhardia, which exhibits
characteristics of an obligate symbiont (Klein et al. 2015). Populations of *C. obscurior* from
Brazil (BR) and Japan (JP) show remarkable genomic heterogeneity and differ in a number
of phenotypic traits [10]; nevertheless, mating between queens and males from the two
populations readily takes place in the laboratory. However, a phenotype indicative of
PoMRI was found in hybrid crosses between BR and JP populations, with BR queens
mated to JP males exhibiting shorter lifespans and lower fecundity than BR queens mated
to BR males [11]. At the time, this was attributed to BDM incompatibility of fast evolving
sexual traits such as seminal fluid proteins, as suggested by an up-regulation of genes
involved with immune response in outbred BR queens [11].
115
Here, we show that the negative effect on queen fitness is in fact caused by incompatible
strains of *Wolbachia* harbored by the two populations. By reducing *Wolbachia* levels in
males using antibiotic treatment, we demonstrate that queen fecundity can be rescued.
Genomic analyses of the *Wolbachia* strains isolated from the two populations reveal that
both strains belong to *Wolbachia* supergroup A, which typically contains strains capable of
reproductive manipulation. Surprisingly, CI is only induced by the putatively ancestral
strain, and this may be linked to functional differences in CI-associated loci. By discovering
the first case of CI between two ant populations, our study provides the first description of
functional *Wolbachia* infection dynamics in ants and presents a novel perspective on the
biology of globally distributed ants.
126
Results
Unidirectional hybrid incompatibility
We verified the results from [11] by crossing queens and males from laboratory colonies of
BR and JP populations collected in 2009 and 2010, respectively [10]. We conducted this
experiment once in 2015 with the same outcome (Figure S1, Supplement Methods) and
again in 2017/18, the results of which are presented here. As expected, compared to all
other crosses the combination BR x JP (queen x male) resulted in lower fecundity as
estimated by mean weekly egg number over the first six weeks after initiation of egg
laying, regardless of whether the male was infected with *Cand.* Westeberhardia or not
(Figure 1, Kruskal Wallis rank sum test, $X^2 = 78.04$, df=6, p<0.001, see Table S1 for
Bonferroni-Holm corrected pairwise p-values). Similar results were obtained when
maximum weekly egg numbers were compared (Figure S2, Table S2, Kruskal Wallis rank
sum test, $X^2=81.10$, df=6, p<0.001). Strikingly, the effect was even stronger when using
males naturally uninfected with *Cand.* Westeberhardia, ruling out the possibility that this
endosymbiont plays a role in the induction of reproductive isolation.

We monitored a subset of inter-population crosses for an additional 6 weeks (12 weeks in
total). Of the 28 BR x JP colonies that were monitored for 12 weeks, seven failed to
produce diploid offspring (queen/worker pupae), and four produced no pupae at all. We
dissected the spermathecae of three of the seven male-only producing queens, and all
contained sperm. All of the JP x BR colonies that were monitored for 12 weeks produced
pupae, and they produced significantly more pupae than BR x JP (Figure 2A; Kruskal-
Wallis rank sum test, X^2=27.90, df=2, Bonferroni-Holm corrected p<0.001). Furthermore,
sex ratios of produced pupae were significantly more male-biased in BR x JP colonies
compared to JP x BR colonies (Figure 2B; binomial GLM: $T_{2,52}$=32.81, Bonferroni-Holm
corrected p<0.001), an effect caused entirely by the lack of developing females (Figure
S3; Kruskal Wallis rank sum test, X^2=4.04, df=2, p=0.13). Together, these data show
consistent incompatibility between BR queens and JP males.

Cytoplasmic incompatibility
Cytoplasmic incompatibility (CI) occurs when an endosymbiont favors its transmission by
making infected males incompatible with uninfected females, while infected females can
mate with both infected and uninfected males. A necessary test to verify CI is 'curing' the
host of the putative CI-causing bacteria using antibiotic treatment, which should result in
the 'rescue' of the wild-type phenotype.

We fed JP^{rif+} colonies the antibiotic rifampicin diluted in honey every other week for 10
weeks, which resulted in the complete loss of the main endosymbiont *Cand.*
Westeberhardia (Figure S4; Mann-Whitney-U-Test, W=34, df=1, p<0.001) and a significant
reduction of *Wolbachia* levels in workers (Figure S4; Mann-Whitney-U-Test, W=95, df=1,
p<0.01). Treating colonies with the antibiotic tetracycline did not have a negative effect on
Wolbachia (Figure S5; Mann-Whitney-U-Test, X^2=24, df=1, p=0.052). While *Cand.*
Westeberhardia is permanently eradicated in subsequent generations the negative effect
of rifampicin on *Wolbachia* is transient, with levels returning to normal 6 months after
termination of treatment (Figure S6; *Cand.* Westeberhardia: Mann-Whitney-U-Test, W=64,
df=1, p<0.001; *Wolbachia*: Mann-Whitney-U-Test, W=45, df=1, p=0.19). Five of 28
rifampicin-treated colonies succumbed to the treatment and died during the 10-week
period, while all 28 control colonies survived.

Mating JP^{rif+} males collected shortly after ceasing antibiotic feeding with BR queens
resulted in significantly rescued fecundity, although mean and maximum weekly egg
numbers did not quite reach levels of BR and JP inbred queens (Figure 1, Table S1). This
rescue effect was evident regardless of whether JP^{rif+} males came from *Cand.*
Westeberhardia infected or uninfected colonies. Rescued BR x JP^{rif+} colonies also
produced significantly more pupae over a 12-week period than their non-rescued BR x JP
counterparts (Figure 2A; Kruskal-Wallis rank sum test, X^2=27.90, df=2, Bonferroni-Holm
corrected p=0.027), and sex ratios among pupae were as female-biased as in JP x BR
crosses (Figure 2B; binomial GLM: $T_{2,52}$=32.81, Bonferroni-Holm corrected p=0.011).
Strongly reduced *Wolbachia* levels in JP^{rif+} males (Figure 3, Mann-Whitney-U-Test, W=163,
df=1, p<0.001) support the role of *Wolbachia* in inducing hybrid incompatibility.

We confirmed unidirectional CI with males from laboratory colonies of a second Asian
population collected in Taiwan (TW), which is infected with a *Wolbachia* strain that
amplifies a *wsp* sequence (coding for the hypervariable *Wolbachia surface protein*)
identical to those from JP samples (Figure S7). We set up crosses with queens from BR (BR
x TW) and JP (JP x TW), and the reciprocal combination (TW x JP). Again, only the
combination BR x TW resulted in reduced fecundity, both when mean weekly egg
numbers (Figure S8; Kruskal Wallis rank sum test, X^2=9.96, df=2, p<0.01, see Table S3 for
Bonferroni-Holm corrected pairwise p-values) and maximum weekly egg numbers over six
weeks (Figure S8; Kruskal Wallis rank sum test, X^2=10.19, df=2, p<0.01, see Table S4 for
Bonferroni-Holm corrected pairwise p-values) were compared.
198
Wolbachia genomics and phylogeny
We assembled the *Wolbachia* genomes from BR (*w*Cobs-BR) and JP (*w*Cobs-JP)
populations of *C. obscurior* using Illumina HiSeq2000 100-bp reads of 200 and 500-bp
insert paired-end libraries [10]. The genomes of both *Wolbachia* were assembled to 148 (BR)
and 182 (JP) scaffolds (Table 1). Both strains showed similar genome sizes and G+C
contents, with an estimated 1,013 (BR) and 1,058 (JP) predicted protein-coding genes. An
analysis on shared genes revealed that both strains share a core of 907 proteins, with 85
unique to *w*Cobs-BR and 109 unique to *w*Cobs-JP. According to Bayesian phylogenetic
analyses, both *Wolbachia* strains belong to supergroup A, which together with
supergroup B contains strains capable of reproductive manipulation. *w*Cobs-JP was found
to be more closely related to *w*Ha and *w*CauA strains, while *w*Cobs-BR was
phylogenetically closer to *w*Au and *w*Mel (Figure 4A). A synteny analysis revealed that
both *w*Cobs-JP and *w*Cobs-BR strains keep general synteny, but with several
rearrangements (Figure 4B). This is consistent with previous observations of group A
Wolbachia strains [12].
214
Wolbachia phenology
To compare infection titers between populations we performed qPCR of the *Wolbachia*
gene *Cytochrome c oxidase subunit 1* (*coxA*) against the ant housekeeping gene
elongation factor 1-alpha 1 (*EF1*). JP queens had higher relative *w*Cobs-JP titers than BR
queens infected with *w*Cobs-BR (Figure S9, Mann-Whitney-U-Test, W=23, df=1, p=0.043).
Likewise, JP workers showed higher infection titers with *w*Cobs–JP compared to BR
workers infected with *w*Cobs-BR (Figure S9, Mann-Whitney-U-Test, W=16, df=1, p=0.017).
222
223 To test for morph-bias in infection we compared *coxA* expression in adult queens,
224 workers, winged males and wingless males using qPCR. *coxA* levels did not differ between
225 sexes and were consistently higher in winged morphs (queens, winged males) compared
226 to wingless morphs (workers, wingless males) (Figure S10, Kruskal Wallis rank sum test,
227 X^2=19.53, df=3, p<0.001).
228
To measure the potential cost of *Wolbachia* infection and to test for a selective loss in
workers, we compared infection levels of workers and queens over time. In contrast to
Cand. Westeberhardia [13], *Wolbachia* levels quickly reached a plateau in both queens and
workers. There was no indication that individuals lose *Wolbachia* over the first few weeks
of their lives, which typically last for up to three (workers) or six months (queens) (Figure
S11, workers: ANOVA, $F_{4,26}$= 2.69, p=0.059; queens: ANOVA, $F_{5,50}$= 15.95, p<0.001).

CI mechanism

CI usually results in the failure to produce viable zygotes but how CI operates functionally is still not clear. The two championed models are based on 1) a sperm modification step during spermatogenesis which is reversed by a rescue factor or 2) the transfer of a toxic product via the sperm and subsequent binding to a female-derived antidote [14]. The current genetic view for CI in *Drosophila* favors a two-by-one model [15], involving two adjacent genes (coined *cifA* and *cifB*) located in a region of prophage origin in *w*Mel [16], which contains several genes with eukaryote function [17]. Expression of both *cifA* and *cifB* in males is required to cause CI, while expression of *cifA* rescues embryo mortality in females [14].

We found orthologues of *cifA* and *cifB* in *Wolbachia* strains from BR and JP. In the latter, we found two additional paralogues. Through Bayesian phylogenetic inference, and following the classification scheme of [16], we found that all copies of *cifA* and *cifB* belong to their corresponding type. The *cifA* and *cifB* genes found in NODE_002 of *w*Cobs-JP are closely related to those found in *w*Cobs-BR and in the cryptic phage of strain *w*Sol isolated from the fig wasp *Ceratosolen solmsi* (Figure 5A, B). The *cifA* and *cifB* genes found in NODE_063 of *w*Cobs-JP show a more distant relationship to their homologues in *w*Cobs-BR. Following [16], we performed a protein remote homology search with the webserver of HHpred's v3.2.0 webserver [18]. While the *cifA* gene of *w*Cobs-BR contained a DUF249 domain, neither *cifA* homologues in *w*Cobs-JP showed homology to known domains (Figure 5C). In contrast, putative domains were found in all *cifB* genes. These shared a putative nuclease domain (PD-(D/E)XK nuclease/DpnII-MboI) found in other type I *cifB* genes. The *cifB* homologue found in NODE_063 in *w*Cobs-JP shared most domains with the *cifB* homologue in *w*Cobs-BR, with an additional peptidase domain (Peptidase_C58; PF03543).

Wolbachia in males

As mentioned above, *Wolbachia*-induced CI has often been linked to sperm modification during spermatogenesis [19] or the transfer of a CI toxin along with or in the sperm. The presence of *Wolbachia* in the testes is thus expected, although their specific localization in spermatocytes or spermatids might not be a prerequisite for inducing CI, as has been observed in the wasp *Nasonia vitripennis* [20]. We used fluorescence in situ hybridization to localize *Wolbachia* in the abdomens of male *C. obscurior*, confirming the presence of the bacteria throughout different tissues including the testes in both JP (Fig. 6A-C) and BR (Figure 6D) individuals. *Wolbachia* was also observed co-infecting bacteriocytes dominated by *Cand*. Westeberhardia, yet in significantly lower densities (Figure 6C).

Discussion

Insect-*Wolbachia* interactions can range from beneficial to detrimental and from facultative associations to rare cases of ultimate mutualisms where hosts have higher fitness with *Wolbachia* than without [5]. This prevalence and diversity of *Wolbachia* associations predictably also occurs in ants; however, our knowledge of ant-*Wolbachia* interactions is limited [9]. In the few ant-*Wolbachia* associations studied so far, infection is facultative (e.g. [21-23]), indicating ongoing arms races between hosts and symbionts [24]. One particularly obvious question to ask is whether *Wolbachia* has the power to control sex

allocation, a trait shaped by kin selection in social Hymenoptera [25]. In this context, facultative *Wolbachia* infections have been reported to cause transient changes in sex allocation (Pontieri et al. 2016), or to have no effect [22,26]. Similarly contrasting results have been obtained regarding effects on colony productivity [22,27]. It has furthermore been speculated that *Wolbachia* can bias caste ratios towards the production of queens [21,28], but current data are scarce and provide little support for this hypothesis [9]. By providing the first functional description of an ant-*Wolbachia* association involving two strains of *Wolbachia*, our study lays the groundwork for in-depth investigations into how *Wolbachia* affect ant reproductive biology and fitness.

Wolbachia clearly play a significant role in the biology of *C. obscurior*. First, all screened individuals were infected regardless of population, caste or sex. *Wolbachia* titers were higher in queens and winged males compared to workers and wingless males, a difference that is likely associated with the relative amounts of reproductive tissue as *C. obscurior* workers lack ovaries and winged males have larger testes than wingless males [29]. Second, rifampicin treatment merely reduced *Wolbachia* titers, and this effect was reversible. This was not the case for *Cand.* Westeberhardia, which was permanently cleared by rifampicin treatment. Third, *Wolbachia* titers did not decline with age in workers. *Wolbachia* infection can be costly and selection may thus favor reduced infection loads in sterile workers, as these do not contribute to vertical transmission [21,28]. Such a mechanism appears to be acting on *Cand.* Westeberhardia, where titers in workers (but not queens) decreased significantly with age, indicating that while the host may benefit from infection during development or in the adult egg-laying queen stage, the symbiont is not required [13]. Consistently high titers in workers indicate that *Wolbachia* do not have deleterious effects on worker performance in *C. obscurior*, unlike *Wolbachia* in *Acromyrmex* ants [30]. To what degree the population-specific strains differ in their effect on host phenotype remains unclear. Higher *Wolbachia* titers in JP individuals suggest that the *w*Cobs-JP strain may be better integrated into the host's biology. Alternatively, these differences may be driven by population-specific competition with *Cand.* Westeberhardia over limited resources.

Sex allocation in *C. obscurior* is female-biased [31] and queens presumably have full control over male production, as investment into males increases in the presence of competing queens [32-34], Accordingly, we found no indication that *Wolbachia* influences sex allocation. Queens from non-CI (JP x BR) crosses produced few males in the first 12 weeks of their lives. This may be a consequence of the short observation period, equaling ~50% of the average queen lifespan [31]. The sex ratios produced by CI (BR x JP) crosses showed a bimodal distribution, with several queens matching sex ratios produced by non-CI queens (albeit with lower overall productivity), while others produced only males. On average, however, CI queens did not produce more males than non-CI queens. There was also no indication that *Wolbachia* altered caste ratios towards production of queens. Concordantly, a previous study comparing investment into sexuals and workers did not find differences between the BR and JP populations [31]. Testing directly for a caste-biasing effect of *Wolbachia* requires complete elimination of the bacterium. While this cannot be done without also killing the ant host, preliminary trials indicate that cross-infection of the two *Wolbachia* strains is possible. In the future, such manipulations may help clarify whether *Wolbachia* have caste-manipulating power in *C. obscurior*.

329
Global exchange of stowaway ants leading to secondary contact between allopatric
populations is common (e.g. in the tropical fire ant *Solenopsis geminata* [35], and the little
fire ant *Wasmannia auropunctata* [36]) and thus the presented scenario, albeit conceived in
the laboratory, is not an unrealistic one. A previous study emphasized the role of BDM
incompatibility between allopatric *C. obscurior* populations in causing a CI-like phenotype
[11]. *Wolbachia*-induced incompatibilities were ruled out because of identity of a partial
sequence of *wsp* extracted from seven individuals (three from BR colonies collected in
2004, four from JP colonies collected in 2005) using the wsp81F/wsp691R primer
combination [37]. Gene expression analyses of allopatrically and sympatrically mated BR
queens pointed towards elevated immune responses and other changes that mimicked a
CI-like phenotype. In *Drosophila*, *Wolbachia* can manipulate expression of key regulatory
genes during spermatogenesis, which leads to embryonic lethality and a CI-like
phenotype [38]. Thus, what appeared to be BDM incompatibility may have been caused by
Wolbachia inducing a CI-like phenotype via interference with queen gene expression. The
alternative and more parsimonious explanation for the CI-like phenotype based on the
results from the present study is a classic unidirectional *Wolbachia*-driven CI effect
involving incompatible *Wolbachia* strains. This also explains why JP x BR crosses did not
elicit a CI phenotype. The overexpression of immunity-related genes in allopatrically
mated BR queens detected by Schrempf et al. 2015 may simply reflect physiological
responses to aborted oogenesis, as opposed to responses to ejaculate quality, or to
injuries caused by non-matching male genitalia. Furthermore, we recently found that some
BR colonies were infected with an additional *Wolbachia* strain displaying partial *wsp*
sequences identical to those of *w*Cobs-JP (data not shown), indicating horizontal
transfection in our rearing facilities, especially in older laboratory colonies.
354
The mechanism of CI in *C. obscurior* appears to align along the *Wolbachia cif*-like gene
axis described in *Drosophila* and *Culex* [15,16,39-41]. This genetic region, which is of prophage
origin and capable of manipulating the eukaryote host, showcases the importance of
using a hologenomic approach to study evolution [42]. *w*Pip of *Culex pipiens* and *w*Yak of *D.*
yakuba have similar prophage-derived regions containing two syntenic pairs of paralog
genes (termed *cidA/cidB* and *cinA/cinB)* [43]. *cidA/cidB* have been suggested to induce and
rescue CI via a toxin-antidote mechanism where *cidB* acts as a modifier and *cidA* as a
rescuer [41]. Surprisingly, the paralog genes *cidA/cifA* in *w*Pip and *w*Mel have different
biochemical functions: both contain ankyrin-like repeats for protein-protein interaction
with eukaryotic DNA [41] but *cifA* is not a putative deubiquitinase [14], as postulated for *cidA*
[39]. Both *Wolbachia* strains isolated from *C. obscurior* have orthologues of *cifA* and *cifB* but
only the *w*Cobs-JP strain has an additional *cifA/B*-like pair, pointing towards a role for the
latter pair in inducing reproductive isolation. Whether the second *cifA/B* containing region
was lost in *w*Cobs-BR, or whether it stems from a duplication event and subsequent
divergence in *w*Cobs-JP as has been described for the *w*AlbB strain of *Aedes albopictus*
[44], remains to be shown. Alternatively, functional differences in *cifA/B* orthologues
between *w*Cobs-JP and *w*Cobs-BR, or an interaction of the two *cif*-like gene products in JP
males could be causing the CI. Irrespective of the exact CI mechanism, it remains unclear
why a mere reduction of *Wolbachia* titers in males was sufficient to rescue queen
fecundity. This requires studying the effects of *Wolbachia* on sperm production. One
possibility is that rifampicin treatment causes ribosomal stress that prevents *Wolbachia*

from modifying sperm. We also do not know why the CI effect was stronger when using
males from colonies naturally uninfected with *Cand*. Westeberhardia, but this points
towards interaction effects between the two symbionts.
379
Our study raises many questions about the evolutionary history of the *C. obscurior-*
Wolbachia association. The putative origin of *C. obscurior* in Southeast Asia, together with
sequence identity between *Wolbachia* strains found in Japan and Taiwan, suggest that
these strains represent the more ancestral association. Geographic patterning separating
New from Old World *Wolbachia* strains has been observed previously in ants [45], and
frequent horizontal transmissions [46,47] can lead to a surprising diversity within species [48]. *C.*
obscurior and its sister species *C. wroughtonii* are the only known representatives of the
100+ described species in the genus *Cardiocondyla* (the so-called heart-node ants) that
have adopted an arboreal lifestyle, as opposed to nesting in soil and rock crevices. This
makes them highly susceptible to horizontal transmission of *Wolbachia* from other plant-
associated insects. How exactly the BR population acquired a novel strain of *Wolbachia* in
the New World habitat, and how this strain subsequently replaced the ancestral strain, is
unclear, however. Although *Wolbachia* from different supergroups may co-exist in a host
[12], it is unlikely that two strains from the same supergroup co-exist without eventually
recombining and merging. Generally, unidirectional CI should facilitate introgression
between populations, while bidirectional CI should promote divergence. This predicts
mixed *C. obscurior* genotypes but only the presence of the *w*Cobs-JP strain in secondary
contact zones, assuming that hybrids perform as well as parental lineages. But how has the
ancestral Asian strain be replaced? *Wolbachia* acquisition and persistence in *C. obscurior*
may be driven by selection on the strain that is better adapted to the local environment,
perhaps analogous to *Wolbachia*-driven thermal preferences in *Drosophila* flies [49,50].
Resolving how *w*Cobs-BR replaced the putatively ancestral *w*Cobs-JP strain (which holds
the power of CI) requires testing how mixed combinations of host and *Wolbachia* perform
under varying ecological conditions. Coupled with a global survey of *Wolbachia-C.*
obscurior associations, this will help unravel how the symbiosis shapes the worldwide
distribution of this tramp ant.
406
Conclusion
One important consequence of unidirectional reproductive isolation is unequal
inheritance, which allows a BDM-causing locus or a reproductive manipulator to quickly
spread in secondary contact zones. Such disruptive factors can be more powerful than
natural selection on other fitness-relevant traits, and lead to the evolution of PoMRI and
hybrid inviability, processes estimated to require several million years in *Drosophila* flies
and *Pogonomyrmex* ants [51,52]. Our discovery of reproductive isolation between allopatric
ant populations, driven by two naturally occurring, closely related *Wolbachia* strains that
differ in their genomic makeup, demonstrates that these evolutionary processes can be
observed in real time and lays the groundwork for detailed studies of *Wolbachia*-ant
biology. In particular, a system in which CI is accompanied by a CI-like phenotype in host
gene expression may prove useful for understanding the function of the *cid/cin* gene
family. Hopefully, hologenomic approaches such as the one presented here will help
provide the comprehensive understanding of species' biology needed to tackle the
challenges associated with human-induced changes to ecological and evolutionary
dynamics.

423
424 Methods

Cardiocondyla obscurior presumably originates from Southeast Asia and is now distributed around the tropics and subtropics. Queens in *C. obscurior* are produced year-round and have a short generation time of ~120 days (from egg to queen to egg). Colonies usually contain 2-5 queens [10] and a single wingless male that can live up to a few months and continually produces sperm [53]. Such males transfer on average 1000 sperm cells per copulation [54], which is sufficient for a lifelong production of diploid offspring by an inseminated queen. Newly hatched queens can decide to stay in the maternal nest or to leave - alone or with a small group of workers - irrespective of mating status. Like all Hymenoptera, ants exhibit haplodiploid sex determination and virgin females can produce males from haploid, unfertilized eggs. Virgin queens in *C. obscurior* can copulate with their own sons (pers. obs.), enabling the production of workers and queens, similar to a related *Cardiocondyla* species [55]. This biology has facilitated the worldwide spread of several species of the genus [56].

In this study, we used inbred laboratory 'populations' derived from colonies collected in Okinawa, Japan in 2010 (JP) and Bahia, Brazil in 2009 (BR). Although the two populations differ in cuticular hydrocarbon composition, queen size and behavior [10], workers are morphologically indiscernible according to exploratory data analyses by hierarchical and nonhierarcical, vector-based forms of Nest Centroid clustering: NC-Ward, NC-part.hclust, NC-part.kmeans, NC-NMDS-kmeans [57,58]. These analyses considered 15 continuous morphometric characters. Specifically, when considering 82 cosmopolitan nest samples of the sister taxa *C. wroughtonii* and *C. obscurior*, all JP and BR samples were allocated to the latter species by all four types of NC-clustering. This result was confirmed with posterior probabilities of $p > 0.98$ if these samples were run as wild-cards in a linear discriminant analysis. Analyses of 32 nest samples representing only the *C. obscurior* cluster could not allocate the JP and BR populations to different intraspecific clusters nor could these analyses demonstrate any intraspecific structuring at all. This suggests that *Wolbachia*-induced reproductive isolation has not yet led to a divergence into morphospecies. In addition to JP and BR colonies, we used colonies collected from a second Asian population in Taipei, Taiwan in 2013 (TW) to confirm the presence of unidirectional CI. For all experiments, colonies were kept in square plaster-bottom nests (100 x 100 x 20 mm, Sarstedt, Germany) and held in climate chambers under a 12h/12h light/dark cycle and a 28°C/22°C temperature cycle. Food (honey and *Drosophila* or pieces of *Periplaneta americana*) and water were provided to stock colonies every 3 days. All animal treatment guidelines applicable to ants under international and German law were followed. Collection of colonies that form the basis of the laboratory population used in this study was permitted by the Brazilian Ministry of Science and Technology (RMX 004/02). No other permits were required for this study.

Antibiotic treatment

Rifampicin is a broad-spectrum antibiotic that acts via inhibition of bacterial RNA polymerase [59]. To produce *Wolbachia*-cured males, we split large *Cand*. Westeberhardia-infected (JP) and uninfected (JP$_{we-}$) stock colonies into two equal halves. Half of these splits were designated as controls (JP: n=13, JP$_{we-}$: n=5), while the other half was treated with antibiotics (JP: n=13, JP$_{we-}$: n=5). We diluted 0.0025 g of solid rifampicin (Sigma-Aldrich,

USA) in 500 µl of a 1:1 honey-water solution for a final antibiotic concentration of 0.5%.
The antibiotic solution was placed on a shaker for 4 hours to ensure complete mixing, and
subsequently covered with aluminum foil and stored at -20°C. Treated colonies were fed
with the antibiotic solution twice per week every other week, for a total duration of ten
weeks. On the days following treatment, the antibiotic solution was removed from the
nest. Between antibiotic feedings (i.e. every other week), colonies were fed with honey
and pieces of autoclaved cockroach (to prevent bacterial re-infections) and water ad
libitum. Control colonies were fed twice a week with honey and cockroaches and received
water ad libitum.
479
480 ***Wolbachia* and *Cand.* Westeberhardia infection**
481 We assessed the presence and titers of bacterial infections using PCR and qPCR,
482 respectively, on DNA extracted from whole bodies (worker pupae & adults, queen pupae
483 & adults, adult males). DNA was extracted from samples using a standard CTAB protocol.
484
Presence of infection
We confirmed infection in a subset of workers from BR and JP colonies. Infection with
Wolbachia was tested with a PCR assay using primers specific for a fragment of the
Wolbachia surface protein gene of the BR (105-bp, *wsp_wCobs_BR-for:* 5-
'TAAATCTTGCATCTGTAACATT-3', *wsp_wCobs_BR-rev:* 5'-
CTGCGGATACTGATACAACTACTG-3') and JP (254-bp, *wsp_wCobs_JP-for:* 5'-
CATTTTGACTACTCACAGCGGTTG-3', *wsp_wCobs_JP-rev:*
5'CTGCGGATACTGATACAACTACTG-3') strains. *Cand.* Westeberhardia infection was
tested by amplifying a 204-bp fragment of the *ribonucleoside-diphosphate reductase 1*
subunit beta gene of *Westeberhardia* (*WEOB_403*) (*nrdB_weCobs-for:* 5'-
GGAAGGAGTCCTAATGTTGCG-3'; nrdB_*weCobs*-rev: 5'-ACC
AGAAATATCTTTTGCACGTT-3'). A 104-bp fragment of the *C. obscurior* gene *elongation*
factor 1-alpha 1 (*Cobs_01649*) (*EF1-for:* 5'-TCACTGGTACCTCGCAAGCCGA-3'; *EF1-rev:*
5'-AGCGTGCTCACGAGTTTGTCCG-3') was used as a positive control. In addition,
infection was assessed in two adult queens from each of five TW colonies to confirm that
this population carries the *w*Cobs-JP strain. Each PCR reaction contained 5 µL Taq
polymerase (GoTaq, Promega), 3 µL H_2O, 0.5 µL forward primer (10 µM), 0.5 µL reverse
primer (10 µM) and 1 µL DNA. PCRs were run at 94°C for 4 minutes followed by 36 cycles
at 94°C for 30s, 57°C for 30s and 72° for 30s, with a final step of 72°C for 10 minutes.
Electrophoresis separation of PCR products was performed on a 1.5% TAE-Gel with 0.5 µl
of 0.001 % Gel-Red for 45 minutes at 60 V and 65 mA.
506
Population-specific Wolbachia titers
We compared *Wolbachia* infection titers of adult workers and queens from BR (n=9
workers, n=10 queens) and JP (n=10 workers, n=9 queens) colonies by amplifying a
portion of the *Cytochrome c oxidase subunit 1* gene (*coxA_wCobs-for*: 5'-
TTGGTCATCCAGAAGTTTACGT-3', *coxA_wCobs-rev*: 5'-TGAGCCCAAACCATAAAGCC-
3') in qPCR. As a relative standard *elongation factor 1-alpha 1* (*Cobs_01649*) was used.
Each qPCR reaction contained 5 µL SYBR (Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1
µL reverse primer (2 µM) and 1 µL DNA. Reactions were performed in duplicates and run
at 95°C for 3 minutes followed by 40 cycles at 95°C for 5s, 60°C for 20s, and 95° for 10s,
and a final melt curve analysis step in which reactions were heated from 65°C to 95°C in

5°C steps. Single-amplicon production was confirmed with melt curve analyses and
relative *Wolbachia* titers calculated with the $2^{-\Delta CT}$ method [60].
519
Age and morph-specific Wolbachia titers
We quantified *Wolbachia* titers in individual queens and workers from BR colonies by
determining relative *coxA_wCobs* copy numbers with qPCR across developmental stages
and adult age classes. In workers, titers were measured in pupae (n=4 white pupae, n=5
dark pupae) and in adults two days (n=5), 14 days (n=7), and 28 days (n=6) post-hatching.
In queens, titers were measured in pupae (n=5 white pupae, n=4 dark pupae), as well as in
adults two days (n=7), 14 days (n=7), 28 days (n=10 virgin queens, n=7 mated queens)
and 48 days (n=9) post-hatching. In addition, we quantified *coxA_wCobs* copy numbers in
adult morphs (n=15 queens, n=12 workers, n=6 winged males, n=7 wingless males). For
all reactions, *elongation factor 1-alpha 1* (*Cobs_01649*) was used as a housekeeper. Each
qPCR reaction contained 5 µL SYBR (Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1 µL
reverse primer (2 µM) and 1 µL DNA. Reactions were performed in triplicates and run as
described above. Single-amplicon production was confirmed with melt curve analyses and
relative quantities of *Wolbachia* calculated with the $2^{-\Delta CT}$ method.
534
Rifampicin effects on infection titers
We verified the efficacy of rifampicin treatment by measuring *Wolbachia* and
Westeberhardia titers in dark worker pupae produced by antibiotic-treated (n=21 from 9
colonies) and control colonies (n=21 from 8 colonies) using qPCR of *coxA_wCobs and*
nrdB_weCobs. In addition, we measured the *Wolbachia* titers of JP males used in mating
combinations BR x JP and BR x JP^{rif+} (JP: n=15, JP^{rif+}: n=12) with qPCR of a portion of the
JP-specific *wsp* gene (*wsp_wCobs_JP*). For all samples, *elongation factor 1-alpha 1*
(*Cobs_01649*) was used as a housekeeper. Each qPCR reaction contained 5 µL SYBR
(Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1 µL reverse primer (2 µM) and 1 µL DNA.
Reactions were performed in duplicates (males) or triplicates (worker pupae) and run as
described above (except with an annealing temperature of 57°C for male pupae). Single-
amplicon production was confirmed with melt curve analyses and relative quantities of
Wolbachia and *Westeberhardia* calculated with the $2^{-\Delta CT}$ method.
548
Experimental crosses
We designed seven mating combinations between queens and males from the two
populations: 1) JP x JP (n=7), 2) JP x BR (n=26), 3) BR x BR (n=11), 4) BR x JP (n=36), 5) BR
x JP$_{we-}$ (n=29), 6) BR x JP^{rif+} (n=13) and 7) BR x JP$_{we-}^{rif+}$ (n=5). Each experimental colony was
set up with one dark queen pupa, one freshly hatched male, 10 workers and brood (10
eggs, 10 larvae, 10 worker pupae). All individuals were collected from un-manipulated
stock colonies, except for JP^{rif+} and JP$_{we-}^{rif+}$ males used in combinations #6 and #7, which
were collected from rifampicin-treated colonies. To avoid manipulation effects, all
combinations were set up in parallel. In such experimental colonies, mating between
queens and males usually occurs within three weeks, after which queens shed their wings
and being laying eggs. We monitored colonies for queen wing loss once per week. Once
queens had shed their wings, we removed any brood placed in the colony during set up
and standardized worker number to 20. Colonies were then monitored once a week for six
weeks and all eggs and pupae (queen, worker, wingless male, winged male) counted. A
subset of colonies was monitored for a total of 12 weeks (JP x BR: n=19, BR x JP: n=28, BR

x JP[rif+]: n=9). In each colony, the number of adult workers was kept constant at 20
individuals by removing hatched workers or adding workers from stock colonies. From
monitoring data, we calculated mean and maximum weekly egg numbers, and total
number and sex ratio of produced pupae (male pupae:total pupae).

To verify that CI is specifically induced by the *w*Cobs-JP strain when it encounters the
*w*Cobs-BR strain, we set up three additional mating combinations between queens and
males using a third population from Taiwan (TW) which also harbors the *w*Cobs-JP strain:
1) JP x TW (n=5), 2) TW x JP (n=6), 3) BR x TW (n=5). Experimental colonies were set up
and monitored as described above for six weeks.

DNA sequencing and genome assembly
We extracted DNA from a pool of 26 (JP) and 30 (BR) males from two colonies, collected
in 2009 in Una, Brazil (BR2009-alpha) from aborted fruits of coconut trees and in 2010 in
Naha, Japan (JP2010-OypB) from under bark of coral trees (*Erythrina* sp.). For details see
[10]. Briefly, the reads were right-tail clipped (minimum quality of 20) and all reads with
undefined nucleotides were removed using a combination of FASTX-Toolkit v0.0.14
(http://hannonlab.cshl.edu/fastx_toolkit/, last accessed December 3, 2010) and
PRINSEQ++ v1.2 (10.7287/peerj.preprints.27553v1). The resulting paired end reads were
assembled with SPAdes v3.13.1 (-k 33,55,77 --only-assembler [61]. The resulting contigs
were then binned according to an *ad hoc* BLASTX-based method. First, a proteome
database was built from representative genomes (Table S5). Second, the best hit for each
scaffold was used for assigning a preliminary bin and a coverage cut-off (10x for BR and
30x for JP) was implemented based on the coverage of the longest contigs. Third, each
scaffold was manually inspected using the online NCBI web-server *vs.* BLAST+2.9.0. If the
best hits to proteins were consistently to *Wolbachia* bacteria across the scaffold, the query
sequence was retained. Finally, the assembly graph was loaded into Bandage v0.8.1 [62],
the graph(s) matching *Wolbachia* sequences were identified and the sequences were
extracted. These extracted sequences were then compared to the *Wolbachia* draft bin. In
both cases, the extracted sequences were roughly the same compounded length as the
manually-curated bins and added only a couple of kilo base pairs (kbp), suggesting near
completeness of the genome. Using these *Wolbachia* bins, we performed a mapping
using BOWTIE2 v2.3.4.1 [63] followed by a re-assembly of the mapped reads in SPAdes
(with previously-mentioned parameters). This final reassembly was filtered by both
coverage and manual screening of scaffolds shorter than 3 kbp. Assembled sequences
have been deposited in the European Nucleotide Archive (ENA) under the accessions
CACTIU010000000.1 (*w*Cobs-BR) and CACTIV010000000.1 (*w*Cobs-JP).

Phylogeny and comparative genomics of *Wolbachia* strains
A draft annotation of the scaffolds was done using Prokka v1.14 [64]. For comparative
genomic purposes, we built a subset of the predicted ORFs by filtering out all proteins
annotated as "hypothetical proteins" with genes shorter than 300 base pairs (bp). This
helped us avoid wrongly annotated ORFs overlapping pseudogenes. To infer the
phylogenetic relationship between *Wolbachia* strains found to infect the analysed BR and
JP populations as well as among other *Wolbachia* bacteria, we collected the amino acid
sequences for the ribosomal proteins of 23 *Wolbachia* strains from supergroups A (8), B
(10), C (2), D (1), E (1), and F (1) (Table S6). Using *Anaplasma phagocytophilum* strain HZ

and *Ehrlichia canis* strain Jake as outgroups, we inferred a Bayesian phylogeny using
MrBayes v3.2.6 [65]. We ran two independent analyses with four chains each for 300,000
generations and checked for convergence. The substitution model JTT+I+G+F was used
as suggested by jModelTest v2.1.10.
615
616 To infer the shared protein-coding gene content of *w*Cobs-BR and *w*Cobs-JP strains, we
617 used OrthoMCL v2.0.9 [66,67] with an inflation value of 1.5. The synteny of both strains was
618 assessed using nucmer v3.23 [68] using anchor matches that were unique in both the
619 reference and query (--mum).
620

Cif-family gene annotation and phylogenetics

cif genes were first identified by BLASTP using the protein sequences WP_010962721.1
(*cifA*) and WP_010962722.1 (*cifB*) as references. Predicted start codons were manually
checked for the presence of a Shine-Dalgarno-like sequence. To identify whether the *cif*
genes form each strain belonged to the same type or not, we performed a phylogenetic
analysis following Lindsey *et. al.* 2018. Briefly, the previously used protein sequences were
aligned with the newly acquired ones from both wCobs-BR and wCobs-JP. The alignments
were then back-translated to nucleotides. For *cifB*, the type IV genes were excluded and
the 3'-end of the alignment was manually truncated to remove the longer unmatched
regions of some *cifB* variants. Bayesian phylogenetic inference was performed with
MrBayes v3.2.6 with the GTR+G substitution model (as suggested by jModelTest) running
two independent analyses with four chains each for 1,000,000 generations, and checked
for convergence. Following [16], Cif protein domains were searched for using HHpred's
v3.2.0 webserver (https://toolkit.tuebingen.mpg.de/tools/hhpred) with default parameters
and the following databases: SCOPe70 v2.07, COG/KOG v1.0, Pfam-A v32.0, and SMART
v6.0. hits with a probability >=80% were considered.
637
638 All genomic data files are available at the zenodo repository
639 https://doi.org/10.5281/zenodo.3561160
640

Histological sectioning and fluorescence in situ hybridization

C. obscurior (JP) male abdomens were fixed in 4% formaldehyde in PBS and washed with
water followed by a dehydration series in n-Butanol (30%, 50%, 70%, 80%, 90%, 96%) at
room temperature with shaking for 1 h at each step. Absolute n-butanol was used for the
last three hours at 30°C, exchanging the solution each hour. Dehydrated samples were
embedded in Technovit 8100 cold polymerizing resin (Heraeus Kulzer, Germany)
according to the manufacturer's instructions, cut into semithin sections (8 µm) with a
Microm HM355S microtome (Thermo Fisher Scientific, Germany) and mounted on glass
slides coated with poly-L-lysine (Kindler, Germany). The tissue sections were incubated for
90 min at 50°C in hybridization buffer (0.9 M NaCl, 20 mM Tris/HCl (pH 8.0), 0.01 % SDS)
containing 0.5 µM of the *Wolbachia*-specific probes Wolb_W2_Cy5 (5'-
CTTCTGTGAGTACCGTCATTATC-3') [69] and Wolb_Wol3_Cy5 (5'-
TCCTCTATCCTCTTTCAATC-3'), [70], as well as the *Cand*. Westeberhardia-specific probe
Wcard1_Cy3 (5'- ATCAGTTTCGAACGCCATTC-3') [13]. DAPI (4',6-diamidino-2-
phenylindole) was used for host DNA counterstaining. After hybridization, samples were
washed with buffer (0.1 M NaCl, 20 mM Tris/HCl (pH 8.0), 5mM EDTA, 0.01% SDS) for 20
min at 50°C. Subsequently, the washing buffer was removed and washed with distilled

water for 20 min at 50°C. Samples were air-dried and covered with VectaShield® (Vector
Laboratories, Burlingame, CA, USA) and stored overnight at room temperature. Images
were acquired on an AxioImager.Z1 epifluorescence microscope (Carl Zeiss, Jena,
Germany).
662
Acknowledgements

The authors thank John Wang for help collecting colonies in Taiwan and for providing the
wsp sequence of the TW population. We thank Julia Hacker for help with the BR x TW
crossings and Helena Lowack and Jennifer Wallner for help with lab work. JD Shropshire
helped with the *cid/cif/cin* annotation and Benjamin Weiss assisted with histological
preparations. This work was supported by the Marie-Curie AgreenSkills+ fellowship
program cofunded by the EU's Seventh Framework Programme (FP7-609398) to A.M.M,
and a DFG grant He1623/31 to JH and JO.

671
References

1. Wetterer, J. K., Wild, A. L., Suarez, A. V., Roura-Pascual, N. & Espadaler, X.
Worldwide spread of the Argentine ant, *Linepithema humile* (Hymenoptera:
Formicidae). *Myrmecol News* **12,** 187–194 (2009).
2. Ascunce, M. S. *et al.* Global invasion history of the fire ant *Solenopsis invicta*. *Science*
331, 1066–1068 (2011).
3. Tsutsui, N. D., Suarez, A. V., Holway, D. A. & Case, T. J. Reduced genetic variation
and the success of an invasive species. *Proc Natl Acad Sci U S A* **97,** 5948–5953
(2000).
4. Brucker, R. M. & Bordenstein, S. R. Speciation by symbiosis. *Trends in Ecology &*
Evolution **27,** 443–451 (2012).
5. Zug, R. & Hammerstein, P. Bad guys turned nice? A critical assessment of *Wolbachia*
mutualisms in arthropod hosts. *Biological Reviews* **90,** 89–111 (2014).
6. Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J. H.
How many species are infected with *Wolbachia*?--A statistical analysis of current
data. *FEMS Microbiology Letters* **281,** 215–220 (2008).
7. LePage, D. P. *et al.* Prophage WO genes recapitulate and enhance *Wolbachia*-
induced cytoplasmic incompatibility. *Nature* **543,** 243–247 (2017).
8. Werren, J. H. Biology of *Wolbachia*. *Annu. Rev. Entomol.* **42,** 587–609 (1997).
9. Russell, J. A. The ants (Hymenoptera: Formicidae) are unique and enigmatic hosts of
prevalent *Wolbachia* (Alphaproteobacteria) symbionts. *Myrmecol News* **16,** 7–23
(2012).
10. Schrader, L. *et al.* Transposable element islands facilitate adaptation to novel
environments in an invasive species. *Nature Communications* **5,** 5495 (2014).
11. Schrempf, A. *et al.* Mating with an allopatric male triggers immune response and
decreases longevity of ant queens. *Molecular Ecology* **24,** 3618–3627 (2015).
12. Ellegaard, K. M., Klasson, L., Näslund, K., Bourtzis, K. & Andersson, S. G. E.
Comparative genomics of *Wolbachia* and the bacterial species concept. *PLoS Genet*
9, e1003381 (2013).
13. Klein, A. *et al.* A novel intracellular mutualistic bacterium in the invasive ant
Cardiocondyla obscurior. *ISME J* **10,** 376–388 (2016).
14. Shropshire, J. D. & Bordenstein, S. R. Two-By-One model of cytoplasmic
incompatibility: Synthetic recapitulation by transgenic expression of cifA and cifB in

705   *Drosophila*. *PLoS Genet* **15,** e1008221 (2019).

706 15. Shropshire, J. D. *et al.* Models and Nomenclature for Cytoplasmic Incompatibility:
707   Caution over Premature Conclusions - A Response to Beckmann et al. *Trends Genet.*
708   **35,** 397–399 (2019).

709 16. Lindsey, A. R. I. *et al.* Evolutionary Genetics of Cytoplasmic Incompatibility Genes
710   cifA and cifB in Prophage WO of *Wolbachia*. *Genome Biology and Evolution* **10,**
711   434–451 (2018).

712 17. Perlmutter, J. I. *et al.* The phage gene wmk is a candidate for male killing by a
713   bacterial endosymbiont. *PLoS Pathog* **15,** e1007936 (2019).

714 18. Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics Toolkit with
715   a New HHpred Server at its Core. *J. Mol. Biol.* **430,** 2237–2243 (2018).

716 19. Veneti, Z. *et al.* Cytoplasmic incompatibility and sperm cyst infection in different
717   *Drosophila-Wolbachia* associations. *Genetics* **164,** 545–552 (2003).

718 20. Clark, M. E. *et al. Wolbachia* modification of sperm does not always require
719   residence within developing sperm. *Heredity* **101,** 420–428 (2008).

720 21. Wenseleers, T., Sundstrom, L. & Billen, J. Deleterious *Wolbachia* in the ant *Formica*
721   *truncorum*. *Proceedings of the Royal Society B: Biological Sciences* **269,** 623–629
722   (2002).

723 22. Pontieri, L., Schmidt, A. M., Singh, R., Pedersen, J. S. & Linksvayer, T. A. Artificial
724   selection on ant female caste ratio uncovers a link between female-biased sex ratios
725   and infection by *Wolbachia* endosymbionts. *J Evolution Biol* **30,** 225–234 (2017).

726 23. Shoemaker, D. D., Ross, K. G., Keller, L., Vargo, E. L. & Werren, J. H. *Wolbachia*
727   infections in native and introduced populations of fire ants (*Solenopsis spp.*). *Insect*
728   *Molecular Biology* **9,** 661–673 (2000).

729 24. Correa, C. C. & Ballard, J. W. O. *Wolbachia* Associations with Insects: Winning or
730   Losing Against a Master Manipulator. *Front. Ecol. Evol.* **3,** 506 (2016).

731 25. Trivers, R. L. & Hare, H. Haploidploidy and the evolution of the social insect. *Science*
732   **191,** 249–263 (1976).

733 26. Keller, L. *et al.* Sex ratio and *Wolbachia* infection in the ant *Formica exsecta*. *Heredity*
734   **87,** 227–233 (2001).

735 27. Singh, R. & Linksvayer, T. A. The bacterial endosymbiont *Wolbachia* increases
736   reproductive investment and accelerates the life cycle of ant colonies. **90,** 89

737 28. Treanor, D., Pamminger, T. & Hughes, W. O. H. The evolution of caste-biasing
738   symbionts in the social hymenoptera. *Insect. Soc.* **65,** 513–519 (2018).

739 29. Schrempf, A., Moser, A., Delabie, J. & Heinze, J. Sperm traits differ between winged
740   and wingless males of the ant *Cardiocondyla obscurior*. *Integrative Zoology* **11,**
741   427–432 (2016).

742 30. Andersen, S. B., Boyle, M., Nash, D. R. & Boomsma, J. J. Dynamic *Wolbachia*
743   prevalence in *Acromyrmex* leaf-cutting ants: potential for a nutritional symbiosis. *J*
744   *Evolution Biol* **25,** 1340–1350 (2012).

745 31. Kramer, B. H., Schrempf, A., Scheuerlein, A. & Heinze, J. Ant Colonies Do Not Trade-
746   Off Reproduction against Maintenance. *PLoS ONE* **10,** e0137969 (2015).

747 32. Cremer, S. & Heinze, J. Adaptive production of fighter males: queens of the ant
748   *Cardiocondyla* adjust the sex ratio under local mate competition. *Proc. Biol. Sci.*
749   **269,** 417–422 (2002).

750 33. de Menten, L., Cremer, S., Heinze, J. & Aron, S. Primary sex ratio adjustment by ant
751   queens in response to local mate competition. *Animal Behaviour* **69,** 1031–1035

(2005).

34.   Suefuji, M., Cremer, S., Oettler, J. & Heinze, J. Queen number influences the timing
of the sexual production in colonies of *Cardiocondyla* ants. *Biol. Lett.* **4,** 670–673
(2008).

35.   Gotzek, D., Axen, H. J., Suarez, A. V., Helms Cahan, S. & Shoemaker, D. Global
invasion history of the tropical fire ant: a stowaway on the first global trade routes.
Molecular Ecology **24,** 374–388 (2015).

36.   Foucaud, J. *et al.* Worldwide invasion by the little fire ant: routes of introduction and
eco-evolutionary pathways. *Evol Appl* **3,** 363–374 (2010).

37.   Braig, H. R., Zhou, W., Dobson, S. L. & O'Neill, S. L. Cloning and characterization of a
gene encoding the major surface protein of the bacterial endosymbiont *Wolbachia*
pipientis. *Journal of Bacteriology* **180,** 2373–2378 (1998).

38.   Clark, M. E., Heath, B. D., Anderson, C. L. & Karr, T. L. Induced paternal effects
mimic cytoplasmic incompatibility in *Drosophila*. *Genetics* **173,** 727–734 (2006).

39.   Beckmann, J. F., Ronau, J. A. & Hochstrasser, M. A *Wolbachia* deubiquitylating
enzyme induces cytoplasmic incompatibility. *Nat Microbiol* **2,** 17007–7 (2017).

40.   Beckmann, J. F. *et al.* The Toxin-Antidote Model of Cytoplasmic Incompatibility:
Genetics and Evolutionary Implications. *Trends Genet.* **35,** 175–185 (2019).

41.   Bonneau, M. *et al. Culex pipiens* crossing type diversity is governed by an amplified
and polymorphic operon of *Wolbachia*. *Nature Communications* **9,** 319–10 (2018).

42.   Brucker, R. M. & Bordenstein, S. R. The Hologenomic Basis of Speciation: Gut
Bacteria Cause Hybrid Lethality in the Genus *Nasonia*. *Science* (2013).
doi:10.1126/science.1240659

43.   Cooper, B. S., Vanderpool, D., Conner, W. R., Matute, D. R. & Turelli, M. *Wolbachia*
Acquisition by *Drosophila yakuba*-Clade Hosts and Transfer of Incompatibility Loci
Between Distantly Related *Wolbachia*. *Genetics* **212,** 1399–1419 (2019).

44.   Sinha, A., Li, Z., Sun, L. & Carlow, C. K. S. Complete Genome Sequence of the
Wolbachia wAlbB Endosymbiont of *Aedes albopictus*. *Genome Biology and*
Evolution **11,** 706–720 (2019).

45.   Kautz, S., Rubin, B. E. R. & Moreau, C. S. Bacterial Infections across the Ants:
Frequency and Prevalence of *Wolbachia*, *Spiroplasma*, and *Asaia*. *Psyche: A Journal*
of Entomology **2013,** 1–11 (2013).

46.   Russell, J. A. *et al.* Specialization and geographic isolation among *Wolbachia*
symbionts from ants and lycaenid butterflies. *Evolution* **63,** 624–640 (2009).

47.   Tolley, S. J. A., Nonacs, P. & Sapountzis, P. *Wolbachia* Horizontal Transmission
Events in Ants: What Do We Know and What Can We Learn? *Front Microbiol* **10,**
296 (2019).

48.   Kelly, M., Price, S. L., de Oliveira Ramalho, M. & Moreau, C. S. Diversity of *Wolbachia*
Associated with the Giant Turtle Ant, *Cephalotes atratus*. *Curr Microbiol* **76,** 1330–
1337 (2019).

49.   Arnold, P. A., Levin, S. C., Stevanovic, A. L. & Johnson, K. N. *Drosophila*
melanogaster infected with *Wolbachia* strain wMelCS prefer cooler temperatures.
Ecological Entomology **44,** 287–290 (2018).

50.   Truitt, A. M., Kapun, M., Kaur, R. & Miller, W. J. *Wolbachia* modifies thermal
preference in *Drosophila melanogaster*. *Environ Microbiol* **21,** 3259–3268 (2018).

51.   Coyne, J. A. & Orr, H. A. 'Patterns of speciation in *Drosophila*' revisited. *Evolution*
51, 295–303 (1997).

52. Schwander, T., Suni, S. S., Cahan, S. H. & Keller, L. Mechanisms of reproductive
isolation between an ant species of hybrid origin and one of its parents. *Evolution*
62, 1635–1643 (2008).
53. Heinze, J. & Hölldobler, B. Fighting for a harem of queens: physiology of
reproduction in *Cardiocondyla* male ants. *Proc Natl Acad Sci U S A* **90,** 8412–8414
(1993).
54. Schrempf, A., Darrouzet, E. & Heinze, J. Mating success and potential male-worker
conflict in a male-dimorphic ant. *BMC Evol Biol* **7,** 114 (2007).
55. Schmidt, C. V., Frohschammer, S., Schrempf, A. & Heinze, J. Virgin ant queens mate
with their own sons to avoid failure at colony foundation. *Naturwissenschaften* **101,**
69–72 (2013).
56. Heinze, J., Cremer, S., Eckl, N. & Schrempf, A. Stealthy invaders: the biology of
Cardiocondyla tramp ants. *Insect. Soc.* **53,** 1–7 (2006).
57. Seifert, B., Ritz, M. & Csősz, S. Application of exploratory data analyses opens a new
perspective in morphology-based alpha-taxonomy of eusocial organisms.
Myrmecological News **19,** 1–15 (2013).
58. Seifert, B., Okita, I. & Heinze, J. A taxonomic revision of the *Cardiocondyla nuda*
group (Hymenoptera: Formicidae). *Zootaxa* **4290,** 324–33 (2017).
59. Campbell, E. A. *et al.* Structural mechanism for rifampicin inhibition of bacterial rna
polymerase. *Cell* **104,** 901–912 (2001).
60. Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative CT
method. *Nat Protoc* **3,** 1101–1108 (2008).
61. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications
to single-cell sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).
62. Wick, R. R., Schultz, M. B., Zobel, J. & Holt, K. E. Bandage: interactive visualization of
de novo genome assemblies. *Bioinformatics* **31,** 3350–3352 (2015).
63. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Meth*
9, 357–359 (2012).
64. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30,**
2068–2069 (2014).
65. Ronquist, F. *et al.* MrBayes 3.2: efficient Bayesian phylogenetic inference and model
choice across a large model space. *Systematic Biology* **61,** 539–542 (2012).
66. Li, L. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes.
Genome Research **13,** 2178–2189 (2003).
67. Chen, F., Mackey, A. J., Vermunt, J. K. & Roos, D. S. Assessing performance of
orthology detection strategies applied to eukaryotic genomes. *PLoS ONE* **2,** e383
(2007).
68. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome*
Biol. **5,** R12 (2004).
69. Heddi, A., Grenier, A.-M., Khatchadourian, C., Charles, H. & Nardon, P. Four
intracellular genomes direct weevil biology: Nuclear, mitochondrial, principal
endosymbiont, and *Wolbachia. Proceedings of the National Academy of Sciences*
96, 6814–6819 (1999).
70. Sanguin, H. *et al.* Development and validation of a prototype 16S rRNA-based
taxonomic microarray for Alphaproteobacteria. *Environ Microbiol* **8,** 289–307
(2006).
845

Figure 1: Mean weekly egg numbers produced by intra- and inter-population crosses of the ant *C. obscurior*

Mean weekly egg numbers produced by Brazilian (BR) and Japanese (JP) queens mated to males from their own or from a different population over a period of six weeks. JP ants were collected from colonies that either carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae (JP) or did not (JP_we-). Males used in "CI rescue" crosses were collected from JP colonies with and without *Cand*. Westeberhardia cardiocondylae, which had been treated with the antibiotic rifampicin (JP^rif+; JP_we-^rif+). Numbers below box plots indicate the number of replicates. Differences between groups were tested with pairwise Mann-Whitney-U-tests followed by Bonferroni-Holms correction of p-values. Letters above boxplots indicate statistically significant differences at p<0.05. For Bonferroni-Holms corrected pairwise p-values see Table S1.

Figure 2: Total pupae numbers and pupae sex ratios produced by intra- and inter-population crosses of the ant *C. obscurior*

Total number of pupae (A) and sex ratio of pupae (B) produced in crosses between Brazilian (BR) and Japanese (JP) queens and males over a period of 12 weeks. All JP ants were collected from colonies that carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae (JP). Males used in BR x JP[rif+] crosses were collected from JP colonies that had been treated with the antibiotic rifampicin. Numbers below box plots indicate the number of replicates. For total pupae numbers, differences between groups were tested with pairwise Mann-Whitney-U-tests followed by Bonferroni-Holms correction of p-values. Sex ratios produced by different crosses were compared with a generalized linear model with logit link followed by manual Bonferroni-Holms correction of p-values. Letters above boxplots indicate statistically significant differences at p<0.05.

Figure 3: Effects of rifampicin treatment on *Wolbachia* titres in males
All males were collected from colonies that carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae. Numbers below box plots indicate the number of replicates. Differences in *Wolbachia* titres between control and rifampicin-treated males were tested with a Mann-Whitney-U-Test. Stars indicate statistically significant differences at p<0.001 (***).

Figure 4: *Wolbachia* spp. phylogeny and genome synteny

(A) Bayesian phylogenetic placement of *Wolbachia* strains sequenced from *C. obscurior* (*w*Cobs-BR and *w*Cobs-JP). *Anaplasma phagocytophilum* and *Ehrlichia canis* were used as outgroups. For *Wolbachia*, names at tips correspond to the host from which they were sequenced/isolated. Red dots are used to highlight the newly sequenced strains. Next to the binomial names, strain names are shown in grey and silhouettes for the host are sketched in black. Horizontal bars delimit *Wolbachia* supergroups (A-F). *=1.0 posterior probability. (B) Genome-wise nucleotide-based synteny plot between *w*Cobs-BR and *w*Cobs-JP. Red is used to indicate a direct match and blue a reverse complement match.

Figure 5. Phylogeny and domain structure of *cif* genes

Bayesian phylogenetic trees for (A) *cifA* and (B) *cifB* genes. Colored ovals are used to highlight the different gene "types". *= 1.0 posterior probability. (C) Domain structure for the *cif* genes of *w*Cobs-BR (top) and *w*Cobs-JP (bottom) strains. For the latter, the two loci are shown and indicated with a vertical bar to the right of the sketches. Color codes and designation of the domains are shown in the legend.

Figure 6: Localization of endosymbiotic bacteria in male adult abdomens of *C. obscurior* via fluorescence in situ hybridization.

Wolbachia were specifically stained with two Cy5-labeled probes (magenta), and *Cand.* Westeberhardia with a Cy3-labeled probe (green). Host cell nuclei are counterstained with DAPI (blue). (A) Transversal section of *C. obscurior* JP male abdomen showing the testes (t), the *Cand.* Westeberhardia-containing bacteriomes (B) and the gut (g). The areas indicated by rectangles are displayed in higher magnification showing *Wolbachia*-infected seminiferous tubules (B) and bacteriomes co-infected with *Cand.* Westeberhardia and *Wolbachia* (C). (D) *C. obscurior* BR male testes and gut epithelium infected with *Wolbachia*. Scale bars: 50 µm (A) and 20 µm (B-D).

Table 1: Wolbachia genome assembly statistics

Genome assembly and draft annotation statistics for the newly sequenced *Wolbachia* strains from *C. obscurior* (*w*Cobs-BR and *w*Cobs-JP).

	Wolbachia	
	wCobs-BR	wCobs-JP
Assembled genome size (Mbp)	1.19	1.30
Number of scaffolds	148	182
N50	14,698	16,173
G+C content (%)	35.3	35.1
Predicted CDSs*	1,013	1,058
tRNAs	34	34
rRNAs	3	3
Other ncRNAs genes/motifs	71	54