1   Cytoplasmic incompatibility between Old and New World populations of a tramp ant
2

3   Çiğdem Ün[1]°, Eva Schultner[1]°, Alejandro Manzano-Marín[2], Laura V. Flórez[3], Bernhard
4   Seifert[4], Jürgen Heinze[1], Jan Oettler[1]*
5
6   [1]Zoology/Evolutionary Biology, University of Regensburg, Universitätsstrasse 31, 93053
7   Regensburg, Germany
8
9   [2]Centre for Microbiology and Environmental Systems Science, University of Vienna,
10 Althanstraße 14, 1090 Vienna, Austria
11
12 [3]Institute of Organismic and Molecular Evolution, Evolutionary Ecology Department,
13 Johannes Gutenberg University Mainz, Hanns-Dieter-Hüsch-Weg-15, 55128 Mainz,
14 Germany
15
16 [4] Senckenberg Museum of Natural History Görlitz, Am Museum 1, 02826 Görlitz, Germany
17
18 °contributed equally
19
20 *corresponding author:
21 Jan Oettler
22 joettler@gmail.com
23
Author ORCIDs:
Çiğdem Ün: 0000-0002-0817-3630
Eva Schultner: 0000-0002-5069-9732
Alejandro Manzano-Marín: 0000-0002-0707-9052
Laura V. Flórez: 0000-0002-0761-3729
Jürgen Heinze: 0000-0002-9403-5946
Jan Oettler: 0000-0002-8539-6029

31
32 Classification: Biological Sciences: Evolution
33
34 Key words: *Wolbachia*, social insects, speciation, endosymbiont, antibiotics
35
Contributions
CÜ performed the crosses and qPCR
ES analyzed and visualized the data
AMM assembled *Wolbachia* genomes, performed comparative genomic analyses,
calculated the phylogeny and annotated CI genes
LVF conducted FISH analyses
BS revised the taxonomic status of the populations
JH acquired funding and reviewed and edited the manuscript
JO conceived and supervised the study
JO, ES and CÜ wrote the original draft of the manuscript
All authors revised and approved the final version of the manuscript
47

Abstract

As we enter the Anthropocene, the evolutionary dynamics of species will change drastically, and as yet unpredictably, due to human activity. Already today, increases in global human traffic have resulted in the rapid spread of species to new areas, leading to the formation of geographically isolated populations. These go on to evolve in allopatry, which can lead to reproductive isolation, and potentially, the formation of new species. Surprisingly, little is known about such eco-evolutionary processes in ants, even though numerous invasive ant species are globally distributed in geographically isolated populations. Here, we describe the first case of cytoplasmic incompatibility (CI) between populations of a cosmotropic distributed tramp ant with Asian roots, *Cardiocondyla obscurior*, which has acquired a novel *Wolbachia* strain in the New World. Our study uncovers the first symbiont-induced mechanism of reproductive isolation in ants, providing a novel perspective on the biology of globally distributed ants.

Introduction

Ants are the most abundant group of insects on earth, and numerous ant species are classified as highly invasive on a global scale. Their distribution is, on the one hand, strongly facilitated by humans. This is evidenced, for instance, by the rapid worldwide spread of the Argentine ant and the fire ant from their origin in South America [1,2]. On the other hand, the particular biology of ants, characterized by reproductive division of labor between highly fecund queens and sterile workers, allows the establishment of large populations from just one founding propagule. Introduced populations thus go through a genetic bottleneck, reducing diversity within the population while at the same time increasing diversity between populations [3]. Such populations then evolve in allopatry, and differences caused by genetic drift may be further amplified over time by selection, eventually leading to reproductive isolation. Reproductive isolation, in turn, together with the evolution of mating systems, underlies the formation of new species and, hence, the vast diversity of life. By allowing ants (and other organisms) to establish allopatric populations around the world, humans have inadvertently created a valuable scenario for studying species evolution in real time.

Reproductive isolation between populations that live in allopatry can evolve in the form of post-mating mechanisms of reproductive isolation (PoMRI). Importantly, PoMRI can result from Bateson–Dobzhansky–Muller (BDM) incompatibilities, in which two parental loci that have diverged cause hybrid sterility or inviability. Extending such negative epistatic interactions to the host's microbiome adds three mechanisms that can reduce hybrid fitness: hybrid susceptibility (i.e. inferior immune systems and higher pathogen loads in hybrids), hybrid autoimmunity (i.e. upregulation of immune functions in hybrids because of negative epistasis), and cytoplasmic incompatibility (CI, i.e. hybrid embryo death caused by incompatibilities between members of the parental microbiomes) [4]. PoMRI mechanisms can operate at any developmental stage and can directly affect the fitness of both hybrid offspring and mated parental females. The best-known cases of PoMRI caused by bacteria involve *Wolbachia*, the enigmatic evolutionary omnia rotundiora comprising endosymbionts and endoparasites [5], which infects around 40% of the world's arthropod species [6], with some estimates reaching 66% [7]. *Wolbachia* is famous for manipulating maternal transmission routes, and *Wolbachia* strains capable of reproductive manipulation can sweep through a population by inducing CI, or by favoring vertical maternal

95 transmission via interference with the host's development, resulting in parthenogenesis,
96 feminization, or male death [8].
97
Although numerous ant species are highly invasive and form allopatric populations on a
global scale, nothing is known about PoMRI mechanisms involving endosymbiotic bacteria
in these ecologically dominant insects. In particular, the functional role of *Wolbachia* in ant
biology has thus far remained elusive, in spite of its omnipresence [9]. The globally
distributed tramp ant *Cardiocondyla obscurior* has spread from its putative origin in
Southeast Asia and today occurs in fruit tree plantations, city parks and other disturbed
habitats in the tropics and sub tropics. In addition to *Wolbachia* populations of *C.*
obscurior carry a main bacterial endosymbiont, *Cand.* Westeberhardia, which exhibits
characteristics of an obligate symbiont [10]. Populations of *C. obscurior* from Brazil (BR) and
Japan (JP) show remarkable genomic heterogeneity and differ in a number of phenotypic
traits [11]; nevertheless, mating between queens and males from the two populations readily
takes place in the laboratory. However, a phenotype indicative of PoMRI was found in
hybrid crosses between BR and JP populations, with BR queens mated to JP males
exhibiting shorter lifespans and lower fecundity than BR queens mated to BR males [12]. At
the time, this was attributed to BDM incompatibility of fast evolving sexual traits such as
seminal fluid proteins, as suggested by an up-regulation of genes involved with immune
response in outbred BR queens [12].
115
Here, we show that the negative effect on queen fitness is in fact caused by incompatible
strains of *Wolbachia* harbored by the two populations. By reducing *Wolbachia* levels in
males using antibiotic treatment, we demonstrate that queen fecundity can be rescued.
Genomic analyses of the *Wolbachia* strains isolated from the two populations reveal that
both strains belong to *Wolbachia* supergroup A, which typically contains strains capable
of reproductive manipulation. Surprisingly, CI is only induced by one of the two strains,
and this may be linked to functional differences in CI-associated loci. By discovering the
first case of CI between two ant populations, our study provides the first description of
functional *Wolbachia* infection dynamics in ants and presents a novel perspective on the
biology of globally distributed ants.
126
Results
Unidirectional hybrid incompatibility
We verified the results from [12] by crossing queens and males from laboratory colonies of
BR and JP populations collected in 2009 and 2010, respectively [11]. We conducted this
experiment once in 2015 with the same outcome (Figure S1, Supplement Methods) and
again in 2017/18, the results of which are presented here. As expected, compared to all
other crosses the combination BR x JP (queen x male) resulted in lower fecundity as
estimated by mean weekly egg number over the first six weeks after initiation of egg
laying, regardless of whether the male was infected with *Cand.* Westeberhardia or not
(Figure 1, Kruskal Wallis rank sum test, $X^2 = 78.04$, df=6, p<0.001, see Table S1 for
Bonferroni-Holm corrected pairwise p-values). Similar results were obtained when
maximum weekly egg numbers were compared (Figure S2, Kruskal Wallis rank sum test,
$X^2 = 81.10$, df=6, p<0.001, see Table S2 for Bonferroni-Holm corrected pairwise p-values).
Strikingly, the effect was even stronger when using males naturally uninfected with *Cand.*

Westeberhardia, ruling out the possibility that this endosymbiont plays a role in the
induction of reproductive isolation.
143
We monitored a subset of inter-population crosses for an additional 6 weeks (12 weeks in
total). Of the 53 BR x JP colonies that were monitored for 12 weeks, nine failed to produce
diploid offspring (queen/worker pupae), and eight produced no pupae at all. We
dissected the spermathecae of five of the nine male-only producing queens, and all
contained sperm. All of the JP x BR colonies that were monitored for 12 weeks produced
pupae, and they produced more pupae than BR x JP colonies, irrespective of whether
males were infected with *Cand*. Westeberhardia (Figure 2A; Kruskal-Wallis rank sum test,
X^2=39.80, df=4, p<0.001, see Table S3 for Bonferroni-Holm corrected pairwise p-values).
Furthermore, sex ratios of produced pupae were more male-biased in BR x JP colonies
compared to JP x BR colonies (Figure 2B; binomial GLM: $T_{4,78}$=38.59, p<0.001, see Table
S4 for Bonferroni-Holm corrected pairwise p-values), an effect caused entirely by the lack
of developing females (Figure S3; Kruskal Wallis rank sum test, X^2=5.61, df=4, p=0.23).
Together, these data show consistent incompatibility between BR queens and JP males.
157

Cytoplasmic incompatibility
Cytoplasmic incompatibility (CI) occurs when an endosymbiont favors its transmission by
making infected males incompatible with uninfected females, while infected females can
mate with both infected and uninfected males. A necessary test to verify CI is 'curing' the
host of the putative CI-causing bacteria using antibiotic treatment, which should result in
the 'rescue' of the wild-type phenotype.
164

We fed JP^{rif+} colonies the antibiotic rifampicin diluted in honey every other week for 10
weeks, which resulted in the complete loss of the main endosymbiont *Cand*.
Westeberhardia (Figure S4; Mann-Whitney-U-Test, W=34, df=1, p<0.001) and a significant
reduction of *Wolbachia* levels in workers (Figure S4; Mann-Whitney-U-Test, W=95, df=1,
p<0.01). Treating colonies with the antibiotic tetracycline did not have a negative effect on
Wolbachia (Figure S5; Mann-Whitney-U-Test, X^2=24, df=1, p=0.052). While *Cand*.
Westeberhardia is permanently eradicated in subsequent generations the negative effect
of rifampicin on *Wolbachia* is transient, with levels returning to normal 6 months after
termination of treatment (Figure S6; *Cand*. Westeberhardia: Mann-Whitney-U-Test, W=64,
df=1, p<0.001; *Wolbachia*: Mann-Whitney-U-Test, W=45, df=1, p=0.19). Five of 28
rifampicin-treated colonies succumbed to the treatment and died during the 10-week
period, while all 28 control colonies survived.
177

Mating JP^{rif+} males collected shortly after ceasing antibiotic feeding with BR queens
resulted in significantly rescued fecundity, although mean and maximum weekly egg
numbers did not quite reach levels of BR and JP inbred queens (Figure 1, Table S1). This
rescue effect was evident regardless of whether JP^{rif+} males came from *Cand*.
Westeberhardia infected or uninfected colonies. Rescued BR x JP^{rif+} colonies also
produced more pupae over a 12-week period than their non-rescued BR x JP counterparts
(Figure 2A), and sex ratios among pupae were as female-biased as in JP x BR crosses
(Figure 2B). Strongly reduced *Wolbachia* levels in JP^{rif+} males (Figure 3, Mann-Whitney-U-
Test, W=163, df=1, p<0.001) support the role of *Wolbachia* in inducing hybrid
incompatibility.

We confirmed unidirectional CI with males from laboratory colonies of a second Asian
population collected in Taiwan (TW), which is infected with a *Wolbachia* strain that
amplifies a *wsp* sequence (coding for the hypervariable *Wolbachia surface protein*)
identical to those from JP samples (Figure S7). We set up crosses with queens from BR (BR
x TW) and JP (JP x TW), and the reciprocal combination (TW x JP). Again, only the
combination BR x TW resulted in reduced fecundity, both when mean weekly egg
numbers (Figure S8; Kruskal Wallis rank sum test, X^2=9.96, df=2, p<0.01, see Table S5 for
Bonferroni-Holm corrected pairwise p-values) and maximum weekly egg numbers over six
weeks (Figure S8; Kruskal Wallis rank sum test, X^2=10.19, df=2, p<0.01, see Table S6 for
Bonferroni-Holm corrected pairwise p-values) were compared.

Wolbachia genomics and phylogeny
We assembled the *Wolbachia* genomes from BR (*w*Cobs-BR) and JP (*w*Cobs-JP)
populations of *C. obscurior* using Illumina HiSeq2000 100-bp reads of 200 and 500-bp
insert paired-end libraries [11]. The genomes of both *Wolbachia* were assembled to 148 (BR)
and 182 (JP) scaffolds (Table 1). Both strains showed similar genome sizes and G+C
contents, with an estimated 1,013 (BR) and 1,058 (JP) predicted protein-coding genes. An
analysis on shared genes revealed that both strains share a core of 907 proteins, with 85
unique to *w*Cobs-BR and 109 unique to *w*Cobs-JP. According to Bayesian phylogenetic
analyses of ribosomal proteins from 23 *Wolbachia* strains, both *Wolbachia* strains belong
to supergroup A, which together with supergroup B contains strains capable of
reproductive manipulation. *w*Cobs-JP was found to be more closely related to *w*Ha and
*w*CauA strains, while *w*Cobs-BR was phylogenetically closer to *w*Au and *w*Mel (Figure 4A).
A synteny analysis revealed that both *w*Cobs-JP and *w*Cobs-BR strains keep general
synteny, but with several rearrangements (Figure 4B). This is consistent with previous
observations of group A *Wolbachia* strains [13].

Wolbachia phenology
To compare infection titers between populations we performed qPCR of the *Wolbachia*
gene *Cytochrome c oxidase subunit 1* (*coxA*) against the ant housekeeping gene
elongation factor 1-alpha 1 (*EF1*). JP queens had higher relative *w*Cobs-JP titers than BR
queens infected with *w*Cobs-BR (Figure S9, Mann-Whitney-U-Test, W=23, df=1, p=0.043).
Likewise, JP workers showed higher infection titers with *w*Cobs–JP compared to BR
workers infected with *w*Cobs-BR (Figure S9, Mann-Whitney-U-Test, W=16, df=1, p=0.017).

To test for morph-bias in infection we compared *coxA* expression in adult queens,
workers, winged males and wingless males using qPCR. *coxA* levels did not differ between
sexes and were consistently higher in winged morphs (queens, winged males) compared
to wingless morphs (workers, wingless males) (Figure S10, Kruskal Wallis rank sum test,
X^2=19.53, df=3, p<0.001).

To measure the potential cost of *Wolbachia* infection and to test for a selective loss in
workers, we compared infection levels of workers and queens over time. In contrast to
Cand. Westeberhardia [10], *Wolbachia* levels quickly reached a plateau in both queens and
workers. There was no indication that individuals lose *Wolbachia* over the first few weeks

of their lives, which typically last for up to three (workers) or six months (queens) (Figure
S11, workers: ANOVA, $F_{4,26}$= 2.69, p=0.059; queens: ANOVA, $F_{5,50}$= 15.95, p<0.001).
236
CI mechanism
CI usually results in the failure to produce viable zygotes but how CI operates functionally
is still not clear. The two championed models are based on 1) a sperm modification step
during spermatogenesis which is reversed by a rescue factor or 2) the transfer of a toxic
product via the sperm and subsequent binding to a female-derived antidote [14-17]. The
current genetic view for CI in *Drosophila* favors a two-by-one model [17], involving two
adjacent genes (coined *cifA* and *cifB*) located in a region of prophage origin in *w*Mel [16-21],
which contains several genes with eukaryote function [22]. Expression of both *cifA* and *cifB* in
males is required to cause CI, while expression of *cifA* rescues embryo mortality in females
[21]. Phylogenetic analyses of the Cif proteins have revealed at least 5 distinct phylogenetic
clades referred to as Types 1-5, respectively [18,23,24].
248
We found homologues of *cifA* and *cifB* in *Wolbachia* strains from BR and JP. In the latter,
we found two additional homologues. Through Bayesian phylogenetic inference, and
following the classification scheme of [23], we found that all copies of *cifA* and *cifB* belong to
Type I (Figure 5A, B). The *cifA* and *cifB* genes found in NODE_002 of *w*Cobs-JP are
closely related to those found in *w*Cobs-BR and in the cryptic phage of the strain *w*Sol
isolated from the fig wasp *Ceratosolen solmsi* (Figure 5A, B). The *cifA* and *cifB* genes
found in NODE_063 of *w*Cobs-JP show a more distant relationship to their homologues in
*w*Cobs-BR. Following [23], we performed a protein remote homology search with the
webserver of HHpred's v3.2.0 webserver [25]. While the *cifA* gene of *w*Cobs-BR contained a
DUF249 domain, neither *cifA* homologues in *w*Cobs-JP showed homology to known
domains (Figure 5C). In contrast, putative domains were found in all *cifB* genes. These
shared a putative nuclease domain (PD-(D/E)XK nuclease/DpnII-MboI) found in other type
I *cifB* genes. The *cifB* homologue found in NODE_063 in *w*Cobs-JP shared most domains
with the *cifB* homologue in *w*Cobs-BR, with an additional peptidase domain
(Peptidase_C58; PF03543).
264
Wolbachia in males
As mentioned above, *Wolbachia*-induced CI has often been linked to sperm modification
during spermatogenesis [26] or the transfer of a CI toxin along with or in the sperm. The
presence of *Wolbachia* in the testes is thus expected, although their specific localization in
spermatocytes or spermatids might not be a prerequisite for inducing CI, as has been
observed in the wasp *Nasonia vitripennis* [27]. We used fluorescence in situ hybridization to
localize *Wolbachia* in the abdomens of male *C. obscurior*, confirming the presence of the
bacteria throughout different tissues including the testes in both JP (Fig. 6A-C) and BR
(Figure 6D) individuals. *Wolbachia* was also observed co-infecting bacteriocytes
dominated by *Cand*. Westeberhardia, yet in significantly lower densities (Figure 6C).
275
Discussion
Insect-*Wolbachia* interactions can range from beneficial to detrimental and from
facultative associations to rare cases of ultimate mutualisms where hosts have higher
fitness with *Wolbachia* than without [5]. This prevalence and diversity of *Wolbachia*
associations predictably also occurs in ants; however, our knowledge of ant-*Wolbachia*

interactions is limited [9]. In the few ant-*Wolbachia* associations studied so far, infection is
facultative (e.g. [28-30]), indicating ongoing arms races between hosts and symbionts [31]. One
particularly obvious question to ask is whether *Wolbachia* has the power to control sex
allocation, a trait shaped by kin selection in social Hymenoptera [32]. In this context,
facultative *Wolbachia* infections have been reported to cause transient changes in sex
allocation [29], or to have no effect [33]. Similarly contrasting results have been obtained
regarding effects on colony productivity [29,34]. It has furthermore been speculated that
Wolbachia can bias caste ratios towards the production of queens [28,35], but current data
are scarce and provide little support for this hypothesis [9]. By providing the first functional
description of an ant-*Wolbachia* association involving two strains of *Wolbachia*, our study
lays the groundwork for in-depth investigations into how *Wolbachia* affect ant
reproductive biology and fitness.
293
Wolbachia clearly play a significant role in the biology of *C. obscurior*. First, all screened
individuals were infected regardless of population, caste or sex. *Wolbachia* titers were
higher in queens and winged males compared to workers and wingless males, a
difference that is likely associated with the relative amounts of reproductive tissue as *C.
obscurior* workers lack ovaries and winged males have larger testes than wingless males
[36]. Second, rifampicin treatment merely reduced *Wolbachia* titers, and this effect was
reversible. This was not the case for *Cand.* Westeberhardia, which was permanently
cleared by rifampicin treatment. Third, *Wolbachia* titers did not decline with age in
workers. *Wolbachia* infection can be costly and selection may thus favor reduced infection
loads in sterile workers, as these do not contribute to vertical transmission [28,35]. Such a
mechanism appears to be acting on *Cand.* Westeberhardia, where titers in workers (but
not queens) decreased significantly with age, indicating that while the host may benefit
from infection during development or in the adult egg-laying queen stage, the symbiont is
not required [10]. Consistently high titers in workers indicate that *Wolbachia* do not have
deleterious effects on worker performance in *C. obscurior*, unlike *Wolbachia* in
Acromyrmex ants [37]. To what degree the population-specific strains differ in their effect on
host phenotype remains unclear. Higher *Wolbachia* titers in JP individuals suggest that the
*w*Cobs-JP strain may be better integrated into the host's biology. Alternatively, these
differences may be driven by population-specific competition with *Cand.* Westeberhardia
over limited resources.
314
Sex allocation in *C. obscurior* is female-biased [38] and queens presumably have full control
over male production, as investment into males increases in the presence of competing
queens [39-41], Accordingly, we found no indication that *Wolbachia* influences sex allocation.
Queens from non-CI (JP x BR) crosses produced few males in the first 12 weeks of their
lives. This may be a consequence of the short observation period, equaling ~50% of the
average queen lifespan [38]. The sex ratios produced by CI (BR x JP) crosses showed a
bimodal distribution, with several queens matching sex ratios produced by non-CI queens
(albeit with lower overall productivity), while others produced only males. On average,
however, CI queens did not produce more males than non-CI queens. There was also no
indication that *Wolbachia* altered caste ratios towards production of queens (data not
shown). Concordantly, a previous study comparing investment into sexuals and workers
did not find differences between the BR and JP populations [38]. Testing directly for a caste-
biasing effect of *Wolbachia* requires complete elimination of the bacterium. While this

cannot be done without also killing the ant host, preliminary trials indicate that cross-
infection of the two *Wolbachia* strains is possible. In the future, such manipulations may
help clarify whether *Wolbachia* have caste-manipulating power in *C. obscurior*.
331
Global exchange of stowaway ants leading to secondary contact between allopatric
populations is common (e.g. in the tropical fire ant *Solenopsis geminata* [42], and the little
fire ant *Wasmannia auropunctata* [43]) and thus the presented scenario, albeit conceived in
the laboratory, is not an unrealistic one. A previous study emphasized the role of BDM
incompatibility between allopatric *C. obscurior* populations in causing a CI-like phenotype
[12]. *Wolbachia*-induced incompatibilities were ruled out because of identity of a partial
sequence of *wsp* extracted from seven individuals (three from BR colonies collected in
2004, four from JP colonies collected in 2005) using the wsp81F/wsp691R primer
combination [44]. Gene expression comparisons of allopatrically and sympatrically mated BR
queens pointed towards elevated immune responses and other changes that mimicked a
CI-like phenotype. In *Drosophila*, *Wolbachia* can manipulate expression of key regulatory
genes during spermatogenesis, which leads to embryonic lethality and a CI-like
phenotype [45]. Thus, what appeared to be BDM incompatibility may have been caused by
Wolbachia inducing a CI-like phenotype via interference with queen gene expression. The
alternative and more parsimonious explanation for the CI-like phenotype based on the
results from the present study is a classic unidirectional *Wolbachia*-driven CI effect
involving incompatible *Wolbachia* strains. This also explains why JP x BR crosses did not
elicit a CI phenotype. The overexpression of immunity-related genes in allopatrically
mated BR queens detected by Schrempf et al. 2015 may simply reflect physiological
responses to aborted oogenesis, as opposed to responses to ejaculate quality, or to
injuries caused by non-matching male genitalia. Furthermore, we recently found that some
BR colonies were infected with an additional *Wolbachia* strain displaying partial *wsp*
sequences identical to those of *w*Cobs-JP (data not shown), indicating horizontal
transfection in our rearing facilities, especially in older laboratory colonies.
356
The mechanism of CI in *C. obscurior* appears to align along the *Wolbachia cif* gene axis
described in *Drosophila* fruit flies and *Culex* mosquitoes [15,17,19,23,46]. This genetic region,
which is of prophage origin and capable of manipulating the eukaryote host [18], showcases
the importance of using a hologenomic approach to study evolution [47]. Similar prophage-
derived regions containing pairs of homolog *cif*/*cif*-like genes have since been described
in a number of *Wolbachia* strains [23,24]. Surprisingly, the homologs of *cifB* in *w*Pip and *w*Mel
have different biochemical functions: both putatively contain ankyrin-like repeats for
protein-protein interaction with eukaryotic DNA [46] but $cifB_{wMel}$ is not a putative
deubiquitinase [16], as postulated for $cifB_{wPip}$[19]. Both *Wolbachia* strains isolated from *C.*
obscurior have homologs of *cifA* and *cifB* but only the *w*Cobs-JP strain has an additional
cifA/*B* pair, pointing towards a role for the latter pair in inducing reproductive isolation.
Whether the second *cifA*/*B* containing region was lost in *w*Cobs-BR, or whether it stems
from a duplication event and subsequent divergence in *w*Cobs-JP as has been described
for the *w*AlbB strain of *Aedes albopictus* [48], remains to be shown. Alternatively, functional
differences in *cifA*/*B* homologs between *w*Cobs-JP and *w*Cobs-BR, or an interaction of the
two *cif* gene products in JP males could be causing CI. Irrespective of the exact CI
mechanism, we do not know why a mere reduction of *Wolbachia* titers in males was
sufficient to rescue queen fecundity. This requires studying the effects of *Wolbachia* and

antibiotic treatment on sperm production and development. One possibility is that
rifampicin treatment causes ribosomal damage that prevents *Wolbachia* from modifying
sperm. At the same time, antibiotic-induced damage to sperm mitochondria [49] may
explain the slightly lower productivity of "CI rescue" crosses compared to non-CI crosses.
It is also unclear why the CI effect was stronger on egg numbers but not pupae numbers
when using males from colonies naturally uninfected with *Cand.* Westeberhardia, but this
could point toward interaction effects between the two symbionts.
382
Our study raises many questions about the evolutionary history of the *C. obscurior-*
Wolbachia association. The putative origin of *C. obscurior* in Southeast Asia, together with
sequence identity between *Wolbachia* strains isolated from populations found in Japan
and Taiwan, suggest that these strains represent the ancestral association. Geographic
patterning separating New from Old World *Wolbachia* strains has been observed
previously in ants [50], and frequent horizontal transmissions [51,52] can lead to a surprising
diversity within species [53]. *C. obscurior* and its sister species *C. wroughtonii* are the only
known representatives of the 100+ described species in the genus *Cardiocondyla* (the so-
called heart-node ants) that have adopted an arboreal lifestyle, as opposed to nesting in
soil and rock crevices. This makes them highly susceptible to horizontal transmission of
Wolbachia from other plant-associated insects. Although *Wolbachia* from different
supergroups may co-exist in a host [13], it is unlikely that two strains from the same
supergroup co-exist without eventually recombining and merging. Generally,
unidirectional CI should facilitate introgression between populations, while bidirectional
CI should promote divergence. This predicts mixed *C. obscurior* genotypes but only the
presence of the *w*Cobs-JP strain in secondary contact zones, assuming that hybrids
perform as well as parental lineages. How then has the ancestral Asian strain been
replaced? *Wolbachia* acquisition and persistence in *C. obscurior* may be driven by
selection on the strain that is better adapted to the local environment, perhaps analogous
to *Wolbachia*-driven thermal preferences in *Drosophila* flies [54,55]. Resolving how *w*Cobs-BR
replaced the putatively ancestral *w*Cobs-JP strain, which holds the power of CI, requires
testing how mixed combinations of host and *Wolbachia* perform under varying ecological
conditions. Coupled with a global survey of *Wolbachia-C. obscurior* associations, this will
help unravel how the symbiosis shapes the worldwide distribution of this tramp ant.
407
Conclusion
One important consequence of unidirectional reproductive isolation is unequal
inheritance, which allows a BDM-causing locus or a reproductive manipulator to quickly
spread in secondary contact zones. Such disruptive factors can be more powerful than
natural selection on other fitness-relevant traits, and lead to the evolution of PoMRI and
hybrid inviability, processes estimated to require several million years in *Drosophila* flies
and *Pogonomyrmex* ants [56,57]. Our discovery of reproductive isolation between allopatric
ant populations, driven by two naturally occurring, closely related *Wolbachia* strains that
differ in their genomic makeup, demonstrates that these evolutionary processes can be
observed in real time and lays the groundwork for detailed studies of *Wolbachia*-ant
biology. In particular, a system in which CI is accompanied by a CI-like phenotype in host
gene expression may prove useful for understanding the function of the *cif* gene family.
More generally, hologenomic approaches such as the one presented here will help tackle

the challenges associated with human-induced changes to ecological and evolutionary
dynamics.

Methods

Cardiocondyla obscurior presumably originates from Southeast Asia and is now
distributed around the tropics and subtropics. Queens in *C. obscurior* are produced year-
round and have a short generation time of ~120 days (from egg to queen to egg).
Colonies usually contain 2-5 queens [11] and a single wingless male that can live up to a few
months and continually produces sperm [58]. Such males transfer on average 1000 sperm
cells per copulation [59], which is sufficient for a lifelong production of diploid offspring by
an inseminated queen. Newly hatched queens can decide to stay in the maternal nest or
to leave - alone or with a small group of workers - irrespective of mating status. Like all
Hymenoptera, ants exhibit haplodiploid sex determination and virgin females can
produce males from haploid, unfertilized eggs. Virgin queens in *C. obscurior* can copulate
with their own sons (pers. obs.), enabling the production of workers and queens, similar to
a related *Cardiocondyla* species [60]. This biology has facilitated the worldwide spread of
several species of the genus [61].

In this study, we used inbred laboratory 'populations' derived from colonies collected in
Bahia, Brazil (BR) in 2009 (Brazilian Ministry of Science and Technology, permits 20324-
1/40101-1) and Okinawa, Japan (JP) in 2010. The JP population exhibits a natural
polymorphism regarding the presence of the endosymbiont *Candidatus* Westeberhardia
cardiocondylae, with some colonies being naturally uninfected (JP$_{we-}$) [10]. Although the BR
and JP populations differ in cuticular hydrocarbon composition, queen size and behavior
[11], workers are morphologically indiscernible according to exploratory data analyses by
hierarchical and nonhierarcical, vector-based forms of Nest Centroid clustering: NC-Ward,
NC-part.hclust, NC-part.kmeans, NC-NMDS-kmeans [62,63]. These analyses considered 15
continuous morphometric characters. Specifically, when considering 82 cosmopolitan nest
samples of the sister taxa *C. wroughtonii* and *C. obscurior*, all JP and BR samples were
allocated to the latter species by all four types of NC-clustering. This result was confirmed
with posterior probabilities of $p > 0.98$ if these samples were run as wild-cards in a linear
discriminant analysis. Analyses of 32 nest samples representing only the *C. obscurior*
cluster could not allocate the JP and BR populations to different intraspecific clusters nor
could these analyses demonstrate any intraspecific structuring at all. This suggests that
Wolbachia-induced reproductive isolation has not yet led to a divergence into
morphospecies. In addition to JP and BR colonies, we used colonies collected from a
second Asian population in Taipei, Taiwan in 2013 (TW) to confirm the presence of
unidirectional CI. For all experiments, colonies were kept in square plaster-bottom nests
(100 x 100 x 20 mm, Sarstedt, Germany) and held in climate chambers under a 12h/12h
light/dark cycle and a 28°C/22°C temperature cycle. Food (honey and *Drosophila* or
pieces of *Periplaneta americana*) and water were provided to stock colonies every 3 days.
All animal treatment guidelines applicable to ants under international and German law
were followed. Collection of colonies that form the basis of the laboratory population used
in this study was permitted by the Brazilian Ministry of Science and Technology (RMX
004/02). No other permits were required for this study.

Antibiotic treatment

Rifampicin is a broad-spectrum antibiotic that acts via inhibition of bacterial RNA
polymerase [64]. To produce *Wolbachia*-cured males, we split large *Cand.* Westeberhardia-
infected (JP) and uninfected (JP$_{we-}$) stock colonies into two equal halves. Half of these splits
were designated as controls (JP: n=13, JP$_{we-}$: n=5), while the other half was treated with
antibiotics (JP: n=13, JP$_{we-}$: n=5). We diluted 0.0025 g of solid rifampicin (Sigma-Aldrich,
USA) in 500 µl of a 1:1 honey-water solution for a final antibiotic concentration of 0.5%.
The antibiotic solution was placed on a shaker for 4 hours to ensure complete mixing, and
subsequently covered with aluminum foil and stored at -20°C. Treated colonies were fed
with the antibiotic solution twice per week every other week, for a total duration of ten
weeks. On the days following treatment, the antibiotic solution was removed from the
nest. Between antibiotic feedings (i.e. every other week), colonies were fed with honey
and pieces of autoclaved cockroach (to prevent bacterial re-infections) and water ad
libitum. Control colonies were fed twice a week with honey and cockroaches and received
water ad libitum.
482
483 **_Wolbachia_ and _Cand._ Westeberhardia infection**
484 We assessed the presence and titers of bacterial infections using PCR and qPCR,
485 respectively, on DNA extracted from whole bodies (worker pupae & adults, queen pupae
486 & adults, adult males). DNA was extracted from samples using a standard CTAB protocol.
487
Presence of infection
We confirmed infection in a subset of workers from BR and JP colonies. Infection with
Wolbachia was tested with a PCR assay using primers specific for a fragment of the
Wolbachia surface protein gene of the BR (105-bp, *wsp_wCobs_BR-for:* 5-
'*TAAATCTTGCATCTGTAACATT-3'*, *wsp_wCobs_BR-rev:* 5'-
CTGCGGATACTGATACAACTACTG-3') and JP (254-bp, *wsp_wCobs_JP-for:* 5'-
CATTTTGACTACTCACAGCGGTTG-3', *wsp_wCobs_JP-rev:*
5'*CTGCGGATACTGATACAACTACTG-3'*) strains. *Cand.* Westeberhardia infection was
tested by amplifying a 204-bp fragment of the *ribonucleoside-diphosphate reductase 1*
subunit beta gene of *Westeberhardia* (*WEOB_403*) (*nrdB_weCobs-for:* 5'-
GGAAGGAGTCCTAATGTTGCG-3'; *nrdB_weCobs*-rev: 5'-ACC
AGAAATATCTTTTGCACGTT-3'). A 104-bp fragment of the *C. obscurior* gene *elongation*
factor 1-alpha 1 (*Cobs_01649*) (*EF1-for:* 5'-TCACTGGTACCTCGCAAGCCGA-3'; *EF1-rev:*
5'-AGCGTGCTCACGAGTTTGTCCG-3') was used as a positive control. In addition,
infection was assessed in two adult queens from each of five TW colonies to confirm that
this population carries the *w*Cobs-JP strain. Each PCR reaction contained 5 µL Taq
polymerase (GoTaq, Promega), 3 µL H$_2$O, 0.5 µL forward primer (10 µM), 0.5 µL reverse
primer (10 µM) and 1 µL DNA. PCRs were run at 94°C for 4 minutes followed by 36 cycles
at 94°C for 30s, 57°C for 30s and 72° for 30s, with a final step of 72°C for 10 minutes.
Electrophoresis separation of PCR products was performed on a 1.5% TAE-Gel with 0.5 µl
of 0.001 % Gel-Red for 45 minutes at 60 V and 65 mA.
509
Population-specific Wolbachia titers
We compared *Wolbachia* infection titers of adult workers and queens from BR (n=9
workers, n=10 queens) and JP (n=10 workers, n=9 queens) colonies by amplifying a
portion of the *Cytochrome c oxidase subunit 1* gene (*coxA_wCobs-for:* 5'-
TTGGTCATCCAGAAGTTTACGT-3', *coxA_wCobs-rev:* 5'-TGAGCCCAAACCATAAAGCC-

3') in qPCR. As a relative standard *elongation factor 1-alpha 1* (*Cobs_01649*) was used.
Each qPCR reaction contained 5 µL SYBR (Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1
µL reverse primer (2 µM) and 1 µL DNA. Reactions were performed in duplicates and run
at 95°C for 3 minutes followed by 40 cycles at 95°C for 5s, 60°C for 20s, and 95° for 10s,
and a final melt curve analysis step in which reactions were heated from 65°C to 95°C in
5°C steps. Single-amplicon production was confirmed with melt curve analyses and
relative *Wolbachia* titers calculated with the $2^{-\Delta CT}$ method [65].
522
Age and morph-specific Wolbachia titers
We quantified *Wolbachia* titers in individual queens and workers from BR colonies by
determining relative *coxA_wCobs* copy numbers with qPCR across developmental stages
and adult age classes. In workers, titers were measured in pupae (n=4 white pupae, n=5
dark pupae) and in adults two days (n=5), 14 days (n=7), and 28 days (n=6) post-hatching.
In queens, titers were measured in pupae (n=5 white pupae, n=4 dark pupae), as well as in
adults two days (n=7), 14 days (n=7), 28 days (n=10 virgin queens, n=7 mated queens)
and 48 days (n=9) post-hatching. In addition, we quantified *coxA_wCobs* copy numbers in
adult morphs (n=15 queens, n=12 workers, n=6 winged males, n=7 wingless males). For
all reactions, *elongation factor 1-alpha 1* (*Cobs_01649*) was used as a housekeeper. Each
qPCR reaction contained 5 µL SYBR (Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1 µL
reverse primer (2 µM) and 1 µL DNA. Reactions were performed in triplicates and run as
described above. Single-amplicon production was confirmed with melt curve analyses and
relative quantities of *Wolbachia* calculated with the $2^{-\Delta CT}$ method.
537
Rifampicin effects on infection titers
We verified the efficacy of rifampicin treatment by measuring *Wolbachia* and
Westeberhardia titers in dark worker pupae produced by antibiotic-treated (n=21 from 9
colonies) and control colonies (n=21 from 8 colonies) using qPCR of *coxA_wCobs and*
nrdB_weCobs. In addition, we measured the *Wolbachia* titers of JP males used in mating
combinations BR x JP and BR x JP^{rif+} (JP: n=15, JP^{rif+}: n=12) with qPCR of a portion of the
JP-specific *wsp* gene (*wsp_wCobs_JP*). For all samples, *elongation factor 1-alpha 1*
(*Cobs_01649*) was used as a housekeeper. Each qPCR reaction contained 5 µL SYBR
(Peqlab), 2 µL H_2O, 1 µL forward primer (2 µM), 1 µL reverse primer (2 µM) and 1 µL DNA.
Reactions were performed in duplicates (males) or triplicates (worker pupae) and run as
described above (except with an annealing temperature of 57°C for male pupae). Single-
amplicon production was confirmed with melt curve analyses and relative quantities of
Wolbachia and *Westeberhardia* calculated with the $2^{-\Delta CT}$ method.
551
Experimental crosses
We designed seven mating combinations between queens and males from the two
populations: 1) JP x JP (n=7), 2) JP x BR (n=26), 3) BR x BR (n=11), 4) BR x JP (n=36), 5) BR
x JP$_{we-}$ (n=29), 6) BR x JP^{rif+} (n=13) and 7) BR x JP$_{we-}^{rif+}$ (n=5). Each experimental colony was
set up with one dark queen pupa, one freshly hatched male, 10 workers and brood (10
eggs, 10 larvae, 10 worker pupae). All individuals were collected from un-manipulated
stock colonies, except for JP^{rif+} and JP$_{we-}^{rif+}$ males used in combinations #6 and #7, which
were collected from rifampicin-treated colonies. To avoid manipulation effects, all
combinations were set up in parallel. In such experimental colonies, mating between
queens and males usually occurs within three weeks, after which queens shed their wings

and being laying eggs. We monitored colonies for queen wing loss once per week. Once
queens had shed their wings, we removed any brood placed in the colony during set up
and standardized worker number to 20. Colonies were then monitored once a week for six
weeks and all eggs and pupae (queen, worker, wingless male, winged male) counted. A
subset of colonies was monitored for a total of 12 weeks (JP x BR: n=19, BR x JP: n=28, BR
x JP$_{we-}$: n=25, BR x JP$_{we-}^{rif+}$: n=9, BR x JP$_{we-}^{rif+}$: n=5). In each colony, the number of adult
workers was kept constant at 20 individuals by removing hatched workers or adding
workers from stock colonies. From monitoring data, we calculated mean and maximum
weekly egg numbers, and total number and sex ratio of produced pupae (male
pupae:total pupae).
572
573 To verify that CI is specifically induced by the *w*Cobs-JP strain when it encounters the
574 *w*Cobs-BR strain, we set up three additional mating combinations between queens and
575 males using a third population from Taiwan (TW) which also harbors the *w*Cobs-JP strain:
576 1) JP x TW (n=5), 2) TW x JP (n=6), 3) BR x TW (n=5). Experimental colonies were set up
577 and monitored as described above for six weeks.
578
DNA sequencing and genome assembly
We extracted DNA from a pool of 26 (JP) and 30 (BR) males from two colonies, collected
in 2009 in Una, Brazil (BR2009-alpha) from aborted fruits of coconut trees and in 2010 in
Naha, Japan (JP2010-OypB) from under bark of coral trees (*Erythrina* sp.). For details see
[11]. Briefly, the reads were right-tail clipped (minimum quality of 20) and all reads with
undefined nucleotides were removed using a combination of FASTX-Toolkit v0.0.14
(http://hannonlab.cshl.edu/fastx_toolkit/, last accessed December 3, 2010) and
PRINSEQ++ v1.2 (10.7287/peerj.preprints.27553v1). The resulting paired end reads were
assembled with SPAdes v3.13.1 (-k 33,55,77 --only-assembler [66]. The resulting contigs
were then binned according to an *ad hoc* BLASTX-based method. First, a proteome
database was built from representative genomes (Table S7). Second, the best hit for each
scaffold was used for assigning a preliminary bin and a coverage cut-off (10x for BR and
30x for JP) was implemented based on the coverage of the longest contigs. Third, each
scaffold was manually inspected using the online NCBI web-server *vs.* BLAST+2.9.0. If the
best hits to proteins were consistently to *Wolbachia* bacteria across the scaffold, the query
sequence was retained. Finally, the assembly graph was loaded into Bandage v0.8.1 [67],
the graph(s) matching *Wolbachia* sequences were identified and the sequences were
extracted. These extracted sequences were then compared to the *Wolbachia* draft bin. In
both cases, the extracted sequences were roughly the same compounded length as the
manually-curated bins and added only a couple of kilo base pairs (kbp), suggesting near
completeness of the genome. Using these *Wolbachia* bins, we performed a mapping
using BOWTIE2 v2.3.4.1 [68] followed by a re-assembly of the mapped reads in SPAdes
(with previously-mentioned parameters). This final reassembly was filtered by both
coverage and manual screening of scaffolds shorter than 3 kbp. Assembled sequences
have been deposited in the European Nucleotide Archive (ENA) under the accessions
CACTIU010000000.1 (*w*Cobs-BR) and CACTIV010000000.1 (*w*Cobs-JP).
605
606 Phylogeny and comparative genomics of *Wolbachia* strains
607 A draft annotation of the scaffolds was done using Prokka v1.14 [69]. For comparative
608 genomic purposes, we built a subset of the predicted ORFs by filtering out all proteins

annotated as "hypothetical proteins" with genes shorter than 300 base pairs (bp). This helped us avoid wrongly annotated ORFs overlapping pseudogenes. To infer the phylogenetic relationship between *Wolbachia* strains found to infect the analysed BR and JP populations as well as among other *Wolbachia* bacteria, we collected the amino acid sequences for the ribosomal proteins of 23 *Wolbachia* strains from supergroups A (8), B (10), C (2), D (1), E (1), and F (1) (Table S8). Using *Anaplasma phagocytophilum* strain HZ and *Ehrlichia canis* strain Jake as outgroups, we inferred a Bayesian phylogeny using MrBayes v3.2.6 [70]. We ran two independent analyses with four chains each for 300,000 generations and checked for convergence. The substitution model JTT+I+G+F was used as suggested by jModelTest v2.1.10.

To infer the shared protein-coding gene content of *w*Cobs-BR and *w*Cobs-JP strains, we used OrthoMCL v2.0.9 [71,72] with an inflation value of 1.5. The synteny of both strains was assessed using nucmer v3.23 [73] using anchor matches that were unique in both the reference and query (--mum).

Cif-family gene annotation and phylogenetics
cif genes were first identified by BLASTP using the protein sequences WP_010962721.1 (*cifA*) and WP_010962722.1 (*cifB*) as references. Predicted start codons were manually checked for the presence of a Shine-Dalgarno-like sequence. To identify whether the *cif* genes form each strain belonged to the same type or not, we performed a phylogenetic analysis following Lindsey *et. al.* 2018. Briefly, the previously used protein sequences were aligned with the newly acquired ones from both wCobs-BR and wCobs-JP. The alignments were then back-translated to nucleotides. For *cifB*, the type IV genes were excluded and the 3'-end of the alignment was manually truncated to remove the longer unmatched regions of some *cifB* variants. Bayesian phylogenetic inference was performed with MrBayes v3.2.6 with the GTR+G substitution model (as suggested by jModelTest) running two independent analyses with four chains each for 1,000,000 generations, and checked for convergence. Following [23], Cif protein domains were searched for using HHpred's v3.2.0 webserver (https://toolkit.tuebingen.mpg.de/tools/hhpred) with default parameters and the following databases: SCOPe70 v2.07, COG/KOG v1.0, Pfam-A v32.0, and SMART v6.0. hits with a probability >=80% were considered.

All genomic data files are available at the zenodo repository https://doi.org/10.5281/zenodo.3561160

Histological sectioning and fluorescence in situ hybridization
C. obscurior (JP) male abdomens were fixed in 4% formaldehyde in PBS and washed with water followed by a dehydration series in n-Butanol (30%, 50%, 70%, 80%, 90%, 96%) at room temperature with shaking for 1 h at each step. Absolute n-butanol was used for the last three hours at 30°C, exchanging the solution each hour. Dehydrated samples were embedded in Technovit 8100 cold polymerizing resin (Heraeus Kulzer, Germany) according to the manufacturer's instructions, cut into semithin sections (8 μm) with a Microm HM355S microtome (Thermo Fisher Scientific, Germany) and mounted on glass slides coated with poly-L-lysine (Kindler, Germany). The tissue sections were incubated for 90 min at 50°C in hybridization buffer (0.9 M NaCl, 20 mM Tris/HCl (pH 8.0), 0.01 % SDS) containing 0.5 μM of the *Wolbachia*-specific probes Wolb_W2_Cy5 (5'-

CTTCTGTGAGTACCGTCATTATC-3') [74] and Wolb_Wol3_Cy5 (5'-
TCCTCTATCCTCTTTCAATC-3'), [75], as well as the *Cand*. Westeberhardia-specific probe
Wcard1_Cy3 (5'- ATCAGTTTCGAACGCCATTC-3') [10]. DAPI (4',6-diamidino-2-
phenylindole) was used for host DNA counterstaining. After hybridization, samples were
washed with buffer (0.1 M NaCl, 20 mM Tris/HCl (pH 8.0), 5mM EDTA, 0.01% SDS) for 20
min at 50°C. Subsequently, the washing buffer was removed and washed with distilled
water for 20 min at 50°C. Samples were air-dried and covered with VectaShield® (Vector
Laboratories, Burlingame, CA, USA) and stored overnight at room temperature. Images
were acquired on an AxioImager.Z1 epifluorescence microscope (Carl Zeiss, Jena,
Germany).

Acknowledgements
The authors thank J. Wang for help collecting colonies in Taiwan and for providing the
wsp sequence of the TW population. We thank J. Hacker for help with the BR x TW
crossings and H. Lowack and J. Wallner for help with lab work. J.D. Shropshire helped with
the *cif* annotation and provided useful comments on a previous version of the manuscript.
B. Weiss assisted with histological preparations. This work was supported by the Marie-
Curie AgreenSkills+ fellowship program cofunded by the EU's Seventh Framework
Programme (FP7-609398) to A.M.M, and a DFG grant He1623/31 to J.H. and J.O.

References
1.   Wetterer, J. K., Wild, A. L., Suarez, A. V., Roura-Pascual, N. & Espadaler, X.
679   Worldwide spread of the Argentine ant, *Linepithema humile* (Hymenoptera:
680   Formicidae). *Myrmecol News* **12,** 187–194 (2009).
681   2.   Ascunce, M. S. *et al.* Global invasion history of the fire ant *Solenopsis invicta*.
682   *Science* **331,** 1066–1068 (2011).
683   3.   Tsutsui, N. D., Suarez, A. V., Holway, D. A. & Case, T. J. Reduced genetic variation
684   and the success of an invasive species. *Proc Natl Acad Sci U S A* **97,** 5948–5953
685   (2000).
686   4.   Brucker, R. M. & Bordenstein, S. R. Speciation by symbiosis. *Trends in Ecology &*
687   *Evolution* **27,** 443–451 (2012).
688   5.   Zug, R. & Hammerstein, P. Bad guys turned nice? A critical assessment of Wolbachia
689   mutualisms in arthropod hosts. *Biological Reviews* **90,** 89–111 (2014).
690   6.   Zug, R. & Hammerstein, P. Still a Host of Hosts for Wolbachia: Analysis of Recent
691   Data Suggests That 40% of Terrestrial Arthropod Species Are Infected. *PLoS ONE* **7,**
692   e38544–3 (2012).
693   7.   Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J. H.
694   How many species are infected with Wolbachia?-A statistical analysis of current
695   data. *FEMS Microbiology Letters* **281,** 215–220 (2008).
696   8.   Werren, J. H. Biology of Wolbachia. *Annu. Rev. Entomol.* **42,** 587–609 (1997).
697   9.   Russell, J. A. The ants (Hymenoptera: Formicidae) are unique and enigmatic hosts of
698   prevalent Wolbachia (Alphaproteobacteria) symbionts. *Myrmecol News* **16,** 7–23
699   (2012).
700   10.   Klein, A. *et al.* A novel intracellular mutualistic bacterium in the invasive ant
701   Cardiocondyla obscurior. *ISME J* **10,** 376–388 (2015).
702   11.   Schrader, L. *et al.* Transposable element islands facilitate adaptation to novel

703   environments in an invasive species. *Nature Communications* **5,** 5495 (2014).

12. Schrempf, A. *et al.* Mating with an allopatric male triggers immune response and
decreases longevity of ant queens. *Molecular Ecology* **24,** 3618–3627 (2015).

13. Ellegaard, K. M., Klasson, L., Näslund, K., Bourtzis, K. & Andersson, S. G. E.
Comparative genomics of Wolbachia and the bacterial species concept. *PLoS*
Genet **9,** e1003381 (2013).

14. Beckmann, J. F. *et al.* Caution Does Not Preclude Predictive and Testable Models of
Cytoplasmic Incompatibility: A Reply to Shropshire et al. *Trends Genetics* **35,** 399–
400 (2019).

15. Beckmann, J. F. *et al.* The Toxin-Antidote Model of Cytoplasmic Incompatibility:
Genetics and Evolutionary Implications. *Trends Genet.* **35,** 175-185 (2019).

16. Shropshire, J. D. & Bordenstein, S. R. Two-By-One model of cytoplasmic
incompatibility: Synthetic recapitulation by transgenic expression of cifA and cifB in
Drosophila. *PLoS Genet* **15,** e1008221 (2019).

17. Shropshire, J. D. *et al.* Models and Nomenclature for Cytoplasmic Incompatibility:
Caution over Premature Conclusions - A Response to Beckmann et al. *Trends*
Genet. **35,** 397–399 (2019).

18. LePage, D. P. *et al.* Prophage WO genes recapitulate and enhance Wolbachia-
induced cytoplasmic incompatibility. *Nature* **543,** 243-247 (2017).

19. Beckmann, J. F., Ronau, J. A. & Hochstrasser, M. A Wolbachia deubiquitylating
enzyme induces cytoplasmic incompatibility. *Nat Microbiol* **2,** 17007-7 (2017).

20. Chen, H., Ronau, J. A., Beckmann, J. F. & Hochstrasser, M. A Wolbachia nuclease
and its binding partner provide a distinct mechanism for cytoplasmic
incompatibility. *Proc Natl Acad Sci U S A* 1–8 (2019). doi:10.1073/pnas.1914571116

21. Shropshire, J. D., On, J., Layton, E. M., Zhou, H. & Bordenstein, S. R. One prophage
WO gene rescues cytoplasmic incompatibility in Drosophila melanogaster. *Proc*
Natl Acad Sci U S A **115,** 4987-4991 (2018).

22. Perlmutter, J. I. *et al.* The phage gene wmk is a candidate for male killing by a
bacterial endosymbiont. *PLoS Pathog* **15,** e1007936 (2019).

23. Lindsey, A. R. I. *et al.* Evolutionary Genetics of Cytoplasmic Incompatibility Genes
cifA and cifB in Prophage WO of Wolbachia. *Genome Biology and Evolution* **10,**
434-451 (2018).

24. Bing, X.-L., Zhao, D.-S., Sun, J.-T., Zhang, K.-J. & Hong, X.-Y. Genomic Analysis of
Wolbachia from Laodelphax striatellus (Delphacidae, Hemiptera) Reveals Insights
into Its 'Jekyll and Hyde' Mode of Infection Pattern. *Genome Biology and Evolution*
12, 3818-3831 (2020).

25. Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics Toolkit with
a New HHpred Server at its Core. *J. Mol. Biol.* **430,** 2237-2243 (2018).

26. Veneti, Z. *et al.* Cytoplasmic incompatibility and sperm cyst infection in different
Drosophila-Wolbachia associations. *Genetics* **164,** 545-552 (2003).

27. Clark, M. E. *et al.* Wolbachia modification of sperm does not always require
residence within developing sperm. *Heredity* **101,** 420-428 (2008).

28. Wenseleers, T., Sundstrom, L. & Billen, J. Deleterious Wolbachia in the ant Formica
truncorum. *Proceedings of the Royal Society B: Biological Sciences* **269,** 623–629
(2002).

29. Pontieri, L., Schmidt, A. M., Singh, R., Pedersen, J. S. & Linksvayer, T. A. Artificial
selection on ant female caste ratio uncovers a link between female-biased sex ratios

and infection by Wolbachia endosymbionts. *J Evolution Biol* **30,** 225–234 (2017).

30.   Shoemaker, D. D., Ross, K. G., Keller, L., Vargo, E. L. & Werren, J. H. Wolbachia
infections in native and introduced populations of fire ants (Solenopsis spp.). *Insect*
Molecular Biology **9,** 661–673 (2000).

31.   Correa, C. C. & Ballard, J. W. O. Wolbachia Associations with Insects: Winning or
Losing Against a Master Manipulator. *Front. Ecol. Evol.* **3,** 506 (2016).

32.   Trivers, R. L. & Hare, H. Haploidploidy and the evolution of the social insect. *Science*
191, 249–263 (1976).

33.   Keller, L. *et al.* Sex ratio and Wolbachia infection in the ant Formica exsecta.
Heredity **87,** 227–233 (2001).

34.   Singh, R. & Linksvayer, T. A. The bacterial endosymbiont Wolbachia increases
reproductive investment and accelerates the life cycle of ant colonies. **90,** 89

35.   Treanor, D., Pamminger, T. & Hughes, W. O. H. The evolution of caste-biasing
symbionts in the social hymenoptera. *Insect. Soc.* **65,** 513–519 (2018).

36.   Schrempf, A., Moser, A., Delabie, J. & Heinze, J. Sperm traits differ between winged
and wingless males of the ant *Cardiocondyla obscurior. Integrative Zoology* **11,**
427–432 (2016).

37.   Andersen, S. B., Boyle, M., Nash, D. R. & Boomsma, J. J. Dynamic Wolbachia
prevalence in Acromyrmex leaf-cutting ants: potential for a nutritional symbiosis. *J*
Evolution Biol **25,** 1340–1350 (2012).

38.   Kramer, B. H., Schrempf, A., Scheuerlein, A. & Heinze, J. Ant Colonies Do Not Trade-
Off Reproduction against Maintenance. *PLoS ONE* **10,** e0137969 (2015).

39.   Cremer, S. & Heinze, J. Adaptive production of fighter males: queens of the ant
Cardiocondyla adjust the sex ratio under local mate competition. *Proc. Biol. Sci.*
269, 417–422 (2002).

40.   de Menten, L., Cremer, S., Heinze, J. & Aron, S. Primary sex ratio adjustment by ant
queens in response to local mate competition. *Animal Behaviour* **69,** 1031–1035
(2005).

41.   Suefuji, M., Cremer, S., Oettler, J. & Heinze, J. Queen number influences the timing
of the sexual production in colonies of *Cardiocondyla* ants. *Biol. Lett.* **4,** 670–673
(2008).

42.   Gotzek, D., Axen, H. J., Suarez, A. V., Helms Cahan, S. & Shoemaker, D. Global
invasion history of the tropical fire ant: a stowaway on the first global trade routes.
Molecular Ecology **24,** 374–388 (2015).

43.   Foucaud, J. *et al.* Worldwide invasion by the little fire ant: routes of introduction and
eco-evolutionary pathways. *Evol Appl* **3,** 363–374 (2010).

44.   Braig, H. R., Zhou, W., Dobson, S. L. & O'Neill, S. L. Cloning and characterization of a
gene encoding the major surface protein of the bacterial endosymbiont Wolbachia
pipientis. *Journal of Bacteriology* **180,** 2373–2378 (1998).

45.   Clark, M. E., Heath, B. D., Anderson, C. L. & Karr, T. L. Induced paternal effects
mimic cytoplasmic incompatibility in Drosophila. *Genetics* **173,** 727–734 (2006).

46.   Bonneau, M. *et al.* Culex pipiens crossing type diversity is governed by an amplified
and polymorphic operon of Wolbachia. *Nature Communications* **9,** 319–10 (2018).

47.   Brucker, R. M. & Bordenstein, S. R. The Hologenomic Basis of Speciation: Gut
Bacteria Cause Hybrid Lethality in the Genus Nasonia. *Science* (2013).
doi:10.1126/science.1240659

48.   Sinha, A., Li, Z., Sun, L. & Carlow, C. K. S. Complete Genome Sequence of the

797   Wolbachia wAlbB Endosymbiont of Aedes albopictus. *Genome Biology and*
798   *Evolution* **11,** 706-720 (2019).

799 49. Chatzispyrou, I. A., Held, N. M., Mouchiroud, L., Auwerx, J. & Houtkooper, R. H.
800   Tetracycline antibiotics impair mitochondrial function and its experimental use
801   confounds research. *Cancer Research* **75,** 4446-4449 (2015).

802 50. Kautz, S., Rubin, B. E. R. & Moreau, C. S. Bacterial Infections across the Ants:
803   Frequency and Prevalence of Wolbachia, Spiroplasma, and Asaia. *Psyche: A Journal*
804   *of Entomology* **2013,** 1-11 (2013).

805 51. Russell, J. A. *et al.* Specialization and geographic isolation among Wolbachia
806   symbionts from ants and lycaenid butterflies. *Evolution* **63,** 624-640 (2009).

807 52. Tolley, S. J. A., Nonacs, P. & Sapountzis, P. Wolbachia Horizontal Transmission
808   Events in Ants: What Do We Know and What Can We Learn? *Front Microbiol* **10,**
809   296 (2019).

810 53. Kelly, M., Price, S. L., de Oliveira Ramalho, M. & Moreau, C. S. Diversity of Wolbachia
811   Associated with the Giant Turtle Ant, Cephalotes atratus. *Curr Microbiol* **76,** 1330-
812   1337 (2019).

813 54. Arnold, P. A., Levin, S. C., Stevanovic, A. L. & Johnson, K. N. Drosophila
814   melanogaster infected with Wolbachia strain wMelCS prefer cooler temperatures.
815   *Ecological Entomology* **44,** 287-290 (2018).

816 55. Truitt, A. M., Kapun, M., Kaur, R. & Miller, W. J. Wolbachia modifies thermal
817   preference in Drosophila melanogaster. *Environ Microbiol* **21,** 3259-3268 (2018).

818 56. Coyne, J. A. & Orr, H. A. 'Patterns of speciation in *Drosophila*' revisited. *Evolution*
819   **51,** 295-303 (1997).

820 57. Schwander, T., Suni, S. S., Cahan, S. H. & Keller, L. Mechanisms of reproductive
821   isolation between an ant species of hybrid origin and one of its parents. *Evolution*
822   **62,** 1635-1643 (2008).

823 58. Heinze, J. & Hölldobler, B. Fighting for a harem of queens: physiology of
824   reproduction in Cardiocondyla male ants. *Proc Natl Acad Sci U S A* **90,** 8412-8414
825   (1993).

826 59. Schrempf, A., Darrouzet, E. & Heinze, J. Mating success and potential male-worker
827   conflict in a male-dimorphic ant. *BMC Evol Biol* **7,** 114 (2007).

828 60. Schmidt, C. V., Frohschammer, S., Schrempf, A. & Heinze, J. Virgin ant queens mate
829   with their own sons to avoid failure at colony foundation. *Naturwissenschaften* **101,**
830   69-72 (2013).

831 61. Heinze, J., Cremer, S., Eckl, N. & Schrempf, A. Stealthy invaders: the biology of
832   Cardiocondyla tramp ants. *Insect. Soc.* **53,** 1-7 (2006).

833 62. Seifert, B., Ritz, M. & Csősz, S. Application of exploratory data analyses opens a new
834   perspective in morphology-based alpha-taxonomy of eusocial organisms.
835   *Myrmecological News* **19,** 1-15 (2013).

836 63. Seifert, B., Okita, I. & Heinze, J. A taxonomic revision of the *Cardiocondyla nuda*
837   group (Hymenoptera: Formicidae). *Zootaxa* **4290,** 324-33 (2017).

838 64. Campbell, E. A. *et al.* Structural mechanism for rifampicin inhibition of bacterial rna
839   polymerase. *Cell* **104,** 901-912 (2001).

840 65. Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative CT
841   method. *Nat Protoc* **3,** 1101-1108 (2008).

842 66. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications
843   to single-cell sequencing. *J. Comput. Biol.* **19,** 455-477 (2012).

844   67.   Wick, R. R., Schultz, M. B., Zobel, J. & Holt, K. E. Bandage: interactive visualization of
845   de novo genome assemblies. *Bioinformatics* **31,** 3350–3352 (2015).

846   68.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Meth*
847   **9,** 357–359 (2012).

848   69.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30,**
849   2068–2069 (2014).

850   70.   Ronquist, F. *et al.* MrBayes 3.2: efficient Bayesian phylogenetic inference and model
851   choice across a large model space. *Systematic Biology* **61,** 539–542 (2012).

852   71.   Li, L. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes.
853   *Genome Research* **13,** 2178–2189 (2003).

854   72.   Chen, F., Mackey, A. J., Vermunt, J. K. & Roos, D. S. Assessing performance of
855   orthology detection strategies applied to eukaryotic genomes. *PLoS ONE* **2,** e383
856   (2007).

857   73.   Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome*
858   *Biol.* **5,** R12 (2004).

859   74.   Heddi, A., Grenier, A.-M., Khatchadourian, C., Charles, H. & Nardon, P. Four
860   intracellular genomes direct weevil biology: Nuclear, mitochondrial, principal
861   endosymbiont, and Wolbachia. *Proceedings of the National Academy of Sciences*
862   **96,** 6814–6819 (1999).

863   75.   Sanguin, H. *et al.* Development and validation of a prototype 16S rRNA-based
864   taxonomic microarray for Alphaproteobacteria. *Environ Microbiol* **8,** 289–307
865   (2006).

Figure 1: Mean weekly egg numbers produced by intra- and inter-population crosses of the ant *C. obscurior*

Mean weekly egg numbers produced by Brazilian (BR) and Japanese (JP) queens mated to males from their own or from a different population over a period of six weeks. JP ants were collected from colonies that either carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae (JP) or did not (JP$_{we-}$). Males used in "CI rescue" crosses were collected from JP colonies with and without *Cand.* Westeberhardia cardiocondylae, which had been treated with the antibiotic rifampicin (JP^{rif+}; JP$_{we-}^{rif+}$). Numbers below box plots indicate the number of replicates. Differences between groups were tested with pairwise Mann-Whitney-U-tests followed by Bonferroni-Holms correction of p-values. Letters above boxplots indicate statistically significant differences at p<0.05. For Bonferroni-Holms corrected pairwise p-values see Table S1.

Figure 2: Total pupae numbers and pupae sex ratios produced by intra- and inter-population crosses of the ant *C. obscurior*

Total number of pupae (A) and sex ratio of pupae (B) produced in crosses between Brazilian (BR) and Japanese (JP) queens and males over a period of 12 weeks. JP ants were collected from colonies that either carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae (JP) or did not (JP$_{we-}$). Males used in "CI rescue" crosses were collected from JP colonies with and without *Cand.* Westeberhardia cardiocondylae, which had been treated with the antibiotic rifampicin (JP^{rif+}; JP$_{we-}^{rif+}$). Numbers below box plots indicate the number of replicates. Total pupae numbers are plotted as log +1 for better visualization and differences between groups were tested with pairwise Mann-Whitney-U-tests followed by Bonferroni-Holms correction of p-values. For Bonferroni-Holms corrected pairwise p-values see Table S3. Sex ratios produced by different crosses were compared with a generalized linear model with logit link followed by manual Bonferroni-Holms correction of p-values. For Bonferroni-Holms corrected pairwise p-values see Table S4. Letters above boxplots indicate statistically significant differences at p<0.05.

Figure 3: Effects of rifampicin treatment on *Wolbachia* titres in males
All males were collected from colonies that carried the main endosymbiont *Candidatus* Westeberhardia cardiocondylae. Numbers below box plots indicate the number of replicates. Differences in *Wolbachia* titres between control and rifampicin-treated males were tested with a Mann-Whitney-U-Test. Stars indicate statistically significant differences at p<0.001 (***).

Figure 4: *Wolbachia* spp. phylogeny and genome synteny
(A) Bayesian phylogenetic placement of *Wolbachia* strains sequenced from *C. obscurior* (*w*Cobs-BR and *w*Cobs-JP). *Anaplasma phagocytophilum* and *Ehrlichia canis* were used as outgroups. For *Wolbachia*, names at tips correspond to the host from which they were sequenced/isolated. Red dots are used to highlight the newly sequenced strains. Next to the binomial names, strain names are shown in grey and silhouettes for the host are sketched in black. Horizontal bars delimit *Wolbachia* supergroups (A-F). *=1.0 posterior probability. (B) Genome-wise nucleotide-based synteny plot between *w*Cobs-BR and *w*Cobs-JP. Red is used to indicate a direct match and blue a reverse complement match.

Figure 5. Phylogeny and domain structure of *cif* genes

Bayesian phylogenetic trees for (A) *cifA* and (B) *cifB* genes. Colored ovals are used to highlight the different gene "types". *= 1.0 posterior probability. (C) Domain structure for the *cif* genes of *w*Cobs-BR (top) and *w*Cobs-JP (bottom) strains. For the latter, the two loci are shown and indicated with a vertical bar to the right of the sketches. Color codes and designation of the domains are shown in the legend.

Figure 6: Localization of endosymbiotic bacteria in male adult abdomens of *C. obscurior* via fluorescence in situ hybridization.

Wolbachia were specifically stained with two Cy5-labeled probes (magenta), and *Cand.* Westeberhardia with a Cy3-labeled probe (green). Host cell nuclei are counterstained with DAPI (blue). (A) Transversal section of *C. obscurior* JP male abdomen showing the testes (t), the *Cand.* Westeberhardia-containing bacteriomes (B) and the gut (g). The areas indicated by rectangles are displayed in higher magnification showing *Wolbachia*-infected seminiferous tubules (B) and bacteriomes co-infected with *Cand.* Westeberhardia and *Wolbachia* (C). (D) *C. obscurior* BR male testes and gut epithelium infected with *Wolbachia*. Scale bars: 50 µm (A) and 20 µm (B-D).

Table 1: Wolbachia genome assembly statistics
Genome assembly and draft annotation statistics for the newly sequenced *Wolbachia* strains from *C. obscurior* (*w*Cobs-BR and *w*Cobs-JP).

	Wolbachia	
	wCobs-BR	**wCobs-JP**
Assembled genome size (Mbp)	1.19	1.30
Number of scaffolds	148	182
N50	14,698	16,173
G+C content (%)	35.3	35.1
Predicted CDSs*	1,013	1,058
tRNAs	34	34
rRNAs	3	3
Other ncRNAs genes/motifs	71	54