1   **Human iPSC-based neurodevelopmental models of globoid cell leukodystrophy uncover**

2   **patient- and cell type-specific disease phenotypes**

4   Elisabeth **Mangiameli**[1], Anna **Cecchele**[1], Francesco **Morena**[2], Francesca **Sanvito**[3], Vittoria **Matafora**[4],

5   Lucrezia **Della Volpe**[1], Daniela **Gnani**[1]**,** Marianna **Paulis**[5,6], Lucia **Susani**[5,6], Sabata **Martino**[2], Raffaella **Di**

6   **Micco**[1], Angela **Bachi**[4], and Angela **Gritti**[1]

8   1. San Raffaele Telethon Institute for Gene Therapy (SR-Tiget), IRCCS San Raffaele Scientific

9   Institute, Via Olgettina 60, 20132 Milan, Italy

10   2. Department of Chemistry, Biology and Biotechnology, University of Perugia, Via del Giochetto,

11   06123 Perugia, Italy

3. Pathology Unit, IRCCS San Raffaele Scientific Institute, Via Olgettina 60, 20132 Milan, Italy

4. IFOM-FIRC Institute of Molecular Oncology, Via Adamello 16, 20139 Milan, Italy

5. Humanitas Clinical and Research Center-IRCCS, Rozzano, Milan, Italy.

6. National Research Council (CNR)-IRGB/UOS of Milan, Milan, Italy.

Correspondence:

Angela Gritti, San Raffaele Telethon Institute for Gene Therapy (SR-Tiget), IRCCS San Raffaele Scientific

Institute, Via Olgettina 60, 20132 Milan, Italy; e-mail: gritti.angela@hsr.it

Summary

Globoid Cell Leukodystrophy (GLD, or Krabbe disease) is a rare lysosomal storage disease caused

by inherited deficiency of β-galactocerebrosidase (GALC). The build-up of psychosine and other

undegraded galactosylsphingolipids in the nervous system causes serious demyelination and

neurodegeneration. The molecular mechanisms of GLD are poorly elucidated in neural cells and whether

murine systems faithfully recapitulate critical aspects of the human disease is still to be defined.

Here, we established a collection of GLD patient-specific induced pluripotent stem cell (iPSC)

lines. We differentiated iPSCs from two patients – bearing different disease-causing mutations - into

neural progenitors cells (NPCs) and their neuronal/glial progeny, assessing the impact of GALC deficiency

and lentiviral vector-mediated GALC rescue/overexpression by means of phenotypic, biochemical,

molecular, and lipidomic analysis. We show a progressive increase of psychosine during the

differentiation of GLD NPCs to neurons and glia. We report an early and persistent impairment of glial

and neuronal differentiation in GLD cultures, with peculiar differences observed in the two GLD lines.

GLD cells display a global unbalance of lipid composition during the iPSC to neural differentiation and

early activation of cellular senescence, depending on the disease-causing mutation. Restoration of GALC

activity normalizes the primary pathological hallmarks and partially rescues the differentiation program

of GLD NPCs.

Our results show that multiple mechanisms besides psychosine toxicity concur to CNS pathology

in GLD and highlight the need of a timely regulated GALC expression for proper lineage commitment and

differentiation of human NPCs. These findings have important implications for establishing tailored

cell/gene therapy strategies to enhance disease correction in GLD.

Keywords

Pluripotent stem cells, leukodystrophy, Krabbe disease, gene therapy, neural progenitors,

oligodendrocytes, senescence.

Running title

Human iPSC-based models of GLD

Introduction

Globoid Cell Leukodystrophy (GLD, or Krabbe disease) is a rare lysosomal storage disorder (LSD)

caused by inherited deficiency of β-galactocerebrosidase (GALC), a key enzyme in the catabolism of

galactosphingolipids, e.g. galactosylceramide and galactosylsphingosine (psychosine) (Suzuki and Suzuki,

1970). The infantile forms are characterized by relentless demyelination and neurodegeneration of the

central and peripheral nervous system (CNS, PNS). Allogeneic hematopoietic stem/progenitor cell (HSPC)

transplant - the only therapeutic option for GLD patients - provides a functional GALC enzyme to the

affected organs through the engraftment of HSPC-derived myeloid progeny. However, it fails to fully

correct the severe CNS pathology, even if performed at the asymptomatic stage (Allewelt et al., 2016),

likely because of the insufficient GALC supply provided by HSPC-derived microglia in the brain. *In vivo*

and *ex vivo* gene therapy (GT) approaches using lentiviral vectors (LV) and adeno-associated vectors

(AAV) benefit GLD animal models (Bradbury et al., 2018; Gentner et al., 2010; Lattanzi et al., 2010, 2014;

Lin et al., 2005; Meneghini et al., 2016; Rafi et al., 2015; Ricca et al., 2015; Ungari et al., 2015). The

advantage of *ex vivo* GT with LV-modified HSPCs in infants affected by metachromatic leukodystrophy

(MLD) – a similar LSD (Biffi et al., 2013; Sessa et al., 2016), suggests that a similar approach might be

translated to GLD. Still, the obstacles in reaching safe, widespread and stable expression of therapeutic

GALC levels in the human CNS still limits the clinical development of these GT strategies to GLD.

Studies in GLD animal models suggest that a modest accumulation of psychosine in brain tissues

impairs CNS and PNS development from neural progenitor cells (NPCs) (Castelvetri et al., 2011;

Santambrogio et al., 2012; Teixeira et al., 2014). Psychosine storage is present in nervous tissues before

the onset of symptoms, even if high levels are required to trigger overt cytotoxicity, likely with cell- and

tissue-specific sensitivity. Besides the effects exerted on myelin and axons, psychosine – and other

substrates in the GALC-related sphingolipid pathway (Hannun and Obeid, 2017) - directly or indirectly

affect glial and neuronal cell homeostasis interfering with lipid content and distribution (Giri et al., 2006;

White et al., 2009), altering proteostasis network (Pellegrini et al., 2019), promoting apoptosis (Jatana et

al., 2002), lysosomal dysfunction (Folts et al., 2016; Samie and Xu, 2014), and autophagy (Del Grosso et

al., 2019). Growing evidence points to a previous underestimated association between lysosomal

dysfunction, protein misfolding/aggregation, lipid unbalance, and senescence in multiple organisms and

cell types (Buratta et al., 2017; Irahara-Miyana et al., 2018; Lee et al., 2014; Lizardo et al., 2017; Smith et

al., 2014; Trayssac et al., 2018). Also, recent observations implicate NPCs and oligodendroglial progenitor

cells (OPCs) senescence in the pathogenesis of neurodegenerative disorders with protein aggregation

(Zhang et al., 2019) and neuroinflammatory demyelinating disorders (Nicaise et al., 2019), providing a

strong rationale for investigating the potential implication of these processes in GLD. Altogether, these

complex and still poorly elucidated events may contribute to the multifaceted aspects of GLD pathology

and explain the modest efficacy of conventional or experimental therapies in correcting CNS and PNS

pathology in GLD animal models and patients (Allewelt et al., 2016; Hawkins-Salsbury et al., 2015; Rafi et

al., 2015; Ricca et al., 2015).

More than 100 disease-causing mutations in the human *GALC* gene have been described

(Wenger and Luzi, 2014), among which a 30-kb deletion accounts for 40% of pathogenic alleles in the

Caucasian population (Rafi et al., 1995). The occurrence of this large gene deletion in homozygosis is

associated to the early-onset and most severe disease forms. The genotype-phenotype correlation is less

clear for the many missense/nonsense mutations occurring in homozygosis or compound heterozygosis,

which may be associated to variable residual GALC activity and severe to mild disease variants (Duffner

et al., 2011, 2012; De Gasperi et al., 1996, 1999; Jalal et al., 2012; Tappino et al., 2010). Misfolded GALC

proteins evoke different Unfolded Protein Response (UPR) activation (Irahara-Miyana et al., 2018) and

show mutation-specific alteration of trafficking to lysosomes and processing (Shin et al., 2016),

suggesting the occurrence of pathological cascades that are independent from or act in synergy with

primary lipid storage in determining the complex GLD phenotype. Thus, studying the mechanisms linking

the primary mutation-specific event(s) to progressive cell dysfunction and death is relevant and requires

appropriate experimental models.

Murine, canine and non-human primate models of GLD (Wenger, 2000) are valuable to test the

safety and efficacy of novel therapies; nevertheless, they fail to fully recapitulate the spectrum of

pathological manifestations observed in patients. Also, the most used mouse model (Suzuki, 1983) and

non-human primate model of GLD (Borda et al., 2008; Luzi et al., 1997) carry spontaneous mutations

that are not found in humans. Currently used GLD human cellular models include patient-specific

fibroblasts (Berardi et al., 2014; Gama Sosa et al., 1996; Rafi et al., 1996; Ribbens et al., 2013; Spratley et

al., 2016), human hematopoietic cells (Martino et al., 2009), or epithelial cell lines with induced GALC

mutations (Lee et al., 2010; Shin et al., 2016) that hardly resemble the metabolic and functional features

of NPCs and their differentiated progeny. Thus, there is a strong need for human-derived experimental

models to explore GLD pathogenesis and test correction strategies.

Human induced pluripotent stem cells (iPSCs) (Yamanaka, 2007) offer a unique tool to analyze

the molecular consequences of pathogenic mutations in the context of the genetic background of

individual patients. The differentiation of iPSCs in neural cells has boosted CNS disease modelling and

therapeutic screening (Shi et al., 2017). Human iPSCs are available from a variety of LSDs (Huang et al.,

2012) but there are no reports describing GLD-derived iPSCs.

115   Here, we established GLD patient-specific iPSC lines as a reliable human model for elucidating

GLD pathogenesis and testing the biological efficacy of gene therapy strategies in relevant neural cell

types. To this end, we differentiated iPSCs into NPCs and progeny (oligodendrocytes, neurons, and

astrocytes), and monitored the appearance/progression of cell type- and patient-specific primary and

secondary defects in GLD cells. Finally, we assessed the impact of GALC reconstitution/overexpression

(achieved by lentiviral-mediated gene transfer) in reverting the pathological phenotype and its potential

impact on the biology of human NPCs and progeny.

Results

GLD iPSCs show undetectable GALC activity and psychosine storage but display normal pluripotency

features

127   We generated iPSCs from fibroblasts of 5 GLD patients with distinct biallelic mutations in the

GALC gene (homozygous or compound heterozygous) resulting in the infantile form of the disease.

Control iPSC lines were generated from fibroblasts of two unrelated normal donors (ND1, adult; ND2,

neonatal) and one non-affected relative (ND3, parent of GLD1)(**Table S1**). GALC activity in ND fibroblasts

was 2.42 nmol*h/mg (ND1), 3.89 nmol*h/mg (ND2), and 1.56 nmol*h/mg (ND3), while it was below the

limit of detection in all GLD fibroblasts, indicating that the mutated GALC proteins have no residual

functionality. By using an integration-free, feeder-free microfluidic reprogramming system (Luni et al.,

2016) we obtained several iPSC clones from each patient and ND fibroblast line (reprogramming

efficiency: 2-53%)(**Figure S1A**). After 6-10 subculturing passages, we randomly selected two clones/line

for further analysis (**Figure S1A**). The presence of specific GALC mutations was verified by PCR analysis

and sequencing (**Figure S1B-C**).

138   We assessed the pluripotency of GLD and ND iPSC clones by performing standard assays,

139   including RT-PCR (**Figure S1D**) and immunocytochemistry (**Figure 1A**) for the markers NANOG, SOX2,

140   KLF4, OCT4, and TRA1-60. Analysis of endodermal-, mesodermal-, and ectodermal-specific markers in

141   embryoid bodies (**Figure 1B; Figure S1E**) and histological assessment of teratomas (**Figure S1F**)

142   confirmed the ability of GLD and ND iPSCs to differentiate in multiple cell types and tissues both *in vitro*

143   and *in vivo*.

144   All the GLD iPSC clones displayed undetectable GALC activity (**Figure 1C**) and significant

psychosine storage (50-100-fold the ND levels), which was not observed in the parental fibroblasts

(**Figure 1D).**

Overall, GLD patient iPSCs showed the primary biochemical GLD hallmarks but expressed normal

stem cell properties, justifying their use as human-based GLD model.

LV-mediated gene transfer in GLD iPSCs rescues GALC activity and normalizes psychosine storage

We focused on the mutant lines GLD1 and GLD5 (two clones/line) for further characterization

and gene transfer studies. GLD1 is homozygous for the large 30 Kb *GALC* gene deletion

(c.1161+6532_polyA+9kbdel) responsible for the production of a truncated protein lacking the whole

coding region for the 30kDa subunit and 15% of the coding region for the 50kDa subunit (Luzi et al.,

1995; Tappino et al., 2010). GLD5 is homozygous for a missense mutation (c.1657G>A) that is predicted

to cause GALC protein misfolding (De Gasperi et al., 1996; Tappino et al., 2010). ND1 and ND2 iPSCs (two

clones/lines) were used as controls for all the subsequent experiments (**Table S1 and Figure S1A**).

We selected ND and GLD iPSC clones with normal karyotype (**Figure S2**) and used these cells

within subculturing passages 8-30. To restore GALC activity, we transduced GLD1 and GLD5 iPSC clones

using a laboratory grade/scale VSV-pseudotyped third-generation lentiviral vector (LV) encoding for the

human (h) *GALC* gene tagged with the Myc peptide under the control of the human phosphoglycerate

kinase (PGK) promoter (LV.hGALC) (Meneghini et al., 2016)(**Figure 2A**). Pilot dose-response experiments

showed the proficient transduction of iPSC clones at Multiplicity of Infection (MOI)=2 (**Figure S3A**), which

was selected to transduce GLD1.1, GLD1.3, GLD5.2, GLD5.6 and ND2.2 iPSC clones in all the following

experiments. LV.hGALC-transduced GLD (GLD^{GALC}) and ND (ND^{GALC}) iPSCs showed variable vector copy

number (VCN) (**Figure 2B**), upregulation of GALC mRNA (**Figure 2C**), efficient transgene expression

(**Figure 2D-E),** and increase in enzymatic activity (reaching supraphysiological levels; **Figure 2F**). We

observed a clear correlation between VCN, GALC mRNA expression and enzymatic activity (**Figure S2B**).

The transgenic GALC enzyme was functional, as shown by the normalization of psychosine levels

in GLD iPSCs (**Figure 2G**), which do not show other pathological hallmarks related to GALC deficiency (i.e.

GalCer accumulation, increased LAMP1 expression – index of lysosomal expansion) (**Figure S3C-D).**

Importantly, LV transduction and GALC overexpression did not affect the pluripotency (**Figure S3E**) and

proliferation rate (**Figure S3F**) of ND and GLD iPSC clones.

GLD and ND iPSC-derived NPCs are generated with similar efficiency and display comparable

phenotype

We next assessed the impact of GALC deficiency and LV-mediated GALC restoration in iPSC-

derived neural cells. We differentiated GLD, GLD^GALC, ND, and ND^GALC iPSCs into NPCs using a published

protocol (Chambers et al., 2009)(**Figure 3A**). At the end of the differentiation, iPSC-NPCs showed

downregulation of pluripotency markers (OCT4, NANOG) and upregulation of NPC markers (NESTIN,

PAX6, FOXG1), early neuronal (Doublecortin, DCX) and glial cell (ASCL1) markers **(Figure 3B-C).** We

obtained NPCs with similar efficiency from untransduced (UT) and LV.hGALC-transduced GLD and ND

iPSCs in multiple rounds of differentiation experiments with the only exception of clone GLD1.3^GALC,

which consistently displayed abnormal morphology (not shown) and modest PAX6 upregulation as

compared to other clones (**Figure 3B**). Based on these observations suggesting incomplete/defective

neural induction, we excluded clone GLD1.3^GALC from further analysis. All the selected NPC lines stably

expanded in culture for 5-6 passages in the presence of neural medium (N2) supplemented with EGF,

bFGF and Rock inhibitor (**Figure 3D).** Thus, ND and GLD iPSC-derived NPCs display bona-fide NPC

molecular and phenotypic features.

Time- and cell type-dependent transcriptional regulation of GALC expression and activity during iPSC-

NPC differentiation to neurons and glia

Myelinating cells and neurons are highly susceptible to psychosine-mediated toxicity, but

astrocytes are also affected (Misslin et al., 2017; Potter and Petryniak, 2016). Therefore, we envisaged

that differentiation of iPSC-NPCs into mixed neuronal/glial cultures would better model the composite

GLD pathology. We adapted a published protocol (Frati et al., 2018) to generate cell populations

containing neurons, astrocytes, and oligodendrocytes from ND, GLD, GLD^GALC, and ND^GALC iPSC-NPCs.

Cultures were evaluated at day 0 (d0; NPCs), d7, d14, and d24 of differentiation (**Figure 3E**), to track the

time-course GALC expression/activity and the glial/neuronal cell commitment and maturation by means

of biochemical and molecular analysis.

The GALC transcript was expressed at low levels in both ND and GLD iPSC clones (GLD1 and GLD5

mutations are not predicted to impact on mRNA transcription) and was upregulated during the neural

induction and neuronal/glial differentiation (**Figure 3F**). GALC mRNA expression in GLD^GALC and ND^GALC

iPSCs was constant over time, due to the high and stable expression driven by the constitutive PGK

promoter **(Figure 3F).**

GALC enzymatic activity moderately increased in ND differentiated cultures (d14 and d24) as

compared to the iPSCs and NPCs counterpart (≈1.5 fold) **(Figure 3G).** The GALC activity in GLD^GALC cultures

progressively decreased during the iPSC to neural differentiation to reach physiological levels at d14 and

d24, whereas ND^GALC differentiated cultures maintained supraphysiological GALC activity (≈3-fold the ND

counterpart at d14 and d24). (**Figure 3H**). The stable VCN measured in ND^{GALC} and GLD^{GALC} cultures at

different time points of differentiation (**Figure 3H**) rules out the possibility that the decrease in

enzymatic activity results from the counter selection of cells harboring high VCN over time.

Overall these data show that LV-mediated gene transfer ensures (supra)physiological GALC

expression and activity in iPSC-NPCs and neuronal/glial progeny, and highlight a stringent time- and cell

type-dependent transcriptional regulation of GALC activity during NPCs maturation.

GLD iPSC-NPCs display defective neural and glial differentiation: impact of GALC

rescue/overexpression on psychosine levels and phenotype.

Commitment of NPCs to oligodendrocyte precursors (OPCs) and differentiation in mature

oligodendrocytes (OL) is driven by the timely regulated expression of oligodendroglial genes. GLD cells

displayed an altered pattern of expression of several oligodendroglial master genes (NKX2.2, OLIG2,

SOX10, PDGFRα, MBP) as compared to ND cells, with some differences between GLD1 and GLD5 (e.g.

sustained expression of PDGFRα along the differentiation and increased SOX10 expression in

differentiated GLD5 cells) **(Figure S4A).** These data suggest an altered oligodendroglial differentiation

program, which is only partially normalized in GLD^{GALC} cultures. Interestingly, a similar alteration is also

observed ND^{GALC} cultures (**Figure S4A**). Immunofluorescence and WB analysis confirmed an early and

persistent reduction of oligodendroglial cells (OLIG2+, APC+ cells; **Figure 4A-B)** and a significant

downreguation of OLIG1 protein expression (**Figure S4B**; d24) in both GLD1 and GLD5 NPC-derived

cultures as compared to the ND counterpart, highlighting a partial phenotypic rescue in GLD^{GALC} cells and

a detrimental impact of GALC overexpression in ND cultures (**Figure 4A-B** and **Figure S4B).**

GALC deficiency affected neuronal differentiation, with some differences observed in the two

GLD lines. Real-time PCR analysis revealed similar expression of the neuronal master transcription factor

NeuroD1 in ND and GLD lines. In contrast, doublecortin (DCX) expression was reduced in GLD5 as

compared to GLD1 and ND differentiated cells **(Figure S5A).** In line with these results, GLD5 cultures

showed a significant time-dependent reduction of TUJ1+ neurons (**Figure 4C-D**) and a decreased and/or

delayed TUJ1 and neurofilament (NF)-M protein expression, respectively (**Figure S5B-C).** The major

decrease in the expression of acetylated and polyglutamated tubulin and phosphorylated neurofilaments

observed in GLD5 cultures (d14 and d24) further suggested an important neuronal loss and/or defective

neuronal differentiation/maturation (**Figure S6**). In contrast, GLD1 cultures displayed a normal

proportion of neuronal cells (**Figure 4C-D)** and normal kinetic of expression of neuronal proteins (**Figure**

S5B-C). The quantitative and qualitative neuronal defects observed in GLD5 cultures were rescued in

differentiated GLD^{GALC} cultures (**Figure 4C, Fig S5B-C**), which also showed an early upregulation of NF-M

protein as compared to the ND cultures and untransduced counterparts (**Figure S5C),** suggesting an

accelerated neuronal differentiation associated to GALC overexpression.

GFAP+ astroglia is the less represented and most variable cell population in our differentiated

cultures (<10% in ND cultures at day 24). Nevertheless, RT-PCR (**Figure S7A**), IF (**Figure 4C-D**) and WB

analysis (**Figure S7B**) showed defective glial differentiation in GLD1 cultures, which virtually failed to

generate GFAP+ astrocytes. This defect was not rescued in GLD1^{GALC} cultures (**Figure 4C-D** and **Figure S7**).

In contrast, GLD5 and GLD5^{GALC} cultures showed normal percentages of GFAP+ cells and physiological

GFAP protein expression (**Figure 4C-D** and **Figure S7**). Similarly to what observed in oligodendrocytes,

GALC overexpression negatively affected the pattern of gene expression (**Figure S5, S6, S7**), the number

and morphology of ND neurons (**Figure 4C-D**, **Figure S5)** and astrocytes **(Figure 4C-D, Figure S7**),

particularly at the latest time point in culture. Finally, both GALC deficiency (GLD cultures) and forced

GALC expression (GLD^{GALC} and ND^{GALC} cultures) were associated to higher percentages of cells that

stained negative for the selected lineage-specific markers **(Figure S7C).**

Psychosine levels were low in GLD iPSC-NPCs and increased during differentiation, to reach ≈8

(GLD1) and ≈30 fold (GLD5) the normal levels at d24 (**Figure 5A**). The psychosine build up was not

associated to a significant increase of cleaved caspase 3+ cells (**Figure 5B**) or to enhanced phospholipase

A2 (PLA2) activity (**Figure 5C**), which has been associated to psychosine-induced apoptotic cell death

(Misslin et al., 2017). Psychosine levels were normalized in GLD1^{GALC} cultures and strongly reduced in

GLD5^{GALC} cultures (≈2-fold the normal), in line with the enzymatic reconstitution. We observed moderate

enlargement of lysosomal vesicles (Lysotracker assay) (**Figure 5D**) in GLD NPCs (GLD1>GLD5) and

increased LAMP1 protein expression in GLD and ND^{GALC} differentiated cultures (**Figure 5E**).

Taken together, these data show that: i) GALC deficiency severely impairs the differentiation of

hiPSC-derived NPCs into oligodendrocytes, neurons, and astrocytes, with cell type- and patient-specific

phenotypes; ii) LV-mediated GALC reconstitution in GLD cells reduces psychosine levels and provides

partial phenotypic rescue in GLD neural progeny; iii) GALC-overexpressing ND cells display lysosomal

expansion and a defective neuronal and glial phenotype, suggesting that a tight regulation of GALC

expression is required to ensure proper NPC functionality and preserve multipotency**.**

Early activation of a senescence response in GLD5 iPSC-derived neural cells

Our data suggested a marginal contribution of cell death to the loss of GLD neurons and glial

cells. Therefore, we explored whether the altered differentiation programs of GLD NPCs and progeny

could be caused by an alteration of proliferative potential. By performing and quantifying Ki67/nestin

immunostaining at the different stages of the differentiation protocol, we showed a time-dependent

decrease in the percentage of proliferating cells in all ND and GLD cultures, indicating the lineage

commitment of iPSC-NPCs, their exit of cell cycle and acquisition of a post-mitotic state (**Figure 6A).**

Interestingly, EdU incorporation assay performed at the same time points during differentiation

showed an earlier and more drastic drop in the percentage of cells in active S-phase in GLD5 cultures

as compared to ND and GLD1 counterparts (**Figure 6B).** This proliferation drop was recovered in

GLD5^{GALC} cultures (**Figure 6B**).

We next investigated whether the early proliferation arrest highlighted in GLD5 cultures was

associated with the establishment of cellular senescence, a biological process characterized by stable

growth arrest, accumulation of DNA damage and activation of a pro-inflammatory program

collectively known as SASP, senescence-associated secretory phenotype (He and Sharpless, 2017). We

detected upregulation, at the transcript and protein levels, of the cell cycle inhibitor p21 (**Figure 6C-**

D) and increased nuclear levels of DNA damage response (DDR) markers 53BP1 (**Figure 6E)** and yH2AX

(Figure S8A-B) as early as d7 post-differentiation. The induction of DDR and p21 was specific of GLD5

cultures, not reported in ND or GLD1 cultures, and was partially rescued by in GLD5^{GALC} cultures. We

also observed a dramatic increase in the percentage of Senescence Associated (SA) β-gal positive cells

in GLD5 (but not ND or GLD1 cultures) with a peak at d7 of differentiation (**Figure 6F**), induction of

the senescence-associated marker p16 (**Figure S8C**), and of the SASP pro-inflammatory factors TNF-α

and IL1-β (**Figure S8D-E**). These senescence-associated hallmarks were reduced in GLD5^{GALC} cultures.

Altogether, our data suggest that GLD5 iPSC-derived NPCs activate an early senescence

program that hampers physiological lineage commitment/differentiation. This senescent phenotype

is patient-specific (as it is not evident in GLD1 cells), and is rescued by supplying physiological levels

of a functional GALC enzyme.

GLD cells display an altered lipidomic profile during the iPSC to neural differentiation

Sphingolipids display widespread developmental, physiological and pathological actions in the

CNS, and induce biological effects on various stem/progenitor cell types (Hannun and Obeid, 2017).

Given the central role of GALC in the sphingolipid metabolism, we performed an untargeted global

lipidomic profiling in ND, GLD, and GLD^{GALC} iPSCs, NPCs and neuronal/glial cell progeny (d14 of

differentiation) to elucidate the impact of GALC absence or enhanced expression during the iPSCs to

neural differentiation. Lipid extracts were normalized based on protein concentration and analyzed by

LC-MS-based lipidomics workflow. We determined the lipid species showing reproducible changes in two

independent profiling experiments with two biological replicates for each experiment.

We identified more than 800 lipid species, with a distinct signature and enrichment of specific

lipid classes in iPSCs (e.g. triacylglycerols, TAG; lyso-phosphatidylserine, LPS), NPCs (e.g.

phosphatidylcholine, PC), and differentiated progeny (e.g phosphatidylethanolamine, PE; sphingomyelin,

SM) (**Figure 7A**). We observed profound changes in a number of phospholipid classes in GLD as

compared to ND cells (i.e. TAG; Diacylglycerols, DAG; Monoacylglycerols, MAG; PC; PE;

Lysophosphatidylethanolamine, LPE; PS; SM; Ceramides, Cer; HexosylCeramides, HexCer)(**Figure 7B**).

Within the same lipid class, we detected differences related to the stage of differentiation (iPSCs vs.

NPCs. vs differentiated cells) and genotype (ND vs. GLD1 vs. GLD5). GLD1 and ND iPSCs often showed a

comparable abundance of all these lipid classes, while GLD5 iPSCs displayed a clearly different trend

(**Figure 7B**). The most striking differences between GLD1 and ND cells were observed in NPCs, which

showed decreased levels of PE/LPE and increased Cer and DAG levels. The different lipidomic profile

displayed by GLD1 and GLD5 cells become more evident as cells undergo differentiation. Indeed, when

compared to GLD1 and ND cells, GLD5 cells showed reduced PC, PE, HexCer, and SM, and a strong

accumulation of DAG, TAG, Cer, and PS **(Figure 7B)**. The inverse pattern of Cer and SM expression in

GLD5 cell populations - and particularly in differentiated cells - suggests that excess Cer could result from

the SM-based synthetic pathway in these cells (Young et al., 2012).

LV-mediated rescue of GALC activity partially normalized the lipidomic profiles of GLD1 and GLD5,

as shown by the PCA analysis (**Figure 7C)** and lipid signature **(Figure S9A-B).** In GLD1[GALC] cells we

observed a rescue of PC, PE, HexCer (iPSCs) and TAG (NPCs). In GLD5 [GALC] cells we observed the

normalization of HexCer and SM level, as well as a rescue to normal levels of all DAG, TAG and PC that

were mostly affected in GLD5 differentiated cells. LV-mediated GALC overexpression resulted in a

profound and global impact on the lipidome of ND cells. Coherently with the presence of supranormal

GALC activity, we detected in ND[GALC] iPSCs, NPCs, and differentiated cells decreased levels of HexCer and

variable increase of Cer, but also of DAG and PS in comparison to ND counterparts (**Figure S9C).**

Discussion

Human iPSCs have been differentiated into neurons and/or glial cells to model the neural defects

associated to several lysosomal storage diseases (LSDs) including mucopolysaccharidosis (Bayó-Puxan et

al., 2018; Canals et al., 2015; Kobolák et al., 2019; Lemonnier et al., 2011; Vallejo-Diez et al., 2018),

Niemann−Pick type C (Maetzel et al., 2014; Ordoñez and Steele, 2016; Trilck et al., 2013; Yu et al., 2014),

Pompe disease (Higuchi et al., 2014), Gaucher disease (Panicker et al., 2012; Sun et al., 2015; Tiscornia et

al., 2013), GM1 and GM2 gangliosidosis (Son et al., 2015)(Allende et al., 2018), NCL (Lojewski et al., 2014),

and MLD (Frati et al., 2018; Meneghini et al., 2017). To date, this is the first report describing a human

iPSC-based neural model of GLD.

The five GLD patient-derived fibroblast cell lines that we used for somatic cell reprogramming

have different mutations in the *GALC* gene, all of them associated to the infantile form of the disease. All

the selected GLD iPSC lines displayed normal phenotypic and functional properties despite having

undetectable GALC enzymatic activity and significant psychosine storage. The dispensability of lysosomal

enzymatic activity for somatic cell reprogramming and iPSC maintenance as well as the tolerability to

lipid storage has been described in human iPSCs in other LSD models (Canals et al., 2015; Lojewski et al.,

2014; Meneghini et al., 2017; Trilck et al., 2013). The LV-mediated supply of physiological or slightly

supraphysiological GALC activity (3-4 fold the ND levels) in GLD iPSCs normalizes psychosine levels,

confirming the potential of gene transfer to correct the primary storage in GALC-deficient human cells. In

line with our previous study (Meneghini et al., 2017), we did not experience LV-mediated toxicity in

hiPSCs, which were viable and functional for several passages after transduction, maintaining

pluripotency and stable GALC activity.

The GALC enzyme contributes to preserve NPCs functionality during post-natal murine CNS

development (Santambrogio et al., 2012), but this aspect was never investigated in human NPCs,

neurons, and glial cells. GLD iPSC-derived NPCs had levels of psychosine similar to their ND counterparts,

differently from the parental GLD fibroblasts and iPSCs, in which psychosine is elevated. Interestingly,

the lack of psychosine storage distinguishes human from murine NPCs (Santambrogio et al., 2012). The

differentiation of GLD iPSC-NPCs into oligodendrocytes, neurons, and astrocytes exacerbates the

pathological phenotype, in line with the relevant role of GALC during glial and neuronal differentiation

(Graziano and Cardile, 2015). Specifically, we observed an early and persistent loss of oligodendroglial

cells in both GLD1 and GLD5 NPC-derived cultures. GLD5 cultures also displayed a dramatic loss of

neurons, while astrocytes were less affected. We observed an opposite situation in GLD1 cultures, in

which neurons were normally generated while astrocytes were virtually lost.

The oligodendroglial and neuronal cell dysfunction/loss described in GLD murine models has

been associated to psychosine-mediated toxicity (Hawkins-Salsbury et al., 2013; White et al.,

2011)(Bongarzone et al., 2016; Ricca and Gritti, 2016). Importantly, even a modest psychosine storage

may affect neuronal and glial differentiation (Castelvetri et al., 2011; Santambrogio et al., 2012; Teixeira

et al., 2014). Increased psychosine levels have been detected in the brain of GLD human foetuses (Igisu

and Suzuki, 1984) and elevated psychosine blood concentration is considered a specific marker in

infantile GLD patients (Escolar et al., 2017), who may display early white matter changes and functional

disability (Gupta et al., 2015). The human iPSC-based model described here exacerbates the defective

neuronal/glial development from NPCs reported in GLD murine models, and recapitulates the clinical

observations made on GLD patients. Also, several findings in this study suggest a composite and

mutation-dependent effect associated to GALC deficiency which impacts on NPC

commitment/differentiation besides psychosine toxicity: i) psychosine accumulation starts at early time

points of NPC differentiation and increases as cells differentiate into neurons and glia; still, this

progressive load is not accompanied by a dramatic increase of apoptotic cells; ii) higher percentages of

marker-negative cells are constantly found in GLD differentiated cultures, together with altered

expression of key genes involved in oligodendroglial, neuronal, and astroglial specification/maintenance,

suggesting a block in cell commitment or differentiation; iii) we observe only a partial rescue of the

differentiation program in LV.hGALC-transduced GLD neurons/glia - in which psychosine storage is

cleared - with an evident amelioration in the neuronal compartment but far less improvement in the

glial population.

In the attempt of elucidating the link between GALC deficiency and defective neural

differentiation, the mechanisms of disease correction upon gene transfer, and the mechanisms

underlying the mutation-specific phenotype, we pointed our attention on the lipidome. The

complexity and the many functions of the lipidome match, if not exceed, those of the genome or

proteome (Hannun and Obeid, 2017). Intriguingly, many sphingolipids involved in the GALC pathway,

such as ceramide, sphingosine, and sphingosine-1-phosphate, contribute to proliferation, migration, cell

fate, apoptosis (Hannun and Obeid, 2008, 2017) and, as suggested recently, senescence and autophagy

(Trayssac et al., 2018; Young and Wang, 2018). We previously documented the unbalance of some

bioactive lipids in brain tissues of GLD mice (Santambrogio et al., 2012). Here, we show a distinctive

global lipid profile in iPSCs vs. NPCs. vs. NPC progeny, regardless of the genotype (ND, GLD) or

mutation (GLD1, GLD5), suggesting a robust change in lipid metabolism during iPSC to neural

differentiation in our experimental conditions. In particular, the increase of SM levels during

differentiation in parallel with a decrease of PC might be related to the use of ceramide and PC by SM

synthases to produce SM (Liu et al., 2009).

By comparing the global lipidomic profile in each cell population, we observed that GLD and

ND cells differ for the abundance of various fatty acids and phospolipid classes. The mild unbalance of

lipid composition in GLD NPCs becomes more evident as cells differentiate. Intriguingly, GLD1 and GLD5

cell populations displayed remarkable differences in their lipidomic profile. Specifically, GLD5 iPSCs

and NPC progeny show an accumulation of TAG, DAG, and ceramide, whose role in cell fate, cellular

stress, ageing and replicative senescence has been proposed in multiple tissues and cell types (Hannun

and Obeid, 2002, 2017; Lizardo et al., 2017, 2018; Venable et al., 1995). We envisage that this peculiar

lipid perturbation contributes to the early senescence response and differentiation defects observed in

GLD5 iPSC-derived neural cells. Ceramide accumulation in GLD5 lines is counterintuitive, since

ceramide is the product of GalCer degradation by GALC. We speculate that other major pathways

leading to ceramide production may be induced in GLD5 cells, i.e. activation of sphingomyelinases

and/or ceramide synthases (Claus et al., 2009; Mullen et al., 2012). The SM-ceramide pathway is

particularly intriguing, since it may induce apoptosis, differentiation, and growth arrest (Jana et al., 2009).

SM might be converted into ceramide via sphingomyelinase to compensate GALC deficiency. Moreover

the parallel accumulation of DAG and TAG might be explained by an inhibition of the Kennedy pathway

that is essential for *de novo* glycerophospholipid biosynthesis and the SM cycle in sphingolipid

metabolism (Bieberich, 2012). Our data suggest that normalization of GALC activity rescues or at least

ameliorates the global lipid composition of mutant cells.

The association between lysosomal dysfunction, lipid unbalance, and senescence in NPCs,

neurons, and glial cells and its contribution to neurodegeneration and demyelination is emerging as an

area of active scientific investigations. Specifically, by taking advantage of mouse models or post-mortem

human brain samples it was reported that senescence functionally contributes to multiple sclerosis

pathogenesis (Nicaise et al., 2019) as well as to cognitive defects in Alzheimer disease (Zhang et al.,

2019). Notably, our study highlights the importance of performing this investigation in relevant human

iPSC-based neural models, which recapitulate the patient-specific pathological hallmarks more faithfully

as compared to murine models or non-neural cells.

Our study raises a number of relevant mechanistic questions. First, the respective role and

potential interaction between psychosine load and lipidome perturbation in determining the

different phenotype in GLD5 vs. GLD1 are not fully elucidated. One option would be the detrimental

action exerted by psychosine and ceramide (and possibly other lipid classes, e.g. TAG) preferentially

on oligodendrocyte and neuronal cells, respectively (Jana et al., 2009). Second, while our results

suggest an association between ceramide/TAG/DAG accumulation and senescence, the mechanistic

link between the GLD5 mutation and the occurrence of this specific lipid unbalance is not clear yet.

Possibilities include: i) a direct effect of the mutated GALC protein on the lipidome (the investigation

of the lipid-protein interactions and definition of the molecular mechanisms of pathway and enzyme

regulation are rapidly expanding in the field of bioactive lipids) (Hannun and Obeid, 2017); ii) an

indirect effect of another GLD-associated pathology (i.e. oxidative stress, mitochondrial dysfunction)

(Ricca and Gritti, 2016)(Ballabio and Gieselmann, 2009). It remains to be established which cell type is

mainly affected by these pathological hallmarks. Indeed, cellular stress and senescence programs

affect differently neurons, astrocytes and oligodendrocytes (Crowe et al., 2016).

In summary, our data show that GALC deficiency severely affects the differentiation of hiPSC-

derived neural progenitors into neurons, oligodendrocytes, and astrocytes. We highlight cell type-, and

patient-specific disease hallmarks in GLD cells, including galactosylsphingolipid storage, lipidome

perturbation, and activation of a senescence program. Finally, we show that a timely regulated GALC

expression is critical for proper neural commitment and differentiation. We anticipate these findings to

be starting point for future functional experiments aimed at further dissecting the complex and

mutation-specific GLD pathology. This knowledge will pave the way to tailored gene/cell therapy

strategies to enhance the correction of CNS pathology in GLD.

Acknowledgments

We are grateful to Lucia Sergi Sergi for LV preparation and titration, Andrea Ditadi for providing the ESC

line H9, Wim Kulik for psychosine analyses, Elizabeth Deane for providing the anti-hGALC antibody, all

the members of the Gritti's lab for generous support and helpful discussion.

Part of this work was carried out in ALEMBIC, an advanced microscopy laboratory established by IRCCS

Ospedale San Raffaele and Università Vita-Salute San Raffaele.

We thank the "Diagnosi PrePostnatale Malattie Metaboliche" Laboratory (G.Gaslini Institute) for

providing us with specimens from the "Cell line and DNA bank from patients affected by Genetic

diseases" Biobank - Telethon Genetic Biobank Network (project no. GTB07001A).

This study was funded by grants from Fondazione Telethon (Tiget Core Grant 2016-2021, grant D2) and

European Leukodystrophy Association (grant ELA 2016-010I3) to A.G.

R.D.M was supported by Fondazione Telethon (Tiget Core Grant 2016-2021, grant E5), a Career

Development Award from Human Frontier Science Program (HFSP), an Advanced Research Grant from

the European Hematology Association (EHA), "Pilot and Seed Grant 2015" from the San Raffaele

Hospital, a Hollis Brownstein Research Grant from Leukemia Research Foundation (LRF), the Interstellar

Initiative on Healthy Longevity from NYAS and AMED and by AIRC (MFAG 2019 - ID. 23321 project).

D.G was supported by the IBSA Foundation postdoctoral fellowship.

E.M., A.C., and L.d.V. conducted this study as partial fulfillment of their international Ph.D. in Molecular

Medicine, Vita-Salute San Raffaele University.

Author contribution

Conceptualization, A.G. and E.M.

Methodology, E.M, A.B, R.dM, S.M

Investigation, E.M., A.C, F.M., D.G., L.d.V., V.M., F.S., M.P, L.S.

Writing –Review & Editing, A.G., E.M, R.dM., A.B

Funding Acquisition, A.G.

Resources, S.M., A.B., R.dM., F.S.

Supervision, A.G., R.dM., A.B.

Declaration of interests

The authors declare no competing interests.

Figure titles and legends

Figure 1. GLD iPSCs have undetectable GALC activity and increased psychosine levels (see also Figures

S1 and S2)

A) Representative confocal immunofluorescence pictures of ND and GLD iPSCs expressing OCT4 (green),

NANOG (red), and TRA 1-60 (green). Nuclei are counterstained with Hoechst (blue). Single channels

(OCT4, NANOG, nuclei) and the merged picture (TRA 1-60/nuclei) are shown. Scale bars: 100µm.

B) Representative immunofluorescence pictures showing cells expressing endodermal (AFP),

mesodermal (α-SMA), and ectodermal (TUJ1) markers in ND and GLD embryoid bodies (EB). Lineage

markers in green, nuclei counterstained with Hoechst (blue). Scale bars: 80 µm (AFP) and 200 µm (TUJ1

and α-SMA).

C) GALC enzymatic activity in ND and GLD iPSC clones. Data are expressed as the mean ± SEM and

analyzed using Mann-Whitney test; ***$p < 0.001$.

D) Psychosine levels measured by mass spectrometry in ND1 (white circles), ND2 (grey circles), and GLD

(colored circles) fibroblast cell lines (n=1 sample/line) and the corresponding iPSC clones (n=2 clones/line

in single or duplicate). Data are expressed as the mean ± SEM and analyzed by Mann-Whitney test

(Fibroblasts) and Kruskal Wallis test followed by Dunns' multiple comparison test (iPSCs); ns, not

significant * $p < 0.05$, ***$p < 0.001$ vs ND (all clones). The mean ± SEM of all ND and GLD clones are plotted

in the box and whiskers graph and analyzed by Mann-Whitney test; *$p < 0.05$, ***$p < 0.001$.

Figure 2. LV-mediated gene transfer rescues GALC expression and enzymatic activity in GLD iPSCs (see

also Figure S3)

A) Schematic representation of the LV.hGALC vector. hPGK, human phosphoglycerate kinase promoter;

wpre, Woodchuck hepatitis virus Post-transcriptional Regulatory Element.

B) Vector Copy Number measured in ND and GLD iPSCs clones transduced with LV.hGALC at multiplicity

of infection= 2 (MOI2). Data are expressed as the mean ± SEM (n=2-4 replicates/clone) and analyzed

using Kruskal Wallis test followed by Dunnett's multiple comparison post-test; *$p < 0.05$.

C) GALC mRNA expression in untreated (UT; empty bars) and LV.hGALC-transduced (striped bars). n=3

replicates/clone. Values are normalized on GAPDH and expressed as fold change ($2\text{^}-\Delta\Delta CT$) vs ND

(mean). Data are expressed as mean ± SEM and analysed using one way Anova followed by Turkey

multiple comparison post-test; **** $p < 0.0001$.

D) Representative Western Blot showing the expression of the GALC-myc fusion protein (anti-myc

antibody; expected band size 80 kDa, indicative of the precursor protein) in LV.hGALC transduced (LV;

MOI2) ND and GLD iPSCs clones. β-actin was used as loading control. The 80kDa band is not detectable in

untransduced clones.

E) Representative immunofluorescence pictures showing the expression of the GALC protein (green;

anti-hGALC, recognizing the full-length protein) in untransduced (UT) and LV.hGALC transduced ND and

GLD iPSCs. Nuclei are counterstained with Hoechst (blue). Scale bars= 20μm.

F-G) GALC enzymatic activity (F) and Psychosine levels (G; clones as in F) in untransduced (UT, empty

bars) and LV.hGALC transduced (striped bars) ND, GLD and iPSCs. Data are expressed as the mean ± SEM

(n=2-4 replicates/clone). Data were analyzed by Kruskal Wallis test followed by Dunnett's multiple

comparison post-test (ND selected as control group); **$p < 0.01$; ***$p < 0.001$; ns, not significant.

Figure 3. Differentiation of ND and GLD iPSCs into NPCs, neurons and glial cells

A) Schematic representation of the protocol used to differentiate iPSCs into NPCs.

B) Downregulation of the expression of pluripotency-associated markers (NANOG, OCT4) and

upregulation of the expression of NPC markers (PAX6, DCX, ASCL1) in untransduced (UT) and LV.hGALC-

transduced ND and GLD iPSC-derived NPC populations (assessed by SYBR green RT-PCR). Data are

normalized on GAPDH expression and shown as fold change ($2^{-\Delta\Delta CT}$) on the corresponding iPSC clone.

Data are expressed as the mean ± SEM (n=3-4 independent experiments; 2-3 replicates/clone). The

GLD1.3 clone (box) showed reduced upregulation of PAX6. Data analyzed using Kruskal Wallis test

followed by Dunnett's multiple comparison post-test (ND selected as control group). *$p<0.05$.

C) Representative immunofluorescence pictures showing the expression of the neural markers Nestin,

FOXG1 and PAX6 (green) in ND and GLD iPSC-NPC cultures. Nuclei are counterstained with DAPI (blue).

Scale bars=50μm.

D) Proliferation and stable expansion rate of untransduced and LV.hGALC transduced ND and GLD iPSC-

NPC lines (subculturing passages 0 to 5 shown in the graph; 16 days in vitro, DIV). Each point of the

curves is the mean ± SEM; n=2–3 independent experiments, 1-2 clones/group). Clones used: ND1.2,

ND1.3, ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6.

E) Schematic representation of the differentiation protocol used to differentiate iPSC-NPCs (d0) into

mixed neuronal/glial cultures that were analyzed in different experiments at day 7 (d7), d14, and d24.

GDM, Glial Differentiation Medium; GMM, Glial Maturation Medium.

F) GALC mRNA expression and G) GALC activity in untransduced (UT) and LV.hGALC-transduced iPSCs and

neural progeny (NPCs, d0; differentiated cells, d14 and d24). Values in F) are normalized on GAPDH and

expressed as (2^-ΔCT). Clones used: ND1.2, ND1.3, ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6

Data are expressed as the mean ± SEM (n=2-3 independent experiments; 1-2 replicates/clone). Data

analyzed using two-way ANOVA. To compare the different time points (iPSC, NPC d0, differentiated cells,

d14 and d24) in each group we used Turkey's multiple comparison post-test (*p < 0.05, **p < 0.01,

****p < 0.0001); to compare the different groups at each time point we used Dunnett's multiple

comparison post-test (ND selected as control group). §p < 0.05, §§p < 0.01, §§§§p < 0.0001.

G) GALC enzymatic activity in untransduced (UT) and LV.hGALC-transduced ND and GLD iPSCs and neural

progeny (NPCs, d0; differentiated cells, d14 and d24). Data are expressed as the mean ± SEM (n=2-4

replicates/clone). Data were analyzed by Kruskal Wallis test followed by Dunnett's multiple comparison

post-test (ND selected as control group); *p < 0.01; ****p < 0.0001; ns, not significant.

H) Vector copy number (VCN) measured in LV.hGALC-transduced iPSCs and neural progeny (NPCs, d0;

differentiated cells, d14). Clones used: ND2.2^{GALC}, GLD1.1^{GALC}, GLD5.2^{GALC}, GLD5.6^{GALC}; n= 1-3 independent

experiments, 1-2 clones/group.

Figure 4. GLD NPCs display impaired neuronal and glial differentiation (see also Figures S4, S5, S6, S7)

A) Representative pictures of untransduced (UT) and LV.hGALC-transduced ND and GLD differentiated

neural progeny (d24 of differentiation) expressing Olig2 (red) and APC (green). Nuclei are counterstained

with Hoechst (blue). Scale bar, 50 μm.

B) Quantification of cells expressing Olig2 and APC in UT and LV.hGALC-transduced ND and GLD

differentiated neural progeny (d14 and d24 of differentiation).

C) Representative pictures of UT and LV.hGALC-transduced ND and GLD differentiated neural progeny

(d24 of differentiation) expressing TUJ1 (green; neurons) and GFAP (red; astrocytes). Nuclei are

counterstained with Hoechst (blue). Scale bar, 50 μm.

D) Quantification of cells expressing TUJ1 and GFAP in UT and LV.hGALC-transduced ND and GLD

differentiated neural progeny (d14 and d24 of differentiation).

Data in B and D are expressed as percentage of immunoreactive (IR) cells on total nuclei and represent

the mean+SEM; n= 2-4 independent experiments; 1-3 clones/group with 2–3 replicates/clone for each

time point; 5-10 fields/coverslips were analyzed. Clones used: ND1.2, ND2.1, ND2.2, GLD1.1, GLD1.3),

GLD5.2, GLD5.6, ND2.2^{GALC}), GLD1.1^{GALC}, GLD5.2^{GALC}, GLD5.6^{GALC}). Data were analyzed by two-way ANOVA

followed by Dunnett's multiple comparison post-test (ND at the corresponding time point selected as

control group). *p<0.05, **p<0.01, *** p < 0.001.

Figure 5. Pathological hallmarks in GLD neural progeny

A) Psychosine levels in UT and LV.hGALC-transduced ND and GLD NPCs (d0) and differentiated progeny

(d14 and d24). Data are expressed as mean+SEM, n=2-4 replicates/clone, 1-3 clones/group. Clones used:

ND1.2, ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6, ND2.2^{GALC}, GLD1.1^{GALC}, GLD5.2^{GALC}, GLD5.6^{GALC}.

Data were analyzed by two-way ANOVA followed by Turkey's multiple comparison post-test;

****p<0.0001 GLD vs ND at the corresponding time point; $^{####}$ p<0.0001 GLD vs GLD^{GALC} at the

corresponding time point. p>0.05 GLD^{GALC} vs ND at all time points.

B) Quantification of cells expressing cleaved caspase 3 (CC3) in UT and LV.hGALC-transduced ND and GLD

NPCs (d0) and differentiated progeny (d14 and d24 of differentiation). Data are expressed as percentage

of immunoreactive (IR) cells on total nuclei and represent the mean+SEM; n= 2 independent

experiments; 1-3 clones/group, 2–3 replicates/clone for each time point; 5-10 fields/coverslips were

analyzed. Clones used: ND1.2, ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6, ND2.2^{GALC}, GLD1.1^{GALC},

GLD5.2^{GALC}, GLD5.6^{GALC}. Data were analyzed by two-way ANOVA followed by Dunnett's multiple

comparison post-test (ND at the corresponding time point selected as control group). No differences

between groups (p>0.05).

C) PLA2 activity in untransduced and LV.hGALC-transduced ND and GLD NPCs (d0) and differentiated

progeny (d24 of differentiation). Data are expressed as mean+SEM. n=2 independent experiments; 1-3

clones/group. Clones used: ND1.2, ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6, ND2.2^{GALC},

GLD1.1^{GALC}, GLD5.2^{GALC}, GLD5.6^{GALC}.

D) Representative pictures of lysotracker-positive NPCs (red) and quantification of lysotracker-positive

area (μm^2/cell) in ND1.2, GLD1.3 and GLD5.6 clones. Nuclei are counterstained with Hoechst. Scale

bars= 20μm. Data are expressed as the mean+SEM, n=2 independent experiments, 3 wells/clone, 1.400-

12.000 cells/well were analyzed. Data analysed by two-way ANOVA followed by Dunnett's multiple

comparison post-test (ND selected as control group). *p<0,05.

E) Representative Western Blot and densitometric quantification showing the expression of the LAMP1

protein in UT and LV.hGALC-transduced ND and GLD NPCs (d0) and differentiated progeny (d7, d14 and

d24 of differentiation). β-actin was used as loading control. Data are represented as the mean+SEM; n=

2-3 independent experiments; 1-4 clones/group. Data were analyzed by two-way ANOVA followed by

Dunnett's multiple comparison post-test (ND at the corresponding time point selected as control group).

* p<0.05, **p< 0.01; **** p < 0.0001.

Figure 6. GLD5 NPCs progeny display markers of senescence (see also Figure S8)

A) Representative confocal immunofluorescence pictures and quantification of cells expressing nestin

(green) and Ki67 (red) in ND, GLD, and GLD^GALC NPCs (d0; shown in the pictures) and differentiated

cultures (d7 and d14 of differentiation). Nuclei are counterstained with Hoechst. Scale bars= 50µm. Data

in the graph are expressed as the mean+SEM n= 3 independent experiments; 1-3 clones/group with 1–3

replicates/clone for each time point; 5-10 fields/coverslips were analyzed. Clones used: ND1.2, ND1.3,

ND2.1, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6).

B) Quantification of EdU positive cells (after a 16 hour-pulse) in ND, GLD, and GLD^GALC NPCs (d0) and

differentiated cultures (d7 and d14 of differentiation). Data in the graph are expressed as percentage

over total cells (mean); n= 2 independent experiments for each time point, 1-2 clones/group. Clones

used: ND1.2, ND2.2, GLD1.1, GLD5.2, GLD5.6, GLD1.1^GALC, GLD5.2^GALC.

C) Expression of p21 mRNA in ND, GLD, and GLD^GALC NPCs (d0) and differentiated cultures (d14 of

differentiation). n= 4 independent experiments; 1-2 clones/group. Clones used: ND1.2, ND2.2, GLD1.1,

GLD1.3, GLD5.2, GLD5.6, GLD1.1^GALC, GLD5.2^GALC, GLD5.6^GALC. Data are normalized on the housekeeping

gene glucuronidase β (*GUSB*) and expressed as $2^{-\Delta CT}$. Data are represented as mean+SEM. Data were

analyzed by Kruskal Wallis test followed by Dunns' multiple comparison post-test (ND at the

corresponding time point selected as control group). * $p<0.05$.

D) Representative Western blot and densitometric quantification showing the expression of the p21

protein in ND, GLD, and GLD^GALC NPC differentiated progeny at d7 and d14 of differentiation (d14 shown

in the blot). Calnexin was used as loading control. Data are expressed as fold vs ND at the corresponding

time point (mean of all clones) and represented as the mean+SEM; n= 2 independent experiments; 1-3

clones/group. Clones used: ND1.2, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6, GLD1.1^GALC, GLD5.2^GALC,

GLD5.6^GALC. Data were analyzed by two-way ANOVA followed by Dunnett's multiple comparison post-test

(ND at the corresponding time point selected as control group). * $p<0.05$.

E) Representative confocal immunofluorescence pictures and quantification for 53BP1-positive DDR foci

in ND, GLD, and GLD^GALC NPC (d0) and differentiated cultures (d7 and d14 of differentiation; d14 shown

in the pictures). Nuclei are counterstained with DAPI. Scale Bar = 15 µm (for all panels, shown in ND).

Clones used: ND1.2, ND2.2, GLD1.1, GLD1.3, GLD5.2, GLD5.6, GLD1.1^GALC, GLD5.2^GALC, GLD5.6^GALC. Data in

the graph are expressed as the mean+SEM; n= 3 independent experiments, 2-3 replicates/clone for each

time point. 100 to 600 nuclei per condition were analyzed. Data were analyzed by Kruskal Wallis test

followed by Dunns' multiple comparison post-test (ND at the corresponding time point selected as

control group). * p<0.05, **p < 0.01.

F) Representative pictures (brightfield and immunofluorescence for nuclei) and quantification of SA-

βGal-positive cells in ND, GLD, and GLD^{GALC} NPCs (d0) and differentiated cultures (d7 and d14 of

differentiation; d7 shown in the pictures). Nuclei are counterstained with Hoechst. Scale bars= 50μm (for

all panels, shown in ND). Data in the graph are expressed as the mean+SEM n= 3 independent

experiments; 1-3 clones/group, 1–3 replicates/clone for each time point; 4-10 fields/coverslips were

analyzed. Clones used: ND1.2, ND2.1, ND2.2, GLD1.1, GLD5.2, GLD5.6, GLD1.1^{GALC}, GLD5.2^{GALC},

GLD5.6^{GALC}. Data were analyzed by two-way ANOVA followed by Dunnett's multiple comparison post-test

(ND at the corresponding time point selected as control group). * p<0.05.

Figure 7. Lipidomic profile during iPSC neural differentiation (see also Figure S9)

A) Hierarchical clustering of lipid species differently enriched in iPSCs, NPCs and differentiated cultures

(d14). ANOVA statistical test was applied (FDR<0.05).

B) Hierarchical clustering of lipid species differently enriched in ND, GLD1 and GLD5 iPSCs, NPCs and

differentiated cultures (d14 of differentiation). ANOVA statistical test was applied (FDR<0.05).

C) Principal component analysis (PCA) of lipid species distribution in ND, GLD and LV.hGALC transduced

GLD iPSCs, NPCs and differentiated cultures (d14 of differentiation). Clones used: ND2.2, GLD1.1, GLD5.2,

ND2.2^{GALC}, GLD1.1^{GALC}, GLD5.2^{GALC}. See methods for abbreviation of lipid classes.

Methods

Reagents generated in this study are available upon request from the Lead Contact with a completed

Materials Transfer Agreement.

The lipidomic dataset generated during this study is uploaded at MetaboLights, weblink

https://www.ebi.ac.uk/metabolights/MTBLS1501. The public release date is set to Feb 15 00:00:00 GMT

2021.

Generation of iPSCs from human fibroblasts.

Human dermal fibroblasts from healthy donors (ND1 and ND2) were purchased from Life technologies.

Fibroblasts from GLD patients (GLD1, GLD2, GLD3, GLD4, and GLD5) and of a non-affected relative of

GLD1 (ND3) were obtained from the Cell Line and DNA Bank of Patients affected by Genetic Diseases

(Gaslini Institute, Genova, Italy). Human fibroblasts were cultured in Dulbecco's modified Eagle's

medium (DMEM) supplemented with 10% fetal bovine serum (FBS), L-glutamine, and

penicillin/streptomycin in an humidified atmosphere, 5% CO_2 at 37°C. Fibroblasts at passages (p) 4-5

(GLD1, GLD2, GLD3, ND1, ND2) and p10-11 (GLD4 and GLD5, ND3) were used for reprogramming.

Reprogramming was outsourced to BioERA (Padova, Italy). Reprogramming factors were delivered as

modified mRNAs (mmRNAs) taking advantage of a microfluidic system (Luni et al., 2016). The StemMACS

mRNA Reprogramming Kit (Miltenyi Biotec, Bergisch Gladbach, Germany) was used for mRNA

transfection according to the manufacturer instructions. Colonies positive for NANOG and TRA-1-60

were catalogued as iPSCs with progressive numbering. Six randomly selected colonies/donor were

picked and expanded for 2 passages. Cells were then frozen; two frozen vials/donor were sent to SR-

Tiget. The iPSCs obtained from BioERA were thawed and cultured on matrigel-coated (BD Biosciences,

Erembodegen, Belgium) plastic dishes in StemMACS™ iPS-Brew XF medium (Miltenyi Biotech) in an

humidified atmosphere, 5% O_2, 5% CO_2 at 37°C. The medium was changed daily. For routine passage,

cells were detached in PBS containing 0,5 mM EDTA and split at 1:3 to 1:10 ratio. Accutase (Sigma-

Aldrich, Saint Louis, USA) was used for single cell dissociation of iPSC colonies.

Human cells were used according to the guidelines on human research issued by the San Raffaele

Scientific Institute's ethics committee (protocol TIGET-HPCT).

Karyotype analysis.

Karyotyping was performed as previously described (Meneghini et al., 2017). Briefly, iPSC cultures were

treated with KaryoMAX colcemid 0,1 mg/ml (Life Technologies, Carlsbad, USA) for 2 h at 37°C. After

hypotonic treatment with 0,075 M KCl and fixation in methanol: acetic acid (3:1), the cell suspension was

dropped onto a microscopy slide and air-dried. Chromosome counts and G-banding karyotype analyses

were done on metaphases stained with Vectashield mounting medium with DAPI (Vector Laboratories,

Burlingame, USA). Images were analysed with Applied Imaging Software CytoVision (CytoVision Master

System). We counted >10 metaphases/coverslip. The karyotype is considered normal when >90% of the

counted metaphases are normal (46, XX or 46, XY).

Embryoid Bodies formation.

iPSC colonies were gently detached using a scraper, dissociated in small cell clumps using a 2ml pipette,

plated in Ultra-low attachment plates (Corning, New York, USA) and grown in suspension for 10 days in

EB medium: DMEM-F12+20% Knockout Serum Replacement (KSR), non-essential aminoacids, Sodium

Piruvate, L-glutamine, penicillin/streptomycin and 0,1 mM β-mercaptoethanol. The resulting spheroids

were allowed to reattach on matrigel-coated 6-well plates or glass coverslips to continue differentiation

for additional 10 days. Finally, they were collected for RNA extraction or fixed with 4% paraformaldehyde

(PFA) for immunofluorescence analyses.

iPSCs differentiation into NPCs.

For neural induction of iPSCs we applied a dual-smad inhibition protocol (Chambers et al., 2009, 2011).

Briefly, iPSC colonies were detached using Accutase, suspended as single cells and plated on Matrigel-

coated dishes (75,000 cells/cm^2) in StemMACS™ iPS-Brew XF medium supplemented with 10 µM of

ROCK inhibitor Y-27632 (Sigma-Aldrich). At ≈90% confluence (usually 2 days after plating), culture

medium was replaced with KSR medium (Chambers et al., 2009, 2011) supplemented with 200 ng/ml of

recombinant human Noggin (R&D Systems, Minneapolis, USA) and 10 µM of SB431542 (Sigma-Aldrich).

The medium was changed daily for the next 3 days. At day 4, KSR medium was gradually switched to N2

medium (Chambers et al., 2009, 2011) containing Noggin and SB431542 (75-25%, 50-50%, 25-75% every

two days). After 10 days, cells were detached using Accutase and plated on Matrigel-coated dishes in N2

medium supplemented with 20 ng/ml bFGF, 20 ng/ml EGF and 10 µM Y-27632. In these conditions, the

newly generated NPCs can be expanded for 4-6 passages. For all the differentiation experiments we used

NPCs at passages 2-3.

Assessment of long-term cell proliferation.

iPSCs or NPCs were dissociated with Accutase and plated at a density of 30,000 or 50,000 cells/cm^2,

respectively, in the appropriate medium supplemented with 10 μM Y-27632. After 4 days, or in any case

at full cell confluence, cells were treated with Accutase and counted. Cells were then replated at the

initial density. The procedure was repeated for at least 4 passages.

iPSC-NPCs differentiation into mixed neuronal/glial cultures.

We applied a published protocol (Frati et al., 2018) with some modifications. Cells were detached using

Accutase and plated on Matrigel-coated dishes (20,000 cells/cm^2) in N2 medium with 20 ng/ml bFGF, 20

ng/ml EGF and 10 μM Y-27632 (d0). Two days after plating and for the subsequent 4 days we gradually

replaced N2 medium with increasing amount of Glial Differentiation Medium (GDM) containing 10 ng/ml

PDGF-AA, 10 ng/ml NT3, 10 ng/ml IGF-1, 5 ng/ml HGF, and 60 ng/ml T3. From d4 to the end of the

differentiation protocol (d24) medium was changed every other day. From d14 to day 24 cells were kept

in Glial Maturation Medium (GMM) in absence of growth factors and supplemented with 20 μg/ml of

ascorbic acid (Sigma-Aldrich).

Lentiviral vector-mediated *hGALC* gene transfer in iPSCs.

We used a previously described (Meneghini et al., 2016) third-generation lentiviral vector (LV) encoding

the human (h)GALC cDNA tagged with the myc peptide, under control of the human phosphoglycerate

kinase (PGK) promoter. The titer of the LV batch was 1.6x10^9 TU/ml. For cell transduction, iPSC colonies

were dissociated at single cells using Accutase and plated on Matrigel-coated 6 multiwell plates (10,000

cells/cm^2) in StemMACS™ iPS-Brew XF medium supplemented with 10 μM Y-27632. After 6 hours, cells

were incubated with LV.hGALC vector at different MOI (0-100) in StemMACS™ iPS-Brew XF medium

additioned with 10 μM Y-27632 and 8 μg/ml Polybrene. After an obvernight incubation, we added fresh

culture medium. Cells were passaged as they reached confluency, as described. Transduced cells were

used for the experiments after 3-4 subculturing passages, in order to dilute the non-integrated LV

genome.

Quantification of Psychosine.

Psychosine quantification was performed on frozen cell pellets by TSQ Quantum AM mass spectrometer

(Thermo Fisher Scientific, Waltham, USA) as previously described (Lattanzi et al., 2014).

GALC activity assay

Specific beta-galactocerebrosidase activity was measured by hydrolysis of artificial fluorogenic substrate

4-methylumbelliferone-beta-galactopyranoside in presence of the competitive inhibitor of beta-

galactosidase AgNO$_3$, as previously described (Martino et al., 2009).

PLA2 activity assay

Quantification of Phospholipase A2 (PLA2) activity was performed using the EnzChek Phospholipase A2

Assay Kit (Thermo Fisher). 50 μl of cell lysate/well, corresponding to 20 μg total proteins, was mixed with

50 μl of substrate cocktail to initiate the catalytic reaction. After 10 minutes, fluorescence was measured

using a Victor3 Multilabel Counter (Perkin Elmer, Waltham, USA) at 360 nm (excitation) and 460 nm

(emission). The specific enzymatic activity was calculated for each sample and normalised for protein

content. Values are expressed as relative fluorescent units (RFU)/μg of protein/min.

SA-β-Galactosidase assay (X-gal staining).

Induction of senescence during NPCs differentiation was assessed with a senescence β-galactosidase (SA-

β-Gal) Staining Kit (Cell Signalling Technology, Danvers, USA) according to manufacturer's instructions. In

detail, NPCs were plated on coverslips and fixed at day 0, day 7 and day 14 days of differentiation

with4% PFA for 10 minutes at RT and then incubated for 24 hours with the SA-β-gal staining solution

(pH=6.0) to reveal SA-β-gal activity. Nuclei were then stained with Hoechst. Images were acquired using

a Nikon Eclipse Ni microscope, visualized and analysed using the ImageJ software. Cells displaying blue

signal in the cytosol were counted as positive. At least 850 nuclei/coverslip/timepoint were used for

quantification.

Proliferation analysis (EdU staining)

EdU (5-ethynyl-2'-deoxyuridine), supplied with Click-iT EdU Alexa Fluor 647 Imaging Kit (Thermo-Fisher

Scientific), was diluted in DMSO to a final concentration of 10 mM and kept at −20°C. Cells were

incubated with 2μM EdU overnight in culture. 2x10^5 cells per sample were transferred to cytometry

tubes and then washed with PBS 1% BSA and fixed with 100 μL of Click-iT fixative for 15 minutes. Cells

were washed again with PBS 1% BSA and permeabilized with 100 μL of 1X Click-iT saponin-based solution

for 15 minutes. Detection of EdU-DNA was performed by incubating cells with 500 μL of Click-iT Plus

reaction cocktail for 30 minutes at RT protected from light. Cells were subsequently washed with PBS 1%

BSA and fluorescence was measured by flow cytometry.

High Throughput live cell Microscopy

Determination of lysosomal area was performed using a high content imaging platform in the Advanced Light and Electron Microscopy Bio-Imaging Centre of San Raffaele Scientific Institute (ALEMBIC) using the ArrayScan XTI HCA Reader from Thermo Fisher Scientific. Cells were plated on matrigel-coated 96 Greiner Sensoplate glass bottom multiwell plates (Sigma Aldrich) (30,000 cells/well). The day after plating, cells are incubated with 50 nM LysoTracker Red DND-99 (Thermo Fisher) diluted in culture medium, for 30 minutes at 37°C. Nuclear staining, necessary for the instrument acquisition and analysis, was performed using Hoechst (5 min at room temperature, 10 µg/ml final concentration). Cells are then washed with Krebs-Ringer-Hepes-bicarbonate buffer (KRH) and maintained in the same buffer for the entire duration of the instrument acquisition. Thirty fields were acquired for each well; images were analysed with the Thermo Scientific HCS Studio Cell Analysis Software

Immunofluorescence.

Cells are seeded in 48-multiwell plates on Matrigel-coated 10 mm round sterile glass coverslips, fixed in 4% PFA, rinsed with PBS, incubated with blocking solution containing 10% normal goat serum (NGS) + 0.1% Triton X-100 in PBS for 45 minutes and incubated overnight at 4°C with primary antibodies diluted in blocking solution. After 3 washes (10 minutes each), cells are incubated with species-specific fluorophore-conjugated secondary antibodies diluted in 1% NGS in PBS for 1 hour at room temperature. Cell nuclei are counterstained with Hoechst (10µg/ml, diluted in PBS). Finally, coverslips are mounted on glass slides using Fluorsave mounting medium (Merck Millipore). Samples incubated in blocking solution without primary antibodies were used as negative controls. Images were obtained with a Nikon Eclipse Ni microscope, visualized and analysed using the ImageJ software.

To evaluate DDR activation during NPCs differentiation, cells were fixed with 4% PFA for 20 min at RT and then permeabilized with 0.3% Triton-X100 in PBS for 10 min. After washing, coverslips were blocked in PBG (0.2% cold water fish gelatine, 0.5% BSA in PBS) for 30 min. Cells were then stained with primary antibodies 1 hour at RT followed by secondary antibodies. After DAPI staining (Sigma-Aldrich), slides were mounted with Aqua-Poly/mount (Polysciences Inc, Warrington, USA). Images were obtained with Leica TCS SP5 confocal laser microscope and visualized with Leica Application Suite (LAS) software (Leica Microsystems, Wetzlar, Germany). Quantification of DDR activation in immunofluorescence images was conducted by counting colocalizing foci for 53BP1 and γH2A.X.

Primary and secondary antibodies used:

Primary antibodies		
Antigen	**Host species (company, catalog number)**	**Dilution**
53BP1	Rabbit polyclonal (Bethyl, A300-272A-5)	1: 600
α-fetoprotein	Mouse monoclonal (Abcam, AB3980)	1: 200
α-Smooth Muscle Actin	Mouse monoclonal (Sigma-Aldrich, C6198)	1: 400
APC	Mouse monoclonal (Calbiochem, OP80)	1: 300
β-III tubulin (TUJ1)	Mouse monoclonal (Biolegend, 801202)	1: 1,000
γH2A.X	Mouse monoclonal (Biolegend, 613402)	1: 200
Cleaved Caspase 3	Rabbit polyclonal (Cell Signaling, 9961)	1: 200
FOXG1	Rabbit polyclonal (Abcam, AB 18259)	1: 300
GALC	Mouse monoclonal, kindly provided by Dr. Deane	1: 20
GalCer	Mouse monoclonal (Millipore MAB342)	1: 300
GFAP	Rabbit polyclonal (DAKO, ZO334)	1: 1,000
Ki67	Rabbit polyclonal (Novocastra, NCL-Ki67-p)	1: 200
NANOG	Rabbit polyclonal (Abcam, AB21624)	1: 200
Nestin	Mouse monoclonal (Millipore, MAB353)	1: 500
OCT4	Rabbit monoclonal (Abcam, AB136253)	1: 300
Olig2	Rabbit monoclonal (Abcam, AB136253)	1: 300
PAX6	Rabbit polyclonal (Covance, PRB-278P)	1: 300
TRA 1-60	Mouse monoclonal (Millipore, MAB4360)	1: 100
Secondary antibodies		
Conjugate	**Host species (company, catalog number)**	**Dilution**
Alexa 488	Goat anti-mouse and Goat anti-rabbit IgG (Mol.Probes, A11001 and A11008)	1: 1,000
Alexa 546	Goat anti-mouse and Goat anti-rabbit IgG (Mol.Probes, A11003 and A11010)	1: 2,000
Alexa 568	Goat anti-rabbit IgG (Mol.Probes, A11031)	1: 2,000

Western blot analysis.

Cell pellets were lysed using RIPA buffer (50 mM Tris–HCl pH 7,4, 150 mM NaCl, 0,5% sodium

deoxycholate, 0,1% SDS, 2 mM EDTA) containing protease and phosphatase inhibitors (Roche, Basel,

Switzerland). Protein content was measured using the Bradford reagent (Bio-Rad, Hercules, USA) with

bovine serum albumin (BSA) as the reference standard. SDS-PAGE of protein extracts was performed

using Novex NuPAGE SDS-PAGE system (Thermo Fisher Scientific) according to manufacturer instructions.

Proteins were transferred on PVDF membranes (Millipore, Burlington USA) for 1 hour at 400mA.

Primary and secondary antibodies used:

Primary antibodies		
Antigen	**Host species (company, catalog number)**	**Dilution**
α-tubulin	Mouse monoclonal (Sigma-Aldrich, T6074)	1: 5,000
Acetylated α-tubulin	Mouse monoclonal (Sigma-Aldrich, T7451)	1: 20,000
β-Actin	Rabbit polyclonal (Santa Cruz, sc-1616r)	1: 10,000
β-Actin	Mouse monoclonal (Sigma-Aldrich, A2228)	1: 10,000
β-III tubulin (TUJ1)	Mouse monoclonal (Biolegend,801202)	1: 6,000
Calnexin	Rabbit polyclonal (Sigma-Aldrich, c4731)	1: 10,000
GFAP	Mouse monoclonal (Millipore, MAB3402)	1: 100,000
LAMP1	Rabbit polyclonal (Abcam, AB24170)	1: 6,000
MYC	Rabbit polyclonal (Abcam, AB9106)	1: 5,000
Neurofilament-M (RMO 14.9)	Mouse monoclonal (Cell Signaling, 2838)	1: 1,000
OLIG1	Rabbit polyclonal (Millipore, AB5591)	1: 800
p21	Rabbit monoclonal (Cell Signaling, 2947)	1: 1,000
Polyglutamylation Modification	Mouse monoclonal (Adipogen, AG-20B-0020)	1: 5,000
SMI 311	Mouse monoclonal (Sternberg, SMI-311)	1:1000
SMI 312	Mouse monoclonal (Covance, SMI312R)	1:1000
Secondary antibodies		
Conjugate	**Host species (company, catalog number)**	**Dilution**
HRP	Goat anti-mouse IgG (Chemicon, AP124P)	1: 5,000
HRP	Goat anti-rabbit IgG (Millipore, 12-348)	1: 10,000

Molecular analysis

Sequencing of patient's mutations. Genomic DNA was extracted from iPSCs pellets using Maxwell cell

DNA purification Kit (Promega), following the manufacturers' instructions. DNA concentration was

determined by 260/280nm optical density (OD) reading on a NanoDrop ND-1000 Spectrophotometer

(NanoDrop, Pero, Italy). To confirm the presence of the expected mutations in the *GALC* gene of GLD

patients-derived cells, the regions of interests were amplified by PCR, purified from agarose gels using

the Wizard SV Gel and PCR Clean-Up System (Promega, Madison, USA) and sent for sequencing (to GATC

Services-Eurofin Genomics, Ebersberg, Germany). Sequences were checked using the Snapgene

software. Three primer PCR to confirm 30kb deletion in GLD1 and GLD3 cells was performed as described

by (Liu et al., 2011). Primers used for PCR amplification and sequencing are the following:

Primers for gDNA sequencing	
GALC region	**Sequence**
INTRON 11	Fw 5' - CCTATATGGAAAACAATGTGG - 3',
	Rev 5' - AAGGAGCAACATTTCAGGC - 3' (WT allele)
	Rev 5' - TCAAGTCCTTGATGATCACC - 3' (deleted allele)
EXON 4	Fw 5'- GGTGGGGAGTGAGATGGTC - 3',
	Rev 5' - CAAGGGCAAAGAAAGGATCA - 3'
EXON 8	Fw 5' - TGCTGAGACAAAAGGGCATA - 3',
	Rev 5' - GATGACGCTAACAAGGCAAA - 3'
EXON 9	Fw 5' - TTGGGTGATCCTTTTATTGTCA -3',
	Rev 5' - CACTGGCAAATCTTGCTTAAAA - 3'
EXON 14	Fw 5' - AGCAAGGAGAGCTTCTGAAGGA - 3',
	Rev 5' - AGGTTCTTGAAATAGGAGGACCA - 3'

Gene expression analysis: RT-PCR; SYBR green and TaqMan RT-PCR. Total RNA was extracted using the

RNeasy Mini Kit (Qiagen). RNA was quantified by 260/280nm optical density (OD) reading on a NanoDrop

ND-1000 Spectrophotometer (NanoDrop, Pero, Italy). mRNA reverse transcription (RT) is performed with

QuantiTect reverse transcription kit (Qiagen, Hilden, Germany).

RT-PCR analyses were performed by using FastStart Taq DNA Polymerase, PCR KIT (Roche). For each

reaction, 100 ng of template cDNA and 1 mM of primers were used. PCR products were resolved by

electrophoresis in a 1,5% agarose gel.

Quantitative real time PCR reactions were performed in Optical 96-well Fast Thermal Cycling Plates

(Primo 96-well plates, Euro Clone, Pero, Italy) on Viia7 quantitative real time PCR system (Applied

Biosystem, Foster City, USA), using the following thermal cycling conditions:

- SYBR green qPCR: one cycle at 95°C for 2 minutes, 40 cycles at 95°C for 10 seconds and 60°C for 30

seconds; each sample is run in duplicate in a total volume of 12.5 µl/reaction, containing 6.25 µl 2X Fast

SYBR Green Master Mix (Qiagen), 10 ng of template cDNA, and10µM of custom primers were used. As

internal reference for normalization GAPDH was used

- TaqMan qPCR: one cycle at 95°C for 5 minutes, 40 cycles at 95°C for 50 seconds and 60°C for 1 minute.

Each sample is run in duplicate in a total volume of 12.5 µl/reaction, containing 6.25 µl 2X EagleTaq

Universal Master Mix (Roche), 1µl of sample cDNA and 0.63µl of Probe+Primers mix (TaqMan Gene

Expression Assays, Thermo Fisher). The Viia7 software was used to extract raw data. Relative expression

of mRNA for the target genes was calculated using the comparative Ct (ΔΔCt) method, with a

housekeeping gene as control. Raw data (Ct values) were analyzed according to the comparative Ct

method. For gene expression data, analyses were performed on 2($-^{\Delta Ct}$) values relative to housekeeping

gene or plotted as $2^{-\Delta\Delta Ct}$ (fold change), as specified in figure legends. Primers and probes used for gene

expression analyses are the following:

Primers for RT-PCR	
Gene	**Sequence**
KLF4	Fw 5 ' -ACCCACACAGGTGAGAAACCTT - 3'
	Rev 5' - GTTGGGAACTTGACCATGATTG - 3'
SOX2	Fw 5' - TTACCTCTTCCTCCCACTCCAG - 3'
	Rev 5' - GGGTTTTCTCCATGCTGTTTCT - 3'
NANOG	Fw 5'- GAT CGG GCC CGC CAC CAT GAG TGT GGA TCC AGC TTG-3'
	Rev 5'- GAT CGA GCT CCA TCT TCA CAC GTC TTC AGG TTG - 3'
CD146	Fw 5'- AAG GCA ACC TCA GCC ATG TC - 3'
	Rev 5'- AGA AGA TGC GCT CGT CTT GG - 3'
CXCR4	Fw 5' - CAG CAG GTA GCA AAG TGA CG - 3'
	Rev 5' - TTG TCC GTC ATG CTT CTC AG - 3'
MAP2	Fw 5' - AAG AAG GTC GCC ATC ATA CG - 3'
	Rev 5' - GGG CTT TAG CAT GCT CTC TG - 3
PAX6	Fw 5' - AAC AGA CAC AGC CTT CAC AAA CA - 3'
	Rev 5' - CGG GAA CTT GAA CTG GAA CTG AC - 3'
SOX17	Fw 5' - TTC ATG GTG TGG GCT AAG GAC - 3'
	Rev 5' - AGT TGG GGT GGT CCT GCA TG - 3'
α-SMA	Fw 5' - GTG TTG CCC CTG AAG AGC AT - 3'
	Rev 5' - GCT GGG ACA TTG AAA GTC TCA - 3'
GAPDH	Fw 5' - CAG CCT CAA GAT CAT CAG CA - 3'
	Rev 5' - TGC TGT AGC CAA ATT CGT TG - 3'
Primers for SYBR-green RT-PCR	
Gene	**Sequence**
ASCL1	Fw 5' - TCCCCCAACTACTCCAACGA - 3'
	Rev 5' - GCGATCACCCTGCTTCCAAA - 3'
CDKN1A	Fw 5' - CAGCATGACAGATTTCTACCACTC - 3'
	Rev 5' -CTCGCGCTTCCAGGACTG - 3'
CDKN2A	Fw 5' - CCAACGCACCGAATAGTTACG - 3'
	Rev 5' - GCGCTGCCCATCATCATG - 3'
DCX	Fw 5' - TGGATGAAAATGAATGCCGAGT - 3'
	Rev 5' - AGCTGGAGACTTGCTTCGG - 3'
GAPDH	Fw 5' - TCAAGAAGGTGGTGAAGCAGG - 3'
	Rev 5' - ACCAGGAAATGAGCTTGACAAA - 3'

GUSB	Fw 5' - CTGACACCTCCAAGTATCCCAAG - 3'
	Rev 5' - GTCGTGTACAGAAGTACAGACCGC - 3'
IL1β	Fw 5' - GCTCAAGTGTCTGAAGCAGCC - 3'
	Rev 5' - CAGCTTCAAAGAACAAGTCATCCT - 3'
NANOG	Fw 5' - ATGCCTCACACGGAGACTGT - 3'
	Rev 5' - AAGTGGGTTGTTTGCCTTTG - 3'
OCT4	Fw 5' - TCGAGAACCGAGTGAGAGG - 3'
	Rev 5' - GAACCACACTCGGACCACA - 3'
PAX6	Fw 5' - AGTGAATCAGCTCGGTGGTGTCTT - 3'
	Rev 5' - TGCAGAATTCGGGAAATGTCGC - 3'
TNF-α	Fw 5' – CCAGGGACCTCTCTCTAATCAGC - 3'
	Rev 5' – GGTTTGCTACAACATGGGCTAC - 3'
Probe+Primers for Taqman RT-PCR	
Gene	**Commercial Code**
DCX	Hs00167057_m1
GALC	Hs01012300_m1
GAPDH	Hs02758991_g1
GFAP	Hs00157674_m1
KLF4	Hs00358836_m1
MBP	Hs00921945_m1
NANOG	Hs02387400_g1
NEUROD1	Hs01922995_s1
NKX2.2	Hs0159616_m1
OLIG2	Hs00300164_s1
OCT4	Hs03005111_g1
PDGFRα	Hs00183486_m1
SCL1A3	Hs00904823_g1
SOX2	Hs01053049_s1
SOX10	Hs00366918_m1

Determination of vector copy number (VCN)

Genomic DNA was extracted from cell pelletes using Maxwell cell DNA purification Kit (Promega) or

QIAamp DNA Micro Kit (Quiagen), depending on the number of cells. The number of integrated vector

copies/genome was quantified by TaqMan analysis, as previously described (Lattanzi et al., 2010).

Briefly, quantitative PCR was performed on 50 ng of template DNA using probes and primers that anneal

to the PSI (ψ) sequence of the LV backbone and to a fragment of the human telomerase gene. A

standard curve was set up with gDNA extracted from a human lymphoblast cell line (CEM) carrying 1

copy of integrated LV.GFP, validated by Southern blot analysis. The standard curve, based on different

dilutions of CEM gDNA (from 0,32 ng to 40 ng) was used for both LV and telomerase amplification.

Reactions were carried out in a total volume of 25 µl, on Viia7 quantitative real time PCR system (Applied

Biosystem). Eventually, VCN is calculated as: (ng LV/ng endogenous DNA) × (number of LV integrations in

the standard curve) × cell ploidy. Primers and probes used for VCN analyses are the following:

Primers for VCN assay		
Gene	Probe sequence	Primer sequence
HIV	5'- ATCTCTCTCCTTCTAGCCTC - 3'	For 5' – TACTGACGCTCTCGCACC - 3'
		Rev 5' – TCTCGACGCAGGACTCG - 3'
TELO	5' -TCAGGACGTCGAGTGGACACGGTG- 3'	For 5' – GGCACACGTGGCTTTTCG - 3'
		Rev 5' – GGTGAACCTCGTAAGTTTATGCAA - 3'

Lipidomic analyses

Lipid extraction. Cell pellets were resuspended in 150 mM Ammonium bicarbonate and passed through

a 26G syringe needle to prepare cell lysate. Lipids were extracted starting from an equivalent of 10 µg of

proteins, using a 2 steps extraction protocol (Folch method) with methanol and chloroform in different

proportions (FOLCH et al., 1957). Organic phase fractions were then dried out and resuspended in 50 µL

of CH_3OH for subsequent analysis. Before extraction, samples were spiked in with 16 internal standards:

PC (12:0/13:0) 40 pmol, PE (12:0/13:0) 52 pmol, PG (12:0/13:0) 7.5 pmol, PS (12:0/13:0) 43 pmol, PI

(12:0/13:0) 54 pmol, Cer (d18:1/25:0) 100 pmol, CE(19:0) 100 pmol, GlcCer (d18:1/12:0) 50 pmol, LacCer

(d18:1/12:0) 50 pmol, Sphinganine (d17:0) 50 pmol, Sphingosine-1-P (d17:1) 100 pmol, Sphingosine

(d17:1) 50 pmol, Galactosyl(ß) Sphingosine-d5 20 pmol, d5-TG ISTD Mix I 20 pmol, d5-DG ISTD Mix I 20

pmol, cholesterol (d7) 800 pmol.

Protein quantification. Proteins were extracted form 20 µL of ammonium bicarbonate resuspended

fractions by adding 5 µL of lysis buffer (10% NP40, 2%SDS in PBS) and quantified by BCA protein assay kit

(ThermoFisher Scientific, 23225).

Lipid profiling data acquisition. Lipids extracts were diluted 1:5 with 95% mobile phase A ($CH_3CN:H_2O$

40:60; 5mM NH_4COOCH_3; 0.1% FA) plus 5% mobile phase B ($IPA:H_2O$ 90:10; 5mM NH_4COOCH_3; 0.1%

FA) and 1 µL injected on a liquid chromatography system nLC Ekspert nanoLC400 (Eksigent, 5033460C;

Singapore) coupled with a Triple TOF 6600 (AB Sciex; Singapore). Chromatography was performed using

an in house packed nanocolumn Kinetex EVO C18, 1.7 µm, 100 A, 0.75 x 100 mm. The gradient elution at

flow rate of 150 nL/min was initially started from 5 % of mobile phase B, linearly increased to 100 % B in

5 min, maintained for 45 min, then returned to the initial ratio in 2 min and maintained for 8 min.

Positive mode acquisition in MS was performed with the following parameters: mass over charge (m/z)

range 100–1700, T source 80C, Ion Spray Voltage 2000, Declustering Potential (DP) 80, fixed collision

energy 40 V (+). For IDA analysis (Top 8), range of m/z was set as 200–1800 in positive ion mode, target

ions were excluded for 20 sec after 2 occurrences (Analyst TF 1.7.1).

Data processing and analysis. Lipidview workstation (version 1.3 beta, AB SCIEX, USA) was used for

lipids identification and quantitation. Lipid identification was based on exact mass, retention time, and

MS/MS pattern. Lipid species based on precursor fragment ion pairs were determined using a

comprehensive target list in LipidView (Sciex). Lipid species identification was performed using; the mass

tolerance of 0.05 in MS and 0.02 in MS/MS, s/n of 3 and % peak intensity > 0 for positive ion mode.

Lipid classes included for statistics and downstream analysis were: cholesterol ester (CE), sphingomyelin

(SM), diacylglycerol (DAG), triacylglycerol (TAG), ceramide (Cer) phosphatidylcholine (PC),

phosphatidylethanolamine (PE), phosphatidylglycerol (PG), phosphatidylinositol (PI), phosphatidylserine

(PS) and lysophosphatidylcholine (LPC), lysophosphatidylethanolamine (LPE), lysophosphatidylglycerol

(LPG), lysophosphatidylinositol (LPI), lysophosphatidylserine (LPS), hexosylceramide (HexCer),

dihexosylceramide (Hex2Cer), trihexosylceramide (Hex3Cer), sulphatides (SGalCer), ceramide-phosphate

(CerP), in positive mode.

Statistics

Data were analysed with Graph Pad Prism version 8 for Macintosh and expressed as the mean ± standard

error of the mean (SEM). The number of samples and the statistical tests and post-tests used are

specified in the legends to each figure. p<0.05 was considered statistically significant

For lipid analyses, statistical analysis was performed using Metaboanalyst 4.0 web tool

https://www.metaboanalyst.ca/MetaboAnalyst/faces/home.xhtml (Chong et al., 2018).

References

Allende, M.L., Cook, E.K., Larman, B.C., Nugent, A., Brady, J.M., Golebiowski, D., Sena-Esteves, M., Tifft,

C.J., and Proia, R.L. (2018). Cerebral organoids derived from Sandhoff disease-induced pluripotent

stem cells exhibit impaired neurodifferentiation. J. Lipid Res. *59*, 550-563.

Allewelt, H.B., Page, K., Taskindoust, M., Troy, J.D., Wood, S., Parikh, S., Prasad, V.K., and Kurtzberg, J.

(2016). Long-term functional outcomes following hematopoietic stem cell transplantation for krabbe

disease. Biol. Blood Marrow Transplant. *22*, S102–S103.

Ballabio, A., and Gieselmann, V. (2009). Lysosomal disorders: from storage to cellular damage. Biochim

Biophys Acta *1793*, 684–696.

Bayó-Puxan, N., Terrasso, A.P., Creyssels, S., Simão, D., Begon-Pescia, C., Lavigne, M., Salinas, S., Bernex,

F., Bosch, A., Kalatzis, V., et al. (2018). Lysosomal and network alterations in human

mucopolysaccharidosis type VII iPSC-derived neurons. Sci. Rep. *8*, 16644.

Berardi, A.S., Pannuzzo, G., Graziano, A., Costantino-Ceccarini, E., Piomboni, P., and Luddi, A. (2014).

Pharmacological chaperones increase residual beta-galactocerebrosidase activity in fibroblasts from

krabbe patients. Mol. Genet. Metab. *112*, 294–301.

Bieberich, E. (2012). It's a lipid's world: Bioactive lipid metabolism and signaling in neural stem cell

differentiation. Neurochem. Res. *37*,1208-1229.

Biffi, A., Montini, E., Lorioli, L., Cesani, M., Fumagalli, F., Plati, T., Baldoli, C., Martino, S., Calabria, A.,

Canale, S., et al. (2013). Lentiviral hematopoietic stem cell gene therapy benefits metachromatic

leukodystrophy. Science (80-.). *341*, 1233158.

Bongarzone, E.R. oque, Escolar, M.L. uisa, Gray, S.J. ames, Kafri, T., Vite, C.H. erman, and Sands, M.S.

teven (2016). Insights into the Pathogenesis and Treatment of Krabbe Disease. Pediatr. Endocrinol.

Rev. *13*, 689–696.

Borda, J.T., Alvarez, X., Mohan, M., Ratterree, M.S., Phillippi-Falkenstein, K., Lackner, A. a, and Bunnell, B.

a (2008). Clinical and immunopathologic alterations in rhesus macaques affected with globoid cell

leukodystrophy. Am J Pathol *172*, 98–111.

Bradbury, A.M., Rafi, M.A., Bagel, J.H., Brisson, B.K., Marshall, M.S., Pesayco Salvador, J., Jiang, X., Swain,

G.P., Prociuk, M.L., Odonnell, P.A., et al. (2018). AAVrh10 Gene Therapy Ameliorates Central and

Peripheral Nervous System Disease in Canine Globoid Cell Leukodystrophy (Krabbe Disease). Hum.

Gene Ther. *29,* 785-801.

Buratta, S., Urbanelli, L., Sagini, K., Giovagnoli, S., Caponi, S., Fioretto, D., Mitro, N., Caruso, D., and

Emiliani, C. (2017). Extracellular vesicles released by fibroblasts undergoing H-Ras induced senescence

show changes in lipid profile. PLoS One *12*, e0188840.

Canals, I., Soriano, J., Orlandi, J.G., Torrent, R., Richaud-Patin, Y., Jim??nez-Delgado, S., Merlin, S.,

Follenzi, A., Consiglio, A., Vilageliu, L., et al. (2015). Activity and high-order effective connectivity

alterations in sanfilippo C patient-specific neuronal networks. Stem Cell Reports *5*, 546–557.

Castelvetri, L.C., Givogri, M.I., Zhu, H., Smith, B., Lopez-Rosas, A., Qiu, X., van Breemen, R., and

Bongarzone, E.R. (2011). Axonopathy is a compounding factor in the pathogenesis of Krabbe disease.

Acta Neuropathol *122*, 35-48.

Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L. (2009).

Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. Nat.

Biotechnol. *27*, 275–280.

Chambers, S.M., Mica, Y., Studer, L., and Tomishima, M.J. (2011). Converting human pluripotent stem

cells to neural tissue and neurons to model neurodegeneration. Methods Mol. Biol. *793*, 87–97.

Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., Wishart, D.S., and Xia, J. (2018). MetaboAnalyst

4.0: Towards more transparent and integrative metabolomics analysis. Nucleic Acids Res. *46*, W486-

W494.

Claus, R., Dorer, M., Bunck, A., and Deigner, H. (2009). Inhibition of Sphingomyelin Hydrolysis: Targeting

the Lipid Mediator Ceramide as a Key Regulator of Cellular Fate. Curr. Med. Chem. *16*, 1978-2000.

Crowe, E.P., Tuzer, S.F., Cohen, J., Yetkin, E.C., D'Agostino, L., Sell, C., and Torres, C. (2016) Implications

of Cellular Senescence on Aging and Disease in the Brain. In: Rattan S., Hayflick L. (eds) Cellular Ageing

and Replicative Senescence. Healthy Ageing and Longevity. Springer, Cham

Duffner, P.K., Barczykowski, A., Jalal, K., Yan, L., Kay, D.M., and Carter, R.L. (2011). Early infantile Krabbe

disease: Results of the world-wide Krabbe registry. Pediatr. Neurol. *45*, 141–148.

Duffner, P.K., Barczykowski, A., Kay, D.M., Jalal, K., Yan, L., Abdelhalim, A., Gill, S., Gill, A.L., and Carter, R.

(2012). Later onset phenotypes of Krabbe disease: Results of the world-wide registry. Pediatr. Neurol.

46, 298–306.

Escolar, M.L., Kiely, B.T., Shawgo, E., Hong, X., Gelb, M.H., Orsini, J.J., Matern, D., and Poe, M.D. (2017).

Psychosine, a marker of Krabbe phenotype and treatment effect. Mol. Genet. Metab. *121*, 271-278.

Folch, J., Lees, M., and Sloane Stanley, G.H. (1957). A simple method for the isolation and purification of

total lipides from animal tissues. J. Biol. Chem. *226*, 497-509.

Folts, C.J., Scott-Hewitt, N., Pröschel, C., Mayer-Pröschel, M., and Noble, M. (2016). Lysosomal Re-

acidification Prevents Lysosphingolipid-Induced Lysosomal Impairment and Cellular Toxicity. PLoS

Biol. *14*.

Frati, G., Luciani, M., Meneghini, V., De Cicco, S., Ståhlman, M., Blomqvist, M., Grossi, S., Filocamo, M.,
Morena, F., Menegon, A., et al. (2018). Human iPSC-based models highlight defective glial and
neuronal differentiation from neural progenitor cells in metachromatic leukodystrophy. Cell Death
Dis. *9*, 698.

Gama Sosa, M.A., de Gasperi, R., Undevia, S., Yeretsian, J., Rouse 2nd, S.C., Lyerla, T.A., and Kolodny,
E.H. (1996). Correction of the galactocerebrosidase deficiency in globoid cell leukodystrophy-cultured
cells by SL3-3 retroviral-mediated gene transfer. Biochem Biophys Res Commun *218*, 766–771.

De Gasperi, R., Gama Sosa, M.A., Sartorato, E.L., Battistini, S., MacFarlane, H., Gusella, J.F., Krivit, W., and
Kolodny, E.H. (1996). Molecular heterogeneity of late-onset forms of globoid-cell leukodystrophy.
Am. J. Hum. Genet. *59*, 1233–1242.

De Gasperi, R., Gama Sosa, M.A., Sartorato, E., Battistini, S., Raghavan, S., and Kolodny, E.H. (1999).
Molecular basis of late-life globoid cell leukodystrophy. Hum Mutat *14*, 256–262.

Gentner, B., Visigalli, I., Hiramatsu, H., Lechman, E., Ungari, S., Giustacchini, A., Schira, G., Amendola, M.,
Quattrini, A., Martino, S., et al. (2010). Identification of hematopoietic stem cell-specific miRNAs
enables gene therapy of globoid cell leukodystrophy. Sci. Transl. Med. *2*, 58ra84.

Giri, S., Khan, M., Rattan, R., Singh, I., and Singh, A.K. (2006). Krabbe disease: psychosine-mediated
activation of phospholipase A2 in oligodendrocyte cell death. J Lipid Res *47*, 1478–1492.

Graziano, A.C.E., and Cardile, V. (2015). History, genetic, and recent advances on Krabbe disease. Gene.
555(1), 2-13

Del Grosso, A., Angella, L., Tonazzini, I., Moscardini, A., Giordano, N., Caleo, M., Rocchiccioli, S., and
Cecchini, M. (2019). Dysregulated autophagy as a new aspect of the molecular pathogenesis of
Krabbe disease. Neurobiol. Dis. *129*, 195-207.

Gupta, A., Poe, M.D., Styner, M.A., Panigrahy, A., and Escolar, M.L. (2015). Regional differences in fiber
tractography predict neurodevelopmental outcomes in neonates with infantile Krabbe disease.
NeuroImage Clin. *7*, 792–798.

Hannun, Y.A., and Obeid, L.M. (2002). The ceramide-centric universe of lipid-mediated cell regulation:
Stress encounters of the lipid kind. J. Biol. Chem. *277*, 25847-25850.

Hannun, Y.A., and Obeid, L.M. (2008). Principles of bioactive lipid signalling: Lessons from sphingolipids.
Nat Rev Mol Cell Biol *9*, 139–150.

Hannun, Y.A., and Obeid, L.M. (2017). Sphingolipids and their metabolism in physiology and disease. Nat.
Rev. Mol. Cell Biol. *19*, 175-191.

Hawkins-Salsbury, J. a, Parameswar, A.R., Jiang, X., Schlesinger, P.H., Bongarzone, E., Ory, D.S.,

Demchenko, A. V, and Sands, M.S. (2013). Psychosine, the cytotoxic sphingolipid that accumulates in

globoid cell leukodystrophy, alters membrane architecture. J. Lipid Res. *54*, 3303–3311.

Hawkins-Salsbury, J.A., Shea, L., Jiang, X., Hunter, D.A., Guzman, A.M., Reddy, A.S., Qin, E.Y., Li, Y., Gray,

S.J., Ory, D.S., et al. (2015). Mechanism-based combination treatment dramatically increases

therapeutic efficacy in murine globoid cell leukodystrophy. J. Neurosci. *35*, 6495–6505.

He, S., and Sharpless, N.E. (2017). Senescence in Health and Disease. Cell. *169*, 1000-1011.

Higuchi, T., Kawagoe, S., Otsu, M., Shimada, Y., Kobayashi, H., Hirayama, R., Eto, K., Ida, H., Ohashi, T.,

Nakauchi, H., et al. (2014). The generation of induced pluripotent stem cells (iPSCs) from patients

with infantile and late-onset types of Pompe disease and the effects of treatment with acid-??-

glucosidase in Pompe's iPSCs. Mol. Genet. Metab. *112*, 44–48.

Huang, H.-P., Chuang, C.-Y., and Kuo, H.-C. (2012). Induced pluripotent stem cell technology for disease

modeling and drug screening with emphasis on lysosomal storage diseases. Stem Cell Res. Ther. *3*, 34.

Igisu, H., and Suzuki, K. (1984). Progressive accumulation of toxic metabolite in a genetic leukodystrophy.

Science (80-.). *224*, 753–755.

Irahara-Miyana, K., Otomo, T., Kondo, H., Hossain, M.A., Ozono, K., and Sakai, N. (2018). Unfolded

protein response is activated in Krabbe disease in a manner dependent on the mutation type. J. Hum.

Genet. *63*, 699–706.

Jalal, K., Carter, R., Yan, L., Barczykowski, A., and Duffner, P.K. (2012). Does galactocerebrosidase activity

predict krabbe phenotype? Pediatr. Neurol. *47*, 324–329.

Jana, A., Hogan, E.L., and Pahan, K. (2009). Ceramide and neurodegeneration: Susceptibility of neurons

and oligodendrocytes to cell damage and death. J. Neurol. Sci. *278*, 5-15.

Jatana, M., Giri, S., and Singh, A.K. (2002). Apoptotic positive cells in Krabbe brain and induction of

apoptosis in rat C6 glial cells by psychosine. Neurosci Lett *330*, 183–187.

Kobolák, J., Molnár, K., Varga, E., Bock, I., Jezsó, B., Téglási, A., Zhou, S., Lo Giudice, M., Hoogeveen-

Westerveld, M., Pijnappel, W.P., et al. (2019). Modelling the neuropathology of lysosomal storage

disorders through disease-specific human induced pluripotent stem cells. Exp. Cell Res. *380*, 216-233.

Lattanzi, A., Neri, M., Maderna, C., di Girolamo, I., Martino, S., Orlacchio, A., Amendola, M., Naldini, L.,

and Gritti, A. (2010). Widespread enzymatic correction of CNS tissues by a single intracerebral

injection of therapeutic lentiviral vector in leukodystrophy mouse models. Hum. Mol. Genet. *19*,

2208–2227.

Lattanzi, A., Salvagno, C., Maderna, C., Benedicenti, F., Morena, F., Kulik, W., Naldini, L., Montini, E.,

Martino, S., and Gritti, A. (2014). Therapeutic benefit of lentiviral-mediated neonatal intracerebral

gene therapy in a mouse model of globoid cell leukodystrophy. Hum. Mol. Genet. *23*, 3250–3268.

Lee, S.J., Yi, T., Ahn, S.H., Lim, D.K., Hong, J.Y., Cho, Y.K., Lim, J., Song, S.U., and Kwon, S.W. (2014).

Senescing human bone-marrow-derived clonal mesenchymal stem cells have altered

lysophospholipid composition and functionality. J. Proteome Res. *13*, 1438-1449.

Lee, W.C., Kang, D., Causevic, E., Herdt, A.R., Eckman, E.A., and Eckman, C.B. (2010). Molecular

characterization of mutations that cause globoid cell leukodystrophy and pharmacological rescue

using small molecule chemical chaperones. J Neurosci *30*, 5489–5497.

Lemonnier, T., Blanchard, S., Toli, D., Roy, E., Bigou, S., Froissart, R., Rouvet, I., Vitry, S., Heard, J.M., and

Bohl, D. (2011). Modeling neuronal defects associated with a lysosomal disorder using patient-

derived induced pluripotent stem cells. Hum. Mol. Genet. *20*, 3653–3666.

Lin, D., Fantz, C.R., Levy, B., Rafi, M.A., Vogler, C., Wenger, D.A., and Sands, M.S. (2005). AAV2/5 vector

expressing galactocerebrosidase ameliorates CNS disease in the murine model of globoid-cell

leukodystrophy more efficiently than AAV2. Mol Ther *12*, 422–430.

Liu, J., Zhang, H., Li, Z., Hailemariam, T.K., Chakraborty, M., Jiang, K., Qiu, D., Bui, H.H., Peake, D.A., Kuo,

M.S., et al. (2009). Sphingomyelin synthase 2 is one of the determinants for plasma and liver

sphingomyelin levels in mice. Arterioscler. Thromb. Vasc. Biol. *29*, 850-856.

Liu, Y., Gibson, J., Wheeler, J., Kwee, L.C., Santiago-Turla, C.M., Akafo, S.K., Lichter, P.R., Gaasterland,

D.E., Moroi, S.E., Challa, P., et al. (2011). GALC deletions increase the risk of primary Open-Angle

Glaucoma: The role of mendelian variants in complex disease. PLoS One. *6*, e27134.

Lizardo, D.Y., Lin, Y.L., Gokcumen, O., and Atilla-Gokcumen, G.E. (2017). Regulation of lipids is central to

replicative senescence. Mol. Biosyst. *13*, 498-509.

Lizardo, D.Y., Parisi, L.R., Li, N., and Atilla-Gokcumen, G.E. (2018). Noncanonical Roles of Lipids in

Different Cellular Fates. Biochemistry. *57*, 22-29.

Lojewski, X., Staropoli, J.F., Biswas-Legrand, S., Simas, A.M., Haliw, L., Selig, M.K., Coppel, S.H., Goss, K.A.,

Petcherski, A., Chandrachud, U., et al. (2014). Human iPSC models of neuronal ceroid lipofuscinosis

capture distinct effects of TPP1 and CLN3 mutations on the endocytic pathway. Hum. Mol. Genet. *23*,

2005–2022.

Luni, C., Gagliano, O., and Elvassore, N. (2016). Microfluidic technology enhances the potential of human

pluripotent stem cells. Biochem. Biophys. Res. Commun. *473*, 10–14.

Luzi, P., Rafi, M.A., and Wenger, D.A. (1995). Characterization of the large deletion in the GALC gene

found in patients with Krabbe disease. Hum. Mol. Genet. *4*, 2335–2338.

Luzi, P., Rafi, M.A., Victoria, T., Baskin, G.B., and Wenger, D.A. (1997). Characterization of the rhesus

monkey galactocerebrosidase (GALC) cDNA and gene and identification of the mutation causing
globoid cell leukodystrophy (Krabbe disease) in this primate. Genomics *42*, 319–324.

Maetzel, D., Sarkar, S., Wang, H., Abi-Mosleh, L., Xu, P., Cheng, A.W., Gao, Q., Mitalipova, M., and
Jaenisch, R. (2014). Genetic and chemical correction of cholesterol accumulation and impaired
autophagy in hepatic and neural cells derived from niemann-pick type C patient-specific iPS cells.
Stem Cell Reports *2*, 866–880.

Martino, S., Tiribuzi, R., Tortori, A., Conti, D., Visigalli, I., Lattanzi, A., Biffi, A., Gritti, A., and Orlacchio, A.
(2009). Specific determination of beta-galactocerebrosidase activity via competitive inhibition of
beta-galactosidase. Clin. Chem. *55*, 541–548.

Meneghini, V., Lattanzi, A., Tiradani, L., Bravo, G., Morena, F., Sanvito, F., Calabria, A., Bringas, J., Fisher-
Perkins, J.M., Dufour, J.P., et al. (2016). Pervasive supply of therapeutic lysosomal enzymes in the CNS
of normal and Krabbe-affected non-human primates by intracerebral lentiviral gene therapy. EMBO
Mol. Med. *8*, 489–510.

Meneghini, V., Frati, G., Sala, D., De Cicco, S., Luciani, M., Cavazzin, C., Paulis, M., Mentzen, W., Morena,
F., Giannelli, S., et al. (2017). Generation of Human Induced Pluripotent Stem Cell-Derived Bona Fide
Neural Stem Cells for Ex Vivo Gene Therapy of Metachromatic Leukodystrophy. Stem Cells Transl.
Med. *6*, 352–368.

Misslin, C., Velasco-Estevez, M., Albert, M., O'Sullivan, S.A., and Dev, K.K. (2017). Phospholipase A2 is
involved in galactosylsphingosine-induced astrocyte toxicity, neuronal damage and demyelination.
PLoS One. *12*, e0187217.

Mullen, T.D., Hannun, Y.A., Obeid, L.M., Hannun, Y.A., Obeid, L.M., Gatt, S., Gatt, S., Yavin, E., Gatt, S.,
Sribney, M., et al. (2012). Ceramide synthases at the centre of sphingolipid metabolism and biology.
Biochem. J. *441*, 139–150.

Nicaise, A.M., Wagstaff, L.J., Willis, C.M., Paisie, C., Chandok, H., Robson, P., Fossati, V., Williams, A., and
Crocker, S.J. (2019). Cellular senescence in progenitor cells contributes to diminished remyelination
potential in progressive multiple sclerosis. Proc. Natl. Acad. Sci. U. S. A. *116*, 9030-9039.

Ordoñez, M.P., and Steele, J.W. (2016). Modeling Niemann Pick type C1 using human embryonic and
induced pluripotent stem cells. Brain Res. *1656*, 63-67.

Panicker, L.M., Miller, D., Park, T.S., Patel, B., Azevedo, J.L., Awad, O., Masood, M.A., Veenstra, T.D.,
Goldin, E., Stubblefield, B.K., et al. (2012). Induced pluripotent stem cell model recapitulates
pathologic hallmarks of Gaucher disease. Proc. Natl. Acad. Sci. U. S. A. *109*, 18054–18059.

Pellegrini, D., Del Grosso, A., Angella, L., Giordano, N., Dilillo, M., Tonazzini, I., Caleo, M., Cecchini, M.,

and McDonnell, L.A. (2019). Quantitative microproteomics based characterization of the central and

peripheral nervous system of a mouse model of Krabbe disease. Mol. Cell. Proteomics. *8*, 1227-1241.

Potter, G.B., and Petryniak, M.A. (2016). Neuroimmune mechanisms in Krabbe's disease. J. Neurosci.

Res. *94*, 1341-1348.

Potter, G.B., Santos, M., Davisson, M.T., Rowitch, D.H., Marks, D.L., Bongarzone, E.R., and Petryniak, M.A.

(2013). Missense mutation in mouse GALC mimics human gene defect and offers new insights into

Krabbe disease. Hum Mol Genet *22*, 3397–3414.

Rafi, M.A., Luzi, P., Chen, Y.Q., and Wenger, D.A. (1995). A large deletion together with a point mutation

in the GALC gene is a common mutant allele in patients with infantile krabbe disease. Hum. Mol.

Genet. *4*, 1285–1289.

Rafi, M.A., Fugaro, J., Amini, S., Luzi, P., de Gala, G., Victoria, T., Dubell, C., Shahinfar, M., and Wenger,

D.A. (1996). Retroviral vector-mediated transfer of the galactocerebrosidase (GALC) cDNA leads to

overexpression and transfer of GALC activity to neighboring cells. Biochem Mol Med *58*, 142–150.

Rafi, M.A., Rao, H.Z., Luzi, P., and Wenger, D.A. (2015). Long-term Improvements in Lifespan and

Pathology in CNS and PNS After BMT Plus One Intravenous Injection of AAVrh10-GALC in Twitcher

Mice. Mol. Ther. *23*, 1681–1690.

Ribbens, J., Whiteley, G., Furuya, H., Southall, N., Hu, X., Marugan, J., Ferrer, M., and Maegawa, G.H.B.

(2013). A high-throughput screening assay using Krabbe disease patient cells. Anal. Biochem. *434*, 15–

25.

Ricca, A., and Gritti, A. (2016). Perspective on innovative therapies for globoid cell leukodystrophy. J.

Neurosci. Res. *94*, 1304–1317.

Ricca, A., Rufo, N., Ungari, S., Morena, F., Martino, S., Kulik, W., Alberizzi, V., Bolino, A., Bianchi, F., Del

Carro, U., et al. (2015). Combined gene/cell therapies provide long-term and pervasive rescue of

multiple pathological symptoms in a murine model of globoid cell leukodystrophy. Hum. Mol. Genet.

24, 3372–3389.

Samie, M.A., and Xu, H. (2014). Lysosomal exocytosis and lipid storage disorders. J. Lipid Res. *55*, 995–

1009.

Santambrogio, S., Ricca, A., Maderna, C., Ieraci, A., Aureli, M., Sonnino, S., Kulik, W., Aimar, P., Bonfanti,

L., Martino, S., et al. (2012). The galactocerebrosidase enzyme contributes to maintain a functional

neurogenic niche during early post-natal CNS development. Hum. Mol. Genet. *21*, 4732–4750.

Sessa, M., Lorioli, L., Fumagalli, F., Acquati, S., Redaelli, D., Baldoli, C., Canale, S., Lopez, I.D., Morena, F.,

Calabria, A., et al. (2016). Lentiviral haemopoietic stem-cell gene therapy in early-onset

metachromatic leukodystrophy: an ad-hoc analysis of a non-randomised, open-label, phase 1/2 trial.

Lancet *388*, 476–487.

Shi, Y., Inoue, H., Wu, J.C., and Yamanaka, S. (2017). Induced pluripotent stem cell technology: a decade

of progress. Nat. Rev. Drug Discov. *16*, 115–130.

Shin, D., Feltri, M.L., and Wrabetz, L. (2016). Altered Trafficking and Processing of GALC Mutants

Correlates with Globoid Cell Leukodystrophy Severity. J. Neurosci. *36*, 1858–1870.

Smith, B.R., Santos, M.B., Marshall, M.S., Cantuti-Castelvetri, L., Lopez-Rosas, A., Li, G., Van Breemen, R.,

Claycomb, K.I., Gallea, J.I., Celej, M.S., et al. (2014). Neuronal inclusions of α-synuclein contribute to

the pathogenesis of Krabbe disease. J. Pathol. *232*, 509–521.

Son, M.-Y., Kwak, J.E., Seol, B., Lee, D.Y., Jeon, H., and Cho, Y.S. (2015). A novel human model of the

neurodegenerative disease GM1 gangliosidosis using induced pluripotent stem cells demonstrates

inflammasome activation. J. Pathol. *237*, 98-110.

Spratley, S.J., Hill, C.H., Viuff, A.H., Edgar, J.R., Skjødt, K., and Deane, J.E. (2016). Molecular Mechanisms

of Disease Pathogenesis Differ in Krabbe Disease Variants. Traffic *17*, 908–922.

Sun, Y., Florer, J., Mayhew, C.N., Jia, Z., Zhao, Z., Xu, K., Ran, H., Liou, B., Zhang, W., Setchell, K.D.R., et al.

(2015). Properties of neurons derived from induced pluripotent stem cells of Gaucher disease type 2

patient fibroblasts: Potential role in neuropathology. PLoS One *10*.

Suzuki, K. (1983). The twitcher mouse. A model of human globoid cell leukodystrophy (krabbe's disease).

Am J Pathol *111*, 394–397.

Suzuki, K., and Suzuki, Y. (1970). Globoid cell leucodystrophy (Krabbe's disease): deficiency of

galactocerebroside beta-galactosidase. Proc Natl Acad Sci U S A *66*, 302–309.

Tappino, B., Biancheri, R., Mort, M., Regis, S., Corsolini, F., Rossi, A., Stroppiano, M., Lualdi, S., Fiumara,

A., Bembi, B., et al. (2010). Identification and characterization of 15 novel GALC gene mutations

causing Krabbe disease. Hum. Mutat. *31*, E1894-1914.

Teixeira, C.A., Miranda, C.O., Sousa, V.F., Santos, T.E., Malheiro, A.R., Solomon, M., Maegawa, G.H.,

Brites, P., and Sousa, M.M. (2014). Early axonal loss accompanied by impaired endocytosis, abnormal

axonal transport, and decreased microtubule stability occur in the model of Krabbe's disease.

Neurobiol. Dis. *66*, 92–103.

Tiscornia, G., Lorenzo Vivas, E., Matalonga, L., Berniakovich, I., Barragán Monasterio, M., Eguizábal, C.,

Gort, L., González, F., Ortiz mellet, C., García Fernández, J.M., et al. (2013). Neuronopathic Gaucher's

disease: Induced pluripotent stem cells for disease modelling and testing chaperone activity of small

compounds. Hum. Mol. Genet. *22*, 633–645.

Trayssac, M., Hannun, Y.A., and Obeid, L.M. (2018). Role of sphingolipids in senescence: Implication in

aging and age-related diseases. J. Clin. Invest. *128*, 2702-2712.

Trilck, M., Hübner, R., Seibler, P., Klein, C., Rolfs, A., and Frech, M.J. (2013). Niemann-Pick type C1

patient-specific induced pluripotent stem cells display disease specific hallmarks. Orphanet J. Rare

Dis. *8*, 144.

Ungari, S., Montepeloso, A., Morena, F., Cocchiarella, F., Recchia, A., Martino, S., Gentner, B., Naldini, L.,

and Biffi, A. (2015). Design of a regulated lentiviral vector for hematopoietic stem cell gene therapy of

globoid cell leukodystrophy. Mol. Ther. Methods Clin. Dev. *2*, 15038.

Vallejo-Diez, S., Fleischer, A., Martín-Fernández, J.M., Sánchez-Gilabert, A., and Bachiller, D. (2018).

Generation of two induced pluripotent stem cells lines from a Mucopolysaccharydosis IIIB (MPSIIIB)

patient. Stem Cell Res. *33*, 180-184.

Venable, M.E., Lee, J.Y., Smyth, M.J., Bielawska, A., and Obeid, L.M. (1995). Role of ceramide in cellular

senescence. J. Biol. Chem. *270*, 30701-30708.

Wenger, D.A. (2000). Murine, canine and non-human primate models of Krabbe disease. Mol. Med.

Today *6*, 449–451.

Wenger, D.A., and Luzi, P. (2014). Krabbe Disease: Globoid Cell Leukodystrophy. In Rosenberg's

Molecular and Genetic Basis of Neurological and Psychiatric Disease: Fifth Edition, pp. 337–346.

White, A.B., Givogri, M.I., Lopez-Rosas, A., Cao, H., van Breemen, R., Thinakaran, G., and Bongarzone,

E.R. (2009). Psychosine accumulates in membrane microdomains in the brain of krabbe patients,

disrupting the raft architecture. J Neurosci *29*, 6068–6077.

White, A.B., Galbiati, F., Givogri, M.I., Lopez Rosas, A., Qiu, X., Van Breemen, R., and Bongarzone, E.R.

(2011). Persistence of psychosine in brain lipid rafts is a limiting factor in the therapeutic recovery of a

mouse model for Krabbe disease. J. Neurosci. Res. *89*, 352–364.

Yamanaka, S. (2007). Strategies and new developments in the generation of patient-specific pluripotent

stem cells. Cell Stem Cell *1*, 39–49.

Young, M.M., and Wang, H.G. (2018). Sphingolipids as Regulators of Autophagy and Endocytic

Trafficking. Adv. Cancer Res. *140*, 27-60.

Young, S.A., Mina, J.G., Denny, P.W., and Smith, T.K. (2012). Sphingolipid and ceramide homeostasis:

Potential therapeutic targets. Biochem. Res. Int. *2012*, e248135.

Yu, D., Swaroop, M., Wang, M., Baxa, U., Yang, R., Yan, Y., Coksaygan, T., DeTolla, L., Marugan, J.J.,

Austin, C.P., et al. (2014). Niemann-Pick Disease Type C: Induced Pluripotent Stem Cell-Derived

Neuronal Cells for Modeling Neural Disease and Evaluating Drug Efficacy. J. Biomol. Screen. *19*, 1164–

1173.

Zhang, P., Kishimoto, Y., Grammatikakis, I., Gottimukkala, K., Cutler, R.G., Zhang, S., Abdelmohsen, K.,

Bohr, V.A., Misra Sen, J., Gorospe, M., et al. (2019). Senolytic therapy alleviates Aβ-associated

oligodendrocyte progenitor cell senescence and cognitive deficits in an Alzheimer's disease model.

Nat. Neurosci. *22*, 719-728.

Figure 3

Figure 6

Figure 7