A metabolic life table exemplified by sockeye salmon

Joseph R. Burger[a,b,1*], Chen Hou[c,1], Charles A.S Hall[d,1], and James H. Brown[e,1]

[a] Arizona Institutes for Resilience, University of Arizona, Tucson, AZ 85721 USA;

Corresponding author: ecoevoburger@email.arizona.edu

[b] Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721 USA

[c] Department of Biological Science, Missouri University of Science and Technology, Rolla, MO

65409 USA;

[d] Department of Environmental and Forest Biology and Program in Environmental Science, State

University of New York, – College of Environmental Science and Forestry, Syracuse NY, 13210,

USA;

[e] Department of Biology, University of New Mexico, Albuquerque, NM 87131 USA

[1]All authors contributed equally

Key words: life history; demography; energy budget; ecosystem; conservation; sustainable

harvest; steady-state; metabolic theory of ecology; population

Abstract

1) A metabolic life table (MLT) is a combination of energy budget and life table that quantifies

metabolism and life history over an entire life cycle. It provides a conceptual framework for

integrating data on physiology, demography and ecology that are usually the subject of

discipline- and taxon-specific studies.

2) Our MLT for sockeye salmon revives John Brett's classic data on metabolism, growth, survival

and reproduction to provide a synthetic analysis of metabolic performance and its life history

consequences.

3) The MLT quantifies the energy budget of an average female over her life cycle. The early

stages in fresh water have low rates of growth and mortality. Then juveniles enter the ocean,

feed voraciously, grow rapidly, and accumulate a store of biomass; More than 98% of the total

lifetime assimilation and growth occurs in the ocean. Maturing adults stop feeding, return to

fresh water, expend stored body energy to fuel migration and reproduction, and leave a clutch

of eggs and a depleted carcass.

4) The MLT also quantifies the energetic contribution of salmon to freshwater and marine

ecosystems. Salmon are very efficient: of the food energy in their zooplankton prey, 47% is

expended on respiration to fuel activity, 23% is allocated to growth and reproduction to

produce biomass, and 30% is excreted in feces. Although 96% of lifetime biomass production

occurs in the ocean, about 29% is transported into fresh water in the bodies of maturing adults,

where their carcasses and gametes provide an important "marine subsidy" to the energy and

nutrient budgets of freshwater and terrestrial ecosystems.

5) The MLT highlights some features of the salmon metabolism – variation in rates of

assimilation, respiration and production with body size and temperature – that are qualitatively

similar to other ectotherms and predictions of metabolic theory. But because of the unusual

physiology, life history, ecosystem impacts and socio-economic importance of wild-caught

salmon, reanalyzing Brett's data in the context of a MLT has additional broad applications for

basic and applied ecology.

Introduction

A challenge of contemporary basic and applied ecology is to synthesize the detailed

information available in specialized databases and journals to elucidate general patterns and

processes. Empirical and theoretical metabolic ecology use energy as a common currency to

make explicit linkages among life history, ecology and evolution (Brown, Gillooly, Allen, Savage,

& West, 2004; Burger, Hou, & Brown, 2019; Hall, 1972; Kooijman & Kooijman, 2010; Lotka,

1922; Odum & Pinkerton, 1955; Sibly, Brown, & Kodric-Brown, 2012; Van Valen, 1976). These

connect anatomy, physiology and behavior at the level of individual organisms to ecological

patterns and processes at population, community and ecosystem levels. They also have

important applications to conservation and management. Many studies apply short-term

measurements of resting, active and field metabolism of particular species to address questions

about the ecological costs and benefits and the evolutionary consequences of individual

organism-level traits and ecosystem-level phenomena (e.g., Peters 1983; Brown et al. 2004;

Sibly et al. 2012). Largely missing, however, are comprehensive studies of single species that

can address of how metabolic energy is acquired and expended over an entire life cycle and

interpret them in ecological and evolutionary contexts. Pacific salmon (*Oncorhynchus spp.*)

provide an excellent model organism for two reasons: their semelparous life history simplifies

quantification, obviating the need for detailed data and analyses to address ideterminate

growth and semelparous reproduction (Fig. 1); and they have been studied intensively – long

prized as food for humans, wild salmon fisheries are worth billions of dollars – so there are

relevant physiological and demographic data for the entire life cycle (Knapp, Guettabi, &

Goldsmith, 2013; Quinn, 2018).

Canadian fisheries biologist John "Roly" Brett introduced a "life table of energetics" to

synthesize his life's work on the physiological ecology of the sockeye salmon (*Oncorhynchus*

nerka) that breed in Babine Lake, British Columbia (Brett, 1986; Brett, 1983). His goal was to

"put together a comprehensive estimate of the life energetics of an average sockeye salmon

that commences life as a fertilized egg weighing 0.13 g containing 372 cal … and terminates

growth as a 2,270 g adult containing 4,200 kcal when entering fresh water on its final journey"

(Brett, 1983), p.30). Brett adopted the form of a standard life table to present his data on body

size, energy content, growth and metabolism as a function of age over an entire life cycle (Table

1). Here we resurrect and extend Brett's concept of a "life table of energetics" to construct a

metabolic life table (MLT) that combines the demography of a traditional life table with the

mass-energy balance of physiological ecology. By quantifying energy allocation to components

of fitness – survival, growth and reproduction – as a function of age, the MLT highlights the dual

roles of universal constraints and special adaptations in metabolic ecology and life history of

salmon. We demonstrate the practical utility and broad applicability of the MLT approach, and

show how it goes beyond traditional demography and short-term studies of metabolism to

provide a synthetic picture of the central role of energy in physiology, behavior, ecology and

evolution.

Methods

Brett's "Life Table of Energetics" (Brett, 1983), which gives data on body mass and

composition, and energy acquisition and expenditure for 33 stages of a typical four-year life

cycle is distilled and modified in Table 1. Brett made some simplifications and assumptions:

1) He assumed a steady state population with reproduction balancing mortality. It is well known

that there is considerable variation in the size and composition of the Skeena River runs among

years (e.g., Larkin and McDonald 1968). However, in the 1970s and early 1980s the Babine Lake

sockeye population and fishery was stable, so we followed Brett (1983, 1986) in assuming

steady state.

2) He assumed a simplified four-year life cycle, ignoring the small proportion of the population

returning to breed after three or five years.

3) He estimated average values for the population as a whole, ignoring considerable variation

among individuals and across years.

4) He used laboratory measurements, simplified models, and corrections for environmental

temperature to estimate energy assimilation from feeding and energy expenditure for

maintenance and activity.

5) He used additional information from models and other salmon populations to estimate

parameters (e.g., growth and mortality) for the marine stages for which direct measurements

for the Babine Lake population were not available.

6) He recognized that many parameters were estimated imprecisely, so he rounded off and

reported average values.

We added mortality data from (Brett, 1986) to estimate the age structure for the 33

stages in Fig. 2. Using these values — and ignoring seasonal and interannual variation due to

movement at sea, seasonal fluctuations in ocean temperature and productivity, and differences

among runs — we compiled a MLT for the Babine Lake sockeye population (Table 1). Given the

above assumptions, simplifications, and conversions, our MLT is a faithful distillation and

synthesis of Brett's data.

Metabolic ecology of sockeye salmon

Natural history of the life cycle

Salmon are: 1) anadromous – breeding in fresh water and growing in the ocean, and 2)

semelparous – making a single, end-of-life reproductive effort. The life cycle (Fig. 1) starts in

September when the mature adults have returned to spawn in tributary streams of Babine

Lake. The average female lays 3,000 eggs, each weighing about 0.1 g. Males fertilize the eggs as

females deposit them in the streambed. The eggs are buried in gravel and develop over the

winter. In June ~600 survivors hatch, move to the lake and feed on plankton. After one year

about 84 surviving smolts, each weighing ~5.4 g, migrate downstream and enter the ocean.

During the > two years in marine phase, juveniles move between water masses of varying

temperature and productivity, feed voraciously on zooplankton, grow rapidly, and accumulate

stores of energy-rich lipids and proteins in their distinctive red flesh. In the summer of their

fourth year, the maturing adults weigh ~2700 g when they return to coastal waters, stop

feeding, and start their upstream migration. By September, they reach the spawning grounds,

where females compete to excavate nests in the streambed and males compete to mate with

spawning females. Both sexes die after spawning, leaving a clutch of eggs and depleted

carcasses, having expended most of their body energy for migration and mating.

Life table: survival and reproduction

The first few columns of Table 1 are a life table: a schedule of survival and reproduction

as a function of age, stage and body mass. Each generation starts with 3,000 eggs, lasts four

years, and ends with another 3,000 eggs spawned by two depleted breeders. In between the

eggs hatch, the offspring grow and die as they age, and an average of two survive to breed and

continue the life cycle. The dynamics of the life history as a function of age are shown in Fig. 2.

Until the salmon stop feeding and enter fresh water for the return migration, cumulative

assimilation, respiration and biomass increase with age; deviations from smooth monotonic

curves reflect variations in energy income (assimilation from feeding), expenditure (respiration

for swimming), and associated variation in temperature (Brett, 1972; Brett & Glass, 1973; Brett,

1983). During the last three months of life, the returning adults use most of their stored body

energy on locomotor and reproductive activity, leaving depleted carcasses and a new clutch of

eggs.

Metabolism: body mass and temperature dependence of respiration, growth and assimilation.—

The MLT (Table 1) documents variation in rates of respiration, growth and assimilation with

age, stage, body mass and environmental temperature at monthly intervals over one

generation. We compare these data to predictions from general metabolic theory (Brown et al.,

2004; Calder, 1984; Peters, 1986; Schmidt-Nielsen, 1984; Sibly et al., 2012). Whole-organism

rates, B, characteristically increase with body mass, m, scaling as power laws of the form

$B = B_0 m^\alpha$ 1

where B_O is a normalization constant, and the exponent α is characteristically close to 3/4 for

respiration rate and most other whole-organism rates (Brown et al., 2004; Calder, 1984;

Kleiber, 1932; Peters, 1986; Schmidt-Nielsen, 1984; Sibly et al., 2012; West, Brown, & Enquist,

1997). In ectothermic animals, such as salmon, these rates also vary with body temperature

with $Q_{10} \approx 2.5$ (increasing ~2.5 times for every 10°C increase in temperature (Gillooly, 2001;

Huey & Kingsolver, 2019).

Brett's data for sockeye salmon are qualitatively consistent with predictions from

metabolic scaling theory, but with quantitative caveats. Brett (1983, p. 33) determined that the

temperature dependence was "equivalent to a Q_{i0} of about 2.3." We used this value to

temperature-correct the rates of respiration, growth and assimilation, and analyze their

scalings with body mass (Fig. 3). The scaling relationships are positive as has been found

universally, but the exponents (slopes) are consistently higher than $\alpha \approx 0.75$ expected from

general standard metabolic theory (Brown et al., 2004) and data for most other animals (e.g.,

"Kleiber's rule": (Calder, 1984; Kleiber, 1932; Peters, 1986; Schmidt-Nielsen, 1984; Sibly et al.,

2012). We are uncertain how to interpret these deviations. On the one hand, salmon have an

unusual physiology and ecology as indicated above. On the other hand, the data might be

inaccurate. Brett (Brett, 1983), p. 32) cautions "More often than not when compiling energy

budgets growth is measured, metabolism estimated, assimilation efficiency assumed, and

excretion deduced. …The components of the balanced equation are sensitive to, and respond

differently to, the chief variables of temperature, body size, and activity." Since we have no

objective basis to question or correct Brett's values, we use them here with this caveat.

Energy balance of an individual.—An energy budget for the average individual surviving to

breed accounts for stocks and flows of energy over one generation (Fig. 4):

1) egg: The energy budget starts with a single fertilized egg, weighing 0.10 g and containing 1.25

kJ of energy. The egg lies inactive in the gravel, transforms from an embryo into a fry, and

hatches after about nine months, having lost about one-third of its mass and energy in

respiration.

2) feeding, maintenance and growth: In the three plus years between hatching and breeding,

the offspring assimilates 47,124 kJ of energy by feeding, expends 29,498 kJ (63%) on aerobic

respiration, and produces 17,626 kJ (37%) in body energy reserves. More than 98% of the total

lifetime assimilation and growth occurs in the ocean, where the juveniles swim and feed almost

continuously.

3) migration and breeding: Once a mature salmon enters fresh water, it stops feeding and lives

off body reserves. In the last 45 days of life, an individual expends 9,583

kJ (54%) of its body energy on respiration for expensive migration and spawning.

4) parental investment: A fraction of stored body energy is allocated to parental investment in

gametes. The clutch of 3,000 eggs contains 3,880 kJ (22% and 8.3% of lifetime biomass

production and assimilation, respectively). The male invests less energy, 938 kJ, in sperm.

5) carcass: When the spawned-out adult dies, its weight has decreased from 2,270 to 1,810

g. The carcass, depleted of nearly all fat and much of its protein, contains mostly water. Of the

17,626 kJ stored in the body at the start of migration, 13,463 kJ has been expended, 9,583 kJ on

respiration, 4,163 kJ in the depleted carcass and 3,880 kJ on eggs.

6) production efficiency: The efficiency of individual production, T_{ind}, is the ratio of output over

input or production over assimilation:

$T_{ind} = P_{ind}/A_{ind}$ (2)

It varies with water temperature and age (Table 1), decreasing from >60% for fry and

smolts in fresh water to <25% for the older stages in ocean water. By the time a female stops

feeding and enters fresh water for the return migration, she has accumulated 37% of

assimilated energy in her body as growth. When she dies three months later, after expending

more than half of this store on respiration for migration and spawning, her clutch of eggs and

depleted body contain just 18% of the energy she assimilated over her lifetime.

Energy balance of a cohort.—At least as relevant for life history and ecology is the energy

budget for the population of a representative cohort (Fig. 4B). This accounting includes intake

and expenditure of all offspring produced by a pair of breeders, including those that die before

reproducing. Brett (Brett, 1986) called these losses the "life-cycle deficit" to estimate the "food

conversion efficiency" of the Babine Lake population.

Since the population was approximately constant during Brett's study, we assumed

steady state and used the MLT (Table 1) to compile a balanced energetic budget for a cohort

(Fig. 4B, Table 2):

1) juveniles in fresh water: The energy budget for the cohort starts with 3,000 fertilized eggs

containing 3,880 kJ of energy. By the time the survivors have hatched, fed, grown, migrated

downstream, and transformed into smolts, most of the initial 3,000 offspring have died and left

20,800 kJ of assimilated energy in the ecosystem: 2,916 carcasses containing 6,869 kJ and

13,931 of respired heat energy.

2) juveniles in the ocean: The 84 surviving smolts have assimilated 7,423 kJ of food, expended

5,239 kJ on respiration, and amassed 2,184 kJ of body energy that they take with them to sea.

There they feed and swim almost continuously, assimilating 505,508 kJ and expending 343,286

kJ on respiration. An additional 80 juveniles die in the ocean, leaving 388,016 kJ of assimilated

energy in the ecosystem: 162,226 of biomass in their carcasses and 225,790 kJ of respired heat

energy.

3) migration and breeding: After two years at sea, body energy has increased about 30 times.

The four survivors contain 70,506 kJ of biomass energy when they enter fresh water to start the

return migration. Essentially all of this energy is dissipated in fresh water: 54,408 as respiration,

11,280 kJ in the four carcasses, 3,880 kJ in the new clutch of 3,000 eggs, and 938 kJ in sperm.

4) production and assimilation: We estimate the production and assimilation of the cohort,

P_{cohort} and A_{cohort}, of all offspring of an average female up until a given stage, S_{ref}, as

$P_{cohort} = \sum_{S=0}^{S=S_{ref}} N_d E_d$

and

$A_{cohort} = \sum_{S=0}^{S=S_{ref}} N_d A_d$ (3)

and where N_d is the number of offspring dying before stage, S, and E_d and N_d are respectively,

total body energy content (production) and cumulative assimilation (production plus

respiration) of those offspring when they died. Production efficiency of juveniles is high (32% of

assimilation in fresh water and 46% in the ocean), but at the end of the life cycle only 5% the

total energy assimilated by the cohort over one generation remains in fresh water.

Contribution of sockeye salmon to ecosystem energetics.—Table 2 uses the energy budget of

the cohort (Fig. 4B) and the MLT (Table 1) to quantify the impact of salmon on freshwater and

marine ecosystems. Of the food energy assimilated by juveniles in fresh water, 92% is

dissipated as heat of respiration and biomass of carcasses and 8% is transported into the ocean

by the newly transformed smolts. Of the energy assimilated by juveniles in the ocean, 88% is

dissipated in the marine ecosystem as respiration and mortality and 12% is transported into

streams and lakes in the bodies of the four returning adults. The majority of this stored body

energy (59%) is expended on respiration. The biomass in the carcasses, eggs and sperm is

essentially all consumed by predators and decomposers; it amounts to only 41% of the energy

brought in from the ocean and only 5% of the total energy assimilated by the cohort over one

generation.

The production efficiency of a cohort or species population, T_{cohort}, which Brett (1986)

called the "growth efficiency", is

$T_{cohort} = P/(CK)$ = P/A (4)

where P is the rate of biomass production, C is the rate of biomass consumption, and K is the

energy conversion efficiency for a consumer (e.g., salmon as predator). So T_{cohort} indexes the

efficiency of converting assimilated food eaten by salmon into salmon biomass that is

ultimately consumed by other organisms – predators, scavengers and decomposers – in the

freshwater and marine ecosystems. We calculated T_{cohort} (Table 1) using Brett's (1986)

estimate that K = 0.30 and assuming $C = \frac{P+R}{K} = \frac{A}{K}$. The trophic efficiency of juveniles is

somewhat lower in fresh water (0.22) than in the ocean (0.28). After going without feeding

after re-entering fresh water, the overall trophic efficiency over the entire life cycle is 0.23. For

Babine Lake sockeye population, this efficiency is 23%.

Discussion: broader implications

Physiology, ecology and evolution of salmon.—Oncorhynchus nerka is a an exemplar of how

natural selection has created a unique suite of physiological, behavioral and ecological traits

(Quinn, 2018). These special features beg for explanation in terms of phylogenetic history and

natural selection playing out in temporally and spatially varying environments. They are

typically addressed by different disciplines and presented in specialized publications. The MLT

provides a more synthetic explanation from the perspective of energetics. By combining the

demography of a traditional life table with the physiological ecology of a balanced energy

budget, the MLT applies first principles of physics and biology to integrate and synthesize many

aspects of population biology, physiological ecology, and ecosystem energetics. These topics

are typically addressed by different disciplines and presented in specialized publications. The

MLT for Babine Lake sockeye emphasizes how traits at different levels of organization

contribute to a unified big-picture perspective on salmon biology.

Energetics of migration.—Several features of the MLT reflect the central role of migration. Very

little energy is assimilated and produced by the fresh water stages, even though they account

for most of the individuals and about half of the four-year life cycle. Despite the large clutch of

eggs, because of small size and high mortality of the juvenile freshwater stages, they account

for only a small fraction of cohort assimilation and biomass. Nearly all of the biomass energy

produced over the life cycle (98% for a surviving breeder and 96% for an entire cohort) is

acquired during feeding and growth in the ocean. The last stages of the life cycle are fueled by

stored body energy, transported into fresh water and expended on the respiration for

migration and breeding. By the time they spawn, two returning adults have expended 74% of

their body energy, leaving only 19% in depleted carcasses and 7% in gametes.

Sockeye and kokanee.—Comparisons between closely related salmon populations provide

additional insights. As indicated by fossil history and shared adaptations for breeding in fresh

water, the ancestral salmonids lived in streams and lakes (Hoar, 1976). Multiple lineages

subsequently independently evolved life histories in which juveniles migrate to sea, grow to

maturity, and migrate back to fresh water to breed (McCormick & Saunders, 1987). A

consequence is a diversity of salmonid lifestyles, with closely related migratory and sedentary,

anadromous and "land-locked" populations.

Oncorhynchus nerka is a good example. Sea run sockeye are widely distributed, with

marine stages in the North Pacific and freshwater stages in surrounding lakes and streams.

Another morpho-ecotype, known as kokanee, spends its entire life cycle in fresh water, with

native populations in lakes scattered throughout the range and introduced populations more

broadly distributed (Quinn, 2018). Kokanee and sockeye represent alternative stable

evolutionary strategies: different combinations of traits that confer equal fitness. Sexually

mature landlocked kokanee of the same age are smaller and less fecund (~700 g and ~800 eggs)

than sea-run sockeye (~2,700 g and ~3,000 eggs) (USDA Forest Service, Lake Tahoe Basin

Management Unit, Tahoe Heritage Foundation, 2015). By migrating to sea and feeding on the

abundant plankton, anadromous populations were able to grow larger, produce more offspring,

and occupy a distinctively different ecological niche. Despite such initial advantages that

selected for the derived ocean-run ecomorphotype, however, kokanee and sockeye are now

equally fit, as evidenced by the fact in some locations the two populations coexist and breed

together but rarely hybridize. Compilation of a MLT for a population of kokanee would show

quantitatively how the life history and energetics evolved in successful adaptation to its

landlocked existence and reveal the tradeoffs resulting in equal fitness between kokanee and

anadromous salmon. MLTs could also be used to predict effects of climate change, food

restriction and other human activities on individual performance, cohort energetics, and

ecosystem impacts of salmon and other ectotherms (see Huey & Kingsolver, 2019)

Ecosystem energetics.—One consequence of the anadromous lifestyle is that migrating salmon

transport large quantities of energy and materials from marine to freshwater and surrounding

terrestrial ecosystems (Burger et al., 2012; Quinn, Helfield, Austin, Hovel, & Bunn, 2018;

Schindler, Leavitt, Brock, Johnson, & Quay, 2005). Inorganic compounds in the bodies of salmon

are major sources of nutrients for freshwater algae and land plants. A little more than half of

the organic compounds (lipids, carbohydrates and proteins) are catabolized in aerobic

respiration, generating ATP to fuel migration and reproduction, and releasing carbon dioxide,

water and heat into the environment. A smaller but substantial quantity of the biomass is

consumed by predators, scavengers and decomposers in freshwater and surrounding terrestrial

ecosystems.

The ecological impact of these inputs is substantial. Each returning migrant transports

only a small quantity of biomass, but the large body size and sheer numbers of migrants results

in a significant net effect – a "marine subsidy" of energy and nutrients to freshwater and

terrestrial ecosystems. When Brett did his studies in the late 1970s and early 1980s, the Skeena

River sockeye was in approximate steady state and the annual runs of returning adults

averaged about 3 million (DFO, 2003; Larkin & McDonald, 1968). Multiplying this by the 17,626

kJ of body energy per individual leaving the ocean gives about 53,000 megajoules of ocean

production (or 1,500 tonnes of organic carbon) transported to freshwater and terrestrial

ecosystems. For reference, this energy subsidy from a one-year run is equal to the annual net

primary production, ~25 gC/m^2, of approximately 6,000 ha of taiga forest (Gower et al., 2012).

In watersheds where the number of returning salmon have decreased due to human activities

(see below), the reduced marine subsidies of energy and nutrients have caused substantial

changes in freshwater and terrestrial ecosystems (Burger et al., 2012; Quinn et al., 2018;

Schindler et al., 2005). There is also a "freshwater subsidy" to marine ecosystems due to

juveniles (smolts) entering the ocean. Our MLT based on Brett's data suggests that this is only

about 3% of the marine subsidy (Naiman, Bilby, Schindler, & Helfield, 2002; Moore & Schindler,

2004; Scheuerell, Levin, Zabel, Williams, & Sanderson, 2005).

Use of wild salmon by humans.—Few wild animals have been as important to humans as

salmon. For thousands of years native fishers captured returning breeders in nets and traps at

the mouths of rivers, preserved their flesh by drying and smoking, and budgeted the stored

food to last through the lean months and years between good runs. Exploitation increased after

the arrival of Europeans. Sockeye populations in the U.S. and Canada have declined

precipitously due to overfishing and damming and pollution of rivers. Populations in Alaska and

Asia are still abundant, due in part to supplementation from hatcheries and climate change

(Ruggerone and Irvine 2018), although body sizes are declining (Oke et al., 2020). Ironically, one

of the seriously depleted stocks is the Skeena River population that Brett studied. Annual

returns fluctuated around 3 million until the late 1990s, then dipped to all-time lows. Now

commercial, recreational and Native American fishing has been severely restricted (DFO, 2003;

Hoekstra, 2017).

Theoretical implications.—Recent advances in metabolic ecology raise the prospect for a

general theory that uses energy as a fundamental currency to link tradeoffs in traits of

individual organisms to their ecological and evolutionary manifestations – to show how

anatomy, physiology and behavior affect energy budgets and in turn demography, population

dynamics, community organization, and natural selection underlying biodiversity.

Comparison with dynamic energy budgets.—MLTs are somewhat similar to the dynamic energy

budgets of Kooijman and collaborators (DEBs: e.g. Kooijman & Kooijman, 2010; Sousa,

Domingos, & Kooijman, 2008; Sousa, Domingos, Poggiale, & Kooijman, 2010). Both aim to

contribute to a general theory of biological metabolism, using data to test theoretical

predictions. Both have been applied to empirical model data on Pacific salmon (Nisbet, Jusup,

Klanjscek, & Pecquerie, 2012; Pecquerie, Johnson, Kooijman, & Nisbet, 2011). The primary

difference is that the DEB focuses on how underlying processes at molecular and cellular levels

contribute to metabolic homeostasis at the individual organism level, whereas the MLT focuses

on how metabolism at the whole-organism level affects demography of populations and

energetics of ecosystems. Consequently, DEB models are usually have more details and require

more parameter estimates, whereas MLTs incorporate just a few robust assumptions and

parameters.

The two frameworks offer alternative, but not mutually exclusive theories applying

biophysical laws to understand both species-specific and universal characteristics of living

things. Biophysical laws that apply to all species limit how they acquire and allocate energy and

constrain their life histories. The result is an interesting combination of universal and species-

specific traits. On the one hand, all species are subject to: 1) the demographic constraint that in

each generation at steady state parents are exactly replaced by offspring; 2) the energy balance

constraint that all biomass energy assimilated from the environment is expended on survival,

growth and reproduction; and 3) biological scaling laws that constrain for patterns of co-

variation among traits. On the other hand, the diversity of living things offers examples of

species with a wide variety of life histories: orders-of-magnitude variation in body size, growth

and mortality, number and size of offspring, and reproductive mode (e.g., asexual and sexual,

iteroparous and semelparous, degree of development at birth and subsequent parental care,

etc.). Some of the universality and diversity predicted by existing metabolic theory is supported

by empirical studies. Compilation of MLTs for species in diverse taxonomic and functional

groups and comparisons among them offer exciting opportunities to test and extend metabolic

theory. For example, an important component of energetic fitness is F, the fraction of cohort

biomass production lost to mortality and consumed by other organisms in the ecosystem

(Brown, Hall, & Sibly, 2018; Burger et al., 2019; Burger, Hou, Hall, & Brown, 2020; see below).

In principle, it is straightforward to construct a MLT for any species. Ideally, it would be

put together by a single investigator or group of collaborators with first-hand experience

working on diverse aspects of the biology. In practice, however, MLTs and the basic data

necessary to compile are not available for more than a handful of well-studied species. The

entries for a species in the increasingly available electronic databases suffer from the problems

that often come from multiple studies of different populations and localities that define and

measure the critical traits inconsistently. Our search of the literature for life tables and energy

or mass balance diagrams generated many citations, but most are restricted to only part of the

life cycle or lack critical data (e.g., many life tables do not give fecundity as a function of age).

Our best attempt at compiling MLTs based on body size for eight animals is in Burger et

al., (2020). While this is a good start, it is important to note that most of these contain

incomplete data and unverified assumptions. Comparison among species in this admittedly

small sample suggests that F, the fraction of cohort biomass production reflects the tradeoff

between number and size of offspring. Teleost fish that produce very large numbers of very

small eggs have low values of F (e.g., 0.051 and 0.013 for salmon and Pollock, respectively),

whereas mammals that supply parental care and rear a small number of relatively large

offspring have much higher values (e.g., F = 0.35 and 0.25, for chimpanzee and brown bat,

respectively) (Burger et al. 2020).

In conclusion, the MLT framework that Brett pioneered decades ago to synthesize

information on life history, demography, energetics and physiology of sockeye salmon should

be widely applicable to contemporary ecology. Most existing studies of metabolic ecology are

based on extrapolating short-term measurements of respiratory metabolism – made most

often in the laboratory (for basal, resting, active and maximal metabolic rates) and more rarely

in nature (field metabolic rate). Brett compiled data from such short-term studies to put

together a "life table of energetics" of sockeye salmon that serves as the foundation for the

present paper. The MLT presented here quantifies energy gain from food and loss to heat and

mortality, and energy allocation between respiration to generate ATP to fuel the costs of living

and production of biomass energy passed on to the next generation in the form of offspring

growth and parental gametes. So, it offers a synthetic picture of the central role of energy in

the physiology, behavior, ecology and evolution of this one biologically distinct and

economically important species. It provides an encouraging example of how the compilation of

field data for more species in a MLT framework can allow us to evaluate and extend empirical

and theoretical approaches to life history, ecology and evolution based in energetics (Brown et

al., 2018; Burger et al., 2019; Burger et al. 2020). Doing so will has additional applications for

sustainable resource management, conservation and predicting global change impacts on

biodiversity.

Acknowledgments: JRB acknowledges support from the Bridging Biodiversity and Conservation

Science program in the Arizona Institutes for Resilience at the University of Arizona.

Author contributions

All authors conceived the ideas and designed methodology; collected and analyzed the data;

and wrote the manuscript including contributed critically to the drafts and gave final approval

for publication.

References:

Brett, J. R. (1972). The metabolic demand for oxygen in fish, particularly salmonids, and a

comparison with other vertebrates. *Respiration Physiology*, *14*(1–2), 151–170.

Brett, J. R. (1986). Production energetics of a population of sockeye salmon, Oncorhynchus

nerka. *Canadian Journal of Zoology*, *64*(3), 555–564.

Brett, J. R., & Glass, N. R. (1973). Metabolic rates and critical swimming speeds of sockeye

salmon (Oncorhynchus nerka) in relation to size and temperature. *Journal of the*

Fisheries Board of Canada, *30*(3), 379–387.

Brett, J. R. (1983). Life energetics of sockeye salmon, Oncorhynchus nerka. *Behavioural*

Energetics: The Cost of Survival in Vertebrates, 29–63.

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic

theory of ecology. *Ecology*, *85*(7), 1771–1789.

Brown, J. H., Hall, C. A., & Sibly, R. M. (2018). Equal fitness paradigm explained by a trade-off

between generation time and energy production rate. *Nature Ecology & Evolution*, *2*(2),

262.

Burger, J. R., Allen, C. D., Brown, J. H., Burnside, W. R., Davidson, A. D., Fristoe, T. S., … Okie, J.

G. (2012). The macroecology of sustainability. *PLoS Biology*, *10*(6).

Burger, J. R., Hou, C., & Brown, J. H. (2019). Toward a metabolic theory of life history.

Proceedings of the National Academy of Sciences, *116*(52), 26653–26661.

Burger, J. R., Hou, C., Hall, C., & Brown, J. R. (2020). Universal rules of life: metabolic rates,

biological times and the equal fitness paradigm. *BioRxiv*.

Calder, W.A. (1984). *Size, function, and life history*. Cambridge, MA.: Havard University Press.

DFO. (2003). *Skeena River Sockeye Salmon (updated). DFO Can. Sci. Advis. Sec. Stock Stock

Status Report 2003/047*. Retrieved from https://waves-vagues.dfo-

mpo.gc.ca/Library/342179.pdf

Gillooly, J. F. (2001). Effects of Size and Temperature on Metabolic Rate. *Science*, *293*(5538),

2248–2251. doi: 10.1126/science.1061967

Gower, S. T., krankina, O., Olson, R. J., Apps, M., Linder, S., & Wang, C. (2012). NPP Boreal

Forest: Consistent Worldwide Site Estimates, 1965-1995, R1. *ORNL DAAC*.

Hall, C. A. (1972). Migration and metabolism in a temperate stream ecosystem. *Ecology*, *53*(4),

585–604.

Hoar, W. S. (1976). Smolt transformation: evolution, behavior, and physiology. *Journal of the*

Fisheries Board of Canada, *33*(5), 1233–1252.

Hoekstra, G. (2017, June 23). Skeena River sockeye returns forecast at all-time low, sports

fishing closed. *Vancouver Sun*. Retrieved from

https://vancouversun.com/business/local-business/skeena-river-sockeye-returns-

forecast-at-all-time-low-sports-fishing-closed/

https://en.wikipedia.org/wiki/Aquaculture_of_salmonids. (n.d.).

https://www.google.com/search?q=salmon+life+cycle&rlz=1C1CHFX_enUS555US555&source=l

nms&tbm=isch&sa=X&ved=0ahUKEwj7rZiC3PXWAhWBLmMKHRIiAM8Q_AUICigB&biw=

1067&bih=423#imgrc=WiztTrFdqAKvPM. (n.d.).

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the metabolic

meltdown of ectotherms. *The American Naturalist*, *194*(6), E140–E150.

Kleiber, M. (1932). Body size and metabolism. *Hilgardia*, *6*(11), 315–353.

Knapp, G., Guettabi, M., & Goldsmith, O. S. (2013). *The economic importance of the Bristol Bay*

salmon industry.

Kooijman, B., & Kooijman, S. (2010). *Dynamic energy budget theory for metabolic organisation*.

Cambridge university press.

Larkin, P. A., & McDonald, J. G. (1968). Factors in the population biology of the sockeye salmon

of the Skeena River. *The Journal of Animal Ecology*, 229–258.

Lotka, A. J. (1922). Contribution to the energetics of evolution. *Proceedings of the National*

Academy of Sciences of the United States of America, *8*(6), 147.

McCormick, S. D., & Saunders, R. L. (1987). Preparatory physiological adaptations for marine life

of salmonids: osmoregulation, growth, and metabolism. *Am. Fish. Soc. Symp*, *1*, 1–229.

Moore, J. W., & Schindler, D. E. (2004). Nutrient export from freshwater ecosystems by

anadromous sockeye salmon (Oncorhynchus nerka). *Canadian Journal of Fisheries and*

Aquatic Sciences, *61*(9), 1582–1589.

Naiman, R. J., Bilby, R. E., Schindler, D. E., & Helfield, J. M. (2002). Pacific salmon, nutrients, and

the dynamics of freshwater and riparian ecosystems. *Ecosystems*, *5*(4), 399–417.

Nisbet, R. M., Jusup, M., Klanjscek, T., & Pecquerie, L. (2012). Integrating dynamic energy

budget (DEB) theory with traditional bioenergetic models. *Journal of Experimental*

Biology, *215*(6), 892–902.

Odum, H. T., & Pinkerton, R. C. (1955). Time's speed regulator: the optimum efficiency for

maximum power output in physical and biological systems. *American Scientist*, *43*(2),

331–343.

Oke, K. B., Cunningham, C. J., Westley, P. A. H., Baskett, M. L., Carlson, S. M., Clark, J., … Kibele,

J. (2020). Recent declines in salmon body size impact ecosystems and fisheries. *Nature*

Communications, *11*(1), 1–13.

Pecquerie, L., Johnson, L. R., Kooijman, S. A., & Nisbet, R. M. (2011). Analyzing variations in life-

history traits of Pacific salmon in the context of Dynamic Energy Budget (DEB) theory.

Journal of Sea Research, *66*(4), 424–433.

Peters, R. H., (1986). *The ecological implications of body size* (Vol. 2). Cambridge University

Press.

Quinn, T. P. (2018). *The behavior and ecology of Pacific salmon and trout*. University of

Washington press.

Quinn, T. P., Helfield, J. M., Austin, C. S., Hovel, R. A., & Bunn, A. G. (2018). A multidecade

experiment shows that fertilization by salmon carcasses enhanced tree growth in the

riparian zone. *Ecology*, *99*(11), 2433–2441.

Scheuerell, M. D., Levin, P. S., Zabel, R. W., Williams, J. G., & Sanderson, B. L. (2005). A new

perspective on the importance of marine-derived nutrients to threatened stocks of

Pacific salmon (Oncorhynchus spp.). *Canadian Journal of Fisheries and Aquatic Sciences*,

62(5), 961–964.

Schindler, D. E., Leavitt, P. R., Brock, C. S., Johnson, S. P., & Quay, P. D. (2005). Marine-derived

nutrients, commercial fisheries, and production of salmon and lake algae in Alaska.

Ecology, *86*(12), 3225–3231.

Schmidt-Nielsen, K., & Knut, S.-N. (1984). *Scaling: why is animal size so important?* Cambridge

university press.

Sibly, R. M., Brown, J. H., & Kodric-Brown, A. (2012). *Metabolic ecology: a scaling approach*.

John Wiley & Sons.

Sousa, Tania, Domingos, T., & Kooijman, S. (2008). From empirical patterns to theory: a formal

metabolic theory of life. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *363*(1502), 2453–2464.

Sousa, Tânia, Domingos, T., Poggiale, J.-C., & Kooijman, S. (2010). *Dynamic energy budget*

theory restores coherence in biology. The Royal Society.

USDA Forest Service, Lake Tahoe Basin Management Unit, Tahoe Heritage Foundation. (2015).

Kokanee Samon Conservation Education Program PRE-VISIT PACKET. AN EDUCATOR'S

GUIDE TO TAYLOR CREEK. (pp. 1–17). South Lake Tahoe, CA 96150. Retrieved from

https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5391112.pdf

Van Valen, L. (1976). Energy and evolution. *Evolutionary Theory*, *1*(7), 179–229.

West, G. B., Brown, J. H., & Enquist, B. J. (1997). A general model for the origin of allometric

scaling laws in biology. *Science*, *276*(5309), 122–126.

Figure 1. Life cycle of sockeye salmon after Brett (JOHN R. Brett, 1983). The Babine Lake

population is semelparous – expending energy on a single, end-of-life reproductive effort, and

anadromous – spending about equal time – two years – reproducing in fresh water (green) and

feeding and growing in the ocean (blue). Modified from

("Https://Www.Google.Com/Search?Q=salmon+life+cycle&rlz=1C1CHFX_enUS555US555&sour

ce=lnms&tbm=isch&sa=X&ved=0ahUKEwj7rZiC3PXWAhWBLmMKHRIiAM8Q_AUICigB&biw=106

7&bih=423#imgrc=WiztTrFdqAKvPM," n.d.) and not drawn to scale.

Figure 2. Dynamics of Babine Lake salmon life history and energetics: variation as a function of

age. (A) Survival: number alive out of initial clutch of 3,000 eggs. Deviations from the fitted

exponential relationship are too small to show on this graph. (B) Body energy content. (C)

Cumulative respiration and assimilation.

Figure 3. Plots on logarithmic axes showing power-law scalings of rates of whole-organism

metabolism as functions of body mass before (black symbols) and after temperature (red

symbols and regression lines) correction. (A) Respiration rate (Brett's "total metabolic rate")**:**

$B = 0.00177 m^{0.898}$; R^2 = 0.996; B) Rate of biomass production (from Brett's "growth rate" in

%/day $P = 6.95 x 10^5 m^{1.29}$; R^2 = 0.76; (C) Assimilation rate (sum of the above two rates): $A =$

$0.00133 m^{1.0}$; R^2 = 0.94.

Figure 4. Balanced energy budgets for Babine Lake sockeye salmon: (A) for an individual female. The 4-year life cycle starts and ends with a single egg laid in fresh water; the offspring hatches, grows and enters the ocean, where it grows more rapidly before re-entering fresh water for the return migration; the 17626 kJ of stored body energy is all dissipated in fresh water as respiration, depleted carcass and clutch of eggs. The budget for a male is similar but he allocates less energy to sperm and more to respiration for courtship and mating. (B) for the cohort of all offspring produced by a surviving-to-spawning female. The 3,000 fry suffer high mortality and grow only modesty before entering the ocean. After voracious feeding, rapid growth, and additional mortality at sea, four survivors rapidly deplete their stored body energy on respiration to fuel migration and breeding, leaving a depleted carcass and a new clutch of 3,000 eggs.

Table 1 (Attached). Metabolic life table (MLT) for sockeye salmon in Babine Lake, based on data

in Brett (1983, 1986). The definitions of stages and traits follow Brett, although we have

transformed his data into modern units and calculated additional parameters. Green rows

represent stages in freshwater and blue rows marine. Unlike some life tables that are based

only on females, this one is based on replacement of a male and female parent at steady state.

Table 2 (Attached). Impact of a cohort of Babine Lake sockeye salmon on fresh water and

marine ecosystems. This accounting uses the MLT (Table 1) to quantify the input of food energy

from assimilation (green) and its allocation to biomass production: growth and gametes (blue)

and respiration (red).

Table 1. Metabolic life table (MLT) for sockeye salmon in Babine Lake, based on data in Brett (1983, 1986). The definitions of
stages and traits follow Brett, although we have transformed his data into modern units and calculated additional parameters.
Green rows represent stages in freshwater and blue rows marine. Unlike some life tables that are based only on females, this one
is based on replacement of a male and female parent at steady state.

(°C)	Stage (Months)	Wet weight (g)	Number alive (number)	Body energy content (kJ)	Production rate (watts)	Whole-organism respiration rate (watts)	Assimilation rate (watts)	Energy expended (kJ)	Cumulative energy expended (kJ)	Standing stock of biomass (kJ)	Cohort production lost to mortality (kJ)	Cohort respiration before dying (kJ)	Cohort assimilation before dying (kJ)
6	Egg (0)	0.013	3000	0.13	0.0000	0.0000002	0.0000000	0.00	0.00	386.03	0.00	0.00	0.00
6.0	Fry (9)	0.25	600	1.15	0.00	0.0002144	0.0009733	0.08	2.09	690.69	1535.79	2511.6	4047.39
11.0	Fry (11)	1.00	367	4.97	0.00	0.00078	0.0021052	2.09	9.21	1825.15	713.42	1316.623794	2030.04
9.0	Fry (13)	3.20	224	16.73	0.00	0.00233	0.0060826	7.12	34.74	3755.26	1546.30	3131.982487	4678.29
4.0	Fry (17)	4.50	137	23.51	0.00	0.00246	0.0032261	25.53	54.00	3228.30	1753.78	3868.102532	5621.89
5.0	Smolt yearling (20)	5.00	84	26.00	0.00	0.00254	0.0029052	19.26	62.37	2183.59	1319.87	3102.698087	4422.57
11.0	Smolt yearling (21)	5.40	75	29.39	0.00	0.00366	0.0045643	8.37	84.56	2195.64	257.02	681.9018036	938.93
12.0	Smolt yearling (22)	15.00	66	86.34	0.03	0.02006	0.0547624	22.19	179.16	5738.04	477.72	1088.681233	1566.40
12.5	Smolt yearling (23)	50.00	59	304.74	0.14	0.06203	0.2060148	94.60	400.60	18015.50	1436.05	2128.900796	3564.95
11.0	Smolt yearling (24)	110.00	53	708.65	0.22	0.10927	0.3250237	221.44	733.39	37264.19	3310.00	3703.907398	7013.91
9.0	Smolt yearling (25)	200.00	47	1288.45	0.30	0.15216	0.4489740	332.79	1117.24	60266.29	5802.26	5376.71596	11178.97
6.5	Smolt yearling (26)	250.00	42	1670.21	0.14	0.13447	0.2775480	383.86	1438.31	69490.30	7646.08	6604.313975	14250.39
4.8	Smolt yearling (27)	290.00	37	1937.45	0.11	0.11523	0.2251318	321.07	1740.54	71701.40	8293.05	7307.319567	15600.37
4.5	Smolt yearling (28)	300.00	33	2004.26	0.03	0.10903	0.1345524	302.23	2037.33	65977.58	8059.68	7724.674396	15784.36
4.5	Smolt yearling (29)	310.00	29	2071.07	0.03	0.11266	0.1390374	296.79	2314.44	60643.22	7412.10	7914.886782	15326.98
4.5	Smolt yearling (30)	320.00	26	2137.87	0.03	0.11630	0.1435225	277.11	2666.48	55682.15	6809.22	8058.131768	14867.35

5.3	Smolt yearling (31)	350.00	23	2433.74	0.09	0.14925	0.2352539	352.04	3134.06	56383.71	6578.69	8347.156985	14925.85
7.3	Smolt yearling (32)	380.00	21	2670.67	0.08	0.21913	0.3026008	467.58	3813.86	55035.73	6533.73	8893.456575	15427.18
9.0	2 yr immature (33)	420.00	18	2962.01	0.11	0.29511	0.4082625	679.81	4715.95	54294.69	6413.22	9711.818069	16125.04
11.0	2 yr immature (34)	480.00	16	3374.75	0.18	0.41170	0.5874999	902.08	5986.40	55024.72	6417.62	10838.90796	17256.53
12.0	2 yr immature (35)	590.00	15	4151.67	0.33	0.54893	0.8805507	1270.45	6142.54	60212.21	6780.18	10926.33887	17706.51
11.5	2 yr immature (36)	720.00	13	5074.27	0.39	0.62453	1.0122159	1562.63	7731.96	65460.64	7392.79	11117.6938	18510.48
9.3	2 yr immature (37)	890.00	11	6263.93	0.51	0.60379	1.1186258	1589.42	9151.43	71878.52	8081.41	12033.79686	20115.20
6.5	2 yr immature (38)	1000.00	10	7283.64	0.32	0.46520	0.7883632	1419.47	10315.56	74344.03	8589.14	12342.04889	20931.19
5.0	2 yr immature (39)	1060.00	9	7587.54	0.17	0.43147	0.5983565	1164.13	11380.48	68888.12	8386.47	12235.29054	20621.76
4.5	2 yr immature (40)	1100.00	8	7873.87	0.11	0.36247	0.4718452	1064.92	12349.96	63588.21	7755.83	11903.78294	19659.62
4.5	2 yr immature (41)	1130.00	7	8088.61	0.09	0.36688	0.4526258	969.48	13257.06	58104.24	7122.38	11425.72549	18548.10
4.5	2 yr immature (42)	1180.00	6	8594.70	0.14	0.38311	0.5224005	907.11	14376.40	54917.40	6621.44	10967.44636	17588.88
5.0	2 yr immature (43)	1250.00	6	9104.55	0.20	0.44824	0.6484892	1119.34	15805.08	51746.79	6248.42	10655.06793	16903.49
7.0	3 yr maturing (44)	1400.00	5	10537.00	0.46	0.67841	1.1419302	1428.68	18086.03	53270.57	6167.89	10642.58165	16810.48
9.5	4 yr maturing (45)	1670.00	4	12576.42	0.86	1.07630	1.9352652	2281.79	21685.61	56555.24	6456.11	11109.1341	17565.24
12.5	Adult-mature (46)	2100.00	4	16315.35	1.44	1.80118	3.2365951	3598.70	26813.00	65261.62	7178.38	12049.84547	19228.22
13.0	Adult-mature (46.5)	2270.00	4	17626.41	0.38	2.02399	2.4081649	5127.43	29435.53	70505.64	8485.44	14062.13445	22547.58
12.0	Adult-mature (48)	2200.00	2	4455.00	0.00	2.55858	2.5585824	2622.53	39013.52	8910.00	6682.50	58520.28	65202.78
12.0	Carcass (48.1)	2161.00	0	4169.00	0.00	0.00000	0.0000000	5345.52	39013.52	0.00	8338.00	78027.04	86365.04

Ecosystem	Fresh water			Marine			Fresh Water	
	Dead	Alive	Total	Dead	Alive	Total	Total	Net flow
Assimilation	20800	7423	28223	88016	188002	576018	0	604241
Respiration	13931	5239	19170	225790	117496	343286	41525	403980
Production	6869	2184	9053	162227	70506	232733	0	200261
Imported biomass						2184	70506	
4 Carcasses							24162	
Clutch							3880	
Sperm							938	
Biomass left in fresh water							28980	
Production efficiency	0.23	0.21	0.22	0.29	0.26	0.28	0.0	0.23

Table 2. Impact of a cohort of Babine Lake sockeye salmon on fresh water and marine ecosystems. This accounting uses the MLT (Table 1) to quantify the input of food energy from assimilation (green) and its allocation to biomass production: growth and gametes (blue) and respiration (red).