ICEs are the main reservoirs of the ciprofloxacin-modifying *crpP* gene in

Pseudomonas aeruginosa

João Botelho[1,2,*], Filipa Grosso[3], Luísa Peixe[3]

[1]Antibiotic Resistance Evolution Group, Max-Planck-Institute for Evolutionary Biology, 24306

Plön, Germany.

[2]Department of Evolutionary Ecology and Genetics, Zoological Institute, Christian-Albrechts-

Universität zu Kiel, Am Botanischen Garten 1-9, 24118 Kiel, Germany.

[3]UCIBIO/REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do

Porto, Porto, Portugal.

*Corresponding author: João Botelho, Antibiotic Resistance Evolution Group, Max-Planck-

Institute for Evolutionary Biology, 24306 Plön, Germany; Department of Evolutionary Ecology

and Genetics, Zoological Institute, Christian-Albrechts-Universität zu Kiel, 24118 Kiel, Germany.;

e-mail: botelho@evolbio.mpg.de. Luísa Peixe, Laboratório de Microbiologia. Faculdade de

Farmácia da Universidade do Porto, Rua Jorge Viterbo Ferreira nº 228, 4050-313 Porto, Portugal;

phone: +351 220428500; e-mail: lpeixe@ff.up.pt.

Abstract

The ciprofloxacin-modifying *crpP* gene was recently identified in a plasmid isolated from a

clinical *Pseudomonas aeruginosa* clinical isolate. Homologues of this gene were also identified

in *Escherichia coli* and *Klebsiella pneumoniae*. We set out to explore the mobile genetic elements

involved in the acquisition and spread of this gene in publicly available and complete genomes

of *Pseudomonas*. The *crpP* gene was identified only in *P. aeruginosa*, in more than half of the

complete chromosomes (61.9%, n=133/215) belonging to 52 sequence types, of which the high-

risk clone ST111 was the most frequent. We identified 136 *crpP*-harboring ICEs, with 93.4%

belonging to the mating-pair formation G (MPF$_G$) family. The ICEs were integrated at the end of

a tRNA^{Lys} gene and were all flanked by highly conserved 45-bp direct repeats. The core ICEome

contains 26 genes (2.2% of all genes), which are present in 99% or more of the *crpP*-harboring

ICEs. The most frequently encoded traits on these ICEs include replication, transcription,

intracellular trafficking and cell motility. Our work reveals that ICEs are the main vectors

promoting the dissemination of the ciprofloxacin-modifying *crpP* gene in *P. aeruginosa*.

Introduction

Pseudomonas aeruginosa is a frequent cause of severe nosocomial infections and is one of the

six ESKAPE pathogens [1,2]. Ciprofloxacin is an antibiotic of the fluoroquinolone class that is

active against *P. aeruginosa* infections [3,4]. In this species, a high proportion of clinical isolates

are resistant to ciprofloxacin [5,6]. Commonly reported mechanisms of ciprofloxacin resistance

include mutations in DNA gyrase and topoisomerase IV-encoding genes *gyrA*, *gyrB*, *parC* and

parE and efflux pumps regulatory genes as *nalC* and *nfxB* [6–8].

Besides chromosomal mutations, ciprofloxacin resistance can be mediated by horizontally

transferred genes, such as the quinolone resistance (*qnr*) gene [9]. The ciprofloxacin-modifying

crpP gene induces ATP-dependent phosphorylation of the antibiotic and was originally identified

in a plasmid isolated from a *P. aeruginosa* clinical isolate [10]. This plasmid conferred resistance

to ciprofloxacin when transferred to *P. aeruginosa* PAO1 strain. The authors also cloned the gene

into a shuttle vector and the resulting recombinant plasmid conferred an increased minimum

inhibitory concentration (MIC) to the same antibiotic in *Escherichia coli*. Homologous *crpP* genes

were also identified in *E. coli* and *Klebsiella pneumoniae* clinical isolates from Mexican hospitals

and conferred decreased susceptibility to ciprofloxacin [11,12].

Integrative and conjugative elements (ICEs) are mobile elements involved in vertical and

horizontal transmission of antibiotic resistance genes (ARGs) and other beneficial genes in

bacterial communities [2,13,14]. ICEs can be classified into eight mating-pair formation (MPF)

classes based on the classification of the secretion machinery involved in conjugation [15,16].

Recently, Ruiz verified that several *CrpP* proteins were encoded next to genes typically present

in ICEs [17]. Building on this observation, we set out to trace and characterize the *crpP*-harboring

ICEs present in all complete *Pseudomonas* genomes available on NCBI.

Material and methods

Pangenome and whole-genome analyses

All complete *Pseudomonas* genomes (n=577 chromosomes and 163 plasmids) were downloaded

from NCBI's Refseq library on the 01/02/2020 using ncbi-genome-download v 0.2.11

(https://github.com/kblin/ncbi-genome-download) (**Table S1**). We used mlst v 2.11

(https://github.com/tseemann/mlst) to scan the genomes against the PubMLST *Pseudomonas*

typing schemes (https://pubmlst.org/) [18]. The genomes were annotated using prokka v1.14.5

(https://github.com/tseemann/prokka) [20]. We used the .gff files created by Prokka to

calculate the *P. aeruginosa* pangenome in roary v3.13.0 (https://github.com/sanger-

pathogens/Roary) [21]. The core genome alignment created by Roary was used as input in

fasttree v2.1.10 (http://www.microbesonline.org/fasttree/) [22] to create an approximately-

maximum-likelihood phylogenetic tree using a generalized time-reversible (GTR) model of

nucleotide evolution. We used the newick file created by FastTree to create a phylogenetic tree

in iTOL (https://itol.embl.de/) [23]. We performed functional annotation based on orthology

assignments (COGs) of protein files created by prokka using eggNOG-mapper v2 (http://eggnog-

mapper.embl.de/) [24].

Mining ICEs in complete genomes

We used the ICEfinder standalone version (http://202.120.12.136/ICEfinder/ICEfinder.html) and

manual curation to trace putative ICEs. We searched for ARGs on extracted ICEs using amrfinder

v 3.6.7 and default parameters (50% minimum coverage of the reference protein and 90%

minimum identity) (https://github.com/ncbi/amr/wiki/AMRFinder-database) [25]. Positive ICE

hits for *CrpP*-encoding genes were further characterized; we used fastANI v1.3

(https://github.com/ParBLiSS/FastANI) [19] to compute whole-genome average nucleotide

identity of non-*P. aeruginosa crpP*-positive hits, antismash online tool

(https://antismash.secondarymetabolites.org/#!/start) [26] to look for secondary metabolite

biosynthesis gene clusters, BAGEL (http://bagel4.molgenrug.nl/) [27] to trace bacteriocins,

macsyfinder v1.0.5 (https://github.com/gem-pasteur/macsyfinder) and the CRISPRCasFinder

online tool (https://crisprcas.i2bc.paris-saclay.fr/) to detect CRISPRs and *cas* genes [28]. We

used roary to infer the total *crpP*-harboring ICE content, here referred to as the ICEome.

Results

The *crpP* gene is widespread in *P. aeruginosa*

The *CrpP*-encoding gene was identified in 23.1% of the chromosomes (n=133/577, including 131

P. aeruginosa, 1 *Pseudomonas fluorescens* and 1 *Pseudomonas* sp.) and 1 *P. aeruginosa* plasmid.

However, we compared the ANI between non-*P. aeruginosa* hits (the *P. fluorescens* strain

NCTC10783 and *Pseudomonas* sp. AK6U strains with accession numbers NZ_LR134300.1 and

NZ_CP025229.1, respectively) and the *P. aeruginosa* DSM 50071 type strain (accession number

NZ_CP012001.1) and we realized that these strains belong to the *P. aeruginosa* species, as the

ANI value is above the 95% cutoff for species delineation (**Table S2**). All *crpP*-hits were only

identified in *P. aeruginosa* complete genomes, as so from here on we focused our attention on

P. aeruginosa genomes. Indeed, the percentage increases if we only consider the *P. aeruginosa*

chromosomes (61.9%, n=133/215), meaning that the *crpP* gene is present in more than half of

the *P. aeruginosa* complete genomes (**Figure 1**). The GC content of *crpP*-harboring

chromosomes varies from 65.6 to 66.5% (**Figure 2**) and the sequence length from 6.3 to 7.5 Mb.

We identified several clones (n=52 sequence types), of which the high-risk clone ST111 [29] was

the most frequent (**Table S3**).

Tree scale: 0.01

Figure 1. Approximately-maximum-likelihood phylogenetic tree base on the core genome

alignment of 215 complete *P. aeruginosa* genomes. Branches and labels from *crpP*-positive hits

are colored blue and the branches have twice the standard width. Only bootstrap values from

0.9 to 1 are displayed. The newick file used to create the tree is available in **Additional file 1**.

Figure 2 – Distribution of the GC content of the chromosomes and ICEs. The violin plots were

created using ggplot2.

The majority of *crpP*-harboring ICEs belong to the MPF_G family

The majority of *crpP*-harboring ICEs belong to the MPF$_{G}$ family

A total of 316 putative ICEs were identified among the 133 chromosomes. Each chromosome

carries at least one ICE. Most *crpP* genes present in the chromosome were associated with an

ICE (besides *P. aeruginosa* strains W60856 and B17932 with the accession numbers

NZ_CP008864.2 and NZ_CP034436.1, respectively) with 43.0% of the total putative ICEs

(n=136/316) harboring the ciprofloxacin-modifying gene. In *P. aeruginosa* RW109 strain

(accession number NZ_LT969520.1, position 5629923-5820312 bp), we identified two *crpP*-

harboring ICEs in tandem (**Table S3**). ICE size varied from 81.6 to 145.5 kb, and the GC content

from 56.6 to 61.5% (**Figure 2**). Most of the ICEs (n=129/136) ICEs belong to the MPF$_G$ family. For

the remaining 7, no MPF family could be determined.

The *crpP*-carrying ICEs integrate into a specific hotspot

All *crpP*-carrying ICEs identified in this study were integrated at the end of a tRNA^{Lys} gene, that

was found alone or in a tRNA cluster with tRNA^{Pro} and tRNA^{Asn}. The only exception was observed

in the two contiguous ICEs identified in *P. aeruginosa* RW109 strain, where the second ICE was

integrated at the end of the first one. The tRNA^{Lys} genes shared the exact same sequence. All

ICEs were flanked by 45-bp (5'-TGGTGGGTCGTGTAGGATTCGAACCTACGACCAATTGGTTAAAAG-

3') highly conserved direct repeats (DRs). The 136 integrases identified in this study, however,

only shared 36.1% amino acid identity.

The *P. aeruginosa* pangenome and the *crpP*-carrying ICEome

The total number of genes among the 215 *P. aeruginosa* complete genomes (pangenome) is

33570. The core genome (genes present in 99% or more of the genomes) contains 2674 genes

(8.0% of all genes, **Figure 3** and **Figures S1**). We also identified 11241 unique genes, which are

found in only one strain [30].

Figure 3 – Pie chart of the breakdown of genes and the number of *P. aeruginosa* genomes they
are present in. This figure was created using the contributed Python script roary_plots.py in
https://github.com/sanger-pathogens/Roary/blob/master/contrib/roary_plots/roary_plots.py.

The total number of genes in the 136 *crpP*-harboring ICEs (ICEome) is 1193. The core ICEome
contains 26 genes (2.2% of all genes). The soft core content (genes present in more or less than
95% and less than 99% of ICEs) includes 11 genes, 129 shell genes (between 15 and 95%) and
1027 cloud genes (between zero and 15%). We identified 451 unique genes among the ICEome.

The *crpP*-carrying ICEs encode other beneficial genes

The most frequently encoded traits on *crpP*-harboring ICEome include replication, transcription,
intracellular trafficking, and cell motility (**Figure 4**). However, nearly one-third of the proteins
for which a COG category was attributed (30.7%, n=211/687) encode for an unknown function
(**Table S4**).

The *crpP* gene encoded for proteins ranging from 39 to 68 amino acid and is present in a single
copy, with no flanking mobile units as integrons, insertion sequences and/or transposons or
found on the close vicinity of the gene. We also observed a lack of co-resident antibiotic
resistance genes (ARGs) within the *crpP*-harboring ICEs. Seven ICEs carry a type I-C CRISPR-Cas
system and ten harbor pyocin S5-encoding genes (**Table S3**). The bacteriocins here identified are

171   highly identical, sharing 95% amino acid identity. Besides bacteriocins, no secondary metabolites

172   were identified within the extracted ICEs.

176   **Figure 4**. Functional annotation of proteins encoded by the *crpP*-harboring ICEome.

Discussion

Our results show that ICEs are the main drivers for the spread of the ciprofloxacin-modifying
crpP gene in *P. aeruginosa*. A selective pressure exerted by the use of ciprofloxacin to treat *P.*
aeruginosa infections may have promoted the dissemination of *crpP*-harboring ICEs among
several clones.

Curiously, and in opposition to several ARGs, no integrons, insertion sequences or transposons
were found on the vicinity of the *crpP* gene. Bacteriocins produced by bacteria are important for
niche competition. The pyocin S5 possesses a bactericidal activity against clinical *P. aeruginosa*
isolates [31,32]. The presence of these bacteriocins among some *crpP*-harboring ICEs can help
to eliminate possible competitors and indirectly enhance ICE maintenance.

As expected for foreign DNA, the GC content of the ICEs is lower than that from the
chromosomes. Still, given the high GC content of these mobile elements and the biological
barrier for gene acquisition between donor-recipient pairs having >5% difference in GC values
[33], we suspect that the ancestor of the *crpP* gene derives from GC-rich taxa.

We identified highly conserved DRs that can be used to accurately delimit *crpP*-carrying ICEs in
Pseudomonas. Interestingly, the exact 45-bp sequence was found on the 2 ICEs from *P.*
fluorescens and the *Pseudomonas* sp. AK6U strain. Also, even though the *crpP* gene was first
reported in a plasmid in *P. aeruginosa* [10], we found a perfect match for the 45-bp DR next to
an integrase-encoding gene. The MPF_G class signatures we identified within the boundaries of
this plasmid probably means that the plasmid evolved from an ICE, or that the ICE integrated
within the plasmid. We only found one match for the DR in this plasmid, meaning that the ICE is
degenerated and most likely depends on the plasmid to be mobilized. The same was observed
for the *P. aeruginosa crpP*-carrying plasmid detected in this study (accession number
NZ_CP030914.1).

The *crpP*-carrying ICEs were traced in several clones, taking advantage of the highly conserved
attachment site in the $tRNA^{Lys}$ gene. Interestingly, we found in a previous study that MPF_G class
ICEs carrying specific ARGs were integrated at the end of $tRNA^{Gly}$ genes [14]. In fact, this class of
ICEs is frequently integrated at the end of a tRNA gene [34]. This behavior suggests selection for
the maintenance of these non-coding integration sites, and seamless site-specific integration of
ICEs in the chromosome will incur in a lower burden to the recipient cell and therefore increase
its fitness [2,35]. As so, conferring decreased susceptibility or low levels of ciprofloxacin

resistance without incurring in fitness cost can help to explain the high frequency of *crpP*-
carrying ICEs observed in this study. Future studies should explore the possible synergistic
effects between commonly reported mutations leading to ciprofloxacin resistance and the
presence of *crpP*-harboring ICEs.

One of the caveats of our study is that we restricted the analyses to complete genomes. Given
that the majority of the assemblies present in public databases as NCBI correspond to draft
genomes, we probably missed the identification of several *crpP*-carrying ICEs among *P.*
aeruginosa. However, mobile elements as ICEs tend to be fragmented in genomes sequenced
with short-reads technologies, due to the presence of repetitive regions. Also, the identification
of ICE fragments within the draft genome would require similar ICE sequences to use as a
reference, which would then bias the study. Sequencing bacteria with long-read approaches is
increasing, which will improve the accurate delimitation of an ever-growing number of mobile
elements.

Conclusions

Our work demonstrates that ICEs are the main promoters involved in the dissemination of the
ciprofloxacin-modifying *crpP* gene among clonally-unrelated *P. aeruginosa* strains. We also
identified highly conserved 45-bp DRs than can be used to accurately trace *crpP*-harboring ICEs
on *P. aeruginosa* genomes. Future studies should explore the mobile elements involved in the
acquisition of the *crpP* homologues identified in species other than *P. aeruginosa* and the
possible synergistic effects between commonly reported mutations leading to ciprofloxacin
resistance and the presence of *crpP*-carrying ICEs.

Conflict of interests

The authors declare that there are no conflicts of interest.

Acknowledgments

This work was supported by the Applied Molecular Biosciences Unit - UCIBIO which is financed
by national funds from FCT (UIDB/04378/2020).

References

1   Boucher, H.W. *et al.* (2009) Bad Bugs, No Drugs: No ESKAPE! An Update from the
241   Infectious Diseases Society of America. *Clin. Infect. Dis.* 48, 1–12

242   2   Botelho, J. *et al.* (2019) Antibiotic resistance in Pseudomonas aeruginosa –
243   Mechanisms, epidemiology and evolution. *Drug Resist. Updat.* 44, 100640

244   3   Paulsson, M. *et al.* (2017) Antimicrobial combination treatment including ciprofloxacin
245   decreased the mortality rate of Pseudomonas aeruginosa bacteraemia: a retrospective
246   cohort study. *Eur. J. Clin. Microbiol. Infect. Dis.* 36, 1187–1196

247   4   Klodzińska, S.N. *et al.* Inhalable antimicrobials for treatment of bacterial biofilm-
248   associated sinusitis in cystic fibrosis patients: Challenges and drug delivery approaches.
249   , *International Journal of Molecular Sciences*, 17. 01-Oct-(2016) , MDPI AG

250   5   Pitt, T.L. *et al.* Survey of resistance of Pseudomonas aeruginosa from UK patients with
251   cystic fibrosis to six commonly prescribed antimicrobial agents. , *Thorax*, 58. 01-Sep-
252   (2003) , Thorax, 794–796

253   6   Rehman, A. *et al.* Mechanisms of ciprofloxacin resistance in pseudomonas aeruginosa:
254   New approaches to an old problem. , *Journal of Medical Microbiology*, 68. 01-Jan-
255   (2019) , Microbiology Society, 1–10

256   7   Bruchmann, S. *et al.* (2013) Quantitative contributions of target alteration and
257   decreased drug accumulation to Pseudomonas aeruginosa fluoroquinolone resistance.
258   *Antimicrob. Agents Chemother.* 57, 1361–8

259   8   Cantón, R. and Ruiz-Garbajosa, P. (2011) Co-resistance: an opportunity for the bacteria
260   and resistance genes. *Curr. Opin. Pharmacol.* 11, 477–485

261   9   Belotti, P.T. *et al.* (2015) Description of an original integron encompassing *bla* VIM-2 ,
262   *qnrVC1* and genes encoding bacterial group II intron proteins in *Pseudomonas*
263   *aeruginosa*. *J. Antimicrob. Chemother.* 70, 2237–2240

264   10   Chávez-Jacobo, V.M. *et al.* (2018) *CrpP* is a novel ciprofloxacin-modifying enzyme
265   encoded by the pseudomonas aeruginosa pUM505 plasmid. *Antimicrob. Agents*
266   *Chemother.* 62, e02629-17

267   11   Chávez-Jacobo, V.M. *et al.* (2019) Prevalence of the *crpP* gene conferring decreased
268   ciprofloxacin susceptibility in enterobacterial clinical isolates from Mexican hospitals. *J.*

Antimicrob. Chemother. 74, 1253–1259

12   Pablo, J. *et al.* (2020) Identification of Essential Residues for Ciprofloxacin Resistance of
Ciprofloxacin- - Modifying Enzyme (*CrpP*) of pUM505. *Microbiology* DOI:
10.1099/mic.0.000889

13   Partridge, S.R. *et al.* (2018) Mobile Genetic Elements Associated with Antimicrobial
Resistance. *Clin. Microbiol. Rev.* 31, e00088-17

14   Botelho, J. *et al.* (2018) Carbapenemases on the move: it's good to be on ICEs. *Mob.*
DNA 9, 37

15   Guglielmini, J. *et al.* (2011) The Repertoire of ICE in Prokaryotes Underscores the Unity,
Diversity, and Ubiquity of Conjugation. *PLoS Genet.* 7, e1002222

16   Guglielmini, J. *et al.* (2014) Key components of the eight classes of type IV secretion
systems involved in bacterial conjugation or protein secretion. *Nucleic Acids Res.* 42,
5715–27

17   Ruiz, J. (2019) *CrpP*, a passenger or a hidden stowaway in the Pseudomonas aeruginosa
genome? *J. Antimicrob. Chemother.* DOI: 10.1093/jac/dkz316

18   Jolley, K.A. and Maiden, M.C.J. (2010) BIGSdb: Scalable analysis of bacterial genome
variation at the population level. *BMC Bioinformatics* 11, 595

19   Jain, C. *et al.* (2018) High throughput ANI analysis of 90K prokaryotic genomes reveals
clear species boundaries. *Nat. Commun.* 9, 5114

20   Seemann, T. (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30,
2068–2069

21   Roary: Rapid Large-Scale Prokaryote Pan Genome Analysis - PubMed. . [Online].
Available: https://pubmed.ncbi.nlm.nih.gov/26198102-roary-rapid-large-scale-
prokaryote-pan-genome-analysis/. [Accessed: 23-Feb-2020]

22   FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance
Matrix | Molecular Biology and Evolution | Oxford Academic. . [Online]. Available:
https://academic.oup.com/mbe/article/26/7/1641/1128976. [Accessed: 27-Feb-2020]

23   Letunic, I. and Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res.* 47, W256–W259

24 eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology
resource based on 5090 organisms and 2502 viruses | Nucleic Acids Research | Oxford
Academic. . [Online]. Available:
https://academic.oup.com/nar/article/47/D1/D309/5173662. [Accessed: 24-Feb-2020]

25 Feldgarden, M. *et al.* (2019) Validating the AMRFinder Tool and Resistance Gene
Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a
Collection of Isolates. *Antimicrob. Agents Chemother.* 63,

26 antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline | Nucleic
Acids Research | Oxford Academic. . [Online]. Available:
https://academic.oup.com/nar/article/47/W1/W81/5481154. [Accessed: 21-Feb-2020]

27 van Heel, A.J. *et al.* (2018) BAGEL4: a user-friendly web server to thoroughly mine RiPPs
and bacteriocins. *Nucleic Acids Res.* 46, W278–W281

28 Couvin, D. *et al.* (2018) CRISPRCasFinder, an update of CRISRFinder, includes a portable
version, enhanced performance and integrates search for Cas proteins. *Nucleic Acids*
Res. 46, W246–W251

29 Oliver, A. *et al.* (2015) The increasing threat of Pseudomonas aeruginosa high-risk
clones. *Drug Resist. Updat.* 21–22, 41–59

30 Tettelin, H. *et al.* (2005) Genome analysis of multiple pathogenic isolates of
Streptococcus agalactiae: Implications for the microbial "pan-genome." *Proc. Natl.*
Acad. Sci. U. S. A. 102, 13950–13955

31 Ling, H. *et al.* (2010) A predicted S-type pyocin shows a bactericidal activity against
clinical Pseudomonas aeruginosa isolates through membrane damage. *FEBS Lett.* 584,
3354–3358

32 Elfarash, A. *et al.* (2014) Pore-forming pyocin S5 utilizes the FptA ferripyochelin
receptor to kill Pseudomonas aeruginosa. *Microbiol. (United Kingdom)* 160, 261–269

33 Popa, O. *et al.* (2011) Directed networks reveal genomic barriers and DNA repair
bypasses to lateral gene transfer among prokaryotes. *Genome Res.* 21, 599–609

34 Cury, J. *et al.* (2017) Integrative and conjugative elements and their hosts: composition,
distribution and organization. *Nucleic Acids Res.* 45, 8943–8956

35 Touchon, M. *et al.* The chromosomal accommodation and domestication of mobile

genetic elements. , *Current Opinion in Microbiology*, 22. 01-Dec-(2014) , Elsevier Ltd,

22–29

330