Sulfisoxazole does not inhibit the secretion of small extracellular vesicles

Pamali Fonseka*, Sai V Chitti*, Rahul Sanwlani* and Suresh Mathivanan[#]

Department of Biochemistry and Genetics, La Trobe Institute for Molecular Science, La Trobe University, Melbourne, Victoria, 3086, Australia

*Authors contributed equally.

[#]To whom correspondence should be addressed

Professor Suresh Mathivanan

Department of Biochemistry and genetics,

La Trobe institute for molecular Science,

La Trobe University,

Bundoora, Victoria 3086, Australia

Tel: +61 03 9479 2565

Email: S.Mathivanan@latrobe.edu.au

Abstract

Recently, the study by Im et *al.* focused on blocking the release of extracellular vesicles (EVs) by cancer cells, as a strategy to block metastasis, by deploying a drug repurposing screen. Upon screening the library of FDA approved drugs in breast cancer cells *in vitro*, the authors reported the ability of the antibiotic Sulfisoxazole (SFX) in inhibiting EV biogenesis and secretion. SFX was also effective in reducing breast primary tumor burden and blocking metastasis in immunocompromised and immunocompetent mouse models. As we seek a compound to block EV biogenesis and secretion in our current *in vivo* studies, we intended to use SFX and hence performed *in vitro* characterization as the first step. However, treatment of two cancer cells with SFX did not reduce the amount of EVs as reported by the authors.

Over the last two decades, extracellular vesicles (EVs) have been implicated in intercellular communication and utilised as drug delivery vehicles and as reservoirs of disease biomarkers[1-5]. As they continue to garner interest, several seminal studies have established that EVs regulate various pathophysiological processes in favour of cancer progression, including remodelling the tumour microenvironment, immune evasion, coagulation, vascular leakiness, establishing the pre-metastatic niche, tropism for metastasis and transfer of chemoresistance[6-11]. Hence, there is growing interest in blocking the release of EVs and limiting their systemic circulation as a novel therapeutic avenue to treat cancer[12]. As anti-metastatic therapies are scarce, it is speculated that FDA approved drugs that target EVs could possibly fill the void.

Recently, the study by Im *et al.* focused on blocking the release of EVs by cancer cells, as a strategy to block metastasis, by deploying a drug repurposing screen[13]. The rationale was to screen the existing FDA approved library to identify drugs which can inhibit EV biogenesis or secretion with the obvious advantage of known mode of action, efficacy and toxicity profile and hence has the potential of immediate clinical utility. Upon screening the library of FDA approved drugs in metastatic breast cancer cells *in vitro*, the authors reported the ability of the antibiotic Sulfisoxazole (SFX) in inhibiting EV biogenesis and secretion. The authors also reported that SFX was effective in reducing breast primary tumor burden and blocking metastasis in immunocompromised and immunocompetent mouse models. SFX was proposed to target Endothelin receptor A (ETA) previously in 1994[14] and hence the authors validated that SFX targets ETA which can positively regulate EV biogenies and secretion. The findings in this study thus present SFX as a potential novel EV-targeted therapeutic alternative. As a group interested in EVs, the outcomes proposed in the study were encouraging and attractive as FDA approved drugs targeting EV release are limited. Recently, Datta *et al.* also performed a repurposing screen to identify drugs that modulate the release of EVs in prostate cancer cells[12]. However, the identified drugs are yet to be tested *in vivo*.

As we seek a compound to block EV biogenesis and secretion in our current *in vivo* studies, we intended to use SFX and hence performed *in vitro* characterization as the first step. However, treatment of 4T1 breast cancer cells with SFX did not reduce the amount of EVs as reported by the authors. We acknowledge the fact that our EV

isolation protocol[15] was different from the study[13] (**Fig. 1** – Mathivanan laboratory protocol) and hence could have attributed to the varied results. In order to rule out the possibility of variations in the method of EV isolation or cell-type dependency, three researchers exactly followed the protocol employed by the authors (**Fig. 1** – protocol 1 and 2) to isolate EVs from two different cell types (4T1 and MDA-MB231) that were used by the authors. All these assays were performed with 3 technical replicates (same batch of cells seeded in three different plates) for every biological replicate. Consistent with our previous observations, treatment of the cancer cells with varying concentrations of SFX (50, 100 and 200 µM) did not impede the release of EVs while the positive control ceramide inhibitor GW4869, at low concentration (5 µM), inhibited EV secretion. Upon EV isolation, we quantified the protein amount, particle number and performed Western blotting for EV enriched proteins (TSG101, Alix), all normalised to equal cell number (**Fig. 2**). Contrary to the authors claim of SFX treatment led to a 3-fold decline in EV particle number, we observed a significant increase in particle number upon SFX (200 µM) treatment.

Overall, we report that SFX does not reduce the release of EVs (3 independent researchers) and emphasise caution in using SFX (Merk 31739) as a drug to block EV release. However, we do acknowledge the fact that our findings do not challenge the authors main conclusion of SFX mediated reduction of primary tumour burden and metastasis though our results suggest that the phenotype observed may not be cancer cell-derived EV mediated. Further research is needed to understand as how SFX can reduce primary tumor burden and inhibit metastasis.

Acknowledgements

Suresh Mathivanan is supported by Australian Research Council Future Fellowship (FT180100333). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Fig. 1 has been created with Biorender.com.

Competing interests

The authors declare no competing financial interests.

Author contributions

S.V.C. and S.M. conceived the concept. P.F., S.V.C. and R.S. performed the experiments. P.F. and S.M. made the figures. R.S. and S.M. wrote the manuscript. All authors read and approved the manuscript.

References

Peinado, H. *et al.* Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. *Nat. Med.* **18**, 883-891 (2012).

Kamerkar, S. *et al.* Exosomes facilitate therapeutic targeting of oncogenic KRAS in pancreatic cancer. *Nature* **546**, 498-+, doi:10.1038/nature22341 (2017).

Yoshioka, Y. *et al.* Ultra-sensitive liquid biopsy of circulating extracellular vesicles using ExoScreen. *Nature communications* **5**, 3591, doi:10.1038/ncomms4591 (2014).

Thery, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles* **7**, 1535750, doi:10.1080/20013078.2018.1535750 (2018).

Fonseka, P. *et al.* Exosomes from N-Myc amplified neuroblastoma cells induce migration and confer chemoresistance to non-N-Myc amplified cells: implications of intra-tumor heterogeneity. *J Extracell Vesicles* **8**, 1597614 (2019).

Tkach, M. & Thery, C. Communication by Extracellular Vesicles: Where We Are and Where We Need to Go. *Cell* **164**, 1226-1232, doi:10.1016/j.cell.2016.01.043 (2016).

Al-Nedawi, K. *et al.* Intercellular transfer of the oncogenic receptor EGFRvIII by microvesicles derived from tumour cells. *Nat. Cell Biol.* **10**, 619-624 (2008).

Rajesh, D., Schell, K. & Verma, A. K. Ras mutation, irrespective of cell type and p53 status, determines a cell's destiny to undergo apoptosis by okadaic acid, an inhibitor of protein phosphatase 1 and 2A. *Mol. Pharmacol.* **56**, 515-525 (1999).

Hoshino, A. *et al.* Tumour exosome integrins determine organotropic metastasis. *Nature* **527**, 329-335 (2015).

Kalluri, R. & LeBleu, V. S. The biology, function, and biomedical applications of exosomes. *Science* **367**, doi:10.1126/science.aau6977 (2020).

Anand, S., Samuel, M., Kumar, S. & Mathivanan, S. Ticket to a bubble ride: Cargo sorting into exosomes and extracellular vesicles. *Biochim Biophys Acta Proteins Proteom*, doi:10.1016/j.bbapap.2019.02.005 (2019).

Datta, A. *et al.* High-throughput screening identified selective inhibitors of exosome biogenesis and secretion: A drug repurposing strategy for advanced cancer. *Sci Rep* **8**, 8161, doi:10.1038/s41598-018-26411-7 (2018).

Im, E. J. *et al.* Sulfisoxazole inhibits the secretion of small extracellular vesicles by targeting the endothelin receptor A. *Nature communications* **10**, 1387, doi:10.1038/s41467-019-09387-4 (2019).

Chan, M. F. *et al.* Identification of a new class of ETA selective endothelin antagonists by pharmacophore directed screening. *Biochem. Biophys. Res. Commun.* **201**, 228-234, doi:10.1006/bbrc.1994.1693 (1994).

Kalra, H. *et al.* Extracellular vesicles containing oncogenic mutant beta-catenin activate Wnt signalling pathway in the recipient cells. *J Extracell Vesicles* **8**, 1690217, doi:10.1080/20013078.2019.1690217 (2019).

Figure legend

Figure 1

Schematic representation of cell culture, drug treatment and sEV isolation by two groups

(a) Schematic representation of cell culture and drug treatment approaches highlighting our approach. The cells were seeded and allowed to grow to 70-80% confluency in media supplemented with FCS. Upon attaining desired confluency, cells were washed with 1 x PBS and subjected to drug treatment for 24 h in serum free media. At treatment endpoint, conditioned media was harvested and subjected to sEV isolation. Live cell number was determined to normalise particles released to cell number. Similarly, Baek group seeded cells in a pre-defined range and treated them with drug in serum free media. **(b)** Protocol 1, schematic flow diagram depicting the methodology of sEV isolation and analyses as defined by Baek group in the initial publication (Im et al.) and followed by Fonseka *et al.* Protocol 2, schematic flow diagram depicting the methodology of sEV isolation and analyses as defined by Baek group in the latest version (Lee et al.). The variations in the methodology from the original manuscript (Im et al.) is marked (*). Protocol 2 was also adapted by Fonseka *et al.* to include the variations (*) suggested by Baek group. Protocol 3, standardized and optimized method of sEV isolation in EV depleted serum used in Mathivanan laboratory was also used to collect sEVs post treatment with SFX.

Figure 2

Quantification of EVs released by cells with or without SFX and GW4869

(a) Schematic of EV quantification by three different methods. Firstly, the total EV protein amount was quantified and normalised to equal number of live cells. Secondly, nanoparticle tracking analysis (NTA) was performed to quantify the total number of particles normalised to equal number of live cells. Lastly, Western blot analysis of EV samples obtained from equal number of live cells was performed for EV enriched proteins. **(b)** Relative amount of EV protein normalised to 10^5 MDA-MB-231 cells is shown. **(c)** Relative number of particles normalised to 2.5×10^6 MDA-MB-231 cells is depicted. **(d)** Western blot analysis of EV enriched proteins Alix and TSG101 in EV samples obtained from 15×10^6 MDA-MB-231 cells. Representative image shown, *n=3*. **(e)** Relative amount of EV protein normalised to 10^5 4T1 cells is shown. **(f)**

Relative number of particles normalised to 2.5 x 10^6 4T1 cells is depicted. **(g)** Western blot analysis of EV enriched proteins Alix in EV samples obtained from 15 x 10^6 4T1 cells. Representative image shown, *n=3*. **(h)** Relative number of particles normalised to 2.5 x 10^6 4T1 cells is depicted. All data are represented as mean ± s.e.m. *n=3,* statistical significance was determined by paired two-tailed t-test. EVs were isolated by protocol 1 (**b-g**) and 2 (**h**).

Figure 1

a

Cell culture conditions defined by Fonseka *et al.* (Mathivanan group)

Cells at 70-80% confluency
Three technical replicates per biological replicate
Media - 10% FBS
Volume - 15 mL/plate

Wash the cells twice with 1X PBS to remove any traces of spent media

Drug treatment for 24 h
Media - **serum free media**
Volume - 15 mL/plate

Harvest conditioned media containing EVs secreted in 24 h during drug treatment
Count the live cell number for normalization
Subject the conditioned media to EV isolation steps

Cell culture conditions defined by Lee *et al.* (Baek group)

Seed cells 3-5 x 10^6 cells per plate
Media - 10% FBS
Volume - 20 mL/plate

Wash the cells twice with 1X PBS to remove any traces of spent media

Drug treatment for 24 h
Media - **serum free media**
Volume - 20 mL/plate

Harvest and pool conditioned media containing EVs secreted in 24 h during drug treatement and subject to isolations steps

b

Baek group		Mathivanan group
Protocol 1 Im *et al.* / Fonseka *et al.*	**Protocol 2** Lee *et al.* / Fonseka *et al.*	**Protocol 3**

Protocol 1 (Baek group)

Conditioned media post drug treatment for 24 h in **serum free media**
300 × g 3 min
Transfer supernatant to a fresh tube
Remove cells
2500 × g 20 min
Transfer supernatant to a fresh tube
Remove cell debris
10,000 × g 30 min
Transfer supernatant to a fresh tube and filter (0.2 μm)
Remove large EVs
120,000 × g 90 min
Resuspend sEV pellet in 1 × PBS
120,000 × g 90 min
Resuspend purified sEV pellet in 1 × PBS
sEV → PBS → NTA analysis

Protocol 2 (Baek group)

Conditioned media post drug treatment for 24 h in **serum free media**
300 × g 3 min
Transfer supernatant to a fresh tube
Remove cells
* **2500 × g 15 min**
Transfer supernatant to a fresh tube
Remove cell debris
10,000 × g 30 min
Transfer supernatant to a fresh tube and filter (0.2 μm)
Remove large EVs
120,000 × g 90 min SW 28 rotor
Dry sEV pellet and resuspend in 1 × PBS
* sEV pellet incubated for 1 h at 4°C
sEV, 4 °C, 1 h → PBS → NTA analysis

Protocol 3 (Mathivanan group)

Conditioned media post drug treatment for 24 h in **media with EV depleted serum**
500 × g 10 min
Transfer supernatant to a fresh tube
Remove cells and cell debris
2000 × g 20 min
Transfer supernatant to a fresh tube
Remove cells and cell debris
10,000 × g 30 min SW 28 rotor
Transfer supernatant to a fresh tube
Remove large EVs
100,000 × g 60 min SW 28 rotor
Resuspend sEV pellet in 1 × PBS
100,000 × g 60 min TLA 55 rotor
Purified sEV pellet resuspended in 1 × PBS
sEV → PBS → NTA analysis

Figure 2