Functional Organization of the Maternal and Paternal Human 4D Nucleome

Stephen Lindsly,[1] Wenlong Jia,[2] Haiming Chen,[1] Sijia Liu,[3] Scott Ronquist,[1] Can Chen,[4,7] Xingzhao Wen,[5] Gabrielle A. Dotson,[1] Charles Ryan,[1,8,9] Gilbert S. Omenn,[1,10] Shuai Cheng Li,[2] Max Wicha,[1,6] Alnawaz Rehemtulla,[6] Lindsey Muir,[1] Indika Rajapakse[1,7,*]

---

[1]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI 48109 USA

[2]Department of Computer Science, City University of Hong Kong, Kowloon, Hong Kong

[3]MIT-IBM Watson AI Lab, IBM Research, Cambridge, MA 02142 USA

[4]Department of Electrical Engineering and Computer Science, University of Michigan, Ann Arbor 48109 USA

[5]Department of Bioengineering, University of California San Diego, La Jolla, CA 92093 USA

[6]Department of Hematology/Oncology, University of Michigan, Ann Arbor, MI 48109 USA

[7]Department of Mathematics, University of Michigan, Ann Arbor, MI 48109 USA

[8]Medical Scientist Training Program, University of Michigan, Ann Arbor, MI 48109 USA

[9]Program in Cellular and Molecular Biology, University of Michigan, Ann Arbor, MI 48109 USA

[10]Departments of Internal Medicine, Human Genetics, and School of Public Health, University of Michigan, Ann Arbor, MI 48109, USA

*To whom correspondence should be addressed; E-mail: indikar@umich.edu.

Abstract

Every human somatic cell inherits a maternal and paternal genome, which work together to give rise to cellular phenotypes. However, the allele-specific relationship between gene expression and genome structure through the cell cycle is largely unknown. By integrating haplotype-resolved genome-wide chromosome conformation capture, mature and nascent mRNA, and protein binding data, we investigate this relationship both globally and locally. We introduce the maternal and paternal 4D Nucleome, enabling detailed analysis of the mechanisms and dynamics of genome structure and gene function for diploid organisms. Our analyses find significant coordination between allelic expression biases and local genome conformation, and notably absent expression bias in universally essential cell cycle and glycolysis genes. We propose a model in which coordinated biallelic expression reflects prioritized preservation of essential gene sets.

Introduction

The redundancy inherent in the diploid genome makes it robust to potentially harmful mutations. Dependence on a single allele, for example in genes with monoallelic or allele biased expression (MAE, ABE), may increase vulnerability to disease. In familial cancer syndromes, individuals who inherit genetic loss-of-function mutations have a single functional allele, referred to as loss of heterozygosity (LOH) (*1*). BRCA1 and BRCA2 are quintessential examples, where LOH leads to a significant increase in the likelihood of developing breast cancer (*2, 3*). Imprinted genes are also associated with multiple disease phenotypes such as Angelman and Prader-Willi syndromes (*4, 5*). These observations raise the possibility that other MAE or ABE genes are associated with disease, but the contribution of allelic bias to disease phenotypes remains poorly understood.

A major step towards understanding the contribution of allelic bias to disease is to identify ABE genes and gene modules, recognizing that functionally important biases may be transient and challenging to detect. Differences in allelic expression may result from single nucleotide variants (SNVs), insertions or deletions (InDels), and chromatin modifications (*6–10*). By some estimates, ABE occurs in 4-26% of expressed genes across tissues and individuals (*11*). In addition, higher order chromatin conformation and spatial positioning in the nucleus shape gene expression (*12–15*). Spatial positioning may lead to allele-specific expression differences, as the maternal and paternal alleles have been shown to be distant in the nucleus through techniques such as fluorescent in situ hybridization (FISH) (*16, 17*). Allele-specific expression and 3D structures are frequently not distinguished in genomics methods such as RNA-sequencing and Hi-C (genome-wide chromosome conformation capture), complicating interpretations of structure-function relationships. Complete phasing of the two genomes remains a significant challenge in human genetics.

To uncover ABE and its relationship to genome structure over time, we integrated allele-specific RNA-seq, Bru-seq, and Hi-C across three phases of the cell cycle in human B-lymphocytes (GM12878, **Figure 1**). The RNA-seq and Bru-seq data were separated into their maternal and

paternal components through SNVs/InDels. The Hi-C data were separated using our novel phasing algorithm, HaploHiC, which utilizes SNVs/InDels to impute Hi-C reads of unknown parental origin. We incorporated publicly available allele-specific protein binding data (ChIP-seq) to better understand the mechanisms behind the gene expression and genome structure relationship (*7, 18*). We performed a genome-wide investigation for gene expression biases, chromatin compartment switching, and topologically associated domain (TAD) boundary changes between alleles and cell cycle phases. Along with known XCI and imprinted genes, we found notable patterns of ABE across the cell cycle for hundreds of genes that are not well appreciated in current literature, and corresponding allelic biases in protein binding at many of these loci. Differentially expressed genes were significantly more likely to have changes in their local genomic structure than random allele-specific genes. Additionally, we observed a pronounced lack of ABE in crucial biological pathways and essential genes. Our integrated analysis represents an allele-specific extension of the 4D Nucleome (4DN), which describes the relationship between genome structure, gene expression, and cell phenotype through time (*19–21*). Taken together, our findings demonstrate the value of analyzing ABE in the context of corresponding structural and epigenetic alterations across the cell cycle. This approach will be critical in future investigations of human phenotypic traits and their penetrance, genetic diseases, vulnerability to complex disorders, and tumorigenesis.

Materials and Methods

Cell Culture and Cell Cycle Sorting

Human GM12878 cells were cultivated in RPMI1640 medium supplemented with 10% fetal bovine serum (FBS). Live cells were stained with Hoechst 33342 (Cat #B2261, Sigma-Aldrich), and then sorted by fluorescence-activated cell sorting (FACS) to obtain cell fractions at the corresponding cell cycle phases G1, S, and G2/M (**Figure S2**).

RNA-seq and Bru-seq Sequencing

Total RNA was extracted from sorted live cells for both RNA-seq and Bru-seq. We performed 5'-bromouridine (Bru) incorporation in live cells for 30 minutes, and the Bru-labeled cells were then stained on ice with Hoechst 33342 for 30 min before sorting at 4°C to isolate G1, S, and G2/M phase cells. The sorted cells were immediately lysed in TRizol (Cat # 15596026, ThermoFisher) and frozen. To isolate Bru-labeled RNA, DNAse-treated total RNA was incubated with anti-BrdU antibodies conjugated to magnetic beads (*22*). We converted the transcripts from the RNA-seq and Bru-seq experiments for all samples into cDNA libraries and deep-sequenced at 50-base length on an Illumina HiSeq2500 platform. The RNA-seq and Bru-seq data each consist of three biological replicates. From our RNA-seq replicates, we obtained a total of 193.4, 197.2, and 202.0 million raw reads for G1, S, and G2/M, respectively. From our Bru-seq replicates, we obtained a total of 162.5, 149.9, and 138.0 million raw reads for G1, S, and G2/M, respectively.

Hi-C Sequencing

For cells used in construction of Hi-C libraries, cells were crosslinked with 1% formaldehyde, the reaction was neutralized with 0.125 M glycine, then cells were stained with Hoechst 33342 and sorted into G1, S, and G2/M fractions. Cross-linked chromatin was digested with the restriction enzyme MboI for 12 hours. The restriction enzyme fragment ends were tagged with biotin-dATP and ligated in situ. After ligation, the chromatins were de-cross-linked, and DNA was isolated for fragmentation. DNA fragments tagged by biotin-dATP, in the size range of 300-500 bp, were pulled down for sequencing adaptor ligation and polymerase chain reaction (PCR) products. The PCR products were sequenced on an Illumina HiSeq2500 platform. Respective to G1, S, and G2/M, we obtained 512.7, 550.3, and 615.2 million raw Hi-C sequence reads.

RNA-seq and Bru-seq Data Processing

RNA-seq and Bru-seq analysis were performed as previously described (*23, 24*). Briefly, Bru-seq used Tophat (v1.3.2) to align reads without de novo splice junction calling after checking quality with FastQC (version 0.10.1). A custom gene annotation file was used in which introns are included but preference to overlapping genes is given on the basis of exon locations and stranding where possible (see (*24*) for full details). Similarly for RNA-seq data processing, the raw reads were checked with FastQC. Tophat (version 2.0.11) and Bowtie (version 2.1.0.0) were used to align the reads to the reference transcriptome (HG19). Cufflinks (version 2.2.1) was used for expression quantification, using UCSC hg19.fa and hg19.gtf as the reference genome and transcriptome. A locally developed R script using CummeRbund was used to format the Cufflinks output.

Separation of Maternal and Paternal RNA-seq and Bru-seq Data

To determine allele-specific transcription and gene expression through Bru-seq and RNA-seq, all reads were aligned using GSNAP, a SNV aware aligner (*25*). HG19 and USCS gene annotations were used for the reference genome and gene annotation, respectively. The gene annotations were used to create the files for mapping to splice sites (used with –s option). Optional inputs to perform SNV aware alignment were included. Specifically, –v was used to include the list of heterozygous SNVs and –use-sarray=0 was used to prevent bias against non-reference alleles.

After alignment, the output SAM files were converted to BAM files, sorted and indexed using SAMtools (*26*). SNV alleles were quantified using bam-readcounter which was used to count the number of each base that was observed at each of the heterozygous SNV locations. Allele-specificity of each gene was then assessed by combining all of the SNVs in each gene. For RNA-seq, only exonic SNVs were used. Bru-seq detects nascent transcripts containing both exons and introns, so both exonic and intronic SNVs were used. Maternal and paternal gene expression were calculated by multiplying the genes' overall read counts by the fraction of the

SNV-covering reads that were maternal and paternal, respectively. We identified 266,899 SNVs from the Bru-seq data, compared with only 65,676 such SNVs from RNA-seq data. However in the Bru-seq data, many SNVs have low read coverage depth. We required at least 5 SNV-covering reads for a SNV to be used to separate the maternal and paternal contributions to gene expression. This criterion found that there were similar numbers of informative SNVs (19,394 and 19,998) in the RNA-seq and Bru-seq data, respectively. Any genes which did not contain informative SNVs were divided equally into their maternal and paternal contributions to avoid losing the data.

Allele-specific Differential Expression

For a gene's expression to be considered for differential expression, we require each of the three replicates to have an average of 10 SNV-covering reads mapped at least one of the alleles in all three cell cycle phases. This threshold was found to be sufficient at preventing technical noise from influencing our differential expression results. From the 23,277 Refseq genes interrogated, there were 4,193 genes with at least 10 read counts mapped to either the maternal or paternal allele (or both) in the RNA-seq data. From Bru-seq, there were 5,294 genes using the same criterion. We refer to these genes as "allele-specific" genes. We observed that there were larger variances between samples and lower read counts in the Bru-seq data set than in RNA-seq. We identified differentially expressed genes between alleles and between cell cycle phases for both RNA-seq and Bru-seq using a MATLAB implementation of DESeq (*27*). To reduce the possibility of false positives when determining differential expression, we imposed a minimum FPKM level of 0.1, a false discovery rate adjusted p-value threshold of 0.05, and a fold change cutoff of FC > 2 for both RNA-seq and Bru-seq (*28*).

Separation of Maternal and Paternal Hi-C Data by HaploHiC

Hi-C library construction and Illumina sequencing were performed using established methods (*9*). In this study, we separate the maternal and paternal genomes' contributions to the Hi-C

contact matrices to analyze their similarities and differences in genome structure. In order to determine which Hi-C reads come from which parental origin, we utilize differences in genomic sequence at SNVs/InDels. As these variations are unique to the maternal and paternal genomes, they can be used to quickly and easily separate reads that cover them.

When attempting to separate the maternal and paternal genomes, complications arise when there are sections of DNA that are identical. Due to the cells' ability to correct errors that arise in their DNA, there is a relatively small number of variations. This means our segregated maternal and paternal contact matrices will be extremely sparse if we only separate the small subset of genomic loci with SNVs/InDels. In order to combat this problem, we look to reintroduce contacts of unknown parental origin.

Here we propose HaploHiC, a novel technique for phasing reads of unknown parental origin using a local imputation from known reads. Unlike previous haplotyping methods, HaploHiC uses a data-derived ratio based on the hypothesis that if the maternal and paternal genomes have different 3D structures, we can use the reads with known origin (at SNV/InDel loci) to predict the origin of neighboring unknown reads (**Figure 4A, Supplemental Methods A**) (*29*). For example, if we observe that many contacts between two loci can be directly mapped to the paternal genome but few to the maternal genome, then unphased contacts between those loci are more likely to be from the paternal genome as well.

HaploHiC marks paired-end reads as haplotype-known or -unknown depending on their coverage of heterozygous SNVs and InDels. Haplotype-known reads are directly assigned to their corresponding haplotype, maternal or paternal. HaploHiC uses a local contacts based algorithm to impute the haplotype of haplotype-unknown reads using nearby SNVs and InDels. If the necessary amount of SNVs and InDels for imputation is not reached, HaploHiC randomly assigns the haplotype-unknown reads to be maternal or paternal. Detailed materials and methods for haplotype phasing and Hi-C construction are provided in **Supplemental Methods A**.Evaluation in simulation data shows that HaploHiC performs well, with a 97.66% accuracy of predicting unknown origin reads (**Table S11**). The allelicly separated genomes can be seen

in **Figure 4B**. HaploHiC has been made available through a GitHub repository.

After haplotype assignment through HaploHiC, Hi-C paired-end reads (PE-reads) are distributed to intra-haplotype (P-P and M-M) and inter-haplotype (P-M and M-P) (**Supplemental Methods A.1-A.8**). Juicer is applied on intra-haplotype PE-reads, and outputs maternal and paternal contact matrices which are normalized through the Knight-Ruiz method of matrix balancing (*30, 31*). Inter-haplotype contact matrices are generated by HaploHiC (**Supplemental Methods A.7**). Inter-haplotype contacts shown in **Figure 4B** and **Figure S8**. Both base pair level and fragment level matrices are constructed. The resolution of base pair level matrices are 1 Mb and 100 kb. Gene-level contacts are converted from fragment level matrices by HaploHiC.

Results

Chromosome-Scale Maternal and Paternal Differences

Spatial positioning of genes within the nucleus is known to be associated with transcriptional status (*12–15*). One might expect that the maternal and paternal copies of each chromosome would stay close together to ensure that their respective alleles have equal opportunities for transcription. Imaging of chromosome territories has shown that this is often not the case (**Figure 2A**, **Supplemental Methods B**) (*32–34*). This observation inspired us to investigate whether the two genomes operate in a symmetric fashion, or if allele-specific differences exist between the genomes regarding their respective chromatin organization patterns and gene expression profiles. We analyzed parentally phased whole-chromosome Hi-C and RNA-seq data at 1 Mb resolution to identify allele-specific differences in both chromosome structure and function, respectively. We subtracted each chromosome's paternal Hi-C matrix from the maternal matrix and found the Frobenius norm of the resulting difference matrix. The Frobenius norm here provides a measure of distance between matrices, where equivalent parental genome structures would result in a value of zero. Similarly, we subtracted the phased RNA-seq vectors in log_2 scale and found the Frobenius norm of each difference vector. The Frobenius norms were ad-

justed for chromosome size and normalized for both Hi-C and RNA-seq.

We found that all chromosomes have allelic differences in both structure (Hi-C, blue) and function (RNA-seq, red) (**Figure 2B**). Chromosome X had the largest structural difference, as expected, followed by Chromosomes 9, 21, and 14. Chromosome X had the most extreme functional difference as well, followed by Chromosomes 13, 7, and 9. A threshold was assigned at the median Frobenius norm for Hi-C and RNA-seq (**Figure 2B** green dashed lines). The majority of chromosomes with larger differences than the median for Hi-C are also larger than the median in RNA-seq. We found that there is a positive correlation between chromosome level differences in structure and function, which is statistically significant only when the X Chromosome is included (R = 0.66 and p < 0.05 with the X Chromosome; R = 0.30 and p = 0.17 without the X Chromosome).

Allele-Specific RNA Expression

After confirming broad allelic differences in RNA expression at the chromosomal scale, we examined allele-specific expression of individual genes through RNA-seq and Bru-seq. We hypothesized that the chromosome scale expression differences were not only caused by known cases of ABE such as XCI and imprinting, but also by widespread ABE over many genes. Therefore, we evaluated all allele-specific genes (sufficient reads covering SNVs/InDels) for differential expression across the six settings: maternal and paternal in G1, S, and G2/M (G2/M referred to as G2 for brevity). These settings give rise to seven comparisons which consist of maternal versus paternal within each of the cell cycle phases (three comparisons), as well as G1 versus S and S versus G2 for the maternal and paternal genomes, respectively (two comparisons for each genome).

First, we identified genes with ABE and allele-specific cell cycle-biased expression (CBE) from RNA-seq. From 23,277 RefSeq genes interrogated, there were 4,193 genes with sufficient coverage on SNVs/InDels to reliably determine allele-specific expression (*35*). We performed differential expression analysis for the seven comparisons to identify which of the 4,193 genes

had ABE or CBE (*27*). We identified 615 differentially expressed genes from RNA-seq: 467 ABE genes, 229 CBE genes, and 81 genes with both ABE and CBE (**Tables S2**, **S4**). Both exons and introns containing informative SNVs/InDels were used for our Bru-seq data, from which 5,294 genes had sufficient coverage. We identified 505 differentially expressed genes from Bru-seq: 380 ABE genes, 164 CBE genes, and 39 genes with both ABE and CBE (**Tables S3**, **S5**). We then separated the differentially expressed genes into their respective chromosomes to observe their distribution throughout the genome. From RNA-seq (Bru-seq), we found that autosomes had 3-14% ABE (1-11%) in their allele-specific genes which is comparable to previous findings (*11*). As expected, the X Chromosome had a particularly high percentage of ABE genes at 90% (91%).

We identified 288 genes that have ABE in all three cell cycle phases from RNA-seq (160 paternally biased, 128 maternally biased) and 173 from Bru-seq (129 paternal biased, 44 maternally biased). This is the most common differential expression pattern among ABE genes and these genes form the largest clusters in **Figure 3A**. These clusters include, but are not limited to, XCI, imprinted, and other MAE genes. Known examples within these clusters are highlighted in the 'X-Linked' and 'Imprinted' sections of **Figure 3B**. We also identified hundreds of ABE genes that are not currently appreciated in literature, shown in the 'Autosomal Genes' sections of **Figure 3B** for both mature and nascent RNA. Approximately half of all ABE genes were only differentially expressed in one or two cell cycle phases, or transient allelic biases, leading to the many smaller clusters seen in **Figure 3A**. Examples of genes with transient allelic biases are also presented in the 'Autosomal Genes' section of **Figure 3B**. Transient expression biases like these may be due to coordinated expression of the two alleles in only certain cell cycle phases, though the mechanism behind this behavior is unclear.

Among the ABE genes from RNA-seq analysis, we found 117 MAE genes. In addition to the requirements for differential expression, we impose the thresholds of a FC ≥ 10 and for the inactive allele to have < 0.1 Fragments Per Kilobase of transcript per Million (FPKM), or FC ≥ 50 across all three cell cycle phases. Our analysis confirmed MAE for imprinted and XCI

genes, with examples shown in **Figure 3B**. Imprinted and XCI genes are silenced via transcriptional regulation, which was verified by monoallelic nascent RNA expression (Bru-seq). The *XIST* gene, which is responsible for XCI, was expressed in the maternal allele reflecting the deactivation of the maternal X Chromosome. XCI was also observed from Hi-C through large heterochromatic domains in the maternal X Chromosome, and the absence of these domains in the paternal X Chromosome (**Figure 4C**). The inactive X Chromosome in our cells is opposite of what is commonly seen for the GM12878 cell line in literature (likely due to our specific GM12878 sub-clone), but is consistent between our data modalities (*9, 10, 18*). The MAE genes also include six known imprinted genes, four expressed from the paternal allele (*KCNQ1OT1*, *SNRPN*, *SNURF*, and *PEG10*) and two from the maternal allele (*NLRP2* and *HOXB2*). Some of the known imprinted genes that were confirmed in our data are associated with imprinting diseases, such as Beckwith-Wiedemann syndrome (*KCNQ1OT1* and *NLRP2*), Angelman syndrome (*SNRPN* and *SNURF*), and Prader-Willi syndrome (*SNRPN* and *SNURF*) (*36, 37*). These genes and their related diseases offer further support for allele-specific analysis, as their monoallelic expression could not be detected otherwise.

After observing that approximately half of all ABE genes had transient expression biases, we hypothesized that alleles may have unique dynamics through the cell cycle. We then focused our investigation on allele-specific gene expression through the cell cycle (**Figure S1B**). We compared the expression of each allele between G1 and S as well as between S and G2, which provides insight into the differences between maternal and paternal alleles' dynamics across the cell cycle. In the G1 to S comparison, there are 88 (55) genes in RNA-seq (Bru-seq) which have similar expression dynamics in both alleles. These genes' maternal and paternal alleles are similarly upregulated or downregulated from G1 to S. In contrast, 87 (97) genes in RNA-seq (Bru-seq) have different expression dynamics between alleles. That is, only one allele is up or downregulated in the transition from G1 to S. In the S to G2 comparison, there are 26 (3) genes in RNA-seq (Bru-seq) that have similar expression dynamics in both alleles and 56 (12) genes with different expression dynamics between alleles. From these data, we see a coordination of

expression between many, but certainly not all, alleles through the cell cycle.

Biallelic Expression and Cellular Function

We observed from our analysis of CBE that multiple cell cycle regulatory genes had no instances of ABE (**Figure 3C**). We expanded this set of genes, or biological module, to include all allele-specific genes contained in the KEGG cell cycle pathway (*38*). Again, we observed zero instances of ABE. We hypothesized that genes with crucial cellular functions, like cell cycle regulation, may have coordinated biallelic expression to ensure their sufficient presence. This hypothesis was further motivated by previous findings which showed restricted genetic variation of enzymes in the essential glycolytic pathway (*39*).

We tested additional modules derived from KEGG pathways containing at least five allele-specific genes, with the circadian rhythm module supplemented by a known core circadian gene set (*19*). Examples of modules with varying proportions of ABE are shown in **Table 1** (**Table S7**), where "Percent ABE" refers to the proportion of genes with ABE to the total number of allele-specific genes in that module. We found that there are multiple crucial modules, including the glycolysis/gluconeogenesis and pentose phosphate pathways, which also had zero ABE genes. To explore the possibility of a global phenomenon by which genes essential to cellular fitness are significantly less likely to have biased expression, we analyzed the frequency of ABE in 1,734 genes experimentally determined to be essential in human cells (*40*). Using the 662 allele-specific genes in this set, we found that these essential genes were significantly less likely to have ABE than a random selection of allele-specific genes (5.8% versus 11.1%, p $<$ 0.001, **Supplemental Methods D**), consistent with our hypothesis that critical genes are likely to be expressed by both alleles. In total, we offer a model in which coordinated biallelic expression reflects prioritized preservation of essential gene sets.

Allele-Specific Genome Structure

Motivated by our observations of chromosome level structural differences between the maternal and paternal genomes, we examined the HaploHiC separated data in more detail to determine where these differences reside. The genome is often categorized into two compartments: transcriptionally active euchromatin and repressed heterochromatin. In studies comparing multiple types of cells or cells undergoing differentiation, areas of euchromatin and heterochromatin often switch corresponding to genes that are activated/deactivated for the specific cell type (*41*). We explored this phenomenon in the context of the maternal and paternal Hi-C matrices to determine if the two genomes had differing chromatin compartments. Chromatin compartments can be identified from Hi-C data using methods such as principal component analysis or spectral clustering (**Supplemental Methods C**) (*42*). We applied spectral clustering to every chromosome across all three cell cycle phases at 1 Mb resolution. We found that there were slight changes in chromatin compartments for all chromosomes, but the vast majority of these changes took place on the borders between compartments rather than an entire region switching compartments. These border differences were not enriched for ABE genes. This implies that, although the structures are not exactly the same, the maternal and paternal genomes have similar overall compartmentalization (aside from the X Chromosome).

We next applied spectral clustering recursively to the Hi-C data at 100 kb resolution to determine whether there were differences in TADs between the two genomes throughout the cell cycle (*42*). While the current understanding of genomic structure dictates that TAD boundaries are invariant (between alleles, cell types, etc), it is also known that "intra-TAD" structures are highly variable (*41, 43, 44*). The spectral identification method has an increased ability to discern these subtle structural changes. We found that TAD boundaries were variable between the maternal and paternal genomes and across cell cycle phases in all chromosomes. This supports previous findings of allelic differences in TADs for single cells, and we predict that they are even more variable across cell types (*42, 44*). Differences in TAD boundaries were observed surrounding MAE genes, ABE genes, and genes with coordinated biallelic expression

(**Figure S6**). This indicated that changes in TAD boundaries were not directly related to allelic expression differences.

Although we did not find a significant relationship between TAD boundary differences and ABE genes, we observed during this analysis that the local genome structure around the six imprinted genes had noticeable differences. We then sought to analyze all genes with ABE or CBE to find out if they had corresponding structural differences at a local level. We analyzed the local Hi-C matrices for each of the 615 RNA-seq and 505 Bru-seq differentially expressed genes. Using a 300 kb flanking region centered at the 100 kb bin containing the transcription start site (TSS), we isolated a 7x7 matrix (700 kb) for each differentially expressed gene (**Figure 5A, B**). These matrices represent the local genomic structure of the differentially expressed genes, and are slightly smaller than average TAD size. We then compared the correlation matrices of the log_2-transformed local Hi-C data and determined whether or not the matrices have statistically significant differences ($p < 0.05$) (*45, 46*). We applied this comparison to all genes that were differentially expressed in RNA-seq (Bru-seq) and found that 515 (403) genes had at least one comparison in which both the expression and local structure had significantly changed. While changes in local genome structure and changes in gene expression do not have a one-to-one relationship, we found that both ABE and CBE genes are more likely to have significant architectural differences than randomly sampled allele-specific genes ($p < 0.01$) (**Figure S7** and **Supplemental Methods D**). This lends further support to the idea that there is a relationship between allele-specific differences in gene expression and genome structure.

Allele-Specific Protein Binding

To uncover the mechanisms behind the relationship between allele-specific gene expression and genome structure, we looked to DNA binding proteins such as RNA polymerase II (Pol II), CCCTC-binding factor (CTCF), and 35 other transcription factors. We used publicly available protein binding data from AlleleDB in tandem with RNA-seq and found 114 genes that have an allelic bias in both gene expression and binding of at least one such protein (*18*). We identified

13 genes which have ABE and biased binding of Pol II, with bias agreement in 11 cases (85%). That is, the allelic expression and Pol II binding were biased toward the same allele. For CTCF, 33 of 72 cases have bias agreement (46%), and for all other transcription factors analyzed, 20 of 29 cases have bias agreement (69%) (**Table S6**). The CTCF binding bias agreement of around 50% is expected, based on previous studies (*7*). This is likely due to CTCF's role as an insulator, since an allele could be expressed or suppressed by CTCF's presence depending on the context. To avoid potential inconsistencies between our data and the protein data from AlleleDB, we excluded the X Chromosome when testing for ABE and protein binding biases.

We analyzed the 11 genes with allelic expression and Pol II binding bias agreement further to determine if they also had significant changes in local genome structure. Through local Hi-C comparisons, we found that all 11 of these genes had significant changes in structure in at least one cell cycle phase. 3D models for 6 of these 11 genes are shown in **Figure 5B**, which highlight differences in local genome structure (**Supplemental Methods E**) (*47*). The genes with bias agreement and changes in local genome structure include known imprinted genes such as *SNURF* and *SNRPN*, as well as genes with known allele-specific expression (and suggested imprinting in other cell types) like *ZNF331* (*48*). Additionally, there are multiple genes with known associations with diseases or disorders such as *BMP8A*, *CRELD2*, and *NBPF3* (*49–51*). These findings suggest that changes in local structure often coincide with changes in expression due to the increased or decreased ability of a gene to access the necessary transcriptional machinery within transcription factories. We visualize this relationship for the gene *CRELD2* as an example (**Figure 5C**) (*14*).

The Maternal and Paternal 4D Nucleome

We define the maternal and paternal 4DN as the integration of allele-specific genome structure with gene expression data through time. Many complex dynamical systems are investigated using a network perspective (*52, 53*). A network is a simplified representation of a system, which captures patterns of interactions between its components and how those interactions change

over time. We can consider genome structure as a network, since Hi-C data captures interactions between genomic loci (*12, 15*). In network science, the relative importance of a node in a network is commonly determined using network centrality (*52*). For Hi-C data, we treat genomic loci as nodes and use network centrality to measure the importance of each locus (*46, 54*). We performed a global analysis of the maternal and paternal 4DN by combining RNA-seq with multiple network centrality measures. We found differences between the maternal and paternal genomes and across cell cycle phases, but only the X Chromosome had clear maternal and paternal separation (**Figures S9**, **S10**).

In our earlier analysis, we found a significant relationship between ABE and changes in local genome structure. We also observed that genes in multiple critical biological modules had coordinated biallelic expression. Motivated by these results, we performed a local analysis of structure and function together to determine their allele-specific dynamics. We used the eigenvector centrality (similar to Google's PageRank) to quantify the structure of each sub-network and the average FPKM of the three RNA-seq replicates to quantify the function of each allele in the sub-network (*55*). We utilize the concept of a phase plane to plot the 4DN (4DN phase plane) (**Figure 6**) (*19*). We designate one axis as a measure of structure and the other as a measure of function. Coordinates of each point in the 4DN phase plane are determined from structure data (x-axis, eigenvector centrality derived from Hi-C) and function data (y-axis, FPKM). The 4DN phase plane contains three points for each allele, which represent G1, S, and G2. We define the difference between the maternal and paternal alleles in the 4DN phase plane as the allelic divergence (**Supplemental Methods D**).

We show four phase planes of gene sub-networks with various allelic divergences. Genes which are known to be crucial for cell cycle regulation have a very small average allelic divergence of 0.02 (**Figure 6B**, left). This highlights the robust coordination between the alleles to maintain proper cell cycle function. Given that GM12878 is a B-lymphocyte cell line, we were interested in the allelic divergence of genes which are related to B cell receptor functionality. We found that these genes have a moderate allelic divergence of 0.61 (**Figure 6B**, middle-left). This

is slightly smaller than a randomly selected set of allele-specific genes whose average allelic divergence was 0.65 (**Figure 6B**, middle-right). MAE genes have large average allelic divergence of 4.70, as expected (**Figure 6B**, right). While this analysis does not offer a statistical significance for structural and functional differences, it represents a broadly applicable approach for understanding the similarities and differences of the maternal and paternal genomes through time. This analysis can be applied to any cell type with allele-specific time-series or cell cycle sorted data.

Discussion

In this study, we present evidence for the intimate relationship among allele-specific gene expression, genome structure, and protein binding across the cell cycle. We validated our data and methods using known allele-specific properties such as the monoallelic expression of imprinted and X-linked genes, broad similarities of chromatin compartments between the maternal and paternal genomes, and large heterochromatic domains of the X Chromosome (*9, 56–59*). Unique to this study, we established a coordination of allele-biased expression and changes in local genome structure, which included hundreds of genes not commonly associated with allele-biased expression. In addition, we offer mechanistic evidence of this coordination through allele-specific protein binding biases. Through the integration of these structural and functional data through time, we developed an extension of the 4DN for the maternal and paternal genomes.

Through our analysis of mature (nascent) RNA, we found 467 (380) genes to be differentially expressed between the two alleles and 229 (164) genes with differential expression through the cell cycle. Approximately half of the genes with allele-biased expression are only differentially expressed in certain cell cycle phases, and over half of the genes with CBE are only differentially expressed in one allele or the other. Further research is needed to explore why certain genes have coordinated cell cycle dynamics across both alleles, while other genes have disparate expression in some cell cycle phases. We predict that these transient allelic bi-

ases may be associated with developmental pathologies and tumorigenesis, similar to imprinted and other MAE genes. Conversely, we found no allele-biased expression from genes in multiple biological modules, such as the cell cycle and glycolysis pathways (Table 1). We are not able to establish a statistical significance here due to the limited number of allele-specific genes in these modules, so we surveyed a set of 662 essential genes and found that they are significantly less likely to have allele-biased expression. This supports our hypothesis of highly coordinated biallelic expression in universally essential genes.

We developed a novel phasing algorithm, HaploHiC, which uses Hi-C reads mapped to SNVs/InDels to predict nearby reads of unknown parental origin. This allowed us to decrease the sparsity of our allele-specific contact matrices and increase confidence in our analysis of the parental differences in genome structure. While we found that the overall compartmentalization (euchromatin and heterochromatin) of the two genomes was broadly similar, there were many differences in TAD boundaries and local genome structure between the two genomes and between cell cycle phases. We focused our search of allele-specific differences in genome structure by calculating the similarity of local contacts surrounding the TSS of differentially expressed genes (*45, 46*). We found that differentially expressed genes were significantly more likely to have corresponding changes in local genome structure than random allele-specific genes.

We incorporated publicly available allele-specific protein binding data for Pol II and CTCF to explore the mechanisms behind the gene expression and local genome structure relationship (*7*). In genes that had both allele-biased expression and Pol II binding biases, we found that 85% of these genes had allelic bias agreement. Additionally, we found that these 11 genes with bias agreement also had significant changes in local genome structure. Analysis of the relationships among allele-specific gene expression, genome structure, and protein binding is currently limited by the amount of information available. The ability to separate maternal and paternal gene expression and protein binding is dependent on the presence of a SNV/InDel within the gene body and nearby protein binding motifs. As SNVs/InDels are relatively rare in

the human genome, the number of genes available to study is severely limited. Once we are able to separate the maternal and paternal genomes through advances in experimental techniques, we will be able to fully study these relationships.

Overall, these data support an intimate allele-specific relationship between genome structure and function, coupled through allele-specific protein binding. Changes in genome structure, influenced by the binding of proteins such as CTCF, can affect the ability of transcription factors and transcription machinery to access DNA. This results in changes in the rate of transcription of RNA, captured by Bru-seq. The rate of transcription leads to differential steady state gene expression, captured by RNA-seq. Integration of these data into a comprehensive computational framework led to the development of a maternal and paternal 4DN, which can be visualized using phase planes and quantified using allelic divergence. Allele-specific analysis across the cell cycle will be imperative to discern the underlying mechanisms behind many diseases by uncovering potential associations between deleterious mutations and allelic bias, and may have broad translational impact spanning cancer cell biology, complex disorders of growth and development, and precision medicine.

Description of Supplemental Data

Supplemental Data include additional methods details, 10 figures, and 11 tables.

Declaration of Interests

The authors declare no competing financial interests.

Acknowledgements

We thank the University of Michigan Sequencing Core members for producing high quality Bru-seq, RNA-seq, and Hi-C data. We thank Michele Paulson and Dr. Mats Ljungman for

generating the Bru-seq sequencing libraries and for helpful discussion. We also thank Dr. Jacob Kitzman for providing the GM12878 cell line and assisting in resolving genotype-phasing issues of the cell line. We thank Sam Dilworth and the iReprogram team for processing and formatting the KEGG pathway data. We are grateful to Dr. Thomas Ried for helpful reading of the manuscript. We thank the Air Force Office of Scientific Research (Award No: FA9550-18-1-0028), and the Forbes Institute research fund for supporting our work.

Web Resources

Prototype MATLAB implementation for differential expression analysis can be found here: Identifying Differentially Expressed Genes from RNA-Seq Data. HaploHiC code is available through a GitHub repository: https://github.com/Nobel-Justin/HaploHiC.

Data and Code Availability

The RNA-seq, Bru-seq, and Hi-C data used for this study are available through the Gene Expression Omnibus (GEO) database, https://www.ncbi.nlm.nih.gov/geo/ (accession number GSE159813).

References

1. Alfred G Knudson. Mutation and cancer: statistical study of retinoblastoma. *Proceedings of the National Academy of Sciences*, 68(4):820–823, 1971.

2. Julius Gudmundsson, Gudrun Johannesdottir, Jon Thor Bergthorsson, Adalgeir Arason, Sigurdur Ingvarsson, Valgardur Egilsson, and Rosa Björk Barkardottir. Different tumor types from brca2 carriers show wild-type chromosome deletions on 13q12–q13. *Cancer research*, 55(21):4830–4832, 1995.

3. Kara N Maxwell, Bradley Wubbenhorst, Brandon M Wenz, Daniel De Sloover, John Pluta, Lyndsey Emery, Amanda Barrett, Adam A Kraya, Ioannis N Anastopoulos, Shun Yu, et al. Brca locus-specific loss of heterozygosity in germline brca1 and brca2 carriers. *Nature communications*, 8(1):1–11, 2017.

4. Irina S Zakharova, Alexander I Shevchenko, and Suren M Zakian. Monoallelic gene expression in mammals. *Chromosoma*, 118(3):279–290, 2009.

5. Karin Buiting. Prader–willi syndrome and angelman syndrome. In *American Journal of Medical Genetics Part C: Seminars in Medical Genetics*, volume 154, pages 365–376. Wiley Online Library, 2010.

6. 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*, 526(7571):68, 2015.

7. Joel Rozowsky, Alexej Abyzov, Jing Wang, Pedro Alves, Debasish Raha, Arif Harmanci, Jing Leng, Robert Bjornson, Yong Kong, Naoki Kitabayashi, et al. Alleleseq: analysis of allele-specific expression and binding in a network framework. *Molecular systems biology*, 7(1):522, 2011.

8. Anshul Kundaje, Wouter Meuleman, Jason Ernst, Misha Bilenky, Angela Yen, Alireza Heravi-Moussavi, Pouya Kheradpour, Zhizhuo Zhang, Jianrong Wang, Michael J Ziller, et al. Integrative analysis of 111 reference human epigenomes. *Nature*, 518(7539):317, 2015.

9. Suhas SP Rao, Miriam H Huntley, Neva C Durand, Elena K Stamenova, Ivan D Bochkov, James T Robinson, Adrian L Sanborn, Ido Machol, Arina D Omer, Eric S Lander, et al. A 3d map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell*, 159(7):1665–1680, 2014.

10. Longzhi Tan, Dong Xing, Chi-Han Chang, Heng Li, and X Sunney Xie. Three-dimensional genome structures of single diploid human cells. *Science*, 361(6405):924–928, 2018.

11. Danny Leung, Inkyung Jung, Nisha Rajagopal, Anthony Schmitt, Siddarth Selvaraj, Ah Young Lee, Chia-An Yen, Shin Lin, Yiing Lin, Yunjiang Qiu, et al. Integrative analysis of haplotype-resolved epigenomes across human tissues. *Nature*, 518(7539):350, 2015.

12. Indika Rajapakse and Mark Groudine. On emerging nuclear order. *Journal of Cell Biology*, 192(5):711–721, 2011.

13. Tom Misteli. The inner life of the genome. *Scientific American*, 304(2):66, 2011.

14. Peter R Cook. A model for all genomes: the role of transcription factories. *Journal of molecular biology*, 395(1):1–10, 2010.

15. Tom Misteli. The self-organizing genome: Principles of genome architecture and function. *Cell*, 2020.

16. Brian J Beliveau, Alistair N Boettiger, Maier S Avendaño, Ralf Jungmann, Ruth B McCole, Eric F Joyce, Caroline Kim-Kiselak, Frédéric Bantignies, Chamith Y Fonseka, Jelena Erceg, et al. Single-molecule super-resolution imaging of chromosomes and in situ haplotype visualization using oligopaint fish probes. *Nature communications*, 6(1):1–13, 2015.

17. Thomas Cremer and Marion Cremer. Chromosome territories. *Cold Spring Harbor perspectives in biology*, 2(3):a003889, 2010.

18. Jieming Chen, Joel Rozowsky, Timur R Galeev, Arif Harmanci, Robert Kitchen, Jason Bedford, Alexej Abyzov, Yong Kong, Lynne Regan, and Mark Gerstein. A uniform survey of allele-specific binding and expression over 1000-genomes-project individuals. *Nature communications*, 7:11101, 2016.

19. Haiming Chen, Jie Chen, Lindsey A Muir, Scott Ronquist, Walter Meixner, Mats Ljungman, Thomas Ried, Stephen Smale, and Indika Rajapakse. Functional organization of the

human 4d nucleome. *Proceedings of the National Academy of Sciences*, 112(26):8002–8007, 2015.

20. Thomas Ried and Indika Rajapakse. The 4d nucleome. *Methods (San Diego, Calif.)*, 123:1, 2017.

21. Job Dekker, Andrew S Belmont, Mitchell Guttman, Victor O Leshyk, John T Lis, Stavros Lomvardas, Leonid A Mirny, Clodagh C O'shea, Peter J Park, Bing Ren, et al. The 4d nucleome project. *Nature*, 549(7671):219–226, 2017.

22. Erez Lieberman-Aiden, Nynke L Van Berkum, Louise Williams, Maxim Imakaev, Tobias Ragoczy, Agnes Telling, Ido Amit, Bryan R Lajoie, Peter J Sabo, Michael O Dorschner, et al. Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *science*, 326(5950):289–293, 2009.

23. Laura Seaman, Haiming Chen, Markus Brown, Darawalee Wangsa, Geoff Patterson, Jordi Camps, Gilbert S Omenn, Thomas Ried, and Indika Rajapakse. Nucleome analysis reveals structure–function relationships for colon cancer. *Molecular Cancer Research*, 15(7):821–830, 2017.

24. Michelle T Paulsen, Artur Veloso, Jayendra Prasad, Karan Bedi, Emily A Ljungman, Brian Magnuson, Thomas E Wilson, and Mats Ljungman. Use of bru-seq and bruchase-seq for genome-wide assessment of the synthesis and stability of rna. *Methods*, 67(1):45–54, 2014.

25. Thomas D Wu and Serban Nacu. Fast and snp-tolerant detection of complex variants and splicing in short reads. *Bioinformatics*, 26(7):873–881, 2010.

26. Heng Li, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth, Goncalo Abecasis, and Richard Durbin. The sequence alignment/map format and samtools. *Bioinformatics*, 25(16):2078–2079, 2009.

27. Simon Anders and Wolfgang Huber. Differential expression analysis for sequence count data. *Genome biology*, 11(10):R106, 2010.

28. Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.

29. Siddarth Selvaraj, Jesse R Dixon, Vikas Bansal, and Bing Ren. Whole-genome haplotype reconstruction using proximity-ligation and shotgun sequencing. *Nature biotechnology*, 31(12):1111, 2013.

30. Neva C Durand, Muhammad S Shamim, Ido Machol, Suhas SP Rao, Miriam H Huntley, Eric S Lander, and Erez Lieberman Aiden. Juicer provides a one-click system for analyzing loop-resolution hi-c experiments. *Cell systems*, 3(1):95–98, 2016.

31. Philip A Knight and Daniel Ruiz. A fast algorithm for matrix balancing. *IMA Journal of Numerical Analysis*, 33(3):1029–1047, 2013.

32. Irina Solovei, Antonio Cavallo, Lothar Schermelleh, Françoise Jaunin, Catia Scasselati, Dusan Cmarko, Christoph Cremer, Stanislav Fakan, and Thomas Cremer. Spatial preservation of nuclear chromatin architecture during three-dimensional fluorescence in situ hybridization (3d-fish). *Experimental cell research*, 276(1):10–23, 2002.

33. Andreas Bolzer, Gregor Kreth, Irina Solovei, Daniela Koehler, Kaan Saracoglu, Christine Fauth, Stefan Müller, Roland Eils, Christoph Cremer, Michael R Speicher, et al. Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes. *PLoS Biol*, 3(5):e157, 2005.

34. Scott Ronquist, Walter Meixner, Indika Rajapakse, and John Snyder. Insight into dynamic genome imaging: Canonical framework identification and high-throughput analysis. *Methods*, 123:119–127, 2017.

35. Nuala A O'Leary, Mathew W Wright, J Rodney Brister, Stacy Ciufo, Diana Haddad, Rich McVeigh, Bhanu Rajput, Barbara Robbertse, Brian Smith-White, Danso Ako-Adjei, et al. Reference sequence (refseq) database at ncbi: current status, taxonomic expansion, and functional annotation. *Nucleic acids research*, 44(D1):D733–D745, 2016.

36. Yang Cao, Susan S AlHumaidi, Eissa A Faqeih, Beth A Pitel, Patrick Lundquist, and Umut Aypar. A novel deletion of snurf/snrpn exon 1 in a patient with prader-willi-like phenotype. *European journal of medical genetics*, 60(8):416–420, 2017.

37. J Adams. Imprinting and genetic disease: Angelman, prader-willi and beckwith-weidemann syndromes. *Nat Educ*, 1(1), 2008.

38. Minoru Kanehisa and Susumu Goto. Kegg: kyoto encyclopedia of genes and genomes. *Nucleic acids research*, 28(1):27–30, 2000.

39. PT Wade Cohen, GS Omenn, AG Motulsky, S-H Chen, and ER Giblett. Restricted variation in the glycolytic enzymes of human brain and erythrocytes. *Nature New Biology*, 241(112):229–233, 1973.

40. Vincent A Blomen, Peter Májek, Lucas T Jae, Johannes W Bigenzahn, Joppe Nieuwenhuis, Jacqueline Staring, Roberto Sacco, Ferdy R van Diemen, Nadine Olk, Alexey Stukalov, et al. Gene essentiality and synthetic lethality in haploid human cells. *Science*, 350(6264):1092–1096, 2015.

41. Jesse R Dixon, Inkyung Jung, Siddarth Selvaraj, Yin Shen, Jessica E Antosiewicz-Bourget, Ah Young Lee, Zhen Ye, Audrey Kim, Nisha Rajagopal, Wei Xie, et al. Chromatin architecture reorganization during stem cell differentiation. *Nature*, 518(7539):331, 2015.

42. Jie Chen, Alfred O Hero III, and Indika Rajapakse. Spectral identification of topological domains. *Bioinformatics*, 32(14):2151–2158, 2016.

43. Jesse R Dixon, Siddarth Selvaraj, Feng Yue, Audrey Kim, Yan Li, Yin Shen, Ming Hu, Jun S Liu, and Bing Ren. Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature*, 485(7398):376, 2012.

44. Elizabeth H Finn, Gianluca Pegoraro, Hugo B Brandao, Anne-Laure Valton, Marlies E Oomen, Job Dekker, Leonid Mirny, and Tom Misteli. Extensive heterogeneity and intrinsic variation in spatial genome organization. *Cell*, 176(6):1502–1515, 2019.

45. James A Koziol, Joel E Alexander, Lance O Bauer, Samuel Kuperman, Sandra Morzorati, Sean J O'connor, John Rohrbaugh, Bernice Porjesz, Henri Begleiter, and John Polich. A graphical technique for displaying correlation matrices. *The American Statistician*, 51(4):301–304, 1997.

46. Scott Ronquist, Sijia Liu, Michael Perlman, and Indika Rajapakse. 4dnvestigator: a toolbox for the analysis of time series hi-c and rna-seq data. *bioRxiv*, 2020.

47. Nelle Varoquaux, Ferhat Ay, William Stafford Noble, and Jean-Philippe Vert. A statistical approach for inferring the 3d structure of the genome. *Bioinformatics*, 30(12):i26–i33, 2014.

48. Eyal Ben-David, Shahar Shohat, and Sagiv Shifman. Allelic expression analysis in the brain suggests a role for heterogeneous insults affecting epigenetic processes in autism spectrum disorders. *Human molecular genetics*, 23(15):4111–4124, 2014.

49. Fang-Ju Wu, Ting-Yu Lin, Li-Ying Sung, Wei-Fang Chang, Po-Chih Wu, and Ching-Wei Luo. Bmp8a sustains spermatogenesis by activating both smad1/5/8 and smad2/3 in spermatogonia. *Science signaling*, 10(477), 2017.

50. Yeawon Kim, Sun-Ji Park, Scott R Manson, Carlos AF Molina, Kendrah Kidd, Heather Thiessen-Philbrook, Rebecca J Perry, Helen Liapis, Stanislav Kmoch, Chirag R Parikh, et al. Elevated urinary creld2 is associated with endoplasmic reticulum stress–mediated kidney disease. *JCI insight*, 2(23), 2017.

51. Joseph Petroziello, Andrew Yamane, Lori Westendorf, Melissa Thompson, Charlotte Mc-Donagh, Charles Cerveny, Che-Leung Law, Alan Wahl, and Paul Carter. Suppression subtractive hybridization and expression profiling identifies a unique set of genes over-expressed in non-small-cell lung cancer. *Oncogene*, 23(46):7734–7745, 2004.

52. Mark Newman. *Networks*. Oxford university press, 2018.

53. Steven H Strogatz. Exploring complex networks. *nature*, 410(6825):268–276, 2001.

54. Sijia Liu, Haiming Chen, Scott Ronquist, Laura Seaman, Nicholas Ceglia, Walter Meixner, Pin-Yu Chen, Gerald Higgins, Pierre Baldi, Steve Smale, et al. Genome architecture mediates transcriptional control of human myogenic reprogramming. *iScience*, 6:232–246, 2018.

55. Lawrence Page, Sergey Brin, Rajeev Motwani, and Terry Winograd. The pagerank citation ranking: Bringing order to the web. Technical report, Stanford InfoLab, 1999.

56. Wolf Reik and Jörn Walter. Genomic imprinting: parental influence on the genome. *Nature Reviews Genetics*, 2(1):21, 2001.

57. Tomas Babak, Brian DeVeale, Emily K Tsang, Yiqi Zhou, Xin Li, Kevin S Smith, Kim R Kukurba, Rui Zhang, Jin Billy Li, Derek van der Kooy, et al. Genetic conflict reflected in tissue-specific maps of genomic imprinting in human and mouse. *Nature genetics*, 47(5):544, 2015.

58. Yael Baran, Meena Subramaniam, Anne Biton, Taru Tukiainen, Emily K Tsang, Manuel A Rivas, Matti Pirinen, Maria Gutierrez-Arcelus, Kevin S Smith, Kim R Kukurba, et al. The landscape of genomic imprinting across diverse adult human tissues. *Genome research*, 25(7):927–936, 2015.

59. Federico A Santoni, Georgios Stamoulis, Marco Garieri, Emilie Falconnet, Pascale Ribaux, Christelle Borel, and Stylianos E Antonarakis. Detection of imprinted genes by single-cell

664   allele-specific gene expression. *The American Journal of Human Genetics*, 100(3):444–

665   453, 2017.

 # Figure Titles and Legends

Fig. 1: Experimental and allelic separation workflow. Cell cycle sorted cells were extracted for RNA-seq, Bru-seq, and Hi-C (left to right, respectively). RNA-seq and Bru-seq data were allelicly phased via SNVs/InDels (left). SNV/InDel based imputation and haplotype phasing of Hi-C data using HaploHiC (right). These data provide quantitative measures of structure and function of the maternal and paternal genomes through the cell cycle.

Fig. 2: Genome imaging and chromosome differences. **(A)** Nucleus of a primary human fibroblast imaged using 3D FISH with the maternal and paternal copies of Chromosome 6, 8, and 11 painted red, green, and white, respectively (left). Subsection highlighting the separation between the maternal and paternal copies of Chromosome 11, now colored red (right). **(B)** Normalized chromosome level structural and functional parental differences of GM12878 cells. Structural differences (Δ Structure, blue) represent the aggregate changes between maternal and paternal Hi-C over all 1 Mb loci for each chromosome, adjusted for chromosome size in G1. Functional differences (Δ Function, red) represent the aggregate changes between maternal and paternal RNA-seq over all 1 Mb loci for each chromosome, adjusted for chromosome size in G1. Green dashed lines correspond to the median structural (0.48, chromosome 3) and functional (0.20, chromosome 6) differences, in the top and bottom respectively, and all chromosomes equal to or greater than the threshold are labeled. Scatter plot of maternal and paternal differences in structure and function with best-fit line (R = 0.66 and p < .05).

Fig. 3: Allele-specific mature and nascent RNA expression. **(A)** Differentially expressed genes' maternal and paternal RNA expression through the cell cycle. Expression heatmaps are average FPKM values over three replicates after row normalization. Genes are grouped by their differential expression patterns (**Figure S1**). **(B)** Representative examples of X-linked, imprinted, and other autosomal genes with allelic bias. Top shows mature RNA levels (RNA-seq) and bottom shows nascent RNA expression (Bru-seq) in both (A) and (B). **(C)** Examples of cell cycle regulatory genes' mature RNA levels through the cell cycle. These genes are grouped by their function in relation to the cell cycle and all exhibit CBE but none have ABE. All example genes in (B) and (C) reflect average FPKM values over three replicates, and ABE in a particular cell cycle phase is marked with an orange or purple asterisk for maternal or paternal bias, respectively. G2 includes both G2 and M phase.

Fig. 4: Haplotype phasing of Hi-C data. **(A)** HaploHiC separates paired-end reads into groups based on parental origin determined through SNVs/InDels (left, **Supplemental Methods A.3**). Reads are grouped by: (i) reads with one (sEnd-P/M) or both ends (dEnd-P/M) mapped to a single parent, (ii) reads are inter-haplotype, with ends mapped to both parents (d/sEnd-I), and (iii) reads with neither end mapped to a specific parent (dEnd-U) . An example of a paired-end read (dEnd-U) with no SNVs/InDels has its origin imputed using nearby reads (right, **Supplemental Methods A.6**). A ratio of paternally and maternally mapped reads is found in a dynamically sized flanking region around the haplotype-unknown read's location (**Supplemental Methods A.4**). This ratio then determines the likelihood of the haplotype-unknown read's origin. **(B)** Whole-genome Hi-C of GM12878 cells (top left). Inter- and intra-haplotype chromatin contacts after phasing Hi-C data using HaploHiC (right). Chromosomes 14 and 15 highlight inter- and intra-chromosome contacts within and between genomes (bottom left). Visualized in log_2 scale 1 Mb resolution in G1. **(C)** Haplotype phasing illustrates that the inactive maternal X Chromosome is partitioned into large heterochromatic domains, outlined in dotted black boxes. Visualized in log_2 scale 100 kb resolution in G1.

Fig. 5: Local chromatin structure and transcription factor binding. (**A**) Local regions around differentially expressed genes are tested for significant conformation changes. These regions are modeled to visualize the conformations around each allele through G1, S, and G2 (**Supplemental Methods E**). Example of local chromatin structure extraction is shown for *ZNF331* in G1 phase (center of blue box). Hi-C matrices are shown in log_2 scale 100 kb resolution. (**B**) 3D models of the local genome structure around 6 ABE genes with bias agreement of Pol II and significant changes in local genome structure. (**C**) Schematic representation of allele-specific Pol II and CTCF binding, with highlighted gene *CRELD2*, which had binding biases in both. Table shows extreme binding biases of Pol II and CTCF on *CRELD2* as an example.

Fig. 6: 4DN phase planes reveal a wide range of allelic divergences in gene sub-networks. **(A)** Workflow to obtain structure and function measures. Simplified phase planes are shown with linear relationships between changes in structure and function, changes in structure with no changes in function, and changes in function with no changes in structure. **(B)** 4DN phase planes of cell cycle genes, genes specific to B cell function, random allele-specific genes, and MAE genes, highlighting the similarities and differences between their alleles. Genes such as *BUB1B* and *PIK3AP1* have similar phase planes between alleles, while *RAC1* differs in structure and *WRAP73* differs in function. The bottom plot for each column combines the phase planes of the nine example genes.

 # Tables

Table 1: Allelic bias in biological modules.

Module	Percent ABE
Cell Cycle	0%
Glycolysis/Gluconeogenesis	0%
Pentose Phosphate	0%
BCR Signaling	8%
Circadian Rhythm	9%
p53 Signaling	11%
Wnt Signaling	16%
Hippo Signaling	21%
Whole Genome	**11%**
Pathways with 0% ABE are highlighted	