Whole exome sequencing identifies novel DYT1 dystonia-associated genome variants as potential disease modifiers

Chih-Fen Hu[1]*, G. W. Gant Luxton[2], Feng-Chin Lee[1], Chih-Sin Hsu[3], Shih-Ming Huang[4], Jau-Shyong Hong[5], San-Pin Wu[6]*

[1] Department of Pediatrics, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan
[2] Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis, MN, United States
[3] Center for Precision Medicine and Genomics, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan
[4] Department and Graduate Institute of Biochemistry, National Defense Medical Center, Taipei, Taiwan
[5] Neurobiology Laboratory, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, NC, United States
[6] Reproductive and Developmental Biology Laboratory, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, NC, United States

* Correspondence:
1. Chih-Fen Hu, Department of Pediatrics, Tri-Service General Hospital, National Defense Medical Center, Taipei 114, Taiwan, Email: caperhu@mail.ndmctsgh.edu.tw; caperhu@gmail.com
2. San-Pin Wu, Reproductive and Developmental Biology Laboratory, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, NC 27709, United States, Email: steve.wu@nih.gov

Funding information: This research was funded by Tri-Service General Hospital, grant number TSGH-C108-021 (C.F.H.), TSGH-C108-022 (C.F.H.) and National Institutes of Health GM129374 (G.W.G.L.), Z99-ES999999 (S.P.W.).

Abstract

DYT1 dystonia is a neurological movement disorder characterized by painful sustained muscle contractions resulting in abnormal twisting and postures. In a subset of patients, it is caused by a loss-of-function mutation (ΔE302/303; or ΔE) in the luminal ATPases associated with various cellular activities (AAA+) protein torsinA encoded by the *TOR1A* gene. The low penetrance of the ΔE mutation (~30-40%) suggests the existence of unknown genetic modifiers of DYT1 dystonia. To identify these modifiers, we performed whole exome sequencing (WES) of blood leukocyte DNA isolated from two DYT1 dystonia patients, three asymptomatic carriers of the ΔE mutation, and an unaffected adult relative. A total of 264 DYT1 dystonia-associated variants (DYT1 variants) were identified in 195 genes. Consistent with the emerging view of torsinA as an important regulator of the cytoskeleton, endoplasmic reticulum homeostasis, and lipid metabolism, we found DYT1 variants in genes that encode proteins implicated in these processes. Moreover, 40 DYT1 variants were detected in 32 genes associated with neuromuscular and neuropsychiatric disorders. The DYT1 variants described in this work represent exciting new targets for future studies designed to increase our understanding of the pathophysiology and pathogenesis of DYT1 dystonia.

Keywords: DYT1 dystonia; *TOR1A*; torsinA; neurogenetics; whole exome sequencing

1. INTRODUCTION

Dystonias are a heterogeneous collection of hyperkinetic neurological movement disorders that are characterized by involuntary muscle contractions resulting in abnormal repetitive movements and postures (Alberto Albanese et al., 2013; Fahn, 1988). Dystonias can be acquired as the result of environmental insults (i.e. central nervous system infection, toxins, and traumatic brain injury) (Alberto Albanese et al., 2013; A. Albanese, Di Giovanni, & Lalli, 2019) as well as inherited due to genetic mutations (Weisheit, Pappas, & Dauer, 2018). While several causative genes are known, the mechanisms underlying their contribution to dystonia pathogenesis and/or pathophysiology remain unclear.

Early onset torsion dystonia, or DYT1 dystonia, is the most common and severe inherited dystonia (Petrucci & Valente, 2013). It is a primary torsion dystonia, as dystonia is the only clinical symptom present in patients and it is inherited in a monogenic fashion. The majority of DYT1 dystonia cases are caused by the autosomal dominantly inherited deletion of a GAG codon (c.904_906/907_909ΔGAG) from the *TOR1A* gene, which removes a glutamic acid residue (ΔE302/303; or ΔE) from the C-terminus of the encoded luminal ATPase torsinA (Neuwald, Aravind, Spouge, & Koonin, 1999; L. J. Ozelius et al., 1997). The ΔE mutation is considered a loss-of-function mutation because homozygous torsinA-knockout and homozygous torsinA^{ΔE}-knockin mice both die perinatally and exhibit neurons with abnormal blebbing of the inner nuclear membrane into the perinuclear space of the nuclear envelope (Goodchild, Kim, & Dauer, 2005). In addition, the ΔE mutation impairs the ability of torsinA to interact with its major binding partners the inner nuclear membrane protein lamina-associated polypeptide 1 (LAP1) and the

endoplasmic reticulum/outer nuclear membrane protein luminal domain-like LAP1

(LULL1) (Naismith, Dalal, & Hanson, 2009), which stimulates the ability of torsinA to

hydrolyze ATP above negligible background levels *in vitro* (Zhao, Brown, Chase,

Eisele, & Schlieker, 2013).

Surprisingly, only ~30-40% of individuals heterozygous for the ΔE develop DYT1

dystonia despite the presence of abnormalities in brain metabolism and the

cerebellothalamocortical pathway in all carriers (Argyelan et al., 2009; Eidelberg et

al., 1998; Niethammer, Carbon, Argyelan, & Eidelberg, 2011; Premi et al., 2016;

Trost et al., 2002). Collectively, these clinical findings demonstrate that the presence

of the ΔE mutation results in abnormal brain function regardless of whether or not

an individual develops DYT1 dystonia. Moreover, they suggest the hypothesis that

the penetrance of the ΔE mutation may be influenced by additional as-of-yet

unknown genetic factors.

Consistent with this hypothesis, recent research shows that genetic background

modulates the phenotype of a mouse model of DYT1 dystonia (Tanabe, Martin, &

Dauer, 2012). In addition, expression profiling in peripheral blood harvested from

human DYT1 dystonia patients harboring the ΔE mutation and asymptomatic

carriers revealed a genetic signature that could correctly predict disease state

(Walter et al., 2010). The functional classification of transcripts that were differentially

regulated in DYT1 dystonia patients relative to unaffected carriers identified a variety

of potentially impacted biological pathways, including cell adhesion, cytoskeleton

organization and biogenesis, development of the nervous system, G-protein receptor

signaling, and vesicle-mediated pathway/protein transport. Since these biological

pathways have all been previously associated with torsinA function (Cascalho,

Jacquemyn, & Goodchild, 2017; Gonzalez-Alegre, 2019; Laudermilch & Schlieker,

2016; Weisheit et al., 2018), we hypothesize that the penetrance of the ΔE mutation

and therefore the development of DYT1 dystonia may depend upon the presence or

absence of variants in genes that encode proteins that influence biological pathways

associated with torsinA function. Below, we describe the use of WES to identify

genetic variants in DYT1 dystonia patients but neither unaffected ΔE mutation

carriers nor the unaffected control.

2. MATERIALS AND METHODS

2.1 Human Subjects. This study recruited 11 human subjects, including two patients

from two separate families of Taiwanese ancestry. All subjects (or legal guardians)

gave their written informed consent for participation and the study was approved by

the Institutional Review Board of the Tri-Service General Hospital at the National

Defense Medical Center in Taipei, Taiwan (IRB# 1-107-05-164). Detailed clinical

information was obtained from corresponding clinicians and medical records.

2.2 Purification of genomic DNA from Isolated Human Blood Leukocytes.

Genomic DNA was purified from human leukocytes using the MagPurix® Blood DNA

Extraction Kit LV and run in the MagPurix 24® Nucleic Acid Extraction System

(Labgene Scietific, SA, Châtel-Saint-Denis, Switzerland) following the instructions

provided by the manufacturer.

2.3 Sanger Sequencing of the *TOR1A* gene. The DNA encoding portions of the

TOR1A gene was PCR amplified from the genomic DNA purified from human

leukocytes using the following primer pairs: 1) Transcriptome-Forward (F):

ATCTACCCGCGTCTCTAC and –Reverse (R): ATAATCTAACTTGGTGAACA; 2)

TOR1A c.646G>C, D216H-F: TAATTCAGGATCAGTTACAGTTGTG and –R:

TGCAGGATTAGGAACCAGAT; and 3) *TOR1A* c.904_906/907_909ΔGAG, ΔE-F:

GTGTGGCATGGATAGGTGACCC and –R: GGGTGGAAGTGTGGAAGGAC. The

resulting PCR products were purified using QIAquick PCR Purification Kit (from

company Qiagen®) and subjected to Sanger sequencing, which was performed by

Genomics® (Taipei, Taiwan).

2.4 WES. Purified human genomic DNA was sheared into ~150-200 base-pair

fragments using the S220 Focused-Ultrasonicator (Covaris, Woburn, Massachusetts)

according to the instructions provided by the manufacturer. SureSelectXT Human

All Exon V6 +UTR (Agilent Technologies, Santa Clara, CA) was then used to perform

exome capture and library preparations The library were then sequenced using a

NovaSeq 6000 System (Illumina, San Diego, CA) with 150 base-pair reads and

output data up to 10 Gb per sample. After sequencing, Genome Analysis Toolkit

(GATK) best practices workflows of germline short variant discovery

(https://software.broadinstitute.org/gatk) was used to perform variant calling with

default parameters (DePristo et al., 2011). Briefly, the Burrows-Wheeler Aligner was

first used to align the sequenced exomes with the most up-to-date human genome

reference build (hg38) ("GRCh38 - hg38 - Genome - Assembly - NCBI".

ncbi.nlm.nih.gov). Next, duplicate reads were removed using Picard after which the

GATK was used to perform local realignment of the sequenced exomes with the

reference genome and base quality recalibration. Then, GATK-HaplotypeCaller was

used to call germline SNPs (single-nucleotide polymorphism) and indels. After

variant calling, ANNOVAR was used to variant annotation (Wang, Li, & Hakonarson,

2010) with database, include refGene, clinvar_20170905

(https://www.ncbi.nlm.nih.gov/clinvar/), avsnp150, dbnsfp33a, gnomad_genome,

dbscsnv11. Annotated variants were selected with the following criteria: (1) filtering

with exonic region, (2) removing synonymous mutation, (3) read depth ≥20. Next,

we categorized these filtered variants according to the principle of inheritance

(Figure S1). Finally, the variants of interest were validated by manually viewing them

in the Integrative Genomics Viewer. All of the whole exome sequencing data

generated in this study are deposited online at GenBank

(https://www.ncbi.nlm.nih.gov/sra/PRJNA523662).

3. RESULTS

3.1 Clinical Observations. Patient 1 was a 12-year-old male of Taiwanese descent

who initially presented with waddling gait at seven years of age, which progressed

to upper limb tremor and pronation within a few months. Over time, the patient

sequentially displayed head tilt, scoliosis, kyphosis, repetitive and active twisting of

his limbs. Five years after the onset of his symptoms, the patient showed generalized

and profound muscle twisting and contraction, including dysarthria and dysphagia.

The patient now presents with a sustained opitoshtonous-like posture and needs full

assistance with executing his daily routines. Unfortunately, the patient did not benefit

greatly from medical treatment and he refused deep brain stimulation due to the risks

associated with the necessary surgical procedure. Neither he nor his family had a

prior history of dystonia-related neurological movement disorders. Medical records

from the hospital where Patient 1 received care prior to this study indicate that the

patient lacks any mutations in his *FXN* or *THAP1* loci, which are both differential

diagnoses of genetic, progressive, and neurodegenerative movement disorders

(Campuzano et al., 1996; L. Ozelius & Lubarr, 1993).

Patient 2 was a 40-year-old male of Taiwanese descent who initially presented with

mild foot dystonia followed by cervical dystonia in his early twenties. He is able to

execute his daily routines as the result of medical treatment. The patient had no prior

history of dystonia-related neurological movement disorders and clinical information

regarding the medical history of his family is unavailable.

3.2 Examination of Known DYT1 Dystonia-Associated Mutations. To determine

if either Patient 1 (subject 1) or Patient 2 (subject 11) harbored known DYT1

dystonia-associated mutations in their genomes, we used Sanger sequencing to

screen their *TOR1A* genes for the presence of the ΔE mutation. Our results show

that both patients are heterozygous for the ΔE mutation (Figures 1A-B,1E-F). We

also sequenced the *TOR1A* genes of nine other family members of Patient 1. No ΔE

mutation was found in the genomes of his unaffected mother (subject 2), male sibling

(subject 4) or other relatives (subject 5, 7, 8,10), while his father (subject 3), paternal

aunt (subject 6), and cousin of Patient 1 (subject 9) were asymptomatic

heterozygotic ΔE mutation carriers (Figures 1A-B). Then, we asked if the previously

described protective modifier mutation D216H was present within the *TOR1A* gene

of the core family of patient 1, including subject 1,2,3,4 (Kock et al., 2006 Hum Mol

Genet). However, none of the family members examined were positive for the

D216H mutation (Figures 1C-D). These results suggest that Patient 1 inherited the

ΔE mutation from the paternal side of his family and that the absence of DYT1

dystonia in his father, paternal aunt, and cousin cannot be attributed to the presence

of the protective D216H mutation.

3.3 Identification of DYT1 Dystonia-Associated Genome Variants in the *TOR1A*

gene. To begin to identify potential genetic modifiers of the penetrance of the ΔE

mutation, we performed WES on genomic DNA purified from blood leukocytes

isolated from Patient 1 and Patient 2 as well as three asymptomatic ΔE mutation

carriers from the first family (i.e. the father, paternal aunt, and cousin) and the mother

of patient 1 who did not harbor the ΔE mutation in her genome. Consistent with the

Sanger sequencing results described above, WES confirmed the presence of a

single copy of the ΔE mutation in the exomes of Patient 1, his father, paternal aunt,

and cousin as well as Patient 2, while demonstrating its absence from the mother of

Patient 1 (Table 1). In addition, WES demonstrated that the D216H mutation was

absent from all six exomes examined. Interestingly, three additional previously

reported *TOR1A* variants, rs13300897, rs2296793 and rs1182 (Siokas et al., 2017;

Vulinovic et al., 2014), were found in the exomes of two of the asymptomatic ΔE

mutation carriers and the mother from the family of Patient 1(Table 1). However, the

absence of these variants from the exomes of either Patient 1 or Patient 2 diminishes

the likelihood that they are genetic modifiers of the penetrance of the ΔE mutation.

Collectively, these findings motivated us to search for genome variants outside of

the *TOR1A* gene that might influence the penetrance of the ΔE mutation.

3.4 Identification of DYT1 dystonia-associated Genome Variants (DYT1

variants) Outside of the *TOR1A* Gene. We hypothesized that candidate modifiers

of the penetrance of the ΔE mutation would be those genome variants that were

present in the exomes of both symptomatic patients and absent from the exomes of

the asymptomatic ΔE mutation carriers and mother of the Patient 1. To begin to test

this hypothesis, we examined the results of our WES for genome variants that fit this

criterion and identified a total of DYT1 variants 264 variants in 195 genes. Based on

their respective allele frequencies (AFs), we further classified these variants into

three inheritance groups: 1) Autosomal recessive (AR); 2) Autosomal dominant (AD);

and 3) *De novo* (DN) mutation (Table 2). The 53 genome variants found in 43 genes

classified as AR had AFs of 1 for both patients, an AF of 0.5 or 1 for the mother of

Patient 1, an AF of 0.5 for the father of Patient 1, and an AF of 0 or 0.5 for the paternal

aunt and cousin of Patient 1. The 201 variants found in 149 genes classified as AD

had AFs of 0.5 or 1 for both patients, and AF of 0.5 or 1 for the mother of Patient 1,

and an AF of 0 for the rest of the family members of Patient 1. Finally, the 10 variants

in 5 genes classified as DN had AFs of 0.5 or 1 for both patients, and were not

present in any of the other exomes examined.

Of the 264 variants identified by our WES-based screen, 26 genome variants were

previously implicated in human disease (Table S1) based on their annotation in the

ClinVar database (https://www.ncbi.nlm.nih.gov/clinvar/). Analysis of the potential

pathological impact of these variants using the bioinformatics tools sorting intolerant

from tolerant (SIFT) and polymorphism phenotyping (PolyPhen) (Flanagan, Patch,

& Ellard, 2010) predicts that 11 variants might be loss-of-function mutations (Table

2). Furthermore, gene ontology analysis performed using the Database for

Annotation, Visualization and Integrated Discovery (DAVID) (Huang da, Sherman, &

Lempicki, 2009) identified clustered annotations of genes in which the DYT1 variants

were identified by our WES-based approach. There are total 30 annotation clusters

generated by this tool as listed in Table S2. Next, we filtered these categories with

enrichment score>1 and *p* value<0.05 and the results were enriched for those that

encode proteins that contain the epidermal growth factor-like domain (ten genes),

have dioxygenase (four genes) or Rho guanyl-nucleotide exchange factor activity

(four genes), or exhibit the ability to interact with the actin cytoskeleton (seven genes)

(Table S2). Notably, genes for ER stress and lipid metabolism, which are linked to

DYT1 functions (discussed below), also shown a trend of enrichment (Table S2).

The enrichment of cytoskeleton-related genes and the known function of TOR1A in

regulation of the mechanical integration of the nucleus and the cytoskeleton

prompted us to look closer on the genes that harbor DYT1 variants via literature

search (Flávia C. Nery et al., 2008; Cosmo A. Saunders et al., 2017; Vander Heyden,

Naismith, Snapp, Hodzic, & Hanson, 2009). There are 45 DYT1 variants in 34 genes

that are associated with cytoskeleton (Table 3). In addition, 17 DYT1 variants in 16

genes are found to be linked to endoplasmic reticulum and protein and lipid

metabolism (Table 4), which TOR1A is known to have functional indications at

(Gonzalez-Alegre, 2019; Shin et al., 2019). Lastly, further reviewing previous studies

identified 40 DYT1 variants in 32 genes that have disease associated with human

neuropsychiatric disorders or neuromuscular diseases (Table 5). Taken together,

our results suggest that potential regulators of the ΔE mutation may participate in

the regulation of the following established cellular functions performed by torsinA:

cytoskeletal organization, endoplasmic reticulum homeostasis, and protein and lipid

metabolism.

4. DISCUSSION

The underlying cause of phenotype variation from the same allele remains largely

unknown in most cases when a particular genotype is inherited. Emerging evidence

indicate that modifier genes may contribute to phenotypic variations (Kammenga,

2017). For example, patients with thalassemia, a disorder caused by defective β-

globin synthesis, have diverse clinical characteristics and variable expressivity. A

number of factors underlie this phenotypic diversity, including the involvement of

numerous modifier genes at other genetic loci that affect the production of β-globin

(Rujito et al., 2016). Similarly, DYT1 dystonia patients have a wide spectrum of

symptom severity, which reflects the incomplete penetrance of the pathogenic ΔE

mutation and the variable expressivity of the disease. For most diseases, variable

expressivity of the disease phenotype is the norm among individuals who carry the

same disease-causing allele or alleles (Nadeau, 2001), despite the causes are not

always being clear.

In this work, we describe the identification of 264 variants in 195 genes that are

associated with DYT1 dystonia. Below, we will discuss the potential implications of

our results on our understanding of the pathogenesis and pathophysiology of DYT1

dystonia. Specifically, we will explore the connections between the DYT1 variants

identified here and the following established cellular functions of torsinA: cytoskeletal

regulation, endoplasmic reticulum stress, and lipid metabolism. In addition, we will

examine the relationship revealed between DYT1 dystonia and the neuromuscular

and neuropschiatric disorders linked with the genes in which we identified DYT1

dystonia-associated genomic variants.

DYT1 variants and the cytoskeleton. Of the 195 genes that we identified as

harboring 264 DYT1 variants, 34 genes encode proteins that constitute or associate

with the cytoskeleton (Table 3). Specifically, we found a total of 23 DYT1 variants in

18 genes that encode proteins involved in the function of the microtubule

cytoskeleton. We also found 12 DYT1 variants in ten genes encoding actin

cytoskeleton-associated proteins as well as ten variants in six genes encoding

intermediate filament cytoskeleton-associated proteins. Moreover, seven of the 34

genes described above were found to harbor at least two DYT1 variants, including

CCDC74B, *DYNC2H1*, *KRT6A, KRT6B, LIMCH1, NRAP,* and *TUBA3E*.

The identification of DYT1 variants in genes encoding proteins related to cytoskeletal

function is consistent with the emerging view of torsinA as a critical regulator of

cellular mechanics. Since its discovery in 1997, torsinA function has been implicated

in the regulation of cytoskeletal dynamics and organization (Breakefield, Kamm, &

Hanson, 2001). The first evidence to suggest that torsinA might be involved in

cytoskeletal regulation was the finding that the nematode torsinA protein OOC-5 was

required for the rotation of the nuclear-centrosome complex during early

embryogenesis (Basham & Rose, 1999, 2001). In addition, the fruit fly torsinA protein

torp4a/dTorsin was implicated in the regulation of the actin cytoskeleton (Muraro &

Moffat, 2006). Furthermore, the over-expression of a torsinA construct containing

the ΔE mutation was shown to inhibit neurite extension in human neuroblastoma

cells and to increase the density of vimentin intermediate filaments around the

nucleus (J. W. Hewett, Zeng, Niland, Bragg, & Breakefield, 2006). The relationship

between torsinA and the cytoskeleton is further strengthened by reports of the

impaired migration of dorsal forebrain neurons and fibroblasts from torsinA-knockout mice as well as DYT1 dystonia patient-derived fibroblasts (McCarthy, Gioioso, Zhang, Sharma, & Bhide, 2012; Flavia C. Nery et al., 2014; Flávia C. Nery et al., 2008).

More recently, torsinA was identified as a key regulator of the mechanical integration of the nucleus and the cytoskeleton via the conserved nuclear envelope-spanning linker of nucleoskeleton and cytoskeleton (LINC) complex (Flávia C. Nery et al., 2008; Cosmo A. Saunders et al., 2017; Vander Heyden et al., 2009). The core of LINC complexes is formed by the transluminal interaction between the outer and inner nuclear membrane Klarischt/ANC-1/SYNE homology (KASH) and Sad1/UNC-84 (SUN) proteins, respectively (Crisp et al., 2006). KASH proteins interact with the cytoskeleton and signaling proteins within the cytoplasm (Luxton & Starr, 2014), whereas SUN proteins interact with chromatin, other inner nuclear membrane proteins, and the nuclear lamina within the nucleoplasm (Chang, Worman, & Gundersen, 2015).

While the precise mechanism of torsinA-mediated LINC complex regulation remains unclear, torsinA interacts with the luminal domains of both KASH and SUN proteins (Chalfant, Barber, Borah, Thaller, & Lusk, 2019; Flávia C. Nery et al., 2008). The ability of torsinA to interact with KASH and SUN proteins is thought to promote the disassembly of LINC complexes given the fact that most AAA+ proteins act as molecular chaperones that disassemble protein complexes (Hanson & Whiteheart, 2005; C. A. Saunders & Luxton, 2016). This hypothesis is supported by the finding that torsinA loss elevates LINC complex levels in the mouse brain, which impairs

brain morphogenesis (Dominguez Gonzalez et al., 2018). More recently, fibroblasts

isolated from DYT1 dystonia patients were shown to have increased deformability

similar to that of fibroblasts harvested from mice lacking the two major SUN proteins

SUN1 and SUN2 (Gill et al., 2019).

DYT1 dystonia patient-derived fibroblasts were also shown to have increased

susceptibility to damage by mechanical forces (Gill et al., 2019) strongly suggests

that cellular mechanics may impact the pathogenesis and/or pathophysiology of

DYT1 dystonia. All cells, including neurons, adapt their mechanical properties by

converting extracellular mechanical stimuli into biochemical signals and altered gene

expression through the process of mechanotransduction (Barnes, Przybyla, &

Weaver, 2017; Franze, 2013). Since mechanotransduction instructs neuronal

differentiation, proliferation, and survival (Iwashita, Kataoka, Toida, & Kosodo, 2014;

Koser et al., 2016), it is possible that defective mechanotransduction of neurons in

the developing brain may contribute to the pathogenesis and/or pathophysiology of

DYT1 dystonia. Based on the information provided above, it is intriguing that we

identified DYT1 variants in the KASH protein nesprin-2-encoding *SYNE2* gene and

the *NUP58* gene, which encodes the nuclear pore complex protein nup58 (Table 3

and Figure 2). In the future, it will be interesting to test if the DYT1 variants found in

SYNE2 and *NUP58* negatively impact LINC complex-dependent nuclear-

cytoskeletal coupling and/or mechanotransduction.

It is tempting to speculate that the impairment of the microtubule cytoskeleton is

particularly relevant to dystonia pathogenesis given the enrichment of DYT variants

that we found in genes that encode microtubule-associated proteins. Microtubules

are fundamentally important for the structure and function of neurons, which are

some of the most highly polarized cells in the human body (Kelliher, Saunders, &

Wildonger, 2019). Microtubules establish the polarized architecture of neurons and

serve as tracks for microtubule motor proteins as they carry proteins and lipids to

where they are needed for proper neuronal function. Thus, defects in microtubule

dynamics and organization underly a wide array of neurological and neuropsychiatric

disorders (Matamoros & Baas, 2016; Muñoz-Lasso, Romá-Mateo, Pallardó, &

Gonzalez-Cabo, 2020; Sleigh, Rossor, Fellows, Tosolini, & Schiavo, 2019).

Consistent with our identification of 4 DYT1 variants in the *TUBA3E*, which encodes

the protein α-tubulin-3E, mutations in the β-tubulin-4A-encoding *TUBB4A* gene

cause another hereditary dystonia, Whispering dysphonia or DYT4 dystonia

(Hersheson et al., 2013; Lohmann et al., 2013). These mutations result in the

formation of disorganized microtubule networks and the impaired growth of neuronal

processes similar to the clinical phenotypes observed in DYT4 dystonia patients

(Curiel et al., 2017; Watanabe et al., 2018). Future experiments designed to test the

impact of the DYT1 variants in *TUBA3E* on the organization and function of neuronal

microtubules will help elucidate the role of the microtubule cytoskeleton to the

manifestation of DYT1 dystonia.

DYT1 variants in association with protein synthesis and transport and ER

homeostasis. Accumulating evidence indicate a role of TOR1A in the cellular

protein quality control system in which TOR1A could be both substrate and effector

(Gonzalez-Alegre, 2019). In the 264 genome variants, we observed six variants in

five genes, *CHGB, DOP1B, MTMR6, P2RY13* and *PPP1R15A*, that are annotated

with protein synthesis and transport functions (Table 4). Notably, CHGB and

PPP1R15A has also been linked to ER stress (Crespillo-Casado, Chambers, Fischer,

Marciniak, & Ron, 2017; Ohta et al., 2016; Zhou, Brush, Choy, & Shenolikar, 2011).

These findings support the previously proposed hypothesis that elevated levels of

endoplasmic reticulum stress contributes to DYT1 dystonia pathogenesis (G.

Beauvais et al., 2016; Genevieve Beauvais et al., 2018; Chen et al., 2010; Gordon,

Glenn, & Gonzalez-Alegre, 2011; Hettich et al., 2014; J. Hewett et al., 2003; Kim et

al., 2015; Flávia C. Nery et al., 2011; Thompson et al., 2014).

TorsinA functions to protect against insults from protein aggregates in the neural

system (G. Beauvais et al., 2016). Protein aggregates are products of protein

misfolding commonly seen in neurodegenerative diseases such as Alzheimer's

disease, Parkinson's disease, amyotrophic lateral sclerosis and prion disease, which

triggers endoplasmic reticulum stress response (De Mattos et al., 2020; Scheper &

Hoozemans, 2015). In the TOR1A ΔE mutation background, we identified six

candidate modifier genome variants in five genes that have known functions in

endoplasmic reticulum for protein post translational modification, protein

translocation and ER stress response (Table 4). Among them, DOP1B has

neurological roles in both human and mice (Dickinson et al., 2016; Rachidi,

Delezoide, Delabar, & Lopes, 2009). Whether DOP1B's endoplasmic reticulum

cellular function has a causal effect on its neurological role remains to be

investigated. Collectively, our data provide clinical indications of candidate genes

and genome variants for further investigation on the underlying mechanisms of

TOR1A dependent ER dysfunction in DYT1 dystonia.

DYT1 variants and lipid metabolism.TOR1A also has a pivotal role in lipid

metabolism as demonstrated by the hepatic steatosis of liver-specific torsinA-

knockout mouse model (Shin et al., 2019) and the requirement for the Drosophila

torsinA homologue for proper lipid metabolism in adipose tissue (Grillet et al., 2016).

Because of its functional indication in lipid metabolism, TorsinA is thought to promote

membrane biogenesis (Cascalho et al., 2017) and synaptic physiology (Lauwers,

Goodchild, & Verstreken, 2016). There are 11 DYT1 dystonia associated genome

variants identified in ten lipid metabolism genes *ALOXE3, APOB, CYP1B1, CYP2A7,*

FAM135B, GAL3ST1, GPAM, MTMR6, PLA2G4F and *PLCL1* (Table 4), which

suggest potential genetic interactions between the ΔE mutation and genome variants

that might change membrane homeostasis.

TOR1A regulates lipid metabolism in both fruit flies and mammals (Grillet et al., 2016;

Shin et al., 2019). TOR1A facilitates cell growth, raises lipid content of cellular

membrane and is involved in membrane expansion (Grillet et al., 2016). The linkage

between the TOR1A ΔE mutation and 10 lipid metabolic genes suggest the impact

on lipid metabolism associated cellular functions could be amplified by clustered

mutations and genome variants. Two genes in this category have known functions

in the neural system. The *GAL3ST1* gene encodes galactose-3-O-sulfotransferase

1 that involves in the synthesis of a major lipid component of the myelin sheath

galactosylceramide sulfate (Boggs, 2014). *Gal3st1* deficient mice develop tremor,

progressive ataxia, hind limb weakness, aberrant limb posture and impaired limb

coordination with morphological defects in the neural system (Honke et al., 2002).

PLCL1 Involves in an inositol phospholipid-based intracellular signaling cascade.

PLCL1 is phospholipase C like protein lacking the catalytic activity. PLCL1 binds and

sequesters inositol triphosphates to blunt the downstream calcium signaling (Kanematsu, Takeuchi, Terunuma, & Hirata, 2005). PLCL1 has been linked to the trafficking and turnover of GABAA receptors in neurons (Kanematsu et al., 2007; Mizokami et al., 2007). Physiologically, loss of PLCL1 increases the incidence of chemically induced seizure in mice (Yamaguchi et al., 2004). These findings indicate an essential role of PLCL1 in controlling the neural signaling transduction. While the functional impact of the genome variants on GAL3ST1 and PLCL1 awaits further investigation, their association with the TOR1A ΔE mutation suggests potential functional interactions between these molecules in DYT1 dystonia.

Connections between the genes harboring DYT1 variants and their implicated neuromuscular and neuropsychiatric disorders. The loss of torsinA function in either the cerebral cortex or cerebellum result in motor dysfunction (DeSimone et al., 2017; Fremont, Tewari, Angueyra, & Khodakhah, 2017; Yokoi, Dang, Mitsui, Li, & Li, 2008), indicating a neuronal component of TOR1A's function in dystonia. Based on these observations, we examined the 195 genes that carry candidate ΔE mutation modifiers for their association with neuropsychiatric and neuromuscular disorders. Such link was identified in 32 genes with 40 genome variants (Table 5). These include the *AHNAK2*, *ARHGEF3*, *CDRT1*, *GBE1* and *NRG2* genes in associated with peripheral neuropathy (Charcot-Marie-Tooth disease and Polyglucosan body neuropathy, adult form). The *AMPD2*, *ATXN7* and *MICAL3* genes are linked to cerebellar diseases (Pontocerebellar Hypoplasia, type 9 and spastic paraplegia 63, autosomal recessive; Spinocerebellar ataxia 7; Joubert syndrome (cerebelloparenchymal disorder)). Lastly, the *IRF3*, *TRAF3* and *LIPT2* genes are associated with encephalopathy (acute, infection-induced; encephalopathy, neonatal severe, with lactic acidosis and brain abnormalities and lipoic acid

biosynthesis defects. Overall, more than 16% of the identified 195 genes are in

association with neuropsychiatric and neuromuscular diseases related disorders,

demonstrating the significance of the linkage between DYT1 dystonia and these

diseases.

Study Limitations. The present study examined five individuals who have the

TOR1A ΔE mutation. Among them, two have disease presentation and three are

asymptomatic carriers. Furthermore, one affected patient and the three

asymptomatic carriers are in the same family, which is an advantage to have a

relatively close genetic background for modifier screening. Data from this family

identified 1725 of genome variants as candidate modifiers. With the addition of the

second affected patient, the number of candidate modifier variants were further

narrowed down to 264. This number could have been reduced if data from more

affected patients or asymptomatic carriers are available. Unfortunately, family

members of the second affected patient declined to participate in the study. Due to

the rareness of DYT1 dystonia in Taiwan, it is difficult to increase sample size within

the Taiwanese population in foreseeable future. Alternatively, meta-analysis of our

dataset with WES results from other populations across the world, once publicly

available, may help to identify the common modifiers in the general population

(Rodriguez-Quiroga, Gonzalez-Moron, Espay, & Kauffman, 2018; Zech et al., 2017).

The WES data allows identification of candidate modifiers in the coding genome.

However, majority of the GWAS signals are mapped to the noncoding regions of the

genome and accumulating evidence point to disease associations with the

noncoding genome (Zhang & Lupski, 2015). Mutations in the noncoding genome

may impact cis-acting element functions and chromatin conformations that direct

gene expression. Future inclusion of the whole genome sequencing assay may help

to identify additional modifiers for the DYT1 dystonia.

5. CONCLUSIONS

In summary, we propose that genome variants within nuclear-cytoskeletal coupling

network may constitute potential modifier variants, which could synergistically

reduce the threshold of disease onset of DYT1 dystonia and accelerates the clinical

symptoms and signs of dystonia. We believe that this study provided a path to

unravel candidate genome variants as modifiers. Our findings not only echo the

previous research highlighting the defect of mechanosensing and

mechanotransduction regulated by TOR1A (Gill et al., 2019), but provide knowledge

for further understanding the disease origin of the DYT1 dystonia as well. We will

recommend the physicians to test these variants once the TOR1A ΔE mutation

patient show normal alleles within other *TOR1A* locus and other major binding

proteins in their study. We also provide a list of candidate genes and genome

variants for future mechanistic studies on DYT1 dystonia.

Acknowledgments: We would like to thank the first patient and his family members

who provided the DNAs and clinical information necessary for this research study.

We would also like to thank Dr. Chin-Hsien Lin (Department of Neurology, National

Taiwan University Hospital, Taipei, Taiwan) who kindly provided the DNA sample

and clinical information of the second patient.

Author Contribution

Conceptualization, FCL and CFH; Data curation, GWGL and CSH; Formal analysis

FCL and CSH; Investigation, CFH and GWGL; Methodology, FCL and CFH; Project

administration, CFH; Resources, SPW, JSH and GWGL; Supervision, SMH;

Validation, CFH and SPW; Writing-original draft, CFH and GWGL; Writing-review

and editing, CFH, GWGL and SPW.

Conflict of interest: The authors declare no conflict of interest.

References

Albanese, A., Bhatia, K., Bressman, S. B., Delong, M. R., Fahn, S., Fung, V. S. C., . . .
Teller, J. K. (2013). Phenomenology and classification of dystonia: a consensus
update. *Movement disorders : official journal of the Movement Disorder Society,*
28(7), 863-873. doi:10.1002/mds.25475

Albanese, A., Di Giovanni, M., & Lalli, S. (2019). Dystonia: diagnosis and management.
European journal of neurology, 26(1), 5-17. doi:10.1111/ene.13762

Argyelan, M., Carbon, M., Niethammer, M., Ulug, A. M., Voss, H. U., Bressman, S. B., . . .
Eidelberg, D. (2009). Cerebellothalamocortical connectivity regulates penetrance in
dystonia. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience, 29(31), 9740-9747. doi:10.1523/JNEUROSCI.2300-09.2009

Barnes, J. M., Przybyla, L., & Weaver, V. M. (2017). Tissue mechanics regulate brain
development, homeostasis and disease. *J Cell Sci, 130*(1), 71-82.
doi:10.1242/jcs.191742

Basham, S. E., & Rose, L. S. (1999). Mutations in ooc-5 and ooc-3 disrupt oocyte
formation and the reestablishment of asymmetric PAR protein localization in two-
cell Caenorhabditis elegans embryos. *Developmental biology, 215*(2), 253-263.
doi:10.1006/dbio.1999.9447

Basham, S. E., & Rose, L. S. (2001). The Caenorhabditis elegans polarity gene ooc-5
encodes a Torsin-related protein of the AAA ATPase superfamily. *Development*
(Cambridge, England), 128(22), 4645-4656. Retrieved from
https://pubmed.ncbi.nlm.nih.gov/11714689

Beauvais, G., Bode, N. M., Watson, J. L., Wen, H., Glenn, K. A., Kawano, H., . . .
Gonzalez-Alegre, P. (2016). Disruption of Protein Processing in the Endoplasmic
Reticulum of DYT1 Knock-in Mice Implicates Novel Pathways in Dystonia
Pathogenesis. *J Neurosci, 36*(40), 10245-10256. doi:10.1523/jneurosci.0669-
16.2016

Beauvais, G., Rodriguez-Losada, N., Ying, L., Zakirova, Z., Watson, J. L., Readhead,
B., . . . Gonzalez-Alegre, P. (2018). Exploring the Interaction Between eIF2α
Dysregulation, Acute Endoplasmic Reticulum Stress and DYT1 Dystonia in the
Mammalian Brain. *Neuroscience, 371*, 455-468.
doi:10.1016/j.neuroscience.2017.12.033

Boggs, J. M. (2014). Role of galactosylceramide and sulfatide in oligodendrocytes and

570   CNS myelin: formation of a glycosynapse. *Adv Neurobiol, 9*, 263-291.

571   doi:10.1007/978-1-4939-1154-7_12

572   Breakefield, X. O., Kamm, C., & Hanson, P. I. (2001). TorsinA: movement at many levels.

573   *Neuron, 31*(1), 9-12. doi:10.1016/s0896-6273(01)00350-6

574   Campuzano, V., Montermini, L., Moltò, M. D., Pianese, L., Cossée, M., Cavalcanti, F., . . .

575   Pandolfo, M. (1996). Friedreich's ataxia: autosomal recessive disease caused by

576   an intronic GAA triplet repeat expansion. *Science (New York, N.Y.), 271*(5254),

577   1423-1427. doi:10.1126/science.271.5254.1423

578   Cascalho, A., Jacquemyn, J., & Goodchild, R. E. (2017). Membrane defects and genetic

579   redundancy: Are we at a turning point for DYT1 dystonia? *Movement disorders :*

580   *official journal of the Movement Disorder Society, 32*(3), 371-381.

581   doi:10.1002/mds.26880

582   Chalfant, M., Barber, K. W., Borah, S., Thaller, D., & Lusk, C. P. (2019). Expression of

583   TorsinA in a heterologous yeast system reveals interactions with lumenal domains

584   of LINC and nuclear pore complex components. *Mol Biol Cell, 30*(5), 530-541.

585   doi:10.1091/mbc.E18-09-0585

586   Chang, W., Worman, H. J., & Gundersen, G. G. (2015). Accessorizing and anchoring the

587   LINC complex for multifunctionality. *The Journal of cell biology, 208*(1), 11-22.

588   doi:10.1083/jcb.201409047

589   Chen, P., Burdette, A. J., Porter, J. C., Ricketts, J. C., Fox, S. A., Nery, F. C., . . . Caldwell,

590   G. A. (2010). The early-onset torsion dystonia-associated protein, torsinA, is a

591   homeostatic regulator of endoplasmic reticulum stress response. *Human molecular*

592   *genetics, 19*(18), 3502-3515. doi:10.1093/hmg/ddq266

593   Crespillo-Casado, A., Chambers, J. E., Fischer, P. M., Marciniak, S. J., & Ron, D. (2017).

594   PPP1R15A-mediated dephosphorylation of eIF2α is unaffected by Sephin1 or

595   Guanabenz. *Elife, 6*, e26109. doi:10.7554/eLife.26109

596   Crisp, M., Liu, Q., Roux, K., Rattner, J. B., Shanahan, C., Burke, B., . . . Hodzic, D. (2006).

597   Coupling of the nucleus and cytoplasm: role of the LINC complex. *The Journal of*

598   *cell biology, 172*(1), 41-53. doi:10.1083/jcb.200509124

599   Curiel, J., Rodríguez Bey, G., Takanohashi, A., Bugiani, M., Fu, X., Wolf, N. I., . . .

600   Vanderver, A. (2017). TUBB4A mutations result in specific neuronal and

601   oligodendrocytic defects that closely match clinically distinct phenotypes. *Human*

602   *molecular genetics, 26*(22), 4506-4518. doi:10.1093/hmg/ddx338

603   De Mattos, E. P., Wentink, A., Nussbaum-Krammer, C., Hansen, C., Bergink, S., Melki, R.,

604   & Kampinga, H. H. (2020). Protein Quality Control Pathways at the Crossroad of

605   Synucleinopathies. *J Parkinsons Dis.* doi:10.3233/jpd-191790

606   DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., . . . Daly,

607   M. J. (2011). A framework for variation discovery and genotyping using next-

608   generation DNA sequencing data. *Nat Genet, 43*(5), 491-498. doi:10.1038/ng.806

609   DeSimone, J. C., Pappas, S. S., Febo, M., Burciu, R. G., Shukla, P., Colon-Perez, L.

610   M., . . . Vaillancourt, D. E. (2017). Forebrain knock-out of torsinA reduces striatal

611   free-water and impairs whole-brain functional connectivity in a symptomatic mouse

model of DYT1 dystonia. *Neurobiol Dis, 106*, 124-132.
doi:10.1016/j.nbd.2017.06.015
Dickinson, M. E., Flenniken, A. M., Ji, X., Teboul, L., Wong, M. D., White, J. K., . . . Murray,
S. A. (2016). High-throughput discovery of novel developmental phenotypes.
Nature, 537(7621), 508-514. doi:10.1038/nature19356
Dominguez Gonzalez, B., Billion, K., Rous, S., Pavie, B., Lange, C., & Goodchild, R.
(2018). Excess LINC complexes impair brain morphogenesis in a mouse model of
recessive TOR1A disease. *Human molecular genetics, 27*(12), 2154-2170.
doi:10.1093/hmg/ddy125
Eidelberg, D., Moeller, J. R., Antonini, A., Kazumata, K., Nakamura, T., Dhawan, V., . . .
Fahn, S. (1998). Functional brain networks in DYT1 dystonia. *Annals of neurology,*
44(3), 303-312. doi:10.1002/ana.410440304
Fahn, S. (1988). Concept and classification of dystonia. *Advances in neurology, 50*, 1-8.
Retrieved from https://pubmed.ncbi.nlm.nih.gov/3041755
Flanagan, S. E., Patch, A.-M., & Ellard, S. (2010). Using SIFT and PolyPhen to predict
loss-of-function and gain-of-function mutations. *Genetic testing and molecular*
biomarkers, 14(4), 533-537. doi:10.1089/gtmb.2010.0036
Franze, K. (2013). The mechanical control of nervous system development. *Development,*
140(15), 3069-3077. doi:10.1242/dev.079145
Fremont, R., Tewari, A., Angueyra, C., & Khodakhah, K. (2017). A role for cerebellum in
the hereditary dystonia DYT1. *Elife, 6*. doi:10.7554/eLife.22775
Gill, N. K., Ly, C., Kim, P. H., Saunders, C. A., Fong, L. G., Young, S. G., . . . Rowat, A. C.
(2019). DYT1 Dystonia Patient-Derived Fibroblasts Have Increased Deformability
and Susceptibility to Damage by Mechanical Forces. *Front Cell Dev Biol, 7*, 103.
doi:10.3389/fcell.2019.00103
Gonzalez-Alegre, P. (2019). Advances in molecular and cell biology of dystonia: Focus on
torsinA. *Neurobiol Dis, 127*, 233-241. doi:10.1016/j.nbd.2019.03.007
Goodchild, R. E., Kim, C. E., & Dauer, W. T. (2005). Loss of the dystonia-associated
protein torsinA selectively disrupts the neuronal nuclear envelope. *Neuron, 48*(6),
923-932. doi:10.1016/j.neuron.2005.11.010
Gordon, K. L., Glenn, K. A., & Gonzalez-Alegre, P. (2011). Exploring the influence of
torsinA expression on protein quality control. *Neurochemical research, 36*(3), 452-
459. doi:10.1007/s11064-010-0363-4
Grillet, M., Dominguez Gonzalez, B., Sicart, A., Pottler, M., Cascalho, A., Billion, K., . . .
Goodchild, R. E. (2016). Torsins Are Essential Regulators of Cellular Lipid
Metabolism. *Dev Cell, 38*(3), 235-247. doi:10.1016/j.devcel.2016.06.017
Hanson, P. I., & Whiteheart, S. W. (2005). AAA+ proteins: have engine, will work. *Nature*
reviews. Molecular cell biology, 6(7), 519-529. doi:10.1038/nrm1684
Hersheson, J., Mencacci, N. E., Davis, M., MacDonald, N., Trabzuni, D., Ryten, M., . . .
Houlden, H. (2013). Mutations in the autoregulatory domain of β-tubulin 4a cause
hereditary dystonia. *Annals of neurology, 73*(4), 546-553. doi:10.1002/ana.23832
Hettich, J., Ryan, S. D., de Souza, O. N., Saraiva Macedo Timmers, L. F., Tsai, S., Atai, N.
A., . . . Nery, F. C. (2014). Biochemical and cellular analysis of human variants of

the DYT1 dystonia protein, TorsinA/TOR1A. *Human mutation, 35*(9), 1101-1113. doi:10.1002/humu.22602

Hewett, J., Ziefer, P., Bergeron, D., Naismith, T., Boston, H., Slater, D., . . . Breakefield, X. O. (2003). TorsinA in PC12 cells: localization in the endoplasmic reticulum and response to stress. *Journal of neuroscience research, 72*(2), 158-168. doi:10.1002/jnr.10567

Hewett, J. W., Zeng, J., Niland, B. P., Bragg, D. C., & Breakefield, X. O. (2006). Dystonia-causing mutant torsinA inhibits cell adhesion and neurite extension through interference with cytoskeletal dynamics. *Neurobiology of disease, 22*(1), 98-111. doi:10.1016/j.nbd.2005.10.012

Honke, K., Hirahara, Y., Dupree, J., Suzuki, K., Popko, B., Fukushima, K., . . . Taniguchi, N. (2002). Paranodal junction formation and spermatogenesis require sulfoglycolipids. *Proc Natl Acad Sci U S A, 99*(7), 4227-4232. doi:10.1073/pnas.032068299

Huang da, W., Sherman, B. T., & Lempicki, R. A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc, 4*(1), 44-57. doi:10.1038/nprot.2008.211

Iwashita, M., Kataoka, N., Toida, K., & Kosodo, Y. (2014). Systematic profiling of spatiotemporal tissue and cellular stiffness in the developing brain. *Development (Cambridge, England), 141*(19), 3793-3798. doi:10.1242/dev.109637

Kammenga, J. E. (2017). The background puzzle: how identical mutations in the same gene lead to different disease symptoms. *Febs j, 284*(20), 3362-3373. doi:10.1111/febs.14080

Kanematsu, T., Fujii, M., Mizokami, A., Kittler, J. T., Nabekura, J., Moss, S. J., & Hirata, M. (2007). Phospholipase C-related inactive protein is implicated in the constitutive internalization of GABAA receptors mediated by clathrin and AP2 adaptor complex. *J Neurochem, 101*(4), 898-905. doi:10.1111/j.1471-4159.2006.04399.x

Kanematsu, T., Takeuchi, H., Terunuma, M., & Hirata, M. (2005). PRIP, a novel Ins(1,4,5)P3 binding protein, functional significance in Ca2+ signaling and extension to neuroscience and beyond. *Mol Cells, 20*(3), 305-314.

Kelliher, M. T., Saunders, H. A., & Wildonger, J. (2019). Microtubule control of functional architecture in neurons. *Current opinion in neurobiology, 57*, 39-45. doi:10.1016/j.conb.2019.01.003

Kim, A. Y., Seo, J. B., Kim, W.-T., Choi, H. J., Kim, S.-Y., Morrow, G., . . . Koh, Y. H. (2015). The pathogenic human Torsin A in Drosophila activates the unfolded protein response and increases susceptibility to oxidative stress. *BMC genomics, 16*(1), 338-338. doi:10.1186/s12864-015-1518-0

Koser, D. E., Thompson, A. J., Foster, S. K., Dwivedy, A., Pillai, E. K., Sheridan, G. K., . . . Franze, K. (2016). Mechanosensing is critical for axon growth in the developing brain. *Nature neuroscience, 19*(12), 1592-1598. doi:10.1038/nn.4394

Laudermilch, E., & Schlieker, C. (2016). Torsin ATPases: structural insights and functional perspectives. *Current opinion in cell biology, 40*, 1-7. doi:10.1016/j.ceb.2016.01.001

Lauwers, E., Goodchild, R., & Verstreken, P. (2016). Membrane Lipids in Presynaptic
Function and Disease. *Neuron, 90*(1), 11-25. doi:10.1016/j.neuron.2016.02.033
Lohmann, K., Wilcox, R. A., Winkler, S., Ramirez, A., Rakovic, A., Park, J.-S., . . . Klein, C.
(2013). Whispering dysphonia (DYT4 dystonia) is caused by a mutation in the
TUBB4 gene. *Annals of neurology, 73*(4), 537-545. doi:10.1002/ana.23829
Luxton, G. W., & Starr, D. A. (2014). KASHing up with the nucleus: novel functional roles
of KASH proteins at the cytoplasmic surface of the nucleus. *Curr Opin Cell Biol, 28*,
69-75. doi:10.1016/j.ceb.2014.03.002
Matamoros, A. J., & Baas, P. W. (2016). Microtubules in health and degenerative disease
of the nervous system. *Brain research bulletin, 126*(Pt 3), 217-225.
doi:10.1016/j.brainresbull.2016.06.016
McCarthy, D. M., Gioioso, V., Zhang, X., Sharma, N., & Bhide, P. G. (2012). Neurogenesis
and neuronal migration in the forebrain of the TorsinA knockout mouse embryo.
Developmental neuroscience, 34(4), 366-378. doi:10.1159/000342260
Mizokami, A., Kanematsu, T., Ishibashi, H., Yamaguchi, T., Tanida, I., Takenaka, K., . . .
Hirata, M. (2007). Phospholipase C-related inactive protein is involved in trafficking
of gamma2 subunit-containing GABA(A) receptors to the cell surface. *J Neurosci,*
27(7), 1692-1701. doi:10.1523/jneurosci.3155-06.2007
Muñoz-Lasso, D. C., Romá-Mateo, C., Pallardó, F. V., & Gonzalez-Cabo, P. (2020). Much
More Than a Scaffold: Cytoskeletal Proteins in Neurological Disorders. *Cells, 9*(2),
E358. doi:10.3390/cells9020358
Muraro, N. I., & Moffat, K. G. (2006). Down-regulation of torp4a, encoding the Drosophila
homologue of torsinA, results in increased neuronal degeneration. *Journal of*
neurobiology, 66(12), 1338-1353. doi:10.1002/neu.20313
Nadeau, J. H. (2001). Modifier genes in mice and humans. *Nat Rev Genet, 2*(3), 165-174.
doi:10.1038/35056009
Naismith, T. V., Dalal, S., & Hanson, P. I. (2009). Interaction of torsinA with its major
binding partners is impaired by the dystonia-associated DeltaGAG deletion. *J Biol*
Chem, 284(41), 27866-27874. doi:10.1074/jbc.M109.020164
Nery, F. C., Armata, I. A., Farley, J. E., Cho, J. A., Yaqub, U., Chen, P., . . . Breakefield, X.
O. (2011). TorsinA participates in endoplasmic reticulum-associated degradation.
Nature communications, 2, 393-393. doi:10.1038/ncomms1383
Nery, F. C., Atai, N. A., da Hora, C. C., Kim, E. Y., Hettich, J., Mempel, T. R., . . . Irimia, D.
(2014). Microfluidic platform to evaluate migration of cells from patients with DYT1
dystonia. *Journal of neuroscience methods, 232*, 181-188.
doi:10.1016/j.jneumeth.2014.05.027
Nery, F. C., Zeng, J., Niland, B. P., Hewett, J., Farley, J., Irimia, D., . . . Breakefield, X. O.
(2008). TorsinA binds the KASH domain of nesprins and participates in linkage
between nuclear envelope and cytoskeleton. *Journal of cell science, 121*(Pt 20),
3476-3486. doi:10.1242/jcs.029454
Neuwald, A. F., Aravind, L., Spouge, J. L., & Koonin, E. V. (1999). AAA+: A class of
chaperone-like ATPases associated with the assembly, operation, and disassembly

of protein complexes. *Genome research, 9*(1), 27-43. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/9927482

Niethammer, M., Carbon, M., Argyelan, M., & Eidelberg, D. (2011). Hereditary dystonia as

a neurodevelopmental circuit disorder: Evidence from neuroimaging. *Neurobiology*

of disease, 42(2), 202-209. doi:10.1016/j.nbd.2010.10.010

Ohta, Y., Soucy, G., Phaneuf, D., Audet, J.-N., Gros-Louis, F., Rouleau, G. A., . . . Julien,

J.-P. (2016). Sex-dependent effects of chromogranin B P413L allelic variant as

disease modifier in amyotrophic lateral sclerosis. *Human molecular genetics,*

25(21), 4771-4786. doi:10.1093/hmg/ddw304

Ozelius, L., & Lubarr, N. (1993). DYT1 Early-Onset Isolated Dystonia. In M. P. Adam, H. H.

Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, & A. Amemiya

(Eds.), *GeneReviews((R)).* Seattle (WA): University of Washington, Seattle

University of Washington, Seattle. GeneReviews is a registered trademark of the

University of Washington, Seattle. All rights reserved.

Ozelius, L. J., Hewett, J. W., Page, C. E., Bressman, S. B., Kramer, P. L., Shalish, C., . . .

Breakefield, X. O. (1997). The early-onset torsion dystonia gene (DYT1) encodes

an ATP-binding protein. *Nat Genet, 17*(1), 40-48. doi:10.1038/ng0997-40

Petrucci, S., & Valente, E. M. (2013). Genetic issues in the diagnosis of dystonias.

Frontiers in neurology, 4, 34-34. doi:10.3389/fneur.2013.00034

Premi, E., Diano, M., Gazzina, S., Cauda, F., Gualeni, V., Tinazzi, M., . . . Borroni, B.

(2016). Functional Connectivity Networks in Asymptomatic and Symptomatic DYT1

Carriers. *Movement disorders : official journal of the Movement Disorder Society,*

31(11), 1739-1743. doi:10.1002/mds.26725

Rachidi, M., Delezoide, A. L., Delabar, J. M., & Lopes, C. (2009). A quantitative

assessment of gene expression (QAGE) reveals differential overexpression of

DOPEY2, a candidate gene for mental retardation, in Down syndrome brain

regions. *Int J Dev Neurosci, 27*(4), 393-398. doi:10.1016/j.ijdevneu.2009.02.001

Rodriguez-Quiroga, S., Gonzalez-Moron, D., Espay, A. J., & Kauffman, M. A. (2018).

Holmes Tremor-Like Phenotype in DYT1 Dystonia. *Mov Disord Clin Pract, 5*(2),

206-208. doi:10.1002/mdc3.12585

Rujito, L., Basalamah, M., Siswandari, W., Setyono, J., Wulandari, G., Mulatsih, S., . . .

Sutaryo, S. (2016). Modifying effect of XmnI, BCL11A, and HBS1L-MYB on clinical

appearances: A study on beta-thalassemia and hemoglobin E/beta-thalassemia

patients in Indonesia. *Hematol Oncol Stem Cell Ther, 9*(2), 55-63.

doi:10.1016/j.hemonc.2016.02.003

Saunders, C. A., Harris, N. J., Willey, P. T., Woolums, B. M., Wang, Y., McQuown, A.

J., . . . Luxton, G. W. G. (2017). TorsinA controls TAN line assembly and the

retrograde flow of dorsal perinuclear actin cables during rearward nuclear

movement. *The Journal of cell biology, 216*(3), 657-674.

doi:10.1083/jcb.201507113

Saunders, C. A., & Luxton, G. W. (2016). LINCing defective nuclear-cytoskeletal coupling

and DYT1 dystonia. *Cell Mol Bioeng, 9*(2), 207-216. doi:10.1007/s12195-016-0432-

0

Scheper, W., & Hoozemans, J. J. (2015). The unfolded protein response in
neurodegenerative diseases: a neuropathological perspective. *Acta Neuropathol,*
130(3), 315-331. doi:10.1007/s00401-015-1462-8

Shin, J. Y., Hernandez-Ono, A., Fedotova, T., Ostlund, C., Lee, M. J., Gibeley, S. B., . . .
Worman, H. J. (2019). Nuclear envelope-localized torsinA-LAP1 complex regulates
hepatic VLDL secretion and steatosis. *J Clin Invest, 130*, 4885-4900.
doi:10.1172/jci129769

Siokas, V., Dardiotis, E., Tsironi, E. E., Tsivgoulis, G., Rikos, D., Sokratous, M., . . .
Hadjigeorgiou, G. M. (2017). The Role of TOR1A Polymorphisms in Dystonia: A
Systematic Review and Meta-Analysis. *PLoS One, 12*(1), e0169934.
doi:10.1371/journal.pone.0169934

Sleigh, J. N., Rossor, A. M., Fellows, A. D., Tosolini, A. P., & Schiavo, G. (2019). Axonal
transport and neurological disease. *Nature reviews. Neurology, 15*(12), 691-703.
doi:10.1038/s41582-019-0257-2

Tanabe, L. M., Martin, C., & Dauer, W. T. (2012). Genetic background modulates the
phenotype of a mouse model of DYT1 dystonia. *PLoS One, 7*(2), e32245-e32245.
doi:10.1371/journal.pone.0032245

Thompson, M. L., Chen, P., Yan, X., Kim, H., Borom, A. R., Roberts, N. B., . . . Caldwell,
G. A. (2014). TorsinA rescues ER-associated stress and locomotive defects in C.
elegans models of ALS. *Disease models & mechanisms, 7*(2), 233-243.
doi:10.1242/dmm.013615

Trost, M., Carbon, M., Edwards, C., Ma, Y., Raymond, D., Mentis, M. J., . . . Eidelberg, D.
(2002). Primary dystonia: is abnormal functional brain architecture linked to
genotype? *Annals of neurology, 52*(6), 853-856. doi:10.1002/ana.10418

Vander Heyden, A. B., Naismith, T. V., Snapp, E. L., Hodzic, D., & Hanson, P. I. (2009).
LULL1 retargets TorsinA to the nuclear envelope revealing an activity that is
impaired by the DYT1 dystonia mutation. *Molecular biology of the cell, 20*(11),
2661-2672. doi:10.1091/mbc.e09-01-0094

Vulinovic, F., Lohmann, K., Rakovic, A., Capetian, P., Alvarez-Fischer, D., Schmidt, A., . . .
Seibler, P. (2014). Unraveling cellular phenotypes of novel TorsinA/TOR1A
mutations. *Hum Mutat, 35*(9), 1114-1122. doi:10.1002/humu.22604

Walter, M., Bonin, M., Pullman, R. S., Valente, E. M., Loi, M., Gambarin, M., . . .
Grundmann, K. (2010). Expression profiling in peripheral blood reveals signature
for penetrance in DYT1 dystonia. *Neurobiol Dis, 38*(2), 192-200.
doi:10.1016/j.nbd.2009.12.019

Wang, K., Li, M., & Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic
variants from high-throughput sequencing data. *Nucleic acids research, 38*(16),
e164-e164. doi:10.1093/nar/gkq603

Watanabe, N., Itakaoka, M., Seki, Y., Morimoto, T., Homma, K., Miyamoto, Y., &
Yamauchi, J. (2018). Dystonia-4 (DYT4)-associated TUBB4A mutants exhibit
disorganized microtubule networks and inhibit neuronal process growth.
Biochemical and biophysical research communications, 495(1), 346-352.
doi:10.1016/j.bbrc.2017.11.038

Weisheit, C. E., Pappas, S. S., & Dauer, W. T. (2018). Inherited dystonias: clinical features
and molecular pathways. *Handbook of clinical neurology, 147*, 241-254.
doi:10.1016/B978-0-444-63233-3.00016-6
Yamaguchi, T., Kubota, T., Kanematsu, T., Nakayama, K., Hirata, M., & Yamamoto, T.
(2004). Hypersensitivity to pentylenetetrazol-induced convulsion in mice lacking the
PLC-related inactive protein-1. *Brain Res, 1025*(1-2), 237-240.
doi:10.1016/j.brainres.2004.08.009
Yokoi, F., Dang, M. T., Mitsui, S., Li, J., & Li, Y. (2008). Motor deficits and hyperactivity in
cerebral cortex-specific Dyt1 conditional knockout mice. *J Biochem, 143*(1), 39-47.
doi:10.1093/jb/mvm191
Zech, M., Boesch, S., Jochim, A., Weber, S., Meindl, T., Schormair, B., . . . Winkelmann, J.
(2017). Clinical exome sequencing in early-onset generalized dystonia and large-
scale resequencing follow-up. *Mov Disord, 32*(4), 549-559. doi:10.1002/mds.26808
Zhang, F., & Lupski, J. R. (2015). Non-coding genetic variants in human disease. *Hum Mol*
Genet, 24(R1), R102-110. doi:10.1093/hmg/ddv259
Zhao, C., Brown, R. S. H., Chase, A. R., Eisele, M. R., & Schlieker, C. (2013). Regulation
of Torsin ATPases by LAP1 and LULL1. *Proceedings of the National Academy of*
Sciences of the United States of America, 110(17), E1545-E1554.
doi:10.1073/pnas.1300676110
Zhou, W., Brush, M. H., Choy, M. S., & Shenolikar, S. (2011). Association with
endoplasmic reticulum promotes proteasomal degradation of GADD34 protein. *The*
Journal of biological chemistry, 286(24), 21687-21696.
doi:10.1074/jbc.M110.212787

Table 1. Genomic Variants in the Exons, Promoter Regions and 3'-UTR of the *TOR1A* Gene between the Patients and the Other Family Members.

Subject Number[†]	Human Subject[†]	Variants in *TOR1A*	Site of Variant	Exonic Function	DNA Change	AA[‡] Change	AF[‡] in Person	AF in Taiwan[§]	AF in Dataset[§]	Interpretation
(1)	Patient	rs80358233	Exon 5	Nonframeshift deletion	904_906/ 907_909Δ GAG	ΔE302/ 303	0.5	Unknown	3.23^10⁻⁵	Pathogenic (Ozelius 2016)
(2)	Healthy Control	rs13300897	Promoter	-	C>T	-	0.5	0.174	0.1683	Polymorphism (Vulinovic 2014)
		rs2296793	Exon 2	Synonymous SNV	246G>A	A82A	0.5	0.1943	0.2253	Polymorphism (Vulinovic 2014)
		rs1182	3'-UTR	-	C>A	-	0.5	0.178	0.1666	Possible modifier (Siokas 2017)
(3)	Asympto-matic Carrier	rs13300897	Promoter	-	C>T	-	0.5	0.174	0.1683	Polymorphism (Vulinovic 2014)
		rs2296793	Exon 2	Synonymous SNV	246G>A	A82A	0.5	0.1943	0.2253	Polymorphism (Vulinovic 2014)
		rs80358233	Exon 5	Nonframeshift deletion	904_906/ 907_909Δ GAG	ΔE302/ 303	0.5	Unknown	3.23^10⁻⁵	Pathogenic (Ozelius 2016)
		rs1182	3'-UTR	-	C>A	-	0.5	0.178	0.1666	Possible modifier (Siokas 2017)
(6)	Asympto-matic Carrier	rs13300897	Promoter	-	C>T	-	0.5	0.174	0.1683	Polymorphism (Vulinovic 2014)
		rs2296793	Exon 2	Synonymous SNV	246G>A	A82A	0.5	0.1943	0.2253	Polymorphism (Vulinovic 2014)
		rs80358233	Exon 5	Nonframeshift deletion	904_906/ 907_909Δ GAG	ΔE302/ 303	0.5	Unknown	3.23^10⁻⁵	Pathogenic (Ozelius 2016)
(9)	Asympto-matic Carrier	rs80358233	Exon 5	Nonframeshift deletion	904_906/ 907_909Δ GAG	ΔE302/ 303	0.5	Unknown	3.23^10⁻⁵	Pathogenic (Ozelius 2016)
(11)	Patient	rs80358233	Exon 5	Nonframeshift deletion	904_906/ 907_909Δ GAG	ΔE302/ 303	0.5	Unknown	3.23^10⁻⁵	Pathogenic (Ozelius 2016)

† Subject 1: first patient, subject 2: the mother of the first patient, subject 3: the father of the first patient, subject 6: the aunt of the first patient, subject 9: the first son of the aunt of the first patient, subject 11: second patient.

‡ AA: amino acid, AF: allele frequency.

§Taiwan biobank, https://taiwanview.twbiobank.org.tw ; gnomAD (genome aggregation database), https://gnomad.broadinstitute.org.

869
870
871

Table 2. WES-identified DYT1 variants.

Autosomal Recessive Inheritance

Variants	Gene Symbol	DNA Change	AA Change	AF in Asia	AF in Patients	SIFT[†]	Poly-Phen2[‡]	Disease Association with Gene (PubMed)
rs36000545	AATK	T3488C	F1163S	0.5378		T	B	Unknown
rs55791176	AHNAK2	A3144C	E1048D	0.382		T	B	Charcot-Marie-Tooth disease
rs3772219	ARHGEF3	T1021G	L341V	0.462		D	B	Charcot-Marie-Tooth disease; osteoporosis
rs11689281	ASIC4	G1847T	R616L	0.6178		T	B	Unknown
rs11695248		T1856C	V619A	0.6191		T	B	
rs3774729	ATXN7	G2149A	V717M	0.4981		T	B	Spinocerebellar ataxia 7
rs61561984	C1orf195	A191T	Y64F	0.5359		N/A	N/A	Unknown
rs10804166	C2orf80	A454G	S152G	0.8426		T	B	46 xy gonadal dysgenesis
rs12269028	ZNF22-AS1	T188A	I63N	0.3994		N/A	N/A	Unknown
rs150150392	CCDC66	8_9insGGGTAAGCA	L3_G4delinsLGX	0.5544		N/A	N/A	Leber congenital amaurosis
rs866149312	CDK11A	T1973C	V658A	0.2041		T	B	Neuroblastoma
rs200982240		G1177A	D393N	0.2278	1,1	T	B	
rs6658335		G1108A	G370R	0.1702		D	B	
rs910122	CHGB	G533A	R178Q	0.5617		T	B	Pheochromocytoma; Glucagonoma
rs236152		C1058G	A353G	0.5614		T	B	
rs2230804	CHUK	G802A	V268I	0.4657		T	B	Cocoon syndrome
rs11715522	CX3CR1	T24G	F8L	0.6658		T	N/A	Age-related macular degeneration
rs4813043	DEFB128	A81T	K27N	0.4956		T	B	Unknown
rs1898883	DISP2	C139G	P47A	0.8517		T	B	Unknown
rs1898882		G167C	C56S	0.8383		D	B	
rs688906	DYNC2H1	A4238G	K1413R	0.7816		T	B	Short-rib thoracic dysplasia with or without polydactyly
rs589623		G8612A	R2871Q	0.7876		T	B	
rs13280444	FAM135B	C1445T	P482L	0.5081		T	B	Spinal and bulbar muscular atrophy
rs448012	FLT4	C2670G	H890Q	0.4763		D	B	Lymphatic malformation; Hemangioma

rs2229519	*GBE1*	A568G	R190G	0.4615		D	B	Glycogen storage disease iv
rs2792751	*GPAM*	A127G	I43V	0.7206		T	B	Unknown
rs3732215	*HJURP*	C629G	S210C	0.3879		T	B	Unknown
rs2286430		G226A	E76K	0.3926		T	B	
rs2072597	*KLF1*	T304C	S102P	0.6002		T	B	Congenital dyserythropoietic anemia, type iv
rs2429051	*KLF17*	G170C	S57T	0.676		T	B	Unknown
rs2485652		A467G	N156S	0.4608		T	B	
rs1064608	*MTCH2*	C841G	P281A	0.5837		D	P	Unknown
rs7995033	*MTMR6*	A955G	I319V	0.6064		T	B	Autosomal dominant Polycystic kidney disease
rs11556093	*NUP58*	G100A	A34T	0.6098		T	B	Unknown
rs6951485	*OR2A25*	G224A	S75N	0.6752		T	B	Unknown
rs2961135		G625C	A209P	0.6588		T	B	
rs1466684	*P2RY13*	C536	T179M	0.842		T	B	Unknown
rs35385129	*PVR*	C1171A	R391S	0.3621		T	B	Paralytic poliomyelitis
rs4795690	*RHBDL3*	G739A	V247M	0.3973		T	B	Unknown
rs1022478	*RIBC2*	C804G	F268L	0.5551		N/A	B	Unknown
rs1506418	*SERPINB11*	G16A	A6T	0.617		N/A	P	Unknown
rs1506419		T37A	W13R	0.6165		N/A	D	
rs12729295	*SLC35E2B*	G934A	V312I	0.4782		N/A	B	Unknown
rs2042791	*SPAG16*	A1083C	Q361H	0.4176		T	B	Rheumatoid arthritis
rs10151658	*SYNE2*	C15556A	L5186M	0.6646		T	B	Emery-dreifuss muscular dystrophy
rs3998860	*TET1*	A3369G	I1123M	0.8407		T	B	Acute myocardial infarction
rs614486	*TEX38*	T312G	D104E	0.7637		T	B	Unknown
rs2274791	*TTLL10*	G1733A	G578D	0.5856		D	B	Unknown
rs4838865	*TUBGCP6*	T1700C	L567S	0.8241		T	B	Chorioretinopathy-microcephaly syndrome
rs1801195	*WRN*	G3222T	L1074F	0.6186		T	B	Werner syndrome; Medulloblastoma
rs7258088	*ZFP28*	C128G	A43G	0.6304		T	B	Unknown
rs8100431	*ZNF414*	A194G	Q65R	0.5379		D	D	Unknown
rs4801177	*ZNF470*	C1253T	T418I	0.6776		T	B	Unknown

53 variants/43 genes								
Autosomal dominant inheritance								
Variants	**Gene Symbol**	**DNA Change**	**AA Change**	**AF in Asia**	**AF in Patients**	**SIFT**	**Poly-Phen2**	**Disease Association with Gene (PubMed)**
rs34400049	*AGGF1*	C2092A	P698T	0.2625	0.5,0.5	T	B	Klippel-trenaunay-weber syndrome
rs377198190	*AHNAK2*	T3424C	F1142L	0.0049		T	B	Charcot-Marie-Tooth disease
rs11845640	*AKAP6*	C4475T	A1492V	0.2563		T	B	Unknown
rs2614668	*AKAP13*	G4264A	A1422T	0.3056		T	B	Familial breast cancer
rs3027232	*ALOXE3*	C32T	P11L	0.4412		T	B	Ichthyosis
rs28362581	*AMPD2*	G244A	A82T	0.3146		D	B	Pontocerebellar hypoplasia
rs1465582	*ANKLE1*	T1933G	L645V	NA		N/A	N/A	Unknown
rs1042034	*APOB*	G13013A	S4338N	0.264		T	B	Hypobetalipoproteinemia
rs6668968	*AQP10*	G44A	R15Q	0.2246		T	B	Unknown
rs6685323		C367T	H123Y	0.2265		T	B	
rs3733662	*ARHGEF37*	C1756A	P586T	0.2975		T	B	Unknown
rs8066889	*ARL16*	A28C	S10R	0.1959		T	B	Unknown
rs33995001	*ATP10D*	C128T	T43I	0.1446		T	B	Unknown
rs3750690	*ATRNL1*	A2822T	Q941L	0.0092		T	D	Unknown
rs115476782	*BTNL10*	G376T	A126S	0.0334		N/A	N/A	Unknown
rs2274067	*C1orf131*	C82G	L28V	0.0854		D	D	Unknown
rs3813728	*C1R*	G781A	G261R	0.0932		T	D	Ehlers-danlos syndrome
rs1801046		C455T	S152L	0.3511		T	B	
rs10951942	*C7orf57*	G220T	A74S	0.2769		D	D	Unknown
rs12657663	*CAMLG*	G232A	V78I	0.2357		T	B	Congenital diaphragmatic hernia
rs6886	*CAPG*	A911G	H304R	0.4653		T	B	Breast Cancer Biomarker for Bone Metastasis
rs1801449	*CAPN3*	G706A	A236T	0.1279		T	B	Limb-girdle muscular dystrophy
rs76069883	*CCDC74A*	G688A	G230S	0.0067		T	B	Unknown
rs3177472	*CCDC74B*	G839A	R280H	0.4827		T	B	Unknown
rs2259332		T728C	V243A	0.4602		T	B	
rs7226091	*CCDC137*	C381G	H127Q	0.1965		T	B	Unknown
rs11546630		G686A	R229Q	0.1952		T	B	

rs11546631		C844T	R282W	0.1955		T	B	
rs6740879	CCDC138	G329A	R110K	0.0458		T	B	Unknown
rs12606658	CCDC178	G124A	A42T	0.0297		D	B	Unknown
rs2230552	CCT6B	T143C	V48A	0.5422		D	D	Pelvic varices
rs3809727	CDRT1	C734G	A245G	0.4412		T	B	Charcot-Marie-Tooth disease
rs16959164	CEACAM20	C1064T	S355L	0.0721		T	B	Unknown
rs3734381	CEP85L	A409G	S137G	0.3075		T	B	Myeloproliferative neoplasm
rs6081901	CFAP61	G1105A	V369I	0.2128		T	B	Unknown
rs16858780	CHRD	A1888C	M630L	0.3296		T	B	Cornelia de lange syndrome
rs3740129	CHST3	G1070A	R 357Q	0.087		T	B	Spondyloepiphyseal Dysplasia with Congenital Joint Dislocations
rs3743193	CHSY1	C1075T	P359S	0.2051		T	B	Brachydactyly
rs34964084	CPZ	C17T	P6L	0.1768		T	B	Unknown
rs3738952	CUL3	G1501A	V501I	0.2714	0.5,1	T	B	Pseudohypoaldosteronism
rs1056827	CYP1B1	G355T	A119S	0.2131		N/A	B	Primary congenita glaucoma
rs10012		C142G	R48G	0.2135		N/A	B	
rs2261144	CYP2A7	T950C	M317T	0.2643		T	B	Unknown
rs2544809	DBN1	A1336G	I446V	0.2506		T	B	Alzheimer disease
rs368539076	DHRS4L2	203_204insA	L68fs	0.2159	0.5,0.5	N/A	N/A	Unknown
rs869801	DOCK1	G5440A	A1814T	0.1191		T	B	Chromosome 10q26 Deletions Syndrome
rs3746866	DOP1B	C3446A	P1149H	0.2389		D	P	Down syndrome
rs7813708	FAM83A	G541A	A181T	0.2311	0.5,1	T	B	Pancreatic cancer
rs17429619	FAM114A1	G706A	V236I	0.1045	0.5,0.5	T	B	Unknown
rs16858529	FCRLB	G812T	R271L	0.448	0.5,1	D	B	IgA nephropathy
rs1042229	FPR1	T576G	N192K	0.2784		T	B	Periodontitis
rs35870000	FREM1	G3634T	A1212S	0.1957	0.5,0.5	T	B	Bifid nose renal agenesis; anorectal malformations; manitoba oculotrichoanal syndrome

rs778805	*FUT6*	C370T	P124S	0.5813		D	B	Fucosyltransferase 6 deficiency;gastrointestinal carcinoma
rs2267161	*GAL3ST1*	G85A	V29M	0.3309		T	B	Unknown
rs61753060	*GEMIN4*	A773G	Q258R	0.058		T	B	Neurodevelopmental Disorder with Microcephaly, Cataracts, And Renal Abnormalities
rs696217	*GHRL*	C178A	L60M	0.1922		T	D	Eating disorder
rs75027378	*GLOD4*	C842A	A281E	0.0589		T	B	Unknown
rs3796130	*GP9*	G466A	A156T	0.2373		T	B	Bernard-Soulier syndrome; Gray platelet syndrome
rs1042303	*GPLD1*	A2080G	M694V	0.2627		T	B	Autism spectrum disorders
rs3841128	*GRIA1*	31dupC	P10fs	0.0309		N/A	N/A	Depression; Status epilepticus
rs56058441	*HGS*	G2197T	A733S	0.198		T	B	Unknown
rs2295778	*HIF1AN*	C121G	P41A	0.2571		T	B	Hypoxia; Nephronophthisis
rs397814627	*HOXD9*	794_795insGCA	P265delinsPQ	0.3988		N/A	N/A	Brachydactyly-syndactyly syndrome
rs2296436	*HPS1*	A1448G	Q483R	0.154		T	B	Hermansky-Pudlak syndrome
rs2296434		C1112G	P371R	0.1908		T	D	
rs34533614		C808T	P270S	0.1912		N/A	N/A	
rs142594836	*HRNR*	G2296A	G766S	0.042		T	B	Unknown
rs12094334	*IGFN1*	G2087A	G696D	0.3513		D	N/A	Polypoidal choroidal vasculopathy
rs12063867		A2947G	M983V	0.3511		D	N/A	
rs7551098		C3257T	A1086V	0.3503		T	N/A	
rs7551538		G3517T	A1173S	0.3449		D	N/A	
rs12070918		A7196G	D2399G	0.3499		T	N/A	
rs35267671	*INPP5B*	G16A	G6S	0.3246		D	B	Lowe oculocerebrorenal syndrome
rs7251	*IRF3*	G461C	S154T	0.3288		T	B	Acute Encephalopathy, Infection-induced
rs75304543	*JADE2*	G42T	L14F	0.3554		N/A	N/A	Unknown
rs17618244	*KLB*	G2183A	R728Q	0.1819		T	B	Unknown
rs198977	*KLK2*	C442T	R148W	0.2004		T	B	Prostate cancer
rs201142403	*KRT6A*	T745C	F249L	0.1457		T	B	Pachyonychia congenita

rs199613662		G722A	G241D	0.1127		D	B	
rs201663666		G721A	G241S	0.1097		T	B	
rs652423		A680G	N227S	0.3876		T	B	
rs61745883	KRT6B	G332A	G111D	0.3331		D	D	Pachyonychia congenita
rs61914500		G262A	G88R	0.2388		T	B	
rs2226548	KRTAP13-4	G175A	A59T	0.2884		T	B	Unknown
rs2832873	KRTAP15-1	C127A	L43M	0.2884		D	P	Unknown
rs2298437	KRTAP19-4	A143G	Y48C	0.3241		T	B	Unknown
rs28622470	LIMCH1	C1238T	T413M	0.0972		D	D	Unknown
rs73135482		C1429A	L477I	0.0969		T	B	
rs60455691	LIPT2	C158T	A53V	0.1792		T	B	Mitochondrial lipoylation Defect associated with severe neonatal encephalopathy
rs3816614		G4937A	R1646Q	0.3798	0.5,1	T	B	Cenani-Lenz syndactyly syndrome; congenital myasthenic syndrome
rs2306029	LRP4	A4660G	S1554G	0.2342		D	B	
rs6485702		A3256G	I1086V	0.2827		T	B	
rs17286758	LRRC2	A247G	T83A	0.0999		T	B	Unknown
rs2042919	LRRC8E	A155G	E52G	0.2382		T	B	Unknown
rs2115108		T182C	M61T	0.2654		T	B	
rs2302607	METTL22	G655A	A219T	0.3574		T	B	Unknown
rs78616323	MICAL3	C3827T	T1276I	0.1596		D	B	Joubert syndrome
rs75658007	MRM3	G427A	V143I	0.0537	0.5,0.5	D	B	Unknown
rs80220493		T428A	V143D	0.0537		D	P	
rs11546280	MRPL12	T313C	S105P	0.1969		T	B	Brain glioblastoma multiforme; Brain Cancer
rs2216662		G29725A	V9909I	0.5808		D	B	
rs1833778		G28504A	A9502T	0.5753		T	B	
rs2547064		A23768C	D7923A	0.2805		T	B	
rs1867691		A21814G	I7272V	0.2801		T	B	
rs2121133		T15133C	S5045P	0.2799		T	B	
rs2591590	MUC16	A14704G	I4902V	0.2782		D	B	Ovarian cancer
rs2591591		C12496A	H4166N	0.2803		D	B	
rs2547068		A12229T	T4077S	0.2803	0.5,1	T	B	
rs2591592		A12100T	I4034F	0.2855		T	P	
rs2591593		G11477A	G3826E	0.28		D	B	
rs2547072		C11363T	T3788I	0.2793		D	B	
rs2591594		G10718A	R3573H	0.282		T	B	

rs2547074		G10588A	V3530I	0.2802		D	B	
rs2547075		A10517C	K3506T	0.279		T	B	
rs2547076		G10496T	R3499M	0.2798		T	B	
rs1862462		C10010T	S3337L	0.2817		D	B	
rs2591597		A9643G	S3215G	0.2818		T	B	
rs73168398	*MUC17*	G12998A	R4333Q	0.1058		T	B	Biliary papillomatosis; Colorectal cancer
rs3817552	*MYBPC1*	C1365G	H455Q	0.2979		D	D	Arthrogryposis; Lethal Congenital Contracture Syndrome
rs6421985	*NLRP6*	A487C	M163L	0.2435		T	B	Unknown
rs7482965		A1082T	Y361F	0.254		T	B	
rs56128139	*NME8*	T1478C	I493T	0.1158		D	B	Primary ciliary dyskinesia
rs2270182	*NRAP*	A1451T	N484I	0.1755		T	P	Myopathy, myofibrillar
rs2275799		G844A	A282T	0.1998		T	B	
rs75155858	*NRG1*	G1376T	G459V	0.3453		D	P	Schizophrenia; mucinous lung adenocarcinoma
rs200668592	*NRG2*	1779_1784del	593_595del	0.0883		N/A	N/A	Charcot-Marie-Tooth disease
rs769427	*OR1A1*	C853T	P285S	0.1247	0.5,0.5	D	P	Unknown
rs4836891	*OR1J2*	G494A	R165Q	0.2283		T	B	Unknown
rs142107755	*OR2T33*	G661T	A221S	0.0383		T	B	Unknown
rs200877558		T479C	V160A	0.0149		T	B	
rs227787	*OR3A3*	A949G	K317E	0.3538		D	B	Unknown
rs12885778	*OR4K1*	G266A	R89H	0.2635		T	B	Unknown
rs34394400		C910T	R304C	0.1892		D	B	
rs2318279	*OR4N2*	C397T	P133S	0.1125		T	B	Unknown
rs80295194	*OR10G3*	G875A	R292Q	0.0309		D	P	Unknown
rs7114672	*OR56A5*	G310A	V104M	0.2921		N/A	B	Unknown
rs208294	*P2RX7*	T463C	Y155H	0.3897		N/A	N/A	Extrapulmonary tuberculosis; tularemia
rs726684	*PCDHGA8*	T47G	L16R	0.1383		D	D	Unknown
rs2074912	*PCDHGC5*	A1709G	D570G	0.1293		T	B	Unknown
rs17208425		A2618G	E873G	0.1289		N/A	B	
rs7234309	*PIEZO2*	G4060A	V1354I	0.4988		T	N/A	Marden-Walker syndrome; Distal Arthrogryposis

rs73403546	PLA2G4F	C754G	L252V	0.1179		T	B	Unknown
rs1064213	PLCL1	G1999A	V667I	0.1865	0.5,1	D	D	Unknown
rs7424029	POTEE	G2601T	E867D	0.1611		T	B	Unknown
rs62178369		G2918A	G973D	0.3833		D	D	
rs2599794	POTEF	G2601T	E867D	0.4239		T	B	Unknown
rs2897665		A337G	S113G	0.5779		T	B	
rs143023559	PPFIBP2	893_895 del	298_299 del	0.0718		N/A	N/A	Unknown
rs3786734	PPP1R15A	G94A	A32T	0.1741		D	D	Unknown
rs1769774	PRAMEF1	C652T	P218S	0.3745	0.5,0.5	T	B	Unknown
rs1052908		A423C	R141S	0.4245		D	B	
rs72819488	PROM2	G1537A	G513S	0.2172		T	P	Unknown
rs7260222	PTOV1	C74T	S25L	0.4574		N/A	B	A Marker of Aggressive Diseases in Carcinomas
rs117766916	RAET1L	C79G	R27G	0.0296		T	B	Unknown
rs3744872	RBFA	A733C	N245H	0.4944		T	B	Unknown
rs112636230	RBMXL1	A120G	I40M	0.2333		T	B	Unknown
rs10963	RBP5	G55A	D19N	0.4646	0.5,1	T	B	Total anomalous pulmonary venous return
rs78143373	RHBDF1	A2017G	I673V	0.0867		T	B	Palmoplantar keratoderma
rs148731719	RNF213	G13195A	A4399T	0.0524		T	B	Moyamoya disease; anaplastic large cell lymphoma
rs200943820	RSPH10B	C443T	T148M	0.0975	0.5,0.5	T	D	Unknown
rs2295769	SERPINB6	A310G	M104V	0.2679		T	B	Autosomal recessive non-syndromic sensorineural deafness
rs6092	SERPINE1	G43A	A15T	0.0885		T	B	Plasminogen activator inhibitor-1 deficiency
rs1136287		C215T	T72M	0.5702	0.5,1	T	B	
rs10409962	SIGLEC8	T508C	S170P	0.11		T	B	Unknown
rs11150813	SLC25A10	G892A	V298I	0.3398		N/A	N/A	Unknown
rs13259978	SLC7A2	G82C	D28H	0.0957		T	B	Lysinuric protein intolerance
rs576516	SMAP1	T1247C	M416T	0.2882	0.5,0.5	T	B	Retinitis pigmentosa
rs758896527	SPATA31C1	211_221 del	H71fs	0.0269		N/A	N/A	Unknown
rs61759822	STPG3	C652T	L218F	0.3067		T	P	Unknown
rs10883859	TAF5	T388G	S130A	0.4215		T	B	Unknown

rs1052692	TCF3	G1291A	G431S	0.218		T	B	Agammaglobulinemia
rs12498609	TET2	C86G	P29R	0.2077		D	P	Refractory anemia; Myelodysplastic syndrome
rs1025806	TEX38	C596T	A199V	0.3389	0.5,1	T	B	Unknown
rs3827816	TNC	G1813A	V605I	0.3891	0.5,0.5	T	B	Deafness; Bullous Keratopathy
rs2269495	TNIP2	C938T	A313V	0.3585		T	B	Unknown
rs16847812	TNN	G865A	D289N	0.241	0.5,1	T	B	Dysplastic nevus syndrome
rs6694078		A2575G	M859V	0.3444		T	B	
rs12369033	TNS2	G29C	R10T	0.1079		D	B	Unknown
rs371929937	TPSAB1	A8G	N3S	0.169		T	B	Systemic mastocytosis
rs1131877	TRAF3	T386C	M129T	0.4163		T	B	Acute Encephalopathy, Infection-induced
Chr19: 6751057§	TRIP10	T1513A	S505T	NA	0.5,0.5	N/A	B	Wiskott-aldrich syndrome
rs3762735	TRMT10C	C167G	P56R	0.1462		T	B	Combined oxidative phosphorylation deficiency; mitochondrial metabolism disease
rs2072394	TRUB2	G145C	V49L	0.0605		T	B	Dyskeratosis congenita
rs33970858	TSNARE1	G164C	R55P	0.3317		T	B	Schizophrenia
rs7814359		T52C	F18L	0.3292		T	B	
rs34379910	TSPAN10	T652C	Y218H	0.1994	0.5,1	N/A	D	Unknown
rs1052422	TUBA3E	T1204C	W402R	0.4738		N/A	B	Microlissencephaly
rs13000249		C661A	R221S	0.4252		N/A	P	
rs13000721		C377T	A126V	0.4655	0.5,0.5	N/A	B	
rs3863907		G302A	S101N	0.4698		N/A	B	
rs307658	UBAP2	A1016G	N339S	0.1675		T	B	Unknown
rs2072767	UNC93A	G748A	V250I	0.2382		T	B	Unknown
rs9459921		G1083A	M361I	0.2591		D	B	
rs3744793	USP36	G811A	V271I	0.5318	0.5,1	T	B	Unknown
rs15818	VPS11	A2636G	K879R	0.1788		N/A	B	Hypomyelinating leukodystrophy
rs597371	VWA2	A392G	E131G	0.3567	0.5,0.5	T	B	Type 1 diabetes mellitus; A marker for colon cancer
rs6942733	ZAN	T3035G	L1012R	0.2833	0.5,1	N/A	B	Deafness
rs76337191	ZIC5	C329A	A110E	0.1409	0.5,0.5	T	B	Holoprosencephaly

rs61734609	*ZNF221*	T1598C	L533P	0.1292		T	B	Unknown
rs6509138	*ZNF223*	C412A	L138I	0.3185		T	B	Ovarian carcinomas
rs2249769	*ZSCAN30*	A152C	Q51P	0.2609	0.5,1	T	B	Unknown

201 variants/149 genes

De novo mutation

Variants	Gene Symbol	DNA Change	AA Change	AF in Asia	AF in Patients	SIFT	Poly-Phen2	Disease Association with Gene (PubMed)
rs200227742	*CDK11B*	G337A	G113R	0.0758		N/A	B	Neuroblastoma
rs779250776		1531_1533del	511_511del	0		N/A	N/A	
rs68177477	*GOLGA6L2*	G1509C	Q503H	0		T	B	
rs76062343		C1465G	Q489E	0.0054		T	B	Unknown
rs75486959		A1460T	E487V	0.001		T	B	
rs74565846		G1459A	E487K	0	0.5,0.5	D	B	
Chr11: 71527323§	*KRTAP5-7*	A23G	E8G	0.0008		D	B	Unknown
Chr2: 15940678§	*MYCN*	G92C	R31P	N/A		N/A	N/A	Cerebral primitive neuroectodermal tumor
Chr2: 15940685§		100_101del	G34fs	N/A		N/A	N/A	
rs774139549	*VEGFC*	C571G	P191A	N/A		N/A	P	Lymphedema

10 variants/5 genes

†. SIFT: T: tolerable, D: deleterious, N/A: not applicable.

‡. Poly-Phen2: B: benign, P: possibly damaging, D: damaging, N/A: not applicable.

§. Genomic position without registered SNP ID available.

Table 3. DYT1 variant-harboring genes that encode cytoskeleton-associated proteins.

Microtubules

Variant(s)	Gene Symbol	Protein	AF in the two patients	References
rs3774729	*ATXN7*	Ataxin-7	1,1	PMID: 22100762
rs150150392	*CCDC66*	Coiled-Coil Domain Containing 66	1,1	PMID: 28235840
rs76069883	*CCDC74A*	Coiled-coil domain-containing protein 74A	0.5,0.5	PMID: 31521166
rs3177472 rs2259332	*CCDC74B*	Coiled-coil domain-containing protein 74B	0.5,0.5	
rs6740879	*CCDC138*	Coiled-coil domain-containing protein 138	0.5,0.5	PMID: 31304627
rs3734381	*CEP85L*	Centrosomal protein of 85 kDa-like	0.5,0.5	PMID: 21399614
rs6081901	*CFAP61*	Cilia- and flagella-associated protein 61	0.5,0.5	PMID: 30122541
rs3738952	*CUL3*	Cullin-3	0.5,1	PMID: 19995937
rs869801	*DOCK1*	Dedicator of cytokinesis protein 1	0.5,0.5	PMID: 24637113
rs688906 rs589623	*DYNC2H1*	Dynein Cytoplasmic 2 Heavy Chain 1	1,1	PMID: 25470043
rs78616323	*MICAL3*	Microtubule-associated monoxygenase, calponin, and LIM domain-containg 3	0.5,0.5	PMID: 27528609
rs56128139	*NME8*	Thioredoxin domain-containing protein 3	0.5,0.5	PMID: 17360648
rs35385129	*PVR*	Poliovirus receptor	1,1	PMID: 20964795
rs2042791	*SPAG16*	Sperm-associated antigen 16 protein	1,1	PMID: 21655194
Chr19: 6751057	*TRIP10*	Cdc42-interacting protein 4	0.5,0.5	PMID: 11069762
rs2274791	*TTLL10*	Inactive polyglycylase	1,1	PMID: 19427864
rs1052422 rs13000249 rs13000721	*TUBA3E*	Tubulin alpha-3E chain	0.5,0.5	PMID: 17543498

rs3863907				
rs4838865	*TUBGCP6*	γ-tubulin complex component 6	1,1	PMID: 11694571

23 variants/18 genes

Actin Filaments

Variant(s)	Gene Symbol	Protein	AF in the two patients	References
rs2614668	*AKAP13*	A-kinase anchor protein 13	0.5,0.5	PMID: 24183240
rs6886	*CAPG*	Macrophage-capping protein	0.5,0.5	PMID: 18266911
rs1801449	*CAPN3*	Calpain-3	0.5,0.5	PMID: 11950589
rs2230552	*CCT6B*	Chaperonin Containing TCP1 Subunit 6B	0.5,0.5	PMID: 9013858
rs2544809	*DBN1*	Drebrin	0.5,0.5	PMID: 20215400
rs869801	*DOCK1*	Dedicator of cytokinesis protein 1	0.5,0.5	PMID: 25452388
rs28622470 rs73135482	*LIMCH1*	LIM and calponin homology domains-containing protein 1	0.5,0.5	PMID: 28228547
rs3817552	*MYBPC1*	Myosin-binding protein C, slow-type	0.5,0.5	PMID: 8375400
rs2270182 rs2275799	*NRAP*	Nebulin-related-anchoring protein	0.5,0.5	PMID: 19233165
rs10151658	*SYNE2*	Nesprin-2	1,1	PMID: 22945352

12variants/10 genes

Intermediate Filaments

Variant(s)	Gene Symbol	Protein	AF in the two patients	References
rs201142403 rs199613662 rs201663666	*KRT6A*	Keratin, type II cytoskeletal 6A	0.5,0.5	PMID: 7543104
rs652423 rs61745883 rs61914500	*KRT6B*	Keratin, type II cytoskeletal 6B	0.5,0.5	PMID: 9618173
Chr11: 71527323§	*KRTAP5-7*	Keratin-associated protein 5-7	0.5,0.5	PMID: 31691815

rs2226548	*KRTAP13-4*	Keratin-associated protein 13-4	0.5,0.5	
rs2832873	*KRTAP15-1*	Keratin-associated protein 15-1	0.5,0.5	
rs2298437	*KRTAP19-4*	Keratin-associated protein 19-4	0.5,0.5	
10 variants/6 genes				
Total 45 variants/34 genes				

Table 4. DYT1 variant-harboring genes that encode proteins involved in protein and lipid metabolism and endoplasmic reticulum function.

Endoplasmic Reticulum function and protein metabolism

Variant(s)	Gene Symbol	Protein	AF in the two patients	References
rs910122 rs236152	*CHGB*	Chromogranin B	1,1 1,1	PMID: 31691815
rs3746866	*DOP1B*	DOP1 leucine zipper-like protein B	0.5,0.5	PMID: 16301316
rs7995033	*MTMR6*	Myotubularin-related protein 6	1,1	PMID: 19038970
rs1466684	*P2RY13*	P2Y purinoceptor 13	1,1	PMID: 30576484
rs3786734	*PPP1R15A*	Protein phosphatase 1 regulatory subunit 15A	0.5,0.5	PMID: 21518769

6 variants/5 genes

Lipid Metabolism

Variant(s)	Gene Symbol	Protein	AF in the two patients	References
rs3027232	*ALOXE3*	Hydroperoxide isomerase ALOXE3	0.5,0.5	PMID: 21558561
rs1042034	*APOB*	Apolipoprotein B-100	0.5,0.5	PMID: 15797858
rs1056827 rs10012	*CYP1B1*	Cytochrome P450 1B1	0.5,1	PMID: 15258110
rs2261144	*CYP2A7*	Cytochrome P450 2A7	0.5,0.5	PMID: 21873635
rs13280444	*FAM135B*	Protein FAM135B	1,1	PMID: 21873635
rs2267161	*GAL3ST1*	Galactosylceramide sulfotransferase	0.5,0.5	PMID: 25151383
rs2792751	*GPAM*	Glycerol-3phosphate acyltransferase 1, mitochondrial	1,1	PMID: 18238778
rs7995033	*MTMR6*	Myotubularin-related protein 6	1,1	PMID: 22647598
rs73403546	*PLA2G4F*	Cytosolic phospholipase A2 zeta	0.5,0.5	PMID: 21873635
rs1064213	*PLCL1*	Inactive phospholipase C-like protein 1	0.5,1	PMID: 17254016

11 variants/10 genes

Table 5. DYT1 variant-harboring genes that encode proteins associated with human neuropsychiatric disorders or neuromuscular diseases

Variant(s)	Gene Symbol	AA change	AF in Asia	Disease association	References
rs55791176	*AHNAK2*	E1048D	0.382	Charcot-Marie-Tooth disease, Demyelinating, Type 4F	PMID: 31011849
rs28362581	*AMPD2*	A82T	0.3146	Pontocerebellar hypoplasia, Type 9	PMID: 29463858
rs6755527	*ARHGEF3*	L341V	0.462	Charcot-Marie-Tooth disease, Type 4H	PMID: 14508709
rs3774729	*ATXN7*	V717M	0.4981	Spinocerebellar ataxia 7	PMID: 30473770
rs1801449	*CAPN3*	A236T	0.1279	Limb-girdle Muscular dystrophy	PMID: 31540302
rs200227742	*CDK11B*	G113R	0.0758	Neurobloastoma	PMID: 7777541
rs3809727	*CDRT1*	A245G	0.4412	Charcot-Marie-Tooth disease, Type Ia	PMID: 11381029
rs2544809	*DBN1*	I446V	0.2506	Alzheimer disease	PMID: 28597477
rs3746866	*DOP1B*	P1149H	0.2389	Down syndrome	PMID: 16303751
rs13280444	*FAM135B*	P482L	0.5081	Spinal and bulbar muscular atrophy, X-Linked	PMID: 30391288
rs2229519	*GBE1*	R190G	0.4615	Polyglucosan body neuropathy, adult form	PMID: 25544507
rs61753060	*GEMIN4*	Q258R	0.058	Neurodevelopmental disorder with microcephaly, cataracts, renal abnormalities, and microcephaly	PMID: 25558065
rs1042303	*GPLD1*	M694V	0.2627	Autism spectrum disorders	PMID: 25448322
rs3841128	*GRIA1*	P10fs	0.0309	Depression; Status epilepticus	PMID: 22057216
rs7251	*IRF3*	S154T	0.3288	Acute encephalopathy, Infection-induced	PMID: 26216125
rs60455691	*LIPT2*	A53V	0.1792	Neonatal severe encephalopathy with lactic acidosis and brain abnormalities and lipoic acid	PMID: 28757203

				biosynthesis defect	
rs3816614 rs2300629 rs6485702	*LRP4*	R1646Q S1554G I1086V	0.3798 0.2342 0.2827	Congenital myasthenic syndrome	PMID: 28825343
rs78616323	*MICAL3*	T1276I	0.1596	Joubert syndrome (Cerebelloparenchymal disorder)	PMID: 26485645
rs11546280	*MRPL12*	S105P	0.1969	Brain glioblastoma multiforme and Brain cancer	PMID: 26781422
Chr2: 15940678 15940685	*MYCN*	R31P G34fs	NA	Cerebral primitive neuroectodermal tumor	PMID: 28453467
rs75155858	*NRG1*	G459V	0.3453	Schizophrenia	PMID: 30500411
rs2270182 rs2275799	*NRAP*	N484I A282T	0.1755 0.1998	Myopathy, myofibrillar	PMID: 30986853
rs200668592	*NRG2*	593_595 del	0.0883	Charcot-Marie-Tooth disease, demyelinating form	PMID: 10369162
rs148731719	*RNF213*	A4399T	0.0524	Moyamoya disease	PMID: 29387438
rs35385129	*PVR*	R391S	0.3621	Paralytic poliomyelitis	PMID: 11597452
rs10151658	*SYNE2*	L5186M	0.6646	Emery-Dreifuss Muscular dystrophy	PMID: 21496632
rs1131877	*TRAF3*	M129T	0.4163	Acute encephalopathy, Infection-induced	PMID: 20832341
rs33970858 rs7814359	*TSNARE1*	R55P F18L	0.3317 0.3292	Schizophrenia	PMID: 27668389
rs1052422 rs13000249 rs13000721 rs3863907	*TUBA3E*	W402R R221S A126V S101N	0.4738 0.4252 0.4655 0.4698	Microlissencephaly	PMID: 17571022
rs4838865	*TUBGCP6*	L567S	0.8241	Microcephaly with chorioretinopathy	PMID: 31077665
rs15818	*VPS11*	K879R	0.1788	Hypomyelinating leukodystrophy	PMID: 27473128
rs76337191	*ZIC5*	A110E	0.1409	Holoprosencephaly	PMID: 20531442
Total 40 variants/32 genes					

Figure legends

Figure 1. Two patients and their family pedigrees and Sanger sequencing data. Family pedigree (A,C,E) and Sanger sequencing data (B,D,F) of (A,B) 10 family members of *TOR1A* gene (c.904_906/907_909ΔGAG, p. ΔE302/E303), (C,D) 4 family (core family of the first family) members of *TOR1A* gene (c.646G>C, p.D216H) and (E,F) second patient of *TOR1A* gene (c.904_906/907_909ΔGAG, p. ΔE302/E303). (A,C,E) The arrows point out the two probands. The numbers within parentheses are the order of Sanger sequencing data and the numbers under the box/circle show the age (years old). The question marks within the box/circle indicate the unknown status because we don't have the DNAs sample for study. Triangles denote lack of gender information.

Figure 2. Multiple hits (variants) in the nuclear-cytoskeletal coupling network. TOR1A-LAP1 (or LULL1) heterohexamer regulates the assembly and function of LINC complex. The location of the defects at TOR1A (ΔE302/E303) and variants found in microtubules, actin filaments, intermediate filaments, nesprin-2, nuclear pore complex (NPC), endoplasmic reticulum (ER), and lipid metabolism. LINC complex (the linker of nucleoskeleton and cytoskeleton, consisting of KASH domain and SUN proteins), T (TOR1A), L (LAP1 or LULL1), KASH domain (Klarsicht, ANC-1, and Syne homology domain), SUN 1 and SUN 2 (SUN (Sad1, UNC-84) domain-containing protein 1 and 2), ONM (outer nuclear membrane), INM (inner nuclear membrane).

Figure S1. Workflow to identify genome variants outside the TOR1A Locus from autosomal recessive inheritance, autosomal dominant inheritance, and *de novo* mutation. Six WES samples include patients (subject 1,11), the mother of the patient 1 (subject 2), the father of the patient 1 (subject 3), the aunt of the patient 1 (subject 6), the first son of the aunt of the patient 1 (subject 9)

FIGURE 1

A

TOR1A (c.904_906/907_909ΔGAG, p. ΔE302/E303):
Patient 1 ■ Autosomal dominant carrier ⊡ ⊙

B

(1) c.904_906/907_909 ΔGAG Heterozygous
(2) Wild type
(3) c.904_906/907_909 ΔGAG Heterozygous
(4) Wild type
(5) Wild type
(6) c.904_906/907_909 ΔGAG Heterozygous
(7) Wild type
(8) Wild type
(9) c.904_906/907_909 ΔGAG Heterozygous
(10) Wild type

C

TOR1A (c.646G>C, p.D216H)

D

(1) Wild type
(2) Wild type
(3) Wild type
(4) Wild type

E

P2 → 40
TOR1A (c.904_906/907_909ΔGAG, p. ΔE302/E303):
Patient 2 ■

F

(11) c.904_906/907_909 ΔGAG Heterozygous

FIGURE 2