1 Optimization of skeletal protein preparation for LC-MS/MS sequencing yields
2 additional coral skeletal proteins in *Stylophora pistillata*
3

4 Yanai Peled[1*], Jeana Drake[1*^], Assaf Malik[1], Ricardo Almuly[1], Maya Lalzar[2], David
5 Morgenstern[3], Tali Mass[1+]
6
[1]Marine Biology Department, University of Haifa, Haifa, Israel
[2]Bioinformatics Core Unit, University of Haifa, Haifa, Israel
[3]De Botton Protein Profiling Institute of the Nancy and Stephen Grand Israel National Center for
Personalized Medicine, Weizmann Institute of Science, Israel
*equal contribution
^Corresponding author during review: jeanadrake@g.ucla.edu
+Corresponding author upon publication: tmass@univ.haifa.ac.il

Abstract

Stony corals generate their calcium carbonate exoskeleton in a highly controlled

biomineralization process mediated by a variety of macromolecules including

proteins. Fully identifying and classifying these proteins is crucial to understanding

their role in exoskeleton formation, yet no optimal method to extract and isolate and

characterize coral skeletal proteins has been established and their complete

composition remains obscure. Here, we tested four skeletal protein extraction

protocols using acetone precipitation and ultrafiltration dialysis filters to present a

comprehensive scleractinian coral skeletal proteome. We identified a total of 60

proteins in the coral skeleton, 44 of which were not present in previously published

stony coral skeletal proteomes. Extracted protein treatment protocols carried out in

this study revealed that there is no "one optimal method" and each protocol revealed a

unique set of method-exclusive proteins. To better understand the general mechanism

of skeletal protein transportation, we further examined the proteins' gene ontology,

transmembrane domains, and signal peptides. We found that transmembrane domain

proteins and signal peptide secretion pathways, by themselves, could not explain the

transportation of proteins to the skeleton . We therefore propose that proteins are

transported to the skeletal via vesicles and possibly non-traditional secretion

pathways.

Keywords

Skeleton organic matrix proteins (SOMPs), biomineralization, Scleractinia

Introduction

Scleractinian corals, or stony corals, are the most prolific biomineralizers in

phylum Cnidaria [1]. They are a key components of shallow-water tropical reefs,

often forming massive structures that serve as the foundation of an ecosystem which

hosts some of the more biodiverse communities on the planet [2, 3]. They are amongst

the oldest biomineralizing metazoans, producing calcium carbonate ($CaCO_3$)

exoskeletons in the form of aragonite through biologically-directed mechanisms [4,

5], which makes up $\geq$95% of the entire skeletal mass, the remainder of which is the

skeletal organic matrix (SOM) [6-8].

Biomineralization refers to the ability of a living organism to selectively

exploit elements from its surrounding environment to build a biologically functioning

crystalline structure [9, 10]. Biomineralizers can be found throughout all kingdoms of

life, from bacteria [e.x. 11], to algae [e.x., 12], mollusks [e.x., 13], corals [e.x., 10]

and mammals [e.x., 14]. The minerals formed through this process differ in structure

from their non-biological counterparts and their formation is mediated by a variety of

organic molecules (the SOM), which have been intensively studied since the 1960's in

diverse organisms [reviewed by 10]. The SOM consists of proteins, lipids, and

polysaccharides, which are not present in the geological mineral form [reviewed by

15]. SOM proteins (SOMPs) embedded within the mineral are hypothesized to serve

as a framework for crystal nucleation [reviewed by 16]. The proteins involved in the

process of skeletal biomineralization have been described and studied extensively in

echinoderms [e.x., 17, 18] [and reviewed by 19, 20], mollusks [e.x., 21, 22-24] and

mammals [e.x., 25, 26, 27], among others [28]. At present, the best described SOMP

complex is of mammalian bone and teeth [15].

In stony corals, the most current knowledge of SOMPs is limited and based on

intraskeletal protein extraction [29-33] . It has been suggested that coral SOMPs aid in

the molecular processes of crystallization as well as in the development and

strengthening of the minerals ([34-37] among others). The constant advancement of

mass spectrometry technology has broadened our capability to identify many proteins

in skeletal extracts, even those proteins in low abundance [38]. However, this

technology is sensitive to contamination by organic matter remnants from soft tissue

and cell debris from the study organism, and the little-addressed issue remains of

contamination by researchers during the protein extraction, preparation, and

sequencing steps [39-41]. Upon extraction, the SOMPs are usually divided into two

fractions: acid-soluble matrix and acid-insoluble matrix proteins (ASM and AIM

respectively), based on their solubility in the acid of choice [22, 31-33, 42]. While

some past attention has been directed towards the soluble fraction [30, 43, 44],

Pereira-Mouriès et al. [45] showed that, in the bivalve *Pinctada maxima*, the

classification of AIM and ASM is misleading and that both fractions share common

features. Furthermore, Goffredo et al. [42] found in the stony coral *Balanophyllia*

europaea that both fractions consist of the same macromolecules; they associated the

degree of solubility to the difference in cross-linking. They also showed that each

solubility fraction has a different influence on calcium carbonate crystal morphology,

aggregation, and polymorphism in-vitro. In contrast, Ramos-Silva et al. [32] observed

a different SOMP composition between solubility fractions in the scleractinian coral

Acropora millepora,. Out of 36 SOMPs, only two were found exclusively in the

soluble fraction and twelve were exclusive to the insoluble fraction. These examples

demonstrate the attempts to attribute different properties to the two fractions but the

data remain inconclusive.

84   To date, three major coral skeletal proteomes have been published [31-33]

with each proteome consisting of 30-40 proteins. Of the 30 proteins sequenced from

A. digitifera skeleton, 26 were also detected in *A. millepora* skeleton [32, 33]. They

consist mostly of either transmembrane (TM) domain proteins or secretory proteins

[33]. However, only 12 of the proteins identified in *A. millepora* skeleton matched

those found in *S. pistillata* skeleton [31, 32]. In *A. millepora*, 11 TM domain-

containing proteins were identified, as well as two proteases that were not detected in

S. pistillata [32]. The authors suggested that the proteases' role is in cleaving the

extracellular domain of TM proteins and incorporating them into the skeleton.

93   The coral skeletal proteomes published to date reveal an overlap of several

detected proteins, but at least 10 proteins from each species appear to be unique. It is

currently unknown if this is truly due to species-specific gene expression and protein

localization or to methods in extracting, purifying, and sequencing the proteins. In

this study we analyzed several methods for protein purification to increase the

detection of proteins from cleaned coral skeleton powder. We show that the use of

acetone precipitation versus centrifugal filter washing, and the degree to which each

purification method is performed, affects the numbers and types of proteins that can

be sequenced by mass spectrometry. Further, we suggest that there is no one 'best'

method for coral skeletal protein preparation such that future research projects may

need to utilize several preparation methods to detect the full breadth of proteins

embedded in coral skeleton.

Methods

Sample collection and preparation for protein extraction

The hermatypic coral *Stylophora pistillata* (Esper, 1797) was collected under

a special permit from the Israeli Natural Parks Authority in the waters

in front of the H. Steinitz Marine Biology Laboratory, Eilat, Israel, Red Sea (29°30 N,

34°56 E), using SCUBA diving.

We fragmented one *S. pistillata* colony into small pieces, approximately 2x2

cm, with a diamond band saw. Coral fragments were transferred to 50-ml Falcon

brand conical vials (Falcon tubes) and oxidized with 20 mL 1:1 of 30% H_2O_2: 3%

NaClO solution for 1 hour, during which 1.5 mL of 3% NaClO solution were gently

added to the tubes every 20 minutes and continued the incubation overnight at room

temperature following modified methods of Stoll et al. [46]. Fragments were washed

with double distilled water five times for one minute each time and dried at 60 C°

overnight. We crushed the cleaned fragments to ≤63 µm diameter with a mortar and

pestle. Skeleton powder was then oxidized in sterile Falcon tubes and washed as

above three more times (i.e., four complete rounds of oxidative cleaning) to ensure

that no organic residue remained on the skeletal grains. In each cycle, the removal of

the oxidizing or washes solution was performed by centrifugation at 5,000 x g for 3

min at 4 C°. Cleaned skeletal powder was then dried overnight at 60 C°. We carried

out all the described processes in a laminar flow biological hood (apart from oven

drying) with all preparation tools and surfaces bleached to avoid contamination.

To monitor the removal of proteins from the skeletal fragments, we checked

the cleaning efficiency under SEM after the fourth oxidative cleaning. Samples were

vacuum-coated with 4 nm gold prior to examination under a ZEISS Sigma

TM scanning electron microscope an in-lens detector (5kV, WD = 5-7mm) (SI Figure

1a,b). In addition, we sonicated the powder at 4°C in filter-sterilized phosphate

buffered saline (PBS, pH 7.4) for 30 minutes, pelleted the powder at 5,000 x *g* for 3

minutes at 4˚C, concentrated the supernatant on a 3-kDa cutoff centrifugal filter unit

(Amicon) and loaded samples of supernatant on a 8-16% SDS-PAGE TGX stain free

gels (Bio-Rad) (SI Figure 1c).

Extraction of skeletal proteins

We decalcified approximately 1.3 g cleaned skeleton powder per protein extraction

protocol in 0.5 M acetic acid (30 ml acid/g cleaned skeleton powder) in Falcon tubes

while rotating the tubes at room temperature for 3 hours. Samples were then

centrifuged at 5,000 x g for 5 min at 4^{o}C and supernatant was transferred to a new

tube and stored at 4^{o}C. We continued the decalcification of the undissolved pellets

with a second volume of 0.5 M acetic acid and allowed decalcification to proceed to

completion. We then combined both liquid fractions (70 ml total) for each sample,

froze the total volumes at -80˚C, and dried them by overnight lyophilization. The

dried pellets were stored in -80°C until further processing.

The lyophilized pellets were re-suspended in 12 ml MilliQ water and the

proteins were concentrated on 3 kD cutoff Amicon® Ultra 15 centrifugal filter units

(Merk-Millipore) 5,000 x g at 4˚C to reach a final volume of 0.5 ml. To continue

desalting the samples we diluted them again to 12 ml in MilliQ water and repeated

the concentration procedure as above. At this stage, we observed an insoluble pellet in

all samples.

Because washes of the extracted pellet can reduce the representation of

proteins, in this study we examined two protein concentrating and cleaning methods

performed for previously published coral skeletal proteomes; (i) centrifugation

ultrafiltration (CF methods [32]) and (ii) acetone precipitation (ACT methods, [31])

(Table 1; SI Table 1). Further, for each method we separately examined the ASM and

AIM fractions.

Two filter-concentrated samples containing both soluble and insoluble

fractions were examined for the effects of centrifugal filtration. Sample CF2 was

centrifuged at 5,000 x g for 5 min at 4°C to pellet the AIM fraction. Both the ASM

and AIM of CF2 were divided into separate sterile 1.5 ml microcentrifuge tubes,

lyophilized, and stored at -80°C until further use. Sample CF4 was desalted as

described above for a third time before separating ASM and AIM fractions.

Both ACT samples were centrifuged as above for CF2 to separate ASM and

AIM fractions. To sample ACT1 ASM (concentrated to 0.5 ml by centrifugal

filtration as described above) was added 2 ml 100% ice cold acetone. The sample was

vortexed for 10 seconds, incubated at -20°C for 30 minutes, and centrifuged at 4,300

x g for 30 min at 4°C. Finally, the pellet was washed three more times with 2 mL of

80% ice cold acetone and stored then at -80°C until further use. The AIM fraction was

similarly washed four times with 80% acetone. Both fractions of sample ACT3 were

treated as in ACT1 but with one less washing step of each fraction.

LC MS\MS

S. pistillata skeletal protein samples were dissolved in 5% SDS and digested with

trypsin using the S-trap method overnight at room temperature. We analyzed the

resulting peptides with nanoflow ultra-performance liquid chromatograph

(nanoAcquity) coupled to a high resolution, high mass accuracy mass spectrometer

(Fusion Lumos). The sample was trapped on a Symmetry C18 0.18*20mm trap

column (Waters, Inc) and separated on a HSS T3 0.075*250 mm column (Waters,

Inc.) using a gradient of 4-28% (80% acetonitrile, 0.1% Formic acid) for 150 minutes.

Spray voltage was set to +2kV. The data were acquired in the Fusion Lumos using a

Top Speed Data-Dependent Acquisition method using a cycle time of 3 s. An MS1

scan was performed in the Orbitrap at 120,000 resolution with a maximum injection

time of 60 ms. The data were scanned between 300-1800 m/z. MS2 was selected

using a monoisotopic precursor selection set to peptides, peptide charge states set to

+2-+8 and dynamic exclusion set to 30 s. MS2 was performed using HCD

fragmentation scanned in the Orbitrap, with the first mass set to 130 m/z at a

resolution of 15,000. Maximum injection time was set to 60 ms with automatic gain

control of $5x10^{-4}$ ions as a fill target. The resulting data were searched against the

NCBI *Stylophora pistillata* protein database using the Byonic search engine (Protein

Metrics Inc.) – the first search was carried out without any false discovery rate (FDR)

filtering, to generate a focused database for a second search. The second search was

set to 1% FDR, allowing fixed carbamidomethylation on C and variable oxidation on

MW, deamidation on NQ and protein N-terminal acetylation. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the

Pride [47] partner repository under the dataset identifier PXD017891; reviewer login

username and password have been provided to the reviewers. Data will be made

publicly available without a required login in ProteomeXchange upon publication of

the manuscript.

Data sorting

We used the *S. pistillata* genome database as a reference peptide database for the

mass spectrometry analysis [48] (NCBI BioProjects PRJNA281535 and

PRJNA415215) appended with known NCBI *S. pistillata* skeletal proteins[e.x., 31,

49] *S. pistillata* carbonic anhydrase, ACE95141.1). We also included a common

contaminants database. Despite this inclusion of contaminants databases, several

proteins likely of human origin were sequenced and attributed to *S. pistillata*. To

filter out these potential contaminants from our final list of coral-specific proteins, we

BLASTed all sequences against the 'Primates' database in NCBI using Blast2GO.

We then examined NCBI-generated sequence alignments of coral versus *Homo*

sapiens proteins with e-values lower than e^{-50} and percent mean similarity greater than

50%, all sequences with e-values lower than e^{-100}, and all sequences with percent

similarity greater than 80%, and removed from our final list of coral proteins any

sequences with three or more peptides each of seven or more amino acids in length

that were identical between *S. pistillata* and humans.

All proteins identified by the LC MS\MS analysis were filtered to proteins

with at least two significant spectra and at least one significant unique peptide.

Skeletal proteins were first sorted by fractions and methods (ASM and AIM for each

extraction method). Next, we sorted all skeletal proteins by their gene ontology (GO)

terms [50, 51] to differentiate plasma membrane proteins and extracellular matrix

(ECM) proteins. Finally, we grouped all by terms of interest: protein modification,

transmembrane, and ECM; membrane processing; metal binding and vesicular.

Proteins detected by LC-MS/MS were annotated using the Trinotate pipeline

which relies on both pfam and Uniprot data [52]. GO terms and PFAM annotations

were assigned to *Stylophora pistillata* predicted proteins and transcripts using

Trinotate 3.0.1 (https://github.com/Trinotate).   Transmembrane regions were

predicted using the TMHMM server v2.0 [53]. Signal peptides on the N-termini of

proteins were predicted using SignalP 5.0 [54, 55]. Attachment of proteins to the

exterior of the cell membrane by glycosylphosphatidylinositol (GPI) anchors was

predicted using PredGPI [56]. These computational analyses used program default

settings and cutoffs.   Completeness of protein sequences was determined by

comparing all returned coral proteins to the *Acropora digitifera* genome [57]; NCBI

RefSeq assembly GCF_000222465.1). Detection of various proteins across solubility

and protocol fractions was visualized in Venny 2.1 online software.

We determined orthologous biomineralization genes, known from skeletal

proteomic analysis, across coral taxa. First, we estimated orthology relationships

between all non-redundant genes of selected metazoa species using OrthoFinder. We

included all genes of all Cnidaria species with known genome-based annotations.

OrthoFinder generates orthology-groups (Orthogroups) based on normalized

reciprocal best Blast hits' bit scores [58], and then estimates orthologues genes pairs

within Orthoroups [59]. We then selected all pairs of *Acropora* spp. orthologs to *S.*

pistillata (1:1, 1:many, many:many relationships) ([31-33], this study). From these

pairs we further selected *S. pistillata* spectra-based identified proteins, or skeletal

proteins known in the literature. Since not all skeletal protein annotations from the

literature were included in our reference Orthofinder proteome datasets, we further

found their best matches in the reference orthofinder proteome using blastp.

Results:

After extensive cleaning of the powdered skeleton and acid-extraction of embedded

organic matter we identified in total 60 coral-specific proteins, with two or greater

spectra or one unique peptide with at least 10 spectra and an identification score of

250 or greater, in *S. pistillata* skeleton as predicted by the species genome [48] (Table

2, SI Table 2). Trinotate [60] [52] returned GO annotations [50] for all of these

proteins, although many remain uncharacterized (SI Table 3).

In order to evaluate the efficacy and improve current methods for stony coral

skeletal protein extraction, we examined four different protocols; two centrifugation

ultrafiltration filters (CF) and two further acetone precipitation (ACT) protocols.

Proteomes of the four methodologies differed in composition and variety (Figure 1).

Combining results of all CF fractions identified 52 coral-specific proteins while

combined ACT protocols yielded 13 such proteins (Figure 1A). Moreover,

redundancy between methodologies was low. Only 8.3% of the proteins overlapped

between methods while 78.3% and 13.3% of the proteins were exclusive to combined

CF and combined ACT fraction data, respectively.

To evaluate the extracted protein treatment efficiency of each protocol, we

first compared the number of proteins detected in each method (e,x,, ACT1 vs ACT 3

and CF2 vs CF4). Of the 8 proteins found only in samples from the acetone wash

protocols (ACT 1 and ACT 3), one was observed only in ACT1 while five were

observed only ACT3 (Figure 1, SI Table 2); both ACT1 solubility fractions had one

more wash step than did those in ACT3. In contrast, of the 47 proteins observed only

in CF samples, 11 were found in CF2, which went through two filter centrifugation

steps, while 16 were found in CF4, which went through a third filter centrifugation

step. Further, only two proteins were observed in all four purification fractions in at

least one solubility form: CARP4/SAARP1 and synapsin 2-like. Two additional

proteins were observed in three of the four protocols: STPCA2 and an uncharacterized

proteins.

We next examined the difference in protein composition obtained from the

solubility fractions ASM versus AIM. We identified different distributions of proteins

in the ASM vs. AIM in both purification methods (Figures 1B & C). Notably, when

combining all methods, more proteins were identified in the AIM compared to the

ASM. A total of 57 coral proteins were identified in the AIM fraction compared to a

total of 8 in the ASM fraction (SI Table 2), of which 5 (8.3%) were identify both in

ASM and AIM.

Our data compared to other coral skeletal proteomes

Since our analysis yielded a large amount of new skeletal proteins, we also compared

our results with the three previously published proteome of *S. pistillata, A. digitifera*

and *A. millepora*. Out of our entire identified skeletal proteome containing 60

proteins, using OrthoFinder and BlastP, only 16 were found to be similar to proteins

identified in these studies. Yet, this proportion of overlap (16 out of 60) is

significantly greater than the expected proportion by chance, since the proportion of

known skeletal matrix proteins in the reference coral proteomes is extremely small

(less than ~0.2%). Seven proteins were found to overlap all four proteomes: a

coadhesin-like protein, an EGF and laminin G domain-containing protein, a

hypothetical protein, a MAM and LDL-receptor class A domain-containing protein, a

mucin, aspartic acid-rich protein 2-like, and a ZP domain-containing protein (Table

3).

Skeletal proteome characterization

We interrogated the mechanisms by which proteins may be exported from or attached

to the cell (SI Table 2). Seventeen coral skeletal proteins with likely complete N-

terminus predictions possess signal peptides as a potential mechanism for export from

the cell. Eight proteins contain at least one transmembrane span suggesting that they

are embedded in the cell membrane. Further, 10 proteins likely interact with the

exterior of the cell membrane by GPI anchors. In total, 25 of the 60 sequenced coral

skeletal proteins exhibit documented characteristics for localization in the ECM.

Because the majority of the proteins sequenced from the *S. pistillata* skeleton

do not possess features for signaling their export from the cell, we queried the data set

for further suggestions of positioning the proteins in the membrane or that the proteins

may be exported by vesicles such as those that may be involved in calcium

concentration. To do this, we examined the skeletal proteome annotations and GO

classifications toward finding common features to allow grouping of proteins. Out of

the entire skeletal proteome sequenced in this study (60 proteins), 39 genes were

returned with GO terms that allowed their classification into five groups of interest

based on their cellular component, biological process, and molecular function to

suggest likely cellular locations pertinent to the calcification mechanism, which may

therefore be indicative of their function in this process: lipid\phosphate\glycan related

proteins (i.e., membrane processing); ECM-related, transmembrane, and protein

modification proteins; and metal binding proteins vesicular/secretion related proteins,

(SI Table 3).

Of the proteins with GO terms, 10 proteins are suggested to be involved with

processing of the cell membrane (Figure 2), A much larger number are related to

vesicles/secretion as well as binding metal, with 21 and 25 assigned to each category,

respectively.   We combined these two categories in our proposed cellular location

Figure 2 as some of the skeletal proteins proposed to be found intracellularly in

vesicles are also known to bind calcium [34, 61]. Finally, 19 and 12 are potential

ECM proteins or are involved in protein modification, respectively. It should be

noted that many protein are assigned to multiple categories.

Discussion:

In this study we show the importance of using complementary post-extraction

methods to concentrate coral skeletal proteins for sequencing the full breadth of the

skeletal proteome. Our results show a clear and marked difference in detected proteins

between protein purification methods, with only two proteins observed in all methods.

Centrifugal Filter (CF) methods yielded a much greater abundance and diversity of

proteins than did acetone precipitation (ACT) (Figure 1). Moreover, we found that

protein detection is largely method-exclusive (Figure 1). We hypothesize that the

protein yield differences between extraction methods is based on the different

properties of each method. CF is a mechanical filtration based on size and has a bias

toward hydrophobic proteins, whereas ACT is based mainly on the chemical

interactions of proteins resulting in increased precipitation of hydrophilic proteins

[62]. We speculate that the CF methods yielded more total proteins in our study as

acetone precipitated proteins may be difficult to resolubilize, potentially leading to the

possible loss of many proteins in the pellet which is not transferred to the trypsin

digestion step.

338   A major challenge in working with intra-skeletal proteins is isolating the true

skeletal proteins from soft tissue contamination [40, 41]. In the present study, as in

previous work on *S. pistillata* skeletal proteins [40] we carried out an intensive

oxidative cleaning step on the skeletal powder, in addition to cleaning the skeletal

fragments, to avoid contamination. We did not observe any organic residues by SEM

on intact skeletons following the second cleaning step (SI Figure 1a,b) or in PBS

soaked on cleaned powders and concentrated (SI Figure 1c), and we are therefore

confident that all sequenced proteins are endogenous to the skeleton and are not coral

cellular contamination.

347   In this study, we examined the protein composition of the ASM and AIM

proteins separately. Although we see variations due to the different purification

techniques (Figure 1), our results indicate that, broadly, the AIM is distinguished in

composition from the ASM. Across all purification techniques, the AIM fractions

yielded more total proteins and fraction-exclusive proteins than the ASM fraction did

(Figure 1B, C). Since all extraction and precipitation experiments were carried out

simultaneously, we can rule out batch effect and technical differences. This AIM

versus ASM skeletome fractional composition is similar to previous studies carried

out on other marine biomineralizers such as mollusks [45] as well as to the stony

corals *A. millepora* [32] and *S. pistillata* [31]. However, the overlap between the two

solubility fractions reported for *A. millepora* [32] was 61%, in contrast to our study

which shows an overlap across all purification treatments combined of only 10%

(Figure 1). One plausible explanation for this difference is the use of improved LC-

MS/MS performance over the past several years enabled us to detect more proteins

and obtain a more comprehensive proteome. In previous studies of coral skeletal

proteomes which detailed differences between solubility fractions, the instruments

used were low resolution, low mass accuracy, which tend to result in a lower percent

identification of the data. Additionally, most search engines and FDR-calculating

algorithms struggle with very small datasets [63, 64]. Our use of Byonic helps to

alleviate statistical limitations that can result in false negative results.

Further differences in proteomes, beyond species differences, are the differing

reagents used in precipitation (compared to [31]) and our smaller centrifugal filter

cutoffs (compared to [32]). This second difference is particularly important with

respect to protein degradation. Even when embedded in biominerals so that amino

acids and even short peptides persist, proteins may still succumb to degradation [65].

This result in small peptides fragments that may be lost from centrifugal filter units

with 10 kDa pore sizes and larger. It is reasonable to assume that coral skeletal

proteins go through the same process, and if so, the cutoff of the membrane directly

affects the number of peptide spectra matches (PSM). Using smaller cutoff filters in

this study might have allowed us to capture some of these sheered peptides and led to

higher PSMs.

Traditional protein trafficking by signal peptides and transmembrane domain does

not explain the full extent of protein transportation to skeleton

Corals' skeleton is external to the animal; therefore, proteins in the skeletal matrices

must be transported outside the cells or span the membrane and have an extracellular

portion to reach the skeletal crystallization front. Indeed, recent studies of anthozoans

reveal a significant proportion of TM domain proteins (~35%) in the SOM [32, 33],

similar to that found in the better-studied Echinoidia [66] . Based on these findings,

we examined the hypothesis that many of the skeletal proteins originate in the plasma

membrane.

Our analysis revealed that TM domain proteins are not the major component

of the SOM protein complex; in the present proteome, TMHMM prediction suggests

that eight are embedded in the membrane. Hence we examined a cellular secretion

option. Ramos-Silva et al. [32] reported 15 proteins (41%) with SPs that did not also

possess a TM domain in the *A. millepora* proteome. In their study of the *A. digitifera*

proteome, Takeuchi et al. [33] reported a similar proportion of proteins with SPs but

no TM domains (40%). When examining the combined S. *pistillata* proteomes, 17 out

of 60 (28%) were positive for SPs (SI Table S2, [31]), likely a combination of

incomplete gene prediction and other mechanisms for exporting the proteins to the

calicoblastic space.

Hydrophobic regions of TM domains are similar to those of SPs, making SP

prediction difficult [53, 67]. We therefore cannot say for sure, based on these two

analyses (TMHMM and SignalP) alone, which is the more dominant pathway for

protein transport. While SP by itself can help predict traditional secretory pathways

[67], proteins that do not possess SPs may still leave the cell via other means without

classic SPs. This includes membrane pores [67], ATP-binding cassette transporters

[68] and autophagosome/endosome-based secretion [69, 70]. These are but a few

examples of secretory pathways which bypass the conventional TM/SP signaling

pathway (reviewed in [67, 70, 71]).

A further method by which protein may be exported to the site of calcification

is a vesicular pathway that differs from the conventional SP and TM pathway.

Previous studies have shown Ca^{2+} rich granules in the calicoblastic epithelium

(skeletogenic cells), but not in the other tissue layers, suggesting their role as a Ca^{2+}

reservoirs in the cells. Vesicles were previously identified in corals [72, 73]; however,

their origin and content was not detailed and they are sometimes attributed to

preservation byproducts. These intracellular ion-rich vesicles may endocytose sea

water [74], after which they are enriched in carbonate ions and then form hydrated

ACC and anhydrous ACC precursors stabilized by acidic biomolecules including

CARPs [75]. Using cell cultures, Mass et al. [61] suggest that the vesicles, which

contain Asp rich proteins, then transport their contents to the ECM, releasing their

content by exocytosis. The biomineral then further develops extracellularly, likely

aided by other ECM proteins [30, 35, 49, 76] as well as other biomolecules [77-80].

At present the process of calcium delivery to the skeleton and the role of most of the

proteins in skeletal depositing remain to be determined.

Since TM and SP analysis did not fully explain the mechanism of protein

transportation, we classified the likely functions and locations of the skeletal proteins

according to their GO terms as lipid\phosphate\glycan related proteins, metal binding

proteins, vesicular/secretion related proteins, ECM-related and transmembrane

proteins, and protein modification proteins (Figure 2, SI Table 3). Looking at the

proposed locations or functions of skeletal proteins (Figure 2), we find comparable

numbers of vesicular and metal binding proteins versus TM domain/ECM proteins.

These results are in-line with the trend reported in previous coral skeletal proteomes

[31-33]. While we do not exclude the role of TM proteins in the skeletal deposition

process, we suggest that the biological mechanism of SOM transportation is enhanced

by vesicular secretion and other possible, non-traditional, secretion pathways.

It is difficult to map and characterize proteins in non-model organisms such *S.*

pistillata [81-83]. High quality proteomic mapping requires knowledge of

phosphorylation, glycosylation, proteolytic cite activities and other modifications [31,

84-86], in order to create a more thorough database [81-83]. Further, many of the

coral skeletal proteins reported to date remain uncharacterized. Uncharacterized

proteins were reported in *Acropora* skeletons at a rate of approximately 25% [32, 33]

while, they were previously reported at less than 10% in *Stylophora* skeleton [31].

Our study revealed a much greater proportion of 28% uncharacterized proteins. While

partially attributed to sample size, it is most likely due to quality of genomic data

available, since stony corals are non-model organisms and their genomic libraries are

far from being complete, resulting in incomplete databases on which to map the

proteome and many uncharacterized genes.

Conclusion:

In this study we have considered the differential effects of coral skeletal protein

extract preparation as well as the method by which these proteins, or parts thereof, are

transported from intracellular to extracellular locations. When preparing coral skeletal

proteomes, we propose that a multi-method approach to cleaning, demineralization,

and protein extraction should be used. Our results showed that each protein

preparation protocol yielded exclusive sets of proteins with little overlap between

ACT and CF fractions. While CF protocols yielded many more proteins than did ACT

methods, use of a single protocol to clean and concentrate coral skeletal proteins

results in a significant amount of data loss, and it is therefore of crucial importance to

consider alternative and complementary methods to obtain a fully comprehensive

skeletal proteome. We showed that while the role of TM domain proteins cannot be

overlooked, many of the proteins detected in the *S. pistillata* skeletal proteomes as

well as in that of other species point toward other secretory or vesicular pathways.

Our categorization method, supported by data from other recent studies, also suggests

that corals use an alternative secretory pathway, such as vesicles, and much work is

required in order to determine the calcium deposition pathway and the proteins

involved. Our study provides a large set of new uncharacterized coral skeletal proteins

as well as others of purported function but that have not been observed before in the

coral SOM. These data expand the current knowledge of the SOM in corals and will

help, in future studies, to resolve corals' calcium deposition mechanism and the

various roles of the proteins involved.

Abbreviations

acetone precipitation of proteins (ACT)

acid soluble matrix (ASM)

acid insoluble matrix (AIM)

centrifugation ultrafiltration of proteins (CF)

false discovery rate (FDR)

gene ontology (GO)

glycosylphosphatidylinositol (GPI)

peptide spectra matches (PSM)

phosphate buffered saline (PBS)

skeletal organic matrix (SOM)

skeletal organic matrix proteins (SOMP)

transmembrane (TM)

Declarations

Ethics approval and consent to participate

Coral nubbins were collected under a special permit from the Israeli Natural Parks

Authority.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated during the current study are available in the ProteomeXchange

repository under file number PXD017891 (https://www.ebi.ac.uk/pride/archive). The

reviewer login username and password have been provided to the reviewers. Data

will be made publicly available without a required login in ProteomeXchange upon

publication of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Israel Science Foundation (Grant 312/15), and the

European Research Commission (ERC; Grant # 755876). JLD was supported by the

Zuckerman STEM Leadership Program.

Author contributions

YP, JD, and TM designed the study. YP and RA prepared the samples. YP, RA, AM,

ML, DM, and JD analyzed the data. All authors wrote the manuscript and approve

this submission.

Acknowledgements

We thank staff at the Bioinformatics Core Unit, University of Haifa, and The Crown

Genomics Institute of the Nancy and Stephen Grand Israel National Center for

Personalized Medicine, Weizmann Institute of Science.

References

1. Cairns SD: **Deep-water corals: An overview with special reference to**
diversity and distribution of deep-water scleractinian corals. *Bull Mar Sci*
2007, **81:**311-322.
2. Reaka-Kudla ML: **The global biodiversity of coral reefs: a comparison**
with rain forests. *Biodiversity II: Understanding and Protecting our*
Biological Resources 1997, **2:**551.
3. Knowlton N, Brainard RE, Fisher R, Moews M, Plaisance L, Caley MJ: **Coral**
reef biodiversity. *Life in the world's oceans: Diversity distribution and*
abundance 2010**:**65-74.
4. Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D, Peterson KJ:
The Cambrian conundrum: Early divergence and later ecological success
in the early history of animals. *Science* 2011, **334:**1091-1097.
5. Tambutté S, Tambutté E, Zoccola D, Allemand D: **Organic matrix and**
biomineralization of scleractinian corals. *Handbook of Biomineralization:*
Biological Aspects and Structure Formation 2007**:**243-259.
6. Cuif J-P, Dauphin Y, Berthet P, Jegoudez J: **Associated water and organic**
compounds in coral skeletons: Quantitative thermogravimetry coupled to
infrared absorption spectrometry. *Geochemistry Geophysics Geosystems*
2004, **5:**Q11011.
7. Wainwright SA: **Skeletal organization in the coral, *Pocillopora damicornis*.**
Quarterly Journal of Microscopical Science 1963, **s3-104:**169-183.
8. Von Euw S, Zhang Q, Manichev V, Murali N, Gross J, Feldman LC,
Gustafsson T, Flach C, Mendelsohn R, Falkowski PG: **Biological control of**
aragonite formation in stony corals. *Science* 2017, **356:**933-938.
9. Mann S: *Biomineralization: Principles and Concepts in Bioinorganic*
Materials Chemistry. New York: Oxford University Press; 2001.
10. Weiner S, Dove PM: **An overview of biomineralization processes and the**
problem of the vital effect. *Reviews in Mineralogy and Geochemistry* 2003,
54:1-29.
11. Schüler D, Frankel RB: **Bacterial magnetosomes: microbiology,**
Biomineralization and biotechnological applications. *Appl Microbiol*
Biotechnol 1999, **52:**464-473.
12. Westbroek P, Young J, Linschooten K: **Coccolith production**
(biomineralization) in the marine alga *Emiliania huxleyi*. *The Journal of*
Protozoology 1989, **36:**368-373.
13. Addadi L, Joester D, Nudelman F, Weiner S: **Mollusk shell formation: a**
source of new concepts for understanding biomineralization process.
Chemical European Journal 2006, **12:**980-987.
14. Levine P, Glimcher M: **The isolation and amino acid composition of the**
organic matrix and neutral soluble proteins of developing rodent enamel.
Archives of Oral Biology 1965, **10:**753-IN751.
15. Beniash E: **Biominerals—hierarchical nanocomposites: the example of**
bone. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*
2011, **3:**47-69.
16. Lowenstam HA, Weiner S: *On Biomineralization.* New York, New York:
Oxford University Press; 1989.
17. Karakostis K, Zanella-Cléon I, Immel F, Guichard N, Dru P, Lepage T,
Plasseraud L, Matranga V, Marin F: **A minimal molecular toolkit for**

mineral deposition? Biochemistry and proteomics of the test matrix of adult specimens of the sea urchin *Paracentrotus lividus*. *Journal of Proteomics* 2016, **136:**133-144.

18. Shashikant T, Khor JM, Ettensohn CA: **From genome to anatomy: The architecture and evolution of the skeletogenic gene regulatory network of sea urchins and other echinoderms.** *Genesis* 2018, **56:**e23253.

19. Nebelsick JH, Dynowski JF, Grossmann JN, Tötzke C: **Echinoderms: hierarchically organized light weight skeletons.** In *Evolution of Lightweight Structures*. Springer; 2015: 141-155

20. Killian CE, Wilt FH: **Molecular aspects of biomineralization of the echinoderm endoskeleton.** *Chemical Reviews* 2008, **108:**4463-4474.

21. Weiner S, Hood L: **Soluble protein of the organic matrix of mollusk shells: a potential template for shell formation.** *Science* 1975, **190:**987-989.

22. Mann K, Edsinger-Gonzales E, Mann M: **In-depth proteomic analysis of a mollusc shell: acid-soluble and acid-insoluble matrix of the limpet *Lottia gigantea*.** *Proteome Science* 2012, **10:**1-18.

23. Politi Y, Mahamid J, Goldberg H, Weiner S, Addadi L: **Asprich mollusk shell protein: in vitro experiments aimed at elucidating function in CaCO$_3$ crystallization.** *CrystEngComm* 2007, **9:**1171-1177.

24. Bouyoucef M, Rakic R, Gómez-Leduc T, Latire T, Marin F, Leclercq S, Carreiras F, Serpentini A, Lebel J-M, Galéra P: **Regulation of extracellular matrix synthesis by shell extracts from the marine bivalve *Pecten maximus* in human articular chondrocytes—application for cartilage engineering.** *Mar Biotechnol* 2018**:**1-15.

25. Weiner S, Wagner HD: **The material bone: Structure-mechanical function relations.** *Annual Review of Materials Science* 1998, **28:**271-298.

26. Reid DG, Duer MJ, Murray RC, Wise ER: **The organic–mineral interface in teeth is like that in bone and dominated by polysaccharides: Universal mediators of normal calcium phosphate niomineralization in vertebrates?** *Chemistry of Materials* 2008, **20:**3549-3550.

27. Sharma V, Srinivasan A, Roychoudhury A, Rani K, Tyagi M, Dev K, Nikolajeff F, Kumar S: **Characterization of protein extracts from different types of human teeth and insight in biomineralization.** *Scientific Reports* 2019, **9:**9314.

28. Kamel BS, Voolstra CR, Medina M: **BioMine-DB: A database for metazoan biomineralization proteins.** *PeerJ Preprints* 2016, **4:**e1983v1982.

29. Fukuda I, Ooki S, Fujita T, Murayama E, Nagasawa H, Isa Y, Watanabe T: **Molecular cloning of a cDNA encoding a soluble protein in the coral exoskeleton.** *Biochem Biophys Res Commun* 2003, **304:**11-17.

30. Puverel S, Tambutté E, Pereira-Mouries L, Zoccola D, Allemand D, Tambutté S: **Soluble organic matrix of two Scleractinian corals: Partial and comparative analysis.** *Comparative Biochemistry and Physiology Part B* 2005, **141:**480-487.

31. Drake JL, Mass T, Haramaty L, Zelzion E, Bhattacharya D, Falkowski PG: **Proteomic analysis of skeletal organic matrix from the stony coral *Stylophora pistillata*.** *Proceedings of the National Academy of Sciences* 2013, **110:**3788-3793.

32. Ramos-Silva P, Kaandorp J, Huisman L, Marie B, Zanella-Cleon I, Guichard N, Miller DJ, Marin F: **The skeletal proteome of the coral *Acropora***

millepora: **The evolution of calcification by cooption and domain shuffling.** *Mol Biol Evol* 2013.

33. Takeuchi T, Yamada L, Shinzato C, Sawada H, Satoh N: **Stepwise evolution of coral biomineralization revealed with genome-wide proteomics and transcriptomics.** *PLOS ONE* 2016, **11:**e0156424.

34. Mass T, Drake Jeana L, Haramaty L, Kim JD, Zelzion E, Bhattacharya D, Falkowski Paul G: **Cloning and characterization of four novel coral acid-rich proteins that precipitate carbonates in vitro.** *Curr Biol* 2013, **23:**1126-1131.

35. Puverel S, Tambutté E, Zoccola D, Dumart-Coulon I, Bouchot A, Lotto S, Allemand D, Tambutté S: **Antibodies against the organic matrix in scleractinians: A new tool to study coral biomineralization.** *Coral Reefs* 2005, **24:**149-156.

36. Clode PL, Marshall AT: **Calcium associated with a fibrillar organic matrix in the scleractinian coral *Galaxea fascicularis*.** *Protoplasma* 2003, **220:**153-161.

37. Watanabe T: **Molecular analyses of protein components of the organic matrix in the exoskeleton of two scleractinian coral species.** *Comp Biochem Phys B* 2003, **136:**767 - 774.

38. Scheltema RA, Hauschild J-P, Lange O, Hornburg D, Denisov E, Damoc E, Kuehn A, Makarov A, Mann M: **The Q Exactive HF, a Benchtop mass spectrometer with a pre-filter, high-performance quadrupole and an ultra-high-field Orbitrap analyzer.** *Molecular & Cellular Proteomics* 2014, **13:**3698-3708.

39. Drake JL, Mass T, Falkowski PG: **The evolution and future of carbonate precipitation in marine invertebrates: Witnessing extinction or documenting resilience in the Anthropocene?** *Elementa Science of the Anthropocene* 2014, **2:**000026.

40. Drake JL, Mass T, Haramaty L, Zelzion E, Bhattacharya D, Falkowski PG: **Reply to Ramos-Silva et al.: Regarding coral skeletal proteome.** *Proceedings of the National Academy of Sciences* 2013, **110:**E2147-E2148.

41. Ramos-Silva P, Marin F, Kaandorp J, Marie B: **LETTER: Biomineralization toolkit: the importance of sample cleaning prior to the characterization of biomineral proteomes.** Proceedings of the National Academy of Sciences; 2013.

42. Goffredo S, Vergni P, Reggi M, Caroselli E, Sparla F, Levy O, Dubinsky Z, Falini G: **The skeletal organic matrix from Mediterranean coral *Balanophyllia europaea* influences calcium carbonate precipitation.** *PLoS ONE* 2011, **6**.

43. Dauphin Y: **Comparative studies of skeletal soluble matrices from some scleractinian corals and molluscs.** *International Journal of Biological Macromolecules* 2001, **28:**293-304.

44. Weiner S: **Aspartic acid-rich proteins: major components of the soluble organic matrix of mollusk shells.** *Calcif Tissue Int* 1979, **29:**163-167.

45. Pereira-Mouriès L, Almeida MJ, Ribeiro C, Peduzzi J, Barthélemy M, Milet C, Lopez E: **Soluble silk-like organic matrix in the nacreous layer of the bivalve *Pinctada maxima*: A new insight in the biomineralization field.** *Eur J Biochem* 2002, **269:**4994-5003.

46. Stoll HM, Ruiz Encinar J, Ignacio Garcia Alonso J, Rosenthal Y, Probert I, Klaas C: **A first look at paleotemperature prospects from Mg in coccolith**

carbonate: Cleaning techniques and culture measurements. *Geochemistry,*
Geophysics, Geosystems 2001, **2**.

47. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S,
Kundu DJ, Inuganti A, Griss J, Mayer G, Eisenacher M: **The PRIDE**
database and related tools and resources in 2019: improving support for
quantification data. *Nucleic Acids Res* 2019, **47:**D442-D450.

48. Voolstra CR, Li Y, Liew YJ, Baumgarten S, Zoccola D, Flot J-F, Tambutté S,
Allemand D, Aranda M: **Comparative analysis of the genomes of**
Stylophora pistillata **and** *Acropora digitifera* **provides evidence for**
extensive differences between species of corals. *Scientific Reports* 2017,
7:17583.

49. Bertucci A, Tambutté S, Supuran C, Allemand D, Zoccola D: **A new coral**
carbonic anhydrase in *Stylophora pistillata*. *Mar Biotechnol* 2011, **13:**992-
1002.

50. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP,
Dolinski K, Dwight SS, Eppig JT: **Gene Ontology: Tool for the unification**
of biology. *Nat Genet* 2000, **25:**25-29.

51. Consortium GO: **Expansion of the gene ontology knowledgebase and**
resources. *Nucleic Acids Res* 2017, **45:**D331-D338.

52. Bryant DM, Johnson K, DiTommaso T, Tickle T, Couger MB, Payzin-Dogru
D, Lee TJ, Leigh ND, Kuo T-H, Davis FG: **A tissue-mapped axolotl de novo**
transcriptome enables identification of limb regeneration factors. *Cell*
Reports 2017, **18:**762-776.

53. Krogh A, Larsson B, Heijne Gv, Sonnhammer EL: **Predicting**
transmembrane protein topology with a hidden Markov model:
Application to complete genomes. *J Mol Biol* 2001, **305:**567-580.

54. Armenteros JJA, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak
S, von Heijne G, Nielsen H: **SignalP 5.0 improves signal peptide**
predictions using deep neural networks. *Nat Biotechnol* 2019, **37:**420.

55. Nielsen H, Engelbrecht J, Brunak S, von Heijne G: **Identification of**
prokaryotic and eukaryotic signal peptides and prediction of their
cleavage sites. *Protein Engineering* 1997, **10:**1-6.

56. Poisson G, Chauve C, Chen X, Bergeron A: **FragAnchor: a large-scale**
predictor of glycosylphosphatidylinositol anchors in eukaryote protein
sequences by qualitative scoring. *Genomics, Proteomics & Bioinformatics*
2007, **5:**121-130.

57. Shinzato C, Shoguchi E, Kawashima T, Hamada M, Hisata K, Tanaka M,
Fujie M, Fujiwara M, Koyanagi R, Ikuta T, et al: **Using the** *Acropora*
digitifera **genome to understand coral responses to environmental change.**
Nature 2011, **476:**320-323.

58. Emms DM, Kelly S: **OrthoFinder: Solving fundamental biases in whole**
genome comparisons dramatically improves orthogroup inference
accuracy. *Genome Biology* 2015, **16:**157.

59. Emms DM, Kelly S: **OrthoFinder: Phylogenetic orthology inference for**
comparative genomics. *Genome Biology* 2019, **20:**1-14.

60. Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ,
Robles M, Talón M, Dopazo J, Conesa A: **High-throughput functional**
annotation and data mining with the Blast2GO suite. *Nucleic Acids Res*
2008, **36:**3420-3435.

61. Mass T, Drake JL, Heddleston JM, Falkowski PG: **Nanoscale visualization of**
biomineral formation in coral proto-polyps. *Curr Biol* 2017, **27:**3191-3196.
e3193.
62. Thongboonkerd V, Mcleish KR, Arthur JM, Klein JB: **Proteomic analysis of**
normal human urinary proteins isolated by acetone precipitation or
ultracentrifugation. *Kidney International* 2002, **62:**1461-1469.
63. Bern MW, Kil YJ: **Two-dimensional target decoy strategy for shotgun**
proteomics. *Journal of Proteome Research* 2011, **10:**5296-5301.
64. Fondrie WE, Noble WS: **A machine learning strategy that leverages large**
datasets to boost statistical power in small-scale experiments. *BioRxiv*
2019**:**849331.
65. Sykes GA, Collins MJ, Walton DI: **The significance of a geochemically**
isolated intracrystalline organic fraction within biominerals. *Org Geochem*
1995, **23:**1059-1065.
66. Mann K, Wilt FH, Poustka AJ: **Proteomic analysis of sea urchin**
(*Strongylocentrotus purpuratus*) **spicule matrix.** *Proteome Science* 2010,
8:33.
67. Von Heijne G, Abrahmsèn L: **Species-specific variation in signal peptide**
design Implications for protein secretion in foreign hosts. *FEBS Lett* 1989,
244:439-446.
68. McGrath JP, Varshavsky A: **The yeast STE6 gene encodes a homologue of**
the mammalian multidrug resistance P-glycoprotein. *Nature* 1989,
340:400-404.
69. Nickel W, Rabouille C: **Mechanisms of regulated unconventional protein**
secretion. *Nature Reviews Molecular Cell Biology* 2009, **10:**148-155.
70. Rabouille C: **Pathways of unconventional protein secretion.** *Trends Cell*
Biol 2017, **27:**230-240.
71. Rabouille C, Malhotra V, Nickel W: **Diversity in unconventional protein**
secretion. *J Cell Sci* 2012, **125:**5251-5255.
72. Clode PL, Marshall AT: **Kalisomes in corals: a novel KCl concentrating**
organelle? *Tissue and Cell* 2002, **34:**199-209.
73. Marshall AT, Clode PL, Russell R, Prince K, Stern R: **Electron and ion**
microprobe analysis of calcium distribution and transport in coral tissues.
J Exp Biol 2007, **210:**2453-2463.
74. Mass T, Giuffre AJ, Sun C-Y, Stifler CA, Frazier MJ, Neder M, Tamura N,
Stan CV, Marcus MA, Gilbert PUPA: **Amorphous calcium carbonate**
particles form coral skeletons. *Proceedings of the National Academy of*
Sciences 2017, **114:**E7670-E7678.
75. Neder M, Laissue PP, Akiva A, Akkaynak D, Albéric M, Spaeker O, Politi Y,
Pinkas I, Mass T: **Mineral formation in the primary polyps of**
pocilloporoid corals. *Acta Biomater* 2019.
76. Mass T, Drake JL, Peters EC, Jiang W, Falkowski PG: **Immunolocalization**
of skeletal matrix proteins in tissue and mineral of the coral *Stylophora*
pistillata. *Proceedings of the National Academy of Sciences* 2014, **111:**12728-
12733.
77. Takeuchi T, Plasseraud L, Ziegler-Devin I, Brosse N, Shinzato C, Satoh N,
Marin F: **Biochemical characterization of the skeletal matrix of the**
massive coral, *Porites australiensis*–**The saccharide moieties and their**
localization. *Journal of Structural Biology* 2018, **203:**219-229.

78. Puverel S, Houlbreque F, Tambutté E, Zoccola D, Payan P, Caminiti N,
Tambutté S, Allemand D: **Evidence of low molecular weight components in**
the organic matrix of the reef building coral, *Stylophora pistillata*.
Comparative Biochemistry and Physiology a-Molecular & Integrative
Physiology 2007, **147:**850-856.
79. Farre B, Cuif J-P, Dauphin Y: **Occurrence and diversity of lipids in modern**
coral skeletons. *Zoology* 2010, **113:**250-257.
80. Naggi A, Torri G, Iacomini M, Colombo Castelli G, Reggi M, Fermani S,
Dubinsky Z, Goffredo S, Falini G: **Structure and function of stony coral**
intraskeletal polysaccharides. *ACS Omega* 2018, **3:**2895-2901.
81. Carpentier SC, Panis B, Vertommen A, Swennen R, Sergeant K, Renaut J,
Laukens K, Witters E, Samyn B, Devreese B: **Proteome analysis of**
non-model plants: a challenging but powerful approach. *Mass*
Spectrometry Reviews 2008, **27:**354-377.
82. Dowd WW: **Challenges for biological interpretation of environmental**
proteomics data in non-model organisms. *Integr Comp Biol* 2012, **52:**705-
720.
83. Forné I, Abian J, Cerdà J: **Fish proteome analysis: model organisms and**
non-sequenced species. *Proteomics* 2010, **10:**858-872.
84. Butler WT, Brunn JC, Qin C: **Dentin extracellular matrix (ECM) proteins:**
comparison to bone ECM and contribution to dynamics of
dentinogenesis. *Connect Tissue Res* 2003, **44:**171-178.
85. Qin C, Baba O, Butler WT: **Post-translational modifications of SIBLING**
proteins and their roles in osteogenesis and dentinogenesis. *Critical*
Reviews in Oral Biology and Medicine 2004, **15:**126-136.
86. Sarashina I, Endo K: **Skeletal matrix proteins of invertebrate animals:**
comparative analysis of their amino acid sequences. *Paleontological*
Research 2006, **10:**311-336.

Table and Figure Legends

Tables
Table 1: Summary of methods for protein concentration and cleaning after extraction
in acetic acid. CF – Centrifugation filtration methods; ACT – acetone precipitation
methods; ASM – Acid soluble matrix; AIM – Acid insoluble matrix.

Table 2: 60 coral skeletal proteins detected by LC-MS/MS across all treatments and
solubility fractions. Proteins are listed in order of accession number. Geno ontology
categorization is represented as [a]ECM/transmembrane and protein modification,
[b]membrane processing, and [c]vesicle/secretion and metal binding.

Table 3. Orthologous coral skeletal proteins in the present work and previously
published. Gene ontology categorization is represented as [a]ECM/transmembrane and

797 protein modification, [b]membrane processing, and [c]vesicle/secretion and metal
798 binding.
799
Figures
Figure 1: Distribution of proteins numbers according of the extraction methods.
Distribution of all proteins (SOM and ISOM combined) by extraction methods (A).
SOM and ISOM distribution by acetone precipitation methods (ACT). SOM and
ISOM distribution by centrifugation and ultrafiltration method (CF). Note that CF
methods yielded more proteins in total. ISOM Insoluble Skeletal Organic Matrix;
SOM Skeletal Organic Matrix.
807
808 Figure 2. Suggested cellular and extracellular locations of *S. pistillata* skeletal protein
809 based on GO terms. Large yellow and red circles represent calicoblastic cells with
810 red cell membrane facing the skeleton. Small green circles represent vesicles. And
811 purple squiggles represent transmembrane proteins that may or may not span the
812 width of the calicoblastic space plus ECM proteins.
813

Sample Name	Filter desalting (CF)				Acetone Precipitation (ACT)			
	CF2		CF4		ACT1		ACT3	
Solubility Fraction	ASM	AIM	ASM	AIM	ASM	AIM	ASM	AIM
Filter desalting x 2	+	+	+	+	+	+	+	+
Filter desalting x 3	-	-	+	+	-	-	-	-
Acetone precipitation	-	-	-	-	+	-	+	-
1x 80% acetone wash	-	-	-	-	+	+	+	+
2x 80% acetone wash	-	-	-	-	+	+	+	+
3x 80% acetone wash	-	-	-	-	+	+	-	+
4x 80% acetone wash	-	-	-	-	-	+	-	-

| | | | | Go | Solubility Fraction | |
| | Accession | | | | | |
gi Number	Number	Annotation	Best score	Categorization	ASM	AIM
		Stylophora pistillata genome hit				
1263115389	PFX12726.1	Retrovirus-related Pol polyprotein from transposon 17.6 [Stylophora pistillata]	340.80	-		x
1263115517	PFX12813.1	hypothetical protein AWC38_SpisGene23165 [Stylophora pistillata]	337.40	-		x
1263116757	PFX13778.1	Sacsin [Stylophora pistillata]	267.80	-		x
1263117261	PFX14205.1	Proto-oncogene tyrosine-protein kinase receptor Ret, partial [Stylophora pistillata]	336.70	a,c		x
1263119001	PFX15740.1	Protein FAM208A [Stylophora pistillata]	373.20	-	x	x
1263119725	PFX16398.1	hypothetical protein AWC38_SpisGene19330 [Stylophora pistillata]	273.10	-		x
1263122270	PFX18785.1	Mucin-4 [Stylophora pistillata]	449.50	a,c		x
1263130352	PFX26597.1	Complement C3 [Stylophora pistillata]	283.10	a,b,c		x
1263130510	PFX26751.1	Transmembrane protease serine 9 [Stylophora pistillata]	244.20	a		x
1263131615	PFX27832.1	Poly [ADP-ribose] polymerase 11 [Stylophora pistillata]	320.10	c		x
1263134664	PFX30831.1	hypothetical protein AWC38_SpisGene4366 [Stylophora pistillata]	322.20	-		x
1263134737	PFX30903.1	hypothetical protein AWC38_SpisGene4292 [Stylophora pistillata]	548.90	-		x
1263135656	PFX31810.1	Nidogen-2 [Stylophora pistillata]	286.20	a,c		x
1270028962	XP_022783323.1	phosphopantothenoylcysteine decarboxylase subunit VHS3-like [Stylophora pistillata]	406.60	c		x
1270029544	XP_022786582.1	synapsin-2-like isoform X2 [Stylophora pistillata]	320.50	c	x	x
1270031975	XP_022799089.1	CUB domain-containing protein-like isoform X2 [Stylophora pistillata]	699.70	-		x
1270032917	XP_022804012.1	EGF and laminin G domain-containing protein-like [Stylophora pistillata]	384.10	a		x
1270036141	XP_022779720.1	vitellogenin-like [Stylophora pistillata]	416.00	b,c		x
1270037242	XP_022780303.1	LOW QUALITY PROTEIN: uncharacterized protein LOC111321626 [Stylophora pistillata]	353.60	a,b,c	x	
1270037961	XP_022780690.1	skeletal aspartic acid-rich protein 2-like [Stylophora pistillata]	723.60	a		x
1270037969	XP_022780694.1	CUB and peptidase domain-containing protein 2-like [Stylophora pistillata]	285.80	a	x	x
1270041141	XP_022782398.1	skeletal aspartic acid-rich protein 1-like [Stylophora pistillata]	709.80	-	x	x
1270042352	XP_022783044.1	uncharacterized protein LOC111323869 [Stylophora pistillata]	283.60	-		x
1270043038	XP_022783415.1	coadhesin-like isoform X3 [Stylophora pistillata]	411.20	a		x
1270044042	XP_022783952.1	collagenase 3-like [Stylophora pistillata]	262.90	a,c		x
1270045301	XP_022784623.1	cation channel sperm-associated protein subunit beta-like [Stylophora pistillata]	328.40	a,c		x
1270049598	XP_022786918.1	major yolk protein-like isoform X2 [Stylophora pistillata]	476.50	a,c		x
1270052030	XP_022788227.1	hephaestin-like protein [Stylophora pistillata]	539.80	a,c		x
1270052971	XP_022788730.1	chymotrypsin-like elastase family member 1 [Stylophora pistillata]	264.80	a,c		x
1270054595	XP_022789591.1	endothelin-converting enzyme 1-like isoform X2 [Stylophora pistillata]	94.10	a,b,c		x
1270055217	XP_022789932.1	MAGUK p55 subfamily member 7-like [Stylophora pistillata]	313.70	a		x
1270056169	XP_022790441.1	PHD finger protein 21A-like [Stylophora pistillata]	277.90	c		x
1270059479	XP_022792212.1	ras-like protein 3 [Stylophora pistillata]	289.90	a,c		x
1270063049	XP_022794122.1	galaxin-like isoform X2 [Stylophora pistillata]	309.40	-		x
1270063475	XP_022794351.1	mammalian ependymin-related protein 1-like [Stylophora pistillata]	324.80	a,c		x
1270064196	XP_022794736.1	MAM and LDL-receptor class A domain-containing protein 2-like [Stylophora pistillata]	472.20	-		x
1270068394	XP_022796981.1	uncharacterized skeletal organic matrix protein 8-like [Stylophora pistillata]	664.40	-		x
1270068396	XP_022796982.1	uncharacterized protein LOC111335364 [Stylophora pistillata]	609.30	-		x
1270071953	XP_022798902.1	low-density lipoprotein receptor-related protein 8-like [Stylophora pistillata]	257.10	c		x
1270073139	XP_022799541.1	uncharacterized protein LOC111337489 [Stylophora pistillata]	461.10	b		x
1270080553	XP_022803524.1	digestive cysteine proteinase 1-like [Stylophora pistillata]	262.20	a,c		x
1270081088	XP_022803808.1	deleted in malignant brain tumors 1 protein-like [Stylophora pistillata]	344.70	c		x
1270081207	XP_022803872.1	spore wall protein 2-like isoform X3 [Stylophora pistillata]	338.60	-		x
1270081241	XP_022803894.1	uncharacterized protein LOC111341206 [Stylophora pistillata]	390.70	b		x
1270082891	XP_022804785.1	thioredoxin reductase 1, cytoplasmic-like [Stylophora pistillata]	337.00	c		x
1270084203	XP_022805470.1	uncharacterized protein LOC111342641 [Stylophora pistillata]	256.10	-		x
1270085816	XP_022806326.1	ZP domain-containing protein-like [Stylophora pistillata]	393.90	a		x
1270086467	XP_022806664.1	protein lingerer-like [Stylophora pistillata]	250.10	b,c		x
1270086954	XP_022806928.1	SLIT-ROBO Rho GTPase-activating protein 1-like [Stylophora pistillata]	309.10	-		x
1270087345	XP_022807143.1	condensin-2 complex subunit D3-like [Stylophora pistillata]	278.10	c		x
1270087556	XP_022807256.1	uncharacterized protein LOC111344300 [Stylophora pistillata]	280.40	a,c		x
1270088573	XP_022807807.1	uncharacterized protein LOC111344812 [Stylophora pistillata]	365.30	a,b,c		x
1270089244	XP_022808163.1	uncharacterized protein LOC111345150 [Stylophora pistillata]	310.20	-		x
1270090022	XP_022808576.1	uncharacterized protein LOC111345553 isoform X2 [Stylophora pistillata]	320.10	a,c		x
1270091315	XP_022809269.1	microtubule-associated tumor suppressor 1 homolog isoform X1 [Stylophora pistillata]	259.50	a,c		x
1270091317	XP_022809270.1	microtubule-associated tumor suppressor 1 homolog isoform X2 [Stylophora pistillata]	256.40	-		x
1270093788	XP_022810585.1	von Willebrand factor D and EGF domain-containing protein-like, partial [Stylophora pistillata]	298.80	-		x
1270095516	XP_022778254.1	uncharacterized protein LOC111319781 [Stylophora pistillata]	250.70	c		x
1270095572	XP_022778283.1	uncharacterized protein LOC111319816, partial [Stylophora pistillata]	508.10	-	x	x
190710633	ACE95141.1	carbonic anhydrase [Stylophora pistillata]	495.20	a,c	x	x

NCBI Accession Number	Annotation	gi (NCBI)	Gene model id	RNA id	*Acropora digitifera* skeletal proteins (Takeuchi et al. 2016)	*Acropora millepora* skeletal proteins (Ramos-Silva et al. 2013)	*Stylophora pistillata* skeletal proteins (Drake et al. 2013)
ACE95141.1[a,c]	carbonic anhydrase	190710633	NA	NA			SeriatoporaSp_g7349; mapped to ACE95141.1 (EU532164)
PFX18785.1[a,c]	Mucin-4	1263122270	NA	NA	aug_v2a.09809.t1	JR987773	AGG36340.1
PFX30903.1	hypothetical protein AWC38_SpisGene4292	1263134737	NA	NA	aug_v2a.06327.t1	JR972076.1	AGG36350.1
XP_022779720.1[b,c]	vitellogenin-like	1270036141	gene3329	rna4552	aug_v2a.08856.t1		AGG36347.1, AGG36349.1
XP_022780690.1[a]	skeletal aspartic acid-rich protein 2-like	1270037961	gene4012	rna5498	aug_v2a.11068.t1, JT001945.1, aug_v2a.01441.t1, aug_v2a.01440.t1, JT004498.1, JR991407.1	JR991407.1, JT004498.1	AGG36358.1, AGG36357.1
XP_022780694.1[a]	CUB and peptidase domain-containing protein 2-like	1270037969	gene3981	rna5452		JR970990.1	
XP_022782398.1	skeletal aspartic acid-rich protein 1-like	1270041141	gene5242	rna7162		JT001945.1	AGG36357.1
XP_022783415.1[a]	coadhesin-like isoform X3	1270043038	gene6099	rna8310	aug_v2a.05945.t1	JT016638.1	AGG36341.1, AGG36335.1, AGE45658.1, AGG36343.1
XP_022786918.1[a,c]	major yolk protein-like isoform X2	1270049598	gene8661	rna11721			AGG36338.1
XP_022788227.1[a,c]	hephaestin-like protein	1270052030	gene9579	rna12971	aug_v2a.24015.t1	JT019463.1	
XP_022794122.1	galaxin-like isoform X2	1270063049	gene13931	rna18823	aug_v2a.18631.t1	JR976690.1	
XP_022794736.1	MAM and LDL-receptor class A domain-containing protein 2-like	1270064196	gene14425	rna19450	aug_v2a.09968.t1, aug_v2a.09969.t1	JT011118.1, JR994474.1	AGG36355.1, AGG36356.1, AGG36345.1
XP_022796981.1	uncharacterized skeletal organic matrix protein 8-like	1270068394	gene16153	rna21699		JT014391.1	AGC24391.1
XP_022796982.1	uncharacterized protein LOC111335364	1270068396	gene16156	rna21701			AGC24391.1
XP_022804012.1[a]	EGF and laminin G domain-containing protein-like	1270032917	gene2104	rna2853	aug_v2a.15580.t1, aug_v2a.06123.t1, JR980881.1, aug_v2a.24512.t1, aug_v2a.06122.t1	JR980881.1	AGG36344.1, AGG36352.1
XP_022806326.1[a]	ZP domain-containing protein-like	1270085816	gene23405	rna30870	aug_v2a.07627.t1	JN631095.1	AGG36346.1

ECM/Transmembrane & Protein Modification
Carbonic anhydrase 2-like
Cation channel sperm-associated protein subunit beta-like
Chymotrypsin-like elastase family member 1
Coadhesin-like
Collagenase 3-like
Complement C3
CUB and peptidase domain-containing protein 2-like
Digestive cysteine proteinase 1-like
Endothelin-converting enzyme 1-like
EGF and laminin G domain-containing protein-like
Hephaestin-like protein
MAGUK p55 subfamily member 7-like
Major yolk protein-like
Mammalian ependymin-related protein
Mucin-4
Microtubule-associated tumor suppressor 1 homolog
Nidogen-2
Proto-oncogene tyrosine-protein kinase receptor Ret
Ras-like protein 3
Skeletal aspartic acid-rich protein 2-like
TTransmembrane protease serine 9
Uncharacterized (3)
ZP domain-containing protein-like

Membrane Processing
Complement C3
Endothelin-converting enzyme 1-like
Protein lingerer-like
Uncharacterized (6)
Vitellogenin-like

Vesicle/Secretion & Metal Binding
Carbonic anhydrase 2-like
Cation channel sperm-associated protein subunit beta-like
Chymotrypsin-like elastase family member 1   Collagenase-like
Condensin-2 complex subunit D3-like   Complement C3
Deleted in malignant brain tumors 1 protein-like
Digestive cysteine proteinase 1-like
Endothelin-converting enzyme 1-like   Hephaestin-like protein
Low-density lipoprotein receptor-related protein 8-like   Major yolk protein-like
Mammalian ependymin-related protein 1-like
Microtuble-associated tumor suppressor 1 homolog   Mucin-4
PHD finger protein 21A-like   Nidogen-2
Poly (ADP-ribose) polymerase 11
Phosphopantothenoylcystein decarboxylase subunit VHS3-like
Protein lingerer-like
Proto-oncogene tyrosine-protein kinase receptor Ret Ras-like protein 3
Synapsin-2-like
Thioredoxin reductase 1, cytoplasmic-like
Uncharacterized (5)   Vitellogenin-like

Calicoblastic cell layer

Calcifying medium

Skeleton