1 Optimization of skeletal protein preparation for LC-MS/MS sequencing yields
2 additional coral skeletal proteins in *Stylophora pistillata*
3
4 Yanai Peled[1*], Jeana Drake[1*^], Assaf Malik[1], Ricardo Almuly[1], Maya Lalzar[2], David
5 Morgenstern[3], Tali Mass[1+]
6
[1]Marine Biology Department, University of Haifa, Haifa, Israel
[2]Bioinformatics Core Unit, University of Haifa, Haifa, Israel
[3]De Botton Protein Profiling Institute of the Nancy and Stephen Grand Israel National Center for
Personalized Medicine, Weizmann Institute of Science, Israel
*equal contribution
^Corresponding author during review: jeanadrake@g.ucla.edu
+Corresponding author upon publication: tmass@univ.haifa.ac.il

Abstract

Stony corals generate their calcium carbonate exoskeleton in a highly controlled

biomineralization process mediated by a variety of macromolecules including

proteins. Fully identifying and classifying these proteins is crucial to understanding

their role in exoskeleton formation, yet no optimal method to purify and characterize

the full suite of extracted coral skeletal proteins has been established and hence their

complete composition remains obscure. Here, we tested four skeletal protein

purification protocols using acetone precipitation and ultrafiltration dialysis filters to

present a comprehensive scleractinian coral skeletal proteome. We identified a total of

60 proteins in the coral skeleton, 44 of which were not present in previously published

stony coral skeletal proteomes. Extracted protein purification protocols carried out in

this study revealed that no one method captures all proteins and each protocol

revealed a unique set of method-exclusive proteins. To better understand the general

mechanism of skeletal protein transportation, we further examined the proteins' gene

ontology, transmembrane domains, and signal peptides. We found that

transmembrane domain proteins and signal peptide secretion pathways, by

themselves, could not explain the transportation of proteins to the skeleton . We

therefore propose that some proteins are transported to the skeleton via non-traditional

secretion pathways.

Keywords

Skeleton organic matrix proteins (SOMPs), biomineralization, Scleractinia

Introduction

Scleractinian corals, or stony corals, are the most prolific biomineralizers in phylum Cnidaria [1]. They are a key component of shallow-water tropical reefs, often forming massive structures that serve as the foundation of an ecosystem which hosts some of the more biodiverse communities on the planet [2, 3]. They are amongst the oldest biomineralizing metazoans, producing calcium carbonate ($CaCO_3$) exoskeletons in the form of aragonite through biologically-directed mechanisms [4, 5], which makes up ≥95% of the entire skeletal mass, the remainder of which is the skeletal organic matrix (SOM) [6-8].

Biomineralization refers to the ability of a living organism to selectively exploit elements from its surrounding environment to build a biologically functioning crystalline structure [9, 10]. Biomineralizers can be found throughout all kingdoms of life, from bacteria [e.x. 11], to algae [e.x., 12], mollusks [e.x., 13], corals [e.x., 10] and mammals [e.x., 14]. The minerals formed through this process differ in structure from their non-biological counterparts and their formation is mediated by a variety of organic molecules (the SOM), which have been intensively studied since the 1960's in diverse organisms [reviewed by 10]. The SOM consists of proteins, lipids, and polysaccharides, which are not present in the abiotic mineral form [reviewed by 15]. SOM proteins (SOMPs) embedded within the mineral are hypothesized to serve as a framework for crystal nucleation [reviewed by 16]. The proteins involved in the process of skeletal biomineralization have been described and studied extensively in echinoderms [e.x., 17, 18] [and reviewed by 19, 20], mollusks [e.x., 21, 22-24] and

mammals [e.x., 25, 26, 27], among others [28]. At present, the best described SOMP

complex is of mammalian bone and teeth [15].

In stony corals, the most current knowledge of SOMPs is limited and based on

intraskeletal protein extraction [29-33] . It has been suggested that coral SOMPs aid in

the molecular processes of crystallization as well as in the development and

strengthening of the minerals ([34-37] among others). The constant advancement of

mass spectrometry technology has broadened our capability to identify many proteins

in skeletal extracts, even those proteins in low abundance [38]. However, this

technology is sensitive to contamination by organic matter remnants from soft tissue

and cell debris from the study organism, and the little-addressed issue remains of

contamination by researchers during the protein extraction, preparation, and

sequencing steps [39-41]. Upon extraction, the SOMPs are usually divided into two

fractions: soluble and insoluble matrix proteins (SSOM and ISOM respectively),

based on their solubility in the acid of choice or in water [22, 31-33, 42]. While some

past attention has been directed towards the soluble fraction [30, 43, 44], Pereira-

Mouriès et al. [45] showed that, in the bivalve *Pinctada maxima*, the classification of

SSOM and ISOM is misleading and that both fractions share common features.

Furthermore, Goffredo et al. [42] found in the stony coral *Balanophyllia europaea*

that both fractions consist of the same macromolecules; they associated the degree of

solubility to the difference in cross-linking. They also showed that each solubility

fraction has a different influence on calcium carbonate crystal morphology,

aggregation, and polymorphism in-vitro. In contrast, Ramos-Silva et al. [32] observed

a different SOMP composition between solubility fractions in the scleractinian coral

Acropora millepora,. Out of 36 SOMPs, only two were found exclusively in the

soluble fraction and twelve were exclusive to the insoluble fraction. These examples

demonstrate the attempts to attribute different properties to the two fractions, but the

data remain inconclusive.

To date, three major coral skeletal proteomes have been published [31-33]

with each proteome consisting of 30-40 proteins. Of the 30 proteins sequenced from

A. digitifera skeleton, 26 were also detected in *A. millepora* skeleton [32, 33]. They

consist mostly of either transmembrane (TM) domain proteins or secretory proteins

[33]. However, only 12 of the proteins identified in *A. millepora* skeleton matched

those found in *S. pistillata* skeleton [31, 32]. In *A. millepora*, 11 TM domain-

containing proteins were identified, as well as two proteases that were not detected in

S. pistillata [32]. The authors suggested that the proteases' role is in cleaving the

extracellular domain of TM proteins and incorporating them into the skeleton.

The coral skeletal proteomes published to date reveal an overlap of several

detected proteins, but at least 10 proteins from each species appear to be unique. It is

currently unknown if this is truly due to species-specific gene expression and protein

localization or to methods in extracting, purifying, and sequencing the proteins. In

this study we analyzed several methods for extracted protein purification to increase

the detection of the full suite of SOMPs from cleaned coral skeleton powder. We

show that the use of acetone precipitation versus centrifugal filter washing, and the

degree to which each purification method is performed, affects the numbers and types

of proteins that can be sequenced by mass spectrometry. Further, we suggest that

there is no one 'best' method for coral skeletal protein purification to capture all

SOMPs such that future research projects may need to utilize several preparation

methods to detect the full breadth of proteins embedded in coral skeleton.

Methods

Sample collection and preparation for protein extraction

The hermatypic coral *Stylophora pistillata* (Esper, 1797) was collected under

a special permit from the Israeli Natural Parks Authority in the waters

in front of the H. Steinitz Marine Biology Laboratory, Eilat, Israel, Red Sea (29°30 N,

34°56 E), using SCUBA diving.

We fragmented one *S. pistillata* colony into small pieces, approximately 2x2

cm, with a diamond band saw. Coral fragments were transferred to 50-ml Falcon

brand conical vials (Falcon tubes) and oxidized with 20 mL 1:1 of 30% H_2O_2: 3%

NaClO solution for 1 hour, during which 1.5 mL of 3% NaClO solution were gently

added to the tubes every 20 minutes and continued the incubation overnight at room

temperature following modified methods of Stoll et al. [46]. Fragments were washed

five times with ultra-pure water for one minute each time and dried at 60 C°

overnight. We crushed the cleaned fragments to ≤63 µm diameter with a mortar and

pestle. Skeleton powder, in sterile Falcon tubes, was then oxidized and washed in

ultra-pure water three more times (i.e., four complete rounds of oxidative cleaning) to

ensure that no organic residue remained on the skeletal grains. In each cycle, the

removal of the oxidizing or wash solution was performed by centrifugation at 5,000 x

g for 3 min at 4 C°. Cleaned skeletal powder was then dried overnight at 60 C°. We

carried out all the described processes in a laminar flow biological hood (apart from

oven drying) with all preparation tools and surfaces bleached to avoid contamination.

To monitor the removal of proteins from the skeletal powder, we checked the

cleaning efficiency under SEM after the fourth oxidative cleaning. Samples were

sputter-coated with 4 nm gold prior to examination using a ZEISS Sigma

TM scanning electron microscope with in-lens detector (5kV, WD = 5-7mm) (SI

Figure 1a,b) [17, 47]. In addition, we sonicated the cleaned powder at 4°C in filter-

sterilized phosphate buffered saline (PBS, pH 7.4) for 30 minutes, pelleted the powder

at 5,000 x *g* for 3 minutes at 4˚C, concentrated the supernatant on a 3-kDa cutoff

centrifugal filter unit (Amicon) and loaded samples of supernatant on a 8-16% SDS-

PAGE TGX Stain-Free gels (Bio-Rad) (SI Figure 1c).

Extraction and purification of skeletal proteins

We used four samples of approximately 1.3 g cleaned skeleton powder each to test

four protein purification protocols as described in Figure 1. All samples were

decalcified in 0.5 M acetic acid (30 ml acid/g cleaned skeleton powder) in Falcon

tubes while rotating the tubes at room temperature for 3 hours. Samples were then

centrifuged at 5,000 x *g* for 5 min at 4^{o}C and supernatant was transferred to a new

tube and stored at 4^{o}C. We continued the decalcification of the undissolved pellets

with a second volume of 0.5 M acetic acid and allowed decalcification to proceed to

completion, at which point the pH of the solution was measured at 5.5-6.5. We then

combined both decalcification rounds (70 ml total) for each sample containing both

the acid-soluble and -insoluble fractions, froze the total volumes at -80˚C, and dried

them by overnight lyophilization. The dried pellets were stored at -80°C until further

processing. Because washes of the extracted proteins can reduce the representation of

proteins, in this study we examined two protein concentrating and cleaning methods

similar to those performed for previously published coral skeletal proteomes; (i)

centrifugation ultrafiltration (CF methods [32]) and (ii) acetone precipitation (ACT

methods, [31]) (Figure 1; SI Table 1).

To desalt and concentrate the extracted SOM, the lyophilized pellets of all

samples were re-suspended in 12 ml MilliQ water and the proteins were concentrated

on 3 kDa cutoff Amicon® Ultra 15 centrifugal filter units (Merck-Millipore) 5,000 x

g at 4˚C to reach a final volume of 0.5 ml. This process was repeated once. At this

stage, we observed a water-insoluble pellet (ISOM) in all samples, which for samples

CF2, ACT1, and ACT3 was pelleted from the water-soluble fraction (SSOM) by

centrifugation at 5,000 x g for 5 min at 4°C. This initial wash, concentration, and

solubility fractionation protocol was the terminal step for centrifugal filter sample

CF2. Sample CF4 was resuspended one more time in MilliQ and concentrated before

separation of SSOM and ISOM as above. For acetone precipitation samples ACT1

and ACT3, the two rounds of desalting and concentration were followed upon with

successive washes of the SSOM and ISOM in acetone. To sample ACT1 SSOM was

added 2 ml 100% ice cold acetone. The sample was vortexed for 10 seconds,

incubated at -20°C for 30 minutes, and centrifuged at 4,300 x g for 30 min at 4°C.

The resulting pellet was washed three more times with 2 mL of ice cold 80% acetone.

The ISOM fraction was washed four times with 80% acetone. Both solubility

fractions of sample ACT3 were treated as in ACT1 but with one less washing step of

each fraction. All fractions across all purification treatments were stored at -80°C. An

aliquot of each fraction was analyzed by SDS-PAGE on a 8-16% SDS-PAGE TGX

Stain-Free gels with subsequent silver staining and displays the commonly-observed

smearing of extracted biomineral proteins with minimal observable banding (SI

Figure 2).

LC MS\MS

S. pistillata skeletal protein samples were dissolved in 5% SDS and digested with

trypsin using the S-trap method overnight at room temperature. We analyzed the

resulting peptides using a nanoflow ultra-performance liquid chromatograph

(nanoAcquity) coupled to a high resolution, high mass accuracy mass spectrometer

(Fusion Lumos). The sample was trapped on a Symmetry C18 0.18*20mm trap

column (Waters, Inc) and separated on a HSS T3 0.075*250 mm column (Waters,

Inc.) using a gradient of 4-28% (80% acetonitrile, 0.1% Formic acid) for 150 minutes.

Spray voltage was set to +2kV. The data were acquired in the Fusion Lumos using a

Top Speed Data-Dependent Acquisition method using a cycle time of 3 s. An MS1

scan was performed in the Orbitrap at 120,000 resolution with a maximum injection

time of 60 ms. The data were scanned between 300-1800 m/z. MS2 was selected

using a monoisotopic precursor selection set to peptides, peptide charge states set to

+2-+8 and dynamic exclusion set to 30 s. MS2 was performed using HCD

fragmentation scanned in the Orbitrap, with the first mass set to 130 m/z at a

resolution of 15,000. Maximum injection time was set to 60 ms with automatic gain

control of $5x10^{-4}$ ions as a fill target. The resulting data were searched against the

NCBI *Stylophora pistillata* protein database using the Byonic search engine (Protein

Metrics Inc.) – the first search was carried out without any false discovery rate (FDR)

filtering, to generate a focused database for a second search. The second search was

set to 1% FDR, allowing fixed carbamidomethylation on C and variable oxidation on

MW, deamidation on NQ and protein N-terminal acetylation. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the

Pride partner repository [48], under the dataset identifier PXD017891; reviewer login

username and password have been provided to the reviewers. Data will be made

publicly available without a required login in ProteomeXchange upon publication of

the manuscript.

Data sorting

We used the *S. pistillata* genome database as a reference peptide database for the

mass spectrometry analysis [49] (NCBI BioProjects PRJNA281535 and

PRJNA415215) appended with known NCBI *S. pistillata* skeletal proteins[e.x., 31,

50] *S. pistillata* carbonic anhydrase, ACE95141.1). We also included a common

contaminants database. Despite this inclusion of a contaminants database, several

proteins likely of human origin were sequenced and attributed to *S. pistillata*. To

filter out these potential contaminants from our final list of coral-specific proteins, we

BLASTed all sequences against the 'Primates' database in NCBI using Blast2GO.

We then examined NCBI-generated sequence alignments of coral versus *Homo*

sapiens proteins with e-values lower than e^{-50} and percent mean similarity greater than

50%, all sequences with e-values lower than e^{-100}, and all sequences with percent

similarity greater than 80%, and removed from our final list of coral proteins any

sequences with three or more peptides each of seven or more amino acids in length

that were identical between *S. pistillata* and humans.

All remaining coral-specific proteins identified by the LC MS\MS analysis

were filtered to those with at least two significant peptides or at least one significant

peptide with at least 10 spectra and an identification score of 250 or greater. Skeletal

proteins were first sorted by fractions and methods (i.e., SSOM and ISOM for each

purification method). Next, we sorted all skeletal proteins by their gene ontology

(GO) terms [51, 52]. Finally, we grouped all by terms of interest: protein

modification, transmembrane, and ECM; membrane processing; metal binding and

vesicular.

Proteins detected by LC-MS/MS were annotated using the Trinotate pipeline

which relies on both Pfam and UniProt data [53]. GO terms and Pfam annotations

were assigned to *Stylophora pistillata* predicted proteins and transcripts using

Trinotate 3.0.1 (https://github.com/Trinotate).   Transmembrane regions were

predicted using the TMHMM server v2.0 [54]. Signal peptides on the N-termini of

proteins were predicted using SignalP 5.0 [55, 56]. Attachment of proteins to the

exterior of the cell membrane by glycosylphosphatidylinositol (GPI) anchors was

predicted using PredGPI [57]. These computational analyses used program default

settings and cutoffs.   Completeness of protein sequences was determined by

comparing all returned coral proteins to the *Acropora digitifera* genome [58]; NCBI

RefSeq assembly GCF_000222465.1). Detection of various proteins across solubility

and protocol fractions was visualized in Venny 2.1 online software.

To examine the conserved coral biomineralization proteins, we determined

orthologous biomineralization genes, known from skeletal proteomic analysis, across

coral taxa. First, we estimated orthology relationships between all non-redundant

genes of selected metazoa species using OrthoFinder. We included all genes of all

Cnidaria species with known genome-based annotations. OrthoFinder generates

orthology groups (Orthogroups) based on normalized reciprocal best BLAST hits' bit

scores [59], and then estimates orthologues genes pairs within Orthogroups [60]. We

then selected all pairs of *Acropora* spp. orthologs to *S. pistillata* (1:1, 1:many,

many:many relationships) ([31-33], this study). From these pairs we further selected

S. pistillata spectra-based identified proteins, or skeletal proteins known in the

literature. Since not all skeletal protein annotations from the literature were included

in our reference OrthoFinder proteome datasets, we further found their best matches

in the reference OrthoFinder proteome using BLASTP.

Results:

After extensive cleaning of the powdered skeleton and acid-extraction of embedded

organic matter we identified in total 60 coral-specific proteins meeting our criteria in

S. pistillata skeleton as predicted by the species genome [49] (Table 1, SI Table 2).

Trinotate [53, 61] returned GO annotations [51] for all of these proteins, although

many remain uncharacterized (SI Table 3).

In order to evaluate the efficacy and improve current methods for stony coral

skeletal protein purification, we examined four different protocols; two centrifugation

ultrafiltration filters (CF) and two further acetone precipitation (ACT) protocols.

Proteomes of the four methodologies differed in composition and variety (Figure 2).

Combining results of all CF fractions identified 52 coral-specific proteins while

combined ACT protocols yielded 13 such proteins (Figure 2A). Moreover,

redundancy between methodologies was low. Only 8.3% of the proteins overlapped

between methods while 78.3% and 13.3% of the proteins were exclusive to combined

CF and combined ACT fraction data, respectively.

To evaluate the purification efficiency of each protocol, we first compared the

number of proteins detected in each method (e,x,, ACT1 vs ACT 3 and CF2 vs CF4).

Of the 8 proteins found only in samples from the acetone wash protocols (ACT 1 and

ACT 3), one was observed only in ACT1 while five were observed only ACT3

(Figure 2, SI Table 2); both ACT1 solubility fractions had one more wash step than

did those in ACT3. In contrast, of the 47 proteins observed only in CF samples, 12

were found only in CF2, which went through two filter centrifugation steps, while 16

were found only in CF4, which went through a third filter centrifugation step; the

remaining 19 were found using both CF protocols. Further, only two proteins were

observed in all four purification fractions in at least one solubility form:

CARP4/SAARP1 and synapsin 2-like. Two additional proteins were observed in

three of the four protocols: STPCA2 and an uncharacterized protein.

We next examined the difference in protein composition obtained from the

solubility fractions SSOM versus ISOM. We identified different distributions of

proteins in the SSOM vs. ISOM in both purification methods (Figures 2B & C).

Notably, when combining all methods, more proteins were identified in the ISOM

compared to the SSOM. A total of 57 coral proteins were identified in the ISOM

fraction compared to a total of 8 in the SSOM fraction (SI Table 2), of which 5 (8.3%)

were identify both in SSOM and ISOM.

Our data compared to other coral skeletal proteomes

Since our analysis yielded a large amount of new skeletal proteins, we also compared

our results with the three previously published proteomes of *S. pistillata, A. digitifera*

and *A. millepora* [31-33]. Out of our entire identified skeletal proteome containing 60

proteins, using OrthoFinder and BLASTP, only 16 were found to be similar to

proteins identified in these studies. Yet, this proportion of overlap (16 out of 60) is

significantly greater than the expected proportion by chance, since the proportion of

known skeletal matrix proteins in the reference coral proteomes is extremely small

(less than ~0.2%). Seven proteins were found to overlap all four proteomes: a

coadhesin-like protein, an EGF and laminin G domain-containing protein, a

hypothetical protein, a MAM and LDL-receptor class A domain-containing protein, a

mucin, aspartic acid-rich protein 2-like, and a ZP domain-containing protein (Table

2).

Skeletal proteome characterization

We interrogated the mechanisms by which proteins may be exported from or attached

to the cell (SI Table 2). Seventeen coral skeletal proteins with likely complete N-

terminus predictions possess signal peptides as a potential mechanism for export from

the cell. Eight proteins contain at least one transmembrane span suggesting that they

are embedded in the cell membrane. Further, 10 proteins likely interact with the

exterior of the cell membrane by GPI anchors. In total, 25 of the 60 sequenced coral

skeletal proteins exhibit documented characteristics for localization in the ECM.

Because the majority of the proteins sequenced from the *S. pistillata* skeleton

do not possess features for signaling their export from the cell, we queried the data set

for further suggestions of positioning the proteins in the membrane or that the proteins

may be exported by vesicles such as those that may be involved in calcium

concentration. To do this, we examined the skeletal proteome annotations and GO

classifications toward finding common features to allow grouping of proteins. Out of

the entire skeletal proteome sequenced in this study (60 proteins), 39 genes were

returned with GO terms that allowed their classification into five groups of interest

based on their cellular component, biological process, and molecular function to

suggest likely cellular locations pertinent to the calcification mechanism, which may

therefore be indicative of their function in this process: lipid\phosphate\glycan related

proteins (i.e., membrane processing); ECM-related, transmembrane, and protein

modification proteins; and metal binding proteins vesicular/secretion related proteins,

(SI Table 3).

Of the proteins with GO terms, 10 proteins are suggested to be involved with

processing of the cell membrane (Figure 3), A much larger number are related to

vesicles/secretion as well as binding metal, with 21 and 25 assigned to each category,

respectively.   We combined these two categories in our proposed cellular location

Figure 3, as some of the skeletal proteins proposed to be found intracellularly in

vesicles are also known to bind calcium [34, 62]. Finally, 19 and 12 are potential

ECM proteins or are involved in protein modification, respectively. Of the proteins in

the vesicles/secretion and ECM categories, several appear to be completely predicted,

based on comparison to orthologs in the *A. digitifera* genome, yet lack a signal

peptide (SI Table 2). It should be noted that many proteins are assigned to multiple

categories.

Discussion:

In this study we show the importance of using complementary post-extraction

methods to purify and concentrate coral skeletal proteins for sequencing the full

breadth of the skeletal proteome. Our results show a clear and marked difference in

detected proteins between protein purification methods, with only two proteins

observed in all methods. Centrifugal Filter (CF) methods yielded a much greater

abundance and diversity of proteins than did acetone precipitation (ACT) (Figure 2).

Moreover, we found that protein detection is largely method-exclusive (Figure 2). We

hypothesize that the protein yield differences between purification methods is based

on the different properties of each method. CF is a mechanical filtration based on size

and has a bias toward hydrophobic proteins, whereas ACT is based mainly on the

chemical interactions of proteins resulting in increased precipitation of hydrophilic

proteins [63]. We speculate that the CF methods yielded more total proteins in our

study as acetone precipitation does not recover all proteins [64] and those proteins

which are recovered in a pellet may be difficult to resolubilize due to molecular

interactions formed within the aggregate by protein denaturation [65], potentially

leading to the loss of many proteins in the pellet which are not transferred to the

trypsin digestion step.

A major challenge in working with intra-skeletal proteins is isolating the true

skeletal proteins from soft tissue contamination [40, 41]. In the present study, as in

previous work on *S. pistillata* skeletal proteins [40] we carried out an intensive

oxidative cleaning step on the skeletal powder, in addition to cleaning the skeletal

fragments, to avoid contamination. We did not observe any organic residues on

cleaned powders examined by SEM (SI Figure 1a,b) or in PBS soaked on the

powders and concentrated (SI Figure 1c), and we are therefore confident that all

sequenced proteins are endogenous to the skeleton and are not coral cellular

contamination.

In this study, we examined the protein composition of the SSOM and ISOM

proteins separately. Although we see variations due to the different purification

techniques (Figure 2), our results indicate that, broadly, the ISOM is distinguished in

composition from the SSOM. Across all purification techniques, the ISOM fractions

yielded more total proteins and fraction-exclusive proteins than the SSOM fraction

did (Figure 2B, C). Since all extraction and purification experiments were carried out

simultaneously, we can rule out batch effect and technical differences. This ISOM

versus SSOM skeletome fractional composition is similar to previous studies carried

out on other marine biomineralizers such as mollusks [45] as well as to the stony

corals *A. millepora* [32] and *S. pistillata* [31]. However, the overlap between the two

solubility fractions reported for *A. millepora* [32] was 61%, in contrast to our study

which shows an overlap across all purification treatments combined of only 10%

(Figure 2). One plausible explanation for this difference is the use of improved LC-

MS/MS performance over the past several years enabled us to detect more proteins

and obtain a more comprehensive proteome. In previous studies of coral skeletal

proteomes which detailed differences between solubility fractions, the instruments

used were low resolution, low mass accuracy, which tend to result in a lower percent

identification of the data. Additionally, most search engines and FDR-calculating

algorithms struggle with very small datasets [66, 67]. Our use of Byonic helps to

alleviate statistical limitations that can result in false negative results.

Further differences in proteomes, beyond species differences, are the differing

reagents used in precipitation (compared to [31]) and our smaller centrifugal filter

cutoffs (compared to [32]). This second difference is particularly important with

respect to protein degradation. Even when embedded in biominerals so that amino

acids and even short peptides persist, proteins may still succumb to degradation [68].

This result in small peptide fragments that may be lost from centrifugal filter units

with 10 kDa pore sizes and larger. It is reasonable to assume that coral skeletal

proteins go through the same process, and if so, the cutoff of the membrane directly

affects the number of peptide spectra matches (PSM). SDS-PAGE analysis shows

smearing of extracted skeletal proteins, which in addition to likely differential

addition of glycans, sulfates, and phosphates to proteins [e.x., 30, 69, 70], may

indicate protein degradation (SI Figure 2). Using smaller cutoff filters in this study

might have allowed us to capture some of these sheered peptides and led to higher

PSMs.

Traditional protein trafficking by signal peptides and transmembrane domain does

not explain the full extent of protein transportation to skeleton

Corals' skeleton is external to the animal; therefore, proteins in the skeletal matrices

must be transported outside the cells or span the membrane and have an extracellular

portion to reach the skeletal crystallization front. Indeed, recent studies of anthozoans

reveal a significant proportion of TM domain proteins (~35%) in the SOM [32, 33],

similar to that found in the better-studied Echinoidia [71] . Based on these findings,

we examined the hypothesis that many of the skeletal proteins originate in the plasma

membrane.

Our analysis revealed that TM domain proteins are not the major component

of the SOM protein complex; in the present proteome, TMHMM prediction suggests

that eight are embedded in the membrane. Hence, we examined a cellular secretion

option. Ramos-Silva et al. [32] reported 15 proteins (41%) with SPs that did not also

possess a TM domain in the *A. millepora* proteome. In their study of the *A. digitifera*

proteome, Takeuchi et al. [33] reported a similar proportion of proteins with SPs but

no TM domains (40%). In the present S. *pistillata* proteome, 17 out of 60 (28%) were

positive for SPs (SI Table 2, [31]), likely a combination of incomplete gene prediction

and other mechanisms for exporting the proteins to the calicoblastic space.

Hydrophobic regions of TM domains are similar to those of SPs, making SP

prediction difficult [54, 72]. We therefore cannot say for sure, based on these two

analyses (TMHMM and SignalP) alone, which is the more dominant pathway for

protein transport for many of the SOMPs. Further, while SP by itself can help predict

traditional secretory pathways [72], proteins that do not possess classical SPs or TM

regions may still leave the cell via other means (reviewed in [72-74]). This includes

membrane pores [72], ATP-binding cassette transporters [75] and

autophagosome/endosome-based secretion [73, 76], as well as exosomes [77] which

can transport proteins that lack signal peptides in extracellular vesicles [78]. Since

TM and SP analysis did not fully explain the mechanism of protein transportation, we

classified the likely functions and locations of the skeletal proteins according to their

GO terms as lipid\phosphate\glycan related proteins, metal binding proteins,

vesicular/secretion related proteins, ECM-related and transmembrane proteins, and

protein modification proteins (Figure 3, SI Table 3). Looking at the proposed

locations or functions of skeletal proteins (Figure 3), we find comparable numbers of

vesicular and metal binding proteins versus TM domain/ECM proteins. Further,

several of these proteins appear to be completely predicted at their N-termini yet lack

signal peptides (SI Table 2). These results are in-line with the trend reported in

previous coral skeletal proteomes [31-33]. These proteins may leave the calicoblastic

cells through one of the non-classical mechanisms described above. While we do not

exclude the role of SP and TM proteins in the skeletal deposition process, we suggest

that the biological mechanism of SOM transportation is enhanced by exosomes and

other non-traditional secretion pathways.

A further method by which proteins may be exported to the site of

calcification is a vesicular pathway that differs from the conventional SP and TM

pathway and remains to be fully characterized. Previous studies have shown Ca^{2+} rich

granules in the calicoblastic epithelium (skeletogenic cells), but not in the other tissue

layers, suggesting their role as a Ca^{2+} reservoirs in the cells. Vesicles were previously

identified in corals [79, 80]; however, their origin and content was not detailed, and

they are sometimes attributed to preservation byproducts. These intracellular ion-rich

vesicles may endocytose sea water through macropinocytosis [81, 82], after which

they are enriched in carbonate ions and then form hydrated ACC and anhydrous ACC

precursors stabilized by acidic biomolecules including CARPs [83]. Using cell

cultures, Mass et al. [62] suggest that the vesicles, which contain Asp rich proteins,

then transport their contents to the ECM, releasing their content by exocytosis. The

biomineral then further develops extracellularly, likely aided by other ECM proteins

[30, 35, 50, 84] as well as other biomolecules [85-88]. At present the processes of

calcium delivery to the skeleton and the roles of most of the proteins in coral skeletal

deposition remain to be determined.

It is difficult to map and characterize proteins in non-model organisms such *S.*

pistillata [89-91]. High quality proteomic mapping requires knowledge of

phosphorylation, glycosylation, proteolytic cite activities and other modifications [31,

92-94], in order to create a more thorough database [89-91]. Further, many of the

coral skeletal proteins reported to date remain uncharacterized. Uncharacterized

proteins were reported in *Acropora* skeletons at a rate of approximately 25% [32, 33]

while, they were previously reported at less than 10% in *S. pistillata* skeleton [31].

Our study revealed a greater proportion, of 28% uncharacterized proteins, in *S.*

pistillata skeleton, in line with *Acropora* spp. While partially attributed to sample

size, it is most likely due to quality of genomic data available, since stony corals are

non-model organisms and their genomic libraries are far from complete, resulting in

incomplete databases on which to map the proteome and many uncharacterized genes.

Conclusion:

In this study we have considered the differential effects of coral skeletal protein

extract preparation as well as the method by which these proteins, or parts thereof, are

transported from intracellular to extracellular locations. When preparing coral skeletal

proteomes, we propose that a multi-method approach to cleaning, demineralization,

and protein purification should be used. Our results showed that each protein

preparation protocol yielded exclusive sets of proteins with little overlap between

ACT and CF fractions. While CF protocols yielded many more proteins than did ACT

methods, use of a single protocol to clean and concentrate coral skeletal proteins

results in a significant amount of data loss, and it is therefore of crucial importance to

consider alternative and complementary methods to obtain a fully comprehensive

skeletal proteome. We showed that while the role of TM domain proteins cannot be

overlooked, many of the proteins detected in the *S. pistillata* skeletal proteomes as

well as in that of other species point toward other secretory or vesicular pathways.

Our categorization method, supported by data from other recent studies, also suggests

that corals use an alternative secretory pathway, such as exosomes or non-classical

secretion vesicles, and much work is required in order to determine the calcium

deposition pathway and the proteins involved. Our study provides a large set of new

uncharacterized coral skeletal proteins as well as others of purported function but that

have not been observed before in the coral SOM. These data expand the current

knowledge of the SOM in corals and will help, in future studies, to resolve corals'

calcium deposition mechanism and the various roles of the proteins involved.

Abbreviations

acetone precipitation of proteins (ACT)

centrifugation ultrafiltration of proteins (CF)

false discovery rate (FDR)

gene ontology (GO)

glycosylphosphatidylinositol (GPI)

insoluble skeletal organic matrix (ISOM)

peptide spectra matches (PSM)

phosphate buffered saline (PBS)

skeletal organic matrix (SOM)

skeletal organic matrix proteins (SOMP)

soluble skeletal organic matrix (SSOM)

transmembrane (TM)

Declarations

Ethics approval and consent to participate

Coral nubbins were collected under a special permit from the Israeli Natural Parks

Authority.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated during the current study are available in the ProteomeXchange

repository under file number PXD017891 (https://www.ebi.ac.uk/pride/archive). The

reviewer login username and password have been provided to the reviewers. Data

will be made publicly available without a required login in ProteomeXchange upon

publication of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Israel Science Foundation (Grant 312/15), and the

European Research Commission (ERC; Grant # 755876). JLD was supported by the

Zuckerman STEM Leadership Program.

Author contributions

YP, JD, and TM designed the study. YP and RA prepared the samples. YP, RA, AM,

ML, DM, and JD analyzed the data. All authors wrote the manuscript and approve

this submission.

Acknowledgements

We thank staff at the Bioinformatics Core Unit, University of Haifa, and The Crown

Genomics Institute of the Nancy and Stephen Grand Israel National Center for

Personalized Medicine, Weizmann Institute of Science.

References

1.   Cairns SD: **Deep-water corals: An overview with special reference to**
522   **diversity and distribution of deep-water scleractinian corals.** *Bull Mar Sci*
523   2007, **81:**311-322.
524   2.   Reaka-Kudla ML: **The global biodiversity of coral reefs: A comparison**
525   **with rain forests.** *Biodiversity II: Understanding and Protecting our*
526   *Biological Resources* 1997, **2:**551.
527   3.   Knowlton N, Brainard RE, Fisher R, Moews M, Plaisance L, Caley MJ: **Coral**
528   **reef biodiversity.** *Life in the world's oceans: Diversity distribution and*
529   *abundance* 2010**:**65-74.
530   4.   Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D, Peterson KJ:
531   **The Cambrian conundrum: Early divergence and later ecological success**
532   **in the early history of animals.** *Science* 2011, **334:**1091-1097.
533   5.   Tambutté S, Tambutté E, Zoccola D, Allemand D: **Organic matrix and**
534   **biomineralization of scleractinian corals.** *Handbook of Biomineralization:*
535   *Biological Aspects and Structure Formation* 2007**:**243-259.

536 6. Cuif J-P, Dauphin Y, Berthet P, Jegoudez J: **Associated water and organic**
537   **compounds in coral skeletons: Quantitative thermogravimetry coupled to**
538   **infrared absorption spectrometry.** *Geochemistry Geophysics Geosystems*
539   2004, **5:**Q11011.
540 7. Wainwright SA: **Skeletal organization in the coral,** *Pocillopora damicornis*.
541   *Quarterly Journal of Microscopical Science* 1963, **s3-104:**169-183.
542 8. Von Euw S, Zhang Q, Manichev V, Murali N, Gross J, Feldman LC,
543   Gustafsson T, Flach C, Mendelsohn R, Falkowski PG: **Biological control of**
544   **aragonite formation in stony corals.** *Science* 2017, **356:**933-938.
545 9. Mann S: *Biomineralization: Principles and Concepts in Bioinorganic*
546   *Materials Chemistry.* New York: Oxford University Press; 2001.
547 10. Weiner S, Dove PM: **An overview of biomineralization processes and the**
548   **problem of the vital effect.** *Reviews in Mineralogy and Geochemistry* 2003,
549   **54:**1-29.
550 11. Schüler D, Frankel RB: **Bacterial magnetosomes: Microbiology,**
551   **biomineralization and biotechnological applications.** *Appl Microbiol*
552   *Biotechnol* 1999, **52:**464-473.
553 12. Westbroek P, Young J, Linschooten K: **Coccolith production**
554   **(biomineralization) in the marine alga** *Emiliania huxleyi*. *The Journal of*
555   *Protozoology* 1989, **36:**368-373.
556 13. Addadi L, Joester D, Nudelman F, Weiner S: **Mollusk shell formation: A**
557   **source of new concepts for understanding biomineralization process.**
558   *Chemical European Journal* 2006, **12:**980-987.
559 14. Levine P, Glimcher M: **The isolation and amino acid composition of the**
560   **organic matrix and neutral soluble proteins of developing rodent enamel.**
561   *Archives of Oral Biology* 1965, **10:**753-IN751.
562 15. Beniash E: **Biominerals—hierarchical nanocomposites: The example of**
563   **bone.** *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*
564   2011, **3:**47-69.
565 16. Lowenstam HA, Weiner S: *On Biomineralization.* New York, New York:
566   Oxford University Press; 1989.
567 17. Karakostis K, Zanella-Cléon I, Immel F, Guichard N, Dru P, Lepage T,
568   Plasseraud L, Matranga V, Marin F: **A minimal molecular toolkit for**
569   **mineral deposition? Biochemistry and proteomics of the test matrix of**
570   **adult specimens of the sea urchin** *Paracentrotus lividus*. *Journal of*
571   *Proteomics* 2016, **136:**133-144.
572 18. Shashikant T, Khor JM, Ettensohn CA: **From genome to anatomy: The**
573   **architecture and evolution of the skeletogenic gene regulatory network of**
574   **sea urchins and other echinoderms.** *Genesis* 2018, **56:**e23253.
575 19. Nebelsick JH, Dynowski JF, Grossmann JN, Tötzke C: **Echinoderms:**
576   **Hierarchically organized light weight skeletons.** In *Evolution of Lightweight*
577   *Structures.* Springer; 2015: 141-155
578 20. Killian CE, Wilt FH: **Molecular aspects of biomineralization of the**
579   **echinoderm endoskeleton.** *Chemical Reviews* 2008, **108:**4463-4474.
580 21. Weiner S, Hood L: **Soluble protein of the organic matrix of mollusk shells:**
581   **A potential template for shell formation.** *Science* 1975, **190:**987-989.
582 22. Mann K, Edsinger-Gonzales E, Mann M: **In-depth proteomic analysis of a**
583   **mollusc shell: Acid-soluble and acid-insoluble matrix of the limpet** *Lottia*
584   *gigantea*. *Proteome Science* 2012, **10:**1-18.

23. Politi Y, Mahamid J, Goldberg H, Weiner S, Addadi L: **Asprich mollusk
shell protein:** *In vitro* **experiments aimed at elucidating function in CaCO$_3$**
crystallization. *CrystEngComm* 2007, **9:**1171-1177.

24. Bouyoucef M, Rakic R, Gómez-Leduc T, Latire T, Marin F, Leclercq S,
Carreiras F, Serpentini A, Lebel J-M, Galéra P: **Regulation of extracellular
matrix synthesis by shell extracts from the marine bivalve** *Pecten
maximus* **in human articular chondrocytes—application for cartilage
engineering.** *Mar Biotechnol* 2018**:**1-15.

25. Weiner S, Wagner HD: **The material bone: Structure-mechanical function
relations.** *Annual Review of Materials Science* 1998, **28:**271-298.

26. Reid DG, Duer MJ, Murray RC, Wise ER: **The organic–mineral interface in
teeth is like that in bone and dominated by polysaccharides: Universal
mediators of normal calcium phosphate niomineralization in vertebrates?**
Chemistry of Materials 2008, **20:**3549-3550.

27. Sharma V, Srinivasan A, Roychoudhury A, Rani K, Tyagi M, Dev K,
Nikolajeff F, Kumar S: **Characterization of protein extracts from different
types of human teeth and insight in biomineralization.** *Scientific Reports*
2019, **9:**9314.

28. Kamel BS, Voolstra CR, Medina M: **BioMine-DB: A database for metazoan
biomineralization proteins.** *PeerJ Preprints* 2016, **4:**e1983v1982.

29. Fukuda I, Ooki S, Fujita T, Murayama E, Nagasawa H, Isa Y, Watanabe T:
**Molecular cloning of a cDNA encoding a soluble protein in the coral
exoskeleton.** *Biochem Biophys Res Commun* 2003, **304:**11-17.

30. Puverel S, Tambutté E, Pereira-Mouries L, Zoccola D, Allemand D, Tambutté
S: **Soluble organic matrix of two Scleractinian corals: Partial and
comparative analysis.** *Comparative Biochemistry and Physiology Part B*
2005, **141:**480-487.

31. Drake JL, Mass T, Haramaty L, Zelzion E, Bhattacharya D, Falkowski PG:
**Proteomic analysis of skeletal organic matrix from the stony coral
Stylophora pistillata.** *Proceedings of the National Academy of Sciences* 2013,
110:3788-3793.

32. Ramos-Silva P, Kaandorp J, Huisman L, Marie B, Zanella-Cleon I, Guichard
N, Miller DJ, Marin F: **The skeletal proteome of the coral** *Acropora
millepora***: The evolution of calcification by cooption and domain
shuffling.** *Mol Biol Evol* 2013.

33. Takeuchi T, Yamada L, Shinzato C, Sawada H, Satoh N: **Stepwise evolution
of coral biomineralization revealed with genome-wide proteomics and
transcriptomics.** *PLOS ONE* 2016, **11:**e0156424.

34. Mass T, Drake Jeana L, Haramaty L, Kim JD, Zelzion E, Bhattacharya D,
Falkowski Paul G: **Cloning and characterization of four novel coral acid-
rich proteins that precipitate carbonates in vitro.** *Curr Biol* 2013, **23:**1126-
1131.

35. Puverel S, Tambutté E, Zoccola D, Dumart-Coulon I, Bouchot A, Lotto S,
Allemand D, Tambutté S: **Antibodies against the organic matrix in
scleractinians: A new tool to study coral biomineralization.** *Coral Reefs*
2005, **24:**149-156.

36. Clode PL, Marshall AT: **Calcium associated with a fibrillar organic matrix
in the scleractinian coral** *Galaxea fascicularis***.** *Protoplasma* 2003, **220:**153-
161.

37. Watanabe T: **Molecular analyses of protein components of the organic**
matrix in the exoskeleton of two scleractinian coral species. *Comp Biochem*
Phys B 2003, **136:**767 - 774.
38. Scheltema RA, Hauschild J-P, Lange O, Hornburg D, Denisov E, Damoc E,
Kuehn A, Makarov A, Mann M: **The Q Exactive HF, a Benchtop mass**
spectrometer with a pre-filter, high-performance quadrupole and an
ultra-high-field Orbitrap analyzer. *Molecular & Cellular Proteomics* 2014,
13:3698-3708.
39. Drake JL, Mass T, Falkowski PG: **The evolution and future of carbonate**
precipitation in marine invertebrates: Witnessing extinction or
documenting resilience in the Anthropocene? *Elementa Science of the*
Anthropocene 2014, **2:**000026.
40. Drake JL, Mass T, Haramaty L, Zelzion E, Bhattacharya D, Falkowski PG:
Reply to Ramos-Silva et al.: Regarding coral skeletal proteome.
Proceedings of the National Academy of Sciences 2013, **110:**E2147-E2148.
41. Ramos-Silva P, Marin F, Kaandorp J, Marie B: **LETTER: Biomineralization**
toolkit: the importance of sample cleaning prior to the characterization of
biomineral proteomes. Proceedings of the National Academy of Sciences;
2013.
42. Goffredo S, Vergni P, Reggi M, Caroselli E, Sparla F, Levy O, Dubinsky Z,
Falini G: **The skeletal organic matrix from Mediterranean coral**
***Balanophyllia europaea* influences calcium carbonate precipitation.** *PLoS*
ONE 2011, **6**.
43. Dauphin Y: **Comparative studies of skeletal soluble matrices from some**
scleractinian corals and molluscs. *International Journal of Biological*
Macromolecules 2001, **28:**293-304.
44. Weiner S: **Aspartic acid-rich proteins: Major components of the soluble**
organic matrix of mollusk shells. *Calcif Tissue Int* 1979, **29:**163-167.
45. Pereira-Mouriès L, Almeida MJ, Ribeiro C, Peduzzi J, Barthélemy M, Milet
C, Lopez E: **Soluble silk-like organic matrix in the nacreous layer of the**
bivalve *Pinctada maxima*: A new insight in the biomineralization field. *Eur*
J Biochem 2002, **269:**4994-5003.
46. Stoll HM, Ruiz Encinar J, Ignacio Garcia Alonso J, Rosenthal Y, Probert I,
Klaas C: **A first look at paleotemperature prospects from Mg in coccolith**
carbonate: Cleaning techniques and culture measurements. *Geochemistry,*
Geophysics, Geosystems 2001, **2**.
47. Nothdurft LD, Webb GE: **Earliest diagenesis in scleractinian coral**
skeletons: Implications for palaeoclimate-sensitive geochemical archives.
Facies 2009, **55:**161-201.
48. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S,
Kundu DJ, Inuganti A, Griss J, Mayer G, Eisenacher M: **The PRIDE**
database and related tools and resources in 2019: Improving support for
quantification data. *Nucleic Acids Res* 2019, **47:**D442-D450.
49. Voolstra CR, Li Y, Liew YJ, Baumgarten S, Zoccola D, Flot J-F, Tambutté S,
Allemand D, Aranda M: **Comparative analysis of the genomes of**
***Stylophora pistillata* and *Acropora digitifera* provides evidence for**
extensive differences between species of corals. *Scientific Reports* 2017,
7:17583.

50. Bertucci A, Tambutté S, Supuran C, Allemand D, Zoccola D: **A new coral**
carbonic anhydrase in *Stylophora pistillata*. *Mar Biotechnol* 2011, **13:**992-
1002.
51. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP,
Dolinski K, Dwight SS, Eppig JT: **Gene Ontology: Tool for the unification**
of biology. *Nat Genet* 2000, **25:**25-29.
52. Consortium GO: **Expansion of the gene ontology knowledgebase and**
resources. *Nucleic Acids Res* 2017, **45:**D331-D338.
53. Bryant DM, Johnson K, DiTommaso T, Tickle T, Couger MB, Payzin-Dogru
D, Lee TJ, Leigh ND, Kuo T-H, Davis FG: **A tissue-mapped axolotl de novo**
transcriptome enables identification of limb regeneration factors. *Cell*
Reports 2017, **18:**762-776.
54. Krogh A, Larsson B, Heijne Gv, Sonnhammer EL: **Predicting**
transmembrane protein topology with a hidden Markov model:
Application to complete genomes. *J Mol Biol* 2001, **305:**567-580.
55. Armenteros JJA, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak
S, von Heijne G, Nielsen H: **SignalP 5.0 improves signal peptide**
predictions using deep neural networks. *Nat Biotechnol* 2019, **37:**420.
56. Nielsen H, Engelbrecht J, Brunak S, von Heijne G: **Identification of**
prokaryotic and eukaryotic signal peptides and prediction of their
cleavage sites. *Protein Engineering* 1997, **10:**1-6.
57. Poisson G, Chauve C, Chen X, Bergeron A: **FragAnchor: A large-scale**
predictor of glycosylphosphatidylinositol anchors in eukaryote protein
sequences by qualitative scoring. *Genomics, Proteomics & Bioinformatics*
2007, **5:**121-130.
58. Shinzato C, Shoguchi E, Kawashima T, Hamada M, Hisata K, Tanaka M,
Fujie M, Fujiwara M, Koyanagi R, Ikuta T, et al: **Using the** *Acropora*
digitifera **genome to understand coral responses to environmental change.**
Nature 2011, **476:**320-323.
59. Emms DM, Kelly S: **OrthoFinder: Solving fundamental biases in whole**
genome comparisons dramatically improves orthogroup inference
accuracy. *Genome Biology* 2015, **16:**157.
60. Emms DM, Kelly S: **OrthoFinder: Phylogenetic orthology inference for**
comparative genomics. *Genome Biology* 2019, **20:**1-14.
61. Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ,
Robles M, Talón M, Dopazo J, Conesa A: **High-throughput functional**
annotation and data mining with the Blast2GO suite. *Nucleic Acids Res*
2008, **36:**3420-3435.
62. Mass T, Drake JL, Heddleston JM, Falkowski PG: **Nanoscale visualization of**
biomineral formation in coral proto-polyps. *Curr Biol* 2017, **27:**3191-3196.
e3193.
63. Thongboonkerd V, Mcleish KR, Arthur JM, Klein JB: **Proteomic analysis of**
normal human urinary proteins isolated by acetone precipitation or
ultracentrifugation. *Kidney International* 2002, **62:**1461-1469.
64. Fic E, Kedracka‐Krok S, Jankowska U, Pirog A, Dziedzicka‐Wasylewska M:
Comparison of protein precipitation methods for various rat brain
structures prior to proteomic analysis. *Electrophoresis* 2010, **31:**3573-3579.
65. Fields GB, Alonso DO, Stigter D, Dill KA: **Theory for the aggregation of**
proteins and copolymers. *The Journal of Physical Chemistry* 1992, **96:**3974-
3981.

66. Bern MW, Kil YJ: **Two-dimensional target decoy strategy for shotgun**
proteomics. *Journal of Proteome Research* 2011, **10:**5296-5301.
67. Fondrie WE, Noble WS: **A machine learning strategy that leverages large**
datasets to boost statistical power in small-scale experiments. *BioRxiv*
2019**:**849331.
68. Sykes GA, Collins MJ, Walton DI: **The significance of a geochemically**
isolated intracrystalline organic fraction within biominerals. *Org Geochem*
1995, **23:**1059-1065.
69. Goldberg WM: **Acid polysaccharides in the skeletal matrix and**
calicoblastic epithelium of the stony coral *Mycetophyllia reesi*. *Tissue and*
Cell 2001, **33:**376-387.
70. Cuif J-P, Dauphin Y, Doucet J, Salome M, Susini J: **XANES mapping of**
organic sulfate in three scleractinian coral skeletons. *Geochim Cosmochim*
Acta 2003, **67:**75-83.
71. Mann K, Wilt FH, Poustka AJ: **Proteomic analysis of sea urchin**
(*Strongylocentrotus purpuratus***) spicule matrix.** *Proteome Science* 2010,
8:33.
72. Von Heijne G, Abrahmsèn L: **Species-specific variation in signal peptide**
design Implications for protein secretion in foreign hosts. *FEBS Lett* 1989,
244:439-446.
73. Rabouille C: **Pathways of unconventional protein secretion.** *Trends Cell*
Biol 2017, **27:**230-240.
74. Rabouille C, Malhotra V, Nickel W: **Diversity in unconventional protein**
secretion. *J Cell Sci* 2012, **125:**5251-5255.
75. McGrath JP, Varshavsky A: **The yeast STE6 gene encodes a homologue of**
the mammalian multidrug resistance P-glycoprotein. *Nature* 1989,
340:400-404.
76. Nickel W, Rabouille C: **Mechanisms of regulated unconventional protein**
secretion. *Nature Reviews Molecular Cell Biology* 2009, **10:**148-155.
77. Colombo M, Raposo G, Théry C: **Biogenesis, secretion, and intercellular**
interactions of exosomes and other extracellular vesicles. *Annu Rev Cell*
Dev Biol 2014, **30:**255-289.
78. Whitham M, Parker BL, Friedrichsen M, Hingst JR, Hjorth M, Hughes WE,
Egan CL, Cron L, Watt KI, Kuchel RP: **Extracellular vesicles provide a**
means for tissue crosstalk during exercise. *Cell metabolism* 2018, **27:**237-
251. e234.
79. Clode PL, Marshall AT: **Kalisomes in corals: A novel KCl concentrating**
organelle? *Tissue and Cell* 2002, **34:**199-209.
80. Marshall AT, Clode PL, Russell R, Prince K, Stern R: **Electron and ion**
microprobe analysis of calcium distribution and transport in coral tissues.
J Exp Biol 2007, **210:**2453-2463.
81. Mass T, Giuffre AJ, Sun C-Y, Stifler CA, Frazier MJ, Neder M, Tamura N,
Stan CV, Marcus MA, Gilbert PUPA: **Amorphous calcium carbonate**
particles form coral skeletons. *Proceedings of the National Academy of*
Sciences 2017, **114:**E7670-E7678.
82. Ganot P, Tambutté E, Caminiti-Segonds N, Toullec G, Allemand D, Tambutté
S: **Ubiquitous macropinocytosis in anthozoans.** *Elife* 2020, **9:**e50022.
83. Neder M, Laissue PP, Akiva A, Akkaynak D, Albéric M, Spaeker O, Politi Y,
Pinkas I, Mass T: **Mineral formation in the primary polyps of**
pocilloporoid corals. *Acta Biomater* 2019.

84.   Mass T, Drake JL, Peters EC, Jiang W, Falkowski PG: **Immunolocalization**
783   **of skeletal matrix proteins in tissue and mineral of the coral *Stylophora***
784   ***pistillata**. Proceedings of the National Academy of Sciences* 2014, **111:**12728-
785   12733.
786   85.   Takeuchi T, Plasseraud L, Ziegler-Devin I, Brosse N, Shinzato C, Satoh N,
787   Marin F: **Biochemical characterization of the skeletal matrix of the**
788   **massive coral, *Porites australiensis*–The saccharide moieties and their**
789   **localization.** *Journal of Structural Biology* 2018, **203:**219-229.
790   86.   Puverel S, Houlbreque F, Tambutté E, Zoccola D, Payan P, Caminiti N,
791   Tambutté S, Allemand D: **Evidence of low molecular weight components in**
792   **the organic matrix of the reef building coral, *Stylophora pistillata*.**
793   *Comparative Biochemistry and Physiology a-Molecular & Integrative*
794   *Physiology* 2007, **147:**850-856.
795   87.   Farre B, Cuif J-P, Dauphin Y: **Occurrence and diversity of lipids in modern**
796   **coral skeletons.** *Zoology* 2010, **113:**250-257.
797   88.   Naggi A, Torri G, Iacomini M, Colombo Castelli G, Reggi M, Fermani S,
798   Dubinsky Z, Goffredo S, Falini G: **Structure and function of stony coral**
799   **intraskeletal polysaccharides.** *ACS Omega* 2018, **3:**2895-2901.
800   89.   Carpentier SC, Panis B, Vertommen A, Swennen R, Sergeant K, Renaut J,
801   Laukens K, Witters E, Samyn B, Devreese B: **Proteome analysis of**
802   **non-model plants: A challenging but powerful approach.** *Mass*
803   *Spectrometry Reviews* 2008, **27:**354-377.
804   90.   Dowd WW: **Challenges for biological interpretation of environmental**
805   **proteomics data in non-model organisms.** *Integr Comp Biol* 2012, **52:**705-
806   720.
807   91.   Forné I, Abian J, Cerdà J: **Fish proteome analysis: Model organisms and**
808   **non-sequenced species.** *Proteomics* 2010, **10:**858-872.
809   92.   Butler WT, Brunn JC, Qin C: **Dentin extracellular matrix (ECM) proteins:**
810   **comparison to bone ECM and contribution to dynamics of**
811   **dentinogenesis.** *Connect Tissue Res* 2003, **44:**171-178.
812   93.   Qin C, Baba O, Butler WT: **Post-translational modifications of SIBLING**
813   **proteins and their roles in osteogenesis and dentinogenesis.** *Critical*
814   *Reviews in Oral Biology and Medicine* 2004, **15:**126-136.
815   94.   Sarashina I, Endo K: **Skeletal matrix proteins of invertebrate animals:**
816   **Comparative analysis of their amino acid sequences.** *Paleontological*
817   *Research* 2006, **10:**311-336.
818

819   **Table and Figure Legends**
820
821   *Tables*

| | Stylophora pistillata genome hit | | | | Solubility Fraction | |
gi Number	Accession Number	Annotation	Best score	Go Categorization	ASM	AIM
1263115389	PFX12726.1	Retrovirus-related Pol polyprotein from transposon 17.6 [Stylophora pistillata]	340.80	-		x
1263115517	PFX12813.1	hypothetical protein AWC38_SpisGene23165 [Stylophora pistillata]	337.40	-		x
1263116757	PFX13778.1	Sacsin [Stylophora pistillata]	267.80	-		x
1263117261	PFX14205.1	Proto-oncogene tyrosine-protein kinase receptor Ret, partial [Stylophora pistillata]	336.70	a,c		x
1263119001	PFX15740.1	Protein FAM208A [Stylophora pistillata]	373.20	-	x	x
1263119725	PFX16398.1	hypothetical protein AWC38_SpisGene19330 [Stylophora pistillata]	273.10	-		x
1263122270	PFX18785.1	Mucin-4 [Stylophora pistillata]	449.50	a,c		x
1263130352	PFX26597.1	Complement C3 [Stylophora pistillata]	283.10	a,b,c		x
1263130510	PFX26751.1	Transmembrane protease serine 9 [Stylophora pistillata]	244.20	a		x
1263131615	PFX27832.1	Poly [ADP-ribose] polymerase 11 [Stylophora pistillata]	320.10	c		x
1263134664	PFX30831.1	hypothetical protein AWC38_SpisGene4366 [Stylophora pistillata]	322.20	-		x
1263134737	PFX30903.1	hypothetical protein AWC38_SpisGene4292 [Stylophora pistillata]	548.90	-		x
1263135656	PFX31810.1	Nidogen-2 [Stylophora pistillata]	286.20	a,c		x
1270028962	XP_022783323.1	phosphopantothenoylcysteine decarboxylase subunit VHS3-like [Stylophora pistillata]	406.60	c		x
1270029544	XP_022786582.1	synapsin-2-like isoform X2 [Stylophora pistillata]	320.50	c	x	x
1270031975	XP_022799089.1	CUB domain-containing protein-like isoform X2 [Stylophora pistillata]	699.70	-		x
1270032917	XP_022804012.1	EGF and laminin G domain-containing protein-like [Stylophora pistillata]	384.10	a		x
1270036141	XP_022779720.1	vitellogenin-like [Stylophora pistillata]	416.00	b,c		x
1270037242	XP_022780303.1	LOW QUALITY PROTEIN: uncharacterized protein LOC111321626 [Stylophora pistillata]	353.60	a,b,c	x	
1270037961	XP_022780690.1	skeletal aspartic acid-rich protein 2-like [Stylophora pistillata]	723.60	a		x
1270037969	XP_022780694.1	CUB and peptidase domain-containing protein 2-like [Stylophora pistillata]	285.80	a	x	x
1270041141	XP_022782398.1	skeletal aspartic acid-rich protein 1-like [Stylophora pistillata]	709.80	-	x	x
1270042352	XP_022783044.1	uncharacterized protein LOC111323869 [Stylophora pistillata]	283.60	-		x
1270043038	XP_022783415.1	coadhesin-like isoform X3 [Stylophora pistillata]	411.20	a		x
1270044042	XP_022783952.1	collagenase 3-like [Stylophora pistillata]	262.90	a,c		x
1270045301	XP_022784623.1	cation channel sperm-associated protein subunit beta-like [Stylophora pistillata]	328.40	a,c		x
1270049598	XP_022786918.1	major yolk protein-like isoform X2 [Stylophora pistillata]	476.50	a,c		x
1270052030	XP_022788227.1	hephaestin-like protein [Stylophora pistillata]	539.80	a,c		x
1270052971	XP_022788730.1	chymotrypsin-like elastase family member 1 [Stylophora pistillata]	264.80	a,c		x
1270054595	XP_022789591.1	endothelin-converting enzyme 1-like isoform X2 [Stylophora pistillata]	94.10	a,b,c		x
1270055217	XP_022789932.1	MAGUK p55 subfamily member 7-like [Stylophora pistillata]	313.70	a		x
1270056169	XP_022790441.1	PHD finger protein 21A-like [Stylophora pistillata]	277.90	c		x
1270059479	XP_022792212.1	ras-like protein 3 [Stylophora pistillata]	289.90	a,c		x
1270063049	XP_022794122.1	galaxin-like isoform X2 [Stylophora pistillata]	309.40	-		x
1270063475	XP_022794351.1	mammalian ependymin-related protein 1-like [Stylophora pistillata]	324.80	a,c		x
1270064196	XP_022794736.1	MAM and LDL-receptor class A domain-containing protein 2-like [Stylophora pistillata]	472.20	-		x
1270068394	XP_022796981.1	uncharacterized skeletal organic matrix protein 8-like [Stylophora pistillata]	664.40	-		x
1270068396	XP_022796982.1	uncharacterized protein LOC111335364 [Stylophora pistillata]	609.30	-		x
1270071953	XP_022798902.1	low-density lipoprotein receptor-related protein 8-like [Stylophora pistillata]	257.10	c		x
1270073139	XP_022799541.1	uncharacterized protein LOC111337489 [Stylophora pistillata]	461.10	b		x
1270080553	XP_022803524.1	digestive cysteine proteinase 1-like [Stylophora pistillata]	262.20	a,c		x
1270081088	XP_022803808.1	deleted in malignant brain tumors 1 protein-like [Stylophora pistillata]	344.70	c		x
1270081207	XP_022803872.1	spore wall protein 2-like isoform X3 [Stylophora pistillata]	338.60	-		x
1270081241	XP_022803894.1	uncharacterized protein LOC111341206 [Stylophora pistillata]	390.70	b		x
1270082891	XP_022804785.1	thioredoxin reductase 1, cytoplasmic-like [Stylophora pistillata]	337.00	c		x
1270084203	XP_022805470.1	uncharacterized protein LOC111342641 [Stylophora pistillata]	256.10	-		x
1270085816	XP_022806326.1	ZP domain-containing protein-like [Stylophora pistillata]	393.90	a		x
1270086467	XP_022806664.1	protein lingerer-like [Stylophora pistillata]	250.10	b,c		x
1270086954	XP_022806928.1	SLIT-ROBO Rho GTPase-activating protein 1-like [Stylophora pistillata]	309.10	-		x
1270087345	XP_022807143.1	condensin-2 complex subunit D3-like [Stylophora pistillata]	278.10	c		x
1270087556	XP_022807256.1	uncharacterized protein LOC111344300 [Stylophora pistillata]	280.40	a,c		x
1270088573	XP_022807807.1	uncharacterized protein LOC111344812 [Stylophora pistillata]	365.30	a,b,c		x
1270089244	XP_022808163.1	uncharacterized protein LOC111345150 [Stylophora pistillata]	310.20	-		x
1270090022	XP_022808576.1	uncharacterized protein LOC111345553 isoform X2 [Stylophora pistillata]	320.10	a,c		x
1270091315	XP_022809269.1	microtubule-associated tumor suppressor 1 homolog isoform X1 [Stylophora pistillata]	259.20	a,c		x
1270091317	XP_022809270.1	microtubule-associated tumor suppressor 1 homolog isoform X2 [Stylophora pistillata]	256.40	-		x
1270093788	XP_022810585.1	von Willebrand factor D and EGF domain-containing protein-like, partial [Stylophora pistillata]	298.80	-		x
1270095516	XP_022778254.1	uncharacterized protein LOC111319781 [Stylophora pistillata]	250.70	c		x
1270095572	XP_022778283.1	uncharacterized protein LOC111319816, partial [Stylophora pistillata]	508.10	-	x	x
190710633	ACE95141.1	carbonic anhydrase [Stylophora pistillata]	495.20	a,c	x	x

822
823 Table 1: 60 coral skeletal proteins detected by LC-MS/MS across all treatments and
824 solubility fractions. Proteins are listed in order of accession number. Geno ontology
825 categorization is represented as [a]ECM/transmembrane and protein modification,
826 [b]membrane processing, and [c]vesicle/secretion and metal binding.
827
828

NCBI Accession Number	Annotation	gi (NCBI)	Gene model id	RNA id	*Acropora digitifera* skeletal proteins (Takeuchi et al. 2016)	*Acropora millepora* skeletal proteins (Ramos-Silva et al. 2013)	*Stylophora pistillata* skeletal proteins (Drake et al. 2013)
ACE95141.1[a,c]	carbonic anhydrase	190710633	NA	NA			SeriatoporaSp_g7349; mapped to ACE95141.1 (EU532164)
PFX18785.1[a,c]	Mucin-4	1263122270	NA	NA	aug_v2a.09809.t1	JR987773	AGG36340.1
PFX30903.1	hypothetical protein AWC38_SpisGene4292	1263134737	NA	NA	aug_v2a.06327.t1	JR972076.1	AGG36350.1
XP_022779720.1[b,c]	vitellogenin-like	1270036141	gene3329	rna4552	aug_v2a.08856.t1		AGG36347.1, AGG36349.1
XP_022780690.1[a]	skeletal aspartic acid-rich protein 2-like	1270037961	gene4012	rna5498	aug_v2a.11068.t1, JT001945.1, aug_v2a.01441.t1, aug_v2a.01440.t1, JT004498.1, JR991407.1	JR991407.1, JT004498.1	AGG36358.1, AGG36357.1
XP_022780694.1[a]	CUB and peptidase domain-containing protein 2-like	1270037969	gene3981	rna5452		JR970990.1	
XP_022782398.1	skeletal aspartic acid-rich protein 1-like	1270041141	gene5242	rna7162		JT001945.1	AGG36357.1
XP_022783415.1	coadhesin-like isoform X3	1270043038	gene6099	rna8310	aug_v2a.05945.t1	JT016638.1	AGG36341.1, AGG36335.1, AGE45658.1, AGG36343.1
XP_022786918.1[a,c]	major yolk protein-like isoform X2	1270049598	gene8661	rna11721			AGG36338.1
XP_022788227.1[a,c]	hephaestin-like protein	1270052030	gene9579	rna12971	aug_v2a.24015.t1	JT019463.1	
XP_022794122.1	galaxin-like isoform X2	1270063049	gene13931	rna18823	aug_v2a.18631.t1	JR976690.1	
XP_022794736.1	MAM and LDL-receptor class A domain-containing protein 2-like	1270064196	gene14425	rna19450	aug_v2a.09968.t1, aug_v2a.09969.t1	JT011118.1, JR994474.1	AGG36355.1, AGG36356.1, AGG36345.1
XP_022796981.1	uncharacterized skeletal organic matrix protein 8-like	1270068394	gene16153	rna21699		JT014391.1	AGC24391.1
XP_022796982.1	uncharacterized protein LOC111335364	1270068396	gene16156	rna21701			AGC24391.1
XP_022804012.1[a]	EGF and laminin G domain-containing protein-like	1270032917	gene2104	rna2853	aug_v2a.15580.t1, aug_v2a.06123.t1, JR980881.1, aug_v2a.24512.t1, aug_v2a.06122.t1	JR980881.1	AGG36344.1, AGG36352.1
XP_022806326.1[a]	ZP domain-containing protein-like	1270085816	gene23405	rna30870	aug_v2a.07627.t1	JN631095.1	AGG36346.1

829
830 Table 2. Orthologous coral skeletal proteins in the present work and previously
831 published. Gene ontology categorization is represented as [a]ECM/transmembrane and
832 protein modification, [b]membrane processing, and [c]vesicle/secretion and metal
833 binding.
834
835 *Figures*

836
837 Figure 1. Workflow of methods for skeletal protein purification and concentration
838 after extraction in acetic acid. CF – Centrifugation filtration methods; ACT – acetone
839 precipitation methods; SSOM –soluble skeletal organic matrix; ISOM –insoluble
840 skeletal organic matrix.

841
Figure 2: Distribution of proteins numbers according of the purification methods.
Distribution of all proteins (SOM and ISOM combined) by purification methods (A).
SOM and ISOM distribution by acetone precipitation methods (ACT; B). SOM and
ISOM distribution by centrifugation and ultrafiltration method (CF; C). Note that CF
methods yielded more proteins in total. ISOM Insoluble Skeletal Organic Matrix;
SOM Skeletal Organic Matrix.
848

849
850 Figure 3. Suggested cellular and extracellular locations of *S. pistillata* skeletal protein
851 based on GO terms. Large yellow and red circles represent calicoblastic cells with
852 red cell membrane facing the skeleton. Small green circles represent vesicles. And

853   purple squiggles represent transmembrane proteins that may or may not span the
854   width of the calicoblastic space plus ECM proteins.
855