Multiple approaches for massively parallel sequencing of HCoV-19 genomes directly from clinical samples

Minfeng Xiao[1,7]*, Xiaoqing Liu[2]*, Jingkai Ji[3,1,7]*, Min Li[4,1,7]*, Jiandong Li[4,1,7]*, Lin Yang[5]*, Wanying Sun[4,1,7], Peidi Ren[1,7], Guifang Yang[5], Jincun Zhao[2,8], Tianzhu Liang[1,7], Huahui Ren[1], Tian Chen[5], Huanzi Zhong[1], Wenchen Song[1,7], Yanqun Wang[2], Ziqing Deng[1,7], Yanping Zhao[1,7], Zhihua Ou[1,7], Daxi Wang[1,7], Jielun Cai[1], Xinyi Cheng[1,7,12], Taiqing Feng[5], Honglong Wu[6], Yanping Gong[6], Huanming Yang[1,9], Jian Wang[1,9], Xun Xu[1,10], Shida Zhu[1,11], Fang Chen[5,1], Yanyan Zhang[5]#, Weijun Chen[6,4]#, Yimin Li[2]#, Junhua Li[1,7]#

[1]BGI-Shenzhen, Shenzhen, 518083, China.
[2]State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Health, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China.
[3]School of Future Technology, University of Chinese Academy of Sciences, Beijing 101408, China.
[4]BGI Education Center, University of Chinese Academy of Sciences, Shenzhen, 518083, China.
[5]MGI, BGI-Shenzhen, Shenzhen, 518083, China.
[6]BGI PathoGenesis Pharmaceutical Technology, Shenzhen, China.
[7]Shenzhen Key Laboratory of Unknown Pathogen Identification, BGI-Shenzhen, Shenzhen, 518083, China.
[8]Institute of Infectious disease, Guangzhou Eighth People's Hospital of Guangzhou Medical University, Guangzhou, China.
[9]James D. Watson Institute of Genome Science, Hangzhou, 310008, China.
[10]Guangdong Provincial Key Laboratory of Genome Read and Write, BGI-Shenzhen, Shenzhen, 518120, China.
[11]Shenzhen Engineering Laboratory for Innovative Molecular Diagnostics, BGI-Shenzhen, Shenzhen, 518120, China.
[12]School of Biology and Biological Engineering, South China University of Technology, Guangzhou, China

Correspondence should be addressed to J.L. (lijunhua@genomics.cn), Y.L. (dryiminli@vip.163.com), W.C. (chenwj@genomics.com), and Y.Z. (zhangyanyan@genomics.cn).

*These authors contributed equally to this work.

#These authors jointly supervised this work.

ABSTRACT

COVID-19 has caused a major epidemic worldwide, however, much is yet to be known about the epidemiology and evolution of the virus. One reason is that the challenges underneath sequencing HCoV-19 directly from clinical samples have not been completely tackled. Here we illustrate the application of amplicon and hybrid capture (capture)-based sequencing, as well as ultra-high-throughput metatranscriptomic (meta) sequencing in retrieving complete genomes, inter-individual and intra-individual variations of HCoV-19 from clinical samples covering a range of sample types and viral load. We also examine and compare the bias, sensitivity, accuracy, and other characteristics of these approaches in a comprehensive manner. This is, to date, the first work systematically implements amplicon and capture approaches in sequencing HCoV-19, as well as the first comparative study across methods. Our work offers practical solutions for genome sequencing and analyses of HCoV-19 and other emerging viruses.

INTRODUCTION

As of 14 March 2020, human coronavirus 2019 (HCoV-19) has surpassed severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus (MERS-CoV) in every aspect, infecting over 140,000 people in more than 110 countries, with a mortality of over 5,000[1,2]. So far, coronaviruses have caused three major epidemics in the past two decades, posing a great challenge to global health and economy. Massively parallel sequencing (MPS) of viral genomes has demonstrated enormous capacity as a powerful tool to study emerging infectious diseases, such as SARS, MERS, Zika, and Ebola, in tracing the outbreak origin and drivers, tracking transmission chains, mapping the spread, and monitoring the evolution of the etiological agents[3-8]. Though by 14 March 2020, fewer than 500 HCoV-19 genomes were published on public databases including China National GeneBank DataBase (CNGBdb), NCBI GenBank, the Global initiative on sharing all influenza data (GISAID), etc, and much remains unknown about the epidemiology and evolution of the virus. One possible explanation of the paucity of published HCoV-19 genomes was the challenges posed by sequencing clinical samples with low virus abundance.

The first teams obtained the HCoV-19 genome sequences through metatranscriptomic MPS, supplemented by PCR and Sanger sequencing of a combination of bronchoalveolar-lavage fluid (BALF) and culture[9-11] or from BALF directly[12,13]. Experience from studying SARS-CoV showed that BALF from the lower respiratory tract was an ideal sample type with higher viral load[14]. However, BALF was not routinely collected from every patient, and human airway epithelial (HAE) cell culture is very labor-intensive and time-consuming, taking four to six weeks[10,15]. The University of Hong Kong team managed to get the whole-genome sequences through metatranscriptomic sequencing with Oxford Nanopore platform supplemented by Sanger sequencing from both nasopharyngeal and sputum specimens after single-primer amplification[16]. The United States scientists published the whole-genome sequence using oropharyngeal and nasopharyngeal specimens through Sanger and metatranscriptomic sequencing with both Illumina and MinIon[17]. To date, multiplex PCR-based or hybrid capture-based whole-genome sequencing of HCoV-19, as well as comparative studies between different approaches, have not been reported on peer-reviewed journals.

Besides inter-individual variations, dissecting intra-individual dynamics of viruses also largely promotes our understanding of viral-host interactions, viral evolution and transmission as demonstrated for Ebola, Zika, Influenza, etc[6,18-20]. The analyses of intra-individual single nucleotide variations (iSNVs) and its allele frequency have also contributed to antiviral therapy and drug resistance, e.g., to reveal highly conserved genes during the outbreak that potentially serve as ideal therapeutic targets[19,21]. However, it is a challenge to accurately detect iSNVs from clinical samples, especially when the samples are subjected to extra steps of enrichment and amplification.

Therefore, we aim to comprehensively compare the sensitivity, inter-individual (variant) and intra-individual (iSNV) accuracy, and other general features of different approaches by systematically utilizing ultra-high-throughput metatranscriptomic, hybrid capture-based, and amplicon-based sequencing approaches to obtain genomic information of HCoV-19 from serial dilutions of a cultured isolate and directly from clinical samples. We present a reasonable sequencing strategy that fits into different scenarios, and estimate the minimal amount of sequencing data necessary for downstream HCoV-19 genome analyses. Our study, together with our tailor-made experimental workflows and bioinformatics pipelines, offers very practical solutions to facilitate the studies of HCoV-19 and other emerging viruses in the future and would promote extensive genomic sequencing and analyses of HCoV-19 and other emerging viruses, underpinning more comprehensive real-time virus surveillance and more efficient viral outbreaks managing.

RESULTS

Design of the comparative study. We sampled eight specimens from COVID-19 patients in February 2020, including throat swab, nasal swab, anal swab and sputum specimens, and the corresponding cycle threshold (Ct) value of HCoV-19 qRT-PCR ranges from 18 to 32 (Table 1). We initially tried to boost the coverage and depth of the viral genome by ultra-deep metatranscriptomic sequencing with an average sequencing amount of 1,607,264,105 paired-end reads (Table 1). Although we managed to obtain complete viral genome assemblies for each specimen, the sequencing depth varied across specimens. Only 0.002%-0.003% of the total reads were assigned to the HCoV-19 in three samples (GZMU0014, GZMU0030 and GZMU0031) with Ct between 29-32, resulting in inferior sequencing depth (less than 100X) (Table 1). Isolating viruses and enriching them in cell culture might improve the situation, but this requires high-standard laboratory settings and expertise apart from being time-consuming. Also, unwanted mutations that are not concordant with original clinical samples may be introduced during the culturing process.

To enrich adequate viral content for whole-genome sequencing in a convenient manner, we pursued two other methods: multiplex PCR amplification (amplicon) and hybrid capture (capture) (Fig. 1). We designed a systematic study to comprehensively validate the bias, sensitivity, inter-individual (variant) and intra-individual (iSNV) accuracy of multiple approaches by sequencing serial dilutions of a cultured isolate (unpublished), as well as the eight clinical samples (Fig. 2). We performed qRT-PCR of 10-fold serial dilutions (D1-D7) of the cultured isolate, and the Ct was 17.3, 20.8, 24.5 for, 28.7, 31.8, 35, and 39.9, respectively, indicating the undiluted RNA (D0) of the cultured isolate contained ~1E+08 genome copies per mL. For amplicon sequencing, we designed two set of primers generating PCR products of 300-400 bp and 100-200 bp, respectively. The ~400 bp amplicon-based sequencing was implemented in all samples and analyzed throughout the study, while the ~200 bp amplicon-based sequencing was only applied in the cultured isolate for coverage analysis.

Comparison of evenness and sensitivity. Theoretically, amplicon sequencing should be the most sensitive and economical method among the three, and is particularly suitable in an outbreak where viral isolates are highly related. Although, there are still potential pitfalls, for instance, the 40 cycle-PCR in our workflow might augment trace amounts of HCoV-19 cross-contamination. To ensure the confidence of the datasets, we included serial dilutions of the cultured isolate and negative controls prepared from nuclease-free water and human

nucleic acids since the 1st PCR. All samples in ~400 amplicon-based sequencing exhibited
> 99.5% coverage across the HCoV-19 genome except for 1E+01 (95.23%), GZMU0031
(73.65%), HNA (6.17%), water (60.24%), suggesting the primers were well designed and
the positive datasets were reliable. We also set stringent and method-specific criteria to filter
low-confidence sequencing reads and samples (see Methods), e.g., clinical sample
GZMU0031 was excluded for downstream sensitivity and accuracy validation due to inade-
quate depth in amplicon sequencing (Fig. 3a). Another pitfall is that amplification across the
genome can hardly be unbiased, causing difficulties in complete genome assembly. Indeed,
amplicon sequencing exhibited a higher level of bias compared with metatranscriptomic se-
quencing, in terms of coverage across the viral genomes from the cultural isolate and the
clinical samples tested in our study (Fig. 3b, d, Supplementary Fig. 1). To our surprise, how-
ever, capture sequencing was almost as unbiased as meta sequencing, demonstrating bet-
ter performance than the previous capture method used to enrich ZIKV despite the HCoV-
19 genome is ~3 fold larger than ZIKV[22] (Fig. 3b, c). Two reasons amongst others were
likely to be accountable to this improvement, 1) we utilized 506 pieces of 120 ssDNA probes
covering 2x of the HCoV-19 genome to capture the libraries, 2) we employed the DNBSEQ
sequencing technology that features PCR-free rolling circle replication (RCR) of DNA Nano-
balls (DNBs)[23,24].

The sequencing results of amplicon and capture approaches revealed dramatic increases
in the ratio of HCoV-19 reads out of the total reads compared with meta sequencing, sug-
gesting the enrichment was highly efficient - 5596-fold in capture method and 5710-fold in
amplicon method for each sample on average (Supplementary Table 1-2). To further com-
pare the sensitivity of different methods, we plotted the number of HCoV-19 reads per million
(HCoV-19-RPM) of total sequencing reads against the viral concentration for each sample.
The productivity was similar between the two methods when the input RNA of the cultured
isolate contained 1E+05 genome copies per mL and above (Fig. 3e). However, amplicon
sequencing produced 10-100 fold more HCoV-19 reads than capture sequencing when the
input RNA concentration of the cultured isolate was 1E+04 genome copies per mL and
lower, suggesting amplicon-based enrichment was more efficient than capture for more
challenging samples (conc. ≤ 1E+04 genome copies per mL, or Ct ≥ 28.7) (Fig. 3e). Meta
sequencing - as expected - produced dramatically lower HCoV-19-RPM than the other two
methods among clinical samples tested with a wide range of Ct values, whereas amplicon
and capture were generally comparable to each other (Fig. 3e). Considering the costs for
sequencing, storage, and analysis increase substantially with larger datasets, we tried to

estimate how much sequencing data must be produced for each approach in order to
achieve 10X depth across 95% of the HCoV-19 genome, and the results can be found in
Supplementary Table 3. As a practical, cost-effective guidance for future sequencing, we
also assessed the minimum amount of data required to pass the stringent filters (≥ 95%
coverage and method-specific depth, see Methods) in our pipelines corresponding to differ-
ent viral loads. We estimated that for high-confidence downstream analyses, amplicon se-
quencing requires at least 2,757 to 186 Mega bases (Mb) for samples containing 1E+02 to
1E+06 copies of HCoV-19 genome per mL, while capture sequencing requires 24,474 to 9
Mb for the same situation (Fig. 3g, Supplementary Table 4).

Investigation of inter- and intra-individual variations. To determine the accuracy of dif-
ferent approaches in discovering inter-individual genetic diversity, we tested each method
in calling the single nucleotide variations (SNVs) and verified some of the SNVs with Sanger
sequencing. Two to five SNVs were identified within each clinical sample, and in all the
seven samples, SNVs identified by the three methods were concordant except that capture
missed one SNV at position 16535 in GZMU0014 (Fig. 4a). We then investigated the allele
frequencies of these sites across methods, and found that alleles identified by capture se-
quencing displayed lower frequencies than the other two methods, especially for
GZMU0014, GZMU0030, and GZMU0042 where the viral load was lower (Ct ≥ 29), which
explained why capture sequencing neglected an SNV in our pipeline when the cutoff of SNV
calling was set as 80% allele frequency (Fig. 4b). These data indicate that amplicon se-
quencing is more accurate than capture sequencing in identifying SNVs, especially for chal-
lenging samples.

To further determine the accuracy of different approaches in identifying HCoV-19 iSNVs, we
examined minor allele frequencies in serial dilutions of the cultured HCoV-19 isolate and
clinical samples. For serial dilutions of the cultured isolate, the minor allele frequencies de-
tected in capture sequencing datasets were generally approximate to meta sequencing,
while most allele frequencies in amplicon sequencing datasets deviated with those in meta
sequencing (Fig. 4c) A similar pattern was shown for clinical samples, indicating that am-
plicon sequencing was unreliable of quantifying minor allele frequencies (Fig. 4d). Plotting
allele frequencies against HCoV-19 concentrations supported the above finding, and further
revealed that amplicon sequencing was unreliable of allele frequencies at all concentrations
while capture sequencing was reliable at > 1E+03 genome copies per mL (Supplementary
Fig. 2). Referring to the iSNV identified in clinical samples by meta sequencing, we then
calculated the false positive rate (FPR) and false negative rate (FNR) of minor alleles called

by amplicon and capture methods. The FPR and FNR of minor alleles identified in amplicon sequencing was 0.74% and 66.67%, while that in capture sequencing was 0.02% and 0%, respectively. Together these results suggest amplicon sequencing was not as accurate as capture sequencing in identifying minor alleles, which could be in part due to Matthew effect caused by PCR.

Microbiome in clinical samples. In addition to target viral genome, metatranscriptomic sequencing has also allowed us to investigate RNA expression patterns of the overall microbiome and host content and thus suitable for discovering new viruses, distinguishing co-infections, and dissecting virus-host interactions. To explore the microbiota, we performed further metatranscriptomics analysis of the clinical samples. We were able to identify host nucleic acids in all of the samples, and over 95% of total reads were from the host in sputum, nasal, and throat samples (Supplementary Fig. 3a). Virus contributed to less than 5% of reads in anal swab and throat swab while more than 50% of reads in nasal swab (Supplementary Fig. 3b). These results suggest nasal swab could be the most ideal sample type for viral detection among the four sample types, which agrees with recent clinical evidence[25]. Among the viral reads, over 90% were Coronaviridae, which is consistent with clinical diagnostics (Supplementary Fig. 3c). Reads from other viruses were also identified, indicating further measurements could be taken to confirm if co-infection exists (Supplementary Fig. 3). Bacterial composition was also shown, providing support for scientific research, as well as for further confirmation of bacterial infection and antibiotics prescription (Supplementary Fig. 3d-f).

Guidance for virus sequencing. Taken together, each sequencing scheme elaborated here for massively parallel sequencing of HCoV-19 genomes has its own merits (Table 2). We hereby propose a reasonable, cost-effective strategy for sequencing and analyzing HCoV-19 under different situations: 1) if one wants to study other genetic materials than the target viruses, or the viruses become highly diversified via recombinational events, or the viral load within the RNA sample is high (e.g. conc. ≥ 1E+05 viral genome copies per mL, or Ct ≤ 24.5), meta sequencing can be prioritized; 2) if one focuses on intra-individual variations for more challenging samples (e.g. conc. < 1E+05 viral genome copies per mL, or Ct > 24.5), capture sequencing seems to be a justified choice; and, 3) if identifying SNVs is the main purpose, the most convenient, economical strategy would be amplicon sequencing that can support high-confidence analyses of samples containing viral content as low as 1E+02 viral genome copies per mL, or Ct as high as 35.

DISCUSSION

Sequencing low-titre viruses directly from clinical samples is challenging, which is further exacerbated by the fact that coronavirus genomes are the largest among RNA viruses (~3 fold larger compared with ZIKV). Compared with direct metatranscriptomic sequencing, high sensitivity of hybrid capture and amplicon sequencing methods come at a price of low accuracy, and neither of the two can be used to sequence highly diverse or recombinant viruses because the primers and probes are specific to known viral genomes. Amplicon sequencing compromises its accuracy, while it becomes the most convenient and economical method of all. Either or a combination of the approaches described here can be chosen to cope with various needs of researchers, e.g., metatranscriptomic sequencing data with insufficient coverage and depth can be pooled with hybrid capture data to generate high quality assemblies[22]. Our research, as well as the methods elaborated here, are able to help other researchers to sequence and analyze large viruses from clinical samples and thus benefit investigations on the genomic epidemiology of viruses.

Some pros and cons described above might be specific to the experimental workflows and bioinformatics pipelines tailored in this study, e.g., 1) the bias of amplicon sequencing can be improved by reducing the amount of cycles in the 1st PCR or optimize the molar ratios of primers (Fig. 1a), 2) the amplicon sequencing is particularly convenient compared with previous counterparts since the fragmentation and library construction steps are omitted here by integrating adaptor and barcode ligation in the 2nd PCR and sequencing the amplicons using single-end 400 nt reads (Fig. 1a), 3) using anything less that 506 pieces of 120 ssDNA probes in hybrid capture may attenuate the sequencing coverage while increase the bias, 4) metatranscriptomic sequencing was conducted with an ultra-high-throughput sequencing platform so that the successful rate was substantially higher than usual. 5) the minimal amount of data necessary for analyzing the HCoV-19 genome from clinical samples across methods is higher than that predicted by data from the cultured isolate was probably due to the high nucleic acids background from the host and other microbes (Supplementary Table 3-4, Supplementary Fig. 3). Also, we do not take into account the time spent in sequencing since the workflows can be easily adapted in order to be compatible with various platforms including Illumina and Oxford Nanopore Technologies (ONT), besides DNBSEQ of MGI.

METHODS

Ethics statement

270 The Institutional Review Boards (IRB) of the First Affiliated Hospital of Guangzhou Medical
271 University approved the clinical studies. IRB of BGI-Shenzhen approved the sequencing
272 and downstream studies.

Sampling, RNA extraction, reverse transcription and qRT-PCR

Clinical specimens (including throat swab, nasal swab, anal swab, and sputum) were ob-
tained from confirmed COVID-19 cases at the First Affiliated Hospital of Guangzhou Medical
University. Total RNA of the cultured isolate of HCoV-19 was obtained from the Academy of
Military Medical Science (AMMS), and subjected to 10-fold serial dilutions. Total RNA was
extracted with QiAamp RNeasy Mini Kit (Qiagen, Heiden, Germany) following the manufac-
turer's instructions without modification. Real-time reverse transcription PCR (qRT-PCR)
targeting RdRp gene and N gene of HCoV-19 was used to detect and quantify the viral RNA
within clinical samples and serial dilutions of the cultured isolate using the HCoV-19 Nucleic
Acid Detection Kit following the manufacture's protocol (Geneodx, Shanghai, China，and
BGI-Shenzhen, Shenzhen, China).

Multiplex PCR primers design

For multiplex PCR, Primer3 (v. 0.4.0)[26,27] and BLAST
(https://blast.ncbi.nlm.nih.gov/Blast.cgi) were used to design the primers targeting the
HCoV-19 whole genome, host genome fragments and lambda phage genome fragments
following a proprietary padlock technique (MGI, BGI-Shenzhen, China). Primers were 18 to
35 nucleotides (nt) long and the annealing temperature was between 58°C and 62°C. Adap-
tor sequences for barcoding were attached to the 5'-terminals of primers. 3'-terminals of
primers were designed to avoid hotspots of single nucleotide variants. The primers were
synthesized by GENEWIZ and China National GeneBank. Overlap regions between adja-
cent amplicons were 59-209 bp.

Metatranscriptomic library preparation and sequencing

295 Host DNA was removed from RNA samples using DNase Ⅰ, and the concentration of RNA
296 samples was measured by Qubit RNA HS Assay Kit (Thermo Fisher Scientific, Waltham,
297 MA, USA). DNA-depleted and purified RNA was used to construct the single-stranded cir-
298 cular DNA library with MGIEasy RNA Library preparation reagent set (MGI, Shenzhen,
299 China), as follows: 1) RNA was fragmented by incubating with fragmentation buffer at 87°C

for 6 min; 2) double-stranded (ds) cDNA was synthesized using random hexamers with frag-
mented RNA; 3) ds cDNA was subjected to end repair, adaptor ligation, and 18-cycle PCR
amplification; 4) PCR products were Unique Dual Indexed (UDI), before going through cir-
cularization, and rolling circle replication (RCR) to generate DNA nanoballs (DNBs)-based
libraries. DNBs preps of clinical samples were sequenced on the ultra-high-throughput DNB-
SEQ-T7 platform (MGI, Shenzhen, China) with paired-end 100 nt strategy, generating 321
Gb sequencing data for each sample on average.

Hybrid capture-based enrichment and sequencing

A hybrid capture technique was used to enrich HCoV-19-specific content from the meta-
transcriptomic double-stranded DNA libraries with the 2019-nCoVirus DNA/RNA Capture
Panel (BOKE, Jiangsu, China). Manufacturer's instructions were slightly modified to accom-
modate the MGISEQ-2000 platform, i.e., blocker oligos and PCR primer oligos were re-
placed by MGIEasy exon capture assistive kit (MGI, Shenzhen, China). DNBs-based librar-
ies were constructed and sequenced on the MGISEQ-2000 platform with paired-end 100 nt
strategy using the same protocol described above, generating 37 Gb sequencing data for
each sample on average.

Amplicon-based enrichment and sequencing

Total RNA was reverse transcribed to synthesize the first-strand cDNA with random
hexamers and Superscript II reverse transcriptase kit (Invitrogen, Carlsbad, USA).
Sequencing was attempted on all samples regardless of Ct value including negative controls
prepared from nuclease-free water and NA12878 human gDNA. A two-step HCoV-19
genome amplification was performed with an equimolar mixture of primers by 40 cycles of
using ATOPlex SARS-CoV-2 full-length genome panel (MGI, BGI-Shenzhen, China). 20 µl
of first-strand cDNA was mixed with the components of the first PCR reaction following the
manufacturer's instructions, including lambda phage gDNA unless specified. 2 ng of Human
gDNA was added to each PCR reaction of the cultured isolate. The PCR was performed as
follows: 5 min at 37°C, 10 min at 95°C, 15 cycles of (10 s at 95°C, 1min at 64°C, 1min at
60°C to 10 s at 72°C), 2 min at 72°C. The products were purified with MGI EasyDNA Clean
beads (MGI, BGI-Shenzhen, China) at a 5:4 ratio and cleaned with 80% concentration
ethanol according to the manufacturer's instructions. The 2nd PCR was performed under
the same regimen as the 1st PCR except for 25 cycles, and the beads-purified products
from the first PCR reaction were unique dual indexed. After the 2nd PCR, products were
purified following the same procedures as the 1st PCR and quantified using the Qubit

dsDNA High Sensitivity assay on Qubit 3.0 (Life Technologies). PCR products of samples

yielding sufficient material (> 5 ng/µl) were pooled at equimolar to a total DNA amount of

300 ng before converting to single-stranded circular DNA. DNBs-based libraries were

generated from 20 µl of single-stranded circular DNA pools and sequenced on the MGISEQ-

2000 platform with single-end 400 nt strategy, generating 1.8 Gb sequencing data for each

sample on average.

Identification of HCoV-19-like reads from Massively Parallel Sequencing data

For metatranscriptomic and hybrid capture sequencing data, total reads were first processed

by Kraken v0.10.5[28] (default parameters) with a self-build database of Coronaviridae ge-

nomes (including SARS, MERS and HCoV-19 genome sequences downloaded from

GISAID, NCBI and CNGB) to efficiently identify candidate viral reads with a loose manner.

These candidate reads were further qualified with fastp v0.19.5[29] (parameters: -q 20 -u 20 -

n 1 -l 50) and SOAPnuke v1.5.6[30] (parameters: -l 20 -q 0.2 -E 50 -n 0.02 -5 0 -Q 2 -G -d) to

remove low-quality reads, duplications and adaptor contaminations. Low-complexity reads

were next filtered by PRINSEQ v0.20.4[31] (parameters: -lc_method dust -lc_threshold 7).

After the above process, HCoV-19-like reads generated from metatranscriptomics and

hybrid capture method were obtained.

For amplicon sequencing data, SE 400 reads were processed with fastp v0.19.5[29] (param-

eters: -q 20 -u 20 -n 1 -l 50) to remove low quality-reads and adaptor. Primer sequences

were trimmed with soft-clipping SAM/BAM alignments. After aligning amplicon reads to

HCoV-19 reference genome (GISAID accession: EPI_ISL_402119) by BWA aln (v0.7.16)[32]

with default parameters, the alignment BAM files were marked with BAMClipper v1.1.1[33]

(Parameters: -n 4 -u 21 -d 21) according to BED files of the amplicon primers.

Assembling viral genome

HCoV-19-like reads of metatranscriptomic and hybrid capture sequencing data were *de*

novo assembled with SPAdes (v3.14.0)[34] using the default settings to obtain virus genome

sequences. To reduce the complexity of the assembling process, identified HCoV-19-like

reads of metatranscriptomic and hybrid capture sequencing data were subsampled to the

data amount greater than 100X sequencing depth for the HCoV-19 genome. For the two

metatranscriptomic samples with a sequencing depth lower than 100X for the HCoV-19

genome (GZMU0014 and GZMU0030), all HCoV-19-like reads were used for assembling

viral genomes.

Due to the uneven read coverage in amplicon sequencing of HCoV-19, virus consensus

sequences of amplicon samples were generated by Pilon v1.23[35] (parameters: --changes –

vcf --changes --vcf --mindepth 1 --fix all, amb). Positions with depth less than 100X or lower

five times than negative control samples were masked as ambiguous base N.

Assessment the coverage depth across the viral genome

HCoV-19-like reads of metatranscriptomic and hybrid capture sequencing data were aligned

to the HCoV-19 reference genome (GISAID accession: EPI_ISL_402119) with BWA aln

(v0.7.16)[32].   Duplications   were   identified   by   Picard   Markduplicates

(v2.10.10)(http://broadinstitute.github.io/picard) with default settings. For each sample, we

calculated the depth of coverage at each nucleotide position of the HCoV-19 reference

genome with Samtools (v1.9)[36] and scaled the values to the mean depth. For each

nucleotide position, we calculated the median depth, and 20th and 80th percentiles across

all samples. Read coverage and depth across the HCoV-19 reference genome were plotted

by a 200-nt sliding window with the ggplot2[37] package in R (v3.6.1)[38].

Amplicon sequencing data were processed as described above, except that duplications

were not removed. A heatmap was generated to visualize the viral genome coverage for all

samples sequenced by the amplicon method with the pheatmap[39] package in R (v3.6.1)[38].

The depth at each nucleotide position was binarized, and was shown in pink if the depth

was 100x and above.

Relationships between genome copies and method-dependent minimum amount of

sequencing data

HCoV-19 reads of metatranscriptomic and hybrid capture sequencing data were identified

by aligning the HcoV-19-like reads to the HCoV-19 reference genome (GISAID accession:

EPI_ISL_402119) with BWA in a strict manner of coverage ≥ 95% and identity ≥ 90%. For

comparisons of the coverage and depth of the viral genome across samples and methods,

we normalized the viral reads to total sequencing reads with HCoV-19 Reads Per Million

(HCoV-19-RPM). HCoV-19-RPM for amplicon sequencing data was calculated by the same

pipelines we applied for metatranscriptomic and hybrid capture sequencing data.

To estimate the minimum data requirements for genome assembling and intra-individual

variation analysis, we applied gradient-based sampling to the HCoV-19 genome alignments

(referred to BAM files) to each dataset using Samtools (v1.9)[36]. The effective genome

coverage was set as 95% for all three MPS methods. Considering the distinct

technologies used in different methods, we set method-dependent thresholds of effective
depth as follows: 1) ≥ 10X for metatranscriptomic sequencing; 2) ≥ 20X for hybrid capture
sequencing; and 3) ≥ 100X for amplicon sequencing. We next calculated the coverage and
depth within each subsampled BAM file per sample to determine the minimal BAM file that
could meet the above thresholds of both coverage and sequencing depth. The method-
dependent minimum amount of sequencing data of each sample were estimated accord-
ingly. We assessed the correlations between the HCoV-19 genome copies per mL in diluted
samples of cultured isolates and the minimum amount of sequencing data for amplicon- and
capture-based methods using Pearson correlation coefficient (R) with the function *scatter*
from the R package *ggpubr* (v3.6.1)[40].

Consistency in variants calling performance among methods

Except for amplicon sequencing samples, variants calling in metatranscriptomic and hybrid
capture sequencing samples was performed in the previous BAM files of identified HCoV-
19 reads after removing duplications from alignment output by Picard Markduplicates
(http://broadinstitute.github.io/picard). To accurately identify SNVs from viral sequencing
data of all three methods, we first called SNVs with freebayes (v1.3.1)[41] (parameters: -p 1 -
q 20 -m 60 --min-coverage 10 -V) and then filtered the low-confidence SNVs with snippy-
vcf_filter[42] (parameters: --minqual 100 --mincov 10 --minfrac 0.8). Remaining SNVs post
filtering in VCF files generated by freebayes were annotated in HCoV-19 genome assem-
blies and consensus sequences with SNVeff (v4.3)[43] using default parameters.

Next, we calculated SNV allele frequencies and called iSNVs (intra-host Single Nucleotide
Variations) for each dataset to assess the consistency of variants calling performance
among three methods. We performed pysamstats v1.1.2 (https://github.com/ali-
manfoo/pysamstats) (parameters: -type variation_strand --min-baseq 20 -D 1000000) to
count the number of matches, mismatches, deletions and insertions at each base, estimate
nucleotide percentage and determine allele frequencies of SNVs at reference genome po-
sitions based on the HCoV-19 alignments from BAM files.

For amplicon sequencing data, only base positions covered by ≥100X reads were used for
iSNVs calling. For metatranscriptomic and hybrid capture-based sequencing data, the
thresholds of depth were set as 10X and 20X, respectively. The candidate iSNVs were fur-
ther filtered for quality as follows: 1) frequency filtering, only minor alleles (frequency ≥ 5%
and <50%) and major alleles (frequency ≥ 50% and ≤ 95%) were remained; 2) depth filter-
ing, iSNVs with fewer than five forward or reverse reads were removed; and 3) strand bias

filtering (not applicable to single-end reads of amplicon sequencing), iSNVs were removed if there were more than a 10-fold strand bias or a 5-fold difference between the strand bias of the variant call and that of the reference call.

Taxonomy of clinical samples by unbiased metatranscriptomic sequencing

For metatranscriptomic sequencing of clinical samples, raw sequencing data of a single sequence lane (approximately 60-75 Gb per sample) was used to simultaneously assess the RNA expression patterns of human, bacteria and viruses in clinical samples from COVID-19 patients. We first used software fastp (v0.19.5)[29] to filter low-quality reads and remove adapter with parameters: -5 -3 -q 20 -c -l 30. After QC, we mapped high-quality reads to hg19 and removed human ribosomal RNA (rRNA) reads by SOAP2 v2.21[44] (parameters: -m 0 -x 1000 -s 28 -l 32 -v 5 -r 1), and the remaining RNA reads were then aligned to hg19 by HISAT2[45] with default settings to identify non-rRNA human transcripts as previously described. Next, we applied Kraken 2[46] (version 2.0.8-beta, parameters: --threads 24 --confidence 0) to assign microbial taxonomic ranks to non-human RNA reads against the large reference database MiniKraken2 (April 2019, 8GB) built from the Refseq bacteria, archaea, and viral libraries and the h38 human genome. Bracken[47]
 (Bayesian Reestimation of Abundance with Kraken) was further applied to estimate microbial relative abundances based on taxonomic ranks of reads assigned by Kraken2.

Data availability Sequencing data that support the findings of this study have been deposited in CNGB (https://db.cngb.org/) under Project accession CNP0000951 and CNP0000955, and in GISAID under accession EPI_ISL_414663, EPI_ISL_414686 to EPI_ISL_414692.

ACKNOWLEDGEMENTS

We attribute this work to the amazing people in this land who dedicate themselves to the battle of mankind against viruses. This work is funded by the State Key Research Development Program of China (2019YFC1200501), the National Major Project for Control and Prevention of Infectious Disease in China (2018ZX10301101-004), the emergency grants for prevention and control of SARS-CoV-2 of Ministry of Science and Technology (2020YFC0841400) and Guangdong province (2020B111108001, 2018B020207013),

Guangdong Provincial Key Laboratory of Genome Read and Write (No. 2017B030301011), Guangdong Provincial Academician Workstation of BGI Synthetic Genomics (No. 2017B090904014), and Shenzhen Engineering Laboratory for Innovative Molecular Diagnostics (DRC-SZ[2016]884). We thank China National GeneBank at Shenzhen for providing sequencing service. We appreciate all the authors who have deposited and shared genome data on GISAID.

AUTHOR CONTRIBUTIONS

J.L., W.C. and M.X. conceived the project. X.L., J.Z, Y.W., and Y.L. sampled and processed the clinical specimen. M.X., Ji.L., M.L., and J.L. designed the experiments. L.Y. and Y.Z. developed the multiplex PCR amplicon-based sequencing method. M.L., Ji.L., Y.L, P.R. W.S., G.Y. and T.C. performed multiplex PCR and amplicon sequencing. Ji.L., and P.R. performed metatranscriptomic library construction and hybrid capture experiments. J.J., M.L, W.S., T.L., H.R., and H.Z. processed the sequencing data and conducted bioinformatic analyses. J.L., M.X. H.Z., J.J., M.L., and W.S. interpreted the data. M.X., J.J., M.L., and J.L. wrote and polished the manuscript. H.Z., W.S., L.Y., W.C. and Y.Z. contributed substantially to the manuscript revisions. All other authors provided useful suggestions and comments on the project and the manuscript.

COMPETING INTERESTS

The multiplex PCR panel is under patent application:

PCR PRIMER PAIR AND APPLICATION THEREOF

Patent applicant: MGI Tech Co.,Ltd

Name of inventor(s): Lin Yang,Ya Gao, Guodong Huang, Yicong Wang, Yuqian wang,

Yanyan Zhang, Fang Chen, Na Zhong, Hui Jiang, Xun Xu.

Application number: PCT/CN2017/089195

References

Jiang, S. *et al.* A distinct name is needed for the new coronavirus. *The Lancet* (2020).

Organization, W. H. in *Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/*   (Geneva: WHO, 2020).

Dudas, G. *et al.* Virus genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* **544**, 309-315, doi:10.1038/nature22040 (2017).

Dudas, G., Carvalho, L. M., Rambaut, A. & Bedford, T. Correction: MERS-CoV spillover at the camel-human interface. *Elife* **7**, doi:10.7554/eLife.37324 (2018).

Gardy, J. L. & Loman, N. J. Towards a genomics-informed, real-time, global pathogen surveillance system. *Nat Rev Genet* **19**, 9-20, doi:10.1038/nrg.2017.88 (2018).

Gire, S. K. *et al.* Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. *Science* **345**, 1369-1372, doi:10.1126/science.1259657 (2014).

Grubaugh, N. D. *et al.* Genomic epidemiology reveals multiple introductions of Zika virus into the United States. *Nature* **546**, 401-405, doi:10.1038/nature22400 (2017).

Sabir, J. S. *et al.* Co-circulation of three camel coronavirus species and recombination of MERS-CoVs in Saudi Arabia. *Science* **351**, 81-84, doi:10.1126/science.aac8608 (2016).

Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* **395**, 565-574, doi:10.1016/S0140-6736(20)30251-8 (2020).

Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

Chen, L. *et al.* RNA based mNGS approach identifies a novel human coronavirus from two individual pneumonia cases in 2019 Wuhan outbreak. *Emerg Microbes Infect* **9**, 313-319, doi:10.1080/22221751.2020.1725399 (2020).

Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).

Drosten, C. *et al.* Evaluation of advanced reverse transcription-PCR assays and an alternative PCR target region for detection of severe acute respiratory syndrome-associated coronavirus. *J Clin Microbiol* **42**, 2043-2047, doi:10.1128/jcm.42.5.2043-2047.2004 (2004).

Jonsdottir, H. R. & Dijkman, R. Coronaviruses and the human airway: a universal system for virus-host interaction studies. *Virol J* **13**, 24, doi:10.1186/s12985-016-0479-5 (2016).

Chan, J. F. *et al.* A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* **395**, 514-523, doi:10.1016/S0140-6736(20)30154-9 (2020).

Holshue, M. L. *et al.* First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med* **382**, 929-936, doi:10.1056/NEJMoa2001191 (2020).

McCrone, J. T. *et al.* Stochastic processes constrain the within and between host evolution of influenza virus. *Elife* **7**, e35962 (2018).

Ni, M. *et al.* Intra-host dynamics of Ebola virus during 2014. *Nat Microbiol* **1**, 16151, doi:10.1038/nmicrobiol.2016.151 (2016).

Park, D. J. *et al.* Ebola Virus Epidemiology, Transmission, and Evolution during Seven Months in Sierra Leone. *Cell* **161**, 1516-1526, doi:10.1016/j.cell.2015.06.007 (2015).

Domingo, E., Sheldon, J. & Perales, C. Viral quasispecies evolution. *Microbiol Mol Biol Rev* **76**, 159-216, doi:10.1128/MMBR.05023-11 (2012).

Metsky, H. C. *et al.* Zika virus evolution and spread in the Americas. *Nature* **546**, 411-415, doi:10.1038/nature22402 (2017).

Li, Q. *et al.* Reliable multiplex sequencing with rare index mis-assignment on DNB-based NGS platform. *BMC Genomics* **20**, 215, doi:10.1186/s12864-019-5569-5 (2019).

Xia, Z. *et al.* Advanced Whole Genome Sequencing Using a Complete PCR-free Massively Parallel Sequencing (MPS) Workflow. *bioRxiv* (2019).

Zou, L. *et al.* SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med*, doi:10.1056/NEJMc2001737 (2020).

26 Koressaar, T. & Remm, M. Enhancements and modifications of primer design program
Primer3. *Bioinformatics* **23**, 1289-1291, doi:10.1093/bioinformatics/btm091 (2007).
27 Untergasser, A. *et al.* Primer3--new capabilities and interfaces. *Nucleic Acids Res* **40**,
e115, doi:10.1093/nar/gks596 (2012).
28 Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification using
exact alignments. *Genome Biol* **15**, R46, doi:10.1186/gb-2014-15-3-r46 (2014).
29 Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884-i890, doi:10.1093/bioinformatics/bty560 (2018).
30 Chen, Y. *et al.* SOAPnuke: a MapReduce acceleration-supported software for integrated
quality control and preprocessing of high-throughput sequencing data. *Gigascience* **7**, 1-6,
doi:10.1093/gigascience/gix120 (2018).
31 Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics **27**, 863-864, doi:10.1093/bioinformatics/btr026 (2011).
32 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).
33 Au, C. H., Ho, D. N., Kwong, A., Chan, T. L. & Ma, E. S. K. BAMClipper: removing primers
from alignments to minimize false-negative mutations in amplicon next-generation
sequencing. *Sci Rep* **7**, 1567, doi:10.1038/s41598-017-01703-6 (2017).
34 Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J Comput Biol* **19**, 455-477, doi:10.1089/cmb.2012.0021 (2012).
35 Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection
and genome assembly improvement. *PLoS One* **9**, e112963,
doi:10.1371/journal.pone.0112963 (2014).
36 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079, doi:10.1093/bioinformatics/btp352 (2009).
37 Wickham, H. ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* **3**, 180-185
(2011).
38 Team, R. C. R: A language and environment for statistical computing. (2013).
39 Kolde, R. & Kolde, M. R. Package 'pheatmap'. *R Package* **1** (2015).
40 Kassambara, A. ggpubr:"ggplot2" based publication ready plots. *R package version 0.1* **6**
(2017).
41 Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.
arXiv preprint arXiv:1207.3907 (2012).
42 Seemann, T. in *Snippy: fast bacterial variant calling from NGS reads*   (WWW Document,
2015).
43 Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SNVEff: SNVs in the genome of Drosophila melanogaster strain w1118;
iso-2; iso-3. *Fly (Austin)* **6**, 80-92, doi:10.4161/fly.19695 (2012).
44 Li, R. *et al.* SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics* **25**,
1966-1967, doi:10.1093/bioinformatics/btp336 (2009).
45 Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915,
doi:10.1038/s41587-019-0201-4 (2019).
46 Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol **20**, 257, doi:10.1186/s13059-019-1891-0 (2019).
47 Lu, J., Breitwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: estimating species
abundance in metagenomics data. *PeerJ Computer Science* **3**, e104 (2017).
48 Kircher, M., Sawyer, S. & Meyer, M. Double indexing overcomes inaccuracies in multiplex
sequencing on the Illumina platform. *Nucleic Acids Res* **40**, e3, doi:10.1093/nar/gkr771
(2012).
593
594

595 **Table 1.** Metatranscriptomic sequencing data summary of eight HCoV-19 clinic

596 al samples collected from Guangzhou in February 2020

Sample ID	Sample Type	Ct	# of Sequenc-ing Read Pairs	# of HCoV-19 Read Pairs	% of HCoV-19 Read Pairs	Coverage (%)	Depth (X)
GZMU0047	nasal swab	18	1,547,648,648	85,316,930	5.513	100	113,021
GZMU0016	sputum	21	1,578,573,142	7,489,563	0.474	99.96	12,734
GZMU0048	throat swab	24	1,647,198,588	3,365,330	0.204	99.91	6,508
GZMU0044	nasal swab	26	1,609,367,415	7,275,402	0.452	99.92	12,758
GZMU0030	throat swab	29	1,725,727,056	31,148	0.002	99.87	69
GZMU0014	sputum	30	1,596,713,550	46,199	0.003	99.9	95
GZMU0042	sputum	32	1,481,162,934	567,266	0.038	99.94	1,133
GZMU0031	anal swab	32	1,671,721,507	25,392	0.002	99.89	14

599 **Table 2.** General characteristics of the three approaches employed in this study

	Metatranscriptomic sequencing	Hybrid capture-based sequencing	Multiplex PCR amplicon-based sequencing
Sequencing objective	Microbiome+Human	Target genome	Target genome
2nd strand synthesis	Y	Y	N
Fragmentation	Y	Y	N
Library preparation	Y	Y	N
PCR	18 cycles	18+18 cycles	15+25 cycles
Estimated time for pre-sequencing sample processing	10.5 h	20.5 h	7.5 h
Oligo synthesis	-	120 nt x 506	40-60 nt x 2 x (113+14+10)
Cost estimated for pre-sequencing sample processing	Moderate	High	Low
Estimated minimum data for downstream analyses (Base level)	>10Gb	Mb	Mb
Evenness	High	Moderate	Low
Sensitivity	+	++	+++
Accuracy (SNV)	+++	++	+++
Accuracy (iSNV)	+++	++	+

Figure 1. The general workflow of multiple sequencing approaches adopted in this study. We employed unique dual indexing (UDI) strategy and DNB-based (DNB, DNA Nanoball) PCR-free MPS platform to minimize index hopping and relevant sequencing errors[23,24,48]. **a**, Amplicon-based enrichment, the dual indexing was integrated in the 2nd PCR. Navy, multiplex PCR primers. **b**, Metatracriptomic library preparations, the dual indexing was integrated in the universal PCR. **c**, Library preparations and hybrid capture-based enrichment, the 1st indexing was integrated in the pre-capture PCR while the 2nd indexing was integrated in the post-capture PCR. Ocher, ssDNA probes. Red and green lines represent adaptor sequences; green dots represent phosphate groups.

Figure 2. Overview of the study design. Eight clinical samples (GISAID accession
EPI_ISL_414663, EPI_ISL_414686 to EPI_ISL_414692); 16, GZMU0016; 30, GZMU0030;
31, GZMU0031; 42, GZMU0042; 44, GZMU0044; 47); 16, GZMU0016; 30, GZMU0030;
31, GZMU0031; 42, GZMU0042; 44, GZMU0044; 47); 16, GZMU0016; 30, GZMU0030;
31, GZMU0031; 42, GZMU0042; 44, GZMU0044; 47) and serial dilutions of a cultured
isolate were subjected to direct metatranscriptomic library construction, amplicon-based
enrichment, and hybrid capture-based enrichment, respectively. Libraries generated from
each method were pooled, respectively. DNB, DNA Nanoball. 14, GZMU0014 (GISAID
accession EPI_ISL_414663); 16, GZMU0016; 30, GZMU0030; 31, GZMU0031; 42,
GZMU0042; 44, GZMU0044; 47, GZMU0047; 48, GZMU0048. D0, undiluted sample of the
cultured isolate; D1-D7, seven serial diluted samples of the cultured isolate, ranging from
1E+07 to 1E+01 genome copies per mL, in 10-fold dilution. -, negative controls prepared
from nuclease-free water and human nucleic acids. PE100, paired-end 100-nt reads;
SE400, single-end 100-nt reads.
Su

629
630
Figure 3. Sequencing coverage and depth of the cultured isolate and eight clinical
samples. a, Amplicon sequencing coverage by sample (row) across the HCoV-19
genome. Pink, sequencing depth ≥100×; heatmap (bottom) sums coverage across all
samples. HNA, negative control prepared from human nucleic acids; water, negative
control prepared from nuclease-free water. Green horizontal lines on heatmap, amplicon
locations. Overlap regions between amplicons range from 59-209 bp. **b-d**, Normalized
coverage across viral genomes of the clinical samples across methods. **e**, HCoV-19-RPM
sequenced plotted against genome copies per mL for the cultured isolate. Three
independent experiments were performed for amplicon sequencing. Pink, ~400 bp (base
pairs) amplicon-based sequencing including human and lambda phage nucleic acids
background; red, ~200 bp amplicon-based sequencing; orange, ~400 bp amplicon-based
sequencing excluding human and lambda phage nucleic acids background (NAB); light
blue, capture sequencing. **f**, HCoV-19-RPM (Reads Per Million) sequenced plotted against
qRT–PCR Ct value for the clinical samples. Pink, amplicon; light blue, capture; blue, meta.
g, Estimated minimum amount of bases required by each method for high-confidence
downstream analyses. Pink, amplicon; light blue, capture.

647
648
Figure 4. Between-sample and within-sample variants of HCoV-19 detected across
methods. a, SNVs detected between clinical samples against a reference genome (GISAID
accession: EPI_ISL_402119). Alleles with ≥80% frequencies were called. *, SNVs verified
by Sanger sequencing. **b**, Allele frequencies of the identified SNVs. Pink, amplicon; light
blue, capture; blue, meta. Minor allele frequencies detected in serial dilutions of the cultured
isolate (**c**) and clinical samples (**d**) across methods. Pink, amplicon vs meta; light blue,
capture vs meta. Minor alleles are defined with ≥5% and <50% frequencies. Besides general
quality filter, iSNVs had to pass depth and strand bias filter as described in Methods.