Title: gauseR: Simple methods for fitting Lotka-Volterra models describing Gause's "Struggle for Existence"

Authors: Lina K. Mühlbauer (ORCID: 0000-0002-6493-2742)[1,‡], and Maximilienne Schulze (ORCID: 0000-0002-9619-561X)[1,‡], W. Stanley Harpole (ORCID: 0000-0002-3404-9174)[1,2,3], and Adam T. Clark (ORCID: 0000-0002-8843-3278)[*,2,3,4]

Institutions:
[*]Corresponding Author: adam.tclark@gmail.com
[1]Institute of Biology, Martin Luther University, Halle, Germany; [2]Department of Physiological Diversity, Helmholtz Centre for Environmental Research (UFZ), Leipzig, Germany; [3]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany; [4]Synthesis Centre for Biodiversity Sciences (sDiv), Deutscher Platz 5e, 04103 Leipzig, Germany
[‡]Authors Contributed Equally

Running head: gauseR: Lotka-Volterra and Gause

Key Words: competition; predator-prey; logistic growth; optimization, differential equation, growth rate.
Word count: 3147

Author contributions: L.M., M.S., and A.T.C. wrote the first draft of the paper, designed the functions and R package, and conducted the analyses. L.M. and M.S. contributed equally to this manuscript, and their authorship order was determined by a game of chance. W.S.H. and A.T.C. conceived of the project. All authors contributed significantly to revisions.

Data availability: All data, and corresponding metadata, for the analyses in this manuscript are available through the `gauseR` package, available at github.com/adamtclark/gauseR. Upon publication, the code and data will be uploaded to the CRAN R package repository.

Abstract (262 words)

The ecological models of Alfred J. Lotka and Vito Volterra have had an enormous impact on

ecology over the past century. Some of the earliest – and clearest – experimental tests of these

models were famously conducted by Georgy Gause in the 1930's. Although well known, the

data from these experiments are not widely available, and are often difficult to analyze using

standard statistical and computational tools. Here, we introduce the `gauseR` package, a

collection of tools for fitting Lotka-Volterra models to time series data of one or more species.

The package includes several methods for parameter estimation and optimization, and includes

42 datasets from Gause's species interaction experiments and related work. Additionally, we

include with this paper a short blog post discussing the historical importance of these data and

models, and an R vignette with a walk-through introducing the package methods. The package

is available for download at github.com/adamtclark/gauseR. To demonstrate the package, we

apply it to several classic experimental studies from Gause, as well as two other well-known

datasets on multi-trophic dynamics on Isle Royale, and in spatially structured mite populations.

In almost all cases, models fit observations closely, and fitted parameter values make ecological

sense. Taken together, we hope that the methods, data, and analyses that we present here

provide a simple and user-friendly way to interact with complex ecological data. We are

optimistic that these methods will be especially useful to students and educators who are

studying ecological dynamics, as well as researchers who would like a fast tool for basic

analyses.

Introduction

A century ago, Alfred J. Lotka (1920, 1925) and Vito Volterra (1926) published their canonical

works concerning the mathematics of species interactions. These theories were among the first

to suggest that the dynamics of complex ecological systems could be described through simple

underlying equations (Giorgio 1988). Inspired by the possibility of understanding natural

systems through mathematics, Georgy Gause stated in his "Struggle for Existence" (1934b):

> *"[…] in order to penetrate deeper into the nature of these phenomena we*
>
> *must combine the experimental method with the mathematical theory, a*
>
> *possibility which has been created by the brilliant researches of Lotka and*
>
> *Volterra."*

To match the strong assumptions and narrow scope of Lotka and Volterra's basic equations,

Gause constructed simple experimental microcosms, in which he was able to show that

interactions between pairs of species influenced their abundances and dynamics in ways that

accorded with theoretical predictions (Gause 1934a,b). These experiments laid the groundwork

for a century of experiments based on mathematics as a conceptual framework. It has since

become a primary goal of ecology to explain empirical observations with mathematical theory.

Despite their simplicity, the Lotka-Volterra equations are still the primary models used to

describe species interactions. At their most basic, these models describe the changes per unit

time in the abundance of a species, N_i, as

$$dN_i/dt = N_i(r_i + a_{ii} N_i + \Sigma_j(a_{ij} N_j)) \qquad \text{Eq. (1a)}$$

where r_i is the intrinsic growth rate, which describes the growth of a species at low density and

in the absence of other species, a_{ii} describes the effect of species i on its own growth, and a_{ij}

describes the effect of species j on the growth of species i. The form that we use here differs

somewhat from that often portrayed in some textbooks, as we leave r_i inside the parenthesis

and therefore do not standardize interaction coefficients by the growth rate (thus, we signify

these as *a* rather than *α*). If we divide both sides of Eq. (1a) by N_i to calculate the "per-capita"

growth rate of species *i*, the model can be written as a simple linear equation

$dN_i/N_i dt = r_i + a_{ii} N_i + \Sigma_j(a_{ij} N_j)$ Eq. (1b)

This form is particularly useful for analyzing empirical data, as the parameters can be estimated

using ordinary least squares regression of per-capita growth rates against species abundances,

where r_i is the y-intercept intercept, a_{ii} is the slope with respect to species *i*'s own abundance,

and a_{ij} is the slope with respect to other species (Lehman *et al*. 2020). Note that in a single

species, this relationship can also be expressed in terms of the carrying capacity, K_i, which

describes the equilibrium abundance reached by a species growing in the absence of other

species, and falls on the x-intercept (Fig. 1). For those who are less familiar with regression or

dynamical systems modelling, we have included a more in-depth discussion of this approach in

the vignette accompanying our R package. We can solve for K_i in terms of a_{ii} and r_i by setting

Eq. (1b) equal to zero, and solving for N_i, which yields

$K_i = -r_i/a_{ii}$ Eq. (1c)

61   Because the Lotka-Volterra models make no specific assumptions about the

mechanisms underlying species interactions, they can be parameterized in ways that

approximate any combination of underlying mechanisms, at least locally around equilibrium

(MacArthur 1970; Chesson 2000; Letten & Stouffer 2019). For example, in a system with two

species, if both a_{ij} and a_{ji} are less than zero, species suppress each other's growth, indicating

competition. If both parameters are positive, then both species benefit from the other's

presence, indicating mutualism. Alternatively, if $a_{ij} < 0$ and $a_{ji} > 0$, then species *i* is suppressed

by species *j* whereas species *j* benefits from the presence of species *i*, indicating a predator-

prey relationship.

70   Gause's original experimental work still provides some of the best examples of these

different kinds of interspecific interactions. However, because of the limited analytical tools

available to him in the early twentieth century, Gause was only able to fit models to a small

subset of his experimental data. Moreover, because of the journal formatting conventions of his

time, much of Gause's data was never officially published. Thus, although they are widely used

as teaching tools and widely cited as canonical examples of competitive and predator-prey

interactions, very few of his results have ever been quantitatively analyzed using the Lotka-

Volterra equations.

Here, we introduce the `gauseR` package, which includes a collection of functions for

confronting empirical observations with the classical Lotka-Volterra interaction models. The

package combines methods for data formatting, model simulation, and parameter estimation

into a simple-to-use, automated wrapper function. The package also acts as a digital repository

for Gause's experiments on the "Struggle for Existence", and includes all available data from his

1934 book and associated publications. Below, we introduce the functions and basic workflow

for the package, and explain the underlying theory. To demonstrate the package's functionality,

we analyze two well-known datasets from Gause (1934a), as well as classic examples of multi-

species interactions from McLaren & Peterson (1994) and Huffaker *et al*. (1963). Additionally,

we discuss some limitations and possible extensions of the optimization methods used in the

package. We foresee that `gauseR` will be especially helpful for researchers who seek simple

tools for analyzing their data, as well as a teaching aid for introducing students to the Lotka-

Volterra equations and to Gause's experiments.

Methods

Functions and workflow

The general purpose of the functions in the `gauseR` package is to fit the general Lotka-Volterra

model in Eqs. (1a-b) to observational time series data of one or more interacting species. The

package, along with a walk-through vignette, is available at github.com/adamtclark/gauseR.

Functions are described in detail in Table 1. In general, this procedure will require estimating

values for three types of parameters: starting abundances (N_{0i}), intrinsic growth rates (r_i) and

interaction coefficients (a_{ij}).

For a single species system, dynamics can be directly fit to the Logistic growth model

using the `get_logistic` function. For systems with two or more species, fitting proceeds in

four steps. First, time-lagged abundances for each species are calculated using the `get_lag`

function, yielding pairs of abundance values separated by a fixed time interval for each species.

Second, these time-lagged abundances are used to estimate per-capita growth rates at different

time points, using the `percap_growth` function. This function approximates dynamics following

the linearized equation

$\mathrm{d}N/N\mathrm{d}t = \log(N(t_2)/N(t_1))/(t_2 - t_1)$ Eq. (2)

Third, linear regressions are used to test how per-capita growth rates vary as a function

of conspecific and heterospecific species abundances, from which the parameters in Eq. (1b)

can be estimated. Finally, because parameter estimates from linearized growth rates often do

not match observations for long or complex time series, the `lv_interaction` and `lv_optim`

function are used to directly simulate dynamics and tune parameters to match model outputs to

observations, which can greatly improve model performance (Carrara *et al.* 2015; Rosenbaum &

Rall 2018; Maynard *et al.* 2019). All four of these steps are automated in the `gause_wrapper`

function, which takes observed species abundances as an input, and returns fitted parameter

estimates. We suggest this option for most users.

To facilitate optimization, we include two additional functions. First, the

`lv_interaction_log` function simulates dynamics for log-transformed species abundances.

This approach is helpful in systems where abundances approach zero, which can cause

integration issues (e.g. predictions of negative abundances). This function is the default for

`gause_wrapper`. Lastly, the `ode_prediction` function simulates dynamics for *n* species

given a set of input parameter values, and returns estimates of species abundances as a single

column vector. This function is potentially useful for users who wish to apply other, more

sophisticated optimization routines.

Data and example analyses

The `gauseR` package includes a wide range of digitized data from Gauses' experimental work,

as well as some other well-known multi-species time series. Full metadata and citations for

these datasets are available in the help documentation for the package. Dataset names refer to

the source and figure or table from which the data are drawn. Whenever possible, data were

taken from tables in Gause's publications or appendices. Otherwise data were digitized from

figures using WebPlotDigitizer (Rohatgi 2015).

To demonstrate how to apply our functions, we also conduct example analyses using

four of these datasets. First, `gause_1934_science_f02_03` shows Gause's classic

competition experiment between *Paramecium aurelia* and *P. caudatum*, including both species

grown in monoculture and in mixture (Gause 1934a,b). We use these data to demonstrate

parameter fitting methods for logistic growth and Lotka-Volterra competition. Full source code

for these analyses are available in the help documentation for the `get_logistic` and

`lv_interaction` functions. Second, `gause_1934_science_f01` includes interactions

between the prey species *P. caudatum* and predator species *Didinium nasutum*, with which we

demonstrate the difference between estimates derived from linearized models vs. from the

optimizer fit to the full simulated dynamics. These analyses are shown in the help

documentation for the `lv_optim` function. Third, to demonstrate parameter fitting in a system

with more than two species, we use `mclaren_1994_f03`, which summarizes dynamics of

wolves (*Canis lupus*), moose (*Alces alces*), and fir trees (*Abies balsamea*) on Isle Royale from

McLaren & Peterson (1994). Finally, to demonstrate a case where classic Lotka-Volterra

interaction models are unable to match observations, we analyze dynamics of the prey species

Eotetranychus sexmaculatus and predator species *Typhlodromus occidentalis* from Huffaker's

classic mite experiments (Huffaker *et al*. 1963). Data are available in `huffaker_1963`. Source

code for the analyses of the McLaren and Huffaker datasets are available in the supplement.

Results

In most cases, the simple Lotka-Volterra model in Eqs. (1a-b) accurately describe observed

dynamics, and provide interpretable parameter values. For both logistic growth (Fig. 2) and

competitive interactions between the two *Paramecium* species (Fig. 3), observations closely

matched model predictions. Fitted parameter values suggest that both species are subject to

self-limitation in monoculture (i.e. a_{ii} and $a_{jj} < 0$), and indicate competitive interactions in mixture

(i.e. a_{ij} and $a_{ji} < 0$) (Table 2a-b).

For predator-prey interactions between *D. nasutum* and *P. caudatum*, we found that

parameter estimates from regressions fitted to the linearized growth rates failed to match

observed dynamics (Fig. 4; Table 2c). However, predictions improved substantially after using

the optimizer to tune parameters. For the wolf-moose-fir system of McLaren *et al*. (1994), results

from the optimizer also matched observed dynamics (Fig. 5). Because we expected no direct

interactions between wolves and fir trees, we set these interactions to zero prior to analyses.

Additionally, because McLaren *et al.* hypothesized that moose populations are subject to top-

down control in this system, we set self-limitation for moose to zero (Table 2d).

Lastly, for the Huffaker mite data, we were not able to fit the general Lotka-Volterra

model to match observed dynamics (Fig. 6; Table 2e). In this system, dynamics follow sustained

oscillations, but with periods and amplitudes that vary over time. However, our fitted models

could only achieve either sustained oscillations of a fixed period and amplitude (top panel), or

damped oscillations where the amplitude declined over time (bottom panel), but not both.

Discussion

The `gauseR` package makes it possible to analyse time series data in single- or multi-

species systems using the classic Lotka-Volterra interaction models in an easy and automated

way. For almost all of the examples that we consider here, models were able to accurately

capture observed dynamics, and analyses could be conducted using just the default

`gause_wrapper` function without any modification of the code or additional inputs from the

user. Furthermore, fitted parameter values matched those expected for logistic growth,

competition and predator-prey dynamics, as predicted by Lotka and Volterra. We are therefore

hopeful that this combination of fitting tools, along with the package's repository for Gause's

data, will be especially useful for teaching Lotka-Volterra's basic equations and introducing

Gause's experimental work.

Interpreting the models

In the monoculture experiments, our results demonstrate that *P. caudatum* followed

logistic growth. As expected by theory, the species was able to increase from low abundance

(i.e. $r_i > 0$), limited its own growth as its abundance increased (i.e. $a_{ii} < 0$), and approached a

fixed carrying capacity (K) over time (Lotka 1925; Volterra 1926; Gause 1934a,b). For

competition between *P. caudatum* and *P. aurelia*, fitted parameters indicated self-limitation by

both species, and inhibition of each species by the other (i.e. a_{ii}, a_{jj}, a_{ij} and $a_{ji} < 0$). Again, these

values match theoretical expectations from Lotka and Volterra, and accord with Gause's original

analyses of the data.

For the system with *D. nasutum* and *P. caudatum*, fitted models predicted that prey

abundance increased in the absence of predators (i.e. $r_i > 0$), and that the predators declined

exponentially in the absence of prey (i.e. $r_i < 0$,). Similarly, parameter estimates indicated that

predators had a negative effect on prey, and prey had a positive effect on predators (i.e. $a_{ij} < 0$

and $a_{ji} > 0$, respectively). Jointly, these parameter values match biological expectations - i.e.

that predators consume prey, and starve when prey are absent. However, in contrast to the

classic formulation of the Lotka-Volterra predator-prey model, we also identified self-limitation by

both species (i.e. $a_{ii} < 0$ and $a_{jj} < 0$). These parameter values resulted in damped oscillations in

species abundances, as predicted by theory (Lehman *et al.* 2000). Results were similar for the

wolf-moose-fir tree example, for which fitted parameters indicated self-limitation by wolves and

fir trees, positive effects of moose on wolves and of fir trees on moose, and negative effects of

wolves on moose and of moose on fir trees. Again, predictions from the model closely matched

observations, supporting the hypothesis that top-down control of moose by wolves could explain

dynamics in this system (McLaren & Peterson 1994).

In contrast, none of the models that we fitted were able to match observations from the

Huffaker *et al.* mite experiments. Importantly, this limitation is a general characteristic of the

Lotka-Volterra models, and not a peculiarity of our fitting methods (Lotka 1925; Hatton *et al.*

2015; Lehman *et al.* 2000). When self-limitation is absent for both species, Lotka-Volterra

predator-prey models display "neutrally stable" oscillations, meaning that oscillations maintain a

fixed amplitude and period. When self-limitation exists for either the prey species or the

predator, or both, then oscillations become damped, meaning that over time they decrease in

amplitude and period until they reach stable equilibrium values. Because the Huffaker data

include persistent oscillations for which the amplitude and period vary over time, they cannot be

matched by any parameterization of the Lotka-Volterra equations. Potentially, these complex

dynamics are a result of the spatial structure in the Huffaker experiment, which often cannot be

described with simple equations (Leibold & Chase 2018).

Notes on optimization

For the predator-prey examples that we analyse, our findings show that parameter

estimates generated from the linearized growth rates in Eq. (1b) failed to capture observed

dynamics. This result accords with many other studies, which show that model performance is

typically much better when parameters are tuned to match fully simulated dynamics, i.e. as is

accomplished with the `lv_optim` function (Carrara *et al.* 2015; Rosenbaum & Rall 2018;

Maynard *et al.* 2019). In general, the difference in fit between these two approaches is driven by

the sensitivity of the Lotka-Volterra equations to noisy parameter estimates - even very small

changes in parameter values can lead to large changes in model predictions (Clark &

Neuhauser 2018; Maynard *et al.* 2019). We therefore suggest that in most cases, the

`gause_wrapper` function, which automatically applies the optimization routine to simulated

dynamics, will provide the best fits to empirical data.

Importantly, the `gause_wrapper` function applies several optimization "tricks" that help

improve model performance. First, comparisons between observations and predictions are

carried out on standardized abundances, such that the mean for each species is equal to one.

This prevents abundant species from dominating the fitting process. For example, for the

McLaren & Peterson data, optimization without standardization would closely fit dynamics of

moose, the most common species, at the cost of model performance for wolves and fir trees.

Second, by default, dynamics are simulated for log-transformed abundances, and are then

back-transformed for comparisons to observations. This approach helps prevent computational

errors in the integration from erroneously driving abundances negative. Third, we also log-

transform parameter values during the optimization process, after determining their sign *a priori*

from the linearized functions. This makes it possible to limit the number of positive interaction

coefficients, which otherwise can cause the system to generate unrealistically high abundance

values (e.g. via run-away mutualism) and crash the optimizer (Lehman *et al.* 2020). A trade-off

that comes with this added stability is that this method does not allow for significance testing,

because it constrains the sign of parameters. Thus, the standard errors provided by the

`gause_wrapper` should only be thought of as rough approximations of the degree of

uncertainty for each parameter estimate. Additionally, note that because our approach does not

update predictions based on observations in each time step, it effectively assumes that all

deviations between observations and predictions are due to observation error (i.e. not process

noise) (Rosenbaum & Rall 2018). Users who wish to apply more sophisticated optimization

methods should see the documentation and examples for the `ode_prediction` function.

There are several potential sources of observation error in the data that we analyze

here. Most obviously, data digitized from figures, and especially from hand-made figures such

as those of Gause and Huffaker *et al.*, are necessarily imperfect. More generally, there is

reason to believe that some of Gause's original data and figures contained some errors. In

particular, in a few cases where data are available in both figures and table form, there are

small differences in reported values. Similarly, many of the units reported by Gause, such as

"Volume" or "Amount", are somewhat difficult to interpret and compare across figures and

experiments. Lastly, in some cases, data varies between being reported for individual replicates,

to mean values summed across replicates. These circumstances present a challenge in the

analysis, as they make it difficult to compare values across the entire time series, and exclude

information about potentially meaningful replicate-level dynamics and between-replicate

variation. Wherever possible, we have noted these inconsistencies in the metadata.

Future directions

Given the strong correspondence between the models of Lotka (1920, 1925) and

Volterra (1926) and the experimental data of Gause (1934a,b), it is perhaps not surprising that

these works have had such an enormous impact on ecology over the past century. Together,

these concepts have helped ecologists better describe and model their systems, and have

greatly advanced conceptual understanding of how different arrangements of species and

interaction types are likely to play out. Nevertheless, as the type of systems that ecologists

study grow ever more complex, it seems likely that we may need to move beyond these basic

models to new frameworks that allow more nuanced and dynamic interaction types (Letten &

Stouffer 2019). We therefore hope that the `gauseR` package is useful in better exploring both

the classic insights, and operational limitations, of Lotka, Volterra, and Gause's framework.

Acknowledgements: The authors declare no conflicts or specific funding sources for the work

described in this manuscript.

References

Carrara, Francesco, Andrea Giometto, Mathew Seymour, Andrea Rinaldo, and Florian

Altermatt. 2015. "Inferring Species Interactions in Ecological Communities: A Comparison of

Methods at Different Levels of Complexity." *Methods in Ecology and Evolution* 6:895-906.

Clark, Adam T., and Claudia Neuhauser. 2018. "Harnessing Uncertainty to Approximate

Mechanistic Models of Interspecific Interactions." *Theoretical Population Biology* 123:35-44.

Chesson, Peter. 2000. "Mechanisms of Maintenance of Species Diversity." *Annual Review of*

Ecology and Systematics 31:343-66.

Gause, G. F. 1934. "Experimental Analysis of Vito Volterra's Mathematical Theory of the

Struggle for Existence." *Science* 79:16-17.

Gause, GF. 1934. *The struggle for existence*. Williams and Wilkins. Baltimore. 163 pp.

Hatton, I. A., K. S. McCann, J. M. Fryxell, T. J. Davies, M. Smerlak, A. R. E. Sinclair, and M.

Loreau. 2015. "The Predator-Prey Power Law: Biomass Scaling across Terrestrial and

Aquatic Biomes." Science 349:aac6284–aac6284.

Huffaker, C. B., K. B. Shea, and S. G. Herman. 1963. "Experimental Studies on Predation:

Dispersion Factors and Predator-Prey Oscillations." *Hilgardia* 34:305-330.

Israel, G. 1988. "On the Contribution of Volterra and Lotka to the Development of Modern

Biomathematics". *History and Philosophy of the Life Sciences* 10:37-49.

Leibold, Mathew A, and Jonathan M Chase. 2018. *Metacommunity Ecology, Volume 59*.

Princeton University Press, Princeton, USA.

Lehman, Clarence, Shelby Loberg, A.T. Clark, and Daniel Schmitter. In press. "Unifying the

Basic Models of Ecology to Be More Complete and Easier to Teach." *BioScience*.

Letten, Andrew D., and Daniel B. Stouffer. 2019. "The Mechanistic Basis for Higher-Order

Interactions and Non-Additivity in Competitive Communities." *Ecology Letters* 22:423-36.

Lotka, A. J. (1920). "Analytical Note on Certain Rhythmic Relations in Organic Systems". *PNAS*

6:410-415.

Lotka, A. J. (1925). *Elements of Physical Biology*. Williams and Wilkins. Baltimore. 495 pp.

MacArthur, R. H. 1970. "Species Packing and Competitive Equilibrium for Many Species."

Theoretical Population Biology 1:1–11.

Maynard, Daniel S., J. Timothy Wootton, Carlos Serván, and Stefano Allesina. 2019.

"Reconciling Empirical Interactions and Species Coexistence." *Ecology Letters* 22:1028-37.

McLaren, B. E., and R. O. Peterson. 1994. "Wolves, Moose, and Tree Rings on Isle Royale."

Science 266:1555–58.

Rohatgi, Ankit. 2015. *WebPlotDigitizer*, Version 4.2. https://apps.automeris.io/wpd/. Accessed

11 March 2020.

Rosenbaum, Benjamin, and Björn C. Rall. 2018. "Fitting Functional Responses: Direct

Parameter Estimation by Simulating Differential Equations." *Methods in Ecology and*

Evolution 9:2076–90.

Volterra, V. 1926. "Variazioni e Fluttuazioni Del Numero d'individui in Specie Animali

Conviventi." *Mem. Acad. Lincei Roma*. 2:31-113.

Figures

324
Figure 1: Figure comparing per-capita growth rate (following Eq. 1b) vs. abundance for

Paramecium aurelia grown in monoculture. Note that when plotted in this manner, the

parameter values for the Lotka-Volterra equations can be easily "read" off the graph, which *r* as

the y-intercept, *K* as the x-intercept, and a_{ii} as the slope. In multi-species systems, interaction

coefficients a_{ij} (i.e. effect of species j on species i) can be similarly computed based on the slope

of $dN_i/N_i dt$ vs. N_j.

Goodness of Fit: 0.98

331
332

Figure 2: Logistic growth for size standardized volume of *Paramecium caudatum* grown in

monoculture on Osterhout medium over 24 days, from Gause (1934a). Points show

observations, and line shows fitted logistic growth curve, calculated with the `get_log()`

function, that can be used to fit the logistic growth function to time series of an individual

species. Goodness of fit shows results from the `tests_goodness_of_fit()` function. Recall

that values near 1 imply a close correspondence between observations and predictions. See

Table 1a for parameter values.

340
341

Figure 3: Lotka-Volterra competition for size standardized volume of *Paramecium caudatum*

and *Paramecium aurelia* grown in mixed population over 24 days, from Gause (1934a). Points

show observations, and lines show growth curves fitted using the `lv_interaction()`

function, which can be used to simulate dynamics in multi-species mixtures following the classic

Lotka-Volterra equations. See Table 1b for parameter values.

347
348

Figure 4: Predator-prey interactions between *Didinium nasutum* and *Paramecium caudatum*

grown in mixture over 17 days, from Gause (1934a). Additional individuals of both species were

added to the mixture periodically to prevent local extinction. Points show observations, and lines

show fitted growth curves. Top panel shows results for model fitted using linear regressions of

species abundances vs. per-capita growth rate, whereas bottom panel shows results for

parameters fitted using the simulated differential equations using the `lv_optim()` function.

Note that goodness of fit is varies substantially between the two methods. See Table 1c for

parameter values.

357
358

Figure 5: Multi-trophic dynamics for wolves, moose, and fir trees on Isle Royale from 1960 to

1994, from McLaren & Peterson (1994). Points show observations, and lines show growth

curves fitted using the `lv_optim()` function. The left axis shows wolf and moose abundances,

separated by a ten-fold scaling difference for easier visualization, whereas the right axis shows

tree growth increments. Two methods for testing goodness of fit are shown. In the legend,

values show univariate tests for each species, whereas "Total GOF" shows fit when considered

across all observations simultaneously. Note that these two methods of comparison can lead to

very different conclusions. See Table 1d for parameter values.

367
368

Figure 6: Predator-prey interactions between *Eotetranychus sexmaculatus* and *Typhlodromus*

occidentalis in a spatially structured experiment carried out on a grid of oranges over 60 weeks,

from Huffaker *et al.* (1963). Points show observations, and lines show growth curves fitted using

the `lv_optim()` function. Top panel shows dynamics for a system where neither species

directly inhibits its own growth (i.e. $a_{ii} = a_{jj} = 0$), whereas bottom panel is for a system where only

the predator directly inhibits its own growth. Note that neither option is able to fully capture the

realized dynamics, and that goodness of fit is relatively low in both panels. See Table 1e for

parameter values.

377 **Tables**

Function	Description
get_lag	Calculates time-lagged observations for variable x, separated by treatment.
percap_growth	Calculates per-capita growth rate, using log ratios.
get_logistic	Calculates logistic growth for a population.
lv_interaction	Calculates dn/dt for n species in a Lokta-Volterra system.
lv_interaction_log	Calculates dn/dt for n species in a Lokta-Volterra system, with log-transformed abundances. This form can be helpful for optimization routines where species abundances are close to zero.
lv_optim	Identifies optimal parameter values for a Lotka-Volterra interaction system by simulating dynamics using differential equations.
gause_wrapper	Automatically runs routine for finding starting values and optimal parameter values for a Lotka-Volterra interaction system.
test_goodness_of_fit	Generates an R^2-like goodness of fit index, based on the squared difference between observed and predicted values. Index values of 1 indicate a very close fit, whereas values near or below zero indicate a poor fit.
ode_prediction	Calculates expected abundances for all n species in a community. This function is potentially useful in combination with other optimizer software, e.g. as might be used for hypothesis testing.

378
379 **Table 1:** Brief descriptions of the `gauseR` functions.

a. | **P. caudatum monoculture**

N_0	0.218 (± 0.182)
r	0.959 (± 0.117)
K	202.305 (± 3.033)
(a_{ii})	-0.005 (± 0.001)

b. | **Competition experiment**

	P. caudatum		P. aurelia
N_{0i}	1.595 (± 1.585)	N_{0j}	1.632 (± 1.543)
r_i	1.259 (± 0.911)	r_j	1.026 (± 0.550)

A	P. caudatum	P. aurelia
P. caudatum	-0.005 (± 0.015)	-0.002 (± 0.009)
P. aurelia	-0.008 (± 0.015)	-0.007 (± 0.008)

c. | **Predator-prey experiment**

Linear model

	P. caudatum (prey)		D. nasutum (predator)
r_i	0.998 (± 0.297)	r_j	-0.069 (± 0.538)

A	P. caudatum	D. nasutum
P. caudatum	-0.021 (± 0.013)	0.039 (± 0.013)
D. nasutum	-0.068 (± 0.018)	-0.026 (± 0.02)

ODE optimiser

	P. caudatum (prey)		D. nasutum (predator)
N_{0i}	1.655 (± 1.139)	N_{0j}	0.117 (± 0.11)
r_i	1.099 (± 0.295)	r_j	-0.89 (± 0.358)

A	P. caudatum	D. nasutum
P. caudatum	-0.013 (± 0.006)	0.084 (± 0.024)
D. nasutum	-0.078 (± 0.029)	-0.002 (± 0.004)

d. | **McLaren & Peterson (1994)**

	A. balsamea (producer)		A. alces (herbivore)		C. lupus (predator)
N_{0i}	0.326 (± 0.093)	N_{0j}	699.9 (± 210.3)	N_{0k}	24.312 (± 2.397)
r_i	0.238 (± 0.026)	r_j	2.021 (± 0.354)	r_k	0.01 (± 0.008)

A	A. balsamea	A. alces	C. lupus
A. balsamea	-0.139 (± 0.144)	0.002 (± 0.001)	0
A. alces	-0.0002 (± 0.00002)	0	0.00004 (± 0.000004)
C. lupus	0	-0.088 (± 0.019)	-0.003 (± 0.001)

e. | Huffaker et al. (1963)

	E. sexmaculatus (prey)		T. occidentalis (predator)
N_{0i}	279.142 (± 60.439)	N_{0j}	8.446 (± 0.989)
r_i	0.344 (± 0.097)	r_j	-0.236 (± 0.089)

A	E. sexmaculatus (prey)	T. occidentalis (predator)
E. sexmaculatus	0	0.0005 (± 0.0001)
T. occidentalis	-0.059 (± 0.023)	0

Predator self-limitation

	E. sexmaculatus (prey)		T. occidentalis (predator)
N_{0i}	64.029 (± 5.784)	N_{0j}	5.23 (± 0.597)
r_i	0.187 (± 0.004)	r_j	-0.377 (± 0.022)

A	E. sexmaculatus	T. occidentalis
E. sexmaculatus	0	0.0012 (± 0.0001)
T. occidentalis	-0.028 (± 0.003)	-0.024 (± 0.002)

Table 2: Fitted parameter values for each analysis described in the main text. Values show

parameter estimates ± one standard error. For interaction matrices A, parameters show the

effect of the species listed in the *row* on the species listed in the *column* (i.e. a_{ij} is listed in row j

and column i). Note that standard errors are approximated from the numerically differentiated

Hessian values calculated for the log-transformed parameters, based on the difference between

the mean estimate and the lower standard deviation in log space, and are therefore only meant

as rough approximations of the degree of uncertainty.

Table S1: Summary table of the datasets in the `gauseR` package. Column "name" is the name

of the dataset, and can be used to load it. For all other column metadata, see the help files for

each dataset in the package (i.e. via the `?name` command). See

gauseR_TableS1_dataset_summarytable.csv file for contents of table. Information that is

included in each dataset is marked with "Y".