1

Characterization of the SARS-CoV-2 Spike in an Early Prefusion Conformation

Tingting Li[1,2,#], Qingbing Zheng[1,2,#], Hai Yu[1,2,#], Dinghui Wu[3,#], Wenhui Xue[1,2,#],

Yuyun Zhang[1,2], Xiaofen Huang[1,2], Lizhi Zhou[1,2], Zhigang Zhang[1,2], Zhenghui Zha[1,2],

Tingting Chen[1,2], Zhiping Wang[1,2], Jie Chen[1,2], Hui Sun[1,2], Tingting Deng[1,2], Yingbin

Wang[1,2], Yixin Chen[1,2], Qingjian Zhao[1,2], Jun Zhang[1,2], Ying Gu[1,2,*], Shaowei Li[1,2,*],

Ningshao Xia[1,2,*]

[1] State Key Laboratory of Molecular Vaccinology and Molecular Diagnostics, School

of Life Sciences, School of Public Health, Xiamen University, Xiamen, China 361102

[2] National Institute of Diagnostics and Vaccine Development in Infectious Disease,

Xiamen University, Xiamen, China 361102

[3] Department of Pulmonary Medicine, The First Affiliated Hospital of Xiamen

University, Xiamen, China 361003

[#] T.L., Q.Z., D.W., H.Y. and W.X. contributed equally to this work.

[*] To whom correspondence may be addressed. E-mail: (Y.G.) guying@xmu.edu.cn   or

(S.L.) shaowei@xmu.edu.cn or (N.X.) nsxia@xmu.edu.cn

Running title: SARS-CoV-2 spike in an early prefusion conformation

Abstract

Pandemic coronavirus disease 2019 (COVID-19) is caused by the emerging severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), for which there are no efficacious vaccines or therapeutics that are urgently needed. We expressed three versions of spike (S) proteins—receptor binding domain (RBD), S1 subunit and S ectodomain—in insect cells. RBD appears monomer in solutions, whereas S1 and S associate into homotrimer with substantial glycosylation. The three proteins confer excellent antigenicity with six convalescent COVID-19 patient sera. Cryo-electron microscopy (cryo-EM) analyses indicate that the SARS-CoV-2 S trimer dominate in a unique conformation distinguished from the classic prefusion conformation of coronaviruses by the upper S1 region at lower position ~15 Å proximal to viral membrane. Such conformation is proposed as an early prefusion state for the SARS-CoV-2 spike that may broaden the knowledge of coronavirus and facilitate vaccine development.

Key words: COVID-19, SARS-CoV-2, spike, cryo-electron microscopy, antigenicity, early prefusion conformation

Introduction

The novel coronavirus grouped in betacoronavirus genus has become the third serious virus intruder to human in the coronaviridae, after sever acute respiratory syndrome coronaviruses (SARS-CoV) and middle east respiratory syndrome coronavirus (MERS-CoV), recently named SARS-CoV-2. In the phylogenic tree of the coronaviruses, SARS-CoV-2 is genetically close to some bat coronavirus and SARS-CoV, however, with its origin undefined[1]. SARS-CoV-2 causative disease "Coronavirus disease 2019" (abbreviated "COVID-19") is characterized by high fever, dry cough, difficulty breathing and sever atypical pneumonia, which usually be confirmed by virus RNA positive or pulmonary computed tomography (CT) in clinical practice[2, 3]. In terms of higher human-to-human transmissibility, SARS-CoV-2 has spread over 118 countries and areas, and led to over 125,288 confirmed cases worldwide and at least 4,614 deaths, as of March 12th 2020. The World Health Organization (WHO) has declared the SARS-CoV-2 epidemic as a pandemic of international concern and updates the COVID-19 situation every day.

SARS-CoV-2 is an enveloped, single and positive-stranded RNA virus encapsulated with a genome of ~30 kb. At least three membrane proteins including the surface spike protein (S), an integral membrane protein (M), a membrane protein (E). Like other coronaviruses, S is responsible for initiating the engagement to a specific cellular receptor angiotensin-converting enzyme 2 (ACE2) and mediating the cell-virus membrane fusion by the class I fusion mechanism[4, 5]. Thus, S is the main target for neutralizing antibodies against viral infection and the core immunogen constituent of

vaccine design. S is consisted of S1 and S2 subunits and the cleavage on S1/S2

boundary by protease during biosynthesis is prerequisite for coronaviruses cellular

membrane fusion and subsequent infection[6]. SARS-CoV-2 evolves a 4-residue

insertion (RRAR) as potential furin cleavage site rather than SARS-CoV and other bat

coronaviruses, which may contribute to the higher transmissibility of this novel

coronavirus[6, 7]. Previous studies suggested the infection process of MERS-CoV[8] and

SARS[9] viruses, where S trimer undergoes conformational transition from a prefusion

conformation ready for ACE2 binding to a postfusion conformation for eventual virus-

cell membrane fusion. Structure determination of SARS-CoV and MERS-CoV spike

trimers captured a variety of scenarios in the prefusion conformation showing partial

(one or two) receptor-binding domain (RBD) in the "up" conformation and the rest in

the "down", and all in either "up" or "down". The conformation transition from "down"

to "up" could expose the receptor binding site, and the subsequent receptor engagement

would lead to a substantial conformation rearrangement of S trimer from prefusion

conformation to postfusion. Two recent studies[7, 10] reported cryo-electron microscope

(cryo-EM) structures of SARS-CoV-2 spike trimers in the prefusion conformation with

2 RBDs down and 1 RBD up. In the case of SARS-CoV, this conformational change

during RBDs "down" to "up" was associated with the binding of receptor ACE2 as well

as the recognition of neutralizing monoclonal antibodies [11].

A safe and efficacious vaccine is urgently needed to control and eliminate the

SARS-CoV-2 infection. Various forms of vaccine candidates, mostly aiming to elicit

neutralizing antibodies against S proteins, are under preclinical research or even

subjected to clinical trials[12]. Here, we cloned S ectodomain and its fragments RBD and

S1 into recombinant baculovirus and expressed the proteins in insect cells. We found

that S and S1 formed homotrimer in solutions and the three proteins reacted well with

COVID-19 convalescent human sera. Cryo-EM analysis demonstrated the S trimer

unexpectedly retains a unique conformation distinguished from the classic prefusion

conformation of coronavirus spikes, that may represent an early state rather than the

known prefusion conformation of S spike. These results might broaden the knowledge

on coronavirus virology and provide another protective conformation of S trimer for

structure-based vaccine design against SARS-CoV-2 infection and its causative

COVID-19.

Results

Construct design, expression and purification of SARS-CoV-2 S proteins

To screen a potent immunogen for COVID-19 vaccine development, we designed three

constructs—S ectodomain, S1 and RBD—for the SARS-CoV-2 Spike (S) protein

expression by aligning the SARS-CoV-2 S gene (Genbank accession no. NC_045512.2)

to a SARS-CoV strain (Genbank accession no. NC_004718) S gene sequence in terms

of structure-defined domain profile of the SARS-CoV S protein (Fig. 1A). The gene of

SARS-CoV-2 S ectodomain encoding amino acids (aa) 15-1,213 with removal of its

original signal sequence was cloned to the downstream of the gp67 signal sequence in

pAcgp67B plasmid vector (Fig.1B). and with its C-terminal addition of a thrombin

cleavage site, a T4 trimerization foldon motif and his tag. The segments S1 (aa 15-

680) and RBD (aa 319-541) were cloned similar to S ectodomain, keeping gp67 signal

peptide and his tag to facilitate secretory outside cell and affinity purification,

respectively, but without the thrombin site and T4 foldon (Fig. 1A). The three

constructed plasmids were respectively co-transfected into Sf9 insect cells with v-

cath/chiA gene deficient baculovirus DNA for the generation and amplification of

recombinant baculovirus, which were then harnessed to infect Hive Five insect cells to

eventually produce recombinant proteins, named S, S1 and RBD respectively.

The recombinant proteins were mostly soluble expressed and secreted into the

culture medium. The centrifugation supernatants of cell culture went through metal

affinity chromatography using Ni-NTA resin. S, S1 and RBD proteins were mainly

eluted in a separation fractions under 250 mM imidazole elution, and resolved as

molecular weight (m.w.) of ~180 kDa, 110 kDa and 35 kDa, respectively, in SDS-

PAGE as indicated by a corresponding western blotting (WB) using anti-His antibody

as detection antibody (Fig. 1C). Interestingly, about one half S proteins were cleaved

into S1 (identical migration site to S1 lane in Fig. 1C) and S2 (about 80kDa developed

in anti-His WB) possibly by innate furin of insect cell that was also found in other cases

of enzymatic cleave while protein expression in insect cell, such as Flu HA [13]. The

eluted S fraction was further polished by Superdex 200 to remove contaminative

proteins (Fig. 1D). These peaks fractionated at retention volume 28mL, 36mL, 48mL,

and 65mL, were further harvested and subjected to SDS-PAGE analysis. The results

indicated that S proteins together with cleaved S1/S2 were resolved at peak 1 in size-

exclusion chromatography (Fig. 1D) and showed a high purity of over 95% total

S/S1/S2 in gel (Fig. 1E). Overall, one-step Ni-NTA affinity chromatography produced

RBD with 95% purity and a yield of 30 mg per L cell culture, S1 with about 90% purity

and 10 mg per L yield, while further purification through a size-exclusion

chromatography (SEC), the resultant S sample had over 95% purity regarding intact S

and cleaved S1/S2, and was harvested in a yield of 1 mg per L cell culture. These data

set up a start point for further optimization on expression and purification process of

SARS-CoV-2 S immunogen candidates through insect baculovirus expression vector

system (BEVS).

Physiochemical properties of SARS-CoV-2 S-RBD, S1 and S proteins

We next investigated the physiochemical properties of the recombinant S protein and

its fragments purified from insect cells, including association potential, thermal

stability and glycosylation situation. Firstly, high pressure size-exclusion

chromatography (HPSEC) and sedimentation velocity analytical ultracentrifugation

(SV-AUC) analyses were carried out to measure the oligomerization potential of the

three proteins in solution. RBD, S1 and S all showed single major peak in HPSEC

profiles at elution volume of 9.0 mL, 5.5 and 5.3 mL, respectively (Fig. 2A and Fig.

2B). RBD, S1 and S were further verified by SV-AUC, where RBD sedimented as

single species of 3.1S in c(s) profile, corresponding to apparent molecular weight 22

kDa (Fig. 2D); S1 existed as a dominant species of 11.3 S (estimated as 277 kDa

corresponding to S1 trimer) and a minor aggregate form of 20 S (Fig. 2E); S and cleaved

S1/S2 resolved as 15.2 S, equivalent to 577 kDa, approximately as the theoretical

molecular weight of intact S trimer. The three proteins were further analyzed by

differential scanning calorimetry (DSC) that was usually used to investigate the inner

thermostability of macromolecules or their complexes[14]. RBD and S1 showed one

major peak at comparable thermal denaturation midpoints (Tm) of 46.0 °C and 45.5 °C,

respectively (Fig. 3G and 3H), whereas S sample showed two major peaks at Tm of

45.5°C (identical to Tm of S1) and 64.5°C (Fig. 3I), which might reflect the coexistence

of intact S and cleaved S1/S2.

On the other hand, we investigated the glycosylation extent of the three protein by

enzymatic deglycosylation analysis. Endo H could unleash the chithobiose core of high

mannose and some hybrid oligosaccharides from N-liked glycoproteins, therefore

remove the extended branches of glycans and leave the one N acetylglucosamine

(GlcNAc) on N-linked glycoproteins. While PNGase F would release N-linked glycan

moieties between GlcNAc and ASN residues within a glycoprotein. It should be noted

that glycosylation in insect cells is featured as terminal mannose glycans, unlike

complex sialylated glycans in mammalian cells, and glycosylation is known to correlate

the immunogenicity and broad-coverage protection of a glycoprotein immunogen[15, 16].

After the treatment of either Endo H or PNGase F, RBD showed no discernible decrease

of molecular weight in SDS-PAGE/anti-His WB, S1 and S2 both demonstrated nearly

~10 kDa decrease, and the intact S exhibited substantial shrinkage in molecular weight

of about ~20 kDa decrease (Fig. 2J). The analyses conclude that the glycosylation

extent within S glycoprotein is RBD < S1 ~ S2, consistent to the predicted glycosylation

profile of S polypeptide (Fig. 1A).

Reactivity of SARS-CoV-2 RBD, S1 and S proteins against convalescent COVID-19 human sera

We next evaluated the antigenicity of the three versions of S proteins by WB and ELISA using a panel of six COVID-19 convalescent human sera, which was collected from COVID-19 patients after they recovered from the disease in the First Affiliated Hospital of Xiamen University. Eight reducing SDS gel duplicates of the one depicted in Fig. 1C were prepared for WB analysis using these six convalescent sera and two control sera from health human (Fig. 3A-3H, left panel). As expected, intact S protein bands reacted well with all the six convalescent sera (Fig. 3A-3H, left panel). Unexpectedly, five of six sera showed no or very weak reactivities against RBD, only Serum #6 possessed RBD's activity. Among the five sera with lower RBD-reactivity, Serum #2, #3 and #5 well recognized S1 and the cleaved S1 in lane S, suggesting these sera may specifically react with NTD of S1. S2 demonstrated reaction activity against all the six sera, like the intact S. No detectable reaction was observed in the control sera (Fig. 3G and 3H). Inconsistent to the WB results, RBD, S1 and S shared comparable reactivities against the convalescent sera in ELISA, although the sera per se presented varied reaction titers (represented as ET50) following the reaction sequence: Serum #5 > #2 > #3 > #1 > # 6 > #4 (Fig. 3A-3H, right panel).

Taken together, RBD, S1 and S proteins from insect cells maintain the native-like SARS-CoV-2 epitopes. These epitopes in native virion should be immunogenic in COVID-19 patients and capable of eliciting high antibody titer in the convalescent

phase of SARS-CoV-2 infection. Among these epitopes, most RBD epitopes are strictly

tertiary conformation-dependent sites that are damaged upon the mild denatured

condition of reductant and SDS treatment, NTD within S1 bears some linear epitopes,

whereas S2 part essentially has linear epitopes that are immunogenic in all COVID-19

patients (n=6).

Cryo-EM structures of SARS-CoV-2 S proteins

To examine the structure of the trimeric S ectodomain with native sequence, we

prepared cryo-EM grids using the Ni-NTA purified S proteins and collected 1,513

electron micrograph movies. Most of motion-corrected micrographs demonstrated

plenty of well-dispersed particles with an approximate size as the canonical coronavirus

S trimer (Fig. 4A). A total of 162,645 particles were picked out for multiple rounds of

2D classification, consequently, 37,147 particles grouped into top 10 classes, rendering

typical feature of S trimer in prefusion conformation as recently reported[7, 10], were

selected for further analysis (Fig. 4B). 3D reconstruction (applying 3-fold symmetry)

yielded the density map of prefusion spike (S-pre) at resolution of 5.43 Å (Fig. 3D and

Supplementary Fig. 1A).

Structurally, three S monomers intertwine around each other and associate to

homotrimers with 145 Å height seen from side-view and 160 Å diameter in top-view

(Fig. 4C and 4D). We then recruited the recently reported cryo-EM map of S prefusion

trimer (EMD-21374, at resolution of 3.17 Å, low pass to 5.43 Å prior to structural

comparison) and compared our cryo-EM map at same resolution (Fig. 4D). It was

worthy noted that the compared prefusion SARS-CoV-2 S trimer was engineered with

site-directed mutations to stabilize prefusion conformation and expressed in 239F cells.

The mutant included two stabilizing proline mutations at residues 986, 987 and a

"GSAS" substitution at the furin cleavage site[7]. Surprisingly, the alignment

demonstrated that the two cryo-EM structures share similar mushroom-shaped

architecture in particular nearly identical at stalk moiety (S2 region), but our S-pre

shows the cap part (S1 region) at ~15Å lower position than the reported S trimer in

RBD-down prefusion conformation (Fig. 4D). Regarding to substantial mismatch at the

density of 3 S1 subunits, we respectively fitted 5 individual domains (NTD, RBD, SD1,

SD2 and S2) of the SARS-CoV-2 S structure (PDB code 6VSB) to our S-pre map. In

the fitting map, NTD, RBD, SD2 and S2 could be well placed in the S-pre map,

especially for the latter two, which reflects the aforementioned good match at the stalk

of the mushroom-shape (Supplementary Fig. 2). However, there is no observable

density between RBD and SD2 to accommodate an SD1 model (Supplementary Fig. 2),

which suggests SD1 region is dramatically flexible in our S-pre structure (Fig. 4E and

4F). When the combined model of fitted NTD-RBD-SD2-S2 was superimposed to the

original S protomer structure (PDB code 6VSB, Chain A, RBD in down conformation),

both NTD and RBD in the original S obviously move and rotate up against our

combined model (Fig. 4G). The structural comparison demonstrated that the S-pre

trimer retains a unique conformation different from the prefusion conformation of the

two reported SARS-CoV-2 spike structures (PDB codes 6VSB and 6VXX).

We then compared the conformation of our S-pre structure with that of 21

deposited coronavirus S models. Six representative S structures[7, 17-20] from four known

genus (α-, β-, γ- and δ-genus) in coronaviridae are respectively fitted to the S-pre map

(Supplementary Fig. 3). Five S trimer structures of other coronaviruses share similar

prefusion conformation with the reported SARS-CoV-2 S structure but substantially

distinct with the unique conformation of our S-pre.

Apart from most particles classified as S-pre in our sample, 2D classifications also

showed five classes of few particles (2,951) assuming an elongated rosette-shape

assembly. These particles were further reconstructed and yielded a structure at lower

resolution of 8.40 Å (Supplementary Fig. 1B) that could be considered as post-fusion

spike (S-post), as the structure has similar shape but shorter length (~170 Å) as

compared to the postfusion spike of SARS-CoV[21] and the presumed one observed in

native SARS-CoV-2 virion (BioRxiv, https://doi.org/10.1101/2020.03.02.972927)

(Supplementary Fig. 4) . Fitting the S-post map with the core region structure of SAR-

CoV-2 S2 subunit in post-fusion conformation (PDB code 6LXT) indicated that our S-

post exhibits roughly rod shape similar with the post-fusion structure (Supplementary

Fig. 4).

Conformational transition of SARS-CoV-2 spike from early prefusion to

postfusion

Briefly, we've obtained two conformations of SARS-CoV-2 spike from insect cells.

The dominant one maintains the similar mushroom-shaped trimer as the previous

models, while the S1 region substantially diverges. The other conformation essentially

resembles the postfusion state. However, we could not find the classic prefusion

conformation in our sample. We next tried to figure out at which stage the unique

conformation occurs during the spike conformation change. The space relationship of

NTD or RBD to S2 domain in the unique, RBD-down and RBD-up prefusion

conformations (Fig. 5A) was measured by a reference plant approximately parallel to

the viral membrane. The plane is defined by the positions of three equivalent Cα atoms

(residue 694 be used) from three S2 subunits of the trimer structures. Numerical data

shows that (1) the NTD and RBD in the unique conformation retain the lowest position

in the three prefusion conformations; (2) the NTD and RBD of RBD-down prefusion

stretch upward 16.6° rotation/16 Å elevation, and 13.1°/18 Å, respectively, with respect

to the unique conformation; (3) from RBD-down to RBD-up prefusion state, the RBD

elevates 9 Å with an additional rotation whereas the NTD remaining nearly stationary

(Supplementary Movie S1). The resultant "up" RBD is ready for ACE2 binding and

the spike eventually is rearranged to postfusion state upon RBD-ACE2 interaction[11, 22]

(Fig. 5B). The motion trend of NTD and RBD from prefusion to postfusion state in

conformational transition is away from the viral membrane, suggesting the unique

conformation may occur earlier than RBD-down prefusion conformation, named as

"early prefusion conformation" (Fig. 5). This early prefusion conformation might exist

in other coronaviruses as well.

Discussion

SARS-CoV-2 has crossed the species barrier and sweep over the planet by person-to-person transmission in an R0 ~2.56 rate[23], first wave in China and the second wave booming outside China. WHO has declared the event as another pandemic infectious disease in human history, and the epidemiology of SARS-CoV-2 infection is still in data accumulation. Although the biology and virology of SARS-CoV-2 remain elusive, in terms of knowledge on other coronaviruses, the spikes decorating the SARS-CoV-2 virion play a critical role in viral attachment and entry to host cells. Cryo-EM structures of spikes in the prefusion conformation, and RBD-bound receptor ACE2 have indicated the engagement of SARS-CoV-2 to cellular membrane requires a serial of conformational change of RBDs. The change is presumed from the start point of 3 RBDs down in the prefusion conformation, then RBD(s) up for ACE2 binding, and eventually spike is rearranged to postfusion. In this study, we suggest that the SARS-CoV-2 spike may retain at more precedent state than the classic prefusion conformation that has been determined for other coronaviruses. This early prefusion conformation features that the cap of the mushroom-shaped spike constituted by three S1 subunits is more proximal to viral membrane by 15 Å than in the classic prefusion conformation.

The SARS-CoV-2 spike expressed in insect cells predominantly retains a unique early prefusion conformation, which was repeatable in at least three batches of samples and is ascribed to two possible reasons – native aa sequence used in the S ectodomain construct and over-expression in insect cells. There is about a half of S proteins undergoing cleavage on the S1/S2 boundary site in the purified samples both after the

first Ni-NAT and the second SEC purification (Fig. 1C and 1E). Further analyses

suggest the split between S1 and S2 likely takes no effect on the trimerization of S

trimer. It is known that the insect cells can confer post-translation glycosylation for

protein over-expression as mammalian cells despite the latter can produce more

complex sialyation[24], and thus provide an alternative way to generate glycoprotein in

native conformation. Our results indicate that RBD, NTD and S2 domains of SARS-

CoV-2 demonstrated different glycosylation extent in insect cells, however, RBD, S1

and S proteins comparably react well with six convalescent COVID-19 human sera

albeit they differ in domain composition, polypeptide length and oligomerization.

There are numbers of SARS-CoV-2 vaccine candidates, including inactivated,

vectored, recombinant and nucleotide vaccine forms, under preclinical research.

Various versions of S proteins are the major targets for vaccine immunogen candidate.

In addition to potent neutralizing antibody elicitation upon immunization, potential

antibody-dependent disease enhancement (ADE) is the major concern for an efficacious

SARS-CoV-2 vaccine. ADE has been found in the development of numbers of virus

vaccine candidates, including respiratory syndrome virus (RSV), dengue fever [25, 26],

human immunodeficient virus (HIV), SARS-CoV, MERS-CoV [25-27] and so on. It is

believed that ADE is associated with non-neutralization epitope attribute and / or

specific antibody isotype[28, 29], in which virus-bound antibody would promote the viral

infection to immune cells through Fc fragment targeting γFc receptors on the cellular

surface and enhance the disease severity. Therefore, the strategy of vaccine design

against SARS-CoV-2 should include the consideration of antigen region selection,

glycosylation number/extent and exactly presented prefusion conformation. The

prefusion conformation needed to be maintained is exemplified by the case of RSV

vaccine candidate in which F trimer in prefusion is much potent than postfusion. Hence

the early prefusion conformation proposed for SARS-CoV-2 spike should be drawn an

attention for immunogen design as well as the prefusion one.

In conclusion, we obtain three kinds of S proteins showing excellent antigenicity

and find an early prefusion conformation for SARS-CoV-2 spike. Nevertheless, the

molecular level detail for such conformation and the underlying immunogenicity

should be further investigated, and whether this conformation recapitulates the exact

state of spike in native SARS-CoV-2 virion remains to be determined.

Materials and Methods

Cloning, protein expression and purification

The SARS-CoV-2 S gene (Genbank accession no. NC_045512.2) was synthesized and

cloned into a baculovirus shuttle vector pAcgp67B (BD Biosciences, CA, USA) using

Gibson assembly. The S construct encoding aa 15-1,213 (numbered as original

sequence), contains a thrombin site, a T4 foldon domain to assist in trimerization and a

C-terminal 10-His tag for purification. For S1 construct contains gene encoding aa 15-

680 followed by a 10-His tag. The RBD construct (aa 319-541) also contains 10-his tag

to facilitate purification. In all three constructs, the natural signal peptide (aa 1-14

analyzed by SignalP tool) was replaced with a gp67 secretion signal peptide at N-

terminus.

The expression and purification of proteins were performed as described

previously[30]. All plasmids were co-transfected with linearized 2.0 DNA (deficient in *v-*

cath/chiA genes) (Expression Systems, CA, USA) into *Sf9* insect cells (Thermo Fisher

Scientific, MA, USA), according to the protocol provided by the manufacturer

(Expression Systems). The transfection supernatant was harvested and amplified 2

times to obtain a high titer of the recombinant viruses. Hive Five cells (BTI-TN-5B1-

4) (Thermo Fisher Scientific) were cultured in ESF921 medium (Expression Systems)

and infected with recombinant virus at an multiplicity of infection (MOI) of 5 in the

exponential growth phase (2×10^6 cells/ml, 95% viability) at 28°C for 72 h. The culture

media was centrifuged at 8,000 rpm for 20 min. Then the supernatant was dialyzed

against phosphate-buffered saline (PBS), pH 7.4, and purified with Ni-sepharose fast

flow 6 resin (GE Healthcare, Boston, USA) by the elution with 250 mM imidazole. The

protein concentrations of the final purified samples were measured with Pierce™BCA

Protein Assay Kit (Thermo Fisher Scientific).

SDS-PAGE and western blot

Protein samples were mixed with loading buffer and boiled for 10 min, and subjected

to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Equal

amounts of proteins for each sample were loaded onto two SDS-PAGE gels, one for

western blotting and one for Coomassie staining. The proteins were electrophoresed for

70 min at 80 V in a BioRad MINI-PROTEAN Tetra system (BioRad Laboratories, CA,

USA), and the gel was stained with Coomassie Brilliant Blue R-250 (Bio-Rad) for 30

min at room temperature. For western blotting, separated proteins were transferred onto

a nitrocellulose membrane (Whatman, Dassel, Germany) using a Trans-Blot Turbo

transfer system (Bio-Rad). The membrane was blocked and then incubated for 1 h with

an His-tag-specific mouse mAb antibody (Proteintech, Rosemont, USA) or human sera

(1:500 dilution). Unbound antibody was removed by five 5-min washes and the

membrane was incubated with alkaline phosphatase-conjugated goat anti-mouse

secondary antibody or goat anti-human IgG secondary antibody (Abcam, Cambridge,

UK). Membranes were washed again and then developed using SuperSignal ELISA

Pico Chemiluminescent Substrate Kit (Thermo Fisher Scientific).

Enzyme-Linked Immunosorbent Assay (ELISA)

Purified proteins were coated onto 96-well microtiter plates at 100 ng/well in PBS at

37°C for 4 h. The background was blocked with 1 × Enzyme dilution buffer (PBS +

0.25% casein + 1% gelatin + 0.05% proclin-300) at 37°C for 2 h. Sera were diluted

started at 1:100 followed with three-fold serially dilution, and added to the wells (100 µl)

and incubated at 37°C for 1 h. Horseradish peroxidase (HRP)-labeled mouse anti-

human antibody (Abcam) was used as secondary antibody at 1:5,000 for 30 min. Wells

were washed again and the reaction catalyzed using o-phenylenediamine (OPD)

substrate at 37°C for 10 min. The OD450nm (reference, OD620nm) was measured on

a microplate reader (TECAN, Männedorf, Switzerland), with a cut-off value 0.1. The

Half effective titers (ET50) was calculated by sigmoid trend fitting using GraphPad

Prism software.

Size-Exclusive Chromatography (SEC)

Ni-NTA purified S proteins were further loaded into Superdex200 (GE Healthcare), the fractions were harvested and analyzed by SDS-PAGE. All high-purity RBD, S1 and S proteins were subjected to HPLC (Waters; Milford, MA) analysis using a TSK Gel G5000PWXL7.8 × 300 mm column (TOSOH, Tokyo, Japan) equilibrated in PBS, pH 7.4. The system flow rate was maintained at 0.5 mL/min and eluted proteins were detected at 280 nm.

Analytical Ultracentrifuge (AUC)

The AUC assay was performed using a Beckman XL-Analytical ultracentrifuge (Beckman Coulter, Fullerton, CA), as described elsewhere[31]. The sedimentation velocity (SV) was carried out at 20°C with diluted proteins in PBS. The AN-60 Ti rotor speed was set to 20,000-30,000 rpm according to the molecular weight of the control proteins. Data was collected using SEDFIT computer software, kindly provided by Dr. P. C. Shuck (NIH, Bethesda, MA, USA). Multiple curves were fit to calculate the sedimentation coefficient (S) using continuous sedimentation coefficient distribution model [c(s)], and then the c(s) used to estimate protein molar mass.

Differential scanning calorimetry (DSC)

Differential scanning calorimetry (DSC) was carried out on the S proteins using a MicroCal VP-DSC instrument (GE Healthcare, MicroCal Products Group, Northampton, MA) as described previously[14]. In brief, all samples with a concentration

of 0.2 mg/mL were measured at a heating rate of 1.5°C /min with the scan temperature

ranging from 10°C to 90°C. The melting temperatures (Tm) were calculated using

MicroCal Origin 7.0 (Origin-Lab Corp., Northampton, MA) software assuming a non-

two-state unfolding model.

Endo-H and PNGase-F digestion

The Endo-H (NEB) and PNGase-F (NEB) digestions were performed according to the

protocol offered by instruction. In brief, the deglycosylation reactions were carried out

using 10ug S proteins with 5uL of Endo H or PNGase F and incubated at 37°C

overnights. The reactions were loaded in to SDS-PAGE and analyzed by Western

blotting using anti-His as detecting anybody.

Cryo-EM sample preparation and data collection.

Aliquots (3 µL) of 0.5 mg/mL purified SARS-CoV-2 S protein were loaded onto glow-

discharged (60 s at 20 mA) holey carbon Quantifoil grids (R1.2/1.3, 200 mesh,

Quantifoil Micro Tools) using a Vitrobot Mark IV (ThermoFisher Scientific) at 100%

humidity and 4°C. Data were acquired using the EPU software to control a FEI Tecnai

F30 transmission electron microscope (ThermoFisher Scientific) operated at 300 kV.

and equipped with a ThermoFisher Falcon-3 direct detector. Images were recorded in

the 58-frame movie mode at a nominal magnification of 93,000X with a pixel size of

1.12 Å. The total electron dose was set to 46 e^- $Å^{-2}$ and the exposure time was 1.5 s.

537 micrographs were collected with a defocus range comprised between 1.5 and 2.8

μm.

Cryo-EM data processing

Movie frame alignment and contrast transfer function estimation of each aligned

micrograph were carried out with the programs Motioncor[32] and Gctf[33]. Particles were

picked by the 'Templete picker' session of cryoSPARC v2[34]. Two rounds of reference-

free 2D classification were performed and well-defined particle images were selected

and non-uniform 3D refinement, 3D reconstruction with C3 symmetry were performed

using cryoSPARC v2. The resolutions of the final maps were estimated on the basis of

the gold-standard FSC curve with a cutoff at 0.143[35]. Density-map-based visualization

and segmentation were performed with Chimera[36].

Acknowledgments

448 This work was supported by grants from the National Natural Science Foundation

449 (grant no. U1705283, 31670935, 81971932, 81991491, 31730029), and the Major

450 Project of Fujian Provincial Science Foundation for COVID-19 Research (Grants

451 2020YZ014001).

Financial Disclosure

454 The funders had no role in study design, data collection and analysis, decision to publish,

455 or preparation of the manuscript.

Competing Interest

458   The authors have declared that no competing interests exist.

Author Contributions

Y.G, S.L. and N.X. designed the study. T.L., Q.Zheng., H.Y., D.W., W.X., Y.Z.,

X.H., L.Z., Z.Zhang., Z.Zhai., T.C., Z.W., J.C., H.S. and T.D. performed experiments.

T.L., Q.Z., H.Y., Y.W., Y.C., Q.Zhao., J.Z., Y.G., S.L. and N.X. analyzed data. T.L.,

Q.Z., H.Y., Y.G., and S.L. wrote the manuscript. T.L., Q.Zheng., H.Y., D.W., W.X.,

Q.Zhao., J.Z. Y.G., S.L., and N.X. participated in discussion and interpretation of the

results. All authors contributed to experimental design.

References

1 Chan JF, Kok KH, Zhu Z *et al.* Genomic characterization of the 2019 novel human-pathogenic
coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes*
Infect 2020; **9**:221-236.
2 Zhu N, Zhang D, Wang W *et al.* A Novel Coronavirus from Patients with Pneumonia in China,
2019. *N Engl J Med* 2020; **382**:727-733.
3 Wang D, Hu B, Hu C *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel
Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020.
4 Zhou P, Yang XL, Wang XG *et al.* A pneumonia outbreak associated with a new coronavirus of
probable bat origin. *Nature* 2020; **579**:270-273.
5 Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by novel coronavirus from Wuhan:
An analysis based on decade-long structural studies of SARS. *J Virol* 2020.
6 Millet JK, Whittaker GR. Host cell proteases: Critical determinants of coronavirus tropism and
pathogenesis. *Virus Research*; **202**:120-134.
7 Wrapp D, Wang N, Corbett KS *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science (New York, NY)* 2020.
8 Walls AC, Xiong X, Park Y-JJ *et al.* Unexpected Receptor Functional Mimicry Elucidates Activation
of Coronavirus Fusion. *Cell* 2019; **176**:1026.
9 Yuan Y, Cao D, Zhang Y *et al.* Cryo-EM structures of MERS-CoV and SARS-CoV spike

glycoproteins reveal the dynamic receptor binding domains. *Nature communications* 2017;
8:15092.

10 Walls AC, Park Y-JJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and
Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* 2020.

11 Gui M, Song W, Zhou H *et al.* Cryo-electron microscopy structures of the SARS-CoV spike
glycoprotein reveal a prerequisite conformational state for receptor binding. *Cell research* 2017;
27:119-129.

12 Lu S. Timely development of vaccines against SARS-CoV-2. *Emerg Microbes Infect* 2020; **9**:542-
544.

13 Sui J, Hwang WC, Perez S *et al.* Structural and functional bases for broad-spectrum
neutralization of avian and human influenza A viruses. *Nature structural & molecular biology* 2009;
16:265-273.

14 Zhang X, Wei M, Pan H *et al.* Robust manufacturing and comprehensive characterization of
recombinant hepatitis E virus-like particles in Hecolin(®). *Vaccine* 2014; **32**:4039-4050.

15 Medina RA, Stertz S, Manicassamy B *et al.* Glycosylations in the globular head of the
hemagglutinin protein modulate the virulence and antigenic properties of the H1N1 influenza
viruses. *Science translational medicine* 2013; **5**.

16 Liu W-CC, Jan J-TT, Huang Y-JJ, Chen T-HH, Wu S-CC. Unmasking stem-specific neutralizing
epitopes by abolishing N-linked glycosylation sites of influenza hemagglutinin proteins for vaccine
design. *Journal of virology* 2016.

17 Walls AC, Tortorici MA, Frenz B *et al.* Glycan shield and epitope masking of a coronavirus spike
protein observed by cryo-electron microscopy. *Nat Struct Mol Biol* 2016; **23**:899-905.

18 Yuan Y, Cao D, Zhang Y *et al.* Cryo-EM structures of MERS-CoV and SARS-CoV spike
glycoproteins reveal the dynamic receptor binding domains. *Nature communications* 2017;
8:15092.

19 Shang J, Zheng Y, Yang Y *et al.* Cryo-EM structure of infectious bronchitis coronavirus spike
protein reveals structural and functional evolution of coronavirus spike proteins. *PLoS Pathog* 2018;
14:e1007009.

20 Shang J, Zheng Y, Yang Y *et al.* Cryo-Electron Microscopy Structure of Porcine Deltacoronavirus
Spike Protein in the Prefusion State. *J Virol* 2018; **92**.

21 Song W, Gui M, Wang X, Xiang Y. Cryo-EM structure of the SARS coronavirus spike glycoprotein
in complex with its host cell receptor ACE2. *PLoS Pathog* 2018; **14**:e1007236.

22 Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of the SARS-
CoV-2 by full-length human ACE2. *Science (New York, NY)* 2020.

23 Zhao S, Musa SS, Lin Q *et al.* Estimating the Unreported Number of Novel Coronavirus (2019-
nCoV) Cases in China in the First Half of January 2020: A Data-Driven Modelling Analysis of the
Early Outbreak. *Journal of clinical medicine* 2020; **9**.

24 Palomares LA, Srivastava IK, Ramírez OT, Cox MMJMJ. Glycobiotechnology of the Insect Cell-
Baculovirus Expression System Technology. *Advances in biochemical engineering/biotechnology*
2018.

25 Wang SF, Tseng SP, Yen CH *et al.* Antibody-dependent SARS coronavirus infection is mediated
by antibodies against spike proteins. *Biochem Biophys Res Commun* 2014; **451**:208-214.

26 Wan Y, Shang J, Sun S *et al.* Molecular Mechanism for Antibody-Dependent Enhancement of
Coronavirus Entry. *J Virol* 2020; **94**.

27 Kam YW, Kien F, Roberts A *et al.* Antibodies against trimeric S glycoprotein protect hamsters
against SARS-CoV challenge despite their capacity to mediate FcgammaRII-dependent entry into
B cells in vitro. *Vaccine* 2007; **25**:729-740.

28 Dejnirattisai W, Supasa P, Wongwiwat W *et al.* Dengue virus sero-cross-reactivity drives
antibody-dependent enhancement of infection with zika virus. *Nature immunology* 2016;
17:1102-1108.

29 Rey FAA, Stiasny K, Vaney M-CC, Dellarole M, Heinz FX. The bright and the dark side of human
antibody responses to flaviviruses: lessons for vaccine design. *EMBO reports* 2018; **19**:206-224.

30 Li T, Zhang Z, Zhang Z *et al.* Characterization of native-like HIV-1 gp140 glycoprotein
expressed in insect cells. *Vaccine* 2019; **37**:1418-1427.

31 Li Z, Wang D, Gu Y *et al.* Crystal Structures of Two Immune Complexes Identify Determinants
for Viral Infectivity and Type-Specific Neutralization of Human Papillomavirus. *mBio* 2017; **8**.

32 Zheng SQ, Palovcak E, Armache J-P, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic
correction of beam-induced motion for improved cryo-electron microscopy. *Nature methods*
2017; **14**:331.

33 Zhang K. Gctf: Real-time CTF determination and correction. *Journal of structural biology* 2016;
193:1-12.

34 Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for rapid unsupervised
cryo-EM structure determination. *Nat Methods* 2017; **14**:290-296.

35 Kucukelbir A, Sigworth FJ, Tagare HD. Quantifying the local resolution of cryo-EM density maps.
Nature methods 2014; **11**:63.

36 Pettersen EF, Goddard TD, Huang CC *et al.* UCSF Chimera—a visualization system for
exploratory research and analysis. *Journal of computational chemistry* 2004; **25**:1605-1612.

Figure 1. Schematic map of the SARS-CoV-2 S protein constructs. (A) Linear

representations of the S protein primary structure and construct design. NTD, N-

terminal domain; RBD, receptor binding domain; SD1, subdomain 1; SD2, subdomain

2; HR1, heptad repeat 1; CH, central helix; CD, connector domain; HR2, heptad repeat

2; TM, transmembrane domain; CT, cytoplasmic tail; FD, T4 foldon motif. The

predicted glycosylation sites are indicated above the domain bars. (B) Map of the

cloning vector pAcgp67B. The interest genes were cloned to plasmid pAcgp67B at

BamH I/Not I site to generate transfer vectors. (C) SDS-PAGE and western blotting of

the Ni-NTA purified proteins. RBS, S1 and S were eluted by 250 mM imidazole. Anti-

His antibody was used as detection antibody in western blotting. (D) Size-exclusion

chromatogram of the second-step purification of the S protein. (E) SDS-PAGE of the

four fractions harvested from the chromatography purification as shown in (D).

Figure 2. Characterization of the purified RBD, S1 and S proteins. (A-C) HPSEC

profiles of the purified RBD, S1 and S proteins; (D-F) AUC profiles of RBD, S1 and

S proteins; (G-I) DSC profiles of RBD, S1 and S proteins. (J) Western blotting of

three purified proteins treated with Endo H and PNGase F or untreated as control.

Anti-His antibody was used as detection antibody in western blotting.

Figure 3. Antigenicity of RBD, S1 and S proteins against convalescent sera. (A-F)

The reactivity of the RBD, S1 and S proteins against six COVID-19 convalescent

human sera (#1-#6) by western blotting (left panel) and ELISA (Right panel). (G, H)

Results of two control sera. The gels used for western blotting were duplicates of the

reducing SDS gels same as Fig. 1C.

Figure 4. Cryo-EM structure of the SARS-CoV-2 S trimer. (A) Representative

micrograph of frozen-hydrated SARS-CoV-2 S particles, Scale bar 50nm. (B, C) Ten and five

selected class averages showing the particles along different orientations belonging to prefusion (B)

and postfusion (C) S protein, respectively. (D) 5.43 Å density map of prefusion S trimer (S-pre) that

is colored by protomer. (E) Structural comparison to the reported prefusion SARS-CoV-2 S trimer

(EMD-21374, C3 symmetry, low-pass to 5.43Å) show a different conformation of S-pre (~15Å

shorter in height). (F, G) Each domain of the model of prefusion SARS-CoV-2 S monomer (F) or

trimer (G) (PDB no. 6VSB) were separately fitted in the density map of S-pre. (H) Schematic

592   diagram shows conformational diversities between NTD and RBD of S-pre (NTD: red, RBD:

593   orange) and reported prefusion S (NTD: light green, RBD: light purple).

Figure 5. SARS-CoV-2 S spike at early pre-, pre- and post-fusion states and the

proposed timeline for conformation change. (A) Four conformations of SARS-CoV-

2 S spike. For clarity, only one monomer at early prefusion, RBD down prefusion and

RBD up prefusion conformation is shown in ribbon mode. The model fitted in our S-

pre map is a combined model comprised of individually fitted NTD, RBD, SD2 and S2

domain, which were evicted from the model of prefusion S trimer (PDB no. 6VSB). A

deposited postfusion core of SARS-CoV-2 S2 subunit (PDB no. 6LXT) is fitted to our

S-post map. Domains are designated by different colors, red: NTD, orange: RBD,

yellow: SD1, green: SD2 and cyan: S2. (B) Simplified schematic diagram of the S

monomer interpreting the conformation change across different states.

Supplementary Information

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Movie 1