Selection and hybridization shaped the Africanized honey bee invasion of the Americas

Erin Calfee[1*], Marcelo Nicolás Agra[2], María Alejandra Palacio[2], Santiago R. Ramírez[1], Graham Coop[1]

1 Department of Evolution and Ecology and Center for Population Biology, University of California, Davis
2 Instituto Nacional de Tecnología Agropecuaria (INTA), Balcarce, Argentina

* ecalfee@ucdavis.edu

Abstract

Recent biological invasions offer 'natural' laboratories to understand the genetics and ecology of adaptation, hybridization, and range limits. One of the most impressive and well-documented biological invasions of the 20th century began in Brazil with the escape of introduced African honey bees (*Apis mellifera scutellata*) in 1957. In less than 50 years, populations of 'Africanized' honey bees spread across much of the Americas, hybridizing with and outcompeting resident European honey bees. We use broad geographic sampling and whole genome sequencing of over 300 bees to map the distribution of African ancestry where the northern and southern invasions have presently stalled, forming replicated hybrid zones between Africanized and European bees in California and Argentina. California is much farther from Brazil, yet these hybrid zones occur at very similar latitudes, consistent with the invasion having reached a climate barrier. At these range limits, we observe genome-wide clines for African ancestry, and parallel clines for wing length that span hundreds of kilometers, supporting a smooth transition from climates favoring Africanized bees to climates where they cannot survive winter. We find no large effect loci maintaining exceptionally steep ancestry transitions. Instead, we find most individual loci have concordant ancestry clines across South America, with a build-up of somewhat steeper clines in regions of the genome with low recombination rates, consistent with many loci of small effect contributing to climate-associated fitness trade-offs. Additionally, we find no substantial reductions in genetic diversity associated with rapid expansions nor complete dropout of African ancestry at any individual loci on either continent, which suggests that the competitive fitness advantage of African ancestry at lower latitudes has a polygenic basis and that Africanized bees maintained large population sizes during their invasion. To test for parallel selection across continents, we develop a null model that accounts for drift in ancestry frequencies during the rapid expansion. We identify several peaks within a larger genomic region where selection has pushed African ancestry to high frequency hundreds of kilometers past the present cline centers in both North and South America and that may underlie high-fitness African traits driving the invasion.

Author Summary

Crop pollination around the world relies on local honey bee populations, which vary in their behaviors and climatic ranges. Africanized honey bees ('killer bees') have been

some of the most widely successful; originating in a 1950s experimental breeding program in Brazil, they rapidly came to dominate across most of the Americas. As a recent genetic mixture of imported African bees and European subspecies, Africanized honey bees have a patchwork of ancestry across their genomes, which we leverage to identify loci with an excess of African or European ancestry due to selection. We additionally use the natural replication in this invasion to compare outcomes between North and South America (California and Argentina). We identify several genomic regions with exceptionally high African ancestry across continents and that may underlie favored Africanized bee traits (e.g. Varroa mite resistance). We find evidence that a climatic barrier has dramatically slowed the invasion at similar latitudes on both continents. African-to-European ancestry transition zones span hundreds of kilometers where bees have intermediate African ancestry proportions, which can be used to map the genetic basis of segregating traits (here, wing length), and calls into question the biological basis for binary Africanized vs. European bee classifications.

Introduction ₁

Diverging lineages often spread back into secondary contact before reproductive ₂ isolation is complete, and so can hybridize. In hybrid zones, multiple generations of ₃ admixture and backcrossing create a natural experiment in which genetic variation is ₄ 'tested' in novel ecological and genomic contexts. The mosaic of ancestries in hybrid ₅ zones has allowed researchers to uncover the genetic loci associated with reproductive ₆ barriers (e.g. [1–3]) and to identify rapidly introgressing high-fitness alleles (e.g. [4–7]). ₇ One promising way forward is to compare ancestry patterns across multiple young ₈ hybrid zones and test how repeatable the outcome of hybridization is across these ₉ evolutionary replicates. ₁₀

In this study, we use this powerful comparative framework to better understand the ₁₁ genomic basis of fitness and range limits in the Africanized honey bee invasion, with ₁₂ replicate hybrid zones in North and South America. In 1957, newly introduced honey ₁₃ bees of African ancestry (*Apis mellifera scutellata*) escaped from an experimental ₁₄ breeding program in Brazil and rapidly dispersed. Invading African bees both ₁₅ out-competed and hybridized with European-ancestry populations, creating a rapidly ₁₆ advancing admixed population of 'Africanized' bees that expanded north and south ₁₇ across the Americas at 300-500 km/year [8]. ₁₈

The highly defensive behavior of Africanized honey bees, and the implications for ₁₉ commercial bee management and public health, sparked immediate international ₂₀ concern over the outcome of this invasion. Control efforts have been largely unsuccessful ₂₁ in preventing Africanized honey bees from permeating feral and commercial colonies ₂₂ within their expanding range [8]. However, even without intervention, Africanized honey ₂₃ bees are unlikely to outcompete European bees in the coldest regions of the Americas ₂₄ because of their low overwinter survival in climates where European bees thrive [9, 10]. ₂₅ Models based on winter temperatures and the physiological cost of thermo-regulation ₂₆ predict northern range limits that vary from the Central Californian Coast [11, 12] up to ₂₇ the border with Canada [13]. Thus, the expected impact of Africanized honey bees on ₂₈ agriculture and queen bee production in the United States is still poorly defined. Broad ₂₉ surveys show that African mtDNA and Africanized phenotypes are common in northern ₃₀ Argentina and the southern US, and drop off towards more temperate latitudes, ₃₁ indicating that the rapid spread of these traits has dramatically slowed, if not stopped, ₃₂ on both continents [11, 14–19]. However, we lack a genome-wide view of the range limits ₃₃ of Africanized ancestry and do not know whether individual high-fitness alleles have ₃₄ already introgressed into higher latitudes. ₃₅

A number of studies have corroborated the historical record of this trans-American ₃₆

invasion at the genomic level, showing that honey bees in the Americas are historically admixed between predominately eastern European (C) and western European (M) subspecies, while the recently introduced Africanized honey bees are predominantly a genetic mixture of African (A) and European C & M subspecies [20–24]. A recent study of Africanized bees in Brazil revealed some European alleles at exceptionally high frequency that may have contributed to the success of the invasion, but this work was under-powered to detect high-fitness African alleles due to elevated genome-wide African ancestry (84%) in the Brazilian population [25].

While the key genes remain unknown, bees with African ancestry diverge from European-ancestry bees on a number of traits that may have given Africanized honey bees a selective advantage during the invasion: they defend their colonies more aggressively [26, 27], they have higher reproductive rates (including faster development times, proportionally higher investment in drone production and more frequent swarming to found new colonies [8]), they have higher tolerances to several common pesticides [28], and they prove less susceptible to *Varroa* mites, a major parasite [29–33]. Population monitoring studies show that *Varroa* mites are a strong selective force in the wild and that mite infestations in the 1990s likely contributed to the rapid genetic turnover of feral nest sites from European to Africanized colonies in Arizona and Texas [16, 17, 24].

There are also a number of candidate traits that differ between African and European bees and plausibly contribute to a climate-based range limit for the invasion. Smaller bodies [34] and higher metabolic rates [13], for example, could give Africanized honey bees a competitive advantage in the tropics but come at a cost in cooler climates [12]. In addition to physiological traits, heritable behaviors may also contribute to fitness trade-offs: African ancestry bees preferentially forage for protein-rich pollen (vs. nectar), which supports rapid brood production, but risks insufficient honey stores to thermo-regulate over winter [12, 35, 36].

Here we conduct the first comparative study of the Africanized honey bee invasion in North and South America. First, we use broad geographic sampling and whole genome sequencing to map the present-day ancestry clines on both continents, and assess the evidence for a climatic barrier preventing the further spread of Africanized honey bees. Next, we use genetic diversity within African ancestry to study the shared bottleneck within and amongst populations due to the rapid expansion during the invasion. Finally, we develop a null model that includes recent drift and use this model to test for outlier loci that may underlie high-fitness African traits and climatic barriers.

Results

To survey the current geographic distribution of African ancestry in the Americas, we sampled and sequenced freely foraging honey bees across two latitudinal transects, one in California and one in Argentina, formed from the northern and southern routes of invasion out of Brazil (Fig 1). We generated individual low-coverage whole-genome sequence data for 278 bees, and added to this data set 35 recently published high-coverage bee genomes from 6 additional California populations sampled 3-4 years prior [22]. We inferred genome-wide ancestry proportions for each individual using NGSAdmix [37] assuming a model of 3 mixing populations, which clearly map to the African (A), eastern European (C), and western European (M) reference panels (Figs 1 and S2). We leveraged the fact that admixed Africanized honey bee populations were formed through a recent mixture of well-defined honey bee subspecies to infer the mosaic of African (A) and European (C & M) ancestry tracts across the genome of each bee. For each population, we applied a hidden Markov model that jointly infers the maximum likelihood single-pulse approximation for the generations since mixture and

posterior probabilities for local ancestry state, based on read counts from low-coverage sequence data (ancestry_hmm [38]). The average local ancestry estimates within individuals agree closely with the NGSAdmix genome-wide ancestry estimates (Fig S3, Pearson's $r \geq 0.985$), with the HMM estimating slightly higher minor ancestry for low admixture proportions, likely as a result of some miscalled blocks. Time estimates vary by population, with a median of 47.6 generations in the 62 years since the initial dispersal of Africanized bees out of São Paulo (see Fig S5 for all time estimates). In this section, we first focus on the distribution of genome-wide 'global' ancestry patterns across the two clines, which we will later compare to the variation in local ancestry at individual loci.

Fig 1. Spread of Africanized honey bees. Map of hybrid zones in California and Argentina, with cartoon arrows depicting the two routes of Africanized honey bee invasion out of Rio Claro, São Paulo, Brazil. Dates of first occurrence along the routes of invasion are from [8] [39] and [40], with approximate GPS locations extracted from google maps. Insets zoom in on each hybrid zone to show the mean GPS coordinates for each sampled population. Sampling spanned 646km in California and 878km in Argentina in the north-south direction. Genome-wide A, C, and M ancestry inferred using NGSAdmix for each bee are shown in a bar chart at the bottom, where each vertical bar is one bee and colors indicate proportion ancestry. Populations are arranged by latitude, with samples closest to Brazil on the right. Light fading indicates that a bee comes from the previously published California data set [22] and was collected in the field 3-4 years prior to the bees from this study. These earlier California samples include one island population, Avalon (Catalina Island), indicated by a yellow triangle. Bees from Avalon have majority M ancestry, in contrast to all mainland California bees which have predominantly A and C ancestry.

We observe wide hybrid zones mirrored in North and South America. In Argentina, we find the cline in ancestry spans nearly 900km, from 77% A ancestry in the north to less than 5% to the south in Buenos Aires Province. The current geographic range of Africanized honey bees in South America is broadly consistent with prior studies using a smaller number of genetic markers (e.g. [14, 15, 23, 41]), though the geographic and genetic resolution of these studies is too limited for detailed comparison. In North America, we find that honey bees in California have up to 42% African ancestry in the south, tapering down to approximately 0% in Davis, our northernmost sampling site. In comparison, earlier extrapolations based on mitochondrial surveys may have somewhat overestimated genome-wide African ancestry in California (e.g. 65% of foraging bees in San Diego County [18] and 17% in Monterey County [19] carry African mtDNA haplotypes). We also find excess A-like mtDNA diversity in California. While this finding is potentially consistent with African maternal lines being favored during the expansion into Southern California, this pattern is not strongly replicated in South America and even in North America, African mitochondria do not appear to have introgressed far past the northern range limit for nuclear African ancestry (Fig S26).

Alongside our genomic cline, we find a corresponding phenotypic cline in worker fore wing size: closer to the equator, sampled bees have increasing African ancestry and shorter wings (Fig 2). By fitting a linear model to predict wing length from genome-wide ancestry, we find that African ancestry can explain a difference of -0.72mm, approximately an 8% reduction in wing length ($P = 3.65 \times 10^{-23}, R^2 = 0.31, n = 269$; see Fig S7). We tested for a main effect and an interaction term for the South American continent, and found no significant differences in wing length ($P = 0.81$) or its association with ancestry ($P = 0.86$) between the two clines. Thus, in contrast to the rise of dispersal-enhancing traits in other recent invasions (e.g. [42–45]), we see no evidence of a bias for longer wings at larger dispersal distances (California). Genetic crosses have shown that wing length differences between ancestries have a genetic basis [27] and the wing length patterns we observe here are consistent with expectations of an additive polygenic cline based on genome-wide ancestry alone (Fig 2). However, these phenotypic clines could alternatively be caused purely by developmental plasticity or sorting of within-ancestry genetic variation along a latitudinal gradient. Preliminary evidence that other factors may contribute to the wing length clines observed here comes from a 1991 survey showing that wing length was positively correlated with latitude in California's feral bee populations before the reported arrival of Africanized honey bees [46]. From field-based sampling alone, it remains unclear what portion of the observed phenotypic clines are ancestry-driven. We performed admixture mapping to test for genetic loci underlying ancestry-associated differences in wing length and did not identify any loci meeting genome-wide significance (Fig S9).

Our genomic results indicate that the geographic distribution of Africanized honey bees is presently constrained by climatic barriers, not dispersal. Historical records document an initial rapid spread of Africanized honey bees from their point of origin in Rio Claro, São Paulo, Brazil, followed by the formation of seemingly stable hybrid zones at similar latitudes in North and South America. Yet to reach this same latitude, northern-spreading bees had to travel more than five times the distance as southern-spreading Africanized honey bees.

To more precisely infer the current shape and position of the two hybrid zones, we fit a classic logistic cline model to observed genome-wide individual African ancestry proportions [47–49]:

$$A_i = \frac{M}{1 + e^{-b(x_i - c)}}, \tag{1}$$

where A_i is the genome-wide African ancestry proportion inferred for the i^{th}

Fig 2. Clines across latitude. Genome-wide ancestry estimates (top) and fore wing lengths (bottom) for individual bees, plotted across absolute latitude and colored by continent. Shading indicates the 95% confidence intervals for loess curves of the raw data. We also overlay several model-fitted clines: In the top panel, solid curves show the North and South American logistic cline fits for ancestry predicted by latitude, with dotted vertical lines marking the latitude at which bees have predicted 50% African ancestry, based on these curves. Samples from Avalon are displayed as orange triangles; Catalina Island has a distinct ancestry composition from mainland California populations and low African ancestry for its latitude.

In the bottom panel, dashed curves show the expected phenotypic cline if wing lengths were fully determined by the clines in ancestry depicted in the top panel. To get these predicted wing lengths, we used the mean ancestry cline as input to the best-fit linear model between ancestry and wing length. Note that the y-axis for wing lengths is reversed (smaller wings are higher) to simplify visual comparisons between the top and bottom panels.

individual bee, x_i is their latitude, M is the asymptotic maximum African ancestry ₁₄₆
approaching the equator, which is set at 0.84 (i.e. frequency in Brazil [25]), c is the cline ₁₄₇
center, and $w = |4/b|$ is the cline width (i.e. the inverse of the steepest gradient at the ₁₄₈
center of the cline). ₁₄₉

Each degree latitude corresponds to approximately 111km and presents a natural ₁₅₀
way to compare cline position and shape between the two zones. We fit this model in R ₁₅₁
using non-linear least squares (although maximum likelihood or Bayesian estimation are ₁₅₂
generally preferred when the errors can be fully parameterized, here least squares allows ₁₅₃
for unknown drift variance in addition to binomial sampling variance). We find that the ₁₅₄
two hybrid zones have strikingly similar positions (Fig 2), with cline centers that differ ₁₅₅
by less than half a degree (32.72°N vs. 32.26°S), and no statistically significant ₁₅₆
difference in cline steepness. To better understand the mechanisms underlying this ₁₅₇
parallelism between continents, we tested four possible explanatory climate variables to ₁₅₈
see if we could identify a better predictor for African ancestry across our two zones than ₁₅₉
latitude: Mean annual temperature (°C), mean temperature of the coldest quarter (°C), ₁₆₀
minimum temperature of the coldest month (°C), and mean annual precipitation (cm) ₁₆₁
(downloaded from WorldClim.org [50]). We fit clines for both hybrid zones jointly using ₁₆₂
these four environmental variables in turn as predictors in Eq 1 in place of x_i, and ₁₆₃
compared these results to fits based on absolute latitude and, as a neutral dispersal ₁₆₄
model, distance from São Paulo. ₁₆₅

We find that latitude is the best individual predictor of genome-wide global ancestry, ₁₆₆
and mean annual temperature the second-best predictor, as assessed by AIC (see S2 ₁₆₇
Table). While latitude provides the best-fitting cline, we find it unlikely that latitude or ₁₆₈
daylight per se is the relevant selection gradient. Temperature and precipitation are ₁₆₉
closely coupled to latitude across our transect in Argentina, so nearly all of our ₁₇₀
resolution to disentangle latitude from environmental gradients comes from ₁₇₁
micro-climates within California, and for precipitation, the contrast between continents ₁₇₂
(Fig S6). However, we failed to identify specific environmental variables that may be ₁₇₃
driving the relationship with latitude, either because we did not include the relevant ₁₇₄
environmental variable(s) or because the climate data does not reflect the selection ₁₇₅
environment of sampled bees, e.g. due to mismatches in scale or selective habitat use by ₁₇₆
bees within a foraging range. ₁₇₇

Despite limited resolution on the climate variables driving the latitudinal gradient, ₁₇₈
our comparative framework allows us to firmly reject a neutral model based on distance ₁₇₉
from the point of introduction in Brazil, because a single dispersal rate cannot generate ₁₈₀
predictions that simultaneously fit the clines in North and South America well (see S2 ₁₈₁
Table). ₁₈₂

In addition to these global ancestry estimates, we measure variation in local ancestry ₁₈₃
frequencies across the genome, which are informative about recent evolutionary history. ₁₈₄
African ancestry frequencies at individual loci vary around their genome-wide mean due ₁₈₅
to finite sampling, but also evolutionary processes, including drift and selection. If two ₁₈₆
populations have shared gene flow post-admixture, at loci where one population has ₁₈₇
higher than average African ancestry frequencies, the second population will also tend ₁₈₈
to have higher than average African ancestry. We capture this genetic signature in an ₁₈₉
ancestry covariance matrix, where each entry represents how much a pair of populations ₁₉₀
co-deviate in locus-specific African ancestry away from their individual genome-wide ₁₉₁
means (Fig 3). We expect ancestry co-variances to build up along each route of the ₁₉₂
Africanized honey bee invasion as a result of shared drift post-admixture. Indeed, we do ₁₉₃
observe positive ancestry covariances for nearby populations within each hybrid zone. ₁₉₄
We attribute this pattern to shared demographic history, but also note that weak ₁₉₅
selection for a specific ancestry at many loci genome-wide could also generate these ₁₉₆
positive covariances. Unexpectedly, we find that populations in more temperate North ₁₉₇

and South America, i.e. at opposite ends of the expansion, have higher ancestry correlations with each other than with populations situated between them. This robust signal is a general pattern that holds true on average across chromosomes (Fig S14), and so isn't driven by individual outlier loci, and persists across recombination rate bins (Fig S15). These similar ancestry patterns in geographically distant populations are potentially consistent with a genome-wide signature of convergent selection to cooler climates or convergent selection by beekeepers at higher latitudes. Another possible explanation is recent long-distance migration (e.g. international bee exports); however, we investigated genetic covariance patterns within European and African ancestries and found no clear evidence of gene flow between the high-latitude cline endpoints (see methods).

Genetic basis of the climate barrier

To identify loci that may be contributing to a climate barrier, we looked for loci with steeper than expected ancestry clines across latitude in South America. We estimated best-fitting logistic ancestry clines at $\sim$542k single nucleotide polymorphisms (SNPs) across the genome by re-fitting eqn. 1, where x_i is the population latitude and A_i is the population-mean local ancestry at a focal SNP, and the maximum African ancestry M is 1. Similar cline models have been fit using likelihood methods under some simplifying assumptions about the form of the errors (e.g. [48, 49]). We instead use non-linear least squares to fit cline parameters without specifying a full error model and then quantify the effects of more complex unmodeled errors (including ancestry variances and covariances) through simulation. We simulated data for 100,000 independent loci undergoing drift, which we used to estimate the expected distribution of neutral clines and calculate false-discovery rates. For each simulated locus, we independently drew a vector of population ancestry frequencies from a multivariate-normal model of drift, African ancestry $\sim$ MVN(α, K), where α is the vector of population mean genome-wide A ancestry proportions, and the K matrix measures the expected variance and covariance in ancestry away from this mean (Fig 3), and is empirically calculated using all loci across the genome (see methods for additional details). We limit the analysis of clines at individual loci to South America where, unlike North America, we have samples spanning both halves of the hybrid zone to inform parameter estimates. While cline analyses can be used to identify both adaptive introgression and barriers to introgression (by analysing cline center displacement in addition to cline steepness [51]), here we focus on barrier loci and approach identifying positively selected loci using alternative methods that can be applied to both hybrid zones (see "Scan for ancestry-associated selection").

We find no evidence to support a simple genetic basis or environmental threshold to the climate barrier. Ancestry clines in South America are reasonably concordant across most SNPs; 95% of cline centers fall within a 1.6 degrees latitude range, with a long tail that appears to be due to adaptively introgressing loci (identified as outliers below, see Fig S10). We find no strongly selected individual barrier loci that exceed our 5% false-discovery-threshold for cline steepness, set by MVN simulation of background ancestry patterns. On average, individual SNP clines in South America are 960km wide (w = 8.65 degrees latitude), and the steepest cline in the genome still takes approximately 555km (w = 5 degrees latitude) to fully transition from African to European ancestry. These wide clines, coupled with the evidence for parallel genome-wide clines in North and South America, are consistent with selection tracking smooth climate transitions over broad geographic regions rather than a discrete environmental step. Furthermore, concordance in clines across SNPs in South America suggests that many loci are associated with climate-based fitness trade-offs. Under a polygenic climate barrier, we expect locally-adapted loci to be found across the genome

Fig 3. Correlated ancestry across populations. Shared drift in ancestry shown as an ancestry correlation matrix (see methods). Populations are ordered by latitude and diagonals are left blank (within-population correlations = 1). Low and high African ancestry regions of each hybrid zone are defined relative to the estimated latitude of the cline center for genomewide ancestry. About half of the sampled South American populations, and all of the North American populations, fall in the 'low A' half of their respective hybrid zones.

but steeper clines to be more commonly maintained in regions with low recombination rates. This is because selected loci create stronger barriers to gene flow when there is tight genetic linkage than when selection acts on each locus independently [52]. We test this theoretical prediction in South America and find enrichment for steeper clines in regions of the genome with low recombination. The empirical top 5% steepest clines in South America are found on all 16 chromosomes and are enriched in regions of the genome with low recombination. Steep clines comprise 12.7% CI_{95}[8.4%-16.5%] of loci from the lowest recombination rate quintile vs. only 3.3% CI_{95}[3.0%-3.6%] of loci from the highest recombination rate quintile. The average effect of recombination is a 50km decrease in mean cline width between the highest and lowest recombination rate quintiles ($\Delta b = 0.028$, [0.017-0.038]).

Diversity and rapid expansion

From their point of origin in Brazil, Africanized honey bees invaded much of the Americas in less than 50 years [53]. Such rapid expansion can lead to high rates of drift in the continually bottle-necked populations at the front of the wave of expansion, i.e. those populations sampled furthest from Brazil. To test this expectation, we calculated nucleotide diversity, π, for each sampled population (Fig 4). Despite much further distances traveled to the northern hybrid zone, we do not observe a more pronounced bottleneck in California than in Argentina, suggesting that the expanding wave of Africanized honey bees maintained large population sizes (and did not experience strong 'allele surfing' [54]).

Fig 4. Allelic diversity (π) across the hybrid zones. For each population, we estimated allelic diversity genome-wide and within high-confidence homozygous ancestry states. Horizontal lines show the genome-wide diversity within the reference panels. Vertical lines show the 95% confidence interval for each estimate, based on a simple block bootstrap CI using 1cM blocks. For several populations in the tails of the cline, we do not show A and/or C within-ancestry estimates because these populations have too few high-confidence ancestry blocks for accurate estimation (see methods). The low diversity outlier at 33.34 degrees latitude in the N. American cline is the 2014 Avalon sample, which comes from a small island population off the coast of California.

Africanized bee populations are consistently more diverse than reference bee

populations because they are genetic mixtures of these diverged groups. We do, however, observe a drop in diversity in the tails of both hybrid zones starting at approximately 34.5° latitude from the equator. We tested whether this drop in diversity is necessarily the result of a bottleneck or can be explained solely by a cline in mean ancestry composition from more diverse African and highly admixed genomes to less diverse European stock. To test this alternative, we predicted population diversity from a simple weighted average of A, C, and M reference allele frequencies and the observed population ancestry proportions. We find that based on ancestry composition alone, we do expect a drop in diversity across the hybrid zones, although the observed drop is slightly less than our predictions (Fig S25).

Levels of diversity within European C and M ancestries are similar to the reference panels and stable across latitude, evidence that a diverse population of European ancestry bees hybridized with African bees as they expanded away from Brazil. We also find high diversity within African haplotypes in both hybrid zones, again consistent with no bottleneck associated with the rapid expansion. However, the diversity in the African ancestry background does decline in populations furthest from the equator, which is consistent with either strong filtering of African haplotypes by selection or stochastic haplotype loss due to small African-ancestry population sizes in the tails of the clines.

Scan for ancestry-associated selection.

We identified loci with unusually high African ancestry frequencies, a signal of natural selection, using our MVN simulations of background covariance in ancestry to set a false discovery rate. The ancestry covariances are important to account for when testing for putative selected loci that depart from genome-wide background ancestry patterns, because deviations in ancestry are correlated across populations. Although many population pairs have only small positive ancestry covariances, the cumulative effect on the tails of the distribution of African ancestry frequencies in the larger sample is striking. These covariances can confound outlier tests for selection which only consider variance from sampling (e.g. Poisson-Binomial, e.g. [25]). We find that by incorporating background patterns of shared drift (or weak genome-wide selection) into our null model, we can match the bulk of the observed ancestry distributions across the genome (Fig 5).

Loci important to the successful invasion of Africanized honey bees are likely to have an excess or deficit of African ancestry across both continents. Thus, we tested separately for high and low African ancestry outliers on each continent, and then identified overlapping outliers between the two hybrid zones. We find evidence of selection favoring African ancestry at 0.34% of loci in N. America and 0.13% of loci in S. America, across 14 chromosomes (Fig 6A). From these outliers, we find 13 regions with an excess of African ancestry in both hybrid zones at less than a 10% false-discovery-rate (top right corner of Fig 5). The majority (11/13) of shared outliers co-localize within a ~1.5Mb region on chromosome 1, but within this region outliers separate into multiple distinct peaks (Fig 6B). One way a cluster of African ancestry peaks could form is if favored African alleles experience additional indirect selection from being in linkage disequilibrium with other favored African alleles at nearby loci, thereby increasing the total effective selection in a region [52]. While ancestry-informative markers (AIMs) with fixed or nearly fixed differences between African and both European ancestries are relatively rare, we were able to confirm the highest A peak within this cluster using AIMs outside of the local ancestry inference SNP set (Fig S20). African alleles at this main peak appear to have introgressed to high frequency hundreds of kilometers past the hybrid zone centers in both North and South America, but not reached fixation in any population (Fig 7). The rapid rise and slow fixation of this allele is potentially consistent with dominant fitness benefits. How far these African alleles have introgressed past the present hybrid zones is currently

Fig 5. Local ancestry outliers compared across hybrid zones. We plot mean African ancestry frequency in North vs. South America for 425k SNPs across the honey bee genome. SNPs are binned for visualization, and colored by the number of SNPs within each hexagon. The orange ellipse shows the approximate 99% highest posterior density interval (HPDI) based on the full MVN model, which accounts for drift in ancestry both within and between populations. Using the same axes, we show the marginal histograms of African ancestry for each continent separately (top and right panels). Imposed on these histograms we plot density curves for 3 possible null distributions for ancestry frequencies: the full MVN model, a variance-only MVN model which only accounts for drift within populations, and a Poisson binomial model which only includes sampling variance. Most of the genome is consistent with neutrality under a MVN normal model of drift (98.6% of SNPs fall within the orange ellipse), but there are also some clear outliers. SNPs in the top right, with higher than expected African ancestry proportions in both hybrid zones, are our best candidates for loci underlying adaptive African-ancestry associated traits. Note: While SNPs are thinned for LD, large outlier regions span many SNPs, which creates the streak-like patterns in the scatterplot.

Fig 6. Genomic location of ancestry outliers. (A) Mean African (A) ancestry in each hybrid zone at SNPs across the genome, with outliers colored by their false-discovery-rate. Genome-wide mean African ancestry in each zone is indicated with a dashed line. Shared peaks for high African ancestry are seen on chromosomes 1 and 11; there are no shared peaks for low A ancestry. (B) Zoomed in view of cluster of shared high African ancestry outliers on chromosome 1, with European ancestry separated into eastern (C) and western (M) subtypes. Genome-wide mean frequencies for each ancestry are shown with colored lines. Outlier regions meeting a 10% FDR for high A ancestry are shaded in grey. Shared outliers between continents overlap between the top and bottom panels. (C) Zoomed in view of the high M European ancestry outlier region found in South America. Outlier regions for low African ancestry (<10% FDR) are shaded in grey. Note: The x-axis scale differs between plots.

Our goal was to identify regions of the genome where high fitness is broadly 323
associated with African ancestry, but an alternative explanation for high African 324
ancestry at a locus is that a very recent adaptive mutation just happened to fall on an 325
African haplotype, initiating a classic 'hard sweep'. For shared high African-ancestry 326
outlier regions, we distinguished between these two scenarios using population 327
differentiation (F_{ST}) within A ancestry. We analyzed differentiation across the large 328
cluster of shared high African-ancestry outlier peaks on chromosomes 1 and across a 329
smaller region on chromosome 11 that contains the other two high African ancestry 330
outliers shared between continents. We did not find high allelic differentiation between 331
North and South American A ancestry tracks and the A reference panel (Figs S22 and 332
S23), suggesting that African ancestry in general, and not one particular haplotype, was 333
favored by natural selection at these loci. 334
We used previous literature and gene orthologs to identify possible adaptive 335
functions for regions of the genome where selection has favored African ancestry. There 336

Fig 7. Ancestry clines at outlier SNPs. At two top outlier SNPs, we show the clines for mean population African ancestry across latitude in North (top) and South (bottom) America. To the left, we show the ancestry cline for the SNP with the highest African ancestry in the combined sample, located within the top peak on chromosome 1 for high shared African ancestry across continents. To the right, we show the ancestry cline for the SNP with the lowest African ancestry in South America, located within the large outlier region on chromosome 11 for high M and low A ancestry. Genome-wide mean local ancestry calls for each population are shown for comparison as black circles.

are 3 major quantitative trait loci (QTLs) associated with defense behaviors (e.g. stinging) in genetic crosses of Africanized and European bees [26, 27], none of which overlap any signatures of selection from this study. No studies have mapped the genetic basis of elevated *Varroa* defense in Africanized (vs. European) bees, but we were able to compare our results to quantitative trait loci (QTLs) associated with anti-*Varroa* hygiene behaviors [55–57] and defensive grooming [58] more generally. The cluster of peaks for high shared African ancestry on chromosome 1 overlaps a putative QTL associated with removal of Varroa-infested brood [57], but there are a number of large QTLs in the genome. A total of 104 genes overlap high African ancestry outlier peaks. Predicted functions for these genes (primarily based on fly orthologs) are not significantly enriched for any Gene Ontology (GO) categories, which may simply reflect that many outlier regions are broad and contain many genes, most of which are likely unrelated to their rise in ancestry frequency. A smaller set of 37 genes with high African ancestry have signatures of selection on both continents. For these, we searched the literature and found that two have been associated with *Varroa* in previous studies: a myoneurin (LOC725494) that is overexpressed in the brains of *Varroa* infected worker bees compared to *Nosema* infected bees and controls [59] and an uncharacterized protein (LOC725683) that is over-expressed in parasitized drones compared to non-parasitized drones [60]. While these are potentially intriguing candidates for selection, *Varroa* is only one of many possible selective pressures, and more work is needed to link the signals of selection we find here to adaptive functions.

In contrast to high African ancestry outliers, we do not find any shared outliers for European ancestry. However, the most striking example of a single-zone selection event is a large 1.4Mb region on chromosome 11 with excess European ancestry in South America (bottom middle of Fig 5). This region was previously identified to have low African ancestry in Africanized honey bee populations from Brazil [25]. We independently identify this region as a low African-ancestry outlier across Argentina, using different A/C/M reference bees, ancestry calling algorithm, and bee samples than the previous paper. We find that populations across the South American hybrid zone have reduced African ancestry at this locus, but that North American populations do not appear to have experienced selection (Fig 7). By including C-lineage diversity in our admixture analysis, we additionally show that this region is specifically elevated for M haplotypes, and not European haplotypes more broadly (Fig 6C). This region has many diverged SNPs between the three ancestry groups beyond the SNPs included in our ancestry_hmm analysis, which we use to confirm high rates of M introgression (Fig S21). It does not appear that a new mutation or narrow set of haplotypes was favored within M because we see little differentiation between the M ancestry in this selected region compared to the M reference panel. Additionally, within the selected region we find a peak of high F_{ST} between A, C, and M reference panels (Fig S24), which is consistent with this region having historically been under selection within these ancestry groups. Finally, the California bees do not have a significant deficit of African ancestry like the Argentinian bees do, but they do have two narrow peaks of excess M ancestry within this region, in the top 3% and 7% empirical percentiles for M ancestry genome-wide. Our data support a scenario in which a diversity of M haplotypes carrying the favored allele were driven to high frequency in South America after Africanized honey bees spread north of Brazil. Potential candidate genes specific to this large M-ancestry outlier region on chromosome 11 are previously described by [25]. In total, we find 186 genes that overlap low African ancestry outlier peaks (<10% FDR). These genes are not functionally enriched for any Gene Ontology (GO) categories.

Discussion

The introduction of African honey bees to Brazil in the 1950s sparked one of the largest and best studied biological invasions known to date, spreading from a single point of release over much of the Americas in less than 50 years. We add to this literature the first comparative study of the invasions in North and South America, with genome-wide resolution on the present distribution of African ancestry.

The parallel alignment of the genome-wide cline with latitude in both continents, despite very different length dispersal routes, strongly supports the view that African-ancestry has reached a stable climatic range limit. Because our transect in California only covered the upper half of the North American cline, the full shape for this genome-wide cline is uncertain, and may be asymmetrical because one signature of a moving hybrid zone is elongation of the lagging tail [61]. In contrast, we have strong evidence for convergence in the low-African portions of these two genome-wide clines, which are not expected to be distorted by cline movement, and reflects similar latitudinal range limits for African ancestry in North and South America. Global warming trends could shift the location of the observed clines towards the poles, as has been documented for other hybrid zones sensitive to climate change [62]. While we currently lack temporal data with comparable genomic and geographic resolution, our results can be used as a baseline for future study.

We observe relative uniformity at the climate barrier, with no individual loci showing steeper ancestry clines than what can be produced by a null model accounting for background patterns of variation in ancestry frequencies shared across populations. Nor do we observe any loci that have below 10% frequency of African ancestry in California, despite the large distance and climatic range traveled over by this portion of the invasion. If the invasive ability of Africanized bees were due to a small number of loci we would expect African ancestry to have been swamped out at many unlinked neutral loci in the genome due to interbreeding at the front of the advancing wave of expansion [63]. Instead, relative genomic cohesion points to a polygenic basis for the high fitness and rapid spread of African ancestry as well as the fitness costs in cooler climates underlying the parallel range limits observed across continents. However, we note that the distinction between Africanized and non-Africanized honey bees is likely to further blur over time. Genetic barriers are strengthened when selection is distributed across many loci, but they are still easily permeated by adaptive alleles [64]. Furthermore, given high recombination rates in honey bees, we predict only loci tightly associated with climate-based fitness trade-offs will remain geographically bounded over long periods of time.

Significant effort has been focused on finding an environmental isocline that divides regions where Africanized honey bees are expected to dominate and regions where they cannot overwinter (e.g. [11–14]). However, we observe ancestry clines that are hundreds of kilometers wide, not the narrow clines created by strong selection across a discrete environmental transition. Theoretically, these broad clines could be consistent with neutral diffusion of ancestry by migration over tens of generations. However, a scenario of neutral diffusion is inconsistent with external evidence of the rapid spread of the invasion and strong fitness trade-offs with climate: the high competitive advantage and very rapid advance of Africanized honey bees in the tropics in the face of considerable interbreeding with European bees and, conversely, documented low fitness in cooler climates, with low overwinter survival and maladaptive metabolic efficiency, foraging preferences and nesting behaviors (see [12] for a review). Thus, we conclude that honey bee fitness is more likely to be tracking environmental variables with smooth transitions over broad geographic regions (e.g. climate), which may create intermediate environments where ancestry intermediates have higher fitness, thus broadening the observed hybrid zones. These proposed dynamics are similar to well-studied cases in

other systems where bounded hybrid superiority and/or local adaptation to continuous 438
environments maintain adaptive clines across broad geographic regions (e.g. [65–67]). 439

We sampled an abundance of bees with intermediate African ancestry proportions, 440
which implies that Africanized honey bees cannot be treated as a single genetically and 441
phenotypically cohesive group. Rather, as a null model, we expect phenotypic clines to 442
match the scale of the observed ancestry clines, with smooth transitions in mean 443
phenotype over hundreds of kilometers. Many phenotypes of interest, e.g. defensive 444
behavior or *Varroa* tolerance, are expressed or measured at the colony-level and so we 445
could not assess these in our survey of freely foraging bees. Future phenotypic surveys 446
could be compared with our genomic clines to ascertain if key phenotypes diverge from 447
this expected pattern, e.g. due to strong selection beyond that experienced by the rest 448
of the genome. Indeed we see that wing length, a trait hypothesized to be associated 449
with latitudinal body-size adaptation following Bergmann's rule [46], has a geographic 450
distribution consistent with the genome-wide ancestry cline. This suggests that while 451
wing length, which is strongly correlated with body size [34,68], may well have fitness 452
trade-offs with climate, selection for these traits does not appear to be strong enough 453
compared to average selection for ancestry to deviate from background genomic 454
patterns over a short time scale. 455

Africanized honey bees are a potentially powerful natural system to study the 456
genetic basis of phenotypic adaptation because of the diversity of traits that differ 457
between European and African honeybees and the fine-scale mosaic of ancestry across 458
their genomes, created by over 60 generations of admixture at high recombination rates 459
(24cM/Mb [69]). While we did not identify any large effect loci in our admixture 460
mapping study for wing length, this approach can be readily applied to other 461
measurable traits of interest with genetic differences between European and African 462
honey bees, including defensive behaviors [26,27,70], foraging preferences [26,36], body 463
size [27], overwintering-success [9,10] and *Varroa* tolerance [30,32,33]. 464

Africanized honey bees provide a promising source of genetic variation for breeding 465
in light of the vulnerability of European lineages to current environmental stressors and 466
associated bee declines [71]. Africanized honey bees have high competitive fitness and, 467
we show here, maintained high genetic diversity despite their rapid expansion. However, 468
a targeted approach is needed to identify high fitness African alleles that are promising 469
for breeding within a background that segregates for undesirable traits such as defensive 470
behaviors. In this study, we have taken a first step towards mapping the genetic basis of 471
the high fitness of Africanized honey bees by identifying loci where selection has favored 472
African or European ancestry in both North and South America. We identify several 473
loci with convergently high African ancestry on both continents, and many more across 474
the genome with evidence of selection favoring African ancestry in one hybrid zone. In 475
contrast, with the exception of one striking outlier for high M ancestry in South 476
America, we find little evidence that European ancestry or admixture per se contributed 477
broadly to the success of Africanized honey bees. We attribute this difference in results 478
from a previous study of Africanized honey bees in Brazil [25] to a more appropriate 479
null model that accounts for shared variance in ancestry across populations. While our 480
population genetics approach is trait-blind, our results can be compared to future 481
functional and genetic mapping studies to look for overlap between trait-associated and 482
positively selected loci. Applying similar methods to other systems, especially where 483
replicated hybrid zones can be sampled, holds great promise for revealing loci important 484
to adaptation. 485

Materials and Methods

Statistical results and figures were created in R [72] with use of the tidyverse [73] packages. Other scripts were run using GNU parallel [74].

Sampling

We sampled individual foraging honey bees across two hybrid zones, located at the transitions to temperate climates in North and South America. We sampled at least 10 bees each from 12 populations in California and 21 populations in Argentina (see maps, Fig 1).

For each population, we hand-netted individual foraging bees within a sampling radius of approximately 15km. Because commercial colonies are often temporarily relocated for the spring pollination season, we sampled in summer, when foraging bees are more likely to come from resident populations. We additionally included in our analyses 35 high-coverage published genomes of freely foraging bees collected from 6 populations between September 2014 and January 2015: Davis, Stebbins, Stanislaus, Avalon (Catalina Island), Placerita, and Riverside (Sky Valley and Idyllwild) [22]. While these sampled bees come from an unknown mixture of local feral and domesticated colonies, previous surveys from California have found that freely foraging bees tend to closely match feral sources, based on mtDNA composition [18]. Consistent with this view, eight of our sequenced bees from different populations in Argentina were collected close to a feral nest ($< 5m$), but do not appear to be ancestry outliers for their sampling locations. More specifically, we fit a general linear model (logit(African ancestry) $\sim$ absolute latitude + feral nest) using glm with gaussian errors in R and found no significant effect on African ancestry of sampling near a feral nest ($P = 0.97$). Based on these results and our seasonal timing, the bees in this study are likely sourced primarily from local feral populations, with some contribution from resident domesticated bee colonies.

Lab work and sequencing

We selected a subset of 279 bees from our North and South American hybrid zones for whole genome sequencing, 8-9 bees per sampled population (see S1 Table). For each bee, we dissected wing flight muscles from the thorax and extracted DNA using QIAGEN DNeasy Blood and Tissue kits. We followed a new high-throughput low-volume DNA library preparation protocol (see [75] for details, "Nextera Low Input, Transposase Enabled protocol"). Briefly, we prepared individual Nextera whole-genome shotgun-sequencing DNA libraries using enzymatic sheering and tagmentation. Then we PCR-amplified and barcoded individual libraries using the Kapa2G Robust PCR kit and unique custom 9bp 3' indices. Finally, we pooled libraries within each lane and ran bead-based size-selection for 300-500bp target insert sizes. We targeted 4-6x coverage per bee based on a preliminary analysis of our power to replicate local ancestry calls from one of the published high coverage populations (Riverside 2014) using simulated low coverage data (Fig S4). We multiplexed our samples across 5 Illumina HiSeq4000 lanes for paired-end 2 x 150bp sequencing. In total, we generated 5.1x mean coverage per bee for 278 samples. The 279th sample was excluded from all analyses for having extremely low ($<0.1x$) sequence coverage.

Alignment and SNP set

In addition to the sequence data produced by this study, we downloaded Illumina raw read sequences for 35 previously published California genomes (PRJNA385500 [22]) and

a high-quality reference panel of African (n = 17), eastern European (n = 9) and western European (n = 9) honey bee genomes (PRJNA216922 [76] and PRJNA294105 [77]) from the NCBI Short Read Archive. For all bees, we mapped raw reads to the honey bee reference genome HAv3.1 [78] using Bowtie2 very-sensitive-alignment with default parameters [79]. We then marked and removed duplicate reads with PICARD and capped base quality scores using the 'extended BAQ' option in SAMtools [80]. Using the software ANGSD [81], we identified a set of SNPs with minor allele frequency $\geq 5\%$ in the combined sample based on read counts (-doMajorMinor 2 -doCounts 1). We excluded unplaced scaffolds (<5Mb total) and applied standard quality filters for SNP calling (base quality $\geq$ 20, mapping quality $\geq$ 30, total read depth $\leq$ 5500 ($\sim$2x mean), and coverage across individuals $\geq$ 50%). We calculated the genetic position (cM) for each SNP using a 10kb-scale recombination map [69] and linear interpolation in R (approxfun). We assumed constant recombination rates within windows and extrapolated positions beyond the map using the recombination rate from the nearest mapped window on that chromosome.

We identified SNPs on the mitochondria (HAv3.1 scaffold NC_001566.1) using the same pipeline as nuclear DNA above, but allowing for extra read depth (up to 100000000x). We then called consensus haploid genotypes at these SNPs for all individuals using ANGSD (-dohaplocall 2 -remove_bads 1 -minMapQ 30 -minQ 20 -doCounts 1 -minMinor 2 -maxMis 174).

Global ancestry inference

We estimated genome-wide ancestry proportions for each bee using methods designed for low-coverage sequence data. Briefly, we combined bee genomes from the hybrid zones with reference genomes from Africa (A), eastern Europe (C) and western Europe (M). To reduce linkage disequilibrium (non-independence) between our markers for global ancestry inference, we thinned to every 250th SNP ($\sim$14k SNPs at 19kb mean spacing) before calculating genotype-likelihoods for each bee using the SAMtools method in ANGSD (-GL 1). We first ran a principal components analysis in PCAngsd [82] to confirm that genetic diversity in the hybrid zones is well-described by 3-way admixture between A, C, and M reference panels (Fig S1). We then estimated genome-wide ancestry proportions for all bees using NGSAdmix (K = 3) [37].

Local ancestry inference

We inferred the mosaic of African vs. European ancestry across the genome of each bee using a hidden Markov inference method that can account for African (A), eastern European (C) and western European (M) sources of ancestry within low-coverage Africanized honey bee genomes (ancestry_hmm v0.94 [38]). For local ancestry inference, we enriched for ancestry-informative sites by filtering for ≥ 0.3 frequency in one or more reference population (A, C, or M) and at least 6 individuals with data from each reference population. We subsequently thinned markers to 0.005cM spacing, because at that distance linkage disequilibrium within ancestries is expected to be low ($r^2 < 0.2$ [21]), leaving a final set of 542,655 sites for ancestry calling, or $\sim 1/7$ of the original SNP set. Individual bees sequenced in this study and previously published California bees have 5.42x and 14.5x mean coverage, respectively, across this final SNP set. For each population, we jointly estimated time since admixture and ancestry across the genome of each individual, using read counts from the hybrid zone and allele frequencies for European and African reference populations at each SNP. To generate major/minor allele counts for each reference population, we used ANGSD to call genotypes (-doPost 1) using a minor allele frequency prior (-doMaf 1) and the SAMtools genotype likelihood (-GL 1), after quality filtering (map quality $\geq$ 30, reads matching

major/minor allele $\geq 60\%$, and read depth $\geq$ 6x). As additional inputs to ancestry_hmm, we used NGSAdmix results as a prior for population ancestry proportions and set the effective population size to $N_e = 670,000$ [25]. We modelled a simple three-way admixture scenario: starting with C ancestry, we allowed for a migration pulse from M and a second, more recent, migration pulse from A. Timing of both migration pulses were inferred from the range 2-150 generations, with priors set at 100 and 60 generations. To calculate a point estimate for each individual's ancestry proportion at a SNP, we marginalized over the posterior probabilities for homozygous and heterozygous ancestry from the ancestry_hmm output (i.e. $A = p(AA) + 1/2(p(CA) + p(MA))$.

African ancestry covariance matrix

To explore how populations vary and covary in their African ancestry along the genome we calculated the empirical population ancestry variance-covariance matrix (K), an admixture analog of a genotype coancestry matrix (e.g. [83]). The K matrix is calculated using population African ancestry frequencies inferred by the local ancestry HMM, e.g. for populations i and j with mean ancestry proportions α_i and α_j, and ancestry frequencies at a locus $anc_{i,l}$ and $anc_{j,l}$, their ancestry covariance calculated across all L loci genome-wide is

$$K[i,j] = \frac{1}{L} \sum_{l=1}^{L} (anc_{i,l} - \alpha_i)(anc_{j,l} - \alpha_j).$$

Ancestry correlations between the high-latitude cline endpoints

To more formally test for excess ancestry correlations between more geographically distant (but climatically similar) populations, we grouped populations by dividing each hybrid zone into low- and high-African ancestry regions relative to the estimated latitude for the genome-wide cline center. The southernmost 11 (out of 20) of the South American populations, and all of the sampled North American populations, fall in the 'low A' half of their respective hybrid zones. We calculated mean ancestry covariances (K matrices) separately for each chromosome, using the genome-wide mean ancestry as α, then summarised across populations by taking the mean correlation for each type of pairwise comparison, within and between continents and regions. We tested whether, on average across chromosomes, low-A South American populations share higher ancestry correlations with low-A North American populations than with geographically closer high-A South American populations and repeated this test excluding chromosomes 1 and 11 which contain large outlier regions (Fig S14). We also tested the same group comparison across recombination rate quintiles instead of chromosomes (Fig S15).

To investigate whether recent long-distance migration likely generated the elevated ancestry correlations we observe between low-A South America and low-A North America, we looked at patterns of allelic covariance within ancestry. Specifically, for each ancestry we estimated a genetic covariance matrix in PCAngsd for all individuals sampled from the hybrid zone, based on allelic diversity within high-confidence homozygous ancestry tracts (posterior >0.8). Under recent migration, we would expect the excess African ancestry correlations between the two ends of the hybrid zones to be mirrored by allelic covariances within all three ancestries. Instead, we find that the two most prevalent ancestries, A and C, both have low or negative genetic covariances between continents (Fig S17). In contrast, M ancestry does show an excess of cross-continent covariance, and we followed up to determine if this is uniquely American covariance (i.e. the result of shared drift within the Americas) or could have been imported from Europe. Adding reference individuals to these within-ancestry analyses,

we find that M ancestry in the Americas imported pre-existing structure from Europe, with more Poland-like (*Apis mellifera mellifera*) than Spain-like (*Apis mellifera iberiensis*) M ancestry at the temperate ends of the clines (Figs S18 and S19).

Simulated ancestry frequencies

At various points in the results we compare our outliers to those generated by genome-wide null models of ancestry variation along the genome. We simulated variation in ancestry frequencies at SNPs across the genome under three models: (1) A poisson-binomial model that only accounts for sampling variance, not drift (e.g. [25]); (2) a multivariate-normal model with covariances set to zero, which accounts for effects of both sampling and drift within-populations (e.g. [84]); and (3) a multivariate-normal model with covariances to additionally account for shared drift in ancestry between populations. For each model, we simulated in R neutral African ancestry frequencies at 100,000 independent loci [85, 86]. The full multivariate-normal model is used for comparison to the results, while the first two models are only used to show the effects of ignoring covariances.

In the Poisson-Binomial simulation, for each bee we sampled 2 alleles from a binomial distribution with mean equal to the individual's genome-wide ancestry proportion inferred by ancestry_hmm.

For the variance-only MVN simulation, we empirically calibrated an independent normal distribution for each population that can exceed binomial ancestry variance (e.g. due to drift). This model is equivalent to the full MVN model below, but sets all off-diagonal entries of the K ancestry variance-covariance matrix to zero.

In our full multivariate-normal model, we account for non-independent ancestry within and between our sampled populations:

$$\text{African ancestry} \sim \text{MVN}(\alpha, K),$$

where α is the vector of genome-wide mean population ancestry frequencies and K is the empirical population ancestry variance-covariance matrix. Because the MVN models are not bounded by 0 and 1, but real frequency data is, we set all simulated individual population frequencies exceeding those bounds (5.2% low and 0.09% high) to the bound. Truncation has little effect on the distribution in general and no effect on the frequency of high African ancestry outliers, but does make extremely low outliers (attributed to some populations having simulated negative frequencies) less likely (Fig S11). For more details on model approximations to the observed data, see Figs S12 and S13.

Cline models

To better understand the role of dispersal and selection maintaining the current geographic range limits of Africanized honey bees, we fit a logistic cline model to the individual genome-wide ancestry proportions estimated by NGSAdmix. We estimated continent-specific c and b parameters to test for a difference in cline center (degrees latitude from the equator) and/or slope between the northern and southern invasions. Then we fit a joint model with a single cline to see how well absolute latitude or climate can consistently predict African ancestry frequencies across both continents. Specifically, we tested three environmental variables that likely contribute to varying fitness across space: mean annual temperature (°C), mean temperature of the coldest quarter (°C), and mean annual precipitation (cm). We downloaded mean climate observations for 1960-1990 [50] from WorldClim.org at 30 second map resolution (≈ 1 km^2 at the equator) and then averaged within a 5km radius around each bee's sample coordinates. We compared climate and latitude-based selection models to a neutral dispersal model,

where genome-wide African ancestry is predicted solely based on the distance (km) traveled from Río Claro, São Paulo, Brazil, the point of origin for the Africanized honey bee invasion (estimated from GPS coordinates using "distGeo" in R [87]). For each model, we substituted latitude, distance, or climate for x_i in Eq 1 and we used AIC to compare model fits.

We then fit individual-SNP clines to the mean population ancestry frequencies in South America, where our samples span the full cline. We tested for individual outlier loci that may underlie a climate barrier by fitting the same logistic cline model to a set of simulated population ancestry frequencies for S. America (see MVN simulation), and comparing observed cline slopes to this null distribution. In addition, we tested for enrichment of the empirical top 5% of steep clines in regions of the genome with low recombination rates. We divided the genome into 5 equal-sized recombination rate bins ([0, 2.92], (2.92, 21.6], (21.6, 31.7], (31.7, 38.6] and (38.6, 66.9] cM/Mb) and used 10,000 block bootstraps [88] to calculate basic bootstrap confidence intervals for each recombination rate quintile while accounting for spatial correlation in both cline slopes and recombination rates across the genome. For the bootstrap, we divided the genome into 0.2cM blocks, we re-sampled these blocks with replacement, and for each recombination bin we calculated mean b and the proportion of SNPs in the top 5% steepest slopes from our bootstrap sample. When fitting non-linear least squares in R for both genomewide and individual snp clines, we used multiple random starting values to make sure we searched across all local minima and found the global optimum solution (nls.multstart [89]). Starting values were drawn from uniform distributions: b ~ Unif(-5, 5) and ~ Unif(min, max) across the observed range for latitude and climate variables.

Wing morphology

We imaged a slide-mounted fore wing and measured wing length to the end of the marginal cell using imageJ (Fig S8). We included 269 bees in the wing analysis (only bees sequenced by this study had wings preserved and n = 9 bees were excluded for wing tatter or damage).

We also measured fore wing lengths for A, C, and M reference bees in the Oberursel Collection sampled from their native range (n = 52 [90]). While the effect of ancestry on wing length is similar in magnitude and direction in both datasets, we found that the mean wing lengths for the European reference bees (C & M) fell below the mean for our American bees with close to 100% European ancestry, perhaps reflecting phenotypic plasticity. Thus we do not incorporate these measurements of A, C, and M reference bees into the subsequent analyses.

We tested various models of the relationship between wing length, ancestry and geography. First, we fit a linear model to predict wing length in our sample from genome-wide ancestry. We visually compared our wing measurements to what we would expect if the cline in wing length across latitude were fully described by this linear relationship between ancestry and wing length and our best-fit clines for genome-wide ancestry (Fig 2). We additionally tested for differences between continents by adding a main effect and an interaction term for South America to our linear model.

We performed admixture mapping of wing length to test if the ancestry state at any individual SNP predicts residual variation in wing length. To do this, we first regressed wing length on genome-wide African ancestry, to correct for background ancestry effects. We then took the residual wing lengths from this linear model fit and regressed these on African ancestry allele counts at each locus in turn (using the maximum a posterior probability (MAP) estimates from the local ancestry HMM). We set a genome-wide significance threshold of $p < 1.1 \times 10^{-6}$ to control for multiple testing at a 5% family-wise error rate, using an analytical approximation for admixture mapping, calculated assuming 47.6 generations since admixture [91, 92].

Identifying ancestry outlier regions and genes ₇₁₅

To identify loci underlying ancestry-associated fitness differences, we tested SNPs for an ₇₁₆
excess or deficit African ancestry within each hybrid zone. We calculated 1%, 5% and ₇₁₇
10% false-discovery rates (FDR) by using our MVN simulation results to set the number ₇₁₈
of false-positives we expect under a neutral model for high and low African ancestry ₇₁₉
within each continent separately (one-tailed outlier tests). We then compared the ₇₂₀
overlap in outliers between hybrid zones to identify SNPs with signatures of selection on ₇₂₁
both continents. ₇₂₂

In addition to local ancestry, we used ancestry-informative markers with fixed or ₇₂₃
nearly fixed differences to verify high-introgression regions. We defined ancestry ₇₂₄
informative markers as SNPs with coverage for at least 5 individuals from each reference ₇₂₅
panel and >0.95 allele frequency differences between the focal ancestry and both other ₇₂₆
ancestries. We estimated allele frequencies at each ancestry-informative marker using ₇₂₇
ANGSD, polarized SNPs so the focal ancestry has the highest MAF, and only included ₇₂₈
markers with coverage in all sampled populations. Ancestry-informative markers for A ₇₂₉
(n = 4,302) are relatively rare compared to markers for C (n = 17,384) and M (n = ₇₃₀
15,626) because European populations each experienced a historical bottleneck ₇₃₁
differentiating them from the other two groups. Because of LD-thinning before local ₇₃₂
ancestry inference, 88% of these ancestry-informative markers were not included in the ₇₃₃
ancestry_hmm SNP set, and therefore provide separate support for high-introgression ₇₃₄
regions. ₇₃₅

We identified a set of candidate genes that overlap regions of the genome with ₇₃₆
exceptionally high or low African ancestry (<10% FDR) using BEDtools [93]. For this ₇₃₇
analysis, we downloaded gene annotations for the HAv3.1 genome assembly from NCBI ₇₃₈
(accessed 7/22/19). 72 out of 104 genes overlapping high African ancestry peaks and 131 ₇₃₉
out of 186 genes overlapping low African ancestry peaks have associated BEEBASE ₇₄₀
gene IDs. For these high and low gene sets, we tested for enrichment of Gene Ontology ₇₄₁
(GO) terms compared to a background of all honey bee genes, using DAVID 6.8 [94] and ₇₄₂
a Benjamini-Hochberg corrected FDR of 5% [95]. To find out what is previously known ₇₄₃
about the 37 genes that overlap regions with high African ancestry on both continents, ₇₄₄
we conducted a literature search using the NCBI gene search tool and google. ₇₄₅

We additionally checked if our candidate selected loci overlap regions of the genome ₇₄₆
previously associated with defensive or anti-*Varroa* behavioral traits (QTLs and ₇₄₇
associated marker sequences from [26, 27, 55–58, 96–98]). We estimated genome ₇₄₈
coordinates for QTLs by blasting marker sequences to HAv3.1 and keeping the best ₇₄₉
BLASTn [99] hit with an E-value <0.01 (see S5 Table). When assessing physical ₇₅₀
overlap between genome annotations and ancestry outliers, we assumed ancestry calls ₇₅₁
for a SNP apply to the short genomic window around that SNP, spanning midway to ₇₅₂
the next ancestry call. When visualizing and counting the number of selected regions in ₇₅₃
the genome, we further merged near-adjacent (<10kb) significant ancestry windows into ₇₅₄
contiguous regions. ₇₅₅

Population diversity ₇₅₆

We calculated allelic diversity (π) for each population and our A, C, and M reference ₇₅₇
panels. First, we calculated a simple unbiased population allele frequency in ANGSD ₇₅₈
based on a weighted average of observed read counts (counts -8) for each SNP. For this ₇₅₉
analysis, we included all SNPs ascertained in the combined sample (see 'Alignment and ₇₆₀
SNP set' above) but excluded SNPs from a population's estimate when fewer than two ₇₆₁
individuals had coverage. Using these allele frequency estimates, and a finite-sample ₇₆₂
size correction ($n = 2\times$ number of individuals with data at a site), we calculated mean ₇₆₃
per-SNP heterozygosity. To approximate uncertainty in our estimates, we divided the ₇₆₄

genome into 5,254 non-overlapping 1cM blocks, re-calculated our diversity estimates for
10,000 block bootstrap samples, and calculated a 95% simple bootstrap confidence
interval. Finally, to get per-bp diversity, we scaled our per-SNP diversity estimates by
the density of SNPs in the genome, using the same coverage and depth quality filters in
ANGSD as in our SNP pipeline to count total mappable sites.

For within-ancestry diversity estimates, we used our ancestry calls to identify
contiguous tracts with high posterior probabilities (>0.8) of homozygous A, C, or M
ancestry. We used these tracts to divide the genome into high confidence A, C, and M
ancestry states, and filter for reads that mapped within these states. We then repeated
the estimation and block bootstrap procedure above using only the reads associated
with a particular ancestry. To estimate within-ancestry diversity for a population, we
required data for at least 75 1cM blocks spread across at least 15 of the 16 chromosomes,
which excludes 6 populations with too little A ancestry in the tails of both clines and 8
populations with too little C ancestry in the S. American cline for accurate estimation.
We compared observed and predicted heterozygosity for each population based on
expected allele frequencies calculated by multiplying population-specific admixture
proportions by reference population allele frequencies for each ancestry.

To test whether selection had favored specific haplotypes, or African ancestry more
generally, within shared outlier regions for high African ancestry, we calculated
population differentiation (F_{ST}) between the A reference panel and A haplotypes within
each hybrid zone. We also calculated within-ancestry F_{ST} between the two hybrid zones,
to assess whether the same A haplotypes rose in frequency on both continents. Likewise,
for the large high M outlier on chromosome 11, we calculated pairwise differentiation
within M ancestry between North America, South America, and the M reference panel.
We similarly calculated F_{ST} for all three contrasts between A, C, and M, reference
panels across these outlier regions, to test for signatures of historical selection and
divergence between these ancestry groups. For F_{ST} calculations, we estimated
within-ancestry allele frequencies for North and South America using the same method
described above for within-ancestry π, except pooling individuals by hybrid zone rather
than population. We used Hudson's estimator for F_{ST} (Eq 10 in [100]), calculated the
average per-SNP F_{ST} within sliding 50kb windows stepping every 1kb across ancestry
outlier regions, and only included SNPs with coverage for at least two individuals for
both populations in the contrast and windows with at least 10 SNPs.

Ethics statement

Data availability

Raw Illumina sequence data generated by this study is available through the NCBI
Short Read Archive, PRJNA622776. Access information for all previously published
genomic resources used in this study: HAv3.1 reference genome and gene annotations
(NCBI PRJNA471592), recombination map (available by request from Jones et al.), bee

genomes from California (NCBI PRJNA385500) and A, C, and M reference populations ₈₁₂
(NCBI PRJNA216922 and PRJNA294105). ₈₁₃

Bee metadata, including GPS locations and measured wing lengths are included in ₈₁₄
S1 Table. Wing images generated by this study are available through Data Dryad: ₈₁₅
https://doi.org/10.25338/B8T032. Wing images for museum samples of A, C, and M ₈₁₆
bees are available by request from the Morphometric Bee Data Bank, Institut für ₈₁₇
Bienenkunde, Oberursel, Germany (https://de.institut-fuer-bienenkunde.de). All ₈₁₈
climate data was downloaded from WorldClim.org. ₈₁₉

Scripts are available at https://github.com/ecalfee/bees. ₈₂₀

Acknowledgements ₈₂₁

We thank the National Institute of Agricultural Technology (INTA), Argentina, for the ₈₂₂
use of bee samples from Argentina. Thank you to Stefan Fuchs and the Oberursel ₈₂₃
Collection for sharing wing images of A, C and M reference bees. Thank you to Brock ₈₂₄
Harpur for making marker sequences from previous Hunt lab QTL studies publicly ₈₂₅
available on Data Dryad. Thank you to Julie Cridland and Nicholas Saleh for early tips ₈₂₆
on sampling and California bees, to Philipp Brand, Brenda Cameron, and Anne Lorant ₈₂₇
for sharing protocols and advice on lab work, to Jodie Jacobs and Petra Silverman for ₈₂₈
help phenotyping wing traits, and to Kelsey Lyberger, Roisin McMullen and Jodie ₈₂₉
Jacobs for their help in the field netting bees. Thank you to Daniela Zarate, Jeffrey ₈₃₀
Ross-Ibarra, Michael Turelli, Peter Ralph and to members of the Coop, Ramírez, and ₈₃₁
Ralph labs for insightful discussions and feedback on earlier drafts of the manuscript. ₈₃₂

References

1. Teeter KC, Payseur BA, Harris LW, Bakewell MA, Thibodeau LM, O'Brien JE, et al. Genome-wide patterns of gene flow across a house mouse hybrid zone. Genome Research. 2008;18(1):67–76.

2. Tavares H, Whibley A, Field DL, Bradley D, Couchman M, Copsey L, et al. Selection and gene flow shape genomic islands that control floral guides. Proceedings of the National Academy of Sciences USA. 2018;5:201801832.

3. Powell DL, García-Olazábal M, Keegan M, Reilly P, Du K, Díaz-Loyo AP, et al. Natural hybridization reveals incompatible alleles that cause melanoma in swordtail fish. Science. 2020;368(6492):731–736.

4. Hodgson JA, Pickrell JK, Pearson LN, Quillen EE, Prista A, Rocha J, et al. Natural selection for the Duffy-null allele in the recently admixed people of Madagascar. Proceedings Biological sciences. 2014;281:20140930.

5. Hufford MB, Lubinksy P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J. The genomic signature of crop-wild introgression in maize. PLoS Genetics. 2013;9(5):e1003477.

6. Fitzpatrick BM, Johnson JR, Kump DK, Smith JJ, Voss SR, Shaffer HB. Rapid spread of invasive genes into a threatened native species. Proceedings of the National Academy of Sciences USA. 2010;107(8):3606–3610.

7. Bay RA, Taylor EB, Schluter D. Parallel introgression and selection on introduced alleles in a native species. Molecular Ecology. 2019;28(11):2802–2813.

8. Winston ML. The biology and management of Africanized honey bees. Annual Review of Entomology. 1992;37:173–193.

9. Woyke J. Experiences with Apis mellifera adansonii in Brazil and in Poland. Apiacta. 1973;.

10. Villa JD, Koeniger N, Rinderer TE. Overwintering of Africanized, European, and hybrid honey bees in Germany. Environmental Entomology. 1991;20(1):39–43.

11. Taylor Jr OR, Spivak M. Climatic limits of tropical African honeybees in the Americas. Bee World. 1984;65(1):38–47.

12. Harrison JF, Fewell JH, Anderson KE, Loper GM. Environmental physiology of the invasion of the Americas by Africanized honeybees. Integrative and Comparative Biology. 2006;46(6):1110–1122.

13. Southwick EE, Roubik DW, Williams JM. Comparative energy balance in groups of Africanized and European honey bees: ecological implications. Comparative Biochemistry and Physiology. 1990;97(1):1–7.

14. Sheppard WS, Rinderer TE, Mazzoli JA, Stelzer JA, Shimanuki H. Gene flow between African- and European-derived honey bee populations in Argentina. Nature. 1991;349(6312):782–784.

15. Agra MN, Conte CA, Corva PM, Cladera JL, Lanzavecchia SB, Palacio MA. Molecular characterization of Apis mellifera colonies from Argentina: genotypic admixture associated with ecoclimatic regions and apicultural activities. Entomologia Experimentalis et Applicata. 2018;166(9):724–738.

16. Pinto MA, Rubink WL, Patton JC, Coulson RN, Johnston JS. Africanization in the United States: replacement of feral European honeybees (*Apis mellifera* L.) by an African hybrid swarm. Genetics. 2005;170(4):1653–1665.

17. Loper GM, Fewell J, Smith DR, Sheppard WS, Schiff N. Changes in the genetics of a population of feral honey bees (Apis mellifera L.) in S. Arizona after the impact of tracheal mites (Acarapis woodi), Varroa mites (Varroa jacobsoni) and Africanization. In: Hoopingarner R, Connor L, editors. Apiculture for the 21st Century. Cheshire, CT: Wicwas; 1999. p. 47–51.

18. Kono Y, Kohn JR. Range and frequency of Africanized honey bees in California (USA). PLoS ONE. 2015;10(9):e0137407.

19. Lin W, McBroome J, Rehman M, Johnson BR. Africanized bees extend their distribution in California. PLoS ONE. 2018;13(1):e0190604.

20. Kadri SM, Harpur BA, Orsi RO, Zayed A. A variant reference data set for the Africanized honeybee, Apis mellifera. Scientific Data. 2016;3:160097.

21. Wallberg A, Han F, Wellhagen G, Dahle B, Kawata M, Haddad N, et al. A worldwide survey of genome sequence variation provides insight into the evolutionary history of the honeybee *Apis mellifera*. Nature Genetics. 2014;46(10):1081–1088.

22. Cridland JM, Ramírez SR, Dean CA, Sciligo A, Tsutsui ND. Genome sequencing of museum specimens reveals rapid changes in the genetic composition of honey bees in California. Genome Biology and Evolution. 2018;10(2):458–472.

23. Whitfield CW, Behura SK, Berlocher SH, Clark AG, Johnston JS, Sheppard WS, et al. Thrice out of Africa: ancient and recent expansions of the honey bee, *Apis mellifera*. Science. 2006;314(5799):642–645.

24. Bozek K, Rangel J, Arora J, Tin M, Crotteau E, Loper G, et al. Parallel genomic evolution of parasite tolerance in wild honey bee populations. bioRxiv. 2018;doi:10.1101/498436.

25. Nelson RM, Wallberg A, Simões ZLP, Lawson DJ, Webster MT. Genome-wide analysis of admixture and adaptation in the Africanized honeybee. Molecular Ecology. 2017;26:3603–3617.

26. Hunt GJ, Amdam GV, Schlipalius D, Emore C, Sardesai N, Williams CE, et al. Behavioral genomics of honeybee foraging and nest defense. Die Naturwissenschaften. 2007;94(4):247–267.

27. Hunt GJ, Guzman-Novoa E, Fondrk MK, Page RE. Quantitative trait loci for honey bee stinging behavior and body size. Genetics. 1998;148(3):1203–1213.

28. Danka RG, Rinderer TE, Hellmich RL, Collins AM. Comparative toxicities of four topically applied insecticides to Africanized and European honey bees (Hymenoptera: Apidae). Journal of Economic Entomology. 1986;79(1):18–21.

29. Guzman-Novoa E, Vandame R, Arechavaleta ME. Susceptibility of European and Africanized honey bees (Apis mellifera L.) to Varroa jacobsoni Oud. in Mexico. Apidologie. 1999;30(2-3):173–182.

30. Vandame R, Morand S, Colin ME, Belzunces LP. Parasitism in the social bee Apis mellifera: quantifying costs and benefits of behavioral resistance to Varroa destructor mites. Apidologie. 2002;33(5):433–445.

31. Guerra J, Goncalves LS, De Jong D. Africanized honey bees (Apis mellifera L.) are more efficient at removing worker brood artificially infested with the parasitic mite Varroa jacobsoni Oudemans than are Italian bees or Italian/Africanized hybrids. Genetics and Molecular Biology. 2000;23(1):89–92.

32. Moretto G, de Mello LJ. Varroa jacobsoni infestation of adult Africanized and Italian honey bees (Apis mellifera) in mixed colonies in Brazil. Genetics and Molecular Biology. 1999;22(3):321–323.

33. Medina-Flores CA, Guzman-Novoa E, Hamiduzzaman MM, Aréchiga-Flores CF, López-Carlos MA. Africanized honey bees (*Apis mellifera*) have low infestation levels of the mite Varroa destructor in different ecological regions in Mexico. Genetics and Molecular Research. 2014;13(3):7282–7293.

34. Daly HV, Balling SS. Identification of Africanized honeybees in the Western Hemisphere by discriminant analysis. Journal of the Kansas Entomological Society. 1978;.

35. Danka RG, Hellmich RL, Rinderer TE, Collins AM. Diet-selection ecology of tropically and temperately adapted honey-bees. Animal Behaviour. 1987;35(6):1858–1863.

36. Fewell JH, Bertram SM. Evidence for genetic variation in worker task performance by African and European honey bees. Behavioral Ecology and Sociobiology. 2002;52(4):318–325.

37. Skotte L, Korneliussen TS, Albrechtsen A. Estimating individual admixture proportions from next generation sequencing data. Genetics. 2013;195(3):693–702.

38. Corbett-Detig R, Nielsen R. A Hidden Markov Model Approach for Simultaneously Estimating Local Ancestry and Admixture Time Using Next Generation Sequence Data in Samples of Arbitrary Ploidy. PLoS Genetics. 2017;13(1):e1006529.

39. Kent RB. The introduction and diffusion of the African honeybee in South America. Yearbook of the Association of Pacific Coast Geographers. 1988;50(1):21–43.

40. USDA Agricultural Research Service. Spread of Africanized honey bees by year, by county; 2009. Available from: `https://www.ars.usda.gov/ARSUserFiles/20220500/New%20Bee%20Map09%20compressed.jpg`.

41. Abrahamovich AH, Atela O, De la Rúa P, Galián J. Assessment of the mitochondrial origin of honey bees from Argentina. Journal of Apicultural Research. 2015;46(3):191–194.

42. Simmons AD, Thomas CD. Changes in Dispersal during Species' Range Expansions. American Naturalist. 2015;164(3):378–395.

43. Cwynar LC, MacDonald GM. Geographical Variation of Lodgepole Pine in Relation to Population History. American Naturalist. 1987;129(3):463–469.

44. Phillips BL, Brown GP, Webb JK, Shine R. Invasion and the evolution of speed in toads. Nature. 2006;439(7078):803–803.

45. Hill JK, Thomas CD, Blakeley DS. Evolution of flight morphology in a butterfly that has recently expanded its geographic range. Oecologia. 1999;121(2):165–170.

46. Daly HV, Hoelmer K, Gambino P. Clinal geographic variation in feral honey bees in California, USA. Apidologie. 1991;22(6):591–609.

47. Wang S, Rohwer S, Delmore K, Irwin DE. Cross-decades stability of an avian hybrid zone. Journal of Evolutionary Biology. 2019;32(11):1242–1251.

48. Szymura JM, Barton NH. Genetic analysis of a hybrid zone between the fire-bellied toads, Bombina bombina and B. variegata, near Cracow in southern Poland. Evolution. 1986;40(6):1141.

49. Szymura JM, Barton NH. The genetic structure of the hybrid zone between the fire-bellied toads Bombina bombina and B. variegata: Comparisons between transects and between loci. Evolution. 1991;45(2):237.

50. Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution interpolated climate surfaces for global land areas. International Journal of Climatology. 2005;25(15):1965–1978.

51. Gompert Z, Mandeville EG, Buerkle CA. Analysis of Population Genomic Data from Hybrid Zones. Annual Review of Ecology, Evolution, and Systematics. 2017;48(1):207–229.

52. Barton NH. Multilocus Clines. Evolution. 1983;37(3):454.

53. Schneider SS, DeGrandi-Hoffman G, Smith DR. The African honey bee: Factors contributing to a successful biological invasion. Annual Review Entomology. 2004;49(1):351–376.

54. Excoffier L, Foll M, Petit RJ. Genetic consequences of range expansions. Annual Review of Ecology, Evolution, and Systematics. 2009;40(1):481–501.

55. Tsuruda JM, Harris JW, Bourgeois L, Danka RG, Hunt GJ. High-resolution linkage analyses to identify genes that influence *Varroa* sensitive hygiene behavior in honey bees. PLoS ONE. 2012;7(11):e48276.

56. Oxley PR, Spivak M, Oldroyd BP. Six quantitative trait loci influence task thresholds for hygienic behaviour in honeybees (*Apis mellifera*). Molecular Ecology. 2010;19(7):1452–1461.

57. Spötter A, Gupta P, Nuernberg G, Reinsch N, Bienefeld K. Development of a 44K SNP assay focussing on the analysis of a varroa-specific defence behaviour in honey bees (*Apis mellifera carnica*). Molecular Ecology Resources. 2012;12(2):323–332.

58. Arechavaleta-Velasco ME, Alcala-Escamilla K, Robles-Rios C, Tsuruda JM, Hunt GJ. Fine-scale linkage mapping reveals a small set of candidate genes influencing honey bee grooming behavior in response to *Varroa* mites. PLoS ONE. 2012;7(11):e47269.

59. McDonnell CM, Alaux C, Parrinello H, Desvignes JP, Crauser D, Durbesson E, et al. Ecto- and endoparasite induce similar chemical and brain neurogenomic responses in the honey bee (Apis mellifera). BMC Ecology. 2013;13(1):1–15.

60. Surlis C, Carolan JC, Coffey M, Kavanagh K. Quantitative proteomics reveals divergent responses in Apis mellifera worker and drone pupae to parasitization by Varroa destructor. Journal of Insect Physiology. 2018;107:291–301.

61. Buggs R. Empirical study of hybrid zone movement. Heredity. 2007;99(3):301–312.

62. Taylor SA, Larson EL, Harrison RG. Hybrid zones: windows on climate change. Trends in Ecology & Evolution. 2015;30(7):398–406.

63. Currat M, Ruedi M, Petit RJ, Excoffier L. The hidden side of invasions: massive introgression by local genes. Evolution. 2008;62(8):1908–1920.

64. Barton N, Bengtsson BO. The barrier to genetic exchange between hybridising populations. Heredity. 1986;56:357–376.

65. Good TP, Ellis JC, Annett CA, Pierotti R. Bounded hybrid superiority in an avian hybrid zone: effects of mate, diet, and habitat choice. Evolution. 2000;54(5):1774–1783.

66. De La Torre AR, Wang T, Jaquish B, Aitken SN. Adaptation and exogenous selection in a Picea glauca × Picea engelmannii hybrid zone: implications for forest management under climate change. New Phytologist. 2014;201(2):687–699.

67. Adrion JR, Hahn MW, Cooper BS. Revisiting classic clines in Drosophila melanogaster in the age of genomics. Trends in genetics : TIG. 2015;31(8):434–444.

68. Rinderer TE, Sylvester HA, Brown MA, Villa JD, Pesante D, Collins AM. Field and simplified techniques for identifying Africanized and European honey bees. Apidologie. 1986;17(1):13–48.

69. Jones JC, Wallberg A, Christmas MJ, Kapheim KM, Webster MT, Singh N. Extreme differences in recombination rate between the genomes of a solitary and a social bee. Molecular Biology and Evolution. 2019;36(10):2277–2291.

70. Hunt GJ, Guzmán-Novoa E, Uribe-Rubio JL, Prieto-Merlos D. Genotype–environment interactions in honeybee guarding behaviour. Animal Behaviour. 2003;66(3):459–467.

71. Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science. 2015;347(6229):1255957–1255957.

72. R Core Team. R: A Language and Environment for Statistical Computing; 2019. Available from: https://www.R-project.org/.

73. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to the tidyverse. Journal of Open Source Software. 2019;4(43):1686. doi:10.21105/joss.01686.

74. Tange O. GNU Parallel 2018. Ole Tange; 2018. Available from: https://doi.org/10.5281/zenodo.1146014.

75. Rowan BA, Heavens D, Feuerborn TR, Tock AJ, Henderson IR, Weigel D. An ultra high-density Arabidopsis thaliana crossover map that refines the influences of structural variation and epigenetic features. Genetics. 2019;213(3):771–787.

76. Harpur BA, Kent CF, Molodtsova D, Lebon JMD, Alqarni AS, Owayss AA, et al. Population genomics of the honey bee reveals strong signatures of positive selection on worker traits. Proceedings of the National Academy of Sciences USA. 2014;111(7):2614–2619.

77. Cridland JM, Tsutsui ND, Ramírez SR. The complex demographic history and evolutionary origin of the western honey bee, *Apis mellifera*. Genome Biology and Evolution. 2017;9(2):457–472.

78. Wallberg A, Bunikis I, Pettersson OV, Mosbech MB, Childers AK, Evans JD, et al. A hybrid de novo genome assembly of the honeybee, *Apis mellifera*, with chromosome-length scaffolds. BMC Genomics. 2019;20(1):275.

79. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature methods. 2012;9(4):357–359.

80. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics (Oxford, England). 2009;25(16):2078–2079.

81. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation Sequencing Data. BMC bioinformatics. 2014;15(1):356.

82. Meisner J, Albrechtsen A. Inferring Population Structure and Admixture Proportions in Low-Depth NGS Data. Genetics. 2018;210(2):719–731.

83. Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. Estimating Kinship in Admixed Populations. The American Journal of Human Genetics. 2012;91(1):122–138.

84. Long JC. The genetic structure of admixed populations. Genetics. 1991;127(2):417–428.

85. Venables WN, Ripley BD. Modern Applied Statistics with S. 4th ed. New York: Springer; 2002. Available from: http://www.stats.ox.ac.uk/pub/MASS4.

86. Hong Y. poibin: The Poisson Binomial Distribution; 2019. Available from: https://CRAN.R-project.org/package=poibin.

87. Hijmans RJ. geosphere: Spherical Trigonometry; 2019. Available from: https://CRAN.R-project.org/package=geosphere.

88. Bickel PJ, Boley N, Brown JB, Huang H, Zhang NR. Subsampling methods for genomic inference. The Annals of Applied Statistics. 2010;4(4):1660–1697.

89. Padfield D, Matheson G. nls.multstart: Robust Non-Linear Regression using AIC Scores; 2018. Available from: https://CRAN.R-project.org/package=nls.multstart.

90. Ruttner F. Biogeography and Taxonomy of Honeybees. Berlin: Springer; 1988.

91. Grinde KE, Brown LA, Reiner AP, Thornton TA, Browning SR. Genome-wide significance thresholds for admixture mapping studies. American Journal of Human Genetics. 2019;104(3):454–465.

92. Siegmund D, Yakir B. The Statistics of Gene Mapping. Springer Science & Business Media; 2007.

93. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics (Oxford, England). 2010;26(6):841–842.

94. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nature Protocols. 2009;4(1):44–57.

95. Benjamini, Yoav, Hochberg, Yosef. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society. 1995;57(1):289–300.

96. Arechavaleta-Velasco ME, Hunt GJ, Emore C. Quantitative trait loci that influence the expression of guarding and stinging behaviors of individual honey bees. Behavior Genetics. 2003;33(3):357–364.

97. Solignac M, Mougel F, Vautrin D, Monnerot M, Cornuet JM. A third-generation microsatellite-based linkage map of the honey bee, Apis mellifera, and its comparison with the sequence-based physical map. Genome biology. 2007;8(4):R66.

98. Harpur B. Hunt honey bee markers; 2020. Dryad. Available from: https://doi.org/10.5061/dryad.ns1rn8ppp.

99. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. Journal of molecular biology. 1990;215(3):403–410.

100. Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting FST: the impact of rare variants. Genome Research. 2013;23(9):1514–1521.

101. Honeybee Genome Sequencing Consortium. Insights into social insects from the genome of the honeybee *Apis mellifera*. Nature. 2006;444(7118):512–512.

102. Miller CA, Hampton O, Coarfa C, Milosavljevic A. ReadDepth: A Parallel R Package for Detecting Copy Number Alterations from Short Sequencing Reads. PLoS ONE. 2011;6(1):e16327.

Supporting information

S1 Table.   Sample information. Geographic sampling information (population, location, date, whether collected by a feral nest), approximate sequencing coverage, global ancestry estimates and wing lengths for bees sequenced in this study and reference bees.

S2 Table.   Cline model comparison. Model rankings between logistic cline fits for genome-wide African ancestry predicted by climate and distance variables.

S3 Table.   Ancestry outlier regions. Genome coordinates for outlier regions with high or low African (A) ancestry. Adjacent and near adjacent (within 10kb) ancestry windows meeting <10% FDR have been combined into contiguous regions and are labelled with the lowest FDR within the region. Note that shared high A outlier regions, by definition, will overlap high A South American and high A North American outlier regions, with bp and percent overlap listed. NA signifies not significant for that hybrid zone.

S4 Table.   Ancestry outlier genes. List of genes overlapping ancestry outliers at 1%, 5%, and 10% FDR thresholds. Minimum FDR for each continent listed separately. NA signifies not significant for that hybrid zone.

S5 Table.   Approximate QTL coordinates HAv3.1. Approximate coordinates (HAv3.1) for regions of the genome previously associated with defensive behaviors or *Varroa* tolerance.

S6 Table.   Africanized honey bee invasion historical dates and locations. Approximate locations and dates of first arrival for the spread of Africanized honey bees as plotted in Fig 1. We estimated GPS coordinates for each historical observation using google maps and the available location description.

Fig S1. PCA. Principal components analysis generated in PCAngsd using genotype likelihoods from the same thinned set of 14,044 autosomal SNPs used in global admixture analysis. The major axes of diversity separate out C ancestry (PC1) and M ancestry (PC2). Consistent with 3-way admixture, all sampled bees from North and South America are intermediate on the PCA, in the triangle formed by reference panels for African (A), eastern European (C) and western European (M) honey bees.

Fig S2. Ancestry in reference panels. Results of NGSAdmix global admixture analysis for reference populations from the combined analysis of all populations (K = 3). These results were used to assign the unlabelled ancestry components output by NGSAdmix to "A", "C", and "M" groups, based on a clear mapping to the three reference populations. We see a small amount of admixture between C and M within our reference populations, which is consistent with limited gene flow from secondary contact within Europe.

Fig S3. Comparison of local and global ancestry results. (A) Comparison of the mean genome-wide ancestry estimate from NGSAdmix (x-axis) and ancestry_hmm (y-axis) for each bee, with one-to-one line drawn in grey. The mean for the HMM is calculated by marginalizing the posterior over all ancestry states and taking a mean across SNPs. The individual-level ancestry estimates between the two methods agree strongly (Pearson's correlation: 0.997 A, 0.999 C, 0.985 M), but the HMM estimates slightly higher minor ancestry for bees with low admixture proportions. (B) Population mean summarises for the same comparison of NGSAdmix vs. ancestry_hmm genome-wide ancestry estimates, with one-to-one line drawn in gray. Because the population mean ancestry proportions from NGSAdmix are used as a prior for the population-specific mixing proportions in ancestry_hmm, this panel can also be interpreted as the prior (x-axis) and posterior (y-axis) of the local ancestry HMM for population-level admixture proportions.

Fig S4. Power to call local ancestry. (A) Correlation between high-coverage and low-coverage ancestry calls, across different simulated depths of coverage (1-10x). (B) Proportion of high-confidence ancestry calls from high-coverage data that were replicated in analyses of low-coverage data, with different simulated depths of coverage (1-10x). These results are from a preliminary analysis of the power to call local ancestry accurately, used to inform target sequencing depth for this study. For this preliminary study, we used a published SNP set with data for A, C, and M reference populations [77] based on earlier versions of the honey bee genome (Amel4.5 [101]) and recombination map [21]. We enriched for ancestry-informativeness and thinned for linkage disequilibrium (≥ 0.2 MAF in at least one reference population and $r^2 < 0.4$ within the A reference population), leaving 161k SNPs. First we ran ancestry_hmm [38] using called genotypes from a high-coverage admixed population with intermediate admixture proportions (Riverside 2014 (n=8): 40% C, 20% M, 40% A ancestry). We simulated lower coverage data from this same population by generating a binomial sample of n reads for each locus, based on the individual's genotype. To simulate realistic variance in coverage across the genome, n for each site and individual was generated from a negative binomial distribution with variance 3x the mean [102]. We additionally simulated a 1% sequencing error rate. Running local ancestry inference on the high coverage data, we inferred high confidence ancestry states for 81% of sites. First we calculated a point estimates for African ancestry ($p(AA) + 1/2(p(CA) + p(MA))$) at every site for each individual and used these estimates to calculate a correlation between the high coverage ancestry calls and low coverage ancestry calls. Then we calculated the percent of high confidence calls that were replicated with high confidence (>0.8 posterior) in the low coverage data for the same ancestry state ("correct") or a different ancestry state ("incorrect"). Call to the HMM for simulated low-coverage data: ancestry_hmm -e 3e-3 -a 3 0.4 0.2 0.4 -p 0 100000 0.4 -p 1 -100 0.2 -p 2 -60 0.4 –tmax 150 –tmin 2 –ne 670000. For original high coverage data we used genotype calls rather than read counts (-g) and a lower error rate (-e 1e-3).

Fig S5. Estimated generations post-admixture. Inferred timing of migration pulses from A ancestry (left) and M ancestry (right). Each population's admixture timing is estimated separately, during local ancestry inference (ancestry_hmm), and results are plotted across latitude. We allowed a range of 2-150 generations, so the highest time estimates are truncated at 150 generations. Admixture with African (A) ancestry began in 1956, 62 years before sampling in 2018. We have little prior information about the timing of M into C admixture, which likely varies across the Americas, but in general should pre-date Africanization. The number of generations per year for feral honey bee populations is uncertain.

Fig S6. Climate variables across latitude. Bioclim climate variables for all sample sites plotted against latitude: (A) Mean annual temperature (B) Mean annual precipitation (C) Mean temperature coldest quarter (D) Minimum temperature coldest month. Two adjacent climate outliers in the N. American sample can be seen in the top two panels and represent bees from an inland desert (hot and dry) and a high altitude sampling site (cold and wet) at similar latitudes in Riverside County, CA. Bees from this same high altitude site are also outliers in the bottom two panels, having the coldest mean and minimum winter temperatures of all sites.

Fig S7. Wing length predicted by African ancestry. Individual honey bees are represented as points, with wing lengths plotted along the x-axis and genonewide African ancestry proportions (NGSAdmix results) along the y-axis. We draw the best-fit regression line (slope = -0.72 mm, $F(1, 267) = 119$, $P = 3.65 \times 10^{-23}$, $R^2 = 0.31$, $n = 269$). We also include wing lengths for European (C, M) and African (A) reference bees from the Oberursel Collection, which we assume have none or full A ancestry. These reference bees are plotted slightly outside the range [0,1] and with jitter to facilitate viewing individual points that would otherwise all cluster on the boundaries.

Fig S8. Wing length measurement. Fore wing image cropped and annotated to show length measurement taken. A full length to the tip of the wing is the standard measurement, but we use this alternative because many of our samples have significant wing tatter.

Fig S9. Admixture mapping analysis. We plot the p-value for each SNP across the genome, based on independent tests of association between African ancestry at that SNP and wing length. The red dashed line marks the genome-wide significance threshold for a family-wise error rate of 0.05, using a two-tailed test. In admixture mapping, SNPs are correlated, and the number of independent statistical tests depends on the number of generations recombination has had to break up ancestry blocks. Here we use an analytical approximation for the significance threshold based on 47.65 generations of admixture (population median estimate).

Fig S10. Distribution of ancestry clines in South America across SNPs. A logistic cline model was fit to observed and simulated population ancestry frequencies across latitude for S. America. Estimated cline parameters, center and width ($w = |4/b|$), are presented as violin plots. Units for both cline center and width are degrees latitude. We additionally partition observed SNPs by outlier and non-outlier status, set by 10% FDR for high or low African ancestry in South America. Individual SNP clines were only fit in South America, where we observed the full cline.

Fig S11. Effect of truncating MVN simulated ancestry frequencies. (A) Percent of simulated population A-ancestry frequencies exceeding lower (left) and upper (right) bounds, and thus truncated to [0,1] range. Each population is a point and the populations most affected by truncation are low-A ancestry populations with mean A-ancestry proportions close to the bound at 0. (B) QQ-plot comparing the quantiles for mean A ancestry before and after truncation in N. America (left) and S. America (right). The distribution of mean African ancestry is mostly unaffected by restricting simulated population A ancestry frequencies to the [0,1] range, but truncation does reduce model predictions of very low A-ancestry frequencies in N. America, where mean A ancestry is already low.

Fig S12. Simulated vs. observed African ancestry quantiles. QQ-plots comparing observed quantiles for mean African ancestry in North America (top) and South America (bottom) to the quantiles generated by three simulated distributions (left-to-right): MVN, MVN with zero covariances, and Poisson binomial model. Only the MVN model, allowing for covariances between populations, matches the bulk of the observed distribution.

Fig S13. Overlay of 2D density plot for observed A ancestry frequencies in North and South America compared to simulated A ancestry frequencies under a multivariate-normal model.

Fig S14. Mean ancestry correlation by chromosome across populations.
Mean ancestry covariances (K matrices) were calculated separately for chromosome,
using the genome-wide mean ancestry as α, then correlations were summarised by
taking the mean for each type of population comparison, within and between continents
and low vs. high A regions. Error bars show the normal-approximated 95% confidence
intervals around these means. We divided populations in South America into two
groups based on having greater than ('high A') or less than ('low A') 50% mean African
ancestry. Low A populations are found at higher latitudes and correspondingly cooler
climates. All North American samples come from the low-A side of the cline. On
average across chromosomes, low-A South American populations share higher ancestry
correlations with low-A North American populations than with geographically closer
high-A South American populations (0.032: CI$_{95}$[0.011, .053], P = .005, paired 2-sided
t-test). Two chromosomes harbor large outlier regions consistent with their elevated
correlations shown here: Chromosome 1 has a large cluster of loci with high A ancestry
in North and South America while chromosome 11 has a wide region of low A ancestry
exclusive to South America. The results do not change qualitatively if these two outlier
chromosomes are both removed from the analysis (0.035: CI$_{95}$[0.022, 0.047], P =
$4.8x10^{-5}$, paired 2-sided t-test).

Fig S15. Mean correlation for population pairs by recombination rate.
Mean ancestry covariances (K matrices) were calculated separately for each of the 5 recombination rate quintiles, using the genome-wide mean ancestry as α, then correlations were summarised by taking the mean for each type of population comparison, within and between continents and low vs. high A regions. About half of the South American populations, and all of the sampled North American populations come from the low-A side of the hybrid zone (relative to the estimated cline center). The genomewide mean is additionally shown as an X. On average across recombination bins, low-A South American populations share higher ancestry correlations with low-A North American populations than with geographically closer high-A South American populations (0.049: $CI_{95}[0.021, .078]$, P = .009, paired 2-sided t-test)

Fig S16. Ancestry covariances across populations. African ancestry covariance
matrix (see methods). Populations are ordered by latitude, with high and low A sides of
each hybrid zone defined relative to the estimated genomewide cline center.
Within-population variances (left) are shown separately from between-population
covariances (right) because of the drastically different scales. Populations near the cline
center have higher ancestry variances (most clearly seen in the diagonal elements)
because they have African ancestry proportions closer to 50%. Drift and finite sample
sizes also contribute to the observed ancestry variances. The third lowest latitude
population in the North American cline with exceptionally high ancestry variance is
Avalon, sampled from Catalina Island off the coast of California. The observed excess
covariance between the two distant ends of these clines is unexpected under a simple
model of spread North and South out of Brazil.

Fig S17. Genetic covariance within ancestry. Genetic covariances within A (top), C (middle) and M (bottom) ancestry. Colors represent the population mean genetic covariance between individuals, and the range of values varies by ancestry (note: color bars have different scales). While kinship creates strictly positive covariances, here we observe some negative values because we can only calculate co-variation around the empirical mean combined sample allele frequency, not the true ancestral allele frequency (which is unknown). Population mean covariances were summarised from an individual-by-individual covariance matrix generated using PCAngsd from bam files filtered to only include regions of the genome with high confidence homozygous ancestry calls for the focal ancestry (posterior >0.8 from ancestry_hmm). High and low African sides of each hybrid zone are defined relative to the estimated genomewide cline center.

Fig S18. Principle components analysis of genetic variation within ancestry.
PCA analysis of A (top), C (middle) and M (bottom) ancestry. Analysis was performed
using PCAngsd using all reference samples of the focal ancestry and sequence data from
the hybrid zones filtered to only include regions of the genome with high confidence
homozygous ancestry calls for the focal ancestry (posterior >0.8 from ancestry_hmm).
Each bee is a point, colored by sample location. The two hybrid zones form somewhat
separable clusters for European (C and M) ancestry, but not African (A) ancestry. The
major axis of genetic diversity within M ancestry in the Americas (PC1) mirrors
pre-existing population structure within Europe between *Apis mellifera mellifera*
(Poland) and *Apis mellifera iberiensis* (Spain), two well-known honeybee subspecies
that may have different historical import rates to different regions.

Fig S19. Genetic variation within ancestry by latitude. Here we plot the first principal component for genetic diversity within A (top), C (middle) and M (bottom) ancestry against absolute latitude of sampling location within the hybrid zone (see Fig S18 for original PCA). Each bee is a point and reference bees are plotted to the side (not at their actual latitude). Bees are colored by sample location. Note that bees with very low amounts of the focal ancestry (higher latitudes A ancestry or lower latitudes C ancestry) fall close to zero on PC1 (dashed line), which, despite using a method designed to account for low coverage data (PCAngsd), may simply be an artifact of low information per individual bee for genetic diversity within a low-frequency ancestry. A ancestry shows very little population structure along PC1 by continent or latitude. C is the dominant ancestry at higher latitudes in both zones and shows greater separation between the two ends of the zones (higher absolute latitude for both) than within South America. M ancestry at lower latitudes in South America is more similar to *Apis mellifera iberiensis* (Spain) than M ancestry elsewhere in the Americas.

Fig S20. Ancestry and AIM frequencies for high shared A outliers on chr1.
Zoomed-in view of the region on chromosome 1 with a cluster of high African ancestry
peaks in both North America (left) and South America (right), with shared outlier
regions meeting a 10% FDR for high African ancestry on both continents shaded in grey.
(Top) African (A), western European (M) and eastern European (C) local ancestry
estimates at each HMM marker. (Bottom) Mean frequency of ancestry informative
markers AIMs (see methods), most of which were not included in the ancestry_hmm
inference ('AIM only') due to thinning.

Fig S21. Ancestry and AIM frequencies for low A outlier region on chr11.
Zoomed-in view of the 1.4Mb region on chromosome 11 with high western European
ancestry (M) in South America (right) but not in North America (left), with the outlier
region meeting 10% FDR highlighted in grey. (Top) African (A), western European (M)
and eastern European (C) local ancestry estimates at each HMM marker. (Bottom)
Mean frequency of ancestry informative markers AIMs (see methods), most of which
were not included in the ancestry_hmm inference ('AIM only') due to thinning.

Fig S22. Differentiation across shared high A outliers on chr1. F_{ST} across the region on chromosome 1 with shared high African ancestry outliers. Outlier regions meeting a 10% FDR in both hybrid zones are highlighted in darker grey, while those meeting a 10% FDR in only one hybrid zone are highlighted in lighter grey. Per-SNP F_{ST} is averaged within sliding 50kb windows. In addition to the three ancestry reference panels (A, C, & M), we include contrasts for the subset of individuals in each hybrid zone with high-confidence homozygous A ancestry.

Fig S23. Differentiation across shared high A outliers on chr11. F_{ST} across the region on chromosome 11 with shared high African ancestry outliers. Outlier regions meeting a 10% FDR in both hybrid zones are highlighted in darker grey, while those meeting a 10% FDR in only one hybrid zone are highlighted in lighter grey. Per-SNP F_{ST} is averaged within sliding 50kb windows. In addition to the three ancestry reference panels (A, C, & M), we include contrasts for the subset of individuals in each hybrid zone with high-confidence homozygous A ancestry.

Fig S24. Differentiation across low A outlier region on chr11. F_{ST} across the
1.4Mb region on chromosome 11 with high western European ancestry (M) in South
America but not in North America (left), with the outlier region meeting 10% FDR
highlighted in grey. Per-SNP F_{ST} is averaged within sliding 50kb windows. In addition
to the three ancestry reference panels (A, C, & M), we include contrasts for the subset
of individuals in each hybrid zone with high-confidence homozygous M ancestry.
Windows are dropped if fewer than 10 SNPs have 2 individuals with data, which
produces gaps in the contrasts with N. American M ancestry because this hybrid zone
does not have elevated M ancestry in this region and at many SNPs very few
individuals have high-confidence homozygous M ancestry. Top peaks in M vs. C, M vs.
A, and C vs. A contrasts seen within this region reach the 99.5, 98.0, and 99.8
percentiles (respectively) for 50kb windows genome-wide.

Fig S25. Comparison of observed and predicted diversity. (A) Observed and predicted allelic diversity (π) for each population across latitude. To predict π for a specific population, we calculated the expected allele frequency based on a mixture of A, C, and M reference population allele frequencies, weighted by the population's estimated admixture fractions of these three ancestries. (B) Plot of predicted vs. observed π for each population for direct comparison to the 1-to-1 line (dashed). Avalon (California 2014, marked as a triangle) is a clear outlier, with low diversity for its admixture fraction.

Fig S26. Mitochrondrial clines. Out of 82 SNPs on the mitochondria, we
identified two with more than 80% estimated frequency difference between African (A)
and European (C & M) reference panels. Estimated allele frequencies at these SNPs for
each population in North America (left) and South America (right) are plotted in color.
For comparison, population mean genomewide African ancestry proportions
(NGSAdmix) are plotted as open black circles. At both SNPs, estimated M and C allele
frequencies are zero (not shown) and estimated A allele frequencies are high but not at
fixation (plotted as pink dashed lines). Bees sequenced in this study have low coverage
across most of the mtDNA sequence, which prevented us from constructing a
phylogenetic tree for full mitochondrial haplotypes and creates uncertainty in our allele
frequency estimates here. To reflect this uncertainty, points are shaded by the sample
size (i.e. the number of mtDNA haplotypes in a population for which we were able to
call a consensus base).