Anaerobic sulfur oxidation underlies adaptation of a chemosynthetic symbiont to

oxic-anoxic interfaces

Running title: chemosynthetic ectosymbiont's response to oxygen

Gabriela F. Paredes[1], Tobias Viehboeck[1,2], Raymond Lee[3], Marton Palatinszky[2], Michaela A.

Mausz[4], Sigfried Reipert[5], Arno Schintlmeister[2,6], Jean-Marie Volland[1,‡], Claudia Hirschfeld[7]

Michael Wagner[2,8], David Berry[2,9], Stephanie Markert[7], Silvia Bulgheresi[1,*] and Lena König[1,*]

[1] University of Vienna, Department of Functional and Evolutionary Ecology, Environmental

Cell Biology Group, Vienna, Austria

[2] University of Vienna, Center for Microbiology and Environmental Systems Science, Division

of Microbial Ecology, Vienna, Austria

[3] Washington State University, School of Biological Sciences, Pullman, WA, USA

[4] University of Warwick, School of Life Sciences, Coventry, UK

[5] University of Vienna, Core Facility Cell Imaging and Ultrastructure Research, Vienna,

Austria

[6] University of Vienna, Center for Microbiology and Environmental Systems Science, Large-

Instrument Facility for Environmental and Isotope Mass Spectrometry, Vienna, Austria

[7] University of Greifswald, Institute of Pharmacy, Department of Pharmaceutical

Biotechnology, Greifswald, Germany

[8] Aalborg University, Department of Chemistry and Bioscience, Aalborg, Denmark

[9] Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna,

Vienna, Austria

[‡] Present affiliation: Joint Genome Institute/Global Viral, San Francisco, CA, USA

Corresponding Authors

Silvia Bulgheresi: University of Vienna, Department of Functional and Evolutionary Ecology,

Environmental Cell Biology Group, Althanstrasse 14, 1090 Vienna, Austria, +43-1-4277-

76514, silvia.bulgheresi@univie.ac.at

Lena König: University of Vienna, Department of Functional and Evolutionary Ecology,

Environmental Cell Biology Group, Althanstrasse 14, 1090 Vienna, Austria, +43-1-4277-

76527, lena.koenig@univie.ac.at

* These authors contributed equally to this work.

Competing Interests

The authors declare no competing financial interest.

Abstract

Chemosynthetic symbioses occur worldwide in marine habitats. However, physiological

studies of chemoautotrophic bacteria thriving on animals are scarce. *Stilbonematinae* are

coated by monocultures of thiotrophic Gammaproteobacteria. As these nematodes migrate

through the redox zone, their ectosymbionts experience varying oxygen concentrations.

Here, by applying omics, Raman microspectroscopy and stable isotope labeling, we

investigated the effect of oxygen on the metabolism of *Candidatus* Thiosymbion oneisti.

Unexpectedly, sulfur oxidation genes were upregulated in anoxic relative to oxic conditions,

but carbon fixation genes and incorporation of ^{13}C-labeled bicarbonate were not. Instead,

several genes involved in carbon fixation, the assimilation of organic carbon and

polyhydroxyalkanoate (PHA) biosynthesis, as well as nitrogen fixation and urea utilization

were upregulated in oxic conditions. Furthermore, in the presence of oxygen, stress-related

genes were upregulated together with vitamin and cofactor biosynthesis genes likely

necessary to withstand its deleterious effects.

Based on this first global physiological study of a chemosynthetic ectosymbiont, we

propose that, in anoxic sediment, it proliferates by utilizing nitrate to oxidize reduced sulfur,

whereas in superficial sediment it exploits aerobic respiration to facilitate assimilation of

carbon and nitrogen to survive oxidative stress. Both anaerobic sulfur oxidation and its

decoupling   from   carbon   fixation   represent   unprecedented   adaptations   among

chemosynthetic symbionts.

Introduction

At least six animal phyla and numerous lineages of bacterial symbionts belonging to Alpha-,

Delta-, Gamma- and Epsilonproteobacteria engage in chemosynthetic symbioses, rendering

the evolutionary success of these associations incontestable [1, 2]. Many of these

mutualistic associations rely on sulfur-oxidizing (thiotrophic), chemoautotrophic bacterial

symbionts, that oxidize reduced sulfur compounds for energy generation in order to fix

inorganic carbon (CO_2) for biomass build-up. Particularly in binary symbioses involving

intracorporeal thiotrophic bacteria, it is generally accepted that the endosymbiont's

chemosynthetic metabolism serves to provide organic carbon for feeding the animal host

[reviewed in 1–3]. In addition, some chemosynthetic symbionts have been found capable to

fix atmospheric nitrogen, albeit symbiont-to-host transfer of fixed nitrogen remains unproven

[4, 5]. As for the rarer chemosynthetic bacterial-animal associations in which symbionts

colonize exterior surfaces (ectosymbionts), fixation of inorganic carbon and transfer of

organic carbon to the host has only been demonstrated for the microbial community

colonizing the gill chamber of the hydrothermal vent shrimp *Rimicaris exoculata* [6].

In this study, we focused on *Candidatus* Thiosymbion oneisti, a

Gammaproteobacterium belonging to the basal family of *Chromatiaceae* (also known as

purple sulfur bacteria), which colonizes the surface of the marine nematode *Laxus oneistus*

(*Stilbonematinae*). Curiously, this group of free-living roundworms represents the only known

animals engaging in monospecific ectosymbioses, i.e. each nematode species is

ensheathed by a single *Ca.* Thiosymbion phylotype, and, in the case of *Ca.* T. oneisti, the

bacteria form a single layer on the cuticle of its host [7–11]. Moreover, the rod-shaped

representatives of this bacterial genus, including *Ca.* T. oneisti, divide by FtsZ-based

longitudinal fission, a unique reproductive strategy which ensures continuous and

transgenerational host-attachment [12–14].

Like other chemosynthetic symbionts, *Ca.* Thiosymbion have been considered

chemoautotrophic sulfur oxidizers based on several lines of evidence: stable carbon isotope

ratios of symbiotic nematodes are comparable to those found in other chemosynthetic

symbioses [15]; the key enzyme for carbon fixation via the Calvin-Benson-Bassham cycle

(RuBisCO) along with enzymes involved in sulfur oxidation and elemental sulfur have been

detected [16–18]; reduced sulfur compounds (sulfide, thiosulfate) have been shown to be

taken up from the environment by the ectosymbionts, to be used as energy source, and to

be responsible for the white appearance of the symbiotic nematodes [18–20]; the animals

often occur in the sulfidic zone of marine shallow-water sands [21]. More recently, the

phylogenetic placement and genetic repertoire of *Ca*. Thiosymbion species have equally

been supporting the chemosynthetic nature of the symbiosis [5, 11].

The majority of thioautotrophic symbioses have been described to rely on reduced

sulfur compounds as electron donors and oxygen as terminal electron acceptor [2, 3].

However, given that sulfidic and oxic zones are often spatially separated, also owing to

abiotic sulfide oxidation [22, 23], chemosynthetic symbioses (1) are found where sulfide and

oxygen occur in close proximity (e.g., hydrothermal vents, shallow-water sediments) and (2)

host behavioral, physiological and anatomical adaptations are needed to enable thiotrophic

symbionts to access both substrates. Host-mediated migration across the substrate

gradients is perhaps the most commonly encountered behavioral adaptation and was

proposed for deep-sea crustaceans, as well as for shallow water interstitial invertebrates and

Kentrophoros ciliates [reviewed in 1, 2]. Also the symbionts of *Stilbonematinae* have long

been hypothesized to associate with their motile nematode hosts to maximize sulfur

oxidation-fueled chemosynthesis, by alternately accessing oxygen in superficial sand and

sulfide in deeper, anoxic sand. This hypothesis was based upon the distribution pattern of

Stilbonematinae in sediment cores together with movement patterns in experimental

columns [15, 19, 21]. However, *Ca*. T. oneisti and other chemosynthetic symbionts were

subsequently shown to use nitrate as an alternative electron acceptor, and nitrate respiration

was stimulated by sulfide, suggesting that some of the symbionts are capable of gaining

energy by respiring nitrate instead of oxygen [20, 24–26].

Physiological studies are scarce because chemosynthetic associations remain

difficult to cultivate. Importantly, the impact of anoxic and oxic conditions on central

metabolic processes of the symbionts has not yet been investigated. This knowledge gap is

particularly remarkable in view of the exposed location of ectosymbionts, which leaves them

less sheltered than endosymbionts in the face of fluctuating concentrations of substrates for

energy generation. To understand how oxygen affects thiotrophy, we incubated symbiotic

nematodes associated with *Ca*. T. oneisti under different conditions resembling their natural

environment, and subsequently examined the ectosymbiont's transcriptional responses via

RNA sequencing (RNA-Seq). In combination with complementary methods such as stable isotope-labeling, proteomics, Raman microspectroscopy and lipidomics, we show that the ectosymbiont exhibits specific metabolic responses to anoxic and oxic conditions. Most strikingly, sulfur oxidation but not carbon fixation was upregulated in anoxia. This overturns the current opinion that sulfur oxidation requires oxygen and drives carbon fixation in chemosynthetic symbioses. We finally present a metabolic model of a thiotrophic ectosymbiont experiencing an ever-changing environment and propose that anaerobic sulfur oxidation coupled to denitrification represents the ectosymbiont's preferred metabolism for growth.

Materials and Methods

Sample collection

Laxus oneistus individuals were collected on multiple field trips (2016-2019) at approximately 1 m depth from sand bars off the Smithsonian Field Station, Carrie Bow Cay in Belize (16°48′11.01″N, 88°4′54.42″W). The nematodes were extracted from the sediment by gently stirring the sand and pouring the supernatant seawater through a 212 μm mesh sieve. The retained meiofauna was collected in a petri dish, and single worms of similar size (1-2 mm length, representing adult *L. oneistus*) were handpicked by forceps (Dumont 3, Fine Science Tools, Canada), under a dissecting microscope. Right after collection, the nematodes were used for various incubations as described below.

Sediment cores analysis

The habitat and spatial distribution of *L. oneistus* were characterized by sediment core analyses. Sediment pore-water was collected in July 2017, in 6 cm intervals down to a depth of 30 cm, by using cores of 60 cm length and 60 mm diameter (UWITEC, Mondsee, Austria), connected to rhizon samplers of a diameter of 2.5 mm and mean pore size of 0.15 μm (Rhizosphere Research Products, Wageningen, Netherlands). In total, nine sediment cores

were randomly sampled on different days and at different spots in shallow, submerged

sediment.

Immediately after collection, the sulfide content ($\sum H_2S$, i.e. the sum of H_2S, HS^- and

S^{2-}) was determined by the methylene-blue method [27]. In short, 670 µl of a 2% zinc

acetate solution was mixed with 335 µl sample and subsequently 335 µl 0.5% *N,N*-Dimethyl-

p-phenylenediamine and 17 µl of 10% ferrous ammonium sulfate added and incubated for

30 min in the dark. Surface seawater was used as a blank. Absorbance was measured at

670 nm and concentrations were quantified via calibration (measurement of $\sum H_2S$ standard

solutions in the concentration range from 0 to 0.5 mM $\sum H_2S$). Samples for dissolved

inorganic nitrogen (DIN: nitrate, nitrite, and ammonia) measurements were stored and

transported at -80°C, and analyzed at the University of Vienna, Austria. Nitrate (NO_3^-) and

nitrite (NO_2^-) concentrations were determined according to the Griess method [28] using VCl_3

[29], whereas the concentration of ammonium (NH_4^+) was measured after Solórzano [30].

For the quantification of nitrate, nitrite and, ammonia, freshly prepared KNO_3, $NaNO_2$ and

NH_4Cl solutions ranging from 0 to 20 µM were used to create standard curves, respectively.

Artificial seawater served as a blank [prepared after 31] and all measurements were

performed in triplicates. Rapidly after pore-water sampling, the abundance of *L. oneistus*

was determined within 6 cm-wide subsections of the cores. Each fraction was stirred

separately, and the supernatant decanted through a 212 µm-mesh sieve. The nematodes

were counted under a dissecting microscope. Mean nematode abundances and $\sum H_2S$

concentrations are shown in Figure S1A. All measurements data are listed in Table S1.

Incubations for RNA sequencing (RNA-Seq)

To analyze the physiological response of *Ca*. T. oneisti to oxygenated and reducing

conditions, three batches of 50 freshly collected *L. oneistus* individuals were incubated for

24 h in the dark, in either the presence (oxic) or absence (anoxic) of oxygen, in 13 ml-

exetainers (Labco, Lampeter, Wales, UK) fully filled with 0.2 µm filtered seawater,

respectively (Figure S1B). The oxic incubations consisted of two separate experiments of

low (hypoxic) and high (oxic saturated) oxygen concentrations. Here, all exetainers were kept open, but only the samples with high oxygen concentrations were submerged in an aquarium constantly bubbled with air (air pump plus; Sera, Heinsberg, Germany). Hypoxic samples were exposed to air, and they started with around 130 µM O_2 but reached less than 60 µM O_2 after 24 h of incubation. This likely occurred due to nematode oxygen consumption. On the other hand, the anoxic treatments comprised incubations to which either 11 µM of sodium sulfide (Na_2S*9H_2O; Sigma-Aldrich, St. Louis, MS, USA) was added, or no sulfide was supplied (Figure S1B), and $\sum H_2S$ concentrations were checked at the beginning and at the end (24 h) of each incubation by spectrophotometric determination following the protocol of Cline (see above). Anoxic incubations were achieved with the aid of a polyethylene glove bag (AtmosBag; Sigma-Aldrich) that was flushed with N_2 gas (Fabrigas, Belize City, Belize), together with incubation media and all vials, for at least 1 h before closing. Dissolved oxygen inside the bag, and of every incubation was monitored throughout the 24 h incubation time using a PreSens Fibox 3 trace fiber optic oxygen meter and non-invasive trace oxygen sensor spots attached to the exetainers (PSt6 and PSt3; PreSens, Regensburg, Germany). For exact measurements of $\sum H_2S$ and oxygen see Table S2. Temperature and salinity remained constant throughout all incubations measuring 27-28°C and 33-34 ‰, respectively. After the 24 h incubations, each set of 50 worms was quickly transferred into 2 ml RNA storage solution (13.3 mM EDTA disodium dihydrate pH 8.0, 16.6 mM sodium citrate dihydrate, 3.5 M ammonium sulfate, pH 5.2), kept at 4°C overnight, and finally stored in liquid nitrogen until RNA extraction.

RNA extraction, library preparation and RNA-Seq

RNA was extracted using the NucleoSpin RNA XS Kit (Macherey-Nagel, Düren, Germany). Briefly, sets of 50 worms in RNA storage solution were thawed and the worms were transferred into 90 µl lysis buffer RA1 containing Tris (2-carboxyethyl) phosphine (TCEP) according to the manufacturer. The remaining RNA storage solution was centrifuged to collect any detached bacterial cells (10 min, 4°C, 16 100 x g), pellets were resuspended in

10 µl lysis buffer RA1 (plus TCEP) and then added to the worms in lysis buffer. To further

disrupt cells, suspensions were vortexed for two minutes followed by three cycles of freeze

(-80°C) and thaw (37°C), and homogenization using a pellet pestle (Sigma-Aldrich) for 60 s

with a 15 s break after 30 s. Any remaining biological material on the pestle tips was

collected by rinsing the tip with 100 µl lysis buffer RA1 (plus TCEP). Lysates were applied to

NucleoSpin Filters and samples were processed as suggested by the manufacturer,

including an on-filter DNA digest. RNA was eluted in 20 µl RNase-free water. To remove any

residual DNA, a second DNase treatment was performed using the Turbo DNA-free Kit

(Thermo Fisher Scientific, Waltham, MA, USA), RNA was then dissolved in 17 µl RNase-free

water, and the RNA quality was assessed using a Bioanalyzer (Agilent, Santa Clara, CA,

USA). To check whether all DNA was digested, real-time quantitative PCR using the GoTaq

qPCR Master Mix (Promega, Madison, WI, USA) was performed targeting a 158 bp stretch

of the *sodB* gene using primers specific for the symbiont (sodB-F:

GTGAAGGGTAAGGACGGTTC, sodB-R: AATCCCAGTTGACGATCTCC, 10 µM per

primer). Different concentrations of genomic *Ca.* T. oneisti DNA were used as positive

controls. The program was as follows: 1x 95°C for 2 min, 40x 95°C for 15 s and 60°C for 1

min, 1x 95°C for 15 s, 55°C to 95°C over 20 min. Next, bacterial and eukaryotic rRNA was

removed using the Ribo-Zero Gold rRNA Removal Kit (Epidemiology) (Illumina, San Diego,

CA, USA) following the manufacturer's instructions, but volumes were adjusted for low input

RNA [32]. In short, 125 µl of Magnetic Beads Solution, 32.5 µl Magnetic Bead Resuspension

Solution, 2 µl Ribo-Zero Reaction Buffer and 4 µl Ribo-Zero Removal Solution were used

per sample. RNA was cleaned up via ethanol precipitation, dissolved in 9 µl RNase-free

water, and rRNA removal was evaluated using the Bioanalyzer RNA Pico Kit (Agilent, Santa

Clara, CA, USA). Strand-specific, indexed cDNA libraries were prepared using the SMARTer

Stranded RNA-Seq Kit (Takara Bio USA, Mountain View, CA, USA). Library preparation was

performed according to the instructions, with 8 µl of RNA per sample as input, 3 min

fragmentation time, two rounds of AMPure XP Beads (Beckman Coulter, Brea, CA, USA)

clean-up before amplification, and 18 PCR cycles for library amplification. The quality of the

libraries was assessed via the Bioanalyzer DNA High Sensitivity Kit (Agilent). Libraries were sequenced on an Illumina HiSeq 2500 instrument (single-read, 100 nt) at the next generation sequencing facility of the Vienna BioCenter Core Facilities (VBCF, https://www.viennabiocenter.org/facilities/).

Genome sequencing, assembly and functional annotation

The genome draft of *Ca.* T. oneisti was obtained by performing a hybrid assembly using reads from Oxford Nanopore Technologies (ONT) sequencing and Illumina sequencing. To extract DNA for ONT sequencing and dissociate the ectosymbionts from the host, approximately 800 *Laxus oneistus* individuals were incubated three times for 5 min each in TE‐Buffer (10 mM Tris-HCl pH 8.0, 1 mM disodium EDTA pH 8). Dissociated symbionts were collected by 10 min centrifugation at 7 000 x g and subsequent removal of the supernatant. DNA was extracted from this pellet using the Blood and Tissue Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The eluant was further purified using the DNA Clean & Concentrator-5 kit (Zymo Research, Irvine, CA, USA), and the DNA was eluted twice with 10 µl nuclease free water.

The library for ONT sequencing was prepared using the ONT Rapid Sequencing kit (SQK‐RAD002) and sequenced on a R9.4 flow cell (FLO-MIN106) on a MinION for 48 h. Base calling was performed locally with ONT's Metrichor Agent version 1.4.2, and resulting fastq-files were trimmed using Porechop version 0.2.1 (https://github.com/rrwick/Porechop). Illumina sequencing reads originate from a previous study [5], and were made available by Harald Gruber-Vodicka at the MPI Bremen. Raw reads were filtered: adapters removed and trimmed using bbduk (BBMap version 37.22, https://sourceforge.net/projects/bbmap/), with a minimum length of 36 and a minimum phred score of 2. To only keep reads derived from the symbiont, trimmed reads were mapped onto the available genome draft (NCBI Accession FLUZ00000000.1) using BWA-mem version 0.7.16a-r1181 [33]. Reads that did not map were discarded. The hybrid assembly was performed using SPAdes version 3.11 [34] with flags --careful and the ONT reads supplied as --nanopore. Contigs smaller than 200 bp and

a coverage lower than 5X were filtered out with a custom python script. The genome

completeness was assessed using checkM version 1.0.18 [35] with the

gammaproteobacterial marker gene set using the taxonomy workflow. The genome was

estimated to be 96.63% complete, contain 1.12% contamination, and was 4.35 Mb in length

on 401 contigs with a GC content of 58.7%.

The genome of *Ca.* T. oneisti was annotated using the MicroScope platform [36],

which predicted 5 169 protein-coding genes. To expand the functional annotation provided

by MicroScope, predicted proteins were assigned to KEGG pathway maps using

BlastKOALA and KEGG Mapper-Reconstruct Pathway [37], gene ontology (GO) terms using

Blast2GO version 5 [38] and searched for Pfam domains using the hmmscan algorithm of

HMMER 3.0 [39, 40]. All proteins and pathways mentioned in the manuscript were manually

curated. Functional annotations can be found in Data S1.

Gene expression analyses

Based on quality assessment of raw sequencing reads using FastQC [41] and prinseq [42],

reads were trimmed and filtered using Trimmomatic [43] as follows: 15-24 nucleotides were

removed from the 5-prime end (HEADCROP), Illumina adapters were cut

(ILLUMINACLIP:TruSeq3-SE.fa:2:30:10), and only reads longer than 24 nucleotides were

kept (MINLEN). Mapping and expression analysis were done as previously described [44].

Briefly, reads were mapped to the *Ca.* T. oneisti genome draft using BWA-backtrack [33]

with default settings, only uniquely mapped reads were kept using SAMtools [45], and the

number of strand-specific reads per gene were counted using HTSeq in the union mode of

counting overlaps [46]. On average, 1.3×10^6 (3.5%) reads uniquely mapped to the *Ca.* T.

oneisti genome. For detailed read and mapping statistics see Figure S2A. Gene expression

and differential expression analysis was conducted using the R software environment and

the Bioconductor package edgeR [47–49]. Genes were considered expressed if at least two

reads in at least two replicates of one of the four conditions could be assigned. Replicate 3

of the hypoxic incubations was excluded from further analyses because gene expression

was very different compared to the other two replicates (Figure S2B). Including all four conditions, we found 88.8% of total predicted symbiont protein-encoding genes to be expressed (4 591 genes out of 5 169). $\log_2$TPM (transcripts per kilobase million) values per gene and replicate were calculated and averages were determined per condition to estimate expression strength. Median gene expression was determined from average $\log_2$TPMs to highlight expression trends of entire metabolic processes and pathways. Expression of genes was considered significantly different if their expression changed two-fold between two conditions with a false-discovery rate (FDR) ≤ 0.05 [50]. Throughout the paper, all genes meeting these thresholds are either termed up- or downregulated or differentially expressed. However, most follow-up analyses were conducted only considering differentially expressed genes between the anoxic, sulfidic (AS) condition and both oxygenated conditions combined (O, see Results and Figure S2C). Heatmaps show mean centered expression values to highlight gene expression change.

Bulk δ^{13}C isotopic analysis by Isoprime isotope ratio mass spectrometry (EA-IRMS)

To further analyze the carbon metabolism of *Ca.* T. oneisti, specifically the assimilation of carbon dioxide (CO_2) by the symbionts in the presence or absence of oxygen, batches of 50 freshly collected, live worms were incubated for 24 h in 150 ml of 0.2 μm-filtered seawater, supplemented with 2 mM (final concentration) of either ^{12}C- (natural isotope abundance control) or ^{13}C-labelled sodium bicarbonate (Sigma-Aldrich, St. Louis, MS, USA). In addition, a second control was set up (dead control), in which prior to the 24 h incubation with ^{13}C-labelled sodium bicarbonate, 50 freshly collected worms were chemically killed by fixing them for 12 h in a 2% PFA/water solution.

All three incubations were performed in biological triplicates or quadruplets and set up under anoxic, sulfidic and oxic conditions. Like the RNA-Seq experiment, the oxic incubations consisted of two separate experiments of low (hypoxic) and high (oxic saturated) oxygen concentrations. To prevent isotope dilution through exchange with the atmosphere, both the oxic and anoxic incubations remained closed throughout the 24 h. The procedure

was as follows: 0.2 µm-filtered anoxic seawater was prepared as described above and was

subsequently used for both oxic and anoxic incubations. Then, compressed air (DAN oxygen

kit, Divers Alert Network, USA) and 25 µM of sodium sulfide (Na$_2$S*9H$_2$O; Sigma-Aldrich, St.

Louis, MS, USA) were injected into the oxic and anoxic incubations, respectively, to obtain

concentrations resembling the conditions applied in incubations for the RNA-Seq experiment

(see Table S2 for details about the incubation conditions and a compilation of the

measurement data).

At the end of each incubation (24 h), the nematodes were weighed (0.3-0.7 mg dry

weight) into tin capsules (Elemental Microanalysis, Devon, United Kingdom) and dried at

70°C for at least 24 h. Samples were analyzed using a Costech (Valencia, CA USA)

elemental analyzer interfaced with a continuous flow Micromass (Manchester, UK) Isoprime

isotope ratio mass spectrometer (EA-IRMS) for determination of ^{13}C/^{12}C isotope ratios.

Measurement values are displayed in δ notation [per mil (‰)]. A protein hydrolysate,

calibrated against NIST reference materials, was used as a standard in sample runs. The

achieved precision for δ^{13}C was ± 0.2 ‰ (one standard deviation of 10 replicate

measurements on the standard).

Raman microspectroscopy

Three individual nematodes per EA-IRMS incubation were fixed and stored in 0.1 M Trump's

fixative solution (0.1 M sodium cacodylate buffer, 2.5% GA, 2% PFA, pH 7.2, 1 000 mOsm L$^-$

^{1}) [51], and washed three times for 10 min in 1x PBS (137 mM NaCl, 2.7 mM KCl, 10 mM

Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$, pH 7.4) before their ectosymbionts were dissociated by

sonication for 40 s in 10 µl 1x PBS. 1 µl of each bacterial suspension was spotted on an

aluminum-coated glass slide and measured with a LabRAM HR Evolution Raman

microspectroscope (Horiba, Kyoto, Japan). 50 individual single-cell spectra were measured

from each sample. All spectra were aligned by the phenylalanine peak, baselined using the

Sensitive Nonlinear Iterative Peak (SNIP) algorithm of the R package "Peaks"

(https://www.rdocumentation.org/packages/Peaks/versions/0.2), and normalized by total

spectrum intensity. For calculating the relative sulfur content, the average intensity value for

212-229 wavenumbers (S_8 peak) was divided by the average intensity of the adjacent flat

region of 231-248 wavenumbers [18]. For calculating the relative polyhydroxyalkanoate

(PHA) content, the average intensity value for 1 723-1 758 wavenumbers (PHA peak) was

divided by the average intensity of the adjacent flat region of 1 759-1 793 wavenumbers [52].

Median relative sulfur and PHA content (shown as relative Raman intensities) were

calculated treating all individual symbiont cells per condition as replicates. Statistically

significant differences were determined applying the Wilcoxon rank sum test.

Assessment of the percentage of dividing cells

Samples were fixed and ectosymbionts were dissociated from their hosts as described

above for Raman microspectroscopy. 1.5 µl of each bacterial suspension per condition were

applied to a 1% agarose covered slide [53] and cells were imaged using a Nikon Eclipse NI-

U microscope equipped with an MFCool camera (Jenoptik). Images were obtained using the

ProgRes Capture Pro 2.8.8 software (Jenoptik) and processed with ImageJ [54]. Bacterial

cells were manually counted (> 600 per sample), and grouped into constricted (dividing) and

non-constricted (non-dividing) cells based on visual inspection [14]. The percentage of

dividing cells was calculated by counting the total number of dividing cells and the total

amount of cells per condition. Statistical tests were performed using the Fisher's exact test.

Data availability. The assembled and annotated genome of *Ca*. T. oneisti has been

deposited at DDBJ/ENA/GenBank under the accession JAAEFD000000000. RNA-Seq data

are available at the Gene Expression Omnibus (GEO) database and are accessible through

accession number GSE146081

Results

Hypoxic and oxic conditions induce similar expression profiles

To understand how the movement of the animal host across the chemocline affects symbiont physiology and metabolism, we exposed symbiotic worms to sulfide (thereafter used for $\sum H_2S$) and oxygen concentrations resembling the ones encountered by *Ca.* T. oneisti in its natural habitat. Previous studies showed that *Stilbonematinae* live predominantly in anoxic sediment zones with sulfide concentrations < 50 µM [21]. To assess whether this applies to *Laxus oneistus* (i.e. *Ca.* T. oneisti's host) at our collection site (Carrie Bow Cay, Belize), we determined the nematode abundance relative to the sampling depth and sulfide concentration. We found all *L. oneistus* individuals in pore water containing ≤ 25 µM sulfide and only 2% of them living in non-sulfidic (0 µM sulfide) surface layers (Figure S1A, Table S1). Therefore, we chose anoxic seawater supplemented with ≤ 25 µM sulfide as the optimal incubation medium (AS condition). To assess the effect of oxygen on symbiont physiology, we additionally incubated the nematodes in hypoxic (< 60 µM oxygen) and oxygen-saturated (>100 µM) seawater. Nitrate, nitrite and ammonium could be detected throughout the sediment core including the surface layer (Table S1).

Differential gene expression analysis comparing hypoxic and oxygen-saturated incubations revealed that expression levels of 97.1% of all expressed genes did not significantly differ between these two conditions and, crucially, included genes involved in central metabolic processes such as sulfur oxidation, fixation of carbon dioxide (CO_2), energy generation and stress response (Figure S2, Table S3). Because the presence of oxygen, irrespective of its concentration, resulted in a similar metabolic response, we treated the samples derived from hypoxic and oxygen-saturated incubations as biological replicates, and we will hereafter refer to them as the oxic (O) condition.

As such, we consider the AS condition as a proxy for the environment experienced by *Ca.* T. oneisti in deep pore water, and the O condition as the one experienced by the symbiont in superficial pore water. Differential gene expression analysis revealed that 21.7% of all genes exhibited significantly different expression between these two conditions (Figure S2C), and we will present their expression analysis below.

Sulfur oxidation genes are upregulated in anoxia

Ca. T. oneisti genes encoding for a sulfur oxidation pathway similar to that of the related but free-living purple sulfur bacterium *Allochromatium vinosum* were highly expressed in both conditions compared with other central metabolic processes, albeit median gene expression was higher in AS compared with oxic conditions (Figure 1). Consistently, 22 out of the 23 differentially expressed genes involved in sulfur oxidation, were upregulated in anoxia relative to oxic conditions (Figure 2A). These mostly included genes involved in the cytoplasmic branch of sulfur oxidation, i.e. genes associated with sulfur transfer from sulfur storage globules (*rhd, tusA, dsrE2*), genes encoding for the reverse-acting Dsr (dissimilatory sulfite reductase) system involved in the oxidation of stored elemental sulfur (S^0) to sulfite and, finally, also the genes required for further oxidation of sulfite to sulfate in the cytoplasm by two sets of adenylylsulfate (APS) reductase (*aprAB*) and one membrane-binding subunit (*aprM*) [55]. Genes encoding a quinone-interacting membrane-bound oxidoreductase (*qmoABC*) exhibited the same expression pattern. This is noteworthy, as AprM and QmoABC are hypothesized to have an analogous function, and their co-occurrence is rare among sulfur-oxidizing bacteria [55, 56].

Concerning genes involved in the periplasmic branch of sulfur oxidation such as the two types of sulfide-quinone reductases (*sqrA*, s*qrF*; oxidation of sulfide), the Sox system and the thiosulfate dehydrogenase (*tsdA*) both involved in oxidation of thiosulfate, transcript levels were unchanged between both conditions (Data S1). Only the flavoprotein subunit of the periplasmic flavocytochrome c sulfide dehydrogenase (*fccB*) was downregulated in the absence of oxygen (Figure 2A).

To assess whether upregulation of sulfur oxidation genes under anoxia was due to the absence of oxygen or the presence of sulfide, we performed an additional anoxic incubation without providing sulfide. Differential expression analysis between the anoxic conditions with and without sulfide revealed that transcript levels of 92.3% of all expressed genes did not significantly differ (Figure S2C). Importantly, sulfur oxidation genes were

similarly upregulated, irrespective of the presence of supplemented sulfide (Figure S3). In addition, proteome data derived from incubations with and without oxygen, but no added sulfide showed that one copy of AprA and AprM were among the top expressed proteins under anoxia (Table S4, Supplementary Materials & Methods). Raman microspectroscopy revealed that not only elemental stored sulfur (S^0) was still available at the end of all incubations but there was no significant difference between the S^0 content of the anoxic (no sulfide added) and O conditions (Figures 2B).

Collectively, our data indicate that despite the simultaneous availability of S^0 and oxygen (O condition), anoxia increased the transcription of genes involved in sulfur oxidation to sulfate.

Sulfur oxidation is coupled to denitrification

Given that sulfur oxidation was upregulated in anoxia, we expected this process to be coupled to the reduction of anaerobic electron acceptors, and nitrate respiration has been shown for symbiotic *L. oneistus* [20]. Consistently, genes encoding for components of the four specific enzyme complexes active in denitrification (*nap*, *nir*, *nor*, *nos*) as well as two subunits of the respiratory chain complex III (*petA* and *petB* of the cytochrome bc1 complex, which is known for being involved in denitrification and in the aerobic respiratory chain; [57]) were upregulated under anoxia (Figures 2A and S4).

In accordance with the upregulation of both sulfur oxidation and denitrification genes we observed via RNA-Seq, RT-qPCR targeting *aprA*, *dsrA* and *norB* transcripts showed that these genes were significantly higher expressed under anoxia in the closely related ectosymbiont of *Catanema* sp., which possesses the same central metabolic capabilities as *Ca.* T. oneisti ([11]; Supplementary Materials & Methods, Figures S5 and S6).

Besides nitrate respiration, *Ca.* T. oneisti may also be capable of utilizing dimethyl sulfoxide (DMSO) as a terminal electron acceptor since we observed an upregulation of all three subunits of a putative DMSO reductase (*dmsABC* genes). Concerning other electron acceptors, the symbiont has the genetic potential to carry out fumarate reduction (*frd* genes,

Data S1), and we identified a potential rhodoquinone biosynthesis gene (*rquA*), which acts

as anaerobic electron carrier involved in fumarate reduction in only a few prokaryotic and

eukaryotic organisms [58, 59]. Its upregulation under anoxic conditions indeed suggests an

active role in anaerobic energy generation in *Ca.* T. oneisti (Figure S4).

Intriguingly, lipid profiles of the symbiont revealed a change in lipid composition as

well as significantly higher relative abundances of several lyso-phospholipids under anoxia

(Figure S7, Supplementary Materials & Methods), possibly resulting in altered uptake

behavior and higher membrane permeability for electron donors and acceptors [60–63].

Notably, we also detected lyso-phosphatidylcholine to be significantly more abundant in

anoxia (Figure S7). As the symbiont does not possess known genes for biosynthesis of this

lipid, it may be host-derived. Incorporation of host lipids into symbiont membranes was

reported previously [64, 65].

Our data support that under anoxia, *Ca*. Thiosymbion gain energy by coupling sulfur

oxidation to complete reduction of nitrate to dinitrogen gas. Moreover, the symbiont appears

to exploit oxygen-depleted environments for energy generation by utilizing nitrate, DMSO

and fumarate as electron acceptors.

Sulfur oxidation is decoupled from carbon fixation

Several thioautotrophic symbionts have been shown to utilize the energy generated by sulfur

oxidation for fixation of inorganic carbon [6, 66–70]. Previous studies strongly support that

Ca. T. oneisti is capable of fixing carbon via an energy-efficient Calvin-Benson-Bassham

(CBB) cycle [5, 15, 17, 19, 71]. In this study, bulk isotope-ratio mass spectrometry (IRMS)

conducted with symbiotic nematodes confirmed that they incorporate [13]C-labelled inorganic

carbon. Remarkably, however, we detected incorporation of $^{13}CO_2$ under both anoxic and

oxic conditions to a similar extent in the course of 24 h (Figure 2C). To localize the sites of

$^{13}CO_2$ incorporation, we subjected $^{13}CO_2$-incubated symbiotic nematodes to nanoscale

secondary ion mass spectrometry (NanoSIMS) and detected [13]C enrichment predominantly

within the cells of the symbiont (Figure S8, Supplementary Materials & Methods).

Consistent with the evidence for carbon fixation by the ectosymbiont, all genes

related to the CBB cycle were detected, both on the transcriptome and the proteome level,

with high transcript levels under both anoxic and oxic conditions (Figure 1, Data S1).

However, the upregulation of sulfur oxidation genes observed under anoxia did not coincide

with an upregulation of carbon fixation genes. On the contrary, the median expression level

of all carbon fixation genes was higher in the presence of oxygen (Figure 1). In particular,

the transcripts encoding for the small subunit of the key CO_2-fixation enzyme ribulose-1,5-

bisphosphate carboxylase/oxygenase RuBisCO (*cbbS*) together with the transcripts

encoding its activases (*cbbQ* and *cbbO;* [72]) and the PPi-dependent 6-phosphofructokinase

(PPi-PFK; [73, 74]) were significantly more abundant under oxic conditions (Figure 2A).

Consistently, the large subunit of the RuBisCO protein (CbbL), which was among the top

expressed proteins in both conditions, was slightly more abundant in the presence of oxygen

(Table S4).

In accordance with our omics data, phylogenetic analysis of the *Ca.* T. oneisti

RuBisCO large subunit protein (CbbL) (Figure S9) shows that it clusters within the type I-A

group, whose characterized representatives are adapted to higher oxygen concentrations

[75].

In conclusion, key CO_2 fixation genes are upregulated in the presence of oxygen,

albeit inorganic carbon is incorporated to a similar extent under oxic and anoxic conditions.

In contrast to what has been reported about other thiotrophic bacteria, the upregulation of

sulfur oxidation genes was not accompanied by upregulation of genes involved in CO_2

fixation. Therefore, we hypothesize that (1) under anoxia, the energy derived from sulfur

oxidation is not completely utilized for carbon fixation and that (2) under oxic conditions

another electron donor besides sulfide contributes to energy generation for carbon fixation

(see section below).

Genes involved in the utilization of organic carbon and PHA storage build-up are

upregulated under oxic conditions

As anticipated, the nematode ectosymbiont may exploit additional reduced compounds

besides sulfide for energy generation. Indeed, *Ca.* T. oneisti possesses the genomic

potential to assimilate glyoxylate, acetate and propionate via the partial 3-hydroxypropionate

bi-cycle (like the closely related *Olavius algarvensis* γ1 symbiont; [73]), and furthermore

encodes genes for utilizing additional organic carbon compounds such as glycerol 3-

phosphate, glycolate, ethanol and lactate (Data S1). Figure 1 shows that, overall, the

expression of genes involved in the assimilation of organic carbon was higher under oxic

conditions. Among the upregulated genes were *lutB* (involved in oxidation of lactate to

pyruvate; [76]) and propionyl-CoA synthetase (*prpE,* propanoate assimilation; [77]) (Data

S1).

These gene expression data imply that the nematode ectosymbiont utilizes organic

carbon compounds in addition to CO_2 under oxic conditions, thereby increasing the supply of

carbon Consistent with high carbon availability, genes necessary to synthesize storage

compounds such as polyhydroxyalkanoates (PHA), glycogen and trehalose showed an

overall higher median transcript level under oxic conditions (Figure 1). In particular, two key

genes involved in the biosynthesis of the PHA compound polyhydroxybutyrate (PHB) –

acetyl-CoA acetyltransferase (*phaA*) and a class III PHA synthase subunit (*phaC*) – were

upregulated in the presence of oxygen. Conversely, we observed upregulation of both PHB

depolymerases under anoxia, and Raman microspectroscopy showed that significantly less

PHA was present in symbionts incubated in anoxia (Figure S10).

We propose that under oxic conditions, enhanced mixotrophy (i.e., CBB cycle-

mediated carbon fixation and organic carbon utilization) would result in higher carbon

availability reflected by PHA storage build-up and facilitating chemoorganoheterotrophic

synthesis of ATP via the aerobic respiratory chain.

Upregulation of nitrogen assimilation under oxic conditions

It has been shown that high carbon availability is accompanied by high nitrogen assimilation

[78–80]. Indeed, despite the sensitivity of nitrogenase towards oxygen [81], its key catalytic

MoFe enzymes (*nifD, nifK*; [82]) and several other genes involved in nitrogen fixation were

drastically upregulated in the presence of oxygen (Figures 1 and 3). RT-qPCR revealed an

increase of *nifH* transcripts also in the ectosymbiont of *Catanema* sp. when oxygen was

present (Figure S5, Supplementary Materials & Methods). Moreover, in accordance with a

recent study showing the importance of sulfur assimilation for nitrogen fixation [83], genes

involved in assimilation of sulfate, i.e. the sulfate transporters *sulP* and *cysZ* as well as

genes encoding two enzymes responsible for cysteine biosynthesis (*cysM*, *cysE*) were also

upregulated in the presence of oxygen (Data S1).

Besides nitrogen fixation, genes involved in urea uptake (transporters, *urtCBDE*) and

utilization (urease, *ureF* and *ureG*) were also significantly higher transcribed under oxic

conditions (Figures 1 and 3).

In conclusion, genes involved in assimilation of nitrogen (N_2 and urea) were

consistently upregulated in the presence of oxygen, when (1) carbon assimilation was likely

higher, and when (2) higher demand for nitrogen is expected due to stress-induced

synthesis of vitamins (see section below).

Upregulation of biosynthesis of cofactors and vitamins and global stress response

under oxic conditions

Multiple transcripts and proteins associated with a diverse bacterial stress response were

among the top expressed under oxic conditions (Figure 1 and Table S4). More specifically,

heat-shock proteins Hsp70 and Hsp90 were highly abundant (Table S4), and transcripts of

heat-shock proteins (Hsp15, Hsp40 and Hsp90) were upregulated (Figure 4A). Besides

chaperones, we also detected upregulation of superoxide dismutase (*sodB*), involved in the

scavenging of reactive oxygen species (ROS) [84], a transcription factor which induces

synthesis of Fe-S clusters under oxidative stress (*iscR*; [85, 86]) along with several other

genes involved in Fe-S cluster formation (Figure 4B, [87]), and regulators for redox

homeostasis, like thioredoxins, glutaredoxins and peroxiredoxins [88]. Furthermore, we

observed upregulation of protease genes (*rseP*, *htpX*, *hspQ*; [89–91]), genes required for

repair of double-strand DNA breaks (such as *radA*, *recB* and *mutSY*; [92, 93]) and even

genes that are involved in the response to amino acid, iron and fatty acid starvation (*relA*,

spoT, *sspA*; [94–97]) (Figure 4A).

We hypothesized that the drastic upregulation of stress-related genes observed

under oxic conditions would entail an increase in biosynthesis of vitamins [98–100]. Indeed,

genes involved in biosynthesis of vitamins such as vitamins B_2, B_6, and B_{12} were upregulated

under oxic conditions (Figures 1 and 4A). The proposed upregulation of nitrogen fixation and

urea utilization would support the synthesis of these nitrogen-rich molecules.

The upregulation of stress-related genes under oxic conditions was accompanied by

significantly fewer dividing symbiont cells (19.2% under oxic versus 30.1% under anoxic

conditions) (Figure 4B) and downregulation of early (*ftsEX*, *zipA*, *zapA*) and late (*damX*,

ftsN) cell division genes ([101], Figure 1 and Data S1). Oxygen therefore elicits a stress

response that affects symbiont proliferation.

Discussion

Although transcriptomic analysis was used to study the response of a vent snail

endosymbiont to changes in reduced sulfur species [69], up to this study, no thiotrophic

ectosymbiont has been subjected to global gene expression profiling under variable

environmental conditions. Therefore, our study provides unique insights into the global

physiological response of animal-attached bacteria that must thrive in the face of fluctuating

concentrations of substrates for energy generation.

We performed omics-based gene expression profiling on the nematode ectosymbiont

Ca. T. oneisti exposed to sulfide and oxygen at concentration levels encountered in its

habitat. In particular, the provided sulfide concentrations (< 25 µM) were comparable to

those reported for *Stilbonematinae* inhabiting sand areas in which sulfide concentrations

gradually increase with depth but are relatively low on average [21]. Given that in these so-

called "cool spots" oxygen and sulfide hardly co-occur [102, 103], we did not add any sulfide

to oxygenated seawater in our experimental set-up. However, given that stores of elemental

sulfur were still detected under all conditions after the incubation period and sulfur oxidation

genes were also upregulated in anoxic incubations without added sulfide, the electron donor

was available even when sulfide was not supplied, i.e. the ectosymbiont was not starved of

electron donor. In this study, we detected a stark transcriptional response of *Ca*.

Thiosymbion key metabolic processes to oxygen. The influence of oxygen on physiology

was globally reflected also by shifts in protein abundance and lipid composition.

Anaerobic sulfur oxidation

Genes involved in sulfur oxidation showed high overall expression compared to other central

metabolic processes, indicating that thiotrophy is the predominant energy-generating

process for *Ca.* T. oneisti in both oxic and anoxic conditions (Figure 5). Our data thus

strongly support previous observations of *Stilbonematinae* ectosymbionts performing aerobic

or anaerobic sulfur oxidation [19, 20]. As the majority of genes involved in denitrification

were upregulated in anoxia (Figures 2A and 5) and, importantly, we detected nitrate from the

most superficial down to the deepest pore water zones (Table S1), nitrate likely serves as

terminal electron acceptor for anaerobic sulfur oxidation.

Because sulfur oxidation in chemosynthetic symbioses is commonly described as an

aerobic process that also suits the host animal's physiology [2], anaerobic sulfur oxidation

seems extraordinary. However, many of these symbiotic organisms likely experience periods

of oxygen depletion as would be expected from life at the interface of oxidized and reduced

marine environments. Early studies on the physiology of a few thiotrophic symbionts indeed

describe nitrate reduction in the presence of sulfide under anoxic conditions [24–26].

Moreover, genes for nitrate respiration have been found in thiotrophic symbionts of a variety

of shallow-water and deep-sea vent animals [5, 73, 104–106]. Utilizing nitrate as an

alternative terminal electron acceptor during anaerobic sulfur oxidation to bridge temporary

anoxia could thus be a more important strategy for energy conservation among thiotrophic

symbionts than currently acknowledged.

Although relevant for organisms exposed to highly fluctuating oxygen concentrations,

this is the first study to show differential gene expression of a symbiont's sulfur oxidation

pathway between anoxic and oxic conditions. Specifically, the majority of sulfur oxidation

genes, and in particular genes encoding the cytoplasmic branch of sulfur oxidation, were

strongly upregulated under anoxic conditions (Figure 2). While upregulation of sulfur

oxidation and denitrification genes under anoxia represents no proof for preferential

anaerobic sulfur oxidation, we hypothesize that oxidation of reduced sulfur compounds to

sulfate is more pronounced under anoxia.

Among the upregulated sulfur oxidation genes, we identified *aprM* and the *qmoABC*

complex, both of which are thought to act as electron-accepting units for APS reductase,

and therefore rarely co-occur in thiotrophic bacteria [55]. The presence and expression of

the QmoABC complex could provide a substantial energetic advantage to the ectosymbiont

by mediating electron bifurcation [55], in which the additional reduction of a low-potential

electron acceptor (e.g. ferredoxin, NAD^+) could result in optimized energy conservation

under anoxic conditions. The maximization of sulfur oxidation under anoxia might even

represent a temporary advantage for the host, as it would protect the animal from sulfide

poisoning while crawling in predator-free and detritus-rich sand [20, 107–109]. Due to the

dispensability of oxygen for sulfur oxidation, the ectosymbiont may not need to be shuttled to

superficial sand by their nematode hosts to oxidize sulfur. Host migration into upper zones of

the sediment may therefore primarily reflect the animal's aerobic physiology.

In addition to anaerobic sulfur oxidation, the nematode ectosymbiont's phylogenetic

affiliation with anaerobic, phototrophic sulfur oxidizers such as *Allochromatium vinosum* [5,

110], and also the presence and expression of yet other anaerobic respiratory complexes

(DMSO and fumarate reductase) collectively suggest that *Ca*. Thiosymbion is well-adapted

to anoxic, sulfidic sediment zones.

Symbiont proliferation in anoxia

Pivotal studies addressed the extraordinary strategies *Ca*. T. oneisti evolved to grow and

divide [12, 14, 111]; yet, due to the difficulty in cultivation, the physiological basis of

longitudinal cell division has never been addressed.

In this study, significantly higher numbers of dividing cells were found under AS

conditions (Figure 4B, Data S1) and therefore, sulfur oxidation coupled to denitrification

might represent the ectosymbiont's preferred metabolism for growth. We hypothesize that

aside from sulfur oxidation, the mobilization of PHA could represent an additional source of

ATP (and carbon) supporting symbiont proliferation under anoxia (Figure S10, Figure 5). Of

note, PHA mobilization in anoxia was also shown for *Beggiatoa* spp. [112]. Furthermore,

several lines of research have shown that stress can inhibit growth in bacteria [113–119].

Importantly, studies on growth preferences of thiotrophic symbionts are lacking, and

proliferation of a thiotroph with anaerobic electron acceptors (such as nitrate) has never

been observed before [120–123].

683   Decoupling of sulfur oxidation from carbon fixation

A series of evidences gained by experiments conducted in the presence of oxygen indicate

that several thioautotrophic symbionts fuel carbon fixation by sulfur oxidation-generated

energy [6, 66–70]. Furthermore, (1) the giant tube worm *Riftia pachyptila* was recently

shown to exhibit higher inorganic carbon incorporation rates when incubated with sulfide and

high oxygen concentrations [124], (2) carbon fixation by the clam *Solemya velum* symbiont

was stimulated by sulfide and oxygen but not when nitrate was supplied as electron acceptor

[125] and (3) Schiemer *et al*. [19] showed that sulfide supported the uptake of CO_2 in

symbiotic nematodes incubated in the presence of oxygen. Although our bulk isotope-ratio

mass spectrometry (IRMS) and transcriptomic analyses indicate carbon fixation under both

anoxic and oxic conditions, the mere presence of sulfide did not appear to stimulate

incorporation of inorganic carbon. Instead, key carbon fixation genes of *Ca*. T. oneisti were

upregulated when oxygen was present (Figure 2). Therefore, carbon fixation appears to be

decoupled from anaerobic sulfur oxidation, and to rely on oxygen more than previously

acknowledged. An example of extreme decoupling of sulfur oxidation and carbon fixation was recently reported for *Kentrophoros* ectosymbionts, which were shown to lack genes for autotrophic carbon fixation altogether and thus represent the first heterotrophic sulfur-oxidizing symbionts [71].

Finally, regarding the fate of the fixed carbon, symbiont biomass build-up does not seem to solely rely on autotrophically-derived organic carbon, as the symbiont appeared to proliferate more in anoxia, when expression of carbon fixation genes was lower. Conversely, CO_2 fixed under oxic conditions may not be exclusively used for biomass generation (Figure 5).

Oxic mixotrophy

Besides chemoautotrophy, several chemosynthetic symbionts may engage in mixotrophy [5, 69, 73, 74, 126]. Also the nematode ectosymbiont possesses genes involved in the assimilation of organic carbon such as lactate, propionate, acetate and glycolate, and their expression was more pronounced under oxic conditions (Figure 1). Additional indications for mixotrophy comprise the presence and expression of genes for several transporters for small organic carbon compounds, a pyruvate dehydrogenase complex (*aceE, aceF, lpd*), and a complete TCA cycle, including a 2-oxoglutarate dehydrogenase (*korA*, *korB*), which is often lacking in obligate autotrophs [127] (Data S1). The ectosymbiont thus appears to assimilate both inorganic and organic carbon under oxic conditions and may consequently experience higher carbon availability (Figure 5).

The metabolization of these carbon compounds ultimately yields acetyl-CoA, which in turn could be further oxidized in the TCA cycle and/or utilized for fatty acid and PHA biosynthesis. Our transcriptome and Raman microspectroscopy data suggest that *Ca*. T. oneisti favors PHA build-up over degradation under oxic conditions. Thus, PHA accumulation in the presence of oxygen can be explained by higher carbon availability. In addition, it might play a role in resilience against cellular stress, as there is increasing evidence that PHA biosynthesis is enhanced under unfavorable growth conditions such as

extreme temperatures, UV radiation, osmotic shock and oxidative stress [128–136]. Similar

findings have been obtained for pathogenic [137] and symbiotic bacteria of the genus

Burkholderia [138]; the latter study reports upregulation of stress response genes and PHA

biosynthesis in the presence of oxygen. Finally, oxic biosynthesis of PHA might also prevent

excessive accumulation and breakdown of sugars by glycolysis and oxidative

phosphorylation, which, in turn, would exacerbate oxidative stress [139].

Oxic nitrogen assimilation

Despite the oxygen-sensitive nature of nitrogenase [81], we observed a drastic upregulation

of nitrogen fixation genes (Figures 1 and 3). In addition, urea utilization and uptake genes

were also upregulated. Although the nematode host likely lacks the urea biosynthetic

pathway (L.K., unpublished data), this compound is one of the most abundant organic

nitrogen substrates in marine ecosystems, as well as in animal-inhabited oxygenated sand

layers [140, 141]. Notably, the upregulation of the urea uptake system and urease accessory

proteins has been shown to be a response to nitrogen limitation in other systems [142].

An alternative or additional explanation for the apparent increase in nitrogen

assimilation in the presence of oxygen could be the need to synthesize nitrogen-rich

compounds such as vitamins and cofactors to survive oxidative stress (Figures 1 and 4).

The role of vitamins in protecting cells against the deleterious effects of oxygen has been

shown for animals [143, 144], and the importance of riboflavin for bacterial survival under

oxidative stress has previously been reported [98, 100]. Along this line of thought, oxygen-

exposed *Ca.* T. oneisti upregulated glutathione and thioredoxin, which are known to play a

pivotal role in scavenging reactive oxygen species (ROS) [145]. Their function directly (or

indirectly) requires vitamin B_2, B_6 and B_{12} as cofactors. More specifically, thioredoxin

reductase (*trxB*) requires riboflavin (vitamin B_2) in the form of flavin adenine dinucleotide

(FAD) [146]; cysteine synthase (*cysM*) and glutamate synthases (two-subunit *gltB/gltD,* one-

subunit *gltS*) involved in the biosynthesis of the glutathione precursors L-cysteine and L-

glutamate depend on vitamin B_6, FAD and riboflavin in the form of flavin mononucleotide

(FMN) [147, 148]. As for cobalamin, it was thought that this vitamin only played an indirect

role in oxidative stress resistance [149], by being a precursor of S-adenosyl methionine

(SAM), a substrate involved in the synthesis of glutathione via the methionine metabolism

(and the transulfuration pathway), and in preventing the Fenton reaction [150, 151].

However, its direct involvement in the protection of chemolithoautotrophic bacteria against

oxidative stress has also been illustrated [99].

In summary, in the presence of oxygen, the upregulation of genes involved in

biosynthesis of vitamins B_2, B_6 and B_{12} along with antioxidant systems and their key

precursor genes *cysM*, *gltD* and B_{12}-dependent-methionine synthase *metH*, suggests that

the ectosymbiont requires increased levels of these vitamins to cope with oxidative stress

(Figure 5).

Evolutionary considerations

Anaerobic sulfur oxidation, increased symbiont proliferation and downregulation of stress-

related genes lead us to hypothesize that *Ca.* T. oneisti evolved from a free-living bacterium

that mostly, if not exclusively, inhabited anoxic sand zones. In support of this, the closest

relatives of the nematode ectosymbionts are free-living obligate anaerobes from the family

Chromatiaceae (i.e. *Allochromatium vinosum*, *Thioflavicoccus mobilis*, *Marichromatium*

purpuratum) [5, 152]. Eventually, advantages such as protection from predators or utilization

of host waste products (e.g. fermentation products, ammonia) may have been driving forces

that led to the *Ca*. Thiosymbion-*Stilbonematinae* symbioses. As the association became

more and more stable, the symbiont optimized (or acquired) mechanisms to resist oxidative

stress, as well as metabolic pathways to most efficiently exploit the metabolic potential of

oxygenated sand zones (mixotrophy, nitrogen assimilation, vitamin and cofactor

biosynthesis). From the viewpoint of the host, the acquired "symbiotic skin" enabled *L.*

oneistus to tolerate the otherwise poisonous sulfide and thrive in sand layers virtually devoid

of predators and rich in decomposed organic matter (detritus).

Acknowledgements

This work was supported by the Austrian Science Fund (FWF) grant P28743 (T. V., S. B. and L. K.) and the DK plus: Microbial Nitrogen Cycling (G. F. P.). We are indebted to the staff of the VBCF NGS Unit (Laura-Maria Bayer and Miriam Schalamun) for assistance with Oxford Nanopore MinION sequencing and to Florian Goldenberg, Patrick Hyden and Thomas Rattei (Division of Computational Systems Biology, University of Vienna) for providing and maintaining the Life Science Compute Cluster (LiSC) and help in preparing the MinION sequencing library for *Ca.* T. oneisti at the University of Vienna. Harald Gruber-Vodicka from the MPI Bremen generously provided Illumina raw reads to aid the assemblies of ectosymbiont genomes. We are very grateful to Wiebke Mohr, Nikolaus Leisch and Nicole Dubilier from the MPI for Marine Microbiology (Bremen) for continuous technical support with the stable isotope-based techniques and anaerobic atmosbag and Andreas Maier for DOC measurements (data not shown). We thank Olivier Gros from the Université des Antilles for making all the experiments involving *Catanema sp.* possible. We appreciate Tjorven Hinzke's advice on metaproteome statistics and Carolina Reyes for her input on the nitrogen metabolism. We thank Yin Chen for providing the facilities for lipidomic analysis and Eleonora Silvano for assistance with lipid extractions, and Jana Matulla's and Sebastian Grund's excellent technical work during protein sample preparation and MS analysis, respectively. Also, our sincere gratitude to the Carrie Bow Cay Laboratory, Caribbean Coral Reef Ecosystem Program and his Station Manager Zach Foltz for his help during field work. Finally, we were very much helped and inspired by insightful discussions with Monika Bright, Jörg A. Ott, Christa Schleper, Simon Rittmann, Filipa Sousa and Jillian Petersen.

Competing Interests

The authors declare no competing financial interest.

References

1.   Ott J, Bright M, Bulgheresi S. Marine microbial thiotrophic ectosymbioses. In: Gibson

RN, Atkinson RJA, Gordon JDM (eds). *Oceanography and Marine Biology: An Annual*

Review, 42nd ed. 2004. pp 95–118.

2.   Dubilier N, Bergin C, Lott C. Symbiotic diversity in marine animals: the art of

harnessing chemosynthesis. *Nat Rev Microbiol* 2008; **6**: 725–740.

3.   Stewart FJ, Newton ILG, Cavanaugh CM. Chemosynthetic endosymbioses:

adaptations to oxic–anoxic interfaces. *Trends Microbiol* 2005; **13**: 439–448.

4.   König S, Gros O, Heiden SE, Hinzke T, Thürmer A, Poehlein A, et al. Nitrogen fixation

in a chemoautotrophic lucinid symbiosis. *Nat Microbiol* 2016; **2**: 16193.

5.   Petersen JM, Kemper A, Gruber-Vodicka H, Cardini U, van der Geest M, Kleiner M, et

al. Chemosynthetic symbionts of marine invertebrate animals are capable of nitrogen

fixation. *Nat Microbiol* 2016; **2**: 16195.

6.   Ponsard J, Cambon-Bonavita M-A, Zbinden M, Lepoint G, Joassin A, Corbari L, et al.

Inorganic carbon fixation by chemosynthetic ectosymbionts and nutritional transfers to

the hydrothermal vent host-shrimp *Rimicaris exoculata*. *ISME J* 2013; **7**: 96–109.

7.   Polz MF, Distel DL, Zarda B, Amann R, Felbeck H, Ott JA, et al. Phylogenetic

analysis of a highly specific association between ectosymbiotic, sulfur-oxidizing

bacteria and a marine nematode. *Appl Environ Microbiol* 1994; **60**: 4461–4467.

8.   Bayer C, Heindl NR, Rinke C, Lücker S, Ott J, Bulgheresi S. Molecular

characterization of the symbionts associated with marine nematodes of the genus

Robbea. *Environ Microbiol Rep* 2009; **1**: 136–144.

9.   Williams KP, Gillespie JJ, Sobral BWS, Nordberg EK, Snyder EE, Shallom JM, et al.

Phylogeny of gammaproteobacteria. *J Bacteriol* 2010; **192**: 2305–2314.

10.   Pende N, Leisch N, Gruber-Vodicka HR, Heindl NR, Ott J, den Blaauwen T, et al.

Size-independent symmetric division in extraordinarily long cells. *Nat Commun* 2014;

5: 4803.

11.   Zimmermann J, Wentrup C, Sadowski M, Blazejak A, Gruber-Vodicka HR, Kleiner M,

et al. Closely coupled evolutionary history of ecto- and endosymbionts from two

distantly related animal phyla. *Mol Ecol* 2016; **25**: 3203–3223.

12.   Leisch N, Verheul J, Heindl NR, Gruber-Vodicka HR, Pende N, den Blaauwen T, et al.

Growth in width and FtsZ ring longitudinal positioning in a gammaproteobacterial

symbiont. *Curr Biol* 2012; **22**: R831–R832.

13.   Leisch N, Pende N, Weber PM, Gruber-Vodicka HR, Verheul J, Vischer NOE, et al.

Asynchronous division by non-ring FtsZ in the gammaproteobacterial symbiont of

Robbea hypermnestra. Nat Microbiol 2016; **2**: 1–5.

14.   Pende N, Wang J, Weber PM, Verheul J, Kuru E, Rittmann SK-MR, et al. Host-

polarized cell growth in animal symbionts. *Curr Biol* 2018; **28**: 1039-1051.e5.

15.   Ott JA, Novak R, Schiemer F, Hentschel U, Nebelsick M, Polz M. Tackling the sulfide

gradient: a novel strategy involving marine nematodes and chemoautotrophic

ectosymbionts. *Mar Ecol* 1991; **12**: 261–279.

16.   Powell EN, Crenshaw MA, Rieger RM. Adaptations to sulfide in the meiofauna of the

sulfide system. I. 35S-sulfide accumulation and the presence of a sulfide

detoxification system. *J Exp Mar Bio Ecol* 1979; **37**: 57–76.

17.   Polz MF, Felbeck H, Novak R, Nebelsick M, Ott JA. Chemoautotrophic, sulfur-

oxidizing symbiotic bacteria on marine nematodes: morphological and biochemical

characterization. *Microb Ecol* 1992; **24**: 313–329.

18.   Himmel D, Maurin LC, Gros O, Mansot J-L. Raman microspectrometry sulfur

detection and characterization in the marine ectosymbiotic nematode *Eubostrichus*

dianae (Desmodoridae, Stilbonematidae). *Biol Cell* 2009; **101**: 43–54.

19.   Schiemer F, Novak R, Ott J. Metabolic studies on thiobiotic free-living nematodes and

their symbiotic microorganisms. *Mar Biol* 1990; **106**: 129–137.

20.   Hentschel U, Berger E, Bright M, Felbeck H, Ott J. Metabolism of nitrogen and sulfur

in ectosymbiotic bacteria of marine nematodes (Nematoda, Stilbonematinae). *Mar*

Ecol Prog Ser 1999; **183**: 149–158.

21.   Ott J, Novak R. Living at an interface: Meiofauna at the oxygen/sulfide boundary in

marine sediments. In: Ryland JS, Tyler PA (eds). *Reproduction, Genetics and*

Distributions of Marine Organisms. 1989. Olsen & Olsen, Fredensborg, pp 415–422.

22. Canfield DE, Thamdrup B. Towards a consistent classification scheme for

geochemical environments, or, why we wish the term 'suboxic' would go away.

Geobiology 2009; **7**: 385–392.

23. Luther GW, Findlay AJ, MacDonald DJ, Owings SM, Hanson TE, Beinart RA, et al.

Thermodynamics and kinetics of sulfide oxidation by oxygen: a look at inorganically

controlled reactions and biologically mediated processes in the environment. *Front*

Microbiol 2011; **2**: 62.

24. Wilmot DB, Vetter RD. Oxygen- and nitrogen-dependent sulfur metabolism in the

thiotrophic clam *Solemya reidi. Biol Bull* 1992; **182**: 444–453.

25. Hentschel U, Cary SC, Felbeck H. Nitrate respiration in chemoautotrophic symbionts

of the bivalve *Lucinoma aequizonata. Mar Ecol Prog Ser* . 1993. Inter-Research

Science Center. , **94**: 35–41

26. Hentschel U, Felbeck H. Nitrate respiration in the hydrothermal vent tubeworm *Riftia*

pachyptila. Nature 1993; **366**: 338–340.

27. Cline JD. Spectrophotometric determination of hydrogen sulfide in natural waters.

Limnol Oceanogr 1969; **14**: 454–458.

28. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR.

Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal Biochem* 1982;

126: 131–138.

29. Schnetger B, Lehners C. Determination of nitrate plus nitrite in small volume marine

water samples using vanadium(III)chloride as a reduction agent. *Mar Chem* 2014;

160: 91–98.

30. Solórzano L. Determination of ammonia in natural waters by the phenolhypochlorite

method. *Limnol Oceanogr* 1969; **14**: 799–801.

31. Kester DR, Duedall IW, Connors DN, Pytkowicz RM. Preparation of artificial seawater.

Limnol Oceanogr 1967; **12**: 176–179.

32.   Alberti A, Belser C, Engelen S, Bertrand L, Orvain C, Brinas L, et al. Comparison of

893   library preparation methods reveals their impact on interpretation of

894   metatranscriptomic data. *BMC Genomics* 2014; **15**: 912.

895   33.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

896   transform. *Bioinformatics* 2009; **25**: 1754–1760.

897   34.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:

898   a new genome assembly algorithm and its applications to single-cell sequencing. *J*

899   *Comput Biol* 2012; **19**: 455–477.

900   35.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing

901   the quality of microbial genomes recovered from isolates, single cells, and

902   metagenomes. *Genome Res* 2015; **25**: 1043–1055.

903   36.   Vallenet D, Calteau A, Dubois M, Amours P, Bazin A, Beuvin M, et al. MicroScope: an

904   integrated platform for the annotation and exploration of microbial gene functions

905   through genomic, pangenomic and metabolic comparative analysis. *Nucleic Acids*

906   *Res* 2019; **48**: D579–D589.

907   37.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference

908   resource for gene and protein annotation. *Nucleic Acids Res* 2016; **44**: D457–D462.

909   38.   Götz S, Garcia-Gomez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, et al. High-

910   throughput functional annotation and data mining with the Blast2GO suite. *Nucleic*

911   *Acids Res* 2008; **36**: 3420–3435.

912   39.   Bateman A. The Pfam protein families database. *Nucleic Acids Res* 2002; **30**: 276–

913   280.

914   40.   Finn RD, Clements J, Eddy SR. HMMER web server: interactive sequence similarity

915   searching. *Nucleic Acids Res* 2011; **39**: W29–W37.

916   41.   Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010.

917   42.   Schmieder R, Edwards R. Quality control and preprocessing of metagenomic

918   datasets. *Bioinformatics* 2011; **27**: 863–864.

43.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* 2014; **30**: 2114–2120.

44.   König L, Siegl A, Penz T, Haider S, Wentrup C, Polzin J, et al. Biphasic Metabolism

and Host Interaction of a Chlamydial Symbiont. *mSystems* 2017; **2**: e00202-16.

45.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–2079.

46.   Anders S, Pyl PT, Huber W. HTSeq - A Python framework to work with high-

throughput sequencing data. *Bioinformatics* 2014; **31**: 166–169.

47.   Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, et al.

Bioconductor: open software development for computational biology and

bioinformatics. *Genome Biol* 2004; **5**: R80.

48.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* 2010;

26: 139–140.

49.   R Development Core Team. R: a language and environment for statistical computing.

R Found Stat Comput . 2011. R Foundation for Statistical Computing. , **1**: 409

50.   Rapaport F, Khanin R, Liang Y, Pirun M, Krek A, Zumbo P, et al. Comprehensive

evaluation of differential gene expression analysis methods for RNA-seq data.

Genome Biol 2013; **14**: R95.

51.   Trump BF, Ericsson JL. The effect of the fixative solution on the ultrastructure of cells

and tissues. A comparative analysis with particular attention to the proximal

convoluted tubule of the rat kidney. *Lab Investig* 1965; **14**: 1245–1323.

52.   Samek O, Obruča S, Šiler M, Sedláček P, Benešová P, Kučera D, et al. Quantitative

Raman spectroscopy analysis of polyhydroxyalkanoates produced by *Cupriavidus*

necator H16. *Sensors* 2016; **16**: 1808.

53.   Koppelman C-M, Aarsman MEG, Postmus J, Pas E, Muijsers AO, Scheffers D-J, et

al. R174 of *Escherichia coli* FtsZ is involved in membrane interaction and

protofilament bundling, and is essential for cell division. *Mol Microbiol* 2004; **51**: 645–

657.

54. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* 2012; **9**: 671–675.

55. Dahl C. Sulfur Metabolism in Phototrophic Bacteria. *Modern Topics in the*

Phototrophic Prokaryotes. 2017. Springer International Publishing, Cham, pp 27–66.

56. Ghosh W, Dam B. Biochemistry and molecular biology of lithotrophic sulfur oxidation

by taxonomically and ecologically diverse bacteria and archaea. *FEMS Microbiol Rev*

2009; **33**: 999–1043.

57. Chen J, Strous M. Denitrification and aerobic respiration, hybrid electron transport

chains and co-evolution. *Biochim Biophys Acta - Bioenerg* 2013; **1827**: 136–144.

58. Lonjers ZT, Dickson EL, Chu T-PT, Kreutz JE, Neacsu FA, Anders KR, et al.

Identification of a new gene required for the biosynthesis of rhodoquinone in

Rhodospirillum rubrum. J Bacteriol 2012; **194**: 965–971.

59. Stairs CW, Eme L, Muñoz-Gómez SA, Cohen A, Dellaire G, Shepherd JN, et al.

Microbial eukaryotes have adapted to hypoxia by horizontal acquisitions of a gene

involved in rhodoquinone biosynthesis. *Elife* 2018; **7**: e34292.

60. Mills JK, Needham D. Lysolipid incorporation in dipalmitoylphosphatidylcholine bilayer

membranes enhances the ion permeability and drug release rates at the membrane

phase transition. *Biochim Biophys Acta - Biomembr* 2005; **1716**: 77–96.

61. Arouri A, Mouritsen OG. Membrane-perturbing effect of fatty acids and lysolipids.

Prog Lipid Res . 2013. Pergamon. , **52**: 130–140

62. Koshy C, Schweikhard ES, Gärtner RM, Perez C, Yildiz Ö, Ziegler C. Structural

evidence for functional lipid interactions in the betaine transporter BetP. *EMBO J*

2013; **32**: 3096–3105.

63. Laganowsky A, Reading E, Allison TM, Ulmschneider MB, Degiacomi MT, Baldwin

AJ, et al. Membrane proteins bind lipids selectively to modulate their structure and

function. *Nature* 2014; **510**: 172–175.

64. Wier AM, Nyholm S V, Mandel MJ, Massengo-Tiassé RP, Schaefer AL, Koroleva I, et

al. Transcriptional patterns in both host and bacterium underlie a daily rhythm of

anatomical and metabolic change in a beneficial symbiosis. *Proc Natl Acad Sci U S A*

2010; **107**: 2259–64.

65. Saito HE, Harp JR, Fozo EM. Incorporation of exogenous fatty acids protects

Enterococcus faecalis from membrane-damaging agents. *Appl Environ Microbiol*

2014; **80**: 6527–6538.

66. Cavanaugh CM. Symbiotic chemoautotrophic bacteria in marine invertebrates from

sulphide-rich habitats. *Nature* 1983; **302**: 58–61.

67. Nelson DC, Hagen KD, Edwards DB. The gill symbiont of the hydrothermal vent

mussel *Bathymodiolus thermophilus* is a psychrophilic, chemoautotrophic, sulfur

bacterium. *Mar Biol* 1995; **121**: 487–495.

68. Markert S, Arndt C, Felbeck H, Becher D, Sievert SM, Hügler M, et al. Physiological

proteomics of the uncultured endosymbiont of Riftia pachyptila. *Science* 2007; **315**:

247–250.

69. Seston SL, Beinart RA, Sarode N, Shockey AC, Ranjan P, Ganesh S, et al.

Metatranscriptional response of chemoautotrophic *Ifremeria nautilei* endosymbionts to

differing sulfur regimes. *Front Microbiol* 2016; **7**: 1074.

70. Volland J-M, Schintlmeister A, Zambalos H, Reipert S, Mozetič P, Espada-Hinojosa S,

et al. NanoSIMS and tissue autoradiography reveal symbiont carbon fixation and

organic carbon transfer to giant ciliate host. *ISME J* 2018; **12**: 714–727.

71. Seah BKB, Antony CP, Huettel B, Zarzycki J, Schada von Borzyskowski L, Erb TJ, et

al. Sulfur-oxidizing symbionts without canonical genes for autotrophic CO2 fixation.

MBio 2019; **10**: e01112-19.

72. Tsai YCC, Lapina MC, Bhushan S, Mueller-Cajar O. Identification and

characterization of multiple rubisco activases in chemoautotrophic bacteria. *Nat*

Commun 2015; **6**: 8883.

73. Kleiner M, Wentrup C, Lott C, Teeling H, Wetzel S, Young J, et al. Metaproteomics of

a gutless marine worm and its symbiotic microbial community reveal unusual

pathways for carbon and energy use. *Proc Natl Acad Sci U S A* 2012; **109**: E1173–

E1182.

74.   Jäckle O, Seah BKB, Tietjen M, Leisch N, Liebeke M, Kleiner M, et al.

1006   Chemosynthetic symbiont with a drastically reduced genome serves as primary

1007   energy storage in the marine flatworm *Paracatenula*. *Proc Natl Acad Sci U S A* 2019;

1008   **116**: 8505–8514.

1009   75.   Tabita FR, Satagopan S, Hanson TE, Kreel NE, Scott SS. Distinct form I, II, III, and IV

1010   Rubisco proteins from the three kingdoms of life provide clues about Rubisco

1011   evolution and structure/function relationships. *J Exp Bot* 2007; **59**: 1515–1524.

1012   76.   Chai Y, Kolter R, Losick R. A widely conserved gene cluster required for lactate

1013   utilization in *Bacillus subtilis* and its involvement in biofilm formation. *J Bacteriol* 2009;

1014   **191**: 2423–2430.

1015   77.   Horswill AR, Escalante-Semerena JC. The prpE gene of *Salmonella typhimurium* LT2

1016   encodes propionyl-CoA synthetase. *Microbiology* 1999; **145**: 1381–1388.

1017   78.   Rogers A, Gibon Y, Stitt M, Morgan PB, Bernacchi CJ, Ort DR, et al. Increased C

1018   availability at elevated carbon dioxide concentration improves N assimilation in a

1019   legume. *Plant, Cell Environ* 2006; **29**: 1651–1658.

1020   79.   Elgharably A, Marschner P. Microbial activity and biomass and N and P availability in

1021   a saline sandy loam amended with inorganic N and lupin residues. *Eur J Soil Biol*

1022   2011; **47**: 310–315.

1023   80.   Fellbaum CR, Gachomo EW, Beesetty Y, Choudhari S, Strahan GD, Pfeffer PE, et al.

1024   Carbon availability triggers fungal nitrogen uptake and transport in arbuscular

1025   mycorrhizal symbiosis. *Proc Natl Acad Sci* 2012; **109**: 2666–2671.

1026   81.   Hill S. How is nitrogenase regulated by oxygen? *FEMS Microbiol Lett* 1988; **54**: 111–

1027   129.

1028   82.   Burgess BK, Lowe DJ. Mechanism of molybdenum nitrogenase. *Chem Rev* 1996; **96**:

1029   2983–3012.

1030   83.   Schneider S, Schintlmeister A, Becana M, Wagner M, Woebken D, Wienkoop S.

1031   Sulfate is transported at significant rates through the symbiosome membrane and is

1032   crucial for nitrogenase biosynthesis. *Plant Cell Environ* 2019; **42**: 1180–1189.

1033 84. Hassan HM. Determination of microbial damage caused by oxygen free radicals, and

1034   the protective role of superoxide dismutase. *Methods Enzymol* 1984; **105**: 404–412.

1035 85. Schwartz CJ, Giel JL, Patschkowski T, Luther C, Ruzicka FJ, Beinert H, et al. IscR, an

1036   Fe-S cluster-containing transcription factor, represses expression of *Escherichia coli*

1037   genes encoding Fe-S cluster assembly proteins. *Proc Natl Acad Sci U S A* 2001; **98**:

1038   14895–14900.

1039 86. Yeo W-S, Lee J-H, Lee K-C, Roe J-H. IscR acts as an activator in response to

1040   oxidative stress for the *suf* operon encoding Fe-S assembly proteins. *Mol Microbiol*

1041   2006; **61**: 206–218.

1042 87. Zeller T, Moskvin O V., Li K, Klug G, Gomelsky M. Transcriptome and physiological

1043   responses to hydrogen peroxide of the facultatively phototrophic bacterium

1044   Rhodobacter sphaeroides. *J Bacteriol* 2005; **187**: 7232–7242.

1045 88. Grant CM. Role of the glutathione/glutaredoxin and thioredoxin systems in yeast

1046   growth and response to stress conditions. *Mol Microbiol* 2001; **39**: 533–541.

1047 89. Alba BM, Leeds JA, Onufryk C, Lu CZ, Gross CA. DegS and YaeL participate

1048   sequentially in the cleavage of RseA to activate the sigma E-dependent

1049   extracytoplasmic stress response. *Genes Dev* 2002; **16**: 2156–2168.

1050 90. Sakoh M, Ito K, Akiyama Y. Proteolytic activity of HtpX, a membrane-bound and

1051   stress-controlled protease from *Escherichia coli*. *J Biol Chem* 2005; **280**: 33305–

1052   33310.

1053 91. Puri N, Karzai AW. HspQ functions as a unique specificity-enhancing factor for the

1054   AAA+ Lon protease. *Mol Cell* 2017; **66**: 672-683.e4.

1055 92. Sargentini NJ, Smith KC. Quantitation of the involvement of the *recA, recB, recC,*

1056   *recF, recJ, recN, lexA, radA, radB, uvrD*, and *umuC* genes in the repair of X-ray-

1057   induced DNA double-strand breaks in <i>Es. *Radiat Res* 1986; **107**: 58–72.

1058 93. Au KG, Cabrera M, Miller JH, Modrich P. *Escherichia coli mutY* gene product is

1059   required for specific A-G----C.G mismatch correction. *Proc Natl Acad Sci* 1988; **85**:

1060   9163–9166.

94.   Hansen A-M, Qiu Y, Yeh N, Blattner FR, Durfee T, Jin DJ. SspA is required for acid
       resistance in stationary phase by downregulation of H-NS in *Escherichia coli*. *Mol
       Microbiol* 2005; **56**: 719–734.

95.   Battesti A, Bouveret E. Acyl carrier protein/SpoT interaction, the switch linking SpoT-
       dependent stress response to fatty acid metabolism. *Mol Microbiol* 2006; **62**: 1048–
       1063.

96.   Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Hightower GA, Smith JT, et al.
       The global, ppGpp-mediated stringent response to amino acid starvation in
       Escherichia coli. *Mol Microbiol* 2008; **68**: 1128–1148.

97.   Nguyen D, Joshi-Datar A, Lepine F, Bauerle E, Olakanmi O, Beer K, et al. Active
       starvation responses mediate antibiotic tolerance in biofilms and nutrient-limited
       bacteria. *Science (80-)* 2011; **334**: 982–986.

98.   Chen J, Shen J, Solem C, Jensen PR. Oxidative stress at high temperatures in
       Lactococcus lactis due to an insufficient supply of riboflavin. *Appl Environ Microbiol*
       2013; **79**: 6140–6147.

99.   Ferrer A, Rivera J, Zapata C, Norambuena J, Sandoval Á, Chávez R, et al. Cobalamin
       protection against oxidative stress in the acidophilic iron-oxidizing bacterium
       Leptospirillum group II CF-1. *Front Microbiol* 2016; **7**: 748.

100.   Showman AC, Aranjuez G, Adams PP, Jewett MW. Gene bb0318 is critical for the
       oxidative stress response and infectivity of *Borrelia burgdorferi*. *Infect Immun* 2016;
       84: 3141–3151.

101.   Typas A, Banzhaf M, Gross CA, Vollmer W. From the regulation of peptidoglycan
       synthesis to bacterial growth and morphology. *Nat Rev Microbiol* 2012; **10**: 123–136.

102.   Meyers M, Fossing H, Powell E. Microdistribution of interstitial meiofauna, oxygen and
       sulfide gradients, and the tubes of macro-infauna. *Mar Ecol Prog Ser* 1987; **35**: 223–
       241.

103.   Powell E. Oxygen, sulfide and diffusion: Why thiobiotic meiofauna must be sulfide-

insensitive first-order respirers. *J Mar Res* 1989; **47**: 887–932.

104. Jan C, Petersen JM, Werner J, Teeling H, Huang S, Glöckner FO, et al. The gill

chamber epibiosis of deep-sea shrimp *Rimicaris exoculata*: an in-depth metagenomic

investigation and discovery of *Zetaproteobacteria*. *Environ Microbiol* 2014; **16**: 2723–

2738.

105. Ponnudurai R, Sayavedra L, Kleiner M, Heiden SE, Thürmer A, Felbeck H, et al.

Genome sequence of the sulfur-oxidizing Bathymodiolus thermophilus gill

endosymbiont. *Stand Genomic Sci* 2017; **12**: 50.

106. Ponnudurai R, Kleiner M, Sayavedra L, Petersen JM, Moche M, Otto A, et al.

Metabolic and physiological interdependencies in the *Bathymodiolus azoricus*

symbiosis. *ISME J* 2017; **11**: 463–477.

107. Fenchel T. Studies on the decomposition of organic detritus derived from the turtle

grass *Thalassia testudinum*. *Limnol Oceanogr* 1970; **15**: 14–20.

108. Fenchel TM, Riedl RJ. The sulfide system: a new biotic community underneath the

oxidized layer of marine sand bottoms. *Mar Biol* 1970; **7**: 255–268.

109. Reise K, Ax P. A meiofaunal 'thiobios' limited to the anaerobic sulfide system of

marine sand does not exist. *Mar Biol* 1979; **54**: 225–237.

110. Dahl C, Friedrich C, Kletzin A. Sulfur Oxidation in Prokaryotes. *Encyclopedia of Life*

Sciences. 2008. John Wiley & Sons, Ltd, Chichester, UK.

111. Weber PM, Moessel F, Paredes GF, Viehboeck T, Vischer NOE, Bulgheresi S. A

bidimensional segregation mode maintains symbiont chromosome orientation toward

its host. *Curr Biol* 2019; **29**: 3018-3028.e4.

112. Schwedt A, Kreutzmann AC, Polerecky L, Schulz-Vogt HN. Sulfur respiration in a

marine chemolithoautotrophic *Beggiatoa* strain. *Front Microbiol* 2012; **2**: 276.

113. Nyström T, Neidhardt FC. Cloning, mapping and nucleotide sequencing of a gene

encoding a universal stress protein in *Eschericha coli. Mol Microbiol* 1992; **6**: 3187–

3198.

114. Aldsworth TG, Sharman RL, Dodd CER. Bacterial suicide through stress. *Cell Mol Life*

Sci . 1999. Springer. , **56**: 378–383

115.   Nyström T. Not quite dead enough: on bacterial life, culturability, senescence, and

death. *Arch Microbiol* . 2001. , **176**: 159–164

116.   Nyström T. Translational fidelity, protein oxidation, and senescence: lessons from

bacteria. *Ageing Res Rev* . 2002. Elsevier. , **1**: 693–703

117.   Sat B, Reches M, Engelberg-Kulka H. The *Escherichia coli mazEF* suicide module

mediates thymineless death. *J Bacteriol* 2003; **185**: 1803–1807.

118.   Weber H, Engelmann S, Becher D, Hecker M. Oxidative stress triggers thiol oxidation

in the glyceraldehyde-3- phosphate dehydrogenase of *Staphylococcus aureus. Mol*

Microbiol 2004; **52**: 133–140.

119.   Page R, Peti W. Toxin-antitoxin systems in bacterial growth arrest and persistence.

Nat Chem Biol 2016; **12**: 208–214.

120.   Hoor AT ten. Cell yield and bioenergetics of *Thiomicrospira denitrificans* compared

with *Thiobacillus denitrificans. Antonie Van Leeuwenhoek* 1981; **47**: 231–243.

121.   Sorokin DY, Tourova TP, Antipov AN, Muyzer G, Kuenen JG. Anaerobic growth of the

haloalkaliphilic denitrifying sulfur-oxidizing bacterium *Thialkalivibrio*

thiocyanodenitrificans sp. nov. with thiocyanate. *Microbiology* 2004; **150**: 2435–2442.

122.   Nunoura T, Takaki Y, Kazama H, Kakuta J, Shimamura S, Makita H, et al.

Physiological and genomic features of a novel sulfur-oxidizing

gammaproteobacterium belonging to a previously uncultivated symbiotic lineage

isolated from a hydrothermal vent. *PLoS One* 2014; **9**: e104959.

123.   Justin P, Kelly DP. Growth kinetics of *Thiobacillus denitrificans* in anaerobic and

aerobic chemostat culture. *J Gen Microbiol* 1978; **107**: 123–130.

124.   Mitchell JH, Leonard JM, Delaney J, Girguis PR, Scott KM. Hydrogen does not

appear to be a major electron donor for symbiosis with the deep-sea hydrothermal

vent tubeworm Riftia pachyptila. *Appl Environ Microbiol* 2019; **86**: e01522-19.

125.   Scott KM, Cavanaugh CM. CO2 uptake and fixation by endosymbiotic

chemoautotrophs from the bivalve *Solemya velum. Appl Environ Microbiol* 2007; **73**:

1174–1179.

126.   Woyke T, Teeling H, Ivanova NN, Huntemann M, Richter M, Gloeckner FO, et al.
        Symbiosis insights through metagenomic analysis of a microbial consortium. *Nature*
        2006; **443**: 950–955.

127.   Wood AP, Aurikko JP, Kelly DP. A challenge for 21st century molecular biology and
        biochemistry: what are the causes of obligate autotrophy and methanotrophy? *FEMS*
        Microbiol Rev 2004; **28**: 335–352.

128.   Tal S, Okon Y. Production of the reserve material poly-β-hydroxybutyrate and its
        function in *Azospirillum brasilense* Cd. *Can J Microbiol* 1985; **31**: 608–613.

129.   Natarajan K, Kishore L, Babu CR. Characteristics of NaCl stress associated proteins
        of *Rhizobium* under varying cultural conditions. *J Basic Microbiol* 1995; **35**: 413–420.

130.   Steinbüchel A, Füchtenbusch B. Bacterial and other biological systems for polyester
        production. *Trends Biotechnol* 1998; **16**: 419–427.

131.   Kadouri D, Burdman S, Jurkevitch E, Okon Y. Identification and isolation of genes
        involved in poly(beta-hydroxybutyrate) biosynthesis in *Azospirillum brasilense* and
        characterization of a *phbC* mutant. *Appl Environ Microbiol* 2002; **68**: 2943–2949.

132.   Ayub ND, Pettinari MJ, Ruiz JA, López NI. A polyhydroxybutyrate-producing
        Pseudomonas sp. isolated from antarctic environments with high stress resistance.
        Curr Microbiol 2004; **49**: 170–174.

133.   Kadouri D, Jurkevitch E, Okon Y, Castro-Sowinski S. Ecological and agricultural
        significance of bacterial polyhydroxyalkanoates. *Crit Rev Microbiol* 2005; **31**: 55–67.

134.   Obruca S, Sedlacek P, Krzyzanek V, Mravec F, Hrubanova K, Samek O, et al.
        Accumulation of poly(3-hydroxybutyrate) helps bacterial cells to survive freezing.
        PLoS One 2016; **11**: e0157778.

135.   Obruca S, Sedlacek P, Mravec F, Krzyzanek V, Nebesarova J, Samek O, et al. The
        presence of PHB granules in cytoplasm protects non-halophilic bacterial cells against
        the harmful impact of hypertonic environments. *N Biotechnol* 2017; **39**: 68–80.

136.   Slaninova E, Sedlacek P, Mravec F, Mullerova L, Samek O, Koller M, et al. Light

scattering on PHA granules protects bacterial cells against the harmful effects of UV

radiation. *Appl Microbiol Biotechnol* 2018; **102**: 1923–1931.

137. Kim JK, Won YJ, Nikoh N, Nakayama H, Han SH, Kikuchi Y, et al. Polyester synthesis

genes associated with stress resistance are involved in an insect-bacterium

symbiosis. *Proc Natl Acad Sci U S A* 2013; **110**: E2381–E2389.

138. Hamad MA, Austin CR, Stewart AL, Higgins M, Vázquez-Torres A, Voskuil MI.

Adaptation and antibiotic tolerance of anaerobic *Burkholderia pseudomallei*.

Antimicrob Agents Chemother 2011; **55**: 3313–3323.

139. Claessen D, Errington J. Cell wall deficiency as a coping strategy for stress. *Trends*

Microbiol 2019; **27**: 1025–1033.

140. Antia NJ, Harrison PJ, Oliveira L. The role of dissolved organic nitrogen in

phytoplankton nutrition, cell biology and ecology. *Phycologia* 1991; **30**: 1–89.

141. Walsh P, Wang Y, Campbell C, Boeck DG, Wood C. Patterns of nitrogenous waste

excretion and gill urea transporter mRNA expression in several species of marine fish.

Mar Biol 2001; **139**: 839–844.

142. Beckers G, Bendt AK, Krämer R, Burkovski A. Molecular identification of the urea

uptake system and transcriptional analysis of urea transporter- and urease-encoding

genes in Corynebacterium glutamicum. *J Bacteriol* 2004; **186**: 7645–7652.

143. Ashoori M, Saedisomeolia A. Riboflavin (vitamin B2) and oxidative stress: a review. *Br*

J Nutr 2014; **111**: 1985–1991.

144. Gould RL, Pazdro R. Impact of supplementary amino acids, micronutrients, and

overall diet on glutathione homeostasis. *Nutrients* 2019; **11**: 1056.

145. Carmel-Harel O, Storz G. Roles of the glutathione- and thioredoxin-dependent

reduction systems in the *Escherichia coli* and *Saccharomyces cerevisiae* responses

to oxidative stress. *Annu Rev Microbiol* 2000; **54**: 439–461.

146. Williams CH. Mechanism and structure of thioredoxin reductase from Escherichia coli.

FASEB J 1995; **9**: 1267–1276.

147. Vanoni MA, Curti B. Glutamate synthase: a complex iron-sulfur flavoprotein. *Cell Mol*

Life Sci 1999; **55**: 617–638.

148.   Liang J, Han Q, Tan Y, Ding H, Li J. Current advances on structure-function

relationships of pyridoxal 5′-phosphate-dependent enzymes. *Front Mol Biosci* 2019; **6**:

4.

149.   Banerjee R, Ragsdale SW. The many faces of vitamin B12: catalysis by cobalamin-

dependent enzymes. *Annu Rev Biochem* 2003; **72**: 209–247.

150.   Fontecave M, Atta M, Mulliez E. S-adenosylmethionine: nothing goes to waste.

Trends Biochem Sci 2004; **29**: 243–249.

151.   Caro AA, Cederbaum AI. Antioxidant properties of S-adenosyl-l-methionine in Fe2+-

initiated oxidations. *Free Radic Biol Med* 2004; **36**: 1303–1316.

152.   Imhoff JF. Diversity of Anaerobic Anoxygenic Phototrophic Purple Bacteria. *Modern*

Topics in the Phototrophic Prokaryotes. 2017. Springer International Publishing,

Cham, pp 47–85.

1213

 Figure legends

Figure 1. Median gene expression levels of selected *Ca.* T. oneisti metabolic

processes under anoxic, sulfidic (AS) versus oxic (O) conditions. All genes involved in

a particular process were manually collected and median expression levels ($\log_2$TPMs,

transcripts per kilobase million) per condition and process after 24 h of incubation are shown

(horizontal bold lines). Importantly, metabolic processes include both differentially and

constitutively expressed genes, and the total number of genes considered are indicated at

the bottom of each process. For the specific assignment of genes see Data S1. Note that for

the processes designated as 'Amino acids', 'Storage' and 'Vitamins & cofactors' only the

expression of the biosynthesis genes was considered. Boxes indicate interquartile ranges

(25%-75%), whiskers refer to the minimum and maximum expression values, respectively.

The individual processes are ordered according to the difference in median expression

between AS and O conditions, i.e. denitrification (far left) had the largest difference in

median expression between the two conditions, with higher median expression in the AS

condition, whereas urea utilization (far right) had the largest difference in median expression,

with higher median expression in the O condition. Metabolic processes were considered

highly expressed when their median expression level was above 5.2 $\log_2$TPM (dashed grey

line), which represents the median expression of all expressed protein-coding genes (n = 4

591) under both conditions. CBB cycle: Calvin-Benson-Bassham cycle.

1234

Figure 2. Oxidation of stored sulfur is coupled to denitrification but decoupled from

CO_2 fixation under anoxic conditions. (A) Heatmap visualizing differentially expressed

genes (2-fold change, FDR ≤ 0.05) involved in sulfur oxidation, denitrification and CO_2

fixation via the Calvin-Benson-Bassham cycle between the anoxic, sulfidic (AS) and oxic (O)

conditions after 24 h of incubation. Expression levels are visualized by displaying mean-

centered $\log_2$TPMs (transcripts per kilobase million). Upregulation is indicated in red,

downregulation in blue. Genes are ordered by function in the respective metabolic pathways.

S: sulfur, FCC: flavocytochrome c, Dsr: dissimilatory sulfite reductase, APS: adenylylsulfate,

Qmo: quinone-interacting membrane-bound oxidoreductase, Hdr: heterodisulfide reductase,

Nap: periplasmic nitrate reductase, Nir: cd_1 nitrite reductase, qNor: quinol-dependent nitric

oxide reductase, Nos: nitrous oxide reductase, Siroheme synt.: siroheme biosynthesis

(heme d precursor), RuBisCo su.: ribulose-1,5-bisphosphate carboxylase/oxygenase small

subunit, PPi-PFK: PPi-dependent phosphofructokinase, Ru5P epi.: ribulose-5-phosphate 3-

epimerase. (B) Relative elemental sulfur (S^0) content in ectosymbionts as determined by

Raman microspectroscopy after 24 h incubations under anoxic, sulfidic (AS; red dots),

anoxic without added sulfide (A, black dots) or oxic (O) conditions (O; dark blue: hypoxic,

light blue: oxygen-saturated). Each dot refers to the value obtained from measuring an

individual ectosymbiont cell; 50 cells were measured per condition. Horizontal lines display

medians, boxes show the interquartile ranges (25-75%), whiskers indicate minimum and

maximum values and different lower-case letters indicate significant differences among

conditions ($p < 0.05$, Wilcoxon rank sum test). (C) Relative ^{13}C isotope content of symbiotic

L. oneistus as determined by EA-IRMS after 24 h incubations with ^{13}C-labeled bicarbonate.

Dots refer to the values determined in individual measurements (comprising 50 worms per

measurement, for further details see Table S2). Horizontal lines indicate means, error bars

correspond to standard deviations. The categories ^{12}C-live and ^{13}C-dead refer to the natural

isotope abundance control and the dead control, respectively. Colors were assigned to the

distinct incubations as shown in panel (B). Different lower-case letters indicate significant

differences among conditions (one-way ANOVA, Tukey's post-hoc test, $p < 0.05$).

1263

Figure 3. Nitrogen fixation and urea utilization genes are upregulated under oxic

conditions. The heatmap only shows genes that were differentially expressed (2-fold

change, FDR ≤ 0.05) between anoxic, sulfidic (AS) and oxic (O) conditions after 24 h of

incubation. Expression levels are displayed as mean-centered $\log_2$TPMs (transcripts per

kilobase million). Upregulation is indicated in red, downregulation in blue. Genes are ordered

by function in the respective metabolic pathway. Cofactor synt.: cofactor biosynthesis,
Urease acc. proteins: urease accessory proteins.

1271

Figure 4. Expression of genes involved in various stress responses and vitamin
biosynthesis is upregulated in under oxic conditions and coincides with fewer
dividing symbiont cells. (A) Heatmaps showing stress-related genes and genes involved
in biosynthesis of vitamins and cofactors, as determined on the transcript level. Expression
levels are visualized by displaying mean-centered $\log_2$TPMs (transcripts per kilobase
million). Upregulation is indicated in red, and downregulation in blue. Genes are ordered by
function in the respective metabolic pathways. Heavy metal rest.: heavy metal resistance.
(B) Bars show the mean percentage of dividing *Ca.* T. oneisti cells upon 24 h incubations
under anoxic, sulfidic (AS) or oxic conditions (O). The latter condition includes both, cells
incubated under hypoxic and oxygen-saturated conditions. A total of 658, 1 009, and 1 923
cells were counted for the AS, hypoxic and oxygen-saturated condition, respectively. Error
bars indicate 95% confidence intervals (Fisher's exact test).

1284

Figure 5. Schematic representation of *Ca.* T. oneisti's suggested metabolism in deep
anoxic and superficial sand. Note that the illustrated processes likely occur under both
anoxic, sulfidic (AS) and oxic (O) conditions, but the model highlights those proposed to be
dominant in one condition over the other. Our study suggests that in anoxic, sulfidic
sediment zones (left) the ectosymbiont performs enhanced anaerobic sulfur oxidation
coupled to nitrate reduction to nitrogen gas (dentrification). Other electron acceptors (such
as DMSO) are also being reduced. The storage compound PHA serves as a carbon source
(in addition to CO_2) and as an additional electron donor. Host-derived phospholipids (P-
lipids) may be incorporated into the ectosymbiont's membrane to increase permeability. In
superficial, oxygenated zones (right), oxygen triggers a global stress response that not only
consumes energy and dampens proliferation, but also requires vitamin biosynthesis which
might increase the demand for nitrogen. Small organic carbon compounds (C_{org}), putatively

excreted by the host, contribute to energy generation (and carbon) via aerobic respiration.

Together with autotrophic CO_2 fixation, C_{org} increases carbon availability thereby allowing for

PHA biosynthesis. Heme and other essential nutrients may be directly or indirectly

transferred to the host.

Figure 1

Figure 2

Figure 3

-2 -1 0 1 2
Centered
log$_2$TPM

Figure 4

A

Stress response

Cofactor & vitamin biosynthesis

B

Figure 5