Different temporal trends in vascular plant and bryophyte communities along elevational gradients over four decades of warming

Antoine Becker-Scarpitta*[1,2,3], Diane Auberson-Lavoie[1], Raphael Aussenac[4], Mark Vellend[1]

1: Département de Biologie, Faculté des Sciences, Université de Sherbrooke, Sherbrooke, Québec, Canada.

2: Spatial Foodweb Ecology Group, Faculty of Agriculture and Forestry, Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland.

3: Institute of Botany of the Czech Academy of Sciences, Lidická 971, 60200, Brno, Czech Republic.

4: Univ. Grenoble Alpes, INRAE, LESSEM, 2 rue de la Papeterie-BP 76, St-Martin d'Hères, F-38402, France

Contact: *Corresponding author: Antoine Becker-Scarpitta, Spatial Foodweb Ecology Group, Faculty of Agriculture and Forestry, Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland. antoine.becker.scarpitta@gmail.com

ORCID

Antoine Becker-Scarpitta: 0000-0001-9241-091X

Running title

Temporal change in forest plant communities

Abstract

Despite many studies showing biodiversity responses to warming, the generality of such responses

across taxonomic groups remains unclear. Very few studies have tested for evidence of bryophyte

community responses to warming, even though bryophytes are major contributors to diversity and

functioning in many ecosystems. Here we report an empirical study comparing long-term change of

bryophyte and vascular plant communities in two sites with contrasting long-term warming trends,

using "legacy" botanical records as a baseline for comparison with contemporary resurveys. We

hypothesized that ecological changes would be greater in sites with a stronger warming trend, and

that vascular plant communities, with narrower climatic niches, would be more sensitive than

bryophyte communities to climate warming. For each taxonomic group in each site, we quantified

the magnitude of changes in species' distributions along the elevation gradient, species richness, and

community composition. We found contrasted temporal changes in bryophyte vs. vascular plant

communities, which only partially supported the warming hypothesis. In the area with a stronger

warming trend, we found a significant increase of local diversity and beta-diversity for vascular

plants, but not for bryophytes. Presence absence data did not provide sufficient power to detect

elevational shifts in species distributions. The patterns observed for bryophytes are in accordance

with recent literature showing that local diversity can remain unchanged despite strong changes in

composition. Regardless of whether one taxon is systematically more or less sensitive to

environmental change than another, our results suggest that vascular plants cannot be used as a

surrogate for bryophytes in terms of predicting the nature and magnitude of responses to warming.

Thus, to assess overall biodiversity responses to global change, abundance data from different

taxonomic groups and different community properties need to be synthesized.

Key words

Community ecology, global change ecology, vascular plant, bryophyte, warming, elevational gradient,

historical ecology, legacy data, resurvey, biodiversity, forest, understory vegetation, temporal

dynamics

Introduction

Ecological impacts of global changes on biodiversity can take many different forms (McGill et al. 2015,

Vellend et al. 2017b). In the context of climate warming, montane ecosystems have received

considerable attention due to clear predictions of changes in species' distributions (upslope shifts)

and community composition (favoring warm-adapted species) along elevation gradients. The many

studies to date have focused on relatively few taxonomic groups, in particular vascular plants,

invertebrate and vertebrates (Lenoir et al. 2008, Chen et al. 2011, Pauli et al. 2012, Stockli et al. 2012,

Rumpf et al. 2018, 2019), and there is now a need for more exhaustive multi-taxonomic syntheses

(Lenoir and Svenning 2015, Lenoir et al. 2020). The few studies of bryophyte responses to climate

warming suggest that results might be different than those for vascular plants (Walker et al. 2006,

Bergamini et al. 2009, Hudson and Henry 2010, Raabe et al. 2010, Vanneste et al. 2017, Becker-

Scarpitta et al. 2017).

Understanding variation among taxonomic groups in their responses to environmental change is

crucial for identifying priorities in conservation. However, the spatial variability of climate change

complicates comparisons between studies of different taxa in different regions, and it is difficult to

predict *a priori* differences between taxonomic groups. Thus, our limited knowledge base with which

to identify the most relevant set of metrics or the taxonomic groups most sensitive to environmental

change constrains our ability to set efficient conservation priorities. We operationally define

"sensitivity" here as responsiveness: i.e., the degree to which a given community property changes in

the face of environmental change.

To assess long-term responses of ecological communities to environmental change, "legacy"

ecological records have been used to provide baselines for comparison with contemporary resurveys

(Vellend et al. 2013, Chytrý et al. 2014, Verheyen et al. 2017, Hédl et al. 2017). However, historical

botanical studies are strongly biased towards vascular plants, with few data on bryophytes, due in

part to the difficulty of identification (Gignac 2001; He, He, and Hyvönen 2016; Möls et al. 2013 but

see Becker-Scarpitta et al., 2017; Bergamini et al., 2009; Delgado & Ederra, 2013; Outhwaite, Gregory,

Chandler, Collen, & Isaac, n.d.; Vanneste et al., 2017). This is despite the fact that bryophytes are

significant contributors to diversity in many temperate ecosystem, and a major component of the

vegetation cover in boreal or alpine ecosystems. Bryophytes play a crucial role in ecosystem

functions, such as biomass accumulation, water retention, nutrient cycling, and food web dynamics

(Turetsky 2003, Rydin 2008, Lindo and Gonzalez 2010, Tuba et al. 2011, Patiño and Vanderpoorten

2018).

For several reasons we might expect vascular plants and bryophytes to show different responses to

various sources of environmental change (Lee and La Roi 1979, Möls et al. 2013, Bagella 2014,

Vanneste et al. 2017, Becker-Scarpitta et al. 2017). Compared to vascular plants, bryophytes are

distinguished by their small size, high sensitivity to the moisture and chemistry of their immediate

microenvironment (i.e., they are poikilohydric), lower temperature optima for growth, absence of

roots and an efficient vascular system, and dispersal via spores (Glime 2007). These two groups can

show contrasted spatial patterns of diversity (Lalanne et al. 2008, Mateo et al. 2016). For instance,

vascular plants show a clear latitudinal diversity gradient of decreasing species richness with

increasing latitude, while this is not true for bryophytes, for which temperate latitudes are equally

diverse as tropical latitudes (Geffert et al. 2013, Mateo et al. 2016). Some studies have also observed

different patterns of community β-diversity (Kraft et al. 2011, Mateo et al. 2016) with vascular plant

communities sometimes showing higher β-diversity along elevation gradients than bryophytes,

suggesting a broader tolerance of bryophyte species to temperature (Lee and La Roi 1979, Vittoz et

al. 2010, Glime 2013, Vanneste et al. 2017) or differences in dispersal ability (Lenoir et al. 2012).

General predictions for the effects of warming on vascular plants include declines in the abundance

of cold-adapted species, an upward expansion of elevational range limits for warm-adapted species

(Rumpf et al. 2018), and an increase of local species richness in area without water-stress (Vellend et

al. 2017b, Steinbauer et al. 2018, Harrison 2020). Compared to vascular plants, some studies have

suggested that changes in bryophyte communities are more strongly influenced by stochastic

processes or by micro-environmental variation than macro-environmental conditions (Pharo and Vitt

2000, Raabe et al. 2010, Fenton and Bergeron 2013). Because bryophytes have wider temperature

tolerances and higher affinities to micro-environment than macro-environment, we might expect

bryophytes to show lower sensitivity to global warming than vascular plants. The consequences of

warming for β-diversity are more difficult to predict given a paucity of studies on this topic (Socolar

et al. 2016 but see Nascimbene and Spitale 2017). However, numerical simulations show that species

with high dispersal capacity will be favored in areas experiencing strong environmental changes, in

which case we might predict a decrease of β-diversity (Mouquet and Loreau 2003) and thus biotic

homogenization, at least temporarily (Clavel et al. 2011).

As in many parts of the world, eastern Canada has shown a general warming trend over the past ~50

years, but with a strong east-west gradient in the magnitude of warming in the province of Québec

(Yagouti et al. 2008). Becker-Scarpitta, Vissault, and Vellend (2019) showed that the magnitude of

temporal changes of vascular plant communities in three protected areas generally increased from

east to west in southern Québec, with greater changes in areas of stronger warming in recent decades.

For two of these three parks, the historical data also included bryophytes, thus presenting an

opportunity to test for differential sensitivity among taxonomic groups to warming. Forillon National

Park is located at the eastern tip of the province of Québec where warming has been less pronounced

than in Mont-Mégantic Provincial Park, which is in central Québec where the warming trend has been

stronger (Savage & Vellend, 2015; Yagouti et al., 2008 & Supporting Information 1).

Here we report one of the first studies comparing long-term change of bryophytes and vascular plants

communities in sites with contrasting long-term warming trends. In each of the two parks, we

revisited ~50 legacy vegetation plots initially surveyed in the 1970s, applying the same methods as

the original surveys. To minimize potentially confounding factors, plots were selected in mature

forest ecosystems that have not experienced major natural or anthropogenic disturbances during the

time between surveys. We test two main hypotheses: (**i**) For both bryophytes and vascular plants, the

park with a stronger warming trend (Mont-Mégantic) has experienced greater long-term community

changes than the park with a weaker warming trend (Forillon); (**ii**) Vascular plant communities are

more sensitive than bryophyte communities to climate warming. For each taxonomic group in each

park, we quantified the magnitude of changes in (a) species' distributions along the elevation

gradient, (b) species richness (alpha-diversity), and (c) temporal change in community composition.

Note that these predictions were already tested for vascular plants in Becker-Scarpitta et al. (2019),

to which we add data for bryophytes in this paper.

Methods

Study site

Our two study sites were located in eastern Canada: Forillon National Park in eastern Québec

(48°54′N, 64°21′W), and Mont-Mégantic Provincial Park in central Québec (45°27′N, 71°9′W).

Neither park has experienced logging or forest management over the last ~40 years. Forillon National

Park was created in 1970 and covers 245 km^2; our study plots ranged in elevation from ~50 to 500

m a.s.l. The vegetation at Forillon is dominated by a mixture of northern tree species such as *Abies*

balsamea (L.) Mill., *Picea glauca* (Moench) Voss and *Betula papyrifera* Marsh. at higher elevations, and

Acer saccharum Marsh. and *Betula alleghaniensis* Britt. at lower elevations (Majcen 1981). Mont-

Mégantic Provincial Park was created in 1994 (logging ceased in the 1960s during park planning) and

covers ~55 km². Our study plots span an elevational gradient from ~460 and 1100 m a.s.l., along

which the vegetation gradient strongly resembles that in Parc Forillon: from temperate deciduous

forests dominated by *Acer saccharum.*, *Fagus grandifolia* Ehrh. and *Betula alleghaniensis*, to boreal

forest dominated by *Abies balsamea,* and *Picea rubens* Sargent, Silva. (Marcotte and Grandtner 1974).

Average temperatures at Mont-Mégantic shows an increase of ~0.90°C between 1960 and 2005 while

at Forillon the estimated increase is about ~0.55°C for the same period (Appendix S1).

Data set

Original vegetation surveys at both sites were conducted using phytosociological methods (Marcotte

and Grandtner 1974, Majcen 1981). In each plot, authors listed all species in different strata (canopy

trees, shrubs, herbs and ground bryophytes) and each species was assigned an abundance coefficient

following Braun-Blanquet et al. (1952). For vascular plants, our analyses focused on shrubs and

herbs, which were combined into a single "understorey" stratum. All bryophyte species that were

found on the ground were recorded (i.e., on organic litter and soil surface mineral layers, not including

obvious deadwood, tree trunks and rocks); these surveys did not involve intensive searches for

individual stems of rare species within moss carpets (i.e., some locally rare species were missed).

After consulting with botanists active in Québec in the 1960s and 1970s (C. Ansseau, Z. Majcen,

personal communication, 2014), there remained uncertainty in the definition of the area over which

percent cover was evaluated for bryophytes (microhabitats vs. entire plots). Thus, we are not

confident in comparing Braun-Blanquet indices across time for bryophytes, as was done for vascular

plants in Becker-Scarpitta et al. (2019). To maximize comparability across time and taxonomic group,

we used presence-absence data for both vascular and bryophyte species in all statistical analyses.

Our approach to plot relocation is described in Becker-Scarpitta et al., (2019). In short, original survey

plots were not permanently marked, although locations were reported in maps and/or tables, such

that plots are considered "semi-permanent". In both parks, original surveyors sampled mature forests

where spatial heterogeneity was relatively low. We selected plot locations for recent surveys using

original plot maps and environmental descriptions (elevation, slope, aspect), and observations in the

field to maximize the match of current and historical abiotic conditions. In Mont-Mégantic, recent

surveys included all plots within the current park boundary. Not all plot locations in Forillon were

accessible, and plot selection for the recent surveys used the following criteria: (i) plots occurred in

forest, excluding swamps or bogs; (ii) plots were accessible via <3-4 hours hiking off of established

trails; (iii) plots had not obviously experienced major natural disturbances (e.g., storm, fire, or insect

outbreak); (iv) in the original survey the plots were sampled in mature stands that have since

maintained forest cover (i.e., no early successional dynamics in the intervening period) (Becker-

Scarpitta et al. 2019).

Original surveys in Forillon Park were conducted in June-September 1972 in 256 vegetation plots of

500 m^2 (Majcen 1981). We resurveyed 49 plots during July and August of 2015. Original surveys in

Mont-Mégantic were conducted in 1970 in 94 plots, roughly half of which fall outside the current park

boundaries. Plots were 400 m^2 in coniferous forest and 800 m^2 in broadleaved forests (Marcotte and

Grandtner 1974). We resurveyed the 48 plots falling within the current park limits at Mont-Mégantic

for vascular plants in 2012 (see Becker-Scarpitta et al., 2019) and for bryophytes during June and July

2014 (reported in the present study). We harmonized the taxonomy across both parks and periods

(see below), so the Mont-Mégantic data are not precisely the same as reported in Savage & Vellend,

(2015), but they are exactly the same as in Becker-Scarpitta et al., (2019) except transformed into

presence-absence and subsetted to keep only species with at least two occurrences (see below). Raw

data are stored in the *forestREplot* database (http://www.forestreplot.ugent.be) under the code:

N_AM_005 for Mont-Mégantic and N_AM_008 for Forillon.

Taxonomical database

Our taxonomical reference was the Taxonomic Name Resolution Service v4.0 (assessed in Feb 2017:

http://tnrs.iplantcollaborative.org) for vascular plants and Flore des bryophytes du Québec-

Labrador (Faubert, 2012, 2013, 2014) for bryophytes.

Our data set was collected by four different survey teams: one for each of the two original surveys:

Forillon: Majcen, (1981); Mont-Mégantic: Marcotte & Grandtner, (1974); one for the recent Mont-

Mégantic vascular plant survey: Savage & Velland, (2015); and one for the recent Mont-Mégantic

bryophyte survey and for the recent survey of both taxonomic groups at Forillon (A. Becker-Scarpitta

and assistants). Most plants were identified to the species level in the same way across surveys, so

for these species the only harmonization step was to standardize names. Coarser levels of taxonomic

resolution were used in some but not all surveys for certain species (e.g., a pair of similar species not

identified to the species level), and for other species (e.g., spring ephemeral species) the timing of

different surveys created doubt about the likelihood of comparable detection. In these situations,

comparability was maximized by using the coarser level of resolution applied to all data sets, or by

removing species (see Supplementary material S2 for details on taxonomic standardization). All

specimens identified at the species level were deposited in the Marie-Victorin herbarium (Université

de Montréal, Canada.) and all locations were entered into the GBIF database (GBIF -

https://www.gbif.org/).

Statistical analysis

All statistical analyses were performed in R v.3.4.2 (R Foundation for Statistical Computing 2018). To

minimize the effect of potential bias due to uncertain species identification or very rare species, all

statistical analyses used the subset of species with at least two occurrences in each time period for

each park.

First, to test for upward elevational shifts in species distributions along the elevational gradients, we

calculated the mean elevation across occurrences for each species (Z), in each park, at each time. We

modelled main effects and two- and three-way interactions between park (P), time (Y) and taxonomic

group (T), each as a categorical variable. We thus tested for upward elevational shifts in species

distributions, and how these shifts depend on park and taxonomic group. The model was given by:

$$Z = \left(\beta_1 + b_{sp} + b_{park}\right) + \beta_2 P + \beta_3 T + \beta_4 Y + \beta_5 (P.T) + \beta_6 (P.Y) + \beta_7 (T.Y) + \beta_8 (P.T.Y) + \epsilon$$

Where Z is the mean species elevation, β_{1-8} are parameters to be estimated, b_{sp} and b_{park} are random

intercepts accounting for the effect of species and park, and ϵ is the model residuals assumed to be

normally distributed and independent. Following our first hypothesis, we expected the temporal

change at Mégantic to be stronger than at Forillon (significant β_6 parameter). The second hypothesis

predicted vascular plants to be more sensitive than bryophytes (significant β_7 parameter). All other

terms of the model that are not part of our hypotheses ensure accurate estimation of other

parameters in the model.

Second, we used the exact same model structure and hypotheses to analyze temporal trends of α-

diversity of both bryophytes and vascular plants in the two parks. All linear mixed effect models were

conducted in R, with the function *lmer*, package 'lme4' v.1.1-14, Bates, Mächler, Bolker, & Walker,

2015).

Third, we tested for temporal changes in community composition using two frameworks. In one

framework, we first tested the multivariate homogeneity of group dispersions - i.e. a multivariate

beta-diversity metric - using the Jaccard dissimilarity index (PERMDISP, function *betadisper*, package

'vegan' v.2.4-4; Anderson, Ellingsen, & McArdle, (2006)). Then, we quantified the change in

community composition over time using a permutational analysis of variance (PERMANOVA,

Anderson, 2001) with Jaccard distances using 999 permutations to (functions *adonis*, package 'vegan',

and *pairwise.adonis*, package 'pairwiseAdonis'). We first tested the significance of the three-way

interaction (park * taxonomic group * time) in a full model (F=4.45, p<0.01), and second, we ran

separate models for each combination of taxonomic group and park. The R^2 values computed from

these separate PERMANOVA models were used to compare the magnitude of temporal change

between groups. We used non-metric multidimensional scaling (nMDS) with Jaccard distances for

visualization (function *metaMDS*, package 'vegan').

In a second framework, to focus on the temporal component of change in species composition, we

calculated for each plot the temporal pairwise β-diversity using the binary Jaccard dissimilarity index

(for presence-absence data) between original and recent surveys (function *beta.temp,* package

'betapart' v.1.5.2, Baselga & Orme, 2012). This framework allowed us to decompose the Jaccard

dissimilarity into turnover i.e species replacement between plot and nestedness i.e. species loss

between plot, (Baselga 2010).

As a complementary analysis, we identified for each park separately the indicator species that

characterized each period using the IndVal procedure (function *multipatt*, package 'indicspecies'; De

Cáceres, Legendre, & Moretti, 2010).

Results

Species distributions along elevation gradients

Among the four taxonomic group-site combinations, we failed to detect any significant upward shifts

in mean species elevations over time – i.e., none of the main effects or interaction terms including

time were significant (Table 1a). It is important to note that patterns reported here are not

quantitatively the same as in Becker-Scarpitta et al. (2019), who reported upward elevational shifts

of vascular plants at Mont Mégantic based on abundance-weighted mean elevations. The difference

in this paper is the use of presence-absence data, in addition to our selection of a subset of species

with at least two occurrences per survey.

Despite the lack of trends in mean species elevation for bryophytes, there was substantial variation

among species, more than for vascular plants (Table 1a, Fig. 1). This observation is reflected in the

strength of correlations between original and recent mean species elevations, which was lower for

bryophytes (Forillon Pearson r correlation = 0.33; Mont-Mégantic r = 0.61) than for vascular plants

(Forillon r = 0.52; Mont-Mégantic r = 0.73; see Supplementary material S2 for mean elevations and

sums of occurrences for each species in each survey).

Temporal changes in species richness

At Forillon, the total number of bryophyte species across plots was greater in the recent survey (38

species) than in the original survey (25 species); 6 species from the original survey were not found

while 19 new species were observed, and the two surveys shared 19 species (Supplementary material

S3 for species occurrences). Similarly, at Mont-Mégantic, there was an increase in the overall number

of bryophytes species in the recent survey (original survey = 31 vs. recent = 39 species), with 6 species

lost, 14 gained while 25 species were shared between the two surveys.

Conversely, the total number of vascular plants at Forillon declined over time (original survey = 64

vs. recent = 44 species), with 22 original species not found in the recent survey, 2 new species added

and 42 shared species. While at Mont-Mégantic, the total number of vascular plant species increased

(from 72 species to 77 species), with 8 species lost, 13 gained and 64 shared species.

We detected a significant increase in mean plot-level species richness (α-diversity) only for vascular

plants at Mont-Mégantic (Table 1b). For bryophytes in both parks, and for vascular plants at Forillon

α-diversity showed no changes over time (Table 1b).

Temporal shift in community composition and heterogeneity

We found significant temporal changes in vascular plant β-diversity and community composition at

Mont-Mégantic (Table 2, Fig. 2), where climate warming has been pronounced, as previously reported

(Becker-Scarpitta et al. 2019). However, we found an unexpected significant increase of multivariate

dispersion (β-diversity) over the past 40 years for bryophytes at Forillon but not at Mont-Mégantic

(Table 2, Fig. 2).

For both sites we found significant temporal shifts in bryophyte community composition; for vascular

plants there was significant compositional change only at Mont-Mégantic (Table 2, Fig. 2). As

predicted, the magnitude of the vascular plant compositional shift was greater for Mont-Mégantic

(R^2=0.037) than Forillon (R^2=0.019). However, bryophyte communities experienced an equal

magnitude of compositional shift for both Forillon and Mont-Mégantic (R^2=0.07 and 0.074

respectively). Moreover, compositional shifts were greater for bryophytes than for vascular plants at

both sites (Table 2).

Temporal beta-diversity showed contrasted patterns between taxonomic groups. Jaccard

dissimilarity between original and recent surveys was higher for bryophytes than for vascular plants.

The greater change in temporal beta diversity for bryophytes in the two parks is consistent with the

PERMANOVA results for changes in community composition. For all combinations of taxonomic

group and park, species turnover contributed more than nestedness to overall changes. However,

nestedness was considerably higher for vascular plants than for bryophytes (Table 2). In support of

the beta-diversity analysis, Table 3 shows the indicator species list that characterized each time

period in each park. All combinations of park and time show a complete change in the indicator

species assemblage. Even in the absence of beta-diversity change for bryophytes at Mégantic or for

vascular plants at Forillon, we detected changes in certain species frequencies. Several species

associated with recent time periods suggest possible non-climatic drivers of community change (e.g.,

Dennstaedtia punctilobula and Carex spp. are resistant to deer herbivory), a point we develop further

in the discussion.

Discussion

Most long-term legacy studies have been conducted on vascular plant communities (Verheyen et al.

2017), so it remains unclear whether bryophytes show similar or different responses to

environmental changes. Here we quantified community changes over ~40 years for both bryophytes

and vascular plants in two protected natural areas in eastern Canada with contrasting recent warming

trends. Temporal changes in vascular plant communities contrasted with the warming hypothesis.

As predicted for the area with a strong warming trend (Mont-Mégantic), we found a significant

increase of local diversity of vascular plants but no change for bryophytes (Table 1b), and a stronger

change in β-diversity and community composition for vascular plans as well (Table 2). However, we

did not detect any shift in species distribution along the elevation gradient (Table 1a, Fig. 1) and the

higher magnitude of changes in bryophyte community composition at both sites was not expected

(Table 2, Fig. 2).

Patterns along the gradient of warming trends

At Mont-Mégantic, where the warming trend has been strongest, we found a significant increase of α-

diversity (Table 1b), a significant decrease of β-diversity and a stronger shift in community

composition than in Forillon (Table 2, Fig. 2). At Forillon, where the warming trend has been weaker,

we found neither shifts in elevational distributions (Table 1a, Fig. 1) nor changes of α-diversity for

either vascular plants or bryophytes (Table 1a). The lack of upward shift in elevation of vascular

plants in response to warming is clearly explained by the use of presence/absence data, and contrasts

with our previous results using abundance data (Becker-Scarpitta et al. 2019) in addition to studies

from other sites (Lenoir et al. 2008, Chen et al. 2011, Bertrand et al. 2011),. The temporal increase of

diversity of vascular plants at Mont-Mégantic is also consistent with the prediction that warming

should lead to increased local diversity in areas without severe moisture stress (Vellend et al. 2017b,

Harrison 2020).

Our results suggest that a temperature increase of ~0.9 °C (over the last 45 years at Mont-Mégantic)

does not have as strong an impact on the local diversity of bryophytes as it does for vascular plants,

assuming that the richness increase at Mégantic was indeed due to warming. This interpretation is

also supported by the absence of any relationship between bryophyte richness and elevation in the

two parks (Forillon estimate= 0.0042 ± (SE) 0.0024, t=1.8, p=0.079 ; Mégantic estimate= 0.0022 ±

(SE) 0.0014, t=1.6, p= 0.11) and consistent with previous results in the literature (Bruun et al., 2006;

Grytnes, Heegaard, & Ihlen, 2006; Odland, Reinhardt, & Pedersen, 2014 but see Vittoz et al., 2010).

The decrease of β-diversity over time for vascular plants at Mont-Mégantic is consistent with the

hypothesis that warming might cause biotic homogenization (Urban 2015, Socolar et al. 2016). Note

that this result using presence-absence data was different to the result in Becker-Scarpitta et al.,

(2019) using untransformed abundance data (for which we found no homogenization), but similar to

the finding of homogenization using fourth-root transformed abundances in Savage & Vellend,

(2015). These results collectively illustrate that the locally dominant species – whose influence is

minimized or eliminated via fourth-root or presence-absence transformation – can mask

homogenization created by species of lower abundance. This result highlights how ecological

conclusions are dependent on the data transformation, even with widely used diversity metrics.

Sensitivity of bryophytes vs. vascular plants

We cannot draw strong conclusions about which of bryophytes or vascular plants is more or less

sensitive to environmental change in eastern Canada: results depended on which community

dimension was being investigated. Vascular plants showed more prominent richness increases

(Tables 1b) and a decrease in β-diversity, while bryophytes experienced stronger shifts in community

composition and no difference in the magnitude of compositional changes between the two parks

(Table 2, Fig. 2). We did not detect (with presence-absence data) shifts in distribution along the

elevational gradient for either taxonomic group (Table 1a, Fig. 1), while we previously found a clear

shift in elevation when using vascular plant abundance data at Mont-Mégantic (Becker-Scarpitta et

al. 2019). Our results for elevational shifts here are thus inconclusive, in that we do not know if

analyses using abundance data for bryophytes would have also shown a change in elevation.

Potential non-climatic drivers of vegetation change

Our sites were chosen specifically due to their contrasting warming trends and lack of other obvious

major drivers of vegetation change, but there are certainly other possible drivers of ecological change

that might play a role in this region. Among the indicator species of recent surveys (Table 3), two

were non-native: *Galeopsis tetrahit* and *Epipactis helleborine*, the latter of which has increased

considerably in recent decades throughout its North American range, even in western Canada (Marie-

Victorin 1997, McCune and Vellend 2013). As such, some vegetation changes might be due simply to

protracted periods of non-native species expansions, regardless of local environmental change.

Another potential factor is changes in white-tailed deer browsing, which has increased over the past

century in much of North America (Côté et al. 2004). The indicator species of the recent survey at

Mont-Mégantic (Table 3) include species known to benefit from high levels of deer browsing:

Dennstaedtia punctilobula and *Carex* spp. (de la Cretaz and Kelty 2002, Augustine and Decalesta 2003,

Rooney 2009, Frerker et al. 2014). Interestingly, at Forillon deer are actually thought to have

decreased in abundance due to the expansion of the coyote population in the 1970s (UQCN 2005),

and we found no such species associated with recent surveys in Forillon Park.

Our most difficult result to interpret was the strong species turnover of bryophyte communities at

Forillon, which has not experienced strong long-term increase in temperature, precipitation, or

atmospheric nutrient deposition (Commission Joint International 2014, Hember 2018). We can only

speculate regarding potential non-climatic drivers of bryophyte community change. First, as in all

legacy studies, there is the potential for observer biases due to (i) different sampling effort between

original and recent surveys, or (ii) species' identification errors. It seems highly likely that detection

probabilities and the potential for identification errors are greater for bryophytes than for vascular

plants, although we have no reason to suspect this caused systematic increases or decreases of

particular species frequencies (necessary to explain overall compositional shifts). We paid very close

attention to repeating the original survey methods, focusing on the visually obvious species in each

microsite (i.e., not examining each individual moss stem closely on the field), and the lack of any

difference over time in local richness suggests comparable species' detection abilities in the two

surveys. Although we cannot exclude the possibility that a real richness change was cancelled out by

a change in survey effort, we have no reason to suspect this rather unlikely coincidence. Second, given

uncertainty in the comparability of abundance estimates across time for bryophytes, we also decided

to use presence-absence data. In short, changes in observer effort seem highly unlikely to account for

the temporal change in species composition. Furthermore, our procedure to homogenize taxonomy

across data sets was quite conservative to reduce bias due to misidentifications.

One potential hypothesis to explain compositional change over time in bryophyte communities

relates to the history of park protection. Forillon Park was established (and so protected) only two

years before the original survey was conducted, and parts of the park previously included homesteads

(i.e., non-forest land uses). Mont-Mégantic was established as a park more recently (1994), but

logging activities (the only prominent land use) ceased ~15 years before the original survey. Although

plot selection focused only on non-disturbed forests, metacommunity dynamics involving dispersal

of species from sites undergoing rapid succession may have caused local shifts in composition and

increased β-diversity. It has been previously shown that managed forests tend to have a lower β-

diversity than protected forests (Kaufmann et al. 2017). The increase in bryophyte β-diversity might

partially be due to the recovery of natural forest that occurring in the 1970s.

There is also the possibility that changes in bryophyte communities resulted from interactions with

changing vascular plant communities. Studies have shown that bryophyte diversity and abundance

can be negatively correlated with total vascular plant biomass (Virtanen et al. 2017), cover (Jiang et

al. 2015) or abundances (Jägerbrand et al. 2012). While we have documented an increase of vascular

plant species richness at Mont-Mégantic (Tables 1b), we do not have data on vascular plant biomass.

If bryophytes are highly sensitive to vascular plant community properties, subtle changes for vascular

plants could translate into larger changes in bryophyte communities. This hypothesis is open to

testing via observational and experimental studies of the effect of vascular plants on bryophytes

communities under warming or other environmental changes. Understanding temporal changes in

one component of the community may require more consideration of interactions with other

components (Chesson 2000, HilleRisLambers et al. 2012).

Conclusion and conservation implications

Overall, we found a significant temporal shift in the composition of both taxonomic groups in both

parks but only one significant change in α-diversity. This study shows the limitation of using presence

absence data to detect shift in distribution. While abundance data provide the necessary power to

detect changes in species distributions along the elevational gradient (Becker-Scarpitta et al. 2019),

presence/absence data clearly do not. Our results are in accordance with recent meta-analyses and

syntheses showing that local diversity can remain unchanged (or increase or decrease with equal

likelihood) despite strong changes in composition (Dornelas et al. 2014, Vellend et al. 2017a, Gotelli

et al. 2017, Spaak et al. 2017, Magurran et al. 2018, Blowes et al. 2019). Finally, regardless of whether

one taxonomic group is systematically more or less sensitive to environmental change than another,

our results suggest that one taxonomic group (e.g., vascular plants) cannot be used as a surrogate for

others (e.g., bryophytes) in terms of predicting the nature and magnitude of responses to

environmental change (Bagella 2014, Becker-Scarpitta et al. 2017, Outhwaite et al. 2020). In the same

plots that experienced the same environmental changes, we found that communities of bryophytes

and vascular plants did not predictably change in the same ways (Slack 1977, Lalanne et al. 2008,

Odland et al. 2014, Bagella 2014, Becker-Scarpitta et al. 2017). Thus, to assess overall biodiversity

responses to global change abundance data from different taxonomical groups and different

community properties need to be synthesized.

Acknowledgements

We would like to thank the field assistants who greatly contributed to the collection of data in the field, the identification of species in the laboratory and the realization of the herbarium: Melissa Paquette and Sara Gaignard. This work was made possible thanks to the support of park managers, especially Camille-Antoine Ouimet (at Mont-Mégantic) and Daniel Sigouin (at Forillon). We also thank Guillaume Blanchet, Michael Belluau and Willian Vieira for constructive discussion on this project. The project was funds by the Natural Sciences and Engineering Research Council, Canada and field mission were support by the Quebec Center for Biodiversity Sciences (QCBS).

Data Accessibilty Statement

Data are stored in the *forestREplot* database ([http://forestreplot.ugent.be/)](http://forestreplot.ugent.be/) under the code: N_AM_005 for Mont-Mégantic and N_AM_008 for Forillon. All plant specimens identified at the species level were deposited in the Marie-Victorin herbarium (Université de Montréal, Canada.) and all locations were entered into the GBIF database (GBIF - https://www.gbif.org/).

Supplementary Materials

- Appendix S1 – Climatic temporal trends of the two parks.
- Supplementary material S2 – Taxonomic standardization of bryophytes species
- Supplementary material S3 – Species occurrences and mean species elevations

References

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance. - Austral Ecol. 26: 32–46.

Anderson, M. J. et al. 2006. Multivariate dispersion as a measure of beta diversity. - Ecol. Lett. 9:

683–693.

Augustine, D. J. and Decalesta, D. 2003. Defining deer overabundance and threats to forest

communities: From individual plants to landscape structure. - Écoscience 10: 472–486.

Bagella, S. 2014. Does cross-taxon analysis show similarity in diversity patterns between vascular

plants and bryophytes? Some answers from a literature review. - C. R. Biol. 337: 276–282.

Baselga, A. 2010. Partitioning the turnover and nestedness components of beta diversity. - Glob.

Ecol. Biogeogr. 19: 134–143.

Baselga, A. and Orme, C. D. L. 2012. betapart : an R package for the study of beta diversity. - Methods

Ecol. Evol. 3: 808–812.

Bates, D. et al. 2015. Fitting Linear Mixed-Effects Models Using lme4. - J. Stat. Softw. 67: 1–48.

Becker-Scarpitta, A. et al. 2017. Long-term community change: bryophytes are more responsive

than vascular plants to nitrogen deposition and warming. - J. Veg. Sci. 28: 1220–1229.

Becker-Scarpitta, A. et al. 2019. Four decades of plant community change along a continental

gradient of warming. - Glob. Chang. Biol. 25: 1629–1641.

Bergamini, A. et al. 2009. An elevational shift of cryophilous bryophytes in the last century - an effect

of climate warming? - Divers. Distrib. 15: 871–879.

Bertrand, R. et al. 2011. Changes in plant community composition lag behind climate warming in

lowland forests. - Nature 479: 517–520.

Blowes, S. A. et al. 2019. The geography of biodiversity change in marine and terrestrial

assemblages. - Science (80-.). 366: 339–345.

Braun-Blanquet, J. et al. 1952. Les Groupements Végétaux de la France Méditerranéenne (CNRS,

Ed.). - CNRS.

Bruun, H. H. et al. 2006. Effects of altitude and topography on species richness of vascular plants,

bryophytes and lichens in alpine communities. - J. Veg. Sci. 17: 37–46.

Chen, I.-C. et al. 2011. Rapid Range Shifts of Species Associated with High Levels of Climate

Warming. - Science (80-.). 333: 1024–1026.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. - Annu. Rev. Ecol. Syst. 31: 343–

366.

Chytrý, M. et al. 2014. Assessing vegetation change using vegetation-plot databases: a risky

business. - Appl. Veg. Sci. 17: 32–41.

Clavel, J. et al. 2011. Worldwide decline of specialist species: toward a global functional

homogenization? - Front. Ecol. Environ. 9: 222–228.

Commission Joint International 2014. Canada - United States, Air quality agreement - progress

report.

Côté, S. D. et al. 2004. Ecological impacts of Deer overabundance. - Annu. Rev. Ecol. Evol. Syst. 35:

113–147.

De Cáceres, M. et al. 2010. Improving indicator species analysis by combining groups of sites. - Oikos

119: 1674–1684.

de la Cretaz, A. L. and Kelty, M. J. 2002. Development of tree regeneration in fern-dominated forest

understories after reduction of Deer browsing. - Restor. Ecol. 10: 416–426.

Delgado, V. and Ederra, A. 2013. Long-term changes (1982–2010) in the bryodiversity of Spanish

beech forests assessed by means of Ellenberg indicator values of temperature, nitrogen, light

and pH. - Biol. Conserv. 157: 99–107.

Dornelas, M. et al. 2014. Assemblage time series reveal biodiversity change but not systematic loss. -

Science (80-.). 344: 296–299.

Fenton, N. J. and Bergeron, Y. 2013. Stochastic processes dominate during boreal bryophyte

community assembly. - Ecology 94: 1993–2006.

Frerker, K. et al. 2014. Long-term regional shifts in plant community composition are largely

explained by local deer impact experiments. - PLoS One 9: e115843.

Geffert, J. L. et al. 2013. Global moss diversity: spatial and taxonomic patterns of species richness. - J.

Bryol. 35: 1–11.

Gignac, L. D. 2001. Bryophytes as indicators of climate change. - Bryologist 104: 410–420.

Glime, J. M. 2007. Bryophyte Ecology. Volume 1. Physiological Ecology. - Ebook sponsored by

Michigan Technological University and the International Association of Bryologists.

Glime, J. M. 2013. Life cycles : surviving change, Chap 2.2. In Bryophyte Ecology. Vol. 1 Physiology

and Ecology. - Ebook sponsored by Michigan Technological University and the International

Association of Bryologists.

Gotelli, N. J. et al. 2017. Community-level regulation of temporal trends in biodiversity. - Sci. Adv. 3:

e1700315.

Grytnes, J. A. et al. 2006. Species richness of vascular plants, bryophytes, and lichens along an

altitudinal gradient in western Norway. - Acta Oecologica 29: 241–246.

Harrison, S. 2020. Plant community diversity will decline more than increase under climatic

warming. - Philos. Trans. R. Soc. B Biol. Sci. 375: 20190106.

He, X. et al. 2016. Will bryophytes survive in a warming world? - Perspect. Plant Ecol. Evol. Syst. 19:

49–60.

Hédl, R. et al. 2017. Resurvey of historical vegetation plots: a tool for understanding long-term

dynamics of plant communities. - Appl. Veg. Sci. 20: 161–163.

Hember, R. A. 2018. Spatially and temporally continuous estimates of annual total nitrogen

deposition over North America, 1860–2013. - Data Br. 17: 134–140.

HilleRisLambers, J. et al. 2012. Rethinking community assembly through the lens of coexistence

theory. - Annu. Rev. Ecol. Evol. Syst. 43: 227–248.

Hudson, J. M. G. and Henry, G. H. R. 2010. High Arctic plant community resists 15 years of

experimental warming. - J. Ecol. 98: 1035–1041.

Jägerbrand, A. K. et al. 2012. Effects of neighboring vascular plants on the abundance of bryophytes

in different vegetation types. - Polar Sci. 6: 200–208.

Jiang, Y. et al. 2015. Diversity and distribution of ground bryophytes in broadleaved forests in

Mabian Dafengding National Nature Reserve, Sichuan, China. - Acta Ecol. Sin. 35: 13–19.

Kaufmann, S. et al. 2017. Comparing the plant diversity of paired beech primeval and production

forests: Management reduces cryptogam, but not vascular plant species richness. - For. Ecol.

Manage. 400: 58–67.

Kraft, N. J. B. et al. 2011. Disentangling the drivers of diversity along latitudinal and elevational

gradients. - Science (80-.). 333: 1755–1758.

Lalanne, A. et al. 2008. Opposite responses of vascular plant and moss communities to changes in

humus form, as expressed by the Humus Index. - J. Veg. Sci. 19: 645–652.

Lee, T. D. and La Roi, G. H. 1979. Bryophyte and understory vascular plant beta diversity in relation

to moisture and elevation gradients. - Vegetatio 40: 29–38.

Lenoir, J. and Svenning, J.-C. 2015. Climate-related range shifts - a global multidimensional synthesis

and new research directions. - Ecography (Cop.). 38: 15–28.

Lenoir, J. et al. 2008. A significant upward shift in plant species optimum elevation during the 20th

century. - Science (80-.). 320: 1768–1771.

Lenoir, J. et al. 2012. Dispersal ability links to cross-scale species diversity patterns across the

Eurasian Arctic tundra. - Glob. Ecol. Biogeogr. 21: 851–860.

Lenoir, J. et al. 2020. Species better track climate warming in the oceans than on land. - Nat. Ecol.

Evol. in press.

Lindo, Z. and Gonzalez, A. 2010. The bryosphere: An integral and influential component of the

Earth's biosphere. - Ecosystems 13: 612–627.

Magurran, A. E. et al. 2018. Divergent biodiversity change within ecosystems. - Proc. Natl. Acad. Sci.

115: 1843–1847.

Majcen, Z. 1981. Les forets du parc national Forillon, Gaspésie, Québec, étude phytosociologique.

Marcotte, G. and Grandtner, M. 1974. Étude écologique de la végétsation forestière du Mont

Mégantic.

Marie-Victorin, F. 1997. Flore laurentienne. Troisième édition mise à jour et annotée par L. Brouillet,

SG Hay, I. Goulet, M. Blondeau, J. Cayouette et J. Labrecque. - Les Presses de l'Université de

Montréal.

Mateo, R. G. et al. 2016. The mossy north: an inverse latitudinal diversity gradient in European

bryophytes. - Sci. Rep. 6: 25546.

McCune, J. L. and Vellend, M. 2013. Gains in native species promote biotic homogenization over four

decades in a human-dominated landscape. - J. Ecol. 101: 1542–1551.

McGill, B. J. et al. 2015. Fifteen forms of biodiversity trend in the Anthropocene. - Trends Ecol. Evol.

30: 104–113.

Möls, T. et al. 2013. Global gradients in moss and vascular plant diversity. - Biodivers. Conserv. 22:

1537–1551.

Mouquet, N. and Loreau, M. 2003. Community patterns in source-sink metacommunities. - Am. Nat.

162: 544–557.

Nascimbene, J. and Spitale, D. 2017. Patterns of beta-diversity along elevational gradients inform

epiphyte conservation in alpine forests under a climate change scenario. - Biol. Conserv. 216:

26–32.

Odland, A. et al. 2014. Differences in richness of vascular plants, mosses, and liverworts in southern

Norwegian alpine vegetation. - Plant Ecol. Divers. 00: 1–11.

Outhwaite, C. L. et al. 2020. Complex long-term biodiversity change among invertebrates,

bryophytes and lichens. - Nat. Ecol. Evol. 4: 384–392.

Patiño, J. and Vanderpoorten, A. 2018. Bryophyte Biogeography. - CRC. Crit. Rev. Plant Sci. 37: 175–

209.

Pauli, H. et al. 2012. Recent Plant Diversity Changes on Europe's Mountain Summits. - Science (80-.).

336: 353–355.

Pharo, E. J. and Vitt, D. H. 2000. Local variation in bryophyte and macro-lichen cover and density in

diversity in montane forests of western Canada. - Bryologist 103: 455–466.

Raabe, S. et al. 2010. Drivers of bryophyte diversity allow implications for forest management with a

focus on climate change. - For. Ecol. Manage. 260: 1956–1964.

Rooney, T. P. 2009. High white-tailed deer densities benefit graminoids and contribute to biotic

homogenization of forest ground-layer vegetation. - Plant Ecol. 202: 103–111.

Rumpf, S. B. et al. 2018. Range dynamics of mountain plants decrease with elevation. - Proc. Natl.

Acad. Sci. 115: 1848–1853.

Rumpf, S. B. et al. 2019. Elevational rear edges shifted at least as much as leading edges over the last

century (J Lenoir, Ed.). - Glob. Ecol. Biogeogr. 28: 533–543.

Rydin, H. 2008. Population and community ecology of bryophytes. - In: Bryophyte Biology.

Cambridge University Press, pp. 393–444.

Savage, J. and Velend, M. 2015. Elevational shifts, biotic homogenization and time lags in vegetation

change during 40 years of climate warming. - Ecography (Cop.). 38: 546–555.

Slack, N. G. 1977. Species diversity and community structure in bryophytes: New York State studies.

- New York State Museum, The University of the State of New York.

Socolar, J. et al. 2016. How should beta-diversity inform biodiversity conservation? - Trends Ecol.

Evol. 31: 67–80.

Spaak, J. W. et al. 2017. Shifts of community composition and population density substantially affect

ecosystem function despite invariant richness. - Ecol. Lett. 20: 1315–1324.

Steinbauer, M. J. et al. 2018. Accelerated increase in plant species richness on mountain summits is

linked to warming. - Nature 556: 231–234.

Stockli, V. et al. 2012. Using historical plant surveys to track biodiversity on mountain summits. -

Plant Ecol. Divers. 4: 415–425.

Tuba, Z. et al. 2011. Bryophyte ecology and climate change. - Cambridge University Press.

Turetsky, M. R. 2003. The role of Bryophytes in carbon and nitrogen cycling. - Bryologist 106: 395–

409.

UQCN 2005. Rapport de synthèse - Parc National de Forillon. Importance et impacts des pressions

périphériques sur le maintien de l'intégrité écologique des aires protégées au Québec.

Urban, M. C. 2015. Accelerating extinction risk from climate change. - Science (80-.). 348: 571–573.

Vanneste, T. et al. 2017. Impact of climate change on alpine vegetation of mountain summits in

Norway. - Ecol. Res. 32: 579–593.

Velend, M. et al. 2013. Historical ecology: Using unconventional data sources to test for effects of

global environmental change. - Am. J. Bot. 100: 1294–1305.

Vellend, M. et al. 2017a. Estimates of local biodiversity change over time stand up to scrutiny. -

Ecology 98: 583–590.

Vellend, M. et al. 2017b. Plant biodiversity change across scales during the anthropocene. - Annu.

Rev. Plant Biol. 68: 563–586.

Verheyen, K. et al. 2017. Combining biodiversity resurveys across regions to advance global change

research. - Bioscience 67: 73–83.

Virtanen, R. et al. 2017. Comparing the responses of bryophytes and short-statured vascular plants

to climate shifts and eutrophication (S Power, Ed.). - Funct. Ecol. 31: 946–954.

Vittoz, P. et al. 2010. Subalpine-nival gradient of species richness for vascular plants, bryophytes

and lichens in the Swiss Inner Alps. - Bot. Helv. 120: 139–149.

Walker, M. D. et al. 2006. Plant community responses to experimental warming across the tundra

biome. - Proc. Natl. Acad. Sci. U. S. A. 103: 1342–1346.

Yagouti, A. et al. 2008. Observed changes in daily temperature and precipitation indices for southern

Québec, 1960–2005. - Atmosphere-Ocean 46: 243–256.

Figures and Tables

Figure 1 – Mean elevations of species' distributions in two time periods and two parks. A) Bryophytes,

Forillon = 19 species, Mégantic = 25 species); B) Vascular plants, Forillon = 42 species, Mégantic = 64

species. All species present in both the original and recent surveys were included. Diagonal lines

indicate 1:1 (i.e. no changes over time). See Table 1a for statistical analysis.

Figure 2 – Temporal changes in (A) bryophyte and (B) vascular plant community composition over time in Forillon and Mont-Mégantic, based on Jaccard's dissimilarity and non-metric multidimensional scaling (NMDS). At both sites, bryophytes showed significant shifts in composition over time and a significant increase in beta diversity. For vascular plants significant shifts in community composition and a significant decrease in beta diversity were found at Mont Mégantic but not Forillon (Table 2).

Table 1 - Analyses of temporal changes in (a) mean species elevation, and (b) species richness. For

both models, parameter estimates ± credible intervals (CI) are reported from linear mixed models.

The baseline condition (Intercept) against which effects are tested was bryophytes at Parc Forillon in

the initial survey. R^2m is the marginal coefficient of determination, measuring the proportion of

variance explained by fixed effects i.e., time, park, and taxonomic group; R^2c is the conditional

coefficient of determination, giving the proportion of variance explained by both fixed and random

effects. Names in parentheses refer to the level of the variable tested. In categorical models, the

Intercept is the reference level used to test the effect of other variables. Bold values indicate

statistically significant differences (p<0.05).

	Parameters	Estimates	CI	p	df
(a) Mean species elevation					
Intercept (Forillon, bryo, t0)	β1	189.28	156.35 – 222.22	**<0.001**	216.20
park (Mégantic)	β2	504.98	461.29 – 548.67	**<0.001**	216.20
group_taxo (Vascular)	β3	-8.94	-48.63 – 30.75	0.658	216.20
time (t1)	β4	1.50	-28.33 – 31.33	0.921	146.00
park (Mégantic) * group_taxo (Vascular)	β5	-65.38	-117.55 – -13.21	**0.014**	216.20
park (Mégantic) * time (t1)	β6	-23.48	-63.05 – 16.09	0.243	146.00
group_taxo (Vascular) * time (t1)	β7	9.67	-26.28 – 45.61	0.596	146.00
park (Mégantic) * group_taxo (Vascular) * time (t1)	β8	28.14	-19.11 – 75.39	0.241	146.00
Random Effects					
Variance park:species	bpark, bsp	3141.17			
Variance residual		2163.68			
Marginal R^2 / Conditional R^2		0.905 / 0.961			
(b) Species richness					
Intercept (Forillon, bryo, t0)	β1	5.00	3.28 – 6.73	**<0.001**	348.96
park (Mégantic)	β2	0.28	-2.16 – 2.72	0.820	354.45
group_taxo (Vascular)	β3	12.83	10.61 – 15.05	**<0.001**	275.14
time (t1)	β4	0.89	-1.34 – 3.12	0.431	273.75
park (Mégantic) * group_taxo (Vascular)	β5	2.67	-0.49 – 5.83	0.098	287.09
park (Mégantic) * time (t1)	β6	-1.07	-4.20 – 2.07	0.504	276.07
group_taxo (Vascular) * time (t1)	β7	-2.81	-5.95 – 0.33	0.079	273.75
park (Mégantic) * group_taxo (Vascular) * time (t1)	β8	8.75	4.33 – 13.18	**<0.001**	274.91
Random Effects					
Variance park:plot	bpark, bplot	6.13			
Variance residual		30.18			
Marginal R^2 / Conditional R^2		0.639 / 0.700			

650 Table 2 – Temporal changes in beta diversity and community composition. β-diversity was measured

651 as the multivariate distance of plots to the time-specific centroid in multivariate space. The R^2 of the

652 shift in community composition reflects the proportion of variation in community composition

653 explained by time. Jaccard temporal β-diversity values are decomposed into turnover and nestedness

654 between original and recent surveys. Bold values indicate significant statistical differences p<0.05.

		β-diversity			Composition			Temporal beta diversity		
		Original	Recent	p	F	p.adj	R2	Jaccard	Turnover(%)	Nestedness(%)
Bryophytes										
	Forillon	0.53	0.58	**0.015**	6.93	**0.006**	0.070	0.80±0.02	85.5%	14.5%
	Megantic	0.61	0.60	0.93	7.64	**0.006**	0.074	0.89±0.02	96.4%	3.6%
Vascular plants										
	Forillon	0.45	0.43	0.80	1.86	0.126	0.019	0.52±0.02	69.9%	29.9%
	Megantic	0.50	0.43	**0.004**	3.60	**0.024**	0.037	0.53±0.02	70.8%	29.2%

656   Table 3 - Indicator species of bryophytes and vascular plants associated with original and recent

657   surveys at Forillon and Mont-Mégantic. Indicator values were considered significant if p<0.05

658   (calculated with 999 permutations).

Bryophytes			
Forillon, original	stat	p.value	
Dicranum fuscescens	0.74	0.001	***
Forillon, recent			
Dicranum montanum	0.625	0.002	**
Dicranum polysetum	0.554	0.001	***
Brachythecium campestre	0.484	0.001	***
Pseudoleskeella tectorum	0.413	0.009	**
Plagiochila porelloides	0.357	0.026	*
Mégantic, original			
Brotherella recurvans	0.729	0.001	***
Dicranum fuscescens	0.639	0.001	***
Paraleucobryum longifolium	0.531	0.001	***
Hypnum pallescens	0.484	0.001	***
Bryhnia novae-angliae	0.438	0.035	*
Brachythecium reflexum	0.386	0.006	**
Mégantic, recent			
Atrichum altecristatum	0.600	0.001	***
Hylocomiastrum umbratum	0.543	0.003	**
Dicranum polysetum	0.447	0.002	**
Hypnum curvifolium	0.447	0.001	***

Vascular plants			
Forillon, original	stat	p.value	
Mitella nuda	0.497	0.008	**
Lycopodium annotinum	0.479	0.002	**
Galium triflorum	0.476	0.021	*
Moneses uniflora	0.447	0.029	*
Carex arctata	0.382	0.012	*
Epilobium angustifolium	0.354	0.031	*
Forillon, recent			
Sambucus racemosa	0.476	0.011	*
Epipactis helleborine	0.433	0.003	**

Mégantic, original

Osmunda claytoniana	0.433	0.004	**
Botrychium virginianum	0.408	0.006	**

Mégantic, recent

Dennstaedtia punctilobula	0.677	0.001	***
Carex arctata	0.594	0.001	***
Carex deweyana	0.520	0.001	***
Circaea alpina	0.476	0.017	*
Osmundastrum cinnamomeum	0.424	0.049	*
Epipactis helleborine	0.408	0.007	**
Galeopsis tetrahit	0.408	0.008	**
Lycopodium annotinum	0.354	0.030	*
Habenaria sp	0.354	0.025	*

Appendix S1 - Climatic temporal trends

To model the mean annual temperature trend in both parks, we extracted temperature data from the

ANUSPLINE model (McKenney et al. 2011). The relevant period to study the effect of temperature

change on forest plant communities includes a lag of approximately ~10 years before the time of

the survey. We thus considered the period 1960-2005, which aligns with a previous study showing

a sharp warming gradient in the region (Yagouti et al. 2008). The model includes the effect of the

categorical variable park (with two levels: Forillon and Mégantic), the continuous variable year

from 1960 to 2005, and a random effect on the different measures within each park.

We only report results for annual mean temperature but found similar results for annual minimum

and maximum temperatures, with no temporal change in annual mean precipitation. For the period

1960-2005, we find that Forillon experienced an increase of 0.12 °C ± 0.010°C/decade, while at

Mégantic the increase per decade was almost twice as strong: 0.20 °C ± 0.014°C/decade.

	Mean annual temperature			
	Estimates	*Sdt.Error*	t	*p*
Intercept (Forillon)	-29.04	7.505	-3.87	**<0.001**
Year	0.012	0.0010	11.88	**<0.001**
Year : Mégantic	0.008	0.0014	5.49	**<0.001**
Random Effects				
Variance residual	0.436			
Variance plot:park	0.429			
Variance park	108.725			
Marginal R^2 / Conditional R^2	0.340 / 0.997			