Conscious perception and the modulatory role of dopamine: no effect of the dopamine D2 agonist cabergoline on visual masking, the attentional blink, and probabilistic discrimination

Boonstra, E.A.,[1,2] van Schouwenburg, M.R.,[2] Seth, A.K.,[3,4] Bauer, M.,[5] Zantvoord, J.B.,[6] Kemper, E.M.,[7] Lansink, C.S.[8] & Slagter, H.A.[1,2]

[1] Department of Experimental and Applied Psychology, Vrije Universiteit Amsterdam, Institute for Brain and Behavior Amsterdam (iBBA), The Netherlands
[2] Department of Psychology, University of Amsterdam, Amsterdam Brain and Cognition (ABC), The Netherlands
[3] Sackler Centre for Consciousness Science, Department of Informatics, University of Sussex, Brighton, BN1 9QJ, UK
[4] Canadian Institute for Advanced Research, Azrieli Programme on Brain, Mind, and Consciousness
[5] School of Psychology, University of Nottingham, UK
[6] The Bascule, Academic Centre for Child and Adolescent Psychiatry, Amsterdam, the Netherlands; Department of Child and Adolescent Psychiatry, Amsterdam University Medical Centers, the Netherlands.
[7] Department of Pharmacy, Amsterdam University Medical Centers, the Netherlands
[8] Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam Brain and Cognition (ABC), The Netherlands

Abstract

Rationale. Conscious perception is thought to depend on global amplification of sensory input. In recent years, striatal dopamine has been proposed to be involved in gating information and conscious access, due to its modulatory influence on thalamocortical connectivity.

Objectives. Since much of the evidence that implicates striatal dopamine is correlational, we conducted a double-blind crossover pharmacological study in which we administered cabergoline – a dopamine D2 agonist – and placebo to 30 healthy participants. Under both conditions, we subjected participants to several well-established experimental conscious-perception paradigms, such as backward masking and the attentional blink task.

Results. We found no evidence in support of an effect of cabergoline on conscious perception: key behavioral and event-related potential (ERP) findings associated with each of these tasks were unaffected by cabergoline.

Conclusions. Our results cast doubt on a causal role for dopamine in visual perception. It remains an open possibility that dopamine has causal effects in other tasks, perhaps where perceptual uncertainty is more prominent.

Keywords: dopamine, cabergoline, basal ganglia, striatum, consciousness, backward masking, attentional blink, spontaneous eye blink rate, EEG, event-related potential

Number of words: 10243 (excl. reference list and title page)

50
Funding information
This work was supported by an ERC starting grant to HAS by the H2020 European Research
Council [ERC-2015-STG-679399] and an Amsterdam Brain and Cognition (ABC) grant to
HAS, AKS and CSL. AKS is additionally grateful to the Canadian Institute for Advanced
Research (CIFAR) Azrieli Programme on Brain, Mind, and Consciousness, and to the Dr.
Mortimer and Theresa Sackler Foundation.
57
58 **The authors declare no competing financial interests.**

Introduction

The relationship between consciousness and the brain is often lauded as one of the big
mysteries in contemporary science. How does the brain constrain its own spontaneous
activity as well as the influences it undergoes from outside, in the determination of conscious
awareness? Several influential theories propose that consciousness is related to the
'broadcasting' of sensory information to the whole brain and that thalamocortical circuits
serve as an important mediator of such broadcasting (Crick & Koch, 2003; Dehaene &
Changeux, 2011; Edelman, 2003). The broadcasting of sensory information necessitates the
occurrence of selection or filtering, simply because not everything which takes place in the
brain reaches conscious awareness.
68
With the requirement of selection, we cannot ignore the role of the basal ganglia: a
cluster of subcortical nuclei located deep in the brain, which modulate activity of
thalamocortical circuits (Smith, Raju, Pare, & Sidibe, 2004) and have long been implicated in
action selection; the decision to execute one of several possible behaviors (Redgrave,
Prescott, & Gurney, 1999). Notably, the basal ganglia are connected through parallel loops
via the thalamus not only to the motor cortex, but to many parts of frontal cortex as well
(Alexander, DeLong, & Strick, 1986). Hence, it is capable of modulating a wide range of
cognitive operations. Indeed, the basal ganglia have been implicated in working memory
updating (Frank & O'Reilly, 2006), attention shifting (Cools, 2011), and visual categorization
(Seger, 2008): cognitive acts that support suggestions concerning the common principle
underlying the basal ganglia's operations; namely, selection (Frank, Loughry, & O'Reilly,
2001; Redgrave, Prescott, & Gurney, 1999).
The striatum – the biggest structure constituting the basal ganglia – deserves special
attention on this topic. As the basal ganglia's primary input nucleus, the striatum is well-
positioned to play a pivotal role in the basal ganglia's selective functionalities, as terminal
fields from different cortical regions converge in the striatum (Yeterian & van Hoesen, 1978;
Haber, Kim, Mailly, & Calzavara, 2006; Mailly, Aliane, Groenewegen, Haber, & Deniau,
2013; Heilbronner, Meyer, Choi, & Haber, 2018). While dopaminergic projections are
usually associated with reward prediction error (Schultz, 2016), and the role of reward in
perceptual decision making (Ding & Gold, 2013); less well-studied signaling of the
(dopamine-infused) striatum include saliency, threat, processing of sensory information, and
promoting of behaviors reliant on sensory information (Cox & Witten, 2019).
Despite the suggestion that the basal ganglia may be "mute" as regards to
consciousness (Boly et al., 2017), it has been known for a long time that basal ganglia
structures are involved in sensory and perceptual processes (Alexander & Crutcher, 1990;
Arsalidou, Duerden, & Taylor, 2013; L. L. Brown, Schneider, & Lidsky, 1997; Seger, 2013).
Both the striatum and dopaminergic firing have been implicated in the tight relationship
between perception and action, and consciousness more generally. For example, dopamine-
depleted mice are unable to attend to salient sensory information and choose appropriate
actions, suggestive of a critical role for dopamine in the expression of consciousness
(Palmiter, 2011). In humans as well, it has recently been shown that minimally conscious
patients suffer from a dopaminergic deficit in presynaptic neurons projecting to the striatum
and central thalamus (Fridman, Osborne, Mozley, Victor, & Schiff, 2019).
In humans, the striatum and its irrigation by dopamine have also been implicated in
well-known experimental paradigms used to study the neural correlates of consciousness,
such as the attentional blink and backward masking task. In the attentional blink task, a
deficit occurs when people have to detect two target stimuli (T1 and T2) presented in close
temporal succession among distracter events. Specifically, when T2 follows T1 within 100–
500 ms, it often goes unnoticed. This deficit is called the attentional blink (AB; Shapiro,
Raymond, & Arnell, 1997). Healthy participants with more D2-like receptor binding in the

striatum – as shown with PET – showcased a larger AB (Slagter et al., 2012). In addition,
intracranial EEG recordings in the ventral striatum revealed a short-latency increase in theta-
band oscillatory activity only for consciously perceived target stimuli (Slagter et al., 2017).

In backward masking tasks, processing of target stimuli is interrupted by presenting a
mask in close succession to the target (Breitmeyer, 2007). Studies employing fMRI
consistently show differences in BOLD activity in the striatum and thalamus between seen
and unseen stimuli using backward masking tasks (Bisenius, Trapp, Neumann, & Schroeter,
2015). In another PET study, dopamine D2 binding potential in the right striatum was found
to correlate positively with both objective (task performance) and subjective (seen/unseen)
visibility during backward masking (Van Opstal et al., 2014). These studies collectively
suggest a role for the striatum and dopaminergic activity in the selection of visual
information and the formation of conscious visual percepts.

However, up until now the relationship between striatal dopamine and conscious
perception is based on correlational evidence. As such, in the present study we sought to
manipulate this relationship experimentally, by administering the dopamine D2 agonist
cabergoline to healthy participants. Out of the two main dopamine receptor families, D2
receptors have been found to be more prevalent in the striatum, while D1 receptors are
present more in the prefrontal cortex (Gerfen, 1992). We chose cabergoline because it has
greater affinity for D2 receptors, and it has been reported to have less side-effects compared
to other D2 agonists such as bromocriptine (Frank & O'Reilly, 2006). Cabergoline, at low
doses, has been suggested to preferentially stimulate pre-synaptic D2 autoreceptors, which
have been found to inhibit phasic dopamine bursts in the striatum (Ford, 2014; Frank &
O'Reilly, 2006). Cabergoline has been successfully administered in small dosages (1-1.5 mg)
to manipulate performance in healthy participants on working memory tasks (Broadway,
Frank, & Cavanagh, 2018; Fallon, Zokaei, Norbury, Manohar, & Husain, 2017; Frank &
O'Reilly, 2006), a modified version of the Simon task (Cavanagh, Masters, Bath, & Frank,
2014), as well as action cancellation and error awareness tasks (Nandam et al., 2013). When
cabergoline was administered to Parkinson's patients for longer periods of time, decreases
contrast sensitivity was found (Hutton, Morris, & Elias, 1999). We administered cabergoline
in an attempt to manipulate performance on two paradigms traditionally used to study the
neural correlates of conscious perception: backward masking and the attentional blink task. In
addition, there is increasing evidence that dopamine plays a crucial role in determining the
influence of sensory information in relation to expectations acquired through past experience
(Cassidy et al., 2018; Friston et al., 2012). To investigate this relationship, we subjected
participants to a probabilistic discrimination task in which the probability of stimulus
occurrence varied across blocks of trials, thereby tapping into the learning capabilities the
basal ganglia is implicated in traditionally (Berke, 2018).

The difficulty with manipulating dopamine is that effects have been found to depend
on baseline dopamine levels in accordance with an inverted-U-shape (Cools & D'Esposito,
2011). This means that dopamine is thought to have an optimal level for task performance,
but that dopamine manipulations may either benefit or worsen performance dependent on an
individual's starting point on the U-curve. Two measures used to estimate baseline dopamine
levels are working memory operation span (OSPAN; Broadway et al., 2018; Cools, Gibbs,
Miyakawa, Jagust, & D'Esposito, 2008) and spontaneous eye-blink rate (sEBR; Cavanagh et
al., 2014; Jongkees & Colzato, 2016). We used these measures to analyze and control for
potentially different effects of cabergoline in relation to baseline dopamine levels.

155
156
Methods
Participants

30 native Dutch-speakers were recruited from the University of Amsterdam subject pool to
complete the experiment (mean age = 22, range 18-29, 25 female). Because our study is the
first to investigate the effects of cabergoline on conscious perception, we did not conduct a
power analysis. Instead, our sample size is on the upper end of sample sizes employed by
previous research in which cognitive and neural effects of cabergoline were reported with 12-
30 participants (Cavanagh et al., 2014; Cohen, Krohn-Grimberghe, Elger, & Weber, 2007;
Frank & O'Reilly, 2006; Nandam et al., 2013; Norbury, Manohar, Rogers, & Husain, 2013;
Yousif et al., 2016; Fallon et al., 2017; Broadway et al., 2018), and in which the behavioral
and event-related potential (ERP) effects were reported which we aimed to manipulate with
cabergoline (12-15 subjects; Del Cul et al., 2007; van Opstal et al., 2014; Slagter et al., 2012).
Four participants experienced adverse reactions during the cabergoline session
(dizziness and nausea) that interfered with their ability to participate in the study. In two
participants, nausea was present to the point of vomiting. One participant dropped out due to
headaches in the placebo session. Completers and non-completers did not differ in baseline
age, BMI, heart rate/blood pressure, or baseline dopamine proxies (see below). It should be
noted however that all drop-outs were women.
In total, 124 individuals were considered for inclusion in the study. 113 were
interviewed over the telephone to ensure normal or corrected-to-normal vision, no history of
neurological, psychiatric, or any other relevant medical problems, and abstinence from
psychoactive medication. The use of hormonal contraceptives served as an additional
inclusion criterium for female participants, who were tested outside of their period due to
variability in D2R availability during different phases of the menstrual cycle (Czoty et al.,
2009). 63 individuals were eligible for participation, out of which 35 participants were
available to schedule the required three lab visits (see Procedure). 5 participants did not meet
requirements for inclusion based on the first lab visit. After excluding 5 participants with
adverse reactions, the final sample consisted of 25 participants (mean age = 22, range 18-29,
20 female). In exchange for their participation, participants received 10 euros an hour, with a
minimum of 110 euros in total. The study protocol was approved by the medical ethical
committee of the Academic Medical Centre, Amsterdam (currently Amsterdam University
Medical Centers). All participants provided written informed consent in accordance with the
Declaration of Helsinki.
190
Procedure
Participants came to the lab three times on different days (see Table 1). Once for screening
(duration: 2.5h), and twice for an experimental session in which either placebo or 1.5mg of
cabergoline was administered orally (duration: 4.5h each), as part of a double-blind crossover
design. Previous studies found cognitive and neural effects of cabergoline using a dosage of
1-1.5mg (Cavanagh et al., 2014; Cohen et al., 2007; Frank & O'Reilly, 2006; Nandam et al.,
2013; Norbury et al., 2013; Yousif et al., 2016; Fallon et al., 2017; Broadway et al., 2018).
As with our sample size, we chose a dosage of 1.5mg to be on the upper end of previously
employed dosages. There was at least a day in between screening and the first session, and at
least a week between both sessions.
Screening. The first lab visit took place anywhere between 09:00 and 17:30. After
providing written informed consent, participants answered a series of questions concerning
potential medical conditions. Next, we conducted the M.I.N.I.; a structured screening
interview for DSM-IV axis-I disorders (Sheehan et al., 1998). We subsequently measured
participant's weight, height, BMI, blood pressure (BP), and heart rate. Participants were
included in the experiment only if these measures fell within pre-established bounds (BMI
18-30, diastolic BP < 50 or > 90 mmHg, systolic BP < 95 or > 140 mmHg). Next, six
external electrodes were attached to the participant's face and ears in order to measure

spontaneous eye-blink rate (sEBR) at rest. Finally, participants completed an operation span
(OSPAN) working-memory task (Unsworth, Heitz, Schrock, & Engle, 2005), and a titration
procedure for two behavioral tasks to be completed during both experimental sessions
(backward masking and probabilistic discrimination; see below).
Session. All placebo and cabergoline sessions (the second and third visit) took place
between 08:30 and 14:00. Participants were instructed to abstain from drug and heavy
alcohol use, the day before and during the day of the session. Also, participants were
instructed to abstain from caffeine and nicotine the morning of the session. Compliance to the
instructions was checked by the examiner on arrival, in case of non-compliance the session
was postponed. Female participants completed a midstream pregnancy test. Breakfast was
offered, in order to avoid cabergoline-intake on an empty stomach. Blood pressure and heart-
rate were measured using an Omron® M3 comfort Sphygmomanometer, and participants
filled in a visual analogue scale (VAS, see below) three times during the session: on arrival,
at around 1.5h after placebo or cabergoline intake, and at the end of the session. After the
initial blood pressure/heart-rate/VAS measurement, participants were administered either
placebo or cabergoline in a double-blind fashion (order randomized across participants).
After a 40-minute break, a BioSemi ActiveTwo system (BioSemi Inc., Amsterdam, The
Netherlands) EEG cap and electrodes were fitted. Drug plasma levels have been found to
reach maximum concentration after approximately 1.5-3h (Persiani, Rocchetti, Pacciarini,
Holt, Toon, & Strolin-Benedetti, 1996; Agúndez, Garcia-Martin, Alonso-Navarro, &
Jiménez-Jiménez, 2013). Approximately 1 hour and 20 minutes after drug intake, participants
completed 6 minutes of sEBR recordings, followed by the backward masking task around
1.5h after drug intake (see below), during which EEG was recorded. After this task, the EEG
setup was removed from the participant's head. After a 30-minute lunch, participants
proceeded to the attentional blink task, a simple reaction time task, and the probabilistic
discrimination task (see below). At the end of the experiment, one final blood pressure/heart
rate and VAS measure was undertaken. At the end of the final session, participants were
asked to indicate in which session they believed they had received cabergoline.
237

Table 1. *Experimental procedures for the screening, placebo, and cabergoline sessions.*

Screening	Sessions (placebo/cabergoline)
Medical questionnaire	Questions regarding recent substance use + pregnancy test
M.I.N.I.	Blood pressure/heart rate + VAS 1
Body weight/height	Drug intake
Blood pressure/heart rate	40-minute break
sEBR	EEG setup
OSPAN	sEBR
Titration backward masking	Blood pressure/heart rate + VAS 2
Titration probabilistic discrimination	Backward masking (± 1.5h after drug intake)
	30-minute lunch
	Attentional blink
	Reaction time task
	Probabilistic discrimination
	Blood pressure/heart rate + VAS 3 (± 3.5h after drug intake)

238
Physiological and subjective state measures
Heart rate and blood pressure. Physiological measurements were taken once during
screening, and three times during both sessions; namely, on arrival, at around 1.5h after drug

intake, and on completion of testing (± 3.5h after drug intake) (see Table 1). These
measurements were obtained using an Omron® M3 comfort Sphygmomanometer.

Subjective self-report. A set of sixteen VAS measures were used (Bond & Lader,
1974), to assess the subjective state of the subject before medication intake, at around 1.5h
after drug intake, and on completion of testing (± 3.5h after drug intake). Each scale
consisted of a 100-mm horizontal line, anchored by contrasting states of mind (e.g., happy
versus sad). Subjects were asked to regard each line as a continuum and to rate their feelings
at the time by moving a vertical slider across each line. The scales could then be scored by
measuring the length in millimeters from the positive end of each line to the subject's marked
location. These sixteen VAS measures were summarized as three categories: contentedness,
calmness, and alertness (Bond & Lader, 1974).

Baseline dopamine proxies. Both sEBR and OSPAN are widely-used measures that
have been related to baseline dopamine levels (Cools & D'Esposito, 2011; Jongkees &
Colzato, 2016). Both measures have been used in combination with cabergoline in order to
account for individual differences in baseline dopamine (Broadway et al., 2018; Cavanagh et
al., 2014). Eye blink rate is defined as the number of spontaneous eye blinks per minute. The
measure has high test-retest reliability (Kruis, Slagter, Bachhuber, Davidson, & Lutz, 2016)
and is an often-used biomarker of baseline dopamine D2 receptor functioning (Jongkees &
Colzato, 2016; Karson, 1983; Taylor et al., 1999, but see Sescousse et al., 2018). Subjects
were asked to look at a central fixation cross on a computer screen in a relaxed state for 6
minutes while we measured eye activity from a set of vertical and horizontal electrodes, in
order to detect eye blinks. This procedure was employed during all three lab visits.

Eye blinks were established in two ways. First, through a fully automatic procedure
implemented in the python module MNE (*create_eog_epochs*; Gramfort et al., 2013).
Second, eye-blinks were established through a custom semi-automatic procedure using
EEGLAB for MATLAB (Delorme & Makeig, 2004). If mean sEBR per minute differed more
than 3 blinks between both methods, the semi-automatic procedure was repeated, and an
average was taken of both semi-automatic attempts as the final value. Prior to any repetition
of the semi-automatic method, correlations between the automatic and semi-automatic
method exceeded .95 for measurements during all three lab visits.

OSPAN is a working-memory task with high test-retest reliability (Unsworth et al.,
2005), in which participants are instructed to remember letters, while solving simple
arithmetic problems in between letter presentation (Unsworth et al., 2005). Sets of 3-7 letters
were presented successively at fixation. The OSPAN score was calculated through partial
credit scoring, so that each correctly recalled letter in the appropriate location was counted as
correct, regardless of whether the entire sequence was recalled correctly or not. Scores could
range from zero to 75. OSPAN was measured only during screening.

Reaction time. To assess the effects of cabergoline on alertness we administered a
40-trial simple reaction time (RT) task (Brown et al., 2016). In this task, participants had to
respond as quickly as possible by pressing the spacebar whenever a white circle (subtending
approximately 3.1° of visual angle) appeared at the center of the computer screen against a
black background. Stimulus onset asynchrony was jittered between 500 and 1250 ms, with a
mean of 1000 ms. This task lasted less than 2 minutes.

285
286

287 **Main experimental paradigms**
288 All stimuli were presented on an ASUS VG236H 23-inch LCD screen (refresh rate = 100 Hz,
289 resolution 1920x1080). Participants viewed the screen at a distance of 80 cm.

291
292   **Fig. 1** Experimental procedure for one trial of the backward masking task (adapted from Van Opstal
293   et al., 2014)
294

Backward masking. In the backward masking task, adapted from Van Opstal et al.
(2014), participants had to indicate whether briefly presented masked digits (1, 4, 6, or 9)
were smaller or larger than 5 and rate the confidence in their response (Figure 1). Each trial
started with the presentation of a central fixation cross (30 point Courier New), which
increased in size (106 point Courier New, 150 ms duration), cueing the impending target. The
target stimulus (30 point Courier New) then appeared for 10 ms at one of two positions
centered at the vertical midline (top or bottom, 2.29° from fixation). Both stimulus locations
were equally probable. A mask followed the target (200 ms duration) at a variable stimulus
onset asynchrony (SOA). Due to the employed refresh rate of 100Hz, the SOA could vary
from 10 ms to 100 ms in 10 ms steps. By making the delay between cue and target dependent
on SOA, the delay between cue and mask was held constant at 800 ms. The mask (30 point
Courier New) was composed of two letters "E" and two letters "M", tightly surrounding the
target location without superimposing or touching it. All stimuli were black and presented on
a white background, using the Psychophysics toolbox for MATLAB (Brainard, 1997). The
central fixation cross was visible throughout the experiment.
Participants were instructed to indicate by button press whether the presented digit
was smaller or larger than 5, while simultaneously indicating the confidence in their response
(sure/unsure); resulting in four possible responses (<5 sure, <5 unsure, >5 unsure, >5 sure). In
previous research it was found that the D2 agonist pergolide affected response confidence
(Lou et al., 2011). Responses were given by means of a response box attached to the arm
rests of the participant's chair. Response buttons were counterbalanced across participants,
who were instructed to guess one of two "unsure" buttons if they did not see the target.
317   If the participant's reaction time exceeded 1 second, a message was presented
indicating that their response was too slow for the duration of 1 second, urging a faster
response. An individual threshold for awareness was established during the screening session
(see above), by fitting a logistic model (threshold defined as SOA corresponding to 75%
accuracy; mean threshold = 52.93 ms, min = 31 ms, max = 89 ms, sd = 14.38; Del Cul,
Dehaene, & Leboyer, 2006; Van Opstal et al., 2014). This model was fitted on the basis of
176 trials during screening, where each of 11 SOA durations (from 0-100 ms) was presented
16 times.
Prior to the experiment, participants first completed a practice block (176 trials in
screening, 88 trials during placebo and cabergoline sessions). In both drug sessions,
participants completed 920 trials in total, split by seven possible SOAs between target and
mask: 200 mask-only trials (0 ms SOA), 200 trials each for the main SOAs (10 ms /
awareness threshold / 100 ms), 40 trials surrounding the threshold (threshold minus 10 ms
and threshold plus 10 ms), and 40 trials with a 70 ms SOA. The trials in between individual
thresholds and 100 ms were excluded from analysis (threshold + 10ms and 70 ms), because
some participants arrived at an individual threshold at or above 70 ms. This meant that trials

with a SOA at threshold + 10 ms and 70 ms would fall below or above participant's
individual threshold, depending on the participant. 80 trials per participant were discarded for
this reason, leaving 840 trials.
336

337
338 **Fig. 2** Experimental procedure for one trial of the attentional blink task (adapted from Slagter et al.,
339 2017)
340

Attentional blink. Participants also performed a standard AB task in which they had
to identify two digits (T1 and T2) presented in a rapid stream of centrally presented
distractors (letters and symbols; adapted from Slagter et al., 2012; Slagter et al., 2017; see
Figure 2). T2 followed T1 either in the time window of the AB, after 200 ms (short-interval
trial), or outside the time window of the AB, after 800 ms (long-interval trial). Each trial
started with a central fixation cross (1500 ms), after which the stimulus stream began,
consisting of 22 stimuli. Stimuli were presented on a black background (RGB 70, 70, 70) at
the center of the screen (28 point Arial; 0.85° visual angle) for 50 ms, followed by a 50 ms
blank. Digits were drawn randomly (without replacement) from the set 2–9. Distractors were
randomly drawn (without replacement) from the following set of 30 letters and symbols: W,
E, R, T, Y, U, P, A, D, F, G, H, J, K, L, Z, X, C, V, B, N, M, @, #, $, %, }, &, <, and =.
Participants were asked to indicate sequentially the identity of the targets they saw, using the
numpad on a standard keyboard. If they missed a target, they were instructed to guess.
Stimulus presentation was performed using Presentation (Neurobehavioural Systems).
In both sessions, participants first completed a short practice block (20 trials), in
which the first 8 trials moved at half speed. Next, participants moved on to the main
experiment (222 trials), spread over 6 blocks consisting of 37 trials each.
Probabilistic discrimination. In the probabilistic discrimination task, adapted from
Bauer et al. (2016, September), participants were presented continually with a central fixation
cross (28 point Arial, RGB 0, 0, 0), on top of which an image (6.68° visual angle) of either a
face or house was presented for 120 ms, against a grey background (RGB 128, 128, 128).
Face stimuli were created on the basis of the Park Aging Mind Laboratory, University of
Texas at Dallas (Minear & Park, 2004), while house stimuli were based on the Caltech
University Computational Vision database (http://vision.caltech.edu/archive.html). On each
trial, participants had to report the category of the image with by pressing "Q" or "P" on a
standard keyboard. Responses were counterbalanced across participants, who were instructed
to emphasize accuracy over and above speed. The maximum response interval was 1700 ms,
after which the next stimulus was presented regardless of whether a response was given. The
inter-trial interval was jittered and varied from 800 to 1200 ms.
The difficulty of stimulus discrimination was manipulated in terms of stimulus
coherence (Figure 3). Stimuli from the above-mentioned databases were cropped to the
outlines of faces and houses and a 2-dimensional spatial Fourier transform (on luminance
values for x-/y-coordinates) was calculated. The amplitude (power-) spectra of all 82 face and
house images were averaged and subsequently applied to all individual images, such that all

images (faces and houses of all coherence levels) had an identical power spectrum. In other
words, none of them differed in global contrast or luminance. The phase-spectra of each
individual image (that therefore provided all pictorial information) was retained and was
subsequently superimposed with various levels of (uniform) random noise for each image
(Bauer et al., 2016, September). To account for bias in the circular phase-distribution of
superimposed noise and signal phase-spectra, noise-spectra were sampled following previous
suggestions (Dakin, Hess, Ledgeway, & Achtman, 2002).
Titration consisted of two subsequent procedures, in order to establish three difficulty
levels for each individual participant. First, participants completed 300 trials, spread over 10
staircase blocks, in order to establish difficulty levels corresponding to an accuracy of 75%
for each stimulus category separately. During this 3-up-1-down staircase procedure,
participants received a green thumbs-up (RGB R [56 154 79], border RGB [17 79 22]) or red
thumbs-down (RGB [83 2 5], border RGB [251 84 84]) at the center of the screen as
feedback after each trial (500 ms duration).
389

Fig. 3 Probabilistic discrimination stimuli (adapted from Bauer et al., 2016, September)

Next, a total of 810 trials followed, across 27 blocks, in order to extrapolate the
acquired difficulty level to three difficulty levels. For the second part of the titration
procedure, participants received feedback in the break in between blocks; in order to
counteract the development of a bias for one of two response categories. Difficulty levels
were estimated using the method of constant stimuli (MOCS; Bauer et al., 2016, September).
The psychometric functions obtained through this procedure were used to estimate difficulty
(coherence) levels corresponding to 70, 82, and 95% accuracy.
In both the placebo and cabergoline session, the same three difficulty levels were
employed that were acquired from the titration procedure during screening. Unbeknownst to
participants, the prior probability of each category was manipulated in a block-wise manner
(20/35/50/65/80%), spread over 25 blocks of 40 trials each, for a total of 1000 trials per
session. As such, we manipulated perceptual information (difficulty) and stimulus prior
probability independently of one another. Stimulus presentation for the titration procedure
during the screening was performed using the Psychophysics toolbox for MATLAB
(Brainard, 1997), and Presentation was used to present the task in both experimental sessions
(Neurobehavioural Systems).
409

Behavioral analyses

Physiological and subjective state measures. In order to test whether physiology
and subjective state changed over the course of the experiment, and whether these measures
were influenced by cabergoline, we conducted a repeated-measures analysis-of-variance (RM
ANOVA) for heart rate, diastolic and systolic blood pressure, and each VAS category

separately; across all three time-points and both sessions. We conducted a paired-samples t-
test between sessions for our simple RT task, as an additional measure to assess alertness. In
order to test whether cabergoline exerted influence on sEBR, we conducted a paired-samples
t-test between sEBR under placebo versus cabergoline. Kendall's Tau correlation was
employed to establish the relationship between sEBR sessions, as well as the relationship
between sEBR and OSPAN, as this coefficient is more robust in the case of small samples
and tied ranks (Bonett & Wright, 2000). Correlations were Bonferroni corrected for multiple
comparisons.
Backward masking. The dependent measures in the backward masking task were
accuracy (0/1) and confidence (unsure/sure). For both of these measures, we computed a 2
(Drug; placebo/cabergoline) x 5 (SOA; mask-only/10 ms/threshold – 10 ms/threshold/100
ms) RM ANOVA. Furthermore, we performed an additional analysis including screening
sEBR and OSPAN as covariates in both of these analyses, as we predicted cabergoline effects
may depend on individual baseline dopamine levels, based on previous reports (Broadway et
al., 2018; Cavanagh et al., 2014; Cools & D'Esposito, 2011; Jongkees & Colzato, 2016). We
repeated these analyses for the two-alternative forced choice version of the signal detection
theory parameters d' (Green & Swets, 1966) and meta-d' (Maniscalco & Lau, 2012), for the
three primary SOAs (10 ms, threshold, and 100 ms).
One participant confused the confidence response buttons in one session. We reversed
these confidence scores manually. One participant experienced side effects only near the end
of the last session. This participant thus completed the backward masking task twice without
knowledge about drug condition. In order to maximize statistical power, this participant was
included in all analyses concerning the backward masking task (including EEG), but not in
the analyses of other tasks. It did not matter for our results whether this participant was
included in the backward masking task analyses or not.
Attentional Blink. The dependent measures for the AB task were T1 accuracy and
T2 | T1 accuracy. In other words, T2 accuracy was based only on those trials where T1 was
correctly reported. For each of these measures, we computed a 2 (Drug; placebo/cabergoline)
x 2 (Lag; 2/8) RM ANOVA. For this task as well, we computed additional analyses in order
to include sEBR and OSPAN as covariates. Finally, we computed AB size, in order to
investigate the relationship between AB size and our baseline dopamine measures as some
(Colzato, Slagter, Spapé, & Hommel, 2008) but not other studies (Slagter et al., 2012) have
found.
Probabilistic discrimination. In the case of the probabilistic discrimination task, our
dependent measure of interest was accuracy. As such, we computed a 2 (Drug;
placebo/cabergoline) x 3 (Difficulty; easy/medium/hard) x 5 (Probability; .2/.35/.5/.65/.8)
RM ANOVA. For this analysis as well, we computed additional models including screening
sEBR and OSPAN as covariates.
Our titration procedure was not successful for all participants. As a result, a number
of participants ended up with only two difficulty levels for one out of two stimuli. For this
reason, we repeated the above analysis for both the group with all difficulty levels (N = 16),
and participants who had either two or three difficulty levels as a result of the titration
procedure (N = 24). In this latter analysis, we excluded the medium difficulty trials. One
participant was excluded from both analyses, because this participant ended up with only one
difficulty level for face stimuli.
All covariates in the above-mentioned RM ANOVAs were centered (van Breukelen &
van Dijk, 2007). For all repeated-measures ANOVA analyses, whenever Mauchly's test
suggested a violation of sphericity, we report Geenhouse–Geisser corrected *P*-values, but
uncorrected degrees of freedom. In order to test for order effects, we repeated each of the RM

ANOVAs for our behavioral paradigms including a between-subject factor indicating
whether a participant received either placebo or cabergoline in the first session.
466   **Bayesian statistics.** In order to evaluate evidence in favor of our (null) hypotheses,
we conducted Bayesian statistics. For each reported frequentist test, we report the Bayes
factor corresponding to the inclusion of a factor or interaction within the model in question
(shortened to BF_{incl}), compared to equivalent models stripped of the effect. For example,
$BF_{incl} = 10$ indicates that a model including the factor in question is ten times more likely
given the data compared to a model without the variable. Conversely, $BF_{incl} = .1$ indicates that
a model without said effect is ten times more likely given the data. All Bayesian statistics
were conducted using JASP (2019, version 0.10.0).
All data visualization was performed with the help of raincloud plots (Allen, Poggiali,
Whitaker, Marshall, & Kievit, 2019), which include the mean, individual data points, as well
as the overall distribution of the measure in question.
477
478   **EEG**
479   **Recording and preprocessing.** EEG data, digitized at 512 Hz, were continuously
recorded in both the placebo and cabergoline session during 6 minutes of sEBR and the
backward masking task, using an ActiveTwo system (BioSemi, Amsterdam, the
Netherlands), from 64 scalp electrodes placed according to the 10/20 system, four electro-
oculographic electrodes placed above and below, and to the side of the eyes, and two external
electrodes attached to each earlobe. EEG data were offline referenced to the average activity
recorded at the earlobes, and high-pass 'firws' filtered (default settings) at 0.05 Hz using a
Kaiser window, following previous suggestions (Widmann, Schröger, & Maess, 2015). The
continuous data were subsequently epoched from −1.5 to 1.5 s around stimulus presentation
and baseline corrected to the average activity between −200 ms and 0 ms pre-stimulus.
Epochs containing EMG artifacts or eye blinks surrounding stimulus presentation were
rejected based on visual inspection. Extremely noisy or broken channels were interpolated.
Remaining eye blink artifacts were removed by decomposing the EEG data into independent
sources of brain activity using an Independent Component Analysis, and removing eye blink
components from the data for each subject individually. Epochs were low-pass filtered at 30
Hz for visualization purposes only. Preprocessing was done using the Fieldtrip toolbox
(Oostenveld, Fries, Maris, & Schoffelen, 2011) for Matlab (The MathWorks, Inc. Natick,
MA, USA) using custom-written Matlab scripts.
Analyses. To determine the effect of our manipulations on ERP markers of
information-processing, we examined the effects of SOA and Drug on the amplitude of the
visual-evoked P1 and N1 components, the N2, as well as of the later P3b (Del Cul, Baillet, &
Dehaene, 2007). In line with Del Cul et al (2007), we epoched the ERP data to the onset of
the mask. Next, we subtracted the data from the mask-only SOA condition from all other
SOA conditions. Finally, we shifted ERP onset back to target onset, in order to compute
target-locked ERPs. Visual inspection of the grand- and condition-average ERPs showed that
the P1 and N1 components peaked over lateral occipitoparietal scalp sites (PO7, PO3, O1,
PO4, PO8, O2), the N2 over centroparietal scalp regions (C1, Cz, C2, CP1, CPz, CP2), and
the P3b over central parietal scalp sites (P1, Pz, P2, PO3, POz, PO4). These scalp sites were
used to determine the peak amplitude and latency of these components for each condition of
interest. Specifically, the largest positive voltage value between 75–150 ms post-target, and
the largest voltage negativity within 150–225 ms were selected to determine the amplitude
and latency of the P1 and N1 peaks, respectively, for each subject separately. In the case of
the P2 and N2, these intervals were 175-250 ms, and 250-375 ms, respectively. For the P3, an
interval of 300-450 ms was used. All average amplitude values 15 ms around the peak
sample, as well as individual latencies were entered into separate RM ANOVAs with two

within-subject factors: SOA (10 ms, threshold-10 ms, threshold, 100 ms), and Drug
(placebo/cabergoline). Because the mask-only condition is used to acquire ERP data for the
remaining four SOA conditions (see above), the SOA factor contains four instead of five
levels for ERP analyses.
518
519
520   **Results**
Physiological and subjective state measures
First, we aimed to establish whether cabergoline exerted physiological effects (Figure 4).
Heart rate decreased over time in both placebo and cabergoline conditions ($F_{(2, 48)} = 11.8$, $p <$
$.001$, $\eta^2 = .06$, $BF_{incl} > 100$), and was overall higher in the cabergoline condition ($F_{(1, 24)} = 9$,
$p = .006$, $\eta^2 = .03$, $BF_{incl} = 57.3$), but there was no interaction between Time and Drug ($F_{(2, 48)}$
< 1, $p = .63$, $BF_{incl} = .14$; see Figure 4a). In the case of diastolic blood pressure as well, we
also observed a decrease over time ($F_{(2, 48)} = 21.7$, $p < .001$, $\eta^2 = .14$, $BF_{incl} > 100$), but in this
case an overall lower measurement in the cabergoline condition ($F_{(1, 24)} = 9.9$, $p = .004$, $\eta^2 =$
$.03$, $BF_{incl} = 12.4$), and again no interaction between Time and Drug ($F_{(2, 48)} < 1$, $p = .42$,
$BF_{incl} = .2$). With regards to systolic blood pressure, we also observed a decrease in blood
pressure over time ($F_{(2, 48)} = 36.7$, $p < .001$, $\eta^2 = .22$, $BF_{incl} > 100$) and an overall lower
measurement in the cabergoline condition ($F_{(1, 24)} = 9$, $p = .006$, $\eta^2 = .03$, $BF_{incl} = 31.2$), but
here a significant interaction between Time and Drug was found ($F_{(2, 48)} = 7.2$, $p = .002$,
$BF_{incl} = 5.8$; see Figure 4b). Thus, all three physiological measures decreased over time.
While heart rate was higher, and diastolic blood pressure lower across the cabergoline
session, cabergoline can only be said to have decreased systolic blood pressure, as indicated
by the interaction between Time and Drug.
538

539
540   **Fig. 4 Time course and difference between drug conditions in heart rate and blood pressure.** All
541   three physiological measures decreased over time. Cabergoline only affected systolic blood pressure,
542   while heart rate was higher, and diastolic pressure was lower in the cabergoline versus placebo
543   session
544
545   Cabergoline also affected the subjective state of participants over the course of the
experiment, as indicated by the VAS (visual analog scale, see Figure 5). On the calmness
scale, there was no main effect of Drug ($F_{(1, 24)} < 1$, $p = .63$, $BF_{incl} = .23$), but calmness
increased over the course of the experiment ($F_{(2, 48)} = 14.5$, $p < .001$, $\eta^2 = .04$, $BF_{incl} = 31$). In
addition, we found an interaction between Drug and Time ($F_{(2, 48)} = 5.8$, $p = .005$, $\eta^2 = .02$,
$BF_{incl} = 2$), reflecting the fact that calmness ratings increased less in the cabergoline condition
(Figure 5). In the case of contentedness, there was no main effect of Drug ($F_{(1, 24)} < 1$, $p =$
$.46$, $BF_{incl} = .3$), Time ($F_{(2, 48)} = 2.7$, $p = .08$, $BF_{incl} = .33$), or an interaction between Drug and

Time ($F_{(2, 48)} < 1$,　p = .53, BF_{incl} = .16; see Figure 5). Finally, alertness decreased over the
course of the experiment ($F_{(2, 48)}$ = 10.3,　p = .001, η^2 = .12, BF_{incl} > 100) and was generally
lower in the cabergoline session compared to placebo session ($F_{(1, 24)}$ = 6.3,　p = .02, η^2 = .01,
BF_{incl} = 1.1), but there was no interaction ($F_{(2, 48)}$ = 1.8,　p = .18, BF_{incl} = .21; see Figure 5).
Thus, cabergoline decreased self-reported ratings of calmness and alertness as the experiment
progressed.
559

560
561　**Fig. 5 Time course and difference between drug conditions in VAS scores for calmness,**
562　**contentedness, and alertness.** Participants become calmer and less alert over the course of the
563　experiment. Cabergoline stifled this increase in calmness
564
85% of participants successfully guessed when they received cabergoline at the end of the
experiment. However, despite these effects, we found no significant difference in sEBR
(spontaneous eye blink rate) between placebo (M = 14 blinks per minute, SD = 8.9) and
cabergoline (M = 15.3, SD = 10.6) ($t_{(24)}$ = -1.2,　p = .23, BF_{01} = .41). In addition, we were
unable to replicate the finding by Cavanagh and colleagues (2014) that baseline sEBR
conditioned a cabergoline-induced shift in line with an inverted-U-shape pattern, to the point
where high baseline sEBR (as indicative of high tonic striatal dopamine) is associated with a
reduction in blink rate, whereas low baseline sEBR is linked to an increase in blink rate: we
found that the difference in sEBR between drug conditions was unrelated to screening sEBR
(M = 13.2, SD = 7.8) ($\tau_{(25)}$ = .23, p = .11).
Replicating previous findings (Jongkees & Colzato, 2016; Kruis et al., 2016), sEBR
was correlated across sessions: between the placebo and cabergoline session ($\tau_{(25)}$ = .63, p <
.001), between the screening and cabergoline session ($\tau_{(25)}$ = .37, p = .01), as well as between
the screening and placebo session ($\tau_{(25)}$ = .32, p = .03; see Figure 6a).　The latter two
correlations did not survive a Bonferroni correction for multiple comparisons using a
corrected alpha level of .05 / 6 = .0083, based on all correlations computed in this section.
Together these results lend support to the robustness of the measure when measured at the
same time of day (Barbato et al., 2000; Jongkees & Colzato, 2016). However, despite the
proposed relation between sEBR and OSPAN as a biomarker of striatal dopamine, we found
no relationship between these measures in either the cabergoline ($\tau_{(25)}$ = .06, p = .67) or
placebo session ($\tau_{(25)}$ = .06, p = .67). This relationship was absent even when these measures
were collected in the same session; namely, during screening ($\tau_{(25)}$ = -.11, p = .47; see Figure
6b). Finally, alertness, as indicated by our simple RT task also did not differ between the
placebo (M = 224 ms, SD = 15.8) and cabergoline condition (M = 225 ms, SD = 15.1) ($t_{(25)}$ =
-.54,　p = .59).
Thus, cabergoline did not affect sEBR or our objective measure of alertness.
591

592
593 **Fig. 6 The relationship between sEBR measures among themselves and in relation to OSPAN.**
594 All sEBR measures were found to be positively correlated (**a**), but we found no relationship between
595 sEBR and OSPAN (**b**)
596

Main experimental paradigms

597
We next examined potential effects of cabergoline on our main experimental measures of
interest: target identification accuracy and processing in the backward masking task,
attentional blink size in the attentional blink task, and discrimination accuracy under varying
conditions of difficulty and probability in the probabilistic discrimination task. To
foreshadow our results, cabergoline did not affect any of the key behavioral findings
associated with these tasks, whether sEBR or OSPAN were included as covariates in the
analyses or not. Neither did it affect neural processing of the target in the backward masking
task, as shown by ERP analyses. Yet, importantly, we did replicate all standard findings
typically obtained with these tasks (e.g., effects of masking on target-evoked ERPs, the
attentional blink).

Backward masking

609
Behavior. As is typically observed (e.g., Breitmeyer, 2007) and shown in Figure 7,
targets were more often identified correctly in the backward masking task as the delay
between target and mask (SOA) increased ($F_{(4, 100)} = 187.4$, $p < .001$, $\eta^2 = .79$, $BF_{incl} > 100$).
Yet, in contrast to our main prediction that cabergoline would affect participant's ability to
detect targets, there was no interaction between Drug and SOA ($F_{(4, 100)} < 1$, $p = .56$; $BF_{incl} =$
.03; see Figure 7). We also did not find an overall difference between placebo and
cabergoline on target identification accuracy ($F_{(1, 25)} < 1$, $p = .85$, $BF_{incl} = .13$). Controlling
for screening sEBR, OSPAN, or the combination of both did not change the results. Neither
did the inclusion of a between-subject factor for drug order. We repeated these analyses with
d' as the dependent variable, but results were equivalent.
Similarly, in the case of confidence scores, cabergoline did not affect reported
confidence ratings, as indicated by the lack of an interaction between SOA and Drug ($F_{(4, 100)}$
$= 1.6$, $p = .21$; $BF_{incl} = .06$; see Figure 7). We also found no overall difference between drug
conditions ($F_{(1, 25)} < 1$, $p = .63$; $BF_{incl} = .14$), but participants did report an improvement in
response confidence as SOA increased ($F_{(4, 100)} = 204.8$, $p < .001$, $\eta^2 = .8$; $BF_{incl} > 100$).
Again, correcting for baseline dopamine measures or drug order had no impact on the results.
As with d', repeating these analyses with meta-d' as the dependent variable made no
difference. Thus, we replicated a similar pattern of results with regards to both objective and

subjective aspects of the participant's response during backward masking (Del Cul et al.,
2007), but these patterns were not affected by cabergoline.
630

631
632 **Fig. 7 Mean accuracy and confidence scores in the backward masking task across five possible**
633 **SOAs and both drug conditions.** While both accuracy (**a**) and confidence (**b**) scaled with the
634 duration of SOA, cabergoline had no effect on either of these measures
635
EEG. In addition to these behavioral masking effects, we replicated previous reports
that target-evoked ERP components scale with the duration of target-mask SOA (Del Cul et
al., 2007). We found a strong effect of SOA on the amplitude and peak-latency of the target-
evoked P1, N1, N2, and P3b components. However, in the case of our ERP results as well,
each of these measures was unaffected by cabergoline (see Table 2 and Figure 8). In order to
remain close to previous studies (Del Cul et al., 2007), we repeated these analyses with an
average reference instead of an earlobe reference, but the conclusions belonging to these
results remained the same. Thus, cabergoline did not affect the threshold for conscious
perception by modulating neural target processing.
645

Table 2. *Summary of the statistical analyses performed on ERP data.*

Component	Measure	SOA ($F_{(3, 75)} =$)	Drug ($F_{(1, 25)} =$)	SOA*Drug ($F_{(3, 75)} =$)
P1	Amplitude	51.7, p < .001*	.01, p = .91, BF_{incl} = .16	.62, p = .60, BF_{incl} = .08
	Latency	8.2, p < .001*	.17, p = .68, BF_{incl} = .16	.48, p = .7, BF_{incl} = .08
N1	Amplitude	12.4, p < .001*	.04, p = .85, BF_{incl} = .15	.44, p = .73, BF_{incl} = .06
	Latency	66.8, p < .001*	.33, p = .57, BF_{incl} = .19	.07, p = .98, BF_{incl} = .06
N2	Amplitude	4.7, p = .005*	.04, p = .83, BF_{incl} = .15	.21, p = .89, BF_{incl} = .06
	Latency	99.3, p < .001*	2.9, p = .1, BF_{incl} = .38	.09, p = .97, BF_{incl} = .06
P3b	Amplitude	57.2, p < .001*	.02, p = .9, BF_{incl} = .15	.9, p = .44, BF_{incl} = .07
	Latency	10, p < .001*	.1, p = .74, BF_{incl} = .15	1.7, p = .18, BF_{incl} = .09

* = BFincl > 100

646
647

648
649   **Fig. 8** Effects of cabergoline on target-evoked ERPs in the backward masking task. This figure
650   displays the grand-average target-evoked ERPs for P3b electrodes (P1, Pz, P2, PO3, POz, PO4) for
651   both drug conditions combined (**a**) and separately (**b**), per target-mask SOA condition (10, 40, 50, 60,
652   70, and 100 ms). This figure shows that while ERP amplitudes and latencies generally increased as a
653   function of target-mask SOA, these measures were unaffected by cabergoline.

654
655

Attentional blink
As expected and shown in Figure 9, we found a robust attentional blink: T2|T1 accuracy was
significantly worse when T2 was presented after one distractor (lag 2) compared to seven
distractors (lag 8; $F_{(1, 24)} = 43.9$, $p < .001$, $\eta^2 = .39$; $BF_{incl} > 100$). T2|T1 accuracy was
marginally better in the cabergoline condition (81.9±11%) compared to placebo (79.7±14%;
$F_{(1, 24)} = 4.2$, $p = .052$; $BF_{incl} = .34$), but we found no interaction between Lag and Drug ($F_{(1,}$
$_{24)} = 1.42$, $p = .25$; $BF_{incl} = .3$).

664
Fig. 9 Mean accuracy for T1 and T2|T1 at lag 2 and lag 8. While we replicated behavioral findings
common to the attentional blink paradigm, cabergoline did not affect the attentional blink

When we controlled for baseline dopamine measures (i.e., baseline OSPAN and
sEBR), the inclusion of OSPAN did not affect the results, and controlling for sEBR also did
not reveal a critical interaction between Lag and Drug ($F_{(1, 22)} = 1.3$, $p = .26$; $BF_{incl} = .3$). We
did find a between-subject effect of screening sEBR ($F_{(1, 22)} = 5.2$, $p = .03$; $BF_{incl} = 2.5$),
which exacerbated the main effect of Drug ($F_{(1, 22)} = 5.4$, $p = .029$; $\eta^2 = .004$; $BF_{incl} = .36$). In
addition, we found an interaction between sEBR and Drug ($F_{(1, 22)} = 5.8$, $p = .025$; $\eta^2 = .005$):
sEBR correlated with overall T2|T1 accuracy in the placebo ($\tau_{(25)} = -.31$, $p = .03$), but slightly
less so in the cabergoline ($\tau_{(25)} = -.26$, $p = .07$) condition.

Although the critical three-way interaction between sEBR, Lag and Drug was not
significant, we also found an interaction between screening sEBR and Lag ($F_{(1, 22)} = 5$, p =
.036; η^2 = .004). This interaction is best understood in terms of the relationship between
sEBR and AB size (Colzato et al., 2008; Slagter et al., 2012); namely, the difference in T2|T1
accuracy between lag 2 and 8. AB size ranged from -2.9% to 59.7% in the placebo condition,
and from 0% to 47.5% in the cabergoline condition. The interaction between sEBR and Lag
stems from a small correlation between screening sEBR and AB size in the cabergoline ($\tau_{(25)}$
= .31, p = .03) and placebo condition ($\tau_{(25)}$ = .28, p = .047). We found no other relationships
between dopamine baseline measures and the AB; the difference in AB size between the
cabergoline and placebo session was unrelated to screening sEBR ($\tau_{(25)}$ = .05, p = .71). For
the cabergoline session, there was no relation between AB size and sEBR ($\tau_{(25)}$ = .13, p =
.35), nor between AB size and OSPAN ($\tau_{(25)}$ = .08, p = .57; see Figure 10a). Similarly, for the
placebo session we did not find a relationship either between AB size and sEBR ($\tau_{(25)}$ = .15, p
= .3), nor OSPAN ($\tau_{(25)}$ = .02, p = .89, see Figure 10b). However, none of the correlations
reported in this section survived a Bonferroni correction for multiple comparisons using a
corrected alpha level of .05 / 9 = .0056, based on all correlations computed in this section.
Thus, we found no strong support for a relationship between sEBR and the AB, or OSPAN
and the AB, nor did controlling for these factors reveal drug-related effects on the AB.
When controlling for drug order, we found an interaction between Drug and drug
Order ($F_{(1, 21)} = 5.7$, p = .027; η^2 = .004; BF_{incl} = .43). When participants received placebo
first, they were less accurate overall in identifying T2|T1 in the placebo condition
(77.5±15%), compared to the cabergoline condition (81.6±11%). For participants who
received cabergoline first, this difference was not present (placebo: 82.1±12%, cabergoline:
82.3±10%). No other order effects were present. Together these results indicate that
cabergoline did not affect the attentional blink.
We found no difference in T1 accuracy between the placebo and cabergoline
condition ($F_{(1, 24)} < 1$, p = .49; BF_{incl} = .27), or an interaction between Drug and Lag ($F_{(1, 24)} <$
1, p = .5; BF_{incl} = .32), but T1 accuracy was worse when T2 was presented after only one
distractor (lag 2; $F_{(1, 24)} = 46.6$, p < .001, η^2 = .16; BF_{incl} > 100). Controlling for sEBR,
OSPAN, or drug order did not affect the results for T1 accuracy. Thus, cabergoline also did
not affect T1 identification.
707

708
709 **Fig. 10 The relationship between AB size and striatal-dopamine proxy measures.** Both for the
710 cabergoline and placebo condition, we found a positive relation between AB size and screening

711 sEBR, while such a relationship was not present neither for sEBR measured within the same session,
712 nor for OSPAN
713
714
Probabilistic discrimination
In our final experimental paradigm, participants were tasked with discriminating between
face and house stimuli that varied in difficulty and probability of occurrence (see Methods).
Cabergoline did not affect discrimination accuracy (specifically: hit-rate; $F_{(1, 23)} = 2.4$, $p =$
.13, $BF_{incl} = .35$). Neither did we find an interaction between Drug and Difficulty ($F_{(1, 23)} < 1$,
$p = .91$, $BF_{incl} = .14$), Drug and Probability ($F_{(4, 92)} < 1$, $p = .78$, $BF_{incl} = .02$), or a three-way
interaction ($F_{(4, 92)} < 1$, $p = .48$, $BF_{incl} = .05$; see Figure 11a). Thus, cabergoline did not
significantly affect perception on this task either. Standard effects observed with this task in
non-drug studies were replicated (Bauer et al., 2016, September): accuracy was lower on
difficult ($F_{(1, 23)} = 208.3$, $p < .001$, $\eta^2 = .60$, $BF_{incl} > 100$), and on low-probability trials ($F_{(4,}$
$_{92)} = 22.49$, $p < .001$, $\eta^2 = .04$, $BF_{incl} > 100$). When we controlled for screening sEBR and
OSPAN, this did not change the results. We did find an interaction between drug Order and
Drug ($F_{(4, 88)} = 5.2$, $p = .03$, $BF_{incl} = 1.3$). Opposite to what we found for the attentional blink
task, when participants received placebo first, they were more accurate overall in the placebo
($84\pm7\%$) compared to the cabergoline ($82.4\pm7\%$) condition. When cabergoline was
administered first, this difference was absent (placebo: $83.5\pm5\%$, cabergoline: $83.1\pm4\%$). We
found no other order effects.
When we repeated these analyses for participants with all three difficulty levels for
both stimuli after titration (N = 16; see Methods), the same pattern of results was obtained.
Cabergoline did not affect discrimination accuracy ($F_{(1, 15)} < 1$, $p = .47$, $BF_{incl} = .69$), and we
found no two-way interaction between Drug and Difficulty ($F_{(2, 30)} < 1$, $p = .59$, $BF_{incl} = .05$),
Drug and Probability ($F_{(4, 60)} < 1$, $p = .76$, $BF_{incl} = .02$), or a three-way interaction ($F_{(8, 120)} =$
1.1, $p = .4$, $BF_{incl} = .04$; see Figure 11b). Accuracy was still lower on difficult ($F_{(2, 30)} =$
233.8, $p < .001$, $\eta^2 = .53$, $BF_{incl} > 100$) and low-probability trials ($F_{(4, 60)} = 21.1$, $p < .001$, η^2
$= .08$, $BF_{incl} > 100$). Controlling for screening sEBR and OSPAN did not change the acquired
results. The interaction between drug Order and Drug was maintained ($F_{(1, 14)} = 9.9$, $p = .007$,
$BF_{incl} > 100$), where overall accuracy was higher in the placebo condition ($83\pm7\%$) when
placebo was administered first compared to the cabergoline condition ($80.6\pm6\%$). This
difference was again absent when cabergoline was administered first (placebo: $82.7\pm3\%$,
cabergoline: $83.1\pm3\%$).
Thus, we found no evidence in support of the conclusion that cabergoline affected any
of the key findings in our perceptual tasks.
747

748
Fig. 11 Cabergoline did not affect probabilistic discrimination performance. This figure shows
mean accuracy for each drug condition and across five stimulus probabilities for all participants with
two (**a**; N = 24) and three (**b**; N = 16) difficulty levels after titration. While accuracy was higher for
easy trials and blocks in which the presented stimulus was more likely to occur, these effects were
unaffected by cabergoline. Medium-difficulty trials take up position in between easy and hard trials in
terms of mean accuracy when included

755
756
<center>**Discussion**</center>
This double blind, placebo-controlled, cross-over study tested the hypothesis that striatal
dopamine is involved in conscious perception by administering the dopamine D2 agonist
cabergoline and placebo to healthy participants. To that end 1) we established an effect of
cabergoline on participant's physiological and subjective state, 2) we tested the effect of
cabergoline on two often-used dopamine proxy measures (sEBR and OSPAN), and 3) we
subjected participants to well-known and often-employed experimental paradigms targeting
the neural correlates of consciousness. While we were able to establish an effect of
cabergoline on participant's physiological and subjective state, we did not find an effect of
cabergoline on sEBR. Crucially, while we replicated key behavioral and ERP findings
associated with the paradigms we employed (Del Cul et al., 2007; Slagter et al., 2017; Van
Opstal et al., 2014), none of these findings were affected by cabergoline. Thus, we did not
obtain evidence that the dopamine D2 agonist cabergoline affected conscious perception,
which was also supported by Bayesian statistics.
Just as with positive results, an observed null result could indicate a true effect (i.e.,
no role for striatal dopamine in conscious perception), or it could be due to uncontrolled or
unknown factors. Based on the convergent evidence discussed in the introduction implicating
the striatum and its irrigation by dopamine in conscious perception (e.g., Bisenius et al.,
2015; Slagter et al., 2017; Van Opstal et al., 2014), we predicted that cabergoline would
affect performance on the backward masking and attentional blink tasks. Yet, our Bayesian
results provided evidence against an effect of 1.5mg cabergoline (all $BF_{incl} < .7$; except for

the interaction between Drug and drug Order for discrimination accuracy in the probabilistic
discrimination task). This absence of an effect is further strengthened by the fact that other
studies using different cognitive tasks have previously reported significant effects of
cabergoline on task performance using a similar or an even lower dose of orally administered
cabergoline (Broadway et al., 2018; Cavanagh et al., 2014; Fallon et al., 2017; Frank &
O'Reilly, 2006; Norbury et al., 2013; Nandam et al., 2013; Yousif et al., 2016). These
observations together argue for an interpretation in terms of a true null result. Nonetheless,
there are a number of practical limitations that pertain to the present study.

For one, blinding was unsuccessful as 85% of participants guessed when they
received cabergoline at the end of the experiment. While all participants with adverse
reactions were excluded from any analyses, we cannot exclude the possibility that
participants' ability to tell when they received cabergoline influenced the results. It is
possible this ability stems from the higher dosage (1.5 mg) we employed compared to
previous studies administering cabergoline (1-1.25 mg; Broadway et al., 2018; Cavanagh et
al., 2014; Frank & O'Reilly, 2006; Norbury et al., 2013; Nandam et al., 2013; Yousif et al.,
2016). While a lower dosage may benefit blinding, it may also further reduce the chance of
finding an effect of cabergoline. Another option for future research would be to administer an
antiemetic such as domperidone in combination with cabergoline, in order to mitigate and
mask physical side effects (Fallon et al., 2017; Norbury et al., 2013).

Second, while we titrated performance on the backward masking and probabilistic
discrimination tasks, we did not do so for the attentional blink task, which may have resulted
in reduced sensitivity of this task to our dopamine manipulation. We chose this specific
version of the task in order to stay as close as possible to a recent study in which we found a
relationship between activity in the ventral striatum and the attentional blink (Slagter et al.,
2017). While we cannot rule out that a titration procedure may have resulted in a task more
sensitive to our manipulation, we believe this is unlikely given that our participants showed a
robust attentional blink, indicating that our task was suitable for the effect we intended to
manipulate. In addition, since cabergoline did not affect performance on the two tasks we did
titrate, we believe it unlikely that this would have been different in the case of the attentional
blink task.

Third, a limitation of our study pertains to the use of alleged indirect measures of
dopamine activity: sEBR (Jongkees & Colzato, 2016) and the OSPAN task (Cools, Gibbs,
Miyakawa, Jagust, & D'Esposito, 2008). The functioning of dopamine relies on a relative
equilibrium at the system level (i.e. frontal versus basal ganglia; Wiecki & Frank, 2013;
Wise, Murray, & Gerfen, 1996), within the basal ganglia itself (i.e., direct versus indirect
pathways; Redgrave et al., 2010), and at the level of synapses (i.e., pre- versus postsynaptic
effects; Frank & O'Reilly, 2006; Usiello et al., 2000). The equilibrium at the system- and
pathway-level is in turn implicated in the differing effects at the receptor level (i.e., D1
versus D2 receptors; Shen, Flajolet, Greengard, & Surmeier, 2008). Due to the complexity
and the multitude of interactions between different systems it is especially important to be
able to report on the initial state of the dopamine-system.

Unfortunately, sEBR was unaffected by cabergoline. While this finding goes against
studies with humans showing effects of dopaminergic drugs on sEBR (Blin, Masson, Azulay,
Fondarai, & Serratrice, 1990; Karson, 1983; Strakowski & Sax, 1998), this finding fits with
several pharmacological studies that reported no effect (Depue, Luciana, Arbisi, Collins, &
Leon, 1994; Ebert, 1996; Mohr, Sándor, Landis, Fathi, & Brugger, 2005; van der Post, de
Waal, de Kam, Cohen, & van Gerven, 2004).

Moreover, differences in OSPAN and sEBR were unpredictive for the effect of
cabergoline on performance in our experimental paradigms. These findings are surprising in
light of previous work showing how effects of cabergoline on cognitive function depend on

individual baseline sEBR (Cavanagh et al., 2014) and OSPAN score (Broadway et al., 2018).
However, it should be emphasized that evidence for a relationship between dopamine-proxy
measures and dopamine levels is often correlational, based on studies with small sample sizes
(N < 50; Cools et al., 2008), and that results are mixed (Dang et al., 2017; Sescousse et al.,
2018). It is debatable whether such small-sample studies have adequate power to provide
evidence for this relationship (Cremers, Wager, & Yarkoni, 2017; Rousselet & Pernet, 2012).
Similarly, the correlational findings we report should also be interpreted cautiously. While
measures such as OSPAN and sEBR are easy to administer, the correlational evidence – as
well as inconsistent findings regarding these measures (Dang et al., 2017; Jongkees &
Colzato, 2016; Sescousse et al., 2018) – put pressure on their validity as an index of baseline
dopamine levels.
Despite these practical limitations, the advantage of our study is that our results are
univocal: the administration of cabergoline had no effect on any of our indices of conscious
perception obtained using often-employed experimental paradigms in the study of
consciousness. Of importance, we did consistently replicate all key behavioral and event-
related potential (ERP) findings associated with each of these tasks.
Theoretically, perhaps our null-findings are best explained by an appeal to predictive
processing approaches to brain function (Hohwy, 2012; 2013; de Lange, Heilbron, & Kok,
2018). From the perspective of these theories, the brain is viewed as a prediction machine
which is continuously revising its predictions about the causes of sensory data. On this view,
top-down signals in perceptual hierarchies carry perceptual predictions, while bottom-up
signals convey perceptual prediction errors. Perception becomes a process of continual
minimization of prediction errors across hierarchical levels, instantiating a process of
approximate Bayesian inference on the causes of sensory signals. In this process, dopamine is
thought to regulate the relative precision of top-down predictions and bottom-up prediction
errors, where precision is understood as inverse variance (Friston et al., 2012): a higher
(expected) variance of sensory signals leads to a smaller influence on updating perceptual
predictions. "If true, this means that modulators of synaptic gain (like dopamine) do not
report perceptual content but the context in which percepts are formed. In other words,
dopamine reports the precision or salience of sensorimotor constructs (representations)
encoded by the activity of the synapses they modulate." (Friston et al., 2012, p. 2).
From this perspective, there may not have been enough uncertainty in our
experimental paradigms for our dopamine manipulation to play a determining role. Indeed,
previous studies employing cabergoline in the context of working memory found an effect
under conditions where task-relevant targets were embedded in a larger perceptual field;
namely, in the presence of distractors. In these studies, cabergoline exerted an effect on task
performance in terms of target-detection accuracy and successful recall when a target stood in
competition with other stimuli (Broadway et al., 2018; Cavanagh et al., 2014; Frank &
O'Reilly, 2006). In the case of each of our tasks, targets were consistently surrounded by
distracting information *temporally, but never spatially.* Perhaps the selective functionalities
of the basal ganglia and dopaminergic firing actualize primarily when the system is under
pressure to select sensorimotor constructs in the face of multidimensional uncertainty.
Our data cast doubt on a causal role for dopamine in visual perception, as shown by a
significant lack of effect in several standard perceptual paradigms. Future studies, perhaps
with more direct measures of dopaminergic activity, and more naturalistic paradigms
including more opportunities for selection and/or uncertainty, may yet reveal a more specific
influence of this neurotransmitter on how we perceptually encounter the world.

References

Agúndez, J. A., Garcia-Martin, E., Alonso-Navarro, H., & Jiménez-Jiménez, F. J. (2013). Anti-Parkinson's disease drugs and pharmacogenetic considerations. *Expert Opinion on Drug Metabolism & Toxicology*, *9*(7), 859-874.

Alexander, G. E., DeLong, M. R., & Strick, P. L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annual Review of Neuroscience*, *9*(1), 357-381.

Alexander, G. E., & Crutcher, M. D. (1990). Functional architecture of basal ganglia circuits: Neural substrates of parallel processing. *Trends in Neurosciences*, *13*(7), 266–271. https://doi.org/10.1016/0166-2236(90)90107-L

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019). Raincloud plots: A multi-platform tool for robust data visualization. *Wellcome Open Research*, *4*, 63. https://doi.org/10.12688/wellcomeopenres.15191.1

Arsalidou, M., Duerden, E. G., & Taylor, M. J. (2013). The centre of the brain: Topographical model of motor, cognitive, affective, and somatosensory functions of the basal ganglia. *Human Brain Mapping*, *34*(11), 3031–3054. https://doi.org/10.1002/hbm.22124

Barbato, G., Ficca, G., Muscettola, G., Fichele, M., Beatrice, M., & Rinaldi, F. (2000). Diurnal variation in spontaneous eye-blink rate. *Psychiatry Research*, *93*(2), 145-151.

Bauer, M., Wislowska, M., Veale, T., Morris, P. G., Liddle, P. F., Heekeren, H. R., & Brookes, M. J. (2016). Brain rhythms shape Bayesian integration of prior predictions with precision weighted sensory information for perceptual decisions. Annual Meeting of the Society for Neuroscience, San Diego, CA.

Berke, J. D. (2018). What does dopamine mean? *Nature Neuroscience*, 1–1. https://doi.org/10.1038/s41593-018-0152-y

Bisenius, S., Trapp, S., Neumann, J., & Schroeter, M. L. (2015). Identifying neural correlates of visual consciousness with ALE meta-analyses. *NeuroImage*, *122*, 177–187. https://doi.org/10.1016/j.neuroimage.2015.07.070

Blin, O., Masson, G., Azulay, J., Fondarai, J., & Serratrice, G. (1990). Apomorphine-induced blinking and yawning in healthy volunteers. *British Journal of Clinical Pharmacology*, *30*(5), 769–773. https://doi.org/10.1111/j.1365-2125.1990.tb03848.x

Boly, M., Massimini, M., Tsuchiya, N., Postle, B. R., Koch, C., & Tononi, G. (2017). Are the Neural Correlates of Consciousness in the Front or in the Back of the Cerebral Cortex? Clinical and Neuroimaging Evidence. *The Journal of Neuroscience*, *37*(40), 9603–9613. https://doi.org/10.1523/JNEUROSCI.3218-16.2017

Bonett, D. G., & Wright, T. A. (2000). Sample size requirements for estimating pearson, kendall and spearman correlations. *Psychometrika*, *65*(1), 23–28. https://doi.org/10.1007/BF02294183

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Breitmeyer, B. (2007). Visual masking: Past accomplishments, present status, future developments. *Advances in Cognitive Psychology*, *3*(1–2), 9–20. https://doi.org/10.2478/v10053-008-0010-7

Broadway, J. M., Frank, M. J., & Cavanagh, J. F. (2018). Dopamine D2 agonist affects visuospatial working memory distractor interference depending on individual

differences in baseline working memory span. *Cognitive, Affective, & Behavioral Neuroscience*, *18*(3), 509–520. https://doi.org/10.3758/s13415-018-0584-6

Brown, L. L., Schneider, J. S., & Lidsky, T. I. (1997). Sensory and cognitive functions of the basal ganglia. *Current Opinion in Neurobiology*, *7*(2), 157–163. https://doi.org/10.1016/S0959-4388(97)80003-7

Brown, S. B. R. E., Slagter, H. A., van Noorden, M. S., Giltay, E. J., van der Wee, N. J. A., & Nieuwenhuis, S. (2016). Effects of clonidine and scopolamine on multiple target detection in rapid serial visual presentation. *Psychopharmacology*, *233*(2), 341–350. https://doi.org/10.1007/s00213-015-4111-y

Buzsáki, G. (2019). *The Brain from Inside Out*. Oxford: Oxford University Press.

Cassidy, C. M., Balsam, P. D., Weinstein, J. J., Rosengard, R. J., Slifstein, M., Daw, N. D., … Horga, G. (2018). A Perceptual Inference Mechanism for Hallucinations Linked to Striatal Dopamine. *Current Biology*, *28*(4), 503-514.e4. https://doi.org/10.1016/j.cub.2017.12.059

Cavanagh, J. F., Masters, S. E., Bath, K., & Frank, M. J. (2014). Conflict acts as an implicit cost in reinforcement learning. *Nature Communications*, *5*, 5394–5394. https://doi.org/10.1038/ncomms6394

Cohen, M. X., Krohn‐Grimberghe, A., Elger, C. E., & Weber, B. (2007). Dopamine gene predicts the brain's response to dopaminergic drug. *European Journal of Neuroscience*, *26*(12), 3652-3660.

Colzato, L. S., Slagter, H. A., Spapé, M. M. A., & Hommel, B. (2008). Blinks of the eye predict blinks of the mind. *Neuropsychologia*, *46*(13), 3179–3183. https://doi.org/10.1016/j.neuropsychologia.2008.07.006

Cools, R., Gibbs, S. E., Miyakawa, A., Jagust, W., & D'Esposito, M. (2008). Working Memory Capacity Predicts Dopamine Synthesis Capacity in the Human Striatum. *Journal of Neuroscience*, *28*(5), 1208–1212. https://doi.org/10.1523/JNEUROSCI.4475-07.2008

Cools, Roshan, & D'Esposito, M. (2011). Inverted-U-shaped dopamine actions on human working memory and cognitive control. *Biological Psychiatry*, *69*(12). https://doi.org/10.1016/j.biopsych.2011.03.028

Cools, R. (2011). Dopaminergic control of the striatum for high-level cognition. *Current Opinion in Neurobiology*, *21*(3), 402-407. https://doi.org/10.1016/j.conb.2011.04.002

Cox, J., & Witten, I. B. (2019). Striatal circuits for reward learning and decision-making. *Nature Reviews Neuroscience*. https://doi.org/10.1038/s41583-019-0189-2

Cremers, H. R., Wager, T. D., & Yarkoni, T. (2017). The relation between statistical power and inference in fMRI. *PLOS ONE*, *12*(11), e0184923. https://doi.org/10.1371/journal.pone.0184923

Crick, F., & Koch, C. (2003). A framework for consciousness. *Nature Neuroscience*, *6*(2), 119–126. https://doi.org/10.1038/nn0203-119

Czoty, P. W., Riddick, N. V., Gage, H. D., Sandridge, M., Nader, S. H., Garg, S., … Nader, M. A. (2009). Effect of Menstrual Cycle Phase on Dopamine D2 Receptor Availability in Female Cynomolgus Monkeys. *Neuropsychopharmacology*, *34*(3), 548–554. https://doi.org/10.1038/npp.2008.3

Dakin, S. C., Hess, R. F., Ledgeway, T., & Achtman, R. L. (2002). What causes non-
monotonic tuning of fMRI response to noisy images? *Current Biology*, *12*(14), R476–
R477. https://doi.org/10.1016/S0960-9822(02)00960-0

Dang, L. C., Samanez-Larkin, G. R., Castrellon, J. J., Perkins, S. F., Cowan, R. L.,
Newhouse, P. A., & Zald, D. H. (2017). Spontaneous Eye Blink Rate (EBR) Is
Uncorrelated with Dopamine D2 Receptor Availability and Unmodulated by
Dopamine Agonism in Healthy Adults. *Eneuro*, *4*(5), ENEURO.0211-17.2017.
https://doi.org/10.1523/ENEURO.0211-17.2017

Dehaene, S., & Changeux, J.-P. (2011). Experimental and Theoretical Approaches to
Conscious Processing. *Neuron*, *70*(2), 200–227.
https://doi.org/10.1016/j.neuron.2011.03.018

Del Cul, A., Baillet, S., & Dehaene, S. (2007). Brain dynamics underlying the nonlinear
threshold for access to consciousness. *PLoS Biology*, *5*(10), 2408–2423.
https://doi.org/10.1371/journal.pbio.0050260

Del Cul, A., Dehaene, S., & Leboyer, M. (2006). Preserved Subliminal Processing and
Impaired Conscious Access in Schizophrenia. *Archives of General Psychiatry*,
63(12), 1313–1323. https://doi.org/10.1001/archpsyc.63.12.1313

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-
trial EEG dynamics including independent component analysis. *Journal of
Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Depue, R. A., Luciana, M., Arbisi, P., Collins, P., & Leon, A. (1994). Dopamine and the
structure of personality: Relation of agonist-induced dopamine activity to positive
emotionality. *Journal of Personality and Social Psychology*, *67*(3), 485–498.
https://doi.org/10.1037/0022-3514.67.3.485

Ding, L., & Gold, J. (2013). The basal ganglia's contributions to perceptual decision making.
Neuron, *79*(4), 640–649. https://doi.org/10.1016/j.neuron.2013.07.042

Ebert, D. (1996). Eye-Blink Rates and Depression Is the Antidepressant Effect of Sleep
Deprivation Mediated by the Dopamine System? *Neuropsychopharmacology*, *15*(4),
332–339. https://doi.org/10.1016/0893-133X(95)00237-8

Edelman, G. M. (2003). Naturalizing consciousness: A theoretical framework. *Proceedings
of the National Academy of Sciences*, *100*(9), 5520–5524.
https://doi.org/10.1073/pnas.0931349100

Fallon, S. J., Zokaei, N., Norbury, A., Manohar, S. G., & Husain, M. (2017). Dopamine
Alters the Fidelity of Working Memory Representations according to Attentional
Demands. *Journal of Cognitive Neuroscience*, *29*(4), 728–738.
https://doi.org/10.1162/jocn_a_01073

FitzGerald, T. H. B., Dolan, R. J., & Friston, K. (2015). Dopamine, reward learning, and
active inference. *Frontiers in Computational Neuroscience*, *9*.
https://doi.org/10.3389/fncom.2015.00136

Ford, C. P. (2014). The role of D2-autoreceptors in regulating dopamine neuron activity and
transmission. *Neuroscience*, *282*, 13–22.
https://doi.org/10.1016/j.neuroscience.2014.01.025

Frank, M. J., Loughry, B., & O'Reilly, R. C. (2001). Interactions between frontal cortex and
basal ganglia in working memory: A computational model. *Cognitive, Affective, &*
Behavioral Neuroscience, *1*(2), 137–160. https://doi.org/10.3758/CABN.1.2.137
Frank, Michael J., & O'Reilly, R. C. (2006). A mechanistic account of striatal dopamine
function in human cognition: Psychopharmacological studies with cabergoline and
haloperidol. *Behavioral Neuroscience*, *120*(3), 497–517.
https://doi.org/10.1037/0735-7044.120.3.497
Fridman, E. A., Osborne, J. R., Mozley, P. D., Victor, J. D., & Schiff, N. D. (2019).
Presynaptic dopamine deficit in minimally conscious state patients following
traumatic brain injury. *Brain*. https://doi.org/10.1093/brain/awz118
Friston, K. J., Shiner, T., FitzGerald, T., Galea, J. M., Adams, R., Brown, H., … Bestmann,
S. (2012). Dopamine, affordance and active inference. *PLoS Computational Biology*,
8(1). https://doi.org/10.1371/journal.pcbi.1002327
Gerfen, C. R. (1992). The neostriatal mosaic: multiple levels of compartmental organization.
In *Advances in Neuroscience and Schizophrenia* (pp. 43-59). Springer, Vienna.
Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., …
Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in*
Neuroscience, *7*, 267–267. https://doi.org/10.3389/fnins.2013.00267
Green, D. M., & Swets, J. A. (1966). *Signal Detection Theory and Psychophysics*.
New York: Wiley.
Hohwy, J. (2012). Attention and conscious perception in the hypothesis testing
brain. *Frontiers in Psychology*, *3*, 96.
Hohwy, J. (2013). *The Predictive Mind*. Oxford University Press.
Hutton, J. T., Morris, J. L., & Elias, J. W. (1999). Visual contrast sensitivity in Parkinson's
disease is worsened with cabergoline treatment. *Parkinsonism and Related Disorders*,
5(3), 87–91. https://doi.org/10.1016/S1353-8020(99)00021-8
JASP Team (2019). JASP (Version 0.10.0)[Computer software].
Jongkees, B. J., & Colzato, L. S. (2016). Spontaneous eye blink rate as predictor of
dopamine-related cognitive function—A review. *Neuroscience and Biobehavioral*
Reviews, *71*, 58–82. https://doi.org/10.1016/j.neubiorev.2016.08.020
Karson, C. N. (1983). SPONTANEOUS EYE-BLINK RATES AND DOPAMINERGIC
SYSTEMS. *Brain*, *106*(3), 643–653. https://doi.org/10.1093/brain/106.3.643
de Lange, F. P., Heilbron, M., & Kok, P. (2018). How do expectations shape
perception?. *Trends in cognitive sciences*, *22*(9), 764-779.
Lou, H., Skewes, J., Thomsen, K., Overgaard, M., Lau, H., Mouridsen, K., & Roepstorff, A.
(2011). Lou et al. 2011. Dopaminergic stimulation enhances confidence and accuracy
in seeing rapidly presented words.pdf. *Journal of Vision*, *11*(2), 1–6.
https://doi.org/10.1167/11.2.15.Introduction
Maniscalco, B., & Lau, H. (2012). A signal detection theoretic approach for estimating
metacognitive sensitivity from confidence ratings. *Consciousness and*
Cognition, *21*(1), 422-430.
Minear, M., & Park, D. C. (2004). A lifespan database of adult facial stimuli. *Behavior*
Research Methods, Instruments, & Computers, *36*(4), 630–633.
https://doi.org/10.3758/BF03206543

Mohr, C., Sándor, P. S., Landis, T., Fathi, M., & Brugger, P. (2005). Blinking and schizotypal thinking. *Journal of Psychopharmacology*, *19*(5), 513–520. https://doi.org/10.1177/0269881105056538

Nandam, L. S., Hester, R., Wagner, J., Dean, A. J., Messer, C., Honeysett, A., … Bellgrove, M. A. (2013). Dopamine D 2 Receptor Modulation of Human Response Inhibition and Error Awareness. *Journal of Cognitive Neuroscience*, *25*(4), 649–656. https://doi.org/10.1162/jocn_a_00327

Norbury, A., Manohar, S., Rogers, R. D., & Husain, M. (2013). Dopamine modulates risk-taking as a function of baseline sensation-seeking trait. *Journal of Neuroscience*, *33*(32), 12982-12986.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, 1–9. https://doi.org/10.1155/2011/156869

Palmiter, R. D. (2011). Dopamine signaling as a neural correlate of consciousness. *Neuroscience*, *198*, 213–220. https://doi.org/10.1016/j.neuroscience.2011.06.089

Persiani, S., Rocchetti, M., Pacciarini, M. A., Holt, B., Toon, S., & Strolin-Benedetti, M. (1996). The effect of food on cabergoline pharmacokinetics and tolerability in healthy volunteers. *Biopharmaceutics & Drug Disposition*, *17*(5), 443-455.

Redgrave, P., Prescott, T. J., & Gurney, K. (1999). The basal ganglia: A vertebrate solution to the selection problem? *Neuroscience*, *89*(4), 1009–1023. https://doi.org/10.1016/S0306-4522(98)00319-4

Redgrave, Peter, Rodriguez, M., Smith, Y., Rodriguez-Oroz, M. C., Lehericy, S., Bergman, H., … Obeso, J. A. (2010). Goal-directed and habitual control in the basal ganglia: Implications for Parkinson's disease. *Nature Reviews Neuroscience*, *11*(11), 760–772. https://doi.org/10.1038/nrn2915

Rousselet, G. A., & Pernet, C. R. (2012). Improving standards in brain-behavior correlation analyses. *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00119

Schultz, W. (2016). Dopamine reward prediction-error signalling: A two-component response. *Nature Reviews Neuroscience*, *17*(3), 183–195. https://doi.org/10.1038/nrn.2015.26

Seger, C. A. (2008). How do the basal ganglia contribute to categorization? Their roles in generalization, response selection, and learning via feedback. *Neuroscience & Biobehavioral Reviews*, *32*(2), 265–278. https://doi.org/10.1016/j.neubiorev.2007.07.010

Seger, C. A. (2013). The visual corticostriatal loop through the tail of the caudate: circuitry and function. *Frontiers in Systems Neuroscience*, *7*, 104. https://doi.org/10.3389/fnsys.2013.00104

Sescousse, G., Ligneul, R., van Holst, R. J., Janssen, L. K., de Boer, F., Janssen, M., … Cools, R. (2018). Spontaneous eye blink rate and dopamine synthesis capacity: Preliminary evidence for an absence of positive correlation. *European Journal of Neuroscience*, *47*(9), 1081–1086. https://doi.org/10.1111/ejn.13895

Shapiro, K. L., Raymond, J. E., & Arnell, K. M. (1997). The attentional blink. *Trends in*
Cognitive Sciences, *1*(8), 291–296. https://doi.org/10.1016/S1364-6613(97)01094-2
Shen, W., Flajolet, M., Greengard, P., & Surmeier, D. J. (2008). Dichotomous dopaminergic
control of striatal synaptic plasticity. *Science (New York, N.Y.)*, *321*(5890), 848–851.
https://doi.org/10.1126/science.1160575
Slagter, H. A., Mazaheri, A., Reteig, L. C., Smolders, R., Figee, M., Mantione, M., … Denys,
D. (2017). Contributions of the Ventral Striatum to Conscious Perception: An
Intracranial EEG Study of the Attentional Blink. *The Journal of Neuroscience*, *37*(5),
1081–1089. https://doi.org/10.1523/JNEUROSCI.2282-16.2016
Slagter, H. A., Tomer, R., Christian, B. T., Fox, A. S., Colzato, L. S., King, C. R., …
Davidson, R. J. (2012). PET Evidence for a Role for Striatal Dopamine in the
Attentional Blink: Functional Implications. *Journal of Cognitive Neuroscience*, *24*(9),
1932–1940. https://doi.org/10.1162/jocn_a_00255
Smith, Y., Raju, D. V., Pare, J.-F., & Sidibe, M. (2004). The thalamostriatal system: A highly
specific network of the basal ganglia circuitry. *Trends in Neurosciences*, *27*(9), 520–
527. https://doi.org/10.1016/j.tins.2004.07.004
Stalnaker, T. A., Howard, J. D., Takahashi, Y. K., Gershman, S. J., Kahnt, T., &
Schoenbaum, G. (2019). Dopamine neuron ensembles signal the content of sensory
prediction errors. *Elife*, *8: e49315*. https://doi.org/10.7554/eLife.49315
Strakowski, S. M., & Sax, K. W. (1998). Progressive behavioral response to repeated d-
amphetamine challenge: Further evidence for sensitization in humans. *Biological*
Psychiatry, *44*(11), 1171–1177. https://doi.org/10.1016/S0006-3223(97)00454-X
Taylor, J. R., Elsworth, J. D., Lawrence, M. S., Sladek, J. R., Roth, R. H., & Redmond, D. E.
(1999). Spontaneous Blink Rates Correlate with Dopamine Levels in the Caudate
Nucleus of MPTP-Treated Monkeys. *Experimental Neurology*, *158*(1), 214–220.
https://doi.org/10.1006/exnr.1999.7093
Unsworth, N., Heitz, R. P., Schrock, J. C., & Engle, R. W. (2005). An automated version of
the operation span task. *Behavior Research Methods*, *37*(3), 498–505.
https://doi.org/10.3758/BF03192720
Usiello, A., Baik, J. H., Rougé-Pont, F., Picetti, R., Dierich, A., LeMeur, M., … Borrelli, E.
(2000). Distinct functions of the two isoforms of dopamine D2 receptors. *Nature*,
408(6809), 199–203. https://doi.org/10.1038/35041572
Van Breukelen, G. J. P., & Van Dijk, K. R. A. (2007). Use of covariates in randomized
controlled trials. *Journal of the International Neuropsychological Society*, *13*(05).
https://doi.org/10.1017/S1355617707071147
van der Post, J., de Waal, P. P., de Kam, M. L., Cohen, A. F., & van Gerven, J. M. A. (2004).
No evidence of the usefulness of eye blinking as a marker for central dopaminergic
activity. *Journal of Psychopharmacology*, *18*(1), 109–114.
https://doi.org/10.1177/0269881104042832
Van Opstal, F., Van Laeken, N., Verguts, T., Van Dijck, J. P., De Vos, F., Goethals, I., &
Fias, W. (2014). Correlation between individual differences in striatal dopamine and
in visual consciousness. *Current Biology*, *24*(7), R265–R266.
https://doi.org/10.1016/j.cub.2014.02.001

Widmann, A., Schröger, E., & Maess, B. (2015). Digital filter design for electrophysiological
data—A practical approach. *Journal of Neuroscience Methods*, *250*, 34–46.
https://doi.org/10.1016/j.jneumeth.2014.08.002
Wiecki, T. V., & Frank, M. J. (2013). A computational model of inhibitory control in frontal
cortex and basal ganglia. *Psychological Review*, *120*(2), 329–355.
https://doi.org/10.1037/a0031542
Wise, S. P., Murray, E. A., & Gerfen, C. R. (1996). The Frontal Cortex-Basal Ganglia
System in Primates. *Critical Reviews in Neurobiology*, *10*(3–4), 317–356.
https://doi.org/10.1615/CritRevNeurobiol.v10.i3-4.30
Yousif, N., Fu, R. Z., Abou-El-Ela Bourquin, B., Bhrugubanda, V., Schultz, S. R., &
Seemungal, B. M. (2016). Dopamine Activation Preserves Visual Motion Perception
Despite Noise Interference of Human V5/MT. *Journal of Neuroscience*, *36*(36),
9303–9312. https://doi.org/10.1523/JNEUROSCI.4452-15.2016