Robustness of epithelial sealing is an emerging property of local ERK
feedbacks driven by cell elimination

Léo Valon[1], Anđela Davidović[2], Florence Levillayer[1], Mathilde Chouly[1], Fabiana
Cerqueira-Campos[1] and Romain Levayer[1*]

1. Department of Developmental and Stem Cell Biology, Institut Pasteur, CNRS UMR
3738, 25 rue du Dr. Roux, 75015 Paris

2. Department of Computational Biology, Institut Pasteur, CNRS USR 3756, 28 rue du
Dr. Roux, 75015 Paris

* Lead contact: romain.levayer@pasteur.fr

Abstract:

While the pathways regulating apoptosis and cell extrusion are rather well
described[1,2], what regulates the precise spatio-temporal distribution of cell
elimination in tissues remains largely unknown. This is particularly relevant for
epithelia with high rates of cell elimination, a widespread situation during
embryogenesis[3-6] and epithelial homeostasis[7], where concomitant death of
neighbours could impair the maintenance of epithelial sealing. However, the extent
to which epithelial tissues can cope with concomitant cell death, and whether any
mechanism regulates such occurrence have never been explored so far. Here,
using the *Drosophila* pupal notum (a single layer epithelium) and a new
optogenetic tool to trigger caspase activation and cell extrusion, we first show that
concomitant death of clusters of at least three cells is sufficient to transiently
impair epithelial sealing. Such clustered extrusion was almost never observed *in*
***vivo*, suggesting the existence of a mechanism preventing concomitant elimination**
of neighbours. Statistical analysis and simulations of cell death distribution in the
notum highlighted a transient and local protective phase occurring near every
dying cell. This protection is driven by a transient activation of ERK in the direct
neighbours of extruding cells which reverts caspase activation and prevents
elimination of cells in clusters. Altogether, this study demonstrates that the
distribution of cell elimination in epithelia is an emerging property of transient and

local feedbacks through ERK activation which is required to maintain epithelial sealing in conditions of high rate of cell elimination.

Epithelial cell elimination is driven by extrusion, a succession of remodeling steps removing one cell from the epithelial layer while maintaining epithelial sealing[1,8]. We first asked whether concomitant extrusion of several neighbours impairs the maintenance of epithelial sealing. As such, we developed a UAS-optoDronc *Drosophila* line, which can trigger rapid caspase activation through blue light-induced clustering of Caspase9 (**Figure 1a**, **Figure S1a**). Expression of optoDronc in fly eyes (GMR-gal4 eye-specific driver) is sufficient to trigger cell death upon blue light exposure and can be rescued by expression of the downstream effector caspase inhibitor p35 (**Figure S1b**). We then used the *Drosophila* pupal notum to assess the efficiency of the construct in triggering epithelial cell elimination. Blue light exposure of clones expressing optoDronc triggers elimination of the majority of cells in less than one hour (**Figure 1b, Figure S1c,e**, **movie S1**). OptoDronc-triggered extrusions are similar to physiological extrusions in the pupal notum, albeit slightly faster (**Figure S1f**), and require effector caspase activation (**Figure S1d,e, movie S1**) like physiological extrusions in the notum[9,10]. We then induced concomitant elimination of group of cells of various sizes and shapes by expressing optoDronc in clones. While extrusions of single cells or several cells in lines occurred normally (**Figure 1b,e,f, movie S2 left**), concomitant extrusion of three cells or more in cluster led to aberrant extrusion: cells initiate contraction then relax transiently (**Figure 1c-f**, **movie S2 right**) and eventually close the gap through a process akin to wound healing (**Figure 1c**, E-cad accumulation at vertices, **movie S2 right**). Aberrant extrusions correlated with transient flow of injected extracellular fluorescent Dextran in between cells at the level of adherens junctions (**Figure 1g,i,j**, **movie S3 bottom**), suggesting that epithelial sealing is transiently impaired. Dextran flow however was not observed for cells eliminated in lines (**Figure 1g,h,j**, **movie S3 top**). Altogether, we conclude that concomitant extrusion (<30 min) of three cells or more in cluster leads to transient loss of epithelial sealing and is thus detrimental for the tissue.

Given the rate of cell elimination and assuming that cell eliminations are independent (Poisson process), concomitant elimination of 3 cells or more in cluster should occur

several times per movie (See **Methods** for details, more than 2 events per movie of 20 hours). Yet, we very rarely observed such clusters during notum morphogenesis (<1 case per movie, **Figure 4k**, n=5 pupae). We therefore checked whether cell death distribution was indeed following locally a Poisson process in the pupal notum. We focused on the posterior region of the notum where cell death distribution is rather uniform in time and space (**Figure 2A, movie S4**) to neglect as much as possible the impact of tissue patterning on the rate of cell elimination. We first characterised the distribution of cell death by calculating for every cell elimination the local density of cell death at different distances and different times from the eliminated cell (**Figure 2b**). We obtained a map of local death density for every movie (**Figure 2c**) which was then compared to 200 simulations of the death distribution assuming a Poisson process at the same rate of cell elimination. The difference between the simulated map and experimental map was then used to check local differences in the distribution (**Figure 2c, Figure S2a**) and averaged for 5 nota (**Figure 2d**). Strikingly, there was in our experimental data a significant reduction in the density of cell elimination in the vicinity of each dying cell ($<7\mu m$) in a short time window (starting at 10 minutes and up to 60 minutes) compared to the simulated distributions. This suggests that cell death distribution in the posterior region of the pupal notum does not strictly follow a Poisson process, and that a transient (~one hour) and local (~one cell diameter) refractory phase appears in the vicinity of each dying cell. To confirm this bias, we then used a closest neighbour analysis (**Figure 2e**): for each cell elimination we detected the nearest elimination in different time windows. While the distribution in late time windows overlapped (1h20' to 2h20', 2h20' to 3h20' after cell death **Figure 2 f,g**), there was a significant decrease of the probability of cell death for the first 60 minutes (20 to 80 minutes) following each cell elimination at a distance of 0-10 μm (~one cell diameter, two folds reduction, p=0.008 for distances<5μm). Finally, to confirm the existence of a local dispersion of cell death, we performed a classical p-value tests for dispersion using K-functions (see **Methods** and [11]). This method detects potential temporal and spatial dispersion in a distribution. To validate the approach, we first compared the distribution of one of our experiments with the one of a simulated Poisson process with the same effective death intensity. We also compared the experimental distribution with the one of a simulated process with same intrinsic death intensity (see

Methods) adding a local and transient inhibition of cell death near every dying cell (5

μm range, 40 min inhibition starting 10 min after the initial death, see **Methods**). While no

significant dispersion appears in the Poisson distribution, a significant dispersion peak

appears both for the experimental distribution and the simulated one with local inhibition

(**Figure 2h**, simulated maps are averaged of 20 simulations), and similar dispersion

peaks were found for the other movies analysed (**Figure S2b**, **Figure 2i**). Altogether, we

concluded that cell elimination is followed by a transient and local refractory phase (~one

cell distance, with a delay of ~10-20 minutes and lasting ~60 minutes) that reduces

locally and transiently the probability of cell death (up to 2 folds).

These results suggest the existence of active mechanisms preventing the elimination of

the neighbours and generating this refractory phase. Effector caspase activation

systematically precedes and is required for every cell extrusion in the pupal notum[9,10].

We therefore tracked caspase dynamics using a GFP live sensor of effector caspase

activity (GC3Ai, [12,13]) and used the rate of GFP signal accumulation as a proxy for

caspase activity (see **Methods**). We frequently observed neighbours simultaneously

activating caspases (**Figure 3a**, **movie S5**), however extrusion of the first cell led to

reversion of caspase activity in the neighbours which then remained in the tissue (**Figure**

3a,b,c, caspase activity goes back to basal levels in the neighbours). This suggested that

cell extrusion could inhibit caspase activity and elimination of the direct cell neighbours.

We next asked which mechanism could modulate the dynamics of caspase activation and

bias the distribution of cell elimination. Recently we showed that the EGFR/ERK pathway

is a central regulator of cell elimination and caspase activity in the pupal notum[10], mostly

through the inhibition of the pro-apoptotic gene *hid*[10,14,15]. Moreover, EGFR/ERK can be

activated by tissue stretching and downregulated by tissue compaction[10]. Using a live

sensor of ERK (miniCic-mScarlet, adapted from[10], nuclear accumulation of mScarlet at

low ERK activity and nuclear exclusion at high ERK activity, **Figure 3d**), we monitored

ERK dynamics near extruding cells. Strikingly, we observed systematically a transient

activation of ERK in the direct neighbours of every dying cell (**Figure 3d,f**, **movie S6**,

Figure S3a,b, **movie S7**, activation also observed with EKAR, a ERK FRET sensor[16]).

This activation was concomitant with the transient stretching of the neighbouring cells

triggered by cell extrusion (**Figure 3 e,f**), lasted for ~ 60min and was restricted to the

direct neighbours of the dying cell (**Figure 3f**). ERK activation does not correlate with
pulses of Calcium (**Figure S4a**,**b**, **movie S8**), does not require active secretion of
EGF/Spitz from the dying cells (**Figure S4c**,**d, movie S9**, contrary to enterocytes
elimination in the fly midgut[17]) and can be mimicked by laser induced cell elimination
(**Figure S4e**,**f**, **movie S10**). We previously showed that stretch-induced survival in the
notum required EGFR[10]. Accordingly, ERK activation near dying cells is completely
abolished upon EGFR depletion in the tissue (**Figure 3g,h**, **movie S11**, RNAi previously
validated[10]). Moreover, EGFR is only required in the neighbouring cells and not in the
dying cell (**Figure S3e**). Altogether, this suggested that ERK activation could be driven
by the stretching induced by cell extrusion through EGFR activation, although at this stage
we cannot exclude other contact dependent mechanisms. Importantly, these pulses of
ERK are not restricted to the pupal notum as similar dynamics were observed near dying
larval accessory cells in the pupal abdomen (**Figure S3 c,d**, **movie S12**).

We then asked whether ERK pulses were indeed required for caspase reversion in the
neighbours. We first correlated ERK dynamics with caspase activity. We observed a
significant positive correlation between ERK activity and caspase inhibition with an
averaged lag-time of 15 minutes (**Figure 4a-c**, **movie S13** cross-correlation between
nuclear miniCic and GC3Ai differential, $r^2=0.37$). Moreover, caspase reversion in the
neighbouring cells was abolished upon EGFR depletion (**Figure 4d-f**, **movie S14**). Thus,
ERK pulses precede and are required for caspase inhibition in the neighbouring cells
following each cell death. The localisation and duration of ERK activation (1 row of cells,
~one hour, **Figure 3f**) and the delay with caspases inhibition (15 minutes) were
remarkably similar to the death refractory zone we observed near every cell elimination
($<7\mu m$, 10-20 minutes delay, duration of 60 minutes, **Figure 2**). We therefore checked
whether ERK pulses could modulate the spatio-temporal distribution of cell elimination in
the posterior region of the notum. Strikingly the refractory phase we found in WT pupae
(**Figure 2**) was not observed in EGFR depleted nota either using the local density of cell
death compared to simulations (**Figure 4g,h**, **movie S15**), using the closest neighbours
analysis (**Figure 4i**) or using the p-value dispersion test (no significant dispersion peak,
Figure 4j, **Figure S2d**) . This suggested that EGFR and ERK pulses are indeed required
to generate the transient and local death refractory phase. In absence of local inhibition,

we expected to observe spontaneous occurrence of clusters of cell elimination.
Accordingly, we observed frequent concomitant extrusion of clusters of cells upon EGFR
depletion in the notum (**Figure 4k**, **movie S16**, >3 cells, ~4 events per movie, n=9 pupae).
These clusters undergo abortive extrusion, transient relaxation followed by wound healing
(**Figure 4m**, **movie S16,** E-cad accumulation at vertices), similar to the abortive
extrusions triggered by optoDronc cell clusters (**Figure 1c-f, movie S2**). Altogether, this
showed that ERK pulses are required to reverse caspase activation near dying cells and
to prevent elimination of cells in clusters. This feedback is essential to maintain epithelial
sealing in conditions with a high rate of cell elimination (**Figure 4m**).

High rates of cell elimination are widespread during development and in adult tissues with
high cell turnover. Similar local feedback mechanisms may be required for coherent cell
elimination in the gut to prevent transient and recurrent sealing defects that could lead to
chronic inflammation and inflammatory bowel disease[18]. Importantly, the mechanism we
characterised in this study are conserved in mammals as similar ERK dynamics were
observed near dying MCF10A cells which also generate transient resistance to apoptosis
(personal communication, Olivier Pertz lab). Interestingly, other reports also described
Ca^{2+} and ERK activation waves emanating from extruding MDCK and MCF10A
pretumoral cells or caspase-induced extrusion[19,20]. However in these cases the waves
rather promote extrusion of the dying/extruding cells through collective convergent
movements. In contrast, we still see a high rate of cell extrusion upon depletion of ERK
feedbacks in our tissue (*UAS-EGFR dsRNA*). Strikingly, the range of the communication
seems to be very different between the notum (one cell row) and mammalian cells (10 to
15 cells away), suggesting either that the relays of communication are different, or that
mechanical properties of these cells are very different.

Although at this stage we cannot rule out the contribution of contact-dependent
communication for ERK activation in dying cell neighbours, several evidences point for a
contribution of cell stretching. First, ERK activation dynamics correlate very well with
transient cell stretching (**Fig. 3e,f**) and occurs very fast (few minutes). Moreover, we
previously showed that cell stretching could promote cell survival through EGFR[10].
Accordingly, we found that EGFR depletion completely abolished ERK feedback near

dying cells. We exclude the contribution of calcium signaling or an active secretion of the
ligand from the dying cell, while laser induced wound healing was sufficient to promote
neighbouring cell stretching and ERK activation. Other recent works have characterized
the rapid effect of cell stretching on ERK activation[21,22], which was recently shown to also
rely on EGFR[23]. Further work characterising the molecular mechanism of cell deformation
sensing by ERK will be required to fully test the contribution of mechanics in extrusion-
induced ERK activation.

The occurrence of aberrant extrusion remarkably increases for clusters of three cells or
more. However, we do not know at this stage what triggers these abnormal extrusions.
On the one hand, abortive extrusion may be related to the distance that needs to be
closed by the neighbouring cells: while this can be short for lines of cells (using the
shortest axis), it will be higher for clusters and may prevent fast fusion of neighbouring
cell junctions and gap closure[24]. On the other hand, abortive extrusion may be related to
caspase-induced remodeling of tricellular junctions[25], which are shared by caspase
positive cells only in the case of clustered eliminations. Interestingly, some tricellular
junction components have been shown to be cleaved by caspases[26]. Further
characterisation of extrusion dynamics will be required to test these hypotheses.

Interestingly, 2D simulations of cell disappearance combined with transient death
refractory phase in the neighbours suggest that ERK feedback could have a significant
impact on the total number of dying cells (see **Figure S5a,** ~1.4 folds reduction for a 60
minutes refractory phase and for the rate of cell elimination we observed in the posterior
region of the notum). This buffering effect increases sharply with the rate of cell
elimination (**Figure S5a**). As such, it may be essential in conditions of high stress to
dampen the rate of cell elimination and prevent tissue collapse. Moreover, this feedback
could be sufficient to increase the rate of cell death near clones resistant to apoptosis
when they are located in regions with a high rate of cell death (see **Figure S5b**, ~1.5 folds
increase for cells completely surrounded by cells resistant to apoptosis in our conditions).
This suggests that extrusion-driven ERK feedbacks may be sufficient to recapitulate some
of the features of cell competition: namely a contact and context dependent increase of
cell death near mutant clones[27]. Altogether, we propose that epithelial robustness and

plasticity may be emerging features of local and transient ERK feedbacks driven by cell
death.

Acknowledgements

We thank members of RL lab for critical reading of the manuscript, especially Alexis
Matamoro-Vidal for suggestions on the manuscript organisation. We would like to thank
Jakub Voznica for initiating observations of miniCic in the pupal abdomen during his
internship. We also thank Virgile Andreani for help in improving simulations, statistical
analysis of the data, and some mathematical expressions. We are also grateful to
Magalie Suzanne, the Bloomington Drosophila Stock Center, the Drosophila Genetic
Resource Center, the Vienna Drosophila Resource Center for sharing stocks and
reagents. We also thank B Aigouy for the Packing Analyser software and J. Ellenberg
group for MyPic autofocus macro. LV is supported by a Post-doctoral grant "Aide au
Retour en France" from the FRM (Fondation pour la Recherche Médicale,
ARF20170938651) and a Marie Sklodowska-Curie postdoctoral fellowship (MechDeath,
789573), work in RL lab is supported by the Institut Pasteur (G5 starting package), the
ERC starting grant CoSpaDD (Competition for Space in Development and Disease, grant
number 758457), the Cercle FSER and the CNRS (UMR 3738).

Authors contribution

RL and LV discussed and designed the project. RL did the experiments with GC3Ai and
wrote the manuscript with LV. AD performed the simulations of the theoretical distribution
of cell death, analysed experimental and simulated data using density maps and p-values
maps, wrote theoretical expressions of death rates. LV and AD designed 2D numerical
models to reproduce the effect of the feedback. FL designed miniCic-Scarlet flies and
designed and performed the preliminary test of the optoDronc fly line. LV and MC set up
Dextran injection in the pupal notum and initiated experiments with optoDronc. FCC
provided the Spitz RNAi movies and part of the WT pupae movies. LV performed all the
other analysis and experiments. Every author has commented and edited the manuscript.

Declaration of interests

241   The authors declare no competing interest

Methods

Drosophila melanogaster husbandry

All the experiments were performed with *Drosophila melanogaster* fly lines with regular husbandry techniques. The fly food used contains agar agar (7.6 g/l), saccharose (53 g/l) dry yeast (48 g/l), maize flour (38.4 g/l), propionic acid (3.8 ml/l), Nipagin 10% (23.9 ml/l) all mixed in one liter of distilled water. Flies were raised at 25°C in plastic vials with a 12h/12h dark light cycle at 60% of moisture unless specified in the legends and in the table below (alternatively raised at 18°C or 29°C). Females and males were used without distinction for all the experiments. We did not determine the health/immune status of pupae, adults, embryos and larvae, they were not involved in previous procedures, and they were all drug and test naïve.

Drosophila melanogaster strains

The strains used in this study and their origin are listed in the table below.

Fly line	Chromosome location	Origin (citation)	Source
E-cad-tdTomato (KI)	II	[28]	BDSC_58789
E-cad-GFP (KI)	II	[28]	BDSC_60584
UAS-optoDronc	II (attp40)	This study	This study
Tub-miniCic-mScarlet	II	This study	This study
ubi-EKARnls	II	[16]	Shigeo Hayashi
UAS-EGFR dsRNA	II	VDRC	KK 107130
UAS-spitz dsRNA	II	VDRC	GD 3922
UAS-GcAMPx20	III	Bloomington	BDSC_32236
UAS-GC3Ai	III	[12]	Magalie Suzanne
GMR-gal4	II	Bloomington	BDSC_1104
UAS-his3.3 mIFP-T2A-H01	III	Bloomington	BDSC_64184
Hs-flp22; ; Act<cd2<gal4,UAS-GFP	I; III	[10]	[10]

The exact genotype used for each experiment is listed in the next table. ACI: time After Clone Induction, APF: After Pupal Formation, n: number of pupae/adults.

Figure	Genotype	Heat shock 37°C	Stage	n
1b-j	hs-flp22; E-cad-tdTomato (KI)/UAS-optoDronc ; act<y+<gal4 , UAS-GFP/+	12 min, 2 days ACI	30h APF	10
2a	E-cad-GFP (KI), tub-miniCic-mScarlet ; pnr-gal4, UAS-His3-mIFP	N/A	16h APF	5
3a-c	E-cad-tdTomato (KI)/+; pnr-gal4/UAS-GC3Ai	N/A	20h APF	3
3d-f	E-cad-GFP (KI), tub-miniCic-mScarlet ; pnr-gal4, UAS-His3-mIFP	N/A	20h APF	5 (quantified on 2)
3g	E-cad-GFP (KI), tub-miniCic-mScarlet/UAS-EGFR dsRNA ; pnr-gal4, UAS-His3-mIFP/+	N/A	16h APF	3 (quantified on 2)
4a-c	tub-miniCic-mScarlet ; pnr-gal4/UAS-GC3Ai	N/A	20h APF	3
4d-f	E-cad-tdTomato (KI)/UAS-EGFR dsRNA; pnr-gal4/UAS-GC3Ai	N/A	20h APF	3
4g-l	E-cad-GFP (KI), tub-miniCic-mScarlet/UAS-EGFR dsRNA ; pnr-gal4, UAS-His3-mIFP/+	N/A	16h APF	9
S1a	E-cad-tdTomato (KI)/UAS-optoDronc; pnr-gal4/+	N/A	16h APF	2
S1b	GMR-gal4/UAS-optoDronc (dark)	N/A	Adult	20
S1b	GMR-gal4/UAS-optoDronc (blue light)	N/A	Adult	19
S1b	GMR-gal4/UAS-optoDronc ; UAS-p35/+ (blue light)	N/A	Adult	46
S1c	hs-flp22; E-cad-tdTomato (KI)/UAS-optoDronc ; act<y+<gal4 , UAS-GFP/+	12 min, 2 days ACI	30h APF	3
S1c	hs-flp22, E-cad-tdTomato (KI)/UAS-optoDronc ; act<y+<gal4 , UAS-GFP/UAS-p35	12 min, 2 days ACI	30h APF	3
S1f	hs-flp22, E-cad-tdTomato (KI)/UAS-optoDronc ; act<y+<gal4 , UAS-GFP/+	12 min, 2 days ACI	30h APF	3
S1f	E-cad-GFP (KI), tub-miniCic-mScarlet ; pnr-gal4, UAS-His3-mIFP	N/A	30h APF	5
S2a	E-cad-GFP (KI), tub-miniCic-mScarlet ; pnr-gal4, UAS-His3-mIFP	N/A	16h APF	5
S2b	E-cad-GFP (KI), tub-miniCic-mScarlet/ UAS-EGFR dsRNA ; pnr-gal4, UAS-His3-mIFP	N/A	16h APF	4
S3a,b	tub-miniCic-mScarlet ; pnr-gal4/ubi-EKARnls	N/A	20h APF	4
S3c,d	E-cad-GFP (KI), tub-miniCic-mScarlet	N/A	abdomen (20h APF)	2
S3e	Hs-flp22; E-cad-GFP (KI), tub-miniCic-mScarlet/UAS-EGFR dsRNA ; act<y+<gal4, UAS-His3-mIFP/+	20 min, 2 days ACI	20h APF	3
S4a,b	E-cad-tdTomato (KI)/+; pnr-gal4/UAS-GCaMPx20	N/A	16h APF	2
S4c,d	Hs-flp22; E-cad-GFP (KI), tub-miniCic-mScarlet/UAS-spitz dsRNA ; act<y+<gal4, UAS-His3-mIFP/+	20 min, 2 days ACI	20h APF	2
S4e,f	E-cad-GFP (KI), tub-miniCic-mScarlet ; pnr-gal4, UAS-His3-mIFP	N/A	20h APF	7

Design of optoDronc

The GFP-linker-DRONC-linker-CRY2PHR was first cloned in the pCaspeR4-Tubp-Gal80 vector (addgene 17748) and subsequently cloned in the pJFRC4-3XUAS-IVS-mCD8::GFP (Addgene 26217). Initial cloning was performed by three successive amplifications/ligations. Briefly, GFP was first inserted in pCaspeR4-Tubp by PCR-amplifying GFP from pJFRC19-13XLexAop2-IVS-myr::GFP (Addgene 26224) adding NotI, BglII, XbaI restriction sites and linkers, and eventually ligation into pCasper4-TubP-Gal80 after digestion with NotI and XbaI. Dronc cDNA was then inserted in this plasmid through PCR amplification on the cDNA clone LP09975 (DGRC) adding BglII, NheI and XbaI restriction sites and linkers, and then ligation into pCasper4-TubP-GFP-linker cut with BglII and XbaI. Finally, CRY2PHR was inserted by amplifying residues 1-498 of CRY2 from pGal4BD-CRY2 (Addgene 28243) while adding NheI and XbaI sites and ligation into pCasper4-TubP-GFP-linker-DRONC-linker cut with NheI and XbaI. The GFP-linker-Dronc-linker-CRY2PHR was cut with NotI and XbaI and inserted by ligation in pJFRC4-3XUAS-IVS-mCD8::GFP (Addgene 26217) after digestion with NotI and XbaI. The construct was checked by sequencing and inserted at the attp site attp40A after injection by Bestgene. The primers used for the construct are listed below (restriction sites in bold and linker in italic).

Insert	Forward	Reverse
GFP-linker	TTATA**GCGGCCGC**ATGTCCAAAGGTGAAGAACT	AATT**TCTAGAAGATCT**_GCCGCCTCCTCCGGACCCACCACCTCCAGAGCCACCGCCACCC_TTGTAGAGCTCATCCATGCCGT
DRONC-linker	AATT**AGATCT**ATGCAGCCGCCGGAGCTCGAGATT	AATT**TCTAGAGCTAGC**_GCCGCCTCCTCCGGACCCACCACCTCCAGAGCCACCGCCACCT_TCGTTGAAAAACCCGGGATTG
CRY2PHR	AATT**AGATCT**ATGCAGCCGCCGGAGCTCGAGATT	AATT**TCTAGA**CTATGCTGCTCCGATCATGATCTGT

Induction of cell death using optoDronc

To induce optoDronc in the eye, GMR-gal4 females were crossed with homozygous males UAS-optoDronc. Tubes containing crosses and progeny were either kept in the dark at 25°C (control) or maintained in a cardboard box permanently lighted by a blue

LED array (LIU470A, Thorlab) in the same 25° incubator as the control. Female adult
eyes were then imaged on a Zeiss stereoV8 binocular equipped with a colour camera
(Axiocam Icc5).

For induction of optoDronc in clones in the pupal notum, *hs-flp; E-cad-tdTomato(KI);*
act<cd2<G4 females were crossed with homozygous *UAS-optoDronc* or *UAS-optoDronc;*
UAS-p35. Clones were induced through a 12 minutes heat shock in a 37°C waterbath.
Tubes were then maintained in the dark at 25°C. White pupae were collected 48 hours
after clone induction and aged for 24h at 25°C in the dark. Collection of pupae and
dissection were performed on a binocular with LED covered with home-made red filter
(Lee colour filter set, primary red) after checking that blue light was effectively cut (using
a spectrometer). Pupae were then imaged on a spinning disc confocal (Gataca system).
The full tissue was exposed to blue light using the diode 488 of the spinning disc system
(12% AOTF, 200ms exposure per plane, 1 stack/min). The proportion of remaining cells
was calculated by measuring the proportion of cells remaining in the tissue at time t
compared to t0. Extrusion profiles were obtained by segmenting extruding cells in the
optoDronc clones with E-cad-tdTomato signal (only single cell clones in this case) or WT
cells marked with E-cad-GFP in the posterior region of the notum using Packing
analyser[29]. Curves were aligned on the termination of extrusion (no more apical area
visible) and normalised with the averaged area on the first five points. Clone area profile
(**Figure 1d**, **Figure 4m**) were obtained by segmenting the group of cells in the clone and
calculating the area and the solidity (Area/Convex Area) on Matlab. Categorisation of
normal versus abnormal extrusion was based on the dynamics of clone contraction:
transient relaxation or accumulation of E-cad at vertices was counted as abnormal
extrusion. Large relaxation combined with clear E-cad accumulation at vertices was
counted as "transient holes".

Design of miniCic-mScarlet

The pCaspeR4-Tubp-miniCic-linker-mScarlet-I was obtained by amplifying by PCR
miniCic-linker from pCaspeR4-Tubp-miniCic-linker-mCherry[10] and mScarlet-I from
pmScarlet-i_C1 (Addgene 85044). These two inserts were cloned in the vector
pCaspeR4-TubP-Gal80 linearised by NotI, XbaI digestion (to excise Gal80) using

NEBuilder HiFi DNA Assembly Method. The construct was checked by sequencing and
inserted through P-element after injection by Bestgene. The primers used for the
construct are listed below.

Insert	Forward	Reverse
miniCic-linker	GTGAAGGTACCCGCCCGGGGATCAGATCC	ACAGAACCACCACCAGAACCAC
mScarlet-i	CGGTGGTGGTTCTGGTGGTGGTTCTGTGAGCAAGGGCGAGGCA	CAGAAGTAAGGTTCCTTCACAAAGATCCTCTAGATTACTTGTACAGCTCGTCCATGC

The construct was tested by comparing the dynamics and pattern in the notum compared
to previously characterised dynamics using miniCic-mCherry[10]. Similar dynamics were
also observed at the single cell level between miniCic-mScarlet and the FRET sensor ubi-
EKARnls[16] (**Fig. S3**).

Live imaging and movie preparation

Notum live imaging was performed as followed: the pupae were collected at the white
stage (0 hour after pupal formation), aged at 29°, glued on double sided tape on a slide
and surrounded by two home-made steel spacers (thickness: 0.64 mm, width 20x20mm).
The pupal case was opened up to the abdomen using forceps and mounted with a
20x40mm #1.5 coverslip where we buttered halocarbon oil 10S. The coverslip was then
attached to spacers and the slide with two pieces of tape. Pupae were collected 48 or
72h after clone induction and dissected usually at 16 to 18 hours APF (after pupal
formation). The time of imaging for each experiment is provided in the table above. Pupae
were dissected and imaged on a confocal spinning disc microscope (Gataca systems)
with a 40X oil objective (Nikon plan fluor, N.A. 1.30) or 100X oil objective (Nikon plan fluor
A N.A. 1.30) or a LSM880 equipped with a fast Airyscan using an oil 40X objective (N.A.
1.3), Z-stacks (1 µm/slice), every 5min using autofocus at 25°C. The autofocus was
performed using E-cad-GFP plane as a reference (using a Zen Macro developed by Jan
Ellenberg laboratory, MyPic) or a custom made Metamorph journal on the spinning disc.
Movies were performed in the nota close to the scutellum region containing the midline
and the aDC and pDC macrochaetae. Movies shown are adaptive local Z-projections.
Briefly, E-cad plane was used as a reference to locate the plane of interest on sub

339 windows (using the maximum in Z of average intensity or the maximum of the standard
340 deviation). Nuclear signal was then obtained by projecting maximum of intensity on 7 μm
341 (7 slides) around a focal point which was located 6 μm basal to adherens junctions (see
342 [10] for more details).

Laser ablation

Photo-ablation experiments were performed using a pulse UV-laser (355nm, Teem
photonics, 20kHz, peak power 0.7kW) coupled to a Ilas-pulse module (Gataca-systems)
attached to our spinning disk microscope. The module was first calibrated and then set
to 40-60% laser power. Images were taken every 500ms and 3 to 6 single cells were
ablated 10 images after the beginning of the movie (20 repetitions of "point" ablation
~50ms exposure per cell). Cells were selected in regions with high and homogeneous
nuclear miniCic levels. For each single cell ablation, miniCic signals over time were
extracted in 4 cell nuclei of the first row and second row of cells around the ablated cell.
Nuclei were visualized and clicked by hand using the Histone3-mIFP fluorescent channel
after a local z-projection. Background value for each movies (for miniCic) were extracted
and removed from the signals, then the single cell miniCic signal was normalized to 1
using the first time point.

Dextran injection in pupal notum

30h APF pupae were glued on double sided tape and the notum was dissected with red
filter. Pupae were then injected using home-made needles (borosilicate glass capillary,
outer diameter 1mm, internal diameter 0.5mm, 10cm long, Sutter instruments) pulled with
a Sutter instrument P1000 pipette pulling apparatus. Dextran Alexa 647 10,000 MW
(Thermofisher, D22914) was injected at 2mg/ml in the thorax of the pupae using a
Narishige IM400 injector using a constant pressure differential (continuous leakage) and
depressurisation in between pupae. Imaging of Dextran leakage in live pupae was
performed after local projection using E-cad plane as a reference to measure Dextran
concentration at the junction plane (note that septate junctions are located basally to
adherens junctions). Dextran intensity was measured using a ROI in the center of the
clone after normalization using the first 5 time points and removal of intensity background

(estimated on a region with no signal. Note that Dextran injection cannot be used to track

leakages over long time scales (>4 hours) as the majority of Dextran get rapidly trapped

in fat body cells and in epithelial cell endosomes.

Signaling dynamics and cross-correlation

Measurement of ERK activity

ERK dynamics were measured using the mean nuclear intensity of miniCic. Whenever

possible, a marker of the nucleus (His3-mIFP) was used to track the nucleus manually

(Fiji macro). EKAR FRET signal was obtained after local projection on 6 planes of CFP

and YFP signals. CFP and YFP signals were blurred (Gaussian blur, 2 pixel width) and

the ratio of YFP/CFP signal was then calculated. Raw YFP-nls signal was thresholded

and used as a mask to only keep nuclear FRET signal. For the comparison of miniCic

and EKAR, single curves from the same cell were aligned on the peak of the FRET signal

(maximum value) and eventually averaged for all the cells.

For the measurement of miniCic in neighbours combined with apical area (**Figure 3**), cell

contours were tracked using E-cad-GFP (Packing analyser), while nuclei were manually

tracked on Matlab using His3-mIFP. For each dying cell, the nuclear miniCic intensity and

apical area were averaged for all the neighbours (first row and second row of cells). The

final curves are averaged of each of these curves after alignment at the termination of

cell extrusion (apical area=0). Note that the "n" used for s.e.m. calculation is not the total

number of neighbours, but the number of cell clusters.

Analysis of GC3Ai signal

To analyse caspase activity, we used the differential of GC3Ai signal as a proxy. GC3Ai

becomes fluorescent upon cleavage of a domain by effector caspases which triggers GFP

folding and maturation[12]. The dynamics of GFP signal can be written as follows:

$$\frac{dG}{dt} = C.G_0 - \alpha.G$$

Where G stands for concentration of active GFP, C concentration of active effector

caspase, G_0 concentration of inactive GFP, and α rate of GFP degradation.

On timescales of one hour, we can neglect GFP degradation. Assuming a constant pool
of inactive GC3Ai, G_0 can be considered as a constant. As such:

$$\frac{dG}{dt} = C.G_0$$

Therefore the change of intensity over time should be proportional to effector caspase
activity (C). Practically, GC3Ai signal was obtained after local projection (using E-cad or
nuclear miniCic as a reference) and measurement of mean intensity on small circular ROI
at the center of the cell (Fiji). The differential was calculated using the Diff function of
Matlab after smoothing the intensity curve (10 time points averaging window) and aligning
the curve on the time of disappearance (end of extrusion) of the first cell. Note that the
analysis of caspase reversion was restricted to neighbours having a positive slope before
the elimination of the first cell (in order to see caspase shutdown).

To cross-correlate GC3Ai and ERK dynamics, local projections centered on nuclear
miniCic were used (with GC3Ai plane of interest shifted apically). miniCic nuclear intensity
and GC3Ai cytoplasmic intensity were measured by manually tracking small circular ROI
(20px) in the center of the cells (Fiji home-made macro). The cross-correlation between
GC3Ai differential and miniCic nuclear signal was calculated on Matlab with the "xcorr"
function with the 'coef' option (normalised cross-correlation). All the curves (one per cell)
were then averaged.

Measurement of GCaMP

GCaMP signal was measured on local projections using E-cad-tdTomato as a reference
plane after global correction for bleaching. Each cell extrusion was cropped and the movie
realigned using the Stackreg Fiji plugin. GCaMP signal was then measured using the
mean intensity in a 20x20 pixel square in the center of each cells. Each single cell curve
was aligned (time 0 at the termination of extrusion) and normalised using 5 time values
around time 0.

Statistical analysis of cell death distribution

Calculation of the probability to observe three cells clustered elimination

We wanted to evaluate the probability to observe a cluster of three cells eliminated in less
than 30 minutes. We first considered a hexagonal grid of points in a squared array (N
cells per side) and assumed periodic boundaries. The probability of cell disappearance
over 30 min is p. As such the probability to observe a specific cluster of 3 cells eliminated
in the same 30 min window is p^3. The total number of possible 3 cells configurations is
approximatively given by: $2.N^2$.

Using the posterior region of WT nota, we estimate p~0.04 (probability of cell elimination
in 30 minutes). As such for a group of 400 cells (region of interest used in **Figure 2**)
observed for 20 hours (T= 40 times 30 minutes), the expected number of 3 cell-cluster
elimination is given by: $E(number\ of\ clusters) = 2N^2p^3T$ **= 2.05**.

Therefore, we would expect to observe several occurrences of 3 cell-cluster eliminations
per movie.

Analysis of death distribution

For the analysis of death distribution we selected movies lasting 16 to 20 hours in the
posterior region of the pupal notum. Cell extrusion events were clicked by hand and
checked twice. Extrusion localisations were corrected for global drift (translation) of the
tissue (Matlab procedure). Spatio-temporal distribution of death was first checked by x-
y-t histogram. According to that distribution, for each movie, an area of interest (in time
and space) was selected by hand to obtain an area where distribution was as
homogeneous as possible (double checked by x-y-t histograms). These data sets were
then the basis of three statistical analyses (see below).

1. Distribution of distances

Distances in time and space between all couples of extrusions were computed. For each
data set we plotted the maps of densities of death at a given distance (number of deaths
divided by the area of the disc considered) as a function of spatial and temporal distances
from each dying cell. From each data set we estimated the effective death rate, i.e. the
intensity of the random spatio-temporal process, and used it to simulate the
corresponding Poisson process 200 times. For each simulation we performed the same
analysis as for the experimental data set, namely calculating maps of death densities,

and eventually averaged the 200 maps. Finally, for each movie we calculated the
difference between the averaged simulated map and the corresponding experimental
map. We show in the main figures the average of the "difference map" for every
experiment (5 for WT, 4 for EGFR-RNAi).

2. Closest neighbour analysis

For each death event, we calculated the Euclidean distance to the closest death in a given
time window. Based on the previous analysis, we selected windows of 20' to 1h20', 1h20'
to 2h20' and 2h20' to 3h20'. We excluded the first 20 minutes as they correspond to the
characteristic time of extrusion (where cells cannot be reverted anymore). We then
ordered these shortest distances by size and plotted the cumulative probability of closest
death at a given distance. A value of 40% at a distance of 10 micrometers means than
40% of the closest deaths are localized between 0 and 10 micrometers away.

3. p-values test using "K-functions" and random labeling test
This procedure is used for estimating how far is one spatio-temporal process from a
purely random Poisson process[11]. The test is performed in two main steps. The first step
consists of calculating so-called "K-functions" of the given process, and the second is the
random re-labeling test as explained below.

The space-time K-function of the observed process, $K^0(r,h)$, is defined as the expected
number of additional events within the space-time distance (r,h) of a randomly selected
event. It depends on the intensity λ of the spatio-temporal process and it can be estimated
using only the observed events and without additional assumption on the process.

The complete absence of dispersion or clustering in a given process implies that there is
no relation between the location and timing of events. This is formally given through the
temporal indistinguishability hypothesis. Under this hypothesis, there is an equal
probability to observe our set of events, or events with the same locations in space but
with shuffled times. From here we can perform the random labeling test as follows.

We draw a sample of N random permutations of the times of events. For a given space-
time distance (r,h) we calculate the value of the K-function, $K(r,h)$, for each permutation.
We compute the number $M(r,h)$ of sampled re-labelings with the value of the K-function

smaller than the one of the original labeling, i.e. $K(r,h) < K^0(r,h)$. Then, the probability

of obtaining a value as small as $K^0(r,h)$ is estimated by the space-time dispersion p-

value: $p(r,h) = (M(r,h) + 1)/(N + 1)$. If we repeat the same procedure for all space-time

distances (r,h), we obtain the map of dispersion p-values.

This procedure is implemented in Python. The maximum distance is set to $r_{max} = 25\mu m$,

and the maximum temporal interval to $h_{max} = 150min$. We subdivided these maximum

values in 25 and 30 equal increments ($1\mu m$ and $5min$). We simulated 9999 re-labelings

of times to test for space-time dispersion of the extrusion events. The results of these 750

tests are plotted on a grid and then interpolated to obtain a p-value map. We performed

this procedure for each movie.

Calculation of the effective death rate with caspase resistant neighbours

We assume that the cell has six neighbours, and n neighbours resistant to apoptosis ($n \leq$

6), and that the death of one neighbour will protect the cell for a period T. The probability

of a cell to die at the time t is the probability that none of its $(6 - n)$ sensitive neighbours

died during the period T prior to t, multiplied by its intrinsic probability to die at the moment

t. The death of each isolated sensitive cell is a random event modelled as a temporal

Poisson process, with the event rate λ (expected number of cell deaths per minutes).

By definition, for the Poisson process X with rate λ the probability of the event (death) to

happen in the time interval $[t, t + dt]$, for dt very small, is:

499
$$P\{X \in [t, t + dt]\} = \lambda e^{-\lambda t} dt.$$

500   From here, the probability that an isolated sensitive cell dies during a time period of length

501   T is:

502
$$P\{X \in [0, T]\} = \int_0^T \lambda e^{-\lambda t} dt = 1 - e^{-\lambda T},$$

503   and the probability that it does not die during this time is: $1 - (1 - e^{-\lambda T}) = e^{-\lambda T}$.

504   The probability that the cell dies in a given moment t is defined as the probability to die

505   in the time interval $[t, t + dt]$, for dt very small, under assumption that it didn't die until the

506   time t. It is the same as the probability that it dies in the time interval $[0, dt]$:

507
$$p\colon P\{X \in [t, t+dt] | X \notin (0,t)\} = P\{X \in [0, dt]\} = \lambda e^{-\lambda 0} dt = \lambda dt.$$

508   This is also one of the definitions for the rate of the temporal Poisson process.

509   Thus, for the cell with six neighbours, where n neighbours are resistant to apoptosis ($n \leq$
510   6), the probability to die at the given moment t is:

511
$$p_e = \lambda e^{-\lambda_e T(6-n)} dt = \lambda_e dt.$$

512   Here we approximated the self-inhibiting random process with a Poisson process with a
513   different death rate, that we name the effective rate of death, and we used this rate to
514   express the probability that the neighbouring cells do not die during the period T. Finally,
515   the effective rate is:

516
$$\lambda_e = \frac{L_W(\lambda T(6-n))}{T(6-n)},$$

where L_W is Lambert W function. Note that we neglect here the effect of the delay of
inhibition on the neighbours. We used this expression in **Figure S5b.**

Calculation of the intrinsic death intensity for the self-inhibiting spatio-temporal
process, with continuous time and space
Assume that X is the spatio-temporal Poisson process. Then, by definition, the probability
that the number of events that fall in the spatial region C during the time period $[0, t]$ is
equal to k, is:

526
$$P\{N([0, t] \times C) = k\} = \frac{(\lambda \cdot t \cdot a(C))^k}{k!} e^{-\lambda t a(C)}, k = 0,1,2,3...$$

where λ is the space-time intensity of the process, i.e. it is the expected number of events
per unit of space-time volume (expected number of deaths per minute per micrometer
squared), and $a(C)$ is the surface of the given region C.

Hence, the probability that no event ($k = 0$) happens on a given region C during a period

of time τ is: $e^{-\lambda \tau a(C)}$. If the region C is a disk of radius r, its surface is $r^2\pi$, and this

probability is: $e^{-\lambda \tau r^2 \pi}$.

By definition of the space-time intensity of the process, the probability that the event

happens in a given moment t and that it falls on a position (x, y), is given by:

$$p = \lambda \, dt \, dx \, dy.$$

Now we consider the self-inhibiting spatio-temporal process. Meaning, for each event

there is a feedback that inhibits the process on a disk around the event of radius r, and

during a period of time τ. Such a self-inhibiting process is not anymore a Poisson process,

however it can be approximated as one with a different effective intensity, λ_e. The

probability that the death happens at some time t and on a given location (x, y), is the

probability that there was no event in its spatio-temporal vicinity (characterised by the

spatio-temporal volume of the feedback), multiplied with the probability of an isolated cell

death at this moment and place, or

$$p_e = \lambda e^{-\lambda_e \tau r^2 \pi} dt \, dx \, dy = \lambda_e \, dt \, dx \, dy.$$

Then the observed or effective intensity, λ_e, can be obtained by solving the equation:

$$\lambda_e = \lambda e^{-\lambda_e \tau r^2 \pi}.$$

Finally,

$$\lambda_e = \frac{L_W(\lambda \tau r^2 \pi)}{\tau r^2 \pi},$$

where L_W is Lambert W function. We can also obtain an expression for the intrinsic

intensity:

$$\lambda = \lambda_e e^{\lambda_e \tau r^2 \pi}.$$

Note that with this approximation we neglected an effect of the delay in the feedback.

This means that if we can estimate parameters of the feedback τ and r, and using the

observed intensity, λ_e, we can have an estimate on the intrinsic intensity of the spatio-

temporal self-inhibiting process.

To verify this expression numerically, we perform the following test: assume we want to

observe an effective intensity of the process $\lambda_e = 5.14 \cdot 10^{-5} min^{-1} m^{-2}$ with the self-

inhibiting process whose feedback has the parameters: $\tau = 40min$, $r = 5\mu m$, and delay

$10min$. We first calculate, using the above expression the intrinsic rate of the process to

be $\lambda = 6.04 \cdot 10^{-5}$. Using this rate and given parameters of the feedback, we run a long

simulation ($2 \cdot 10^6 min$) of the self-inhibiting process. We observed the effective intensity

of this process to be: $\lambda_e = 5.12 \cdot 10^{-5} min^{-1} m^{-2}$. For the comparison, we run the same

length of the simulation of the actual Poisson process (without feedback) with the intensity

$\lambda = \lambda_e = 5.14 \cdot 10^{-5} min^{-1} m^{-2}$. The observed intensity of this simulated process is $\lambda =$

$5.13 \cdot 10^{-5} min^{-1} m^{-2}$. This confirms that we have a good approximation for expression of

the intrinsic intensity of the self-inhibiting process. We performed such comparisons for

several values of intrinsic intensity.

We use this expression to estimate an intrinsic intensity of the processes of the

experimental data, and for the simulations of self-inhibiting processes that correspond to

our experiments (**Figure 2h**)

2D simulations of death rate

To evaluate the effect of the duration of the feedback as a function of the intrinsic rate of

death we made Python routine as follows:

We use a hexagonal grid as an approximation of the cell arrangement in the tissue. Each

node in the grid represents a cell, with the intrinsic rate of death λ. Number of cells per

side is n, so in total we have n^2 cells. The cells can be in 2 different states: "regular"

(sensitive to death), or "inhibited for death" (probability to die is 0). When one cell dies it

inhibits itself, and its 6 neighbours to die during some time T (in minutes).

Each death, $e = (t_e, x_e, y_e)$, is described with the time of death, t_e, and its coordinates in

the grid (x_e, y_e). We simulate the times of death for the Poisson process with rate $l = \lambda \cdot$

n^2. For each simulated time, t_e, we assign a random position (x_e, y_e) in the grid. Then we

check if any of its neighbours, or itself, has died in the previous T minutes. If there was

no death recorded in its neighbourhood, the cell is a regular cell and we record a new

death event (t_e, x_e, y_e). If there was a death in its neighbourhood in the last T minutes the

cell is refractory to death and we do not record this event.

At the end of simulation we compute the effective death rate as $\lambda_e = \frac{N}{(T_{max} \cdot n^2)}$, where N is

a total number of recorded deaths and T_{max} is the simulation time.

The parameters that we used in our simulations are the following: $n = 25$ (625 cells),

$T_{max} = 5000 min$. We simulated effects of different inhibition times $T \in$

$[0,5,10,20,\ldots,150] min$ with different intrinsic death rates, and repeated it 5 times per

value. (**Figure S5a**)

Statistics

Data were not analysed blindly. No specific method was used to predetermine the number

of samples. The definition of n and the number of samples is given in each figure legend

and in the table of the Experimental model section. Error bars are standard error of the

mean (s.e.m.). p-values are calculated through t-test if the data passed normality test

(Shapiro-Wilk test), or Mann-Whitney test/Rank sum test if the distribution was not

normal. Statistical tests were performed on Graphpad Prism 8 or Matlab.

References

1   Gudipaty, S. A. & Rosenblatt, J. Epithelial cell extrusion: Pathways and pathologies. *Semin Cell Dev*
Biol, doi:10.1016/j.semcdb.2016.05.010 (2016).
2   Fuchs, Y. & Steller, H. Programmed cell death in animal development and disease. *Cell* **147**, 742-
758, doi:10.1016/j.cell.2011.10.033 (2011).
3   Toyama, Y., Peralta, X. G., Wells, A. R., Kiehart, D. P. & Edwards, G. S. Apoptotic force and tissue
dynamics during Drosophila embryogenesis. *Science* **321**, 1683-1686,
doi:10.1126/science.1157052 (2008).
4   Monier, B. *et al.* Apico-basal forces exerted by apoptotic cells drive epithelium folding. *Nature*
518, 245-248, doi:10.1038/nature14152 (2015).
5   Ninov, N., Chiarelli, D. A. & Martin-Blanco, E. Extrinsic and intrinsic mechanisms directing
epithelial cell sheet replacement during Drosophila metamorphosis. *Development* **134**, 367-379,
doi:10.1242/dev.02728 (2007).
6   Marinari, E. *et al.* Live-cell delamination counterbalances epithelial growth to limit tissue
overcrowding. *Nature* **484**, 542-545, doi:10.1038/nature10984 (2012).

7   Williams, J. M. *et al.* Epithelial cell shedding and barrier function: a matter of life and death at the
small intestinal villus tip. *Vet Pathol* **52**, 445-455, doi:10.1177/0300985814559404 (2015).
8   Ohsawa, S., Vaughen, J. & Igaki, T. Cell Extrusion: A Stress-Responsive Force for Good or Evil in
Epithelial Homeostasis. *Dev Cell* **44**, 532, doi:10.1016/j.devcel.2018.02.007 (2018).
9   Levayer, R., Dupont, C. & Moreno, E. Tissue Crowding Induces Caspase-Dependent Competition
for Space. *Curr Biol* **26**, 670-677, doi:10.1016/j.cub.2015.12.072 (2016).
10   Moreno, E., Valon, L., Levillayer, F. & Levayer, R. Competition for Space Induces Cell Elimination
through   Compaction-Driven   ERK   Downregulation.   *Curr   Biol*   **29**,   23-34   e28,
doi:10.1016/j.cub.2018.11.007 (2019).
11   Smith, T. E. Notebook on Spatial Data Analysis *http://www.seas.upenn.edu/~ese502/#notebook*
(2020).
12   Schott, S. *et al.* A fluorescent toolkit for spatiotemporal tracking of apoptotic cells in living
Drosophila tissues. *Development* **144**, 3840-3846, doi:10.1242/dev.149807 (2017).
13   Zhang, J. *et al.* Visualization of caspase-3-like activity in cells using a genetically encoded
fluorescent   biosensor   activated   by   protein   cleavage.   *Nat   Commun*   **4**,   2157,
doi:10.1038/ncomms3157 (2013).
14   Kurada, P. & White, K. Ras promotes cell survival in Drosophila by downregulating hid expression.
Cell **95**, 319-329 (1998).
15   Bergmann, A., Agapite, J., McCall, K. & Steller, H. The Drosophila gene hid is a direct molecular
target of Ras-dependent survival signaling. *Cell* **95**, 331-341 (1998).
16   Ogura, Y., Wen, F. L., Sami, M. M., Shibata, T. & Hayashi, S. A Switch-like Activation Relay of EGFR-
ERK Signaling Regulates a Wave of Cellular Contractility for Epithelial Invagination. *Dev Cell*,
doi:10.1016/j.devcel.2018.06.004 (2018).
17   Liang, J., Balachandra, S., Ngo, S. & O'Brien, L. E. Feedback regulation of steady-state epithelial
turnover and organ size. *Nature* **548**, 588-591, doi:10.1038/nature23678 (2017).
18   Blander, J. M. Death in the intestinal epithelium-basic biology and implications for inflammatory
bowel disease. *FEBS J* **283**, 2720-2730, doi:10.1111/febs.13771 (2016).
19   Takeuchi,   Y.   *et   al.*   Calcium   Wave   Promotes   Cell   Extrusion.   *Curr   Biol*,
doi:10.1016/j.cub.2019.11.089 (2020).
20   Aikin, T. J., Peterson, A. F., Pokrass, M. J., Clark, H. R. & Regot, S. Collective MAPK Signaling
Dynamics Coordinates Epithelial Homeostasis. *Bioarchive*, doi:https://doi.org/10.1101/826917
(2019).
21   Aoki, K. *et al.* Stochastic ERK activation induced by noise and cell-to-cell propagation regulates cell
density-dependent proliferation. *Mol Cell* **52**, 529-540, doi:10.1016/j.molcel.2013.09.015 (2013).
22   Aoki, K. *et al.* Propagating Wave of ERK Activation Orients Collective Cell Migration. *Dev Cell* **43**,
305-317 e305, doi:10.1016/j.devcel.2017.10.016 (2017).
23   Hino, N. *et al.* ERK-mediated mechanochemical waves direct collective cell polarization.
Bioarchive, doi:https://doi.org/10.1101/2019.12.25.888552 (2019).
24   Staddon, M. F. *et al.* Cooperation of dual modes of cell motility promotes epithelial stress
relaxation   to   accelerate   wound   healing.   *PLoS   Comput   Biol*   **14**,   e1006502,
doi:10.1371/journal.pcbi.1006502 (2018).
25   Higashi, T. & Chiba, H. Molecular organization, regulation and function of tricellular junctions.
Biochim Biophys Acta Biomembr **1862**, 183143, doi:10.1016/j.bbamem.2019.183143 (2020).
26   Janke, S., Mittag, S., Reiche, J. & Huber, O. Apoptotic Fragmentation of Tricellulin. *Int J Mol Sci* **20**,
doi:10.3390/ijms20194882 (2019).
27   Claveria, C. & Torres, M. Cell Competition: Mechanisms and Physiological Roles. *Annu Rev Cell Dev*
Biol **32**, 411-439, doi:10.1146/annurev-cellbio-111315-125142 (2016).

28   Huang, J., Zhou, W., Watson, A. M., Jan, Y. N. & Hong, Y. Efficient ends-out gene targeting in
Drosophila. *Genetics* **180**, 703-707, doi:10.1534/genetics.108.090563 (2008).
29   Etournay, R. *et al.* TissueMiner: A multiscale analysis toolkit to quantify how cellular processes
create tissue dynamics. *Elife* **5**, doi:10.7554/eLife.14334 (2016).

Figure 1: Extrusion of a cluster of more than three cells is sufficient to impair epithelial sealing

a: Schematic of the UAS-optoDronc construct. The cDNA of Dronc (*Drosophila* Caspase9) is fused to eGFP in N-ter and the blue light sensitive protein CRY2-PHR in C-ter. Upon blue light exposure, optoDronc clusterises (green dots) and triggers caspase activation and apoptosis. **b:** Snapshots of a live pupal notum expressing optoDronc in clones (magenta, local z-projection) and E-cad-tdTomato. Most of the clones disappear after 60 minutes of blue light exposure. Scale bar= 10μm. **c:** Elimination of an optoDronc clone of nine cells. Snapshots of inverted E-cad-tdTomato signal. Time 0 is the termination of clone elimination. Clone contour rounds up and relaxes at -32 min and is followed by wound healing. Inset on the right shows E-cad accumulation at tricellular junctions during wound healing. Scale bar= 10μm. **d:** Evolution of the clone area shown in **c**. The grey zone corresponds to the relaxation phase and is followed by wound healing. The lines corresponds to the timing of the images shown in **c**. Inset shows increased of clone solidity (area/convex area) during the relaxation and wound healing phases (grey zone). **e:** Snapshots of clones of different sizes expressing optoDronc upon blue light exposure. Note the transient relaxation at 41 min for 5 cells cluster. Time 0 is the onset of blue light exposure. Scale bars= 10μm. **f:** Quantification of the proportion of normal and aberrant extrusions (extrusions followed by transient relaxation or E-cad accumulation at vertices) and/or transient hole (large relaxation and E-cad accumulation at vertices) observed for clone disappearance of different sizes and topologies (see schematic below). n=number of clones. Error bars are 95% confidence interval. **g:** Injection of far red Dextran 10,000 MW in pupal notum. Bottom shows a transverse view of Dextran signal in the pupae (x- antero-posterior, Z- apical-basal). Schematic showing the localisation of Dextran according to the epithelium (AJs: adherens junctions, SJs: septate junctions). Note that septate junctions are located basally to adherens junctions. **h:** Local projections of optoDronc clones composed of one to several cells in lines after Dextran injection (bottom). Time 0 is the end of clone elimination. No dextran appears at the level of adherens junctions during clone elimination. Scale bar= 10μm. **i:** Local projections of an optoDronc clone composed of four cells in cluster after Dextran injection (bottom). Time 0 is the end of clone elimination. During the relaxation phase (-35min) dextran appears at the level of adherens junctions (white squares). Scale bar= 10μm. **j:** Quantification of the far red Dextran signal during optoDronc clone elimination (in line, black, or in clusters of 3-4 cells, red). The curves are the median +/- s.e.m.. Top inset shows the average clone area during their elimination (note that the elimination is overall slower for clusters, red curve).

a

deaths

time (hours) 5 20

distance y (μm) 0 55

distance x (μm) 0 90

b

Local density of cell death

c

Single experiment   200 simulations   Sim. - exp.

spatial distance (mm)

time distance (min) 0 150

d

Average (Sim. - exp.), 5 nota

spatial distance (mm)

time distance (min) 0 75 150   0 10 50 150

e

Closest death analysis

f

time windows (hours) 0-1 1-2 2-3

5 um

integrated death probability (%)

distance from the extruding cell (μm) 5 10 15

g

**
(p=0.008)

integrated death probability at distance 5 mm

time windows (hours) 0-1 1-2 2-3

h

Dispersion measurments

Single experiment   mean of 20 simulations: without feedback   with 5 μm and 40 min feedback

***

<0.005   ***   *** <0.005

spatial distance (μm)

P value N.S.

time distance (min) 0 75 150

P-values maps from «K» functions

i

Average of 5 experiments   **

<0.05   **

spatial distance (μm)

P value N.S.

time distance (min) 0 75 150

Figure 2: There is a transient and local refractory phase for cell elimination following each cell death.

a: Snapshot of a local projection of a pupal notum expressing E-cad-GFP. Red dots are showing every cell extrusion occurring over 21 hours. A: Anterior, P: Posterior, L: Left, R: Right. Scale bar= 10μm. Orange histograms show the spatial distribution of cell death over left-right axis (left) or along AP axis (bottom). Right histograms show the temporal distribution of cell death (top) and its spatial distribution (middle and bottom) in the yellow rectangle region (used for further analysis). **b:** Scheme explaining the measurement of spatial and temporal distance between death events. For each death event, the density of cell death in a disc (number of death events divided by disc surface) at a given spatial distance is calculated for each time point. **c:** Local cell death density at different spatial (y axis) and temporal distances (x axis) from a dying cell for one movie. The middle map shows the average map obtained for 200 simulations of a Poisson process with the same cell death intensity. The right map shows the difference between the simulated and the experimental distributions ("yellow island" shows that death at short distances are under-represented in the experiment compared to the simulations). **d:** Average of the difference between experimental maps and the corresponding simulated maps (simulation minus experimental distributions, see **Figure S2** for details, 5 movies). The right map is obtained after median filtering. Note the bottom left yellow domain with lower expected number of cell deaths compared to simulations (black square, 7μm~one cell diameter, and from ~ 10 to 60min). **e:** Analysis of cell death distribution through a closest neighbour approach. Time is subdivided in arbitrary windows (here 1 hour, starting from t=20 minutes) and for each time window the spatially closest cell death is found. **f:** Cumulative plots of the probability to find the closest death at a given distance for different time windows (20min to 1hour 20, 1h20 to 2h20, 2h20 to 3h20). The curves are averages of 5 movies, error bars are s.e.m.. Note that only the blue curve detaches from the others, representing what happens 20 to 80 minutes after cell death. **g:** Details of the values obtained for a distance of 5μm (~ 1 cell distance) for each time window (one dot = one movie). There is a two-folds reduction of the

probability to have cell death occurring during the first hour after cell elimination for distances from 0 to 5 μm to the dying cell. **h:** Analysis of dispersion in the death distribution using maps of dispersion p-value calculated with K-functions (see **Methods**). Y-axis: spatial distance between death events, X-axis: time delay between death events. Pseudo colour is the p-value (yellow=significant dispersion, blue=no significant dispersion). The left map corresponds to the experimental distribution used in **2c,** middle map is the mean of the maps from 20 simulations of a Poisson process with the same death intensity; the right map is the mean of the maps of 20 simulations of a random process including a cell death refractory phase following each cell death starting with a delay of 10min and lasting 40min at a distance of 5μm (see **Methods**). **i:** Averaged p-value map for 5 WT movies (see **Figure S2** for details).

Figure 3: Transient ERK activation in cells neighbouring an extruding cell

a: Snapshots of local projections of a pair of cells in the pupal notum tagged with E-cad-tdTomato and expressing the effector caspase sensor GC3Ai (green). Time 0 is the termination of extrusion of the bottom cell (green contour). Scale bar=10μm. **b:** Intensity of GC3Ai in the eliminated cell (green) and its neighbour (purple). Note that GC3Ai signal plateaus in the neigbour after the first cell is eliminated. **c:** Averaged GC3Ai differential (a proxy of effector caspase activity, see **Methods**) for dying cells (green) and its neighbours (purple). Neighbours undergo transient caspase activation followed by reversion. Time 0 is the termination of the first cell extrusion. Light coloured areas show s.e.m. The top inset shows the purple curve with higher magnification. **d:** top: schematic showing the localisation of miniCic (n. : nucleus, c.: cytoplasm) upon modulation of ERK activity. Bottom: Snapshot of the posterior region of the pupal notum with two dying cells (red: miniCic-mScarlet, green: E-cad-GFP, blue: His3-mIFP). Right: snapshots of two dying cells (black stars) showing E-cad signal (inverted colour, top) and miniCic (bottom). Orange zones (red line for miniCic) are the first raw of neighbouring cells, grey zones (white line for miniCic) mark the second raw. Scale bars=10μm. **e:** Averaged cell apical area variation in the first row (red) and second row (black) of cells neighbouring an extruding cell. Time 0 is the termination of extrusion. Error bars are s.e.m.. **f:** Averaged nuclear miniCic intensity in the first row (red) and second row of cells (black). Error bars are s.e.m.. **g:** Snapshots of a *pnr-gal4 ; UAS-EGFRdsRNA* pupae (local projections) expressing E-cad-GFP, miniCic-mScarlet and His3-mIFP (not shown). The blue line shows a dying cell. The white doted circles show miniCic signal in the nuclei of the neighbouring cells. Scale bar=10μm. **h:** Averaged nuclear miniCic intensity (purple) and cell apical area (black) in EGFR depleted nota in the cells neighbouring an extruding cell (Time 0 is the termination of extrusion). Light areas are s.e.m..

a

55 min 80 min 95 min 120 min 145 min

miniCic

GC3Ai

b

miniCic Cell 1 — miniCic Cell2
GC3Ai Cell1 — GC3Ai Cell2

c

Crossscorrelation miniCic Vs caspase activity

before caspase | after caspase
n=20 cells

Time (min)

d

pnr-G4, UAS-EGFRdsRNA

-120 min -55 min -30 min -5 min 20 min 45 min

E-cad

GC3Ai

e

EGFR RNAi

GC3Ai intensity (a.u.)

Time (min)

f

Dying cells (n=40)
Neighbours (n=94)

EGFR RNAi

Effector caspase activity (GC3Ai differential, a.u.)

Time (min)

g

E-cad-GFP dying cells

pnr-G4, UAS-EGFRdsRNA

h

Death density analysis

Averaged (sim. - exp.) n=4

EGFR RNAi   WT

spatial distance (μm)

time distance (min)

i

Closest neighbour analysis

time windows (hours) 0-1 1-2 2-3

WT

integrated death probability (%)

EGFR RNAi

distance from the delaminating cell (μm)

j

Dispersion analysis

EGFR RNAi   WT

10^{-2}

P value

N.S.

spatial distance (μm)

time distance (min)

k

*** (p<0.001)

Number of clusters of death (>=3 cells, 16 to 20h movies)

WT (5 nota)   EGFR RNAi (9 nota)

l

E-cad-GFP

-130 min -70 min -30 min

-20 min -10 min 0 min

pnr-G4, UAS-EGFRdsRNA

m

norm. area

Area/ConvexArea

time (min)

time (min)

n

High ERK
Caspase3 activation
E-cad

ERK feedback

Death-refractory phase (~60 min)

No feedback (EGFR RNAi)

Sealing defect and wound healing

Figure 4: ERK pulses are required for caspase reversion and to prevent clusters of cell elimination

a: Snapshots of local projections of a pupal notum expressing miniCic and GC3Ai. White dotted circles show the first dying cell, white circles show one of its neighours dying later. Time is in agreement with the curves shown in **b**. **b:** Nuclear miniCic signal (magenta) and GC3Ai differential (green, caspase activity) in the first dying cell (dotted lines) and its neighbours (plain lines) shown in **a**. The red region highlights the pulse of ERK (decrease of miniCic) in the neighbours and the subsequent transient reversion of caspase activity. **c:** Averaged normalised cross-correlation between miniCic nuclear signal and caspase activity (GC3Ai differential). Peak at -15 min indicates a 15 minutes delay between ERK activation and caspase reversion. **d:** Snapshots of a local projection of a *pnr-gal4 , UAS-EGFRdsRNA* pupal notum expressing E-cad-tdTomato and GC3Ai. The white stars show three neighbouring cells dying. Green, light and dark purple cell contours correspond to the curves shown in **e**. Scale bar=10μm. **e:** GC3Ai intensity in the three cells marked in **d**. Time 0 is the termination of extrusion of the first dying cell (green contour). Note that there is no inflexion of GC3Ai signal in the neighbours. **f:** Averaged GC3Ai differential signal (caspase activity) in the dying cell (green) and its neighbours (purple) upon depletion of EGFR in the notum. Light areas are s.e.m.. Time 0 is the termination of extrusion of the first dying cell. On average, caspase activity is maintained in the neighbouring cells (differential >0, compare with **Figure 3b,c**). **g:** Local projection of a E-cad-GFP pupae depleted for EGFR (*pnr-gal4 , UAS-EGFR dsRNA*). Red dots show all the dying cells over the course of the movie (21 hours). Scale bar=10μm. The white rectangle show the analysis region (similar to the region used for the WT conditions). **h:** Averaged differences between the experimental distribution of cell death density in four EGFR depleted pupae and the corresponding simulated distribution (assuming independent events and the same death intensity). The inset shows the same analysis for WT pupae (from **Figure 2d**). Note the absence of yellow in the bottom left corner compared to the WT (no reduction of cell death density in EGFR depleted pupae for 10-60 min, <7 μm). **i:** Closest neighbour

analysis of cell death distribution in EGFR depleted pupae. Integrated death probability for different time windows after cell elimination (20min-1h20, 1h20-2h20, 2h20-3h20) at different distances from the dying cells. While the first hour distribution was different in the WT (see top left inset, blue curve, from **Figure 2 e-g**), there is no more differences between curves upon EGFR depletion. **j:** Averaged map of dispersion p-value from the K-functions for 4 *pnr-gal4 EGFR RNAi* movies (pseudoclour: p-value, blue=no significant dispersion). The top right inset shows the averaged map of dispersion p-value for control movies (see **Figure 2i**) **k:** Number of occurrences of clustered elimination (>=3 cells) per movie (16 to 20 hours) in the WT pupae (5 nota) and upon depletion of EGFR (*pnr-gal4, UAS-EGFR dsRNA*, 9 pupae). **l:** Snapshots of E-cad-GFP local projection in a EGFR depleted pupae showing concomitant elimination of three cells (orange area) and an aberrant extrusion (relaxation at -30 and wound healing figure with E-cad accumulation at vertices, bottom left inset, compare with **Figure 1 c,d**). Time 0 is the termination of cell elimination. **m:** Evolution of the clone area shown in **l**. The grey zone corresponds to the relaxation phase and is followed by wound healing. Inset shows increased of clone solidity (area/convex area) during the relaxation and wound healing phases. **n:** Schematic of the local ERK feedback (red) and its impact on the distribution of cell elimination (green cells: caspase activation). Upon EGFR depletion, simultaneous caspase activation and cell elimination can occur, which leads to aberrant extrusion and transient loss of sealing (phase outlined in yellow, white area between green aggregates) followed by wound healing.

a

optoDronc

0 sec   60 sec

Time (sec)

b

GMR-gal4 x
UAS-optoDronc
dark

GMR-gal4 x
UAS-optoDronc
blue light

GMR-gal4 x
UAS-optoDronc ; UAS-p35
blue light

20/20   19/19   46/46

c

Ecad-tdTomato; ayG4 ,UAS-GFPcyto x
UAS-optoDronc

blue light

0min   30min   60min

E-cad-tdTomato

GFPcyto+optoDronc

d

Ecad-tdTomato; ayG4 ,UAS-GFPcyto x
UAS-optoDronc UAS-p35

blue light

0min   30min   60min

E-cad-tdTomato

GFPcyto+optoDronc

e

UAS-optoDronc (3 nota, 214 cells)
UAS-optoDronc; UAS-p35 (3 nota, 350 cells)

Proportion of remaining cells

****   ****

Time (min)

f

Extrusion profiles

optoDRONC (n=16)
WT (n=11)

norm. area

time (min)

Figure S1: optoDronc triggers cell extrusion through effector caspase activation

a: Rapid clustering of UAS-optoDronc (GFP fusion) upon blue light exposure (single plane movie, 1 frame/sec). Scale bar= 10μm. Bottom: kymograph of optoDronc-GFP signal (clusters already appear after 20 sec) **b:** GMR-gal4 adult *Drosophila* eyes expressing UAS-optoDronc from flies raised in a blue light chamber (left), in the dark (middle) or with p35 overexpression (effector caspase inhibitor) in the light chamber. N are numbers of females. **c,d:** Snapshots of pupal nota (local projection) expressing *UAS-optoDronc* (**c**) or *UAS-optoDronc* and *UAS-p35* (**d**) in clones (bottom, UAS-GFP and optoDronc-GFP in the same channel) and E-cad-tdTomato. Green lines show clone contours. White arrowheads show extruding cells. Scale bars=10μm. **e:** Proportion of clonal cells remaining in the epithelium upon blue light exposure expressing *UAS-optoDronc* (green) or *UAS-optoDronc; UAS-p35* (light green). Error bars are 95% confidence interval,-= non-significant, ****=p<10^{-4}. **f:** Averaged and normalised cell apical area during spontaneous extrusion (black curve, WT movie, frame rate 5min) or upon induction of extrusion through light activation of optoDronc in single cell clones (purple curve). Curves are aligned at time 0 min (cell apical area = 0). Error bars are s.e.m.. Note that optoDronc-triggered extrusion is slightly faster than WT extrusion but globally similar.

a *pnr-gal4*

experiment #1 #2 #3 #4 #5

spatial distance (µm)

0 time (min) 150

200 simulations

Sim. - exp.

b Dispersion P-maps

experiment #1 #2 #3 #4 #5

0 time (min) 150

***

10⁻³

P value

N.S.

P-values maps from «K» functions

c *pnr-gal4 ; UAS-EGFR dsRNA*

experiment #1 #2 #3 #4

spatial distance (µm)

0 time (min) 150

200 simulations

Sim. - exp.

d Dispersion P-maps

experiment #1 #2 #3 #4

0 time (min) 150

***

10⁻³

P value

N.S.

P-values maps from «K» functions

Figure S2: Distribution of cell death in control nota and upon depletion of EGFR

a,c: Maps of the local death density near each dying cells at different times (see **Figure 2b-d**) for control pupae (**a**, *pnr-gal4*) or upon depletion of EGFR (**c**, *pnr-gal4; UAS-EGFR dsRNA*). Top: experimental distribution (one map = one pupa), y-axis: spatial distance in μm, x-axis: time distance in minutes. Middle: simulations of the death distribution for the corresponding experiment assuming a Poisson process with the same effective death intensity (average of 200 simulations). Bottom: Difference between experiment and simulation distributions (yellow= low proportion of cell death compared to simulations). **b,d:** Maps of the dispersion p-value calculated using K-functions for control pupae (**b**, *pnr-gal4*) or upon EGFR depletion (**d**, *pnr-gal4; UAS-EGFR dsRNA*). y-axis: spatial distance in μm, x-axis: time distance in minutes. Pseudo-colours show the significance of the dispersion (yellow: significant dispersion, blue: no significant dispersion).

a

pupal notum

miniCic

EKAR (FRET)

high ERK

low ERK

-15 min   0 min   15 min   30 min   45 min   60 min   75 min

b

EKAR FRET, 56 neighbours of 10 dying cells
miniCic, 56 neighbours of 10 dying cells

c

LEC abdomen

E-cad-GFP
miniCic

miniCic

-150 min   -100 min   -50 min   0 min   50 min   100 min   150 min

d

41 neighbours of 10 dying cells

end of extrusion

e

pupal notum

UAS-EGFR dsRNA

miniCic

E-cad-GFP

-30 min   -15 min   0 min   15 min   30 min   45 min   60 min   75 min   90 min

Figure S3: ERK pulses are also present in the abdomen and require EGFR cell-autonomously.

a: Snapshots of local projections of a pupal notum expressing miniCic (top, grey) and a FRET sensor of ERK (EKAR, bottom). Red signal: high ERK, blue signal: low ERK. The white star marks a dying cell. The dotted lines show the miniCic and the FRET signal in the neighbouring cells. Scale bar=10μm. **b:** Averaged nuclear miniCic intensity (red) and FRET signal (purple) in the neighbours of dying cells. The curves were aligned on the peak of the FRET signal. Light areas are s.e.m.. Note that miniCic nuclear exclusion correlates well with the pulses of FRET activity (although there is a delay in the relocation of miniCic to the nucleus). **c:** Snapshots of local projections in the larval accessory cells (LECs) of the pupal abdomen expressing E-cad-GFP and miniCic. Blue stars show an extruding cell, blue arrows show the nuclei of neighbouring cells (which lose transiently miniCic signal). Dotted lines show cell contours. Scale bar=10μm. **d:** Averaged nuclear miniCic signal in the LECs neighbouring extruding cells (Time 0 is the termination of extrusion). Light areas are s.e.m.. **e:** Snapshots of a local projection of a pupal notum with *UAS-EGFR dsRNA* clones (blue, UAS-His3-mIFP) expressing E-cad-GFP and miniCic. White stars show two EGFR RNAi cells extruding. Blue lines show the contour and the nuclei of two EGFR RNAi cells neighbouring the dying cells, white lines show the contours and nuclei of WT cells neighbouring the dying cells. Note that dying EGFR depleted cells can still activate ERK in the WT neighbours (loss of nuclear miniCic) but ERK does not get activated in EGFR depleted cells. Scale bar=10μm.

a

GCaMP
E-cad-tdTomato

-30 min   -20 min   -10 min   0 min   10 min   Ca²⁺ pulse

high Ca²⁺

low Ca²⁺

b

n=58 neighbours of 11 extruding cells

GCaMP (normalised intensity)

Time (min)

c

E-cad
miniCic
His3-mIFP (SpitzdsRNA clones)

-60 min   -25 min   +10 min   +45 min

miniCic

ERK activation

d

n=6 extruding cells, >30 neighbours

miniCic normalised intensity

time (min)

e

E-cad   row1

miniCic   row1 row2

His3-mIFP

0-   0+   25 min   50 min

*Laser burst at time 0

f

n=21 ablations, 7 pupae

miniCic normalised intensity (after bkg removal)

row 2

row 1

time (min)

Figure S4: ERK pulses do not correlate with Ca^{2+} pulses, do not require Spitz expression in the dying cells, and can be mimicked by wound healing-induced stretching.

a: Snapshots of local projections of a pupal notum expressing the Calcium sensor GCaMP (pseudo colour) and E-cad-tdTomato. The white stars show a dying cell. There is no visible change of Ca^{2+} activity in the neighbours or in the dying cell. The right inset show an example of rare spontaneous pulse of Ca^{2+} (one or two per movie, no correlation with cell death). Scale bar=10μm. **b:** Averaged and normalised GCaMP signal in cells neighbouring extruding cells (Time 0, termination of extrusion). Light area is s.e.m.. **c:** Snapshots of a local projection of a pupal notum with *UAS-spitz dsRNA* clones (blue, UAS-His3-mIFP) expressing E-cad-GFP and miniCic. The stars show a dying Spitz RNAi cell, white lines show the contour of the neighbours. Note that the Spitz RNAi effect was validated using the pnr-gal4 driver (nota fusion defects and reduction of ERK assessed with miniCic, not shown). Scale bar=10μm. **d:** Averaged and normalised miniCic nuclear intensity in cells neighbouring Spitz RNAi extruding cells (Time 0, termination of extrusion). Error bars are s.e.m.. **e:** Snapshots of local projection of a pupae expressing E-cad-GFP, miniCic and His3-mIFP. The cell marked with a red star is laser ablated at time 0. Pink line/area show the first neighbours, black line the second row of neighbours. The white circles show miniCic signal in the direct neighbours of the laser ablated cell. Scale bar=10μm. **f:** Averaged and normalised nuclear miniCic signal (after background removal, acquisition on a spinning disc) in the first row (red) and second row (black) of cells neighbouring the laser ablated cells. Time 0 is the ablation time.

a

b

Number of caspase resistant neighbours

Figure S5: ERK pulses can modulate significantly the rate of cell death and could recapitulate features of cell competition

a: 2D simulations of the distribution of cell death assuming a local refractory phase of various times (see **Methods**) and different "intrinsic" rate of cell elimination (elimination rate without refractory phase, from bottom to top curves, λ=0.05, 0.1, 0.2, 0.3, 0.6, 1 x10^{-2} death.min^{-1}). For a refractory phase of 60 minutes (red doted line) the curve showing a rate of cell elimination similar to the one measured in the posterior region of the pupal notum (~0.0013/minute) corresponds to an intrinsic rate of cell death λ=0.002. **b:** Estimation of the effective cell death rate in a cell (grey) neighbouring various number of caspase resistant cells (0 to 6, green cells) with different "intrinsic" rate of cell elimination (from bottom to top, λ=0.05, 0.1, 0.2, 0.3, 0.6, 1 x10^{-2} death.min^{-1}). See **Methods**. In a region with a high rate of cell elimination, clones resistant for apoptosis will increase the rate of neighbouring cell elimination, especially for cells surrounded by several apoptosis resistant neighbours (similar to a cell competition scenario).

Video legends

Video S1: Induction of clone elimination by optoDronc is effector caspase-dependent

Local projections of pupal nota expressing E-cad-tdTomato (green) and optoDronc (magenta, left) or optoDronc and p35 (magenta, right) in clones. Blue light exposure starts at the onset of the movies. Scale bars=10μm.

Video S2: Induction of extrusion by optoDronc in different clone topologies

Local projections of pupal nota expressing E-cad-tdTomato (green) and optoDronc (magenta) in clones of various sizes (one cell, two cells, three cells in raw, three cells in cluster, more than four cells in cluster). Blue light exposure starts at the onset of the movies. Scale bars=10μm.

Video S3: optoDronc clone elimination and Dextran diffusion

Local projections of pupal nota expressing E-cad-tdTomato (green, middle) and optoDronc (magenta) in clones upon injection of far red Dextran (grey levels, same plane as E-cad, right). Top part shows a line of four cells, bottom a cluster of four cells with aberrant extrusion. Blue light exposure starts at the onset of the movies. Scale bars=10μm.

Video S4: Cell extrusion localisation in a WT notum

Local projection of a pupal notum marked with E-cad-GFP showing every cell extrusion (coloured circles). Anterior: left, Posterior: right. Circles appear at the termination of extrusion (orange) and stay for nine frames (progressively becoming blue). Scale bar=10μm.

Video S5: Reversion of caspase activity in a cell neighbouring an extruding cell

Local projection of a pupal notum expressing GC3Ai (green and right, effector caspase activity) and E-cad-tdTomato (magenta). The bottom cell dies, while GC3Ai signal stop increasing in the top cell which remains in the tissue. Scale bar=10μm.

Video S6: Transient activation of ERK in cells neighbouring extruding cells

Local projection of a pupal notum expressing miniCic-mScarlet (magenta, bottom right, ERK sensor), E-cad-GFP (green, top right) and His3-mIFP (grey, bottom left). Cells marked in cyan are extruding. The white lines show the nuclei of the direct neighbouring cells (transient loss of nuclear miniCic= ERK activation). Scale bar=10μm.

Video S7: Transient ERK activation is also visible with a FRET sensor

Local projection of a pupal notum expressing miniCic-mScarlet (grey, left), and EKARnls (ERK FRET sensor, FRET ratio in pseudocolour, red=activation, blue=inhibition). The green circles show a dying cell. Note the transient loss of nuclear miniCic in the direct neighbours and the transient burst of FRET signal. Scale bar=10μm.

Video S8: There is no visible Ca^{2+} pulses near extruding cells

Local projection of a pupal notum expressing GCaMP (green, right), and E-cad-tdTomato (magenta). The cyan contours show an extruding cell. Scale bar=10μm.

Video S9: Spitz is not required in the dying cell to activate ERK in the neighbours

Local projection of a pupal notum expressing miniCic-mScarlet (magenta, bottom right, ERK sensor), E-cad-GFP (green, top right) and His3-mIFP in clones expressing Spitz-dsRNA (grey, bottom left). The cell in cyan is a Spitz RNAi cell dying. The white circles show the nuclei of direct neighbouring WT cells (transient loss of nuclear miniCic = ERK activation). Scale bar=10μm.

Video S10: Activation of ERK driven by single cell laser ablation

Local projection of a pupal notum expressing miniCic-mScarlet (magenta, bottom right, ERK sensor), E-cad-GFP (green, top right) and His3-mIFP (grey, bottom left). The cell in cyan is ablated by a UV pulsed laser at time 0. The white circles show the nuclei of direct neighbouring cells (transient loss of nuclear miniCic = ERK activation). Scale bar=10μm.

Video S11: EGFR is required for extrusion-induced ERK activation

Local projection of a pupal notum depleted for EGFR (*pnr-gal4, UAS-egfrdsRNA*) expressing miniCic-mScarlet (red, bottom left, ERK sensor), E-cad-GFP (green, top right)

and His3-mIFP (blue, bottom right). White contours show the dying cell. The white circles show the nuclei of direct neighbouring cells (no change of nuclear miniCic, no ERK activation). Scale bar=10μm.

Video S12: ERK is also activated by cell extrusion in the pupal abdomen

Local projection of larval accessory cells in the pupal abdomen expressing miniCic-mScarlet (magenta, grey on the right, ERK sensor) and E-cad-GFP (green). The red contours show the dying cell. The blue contours show the direct neighbours. Scale bar=10μm.

Video S13: ERK pulses precede effector caspase inhibition

Local projection of a pupal notum expressing GC3Ai (green and right, effector caspase activity) and miniCic (magenta, middle). Green circles track the nucleus of the first dying cell. Purple circles track the nucleus of a neighbour undergoing transient caspase inhibition (slow-down of GC3Ai signal increase). Scale bar=10μm.

Video S14: EGFR depletion abolishes caspase reversions

Local projection of a pupal notum depleted for EGFR (*pnr-gal4, UAS-egfrdsRNA*) expressing GC3Ai (green and right, effector caspase activity) and E-cad-tdTomato (magenta). Coloured contours show neighbouring cells activating caspase. Scale bar=10μm.

Video S15: Cell extrusion localisation in a EGFR RNAi notum

Local projection of a pupal notum depleted for EGFR (*pnr-gal4, UAS-egfrdsRNA*) marked with E-cad-GFP showing every cell extrusion (coloured circles). Anterior: left, Posterior: right. Circles appear at the termination of extrusion (orange) and stay for nine frames (progressively becoming blue). Scale bar=10μm.

Video S16: Clusters of cell elimination and aberrant extrusions appear upon EGFR depletion

Local projection from different pupal nota depleted for EGFR (*pnr-gal4, UAS-egfrdsRNA*) marked with E-cad-GFP showing four examples of clusters of cell extrusion (cyan, more

than four cells) followed by aberrant extrusions (relaxation, wound healing and E-cad accumulation at vertices). Scale bar=10μm.