1 **Title:** Morphological and functional variability in central and subcentral motor cortex of the
2 human brain
3
4 **Abbreviated title**: Variability in central and subcentral motor cortex
5

6 **Authors:** Nicole Eichert[1,*], Kate E. Watkins[2], Rogier B. Mars[1,3] & Michael Petrides[4,5]
7
[1] Wellcome Centre for Integrative Neuroimaging, Centre for Functional MRI of the Brain
(FMRIB), Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of
Oxford, OX3 9DU, Oxford, United Kingdom
[2] Wellcome Centre for Integrative Neuroimaging, Department of Experimental Psychology,
University of Oxford, OX2 6GG, Oxford, United Kingdom
[3] Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, 6525 AJ
Nijmegen, The Netherlands
[4] Department of Neurology and Neurosurgery, Montreal Neurological Institute and Hospital,
McGill University, 3801 University Street, Montreal H3A 2B4, QC, Canada
[5] Department of Psychology, McGill University, 1205 Dr. Penfield Avenue, Montreal H3A 1B1,
QC, Canada
19
20 * **Corresponding author**: nicole.eichert@psy.ox.ac.uk

21
Number of pages: 40
Number of figures: 9
Number of tables: 3
Number of words for abstract: 161
Number of words for introduction: 646
Number of words for discussion: 1499
28

29 **Conflict of interest**: The authors declare no competing financial interests.
30
Acknowledgments:
N.E. is a Wellcome Trust Doctoral student in Neuroscience at the University of Oxford
[203730/Z/16/Z]. N.E. was supported by a Lazlo and Etelka Kollar Brain@McGill
Graduate/Postgraduate Travel Grant awarded by the Oxford McGill Zürich Partnership in the
Neurosciences. The project was supported by the NIHR Oxford Health Biomedical Research
Centre. The Wellcome Centre for Integrative Neuroimaging is supported by core funding from
the Wellcome Trust [203139/Z/16/Z]. The work of R.B.M. is supported by the Biotechnology
and Biological Sciences Research Council (BBSRC) UK [BB/N019814/1] and the Netherlands
Organization for Scientific Research NWO [452-13-015]. The work of M.P is supported by CIHR
Foundation Grant FDN-143212.

<u>Abstract</u>

There is a long-established link between anatomy and function in the somatomotor system in

the mammalian cerebral cortex. Morphology of the central sulcus predicts the location of

functional activation peaks in individuals, but morphological variation in the subcentral region

and its relationship to functional activation is unknown. Investigating the subcentral region is

particularly important in the context of speech, since control of the larynx during human

speech production activates this region. Here, we examined whether morphological variation

in the central and subcentral region is related to functional activation during movement of the

hand, lips, tongue, and larynx at the individual subject level. We provide a systematic

description of the sulcal pattern of the subcentral and adjacent opercular cortex, including the

inter-individual variability of sulcal morphology. We found a robust relationship between

morphology of the central and subcentral sulcal segments and movement of different

effectors. A surface registration based on sulci revealed that anatomical variability explains, in

part, spatial variability in function.

<u>Significance Statement</u>

There is a long established relationship between structure and function in the somatomotor

system in the mammalian brain. Here, we show that the location of brain activations during

movement involving different effectors relates to morphological landmarks in the central

sulcus and the subcentral region of the cerebral cortex. We provide a systematic description

of the morphological patterns of the subcentral cortical region and the inter-individual

anatomical variability of sulcal segments. We discuss how structural variability can explain

spatial variability in functional activations, which is a critical factor for cross-subject

registrations in group studies.

Introduction

A fundamental challenge in neuroscience is to establish meaningful links between brain

function and structure. One of the clearest cases of such a structure-to-function relationship

is found in the somatomotor system in the central strip of the cerebral cortex (Jackson, 1863;

Hitzig and Fritsch, 1870; Vogt and Vogt, 1919; Penfield and Boldrey, 1937). Different parts of

the body are represented in an orderly and consistent fashion, following a somatotopic

organization, the schematic visualization of which is known as Penfield's 'homunculus'.

In contrast to electrical brain mapping studies, neuroimaging studies typically report average

activations from larger groups. This approach demonstrates effects, which are representative

of the human brain in general, but obscures subject-specific features and inter-individual

variability, limiting sensitivity and functional resolution (Bennett and Miller, 2010; Nieto-

Castañón and Fedorenko, 2012; Woo et al., 2014). Thus, a growing number of neuroimaging

studies now tend to report activation patterns in individuals (Miller et al., 2002; Barch et al.,

2013; Carey et al., 2017). A deeper understanding of structure-to-function relationships,

however, requires examination of inter-individual functional variability together with

anatomical variability.

Several studies have established a coupling between the sulcal/gyral brain anatomy and

functional activation in individuals (e.g., Amiez et al., 2006; Derrfuss et al., 2012; Amiez et al.,

2013; Zlatkina et al., 2016; Bodin et al., 2018). A recent neuroimaging study demonstrated that

individual fMRI activation peaks showed a consistent relationship between the effector being

moved and the morphologically defined segments of the central sulcus (Germann et al., 2019).

The inter-individual variability of the central sulcus segments, however, was not quantified and

this investigation focused exclusively on the central sulcus.

Movement representation in the human brain, however, is not limited to the central sulcus.

The subcentral gyrus, in addition to the ventral region of the central sensorimotor strip, is

involved in speech-related movements (Penfield and Boldrey, 1937; Olthoff et al., 2008;

Grabski et al., 2012; Bouchard et al., 2013). Voluntary control of laryngeal movements during

vocalization evokes brain activity in two distinct regions in lateral motor cortex: a dorsal region

close to the representation of the lips and a ventral region close to the lateral fissure (reviewed

in: Belyk and Brown, 2017; Eichert et al., 2020a). The representation of the larynx in the human

brain, however, remains controversial and inconsistent reports in the literature might relate to

inter-individual variability. Examining functional anatomy of the speech motor system thus

requires investigation both of the central sulcus and the subcentral gyral region at an individual

subject level.

There has been no systematic examination of the morphological variability in the subcentral

gyrus and the adjacent opercular cortex. There are two distinct sulci in the subcentral region:

the anterior subcentral sulcus (*ascs*) and the posterior subcentral sulcus (*pscs*) (Petrides, 2012,

2019). The variability of these two sulci in relation to the central sulcus segments across

individuals requires examination, as well as the relationship of morphological variability to

functional brain activity during movement.

Here, we assessed the structure-to-function relationship in the human motor system on an

individual subject level by examining the morphology of the central sulcus, the subcentral gyrus

and the adjacent central operculum. We performed sulcal labelling in surface space, rather

than in volume space, which is a suitable representation of the intrinsic topology of the

cerebral cortex (Fischl et al., 1999). To visualize and quantify inter-individual variability of the

examined sulcal segments, we derived spatial probability maps in both standard surface and

volume space. In a subset of subjects, we acquired fMRI data to localize brain activity during

movement of the hand, lips and tongue, as well as the larynx during vocalization (Eichert et al.,

2020a). To investigate how this structure-to-function relationship can help to improve

alignment of data, we registered all subjects based on the sulcal surface labels. A decrease in

distances across activation peaks after applying the registration indicates that anatomical

variability can explain some of the functional variability.

Material and Methods

Subjects

Data from two groups of subjects were used. In one study group, both structural and functional

MRI data were acquired; in the other study group, only structural MRI data were available.

Structural and functional MRI data from the performance of motor tasks were acquired from

20 subjects (12 females, 18 – 40 years, 5 left-handers). All subjects were self-reported native

English speakers (two were raised bilingually from infancy and three were fluent in a second

acquired language) and had no history or diagnosis of speech disorders. All had normal hearing,

normal or corrected-to-normal vision, and no neurological impairments. The subjects were

part of a study that had been approved by the Central University Research Ethics Committee

of the University of Oxford (CUREC, R55787/RE001) in accordance with the regulatory

standards of the Code of Ethics of the World Medical Association (Declaration of Helsinki). All

subjects gave informed consent for their participation and were monetarily compensated for

their participation.

In addition, we used cortical brain surface reconstructions from 30 subjects provided by the

Human Connectome Project (HCP), WU-Minn Consortium (Principal Investigators: David Van

Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that

support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for

Systems Neuroscience at Washington University (Van Essen et al., 2013). The minimally pre-

processed datasets of the first 31 subjects (16 female, age range 22-35 years) of the Q2 release

were used. One subject was excluded because of a technical problem in the automatic

FreeSurfer parcellation.

MRI Data Acquisition

MRI data acquisition parameters differed for the two groups of subjects. Data from the

subjects that took part in the functional study were obtained at the Oxford Centre for Human

Brain Activity (OHBA) using a 3T Siemens Prisma scanner with a 32-channel head coil. Two

structural images of the whole brain had been acquired at 1 mm isotropic resolution; a T1w

image (MPRAGE sequence) and a T2w image (SPACE sequence). For task-fMRI, whole head

T2*-weighted echo planar images were acquired at 2.4 mm^3 isotropic resolution (TE = 30 ms,

multiband fact 6, TR = 0.8 s, Casey et al., 2018).

Data acquisition and preprocessing methods of the HCP subjects are detailed in Glasser et al.

(2013) and Uğurbil et al. (2013). T1w images had been acquired using an MPRAGE sequence

at 0.7 mm isotropic resolution.

Experimental Design

The 20 subjects who provided structural and functional data took part in an fMRI study on

speech production and laryngeal motor control. The experimental design, processing, and fMRI

results of this study have been reported elsewhere in detail and are here only briefly described

(Eichert et al., 2020a). In a functional localizer task, subjects were asked to perform repeated

lip protrusion or tongue retraction at a rate of approximately 1-2 reps/s. The subject's

breathing pattern was explicitly controlled using the fixation symbol on the screen, instructing

them to inhale for 1.5 s and exhale for 4 s. A 'breathing only' condition, during which the

subjects followed the same breathing pattern, was acquired as baseline condition. Each task

condition was performed in blocks lasting 22 s followed by a rest period of 8 s with normal

breathing. The conditions were presented in a fixed pseudo-random order following a

balanced Latin-square design wherein each condition was repeated four times.

In a separate task, subjects were instructed to produce a syllable sequence (/la leɪ li la leɪ li/)

under four different conditions: overt speech, silent mouthing, only vowel production and

covert speech. Breathing instructions, task timing and randomization of the four blocks were

the same as described for the localizer task, except that each condition was repeated five

times. In a third task, subjects performed a phonological and semantic judgement task, which

involved button presses with the right index finger to indicate responses. This task was

analyzed as a localizer for the hand region in the left hemisphere.

Structural MRI Data Analysis

Data of the subjects who took part in the functional study were pre-processed using the HCP-

pipeline Glasser et al. (2013). The automatic processing pipeline includes cortical surface

reconstruction using FreeSurfer based on the contrast from the T1w and the T2w images and

automatic assignment of neuroanatomical labels. Cortical surface reconstructions of the HCP

subjects were derived using FreeSurfer based on the T1w scans and directly provided by the

database. A linear transformation (12 degrees of freedom) from FreeSurfer's anatomical to

standard MNI space (nonlinear 6th generation atlas, Fonov et al., 2011) was derived using FSL's

FLIRT (Jenkinson and Smith, 2001; Jenkinson et al., 2002).

Functional MRI Data Analysis and Statistical Analysis

Functional MRI data processing was carried out using FEAT (FMRI Expert Analysis Tool) Version

6.00, part of FSL (FMRIB's Software Library, www.fmrib.ox.ac.uk/fsl) including motion

correction of the images and unwarping using a fieldmap (Jenkinson, 2003). Time-series

statistical analysis was based on a general linear model (GLM) implemented in FILM with local

autocorrelation correction (Woolrich et al., 2001). Standard motion correction parameters and

individual volumes that were motion outliers, determined using fsl_motion_outliers, were

included as separate regressors at the first level for each subject. Registration to the high-

resolution structural scan and standard 2-mm MNI template was carried out using FLIRT.

Registration from high resolution structural to MNI space was then further refined using FNIRT

nonlinear registration (Andersson et al., 2007).

In the functional localizer task for lip and tongue movement, activity during each condition was

assessed relative to the 'breathing only' condition. For the syllable production task, the

conditions were analyzed in a factorial model that allowed separation of the (supralaryngeal)

articulation and the (laryngeal) vocalization components of the task. We refer to the latter as

'vocalization' component (index for laryngeal activity during voice production), while other

studies have referred to it as voicing or phonation. Brain activity associated with the control of

supralaryngeal articulation was defined as ('overt speech' *minus* 'vowel production') *plus*

('silent mouthing' *minus* 'covert speech') and the main contrast for vocalization was derived

by the contrast ('overt speech' *minus* 'silent mouthing') *plus* ('vowel production' *minus* 'covert

speech'). In both tasks described above, the rest blocks with normal breathing served as

baseline, which means that they were not modelled in the GLM. For the hand localizer task,

we derived a contrast of all conditions involving button presses relative to a resting baseline.

Individual Surface Activation Maxima

Activation maxima were derived using the steps described in Eichert et al. (2020a), which are

repeated here. To assess intra-individual variability of the fMRI results, we derived the location

of individual activation maxima for hand, lip, and tongue movement, and larynx activity during

vocalization. Activation maxima were derived in both hemispheres in volume space using

anatomically defined ROIs. For hand (left hemisphere only), lip, and tongue movement, we

used the central sulcus as a volumetric ROI to extract the maximum. The ROI was defined using

FreeSurfer's automatic volumetric labelling based on the Destrieux Atlas.

For the vocalization contrast from the syllable production task, which indicates laryngeal

activity during voice production, we derived two separate activation maxima: one located in a

dorsal area and one in a ventral area of the central sulcus. To account for additional articulation

of the tongue during the syllable production task, a spherical ROI (7 voxels diameter) around

each individual's maximum voxel from the tongue movement localizer was masked out from

the z-statistics image of the vocalization contrast prior to extracting the maximum.

For the dorsal larynx ROI, we used a dorsally and ventrally cropped section of the central sulcus

in volume space (coordinates in MNI space: $z = 50$ to $z = 30$). The ventral larynx ROI was derived

manually based on individual anatomy in surface space to ensure that the ROI did not overlap

with unrelated brain areas such as the subjacent auditory cortex in the temporal lobe or

inferior frontal cortex located anteriorly. A liberal surface ROI was drawn on the individual's

midthickness surface covering the ventral part of the central sulcus and adjacent gyri.

Anteriorly, the ROI was delineated by the inferior portion of the precentral sulcus and

posteriorly the ROI spanned the postcentral gyrus. If present, the lateral portion of the

ascending sulcus in subcentral gyrus was included within the ROI. The dorsal limit of the ROI

was defined by a horizontal line across the gyrus at the level of the usual location of the

posterior ramus of the inferior precentral sulcus. The ventral larynx surface ROI was converted

into a volumetric ROI covering the underlying cortical ribbon using Workbench Command

(wb_command, www.humanconnectome.org/software/connectome-workbench.html).

Individual volumetric ROIs were linearly transformed from FreeSurfer's anatomical to

functional space of the respective task fMRI scan. Within the ROI, the voxel of maximal

intensity was determined from the z-statistics image. The activation maxima were mapped to

the individual's native midthickness surface, then smoothed (FWHM = 1 mm) and binarized to

form a small circular patch. To assess spatial variability of the functional activity, all subjects'

activation peaks were resampled to the same regular 32k surface mesh, prior to any

registration.

Sulcal Labelling

We identified the following sulci and sulcal segments in the structural data from all 50 subjects:

five segments of the central sulcus from dorsal to ventral (*cs_1* to *cs_5*), the lateral and

opercular segments of the anterior subcentral sulcus (*ascs_lat, ascs_op*) and the posterior

subcentral sulcus (*pscs*). Sulcal labels for one example subject are shown in Figure 1. Sulcal

labels were drawn manually onto the native surface mesh (approximately 136,000 vertices) in

Connectome Workbench's wb_view (www.humanconnectome.org/software/connectome-

workbench.html). Surface features of both pial and white matter surface were inspected in

conjunction with the subject's T1w scan. The identification of central sulcus segments was

based on changes in direction of the sulcus and on gyral 'plis de passage', which are small gyral

bridges connecting the postcentral with the precentral gyrus. These gyral bridges can be most

easily identified on the white matter surface (Germann et al., 2019).

Figure 1: Sulcal labelling. **A**: Sulcal labels in one individual displayed onto the native pial surface, white matter surface, and inflated surface. **B**: Sulcal labels of the same subject in MNI volume space.

Cs_1 is the most dorsal segment of the central sulcus, which runs more or less in a vertical straight direction. Its ventral boundary was drawn at the location where a gyral bridge forms a prominent landmark on the posterior bank of the central sulcus. *Cs_2* has a characteristic curvature in the shape of the Greek omega letter (see Figure 1B), which is known as the 'hand knob' (Yousry et al., 1997). This knob is often more pronounced in the left hemisphere and it can comprise two smaller knob-like curves instead of one. The boundary between *cs_2* and *cs_3* was drawn at the location where the central sulcus changes direction and where a gyral passage can be observed on the posterior bank. In some subjects, an additional convexity of

the central sulcus can be observed on the posterior bank in the middle of *cs_3*. The ventral

boundary of *cs_3* was drawn ventral to this convexity, at the location where a small gyral bridge

forms a landmark on the anterior bank of the central sulcus. The last two segments of the

central sulcus, *cs_4* and *cs_5* are smaller in extent, shallower and more variable in their

morphology. The boundary between *cs_4* and *cs_5* was defined based on a gyral bridge on the

posterior bank of the central sulcus. *Cs_5* is the most ventral part of the central sulcus, which

can form an additional curve or run in straight direction.

The labels for *ascs* and *pscs* were assigned based on an atlas of human brain morphology

(Petrides, 2019). For *ascs*, we labelled two distinct segments: a lateral and an opercular

segment (*ascs_lat, ascs_op*). The course of *ascs_lat*, *ascs_op* and *pscs* was found to be highly

variable and a detailed description of the sulcal anatomy in the subcentral region is reported

in the results section.

The morphological patterns of the ventral subcentral region were categorized into five types

depending on the configuration of *ascs_lat*. The classification was based on the location of

ascs_lat on the cortex and its spatial relation to other sulci. Sulcal segments were considered

as 'merged', when there was a clear continuation on the pial surface, although, in some cases,

a discontinuity between the merged sulci was still observed on the white matter surface.

Spatial Probability Maps

In order to characterize the inter-individual morphological variability of the labelled sulci, we

generated probability maps in surface and volume space. To obtain surface probability maps,

all surface labels were resampled from native to a common regular 164k mesh. At each vertex,

the 50 binary surface maps were summed and then normalized to create a surface label with

intensities ranging from 0 % to 100 % at the maximal possible overlap of all 50 subjects. For

visualization, the surface probability maps were displayed onto the average of all subjects'

inflated surfaces.

For the generation of volumetric probability maps, we first deformed the subjects' native pial

surface to MNI space by applying a linear transformation (12 degrees of freedom). Then we

mapped the surface labels from the pial surface to 0.5 mm resolution volume space using

wb_command. The individual volumetric labels were smoothed (Gaussian kernel with FWHM

of 2 mm), thresholded at 0.1, binarized and then summed at each voxel. For visualization,

volume probability maps were overlaid onto the 0.5 mm MNI average brain.

Structure-to-Function Relationship

Next, we examined the spatial relationship between the sulcal segments and the task

activation peaks at the individual subject level. This analysis was performed in the 20 subjects

who contributed both task and structural data. Individual task activation peaks were mapped

onto the individual's cortical surface as described above for sulcal segments. As Extended Data

(Figure 6-1), we show sulcal labels and activation peaks for all subjects, which allows an

assessment of the spatial relationships at the individual subject level.

In order to characterize the structure-to-function relationship at the group level, we aligned all

individual surfaces based on the anatomical surface labels and then we applied the same

registration to individual task activation peaks. This approach allowed us to visualize individual

variability in the spatial distribution of task activation peaks in a common group-level space.

The registration of sulcal labels was driven by the binary labels for *cs_1*, *cs_2*, *cs_3*, *cs_4*, *cs_5*,

ascs_lat, ascs_op and *pscs* and performed using multimodal surface matching (MSM, Robinson

et al. (2014). As target, or reference, for the MSM-based registration, we used the normalized

and thresholded (> 0.4) average labels after projecting all of them to the same regular sphere

(approximately 32,000 vertices). Each subject's sulcal maps and the reference sulcal maps

were merged into a combined file with six data arrays. Then, we derived a registration using

MSM for each subject. We refer to the average of the transformed labels resulting from the

MSM registration as 'group-level segments'.

To characterize the structure-to-function relationship, we quantified the effect of the

registration on the spatial variability of the activation peaks. The spatial variability of peaks in

each task cluster before and after applying the registration was quantified by computing the

median geodesic distances across all 20 activation peaks for each effector on a regular sphere.

We compared this measure to a baseline measure of spatial variability prior to any anatomical

registration, where the functional peaks were resampled to a common sphere. We also derived

two additional anatomical registrations using the same steps as described above: one using

the identical parameters, but based on four FreeSurfer labels (precentral gyrus, central sulcus,

postcentral gyrus, subcentral gyrus/sulcus) and one based on sulcal depth whole-brain maps

using MSM default settings. FreeSurfer labels and sulcal depth maps were provided by the

HCP-processing pipeline.

<u>Results</u>

Sulcal Pattern of the Subcentral Region

The sulcal anatomy of the subcentral region was examined in a total of 100 hemispheres ($n =$

50 brains) and the morphological patterns encountered were classified into five types (Figure

3). The most commonly observed configuration of subcentral sulci (Type 1, 44 % of

hemispheres) is shown in Figure 2. The example in Figure 1 also shows an individual

hemisphere classified as Type 1. In the Type 1 configuration, the central sulcus does not extend

ventrally to meet the lateral fissure. The fifth and most ventral segment of the central sulcus

(*cs_5*) is typically short and forms a curve in an anterior direction. The lateral segment of *ascs*

(*ascs_lat*) is visible on the lateral surface of the brain and is clearly separate from the central

sulcus. Dorsally, the *ascs_lat* extends into the precentral gyrus with variable length. Ventrally,

the *ascs_lat* continues into the opercular cortex, which is hidden within the lateral fissure of

the brain. A small gyral bridge separates the *ascs_lat* from the opercular segment of *ascs*

(*ascs_op*), which continues medially until it reaches the circular insular sulcus (*cris*). The medial

origin of the *ascs_op* can be identified at a curve of *cris*, which is formed by the posterior short

insular gyrus just anterior to the central insular sulcus (*cis*). Posterior to *ascs_op*, the posterior

subcentral gyrus (*pscs*) can be found, which extends laterally towards the lateral fissure. The

medial origin of *pscs* can be identified at a curve of *cris* posterior to *cis*, which is formed by the

anterior long insular gyrus. The number of curvatures of *cris* and the number of subcentral

sulci is variable, but the *ascs_op* and the *pscs* could be reliably identified in every subject

examined. The morphology of all of the mentioned sulci is variable across subjects, but the

description of morphological sub-types below focusses on the configuration of the *ascs_lat*.

Figure 2: Sulcal pattern of the subcentral region. **Left**: Sulcal map of the human cerebral cortex

(Petrides, 2012). **Right**: Type 1 configuration of sulci in the subcentral region. Anatomical variability was

assessed in the sulci marked with colour: *ascs_lat*, lateral segment of the anterior subcentral sulcus;

ascs_op, opercular segment of the anterior subcentral sulcus; *cs_1* to *5*, segments of the central sulcus

(only *cs_4* and *cs_5* are shown in the highlighted region); *pscs*, posterior subcentral sulcus. Grey dashed

line: visible surface boundary between lateral and opercular cortex. *aalf*, ascending anterior ramus of

the lateral fissure; *aplf*, ascending posterior ramus of the lateral fissure; *cis*, central insular sulcus; *cris*,

circular insular sulcus; *ds*, diagonal sulcus; *iprs*, inferior precentral sulcus.

Different Types of Morphological Patterns

411 The subcentral region exhibits high inter-individual variability regarding the configuration of

412 the *ascs_lat*. The observed occurrences of morphological types are reported in Table 1. In

413 addition to the canonical (Type 1, 44 % of hemispheres) configuration described above, it was

414 commonly observed that the *ascs_lat* merged with the central sulcus (Type 2, 20 %) (Figure 2).

415 In these cases, the *ascs_lat* remains mostly on the lateral brain surface and does not reach into

416 the opercular cortex. The *ascs_op* extends further lateral and its extension is visible on the

417 lateral brain surface. Another commonly observed sub-type was identified by a more opercular

418 position of the *ascs_lat* (Type 3, 18 %). In Type 3, the *ascs_lat* is less visible on the lateral

419 surface and the central sulcus extends further ventral towards the lateral fissure. Type 3 is also

420 characterized by a reduced gyral bridge separating the *ascs_lat* from the *ascs_op*. In six cases,

421 the opercular continuation of the the *ascs_lat* curved posteriorly and merged with *pscs* instead

422 of the *ascs_op* (not shown as separate type). In several cases, we also observed the *ascs_lat*

423 to merge with the inferior ramus of the inferior precentral sulcus (*iprs-i*) (Type 4). When

424 *ascs_lat* and *iprs-i* merge, the ventral continuation of *ascs_lat* curves in an anterior rather than

425 a posterior direction and the position of *ascs_lat* is more lateral than opercular. In a few cases

426 (Type 5) the position of the *ascs_lat* was notably further rostral so that it was positioned

427 anterior to *ascs_op* rather than posterior as in the other configurations.

429 Strong hemispheric differences were observed in the occurrences of the morphological types

430 (Table 1). Type 2 (*ascs_lat* merged with *cs_5*) is much more common in left hemispheres (85

431 % vs. 15 %), while Type 4 (*ascs_lat* merged with *iprs*) and Type 5 (*ascs_lat* further anterior),

432 are more common in right hemispheres (75 % vs. 25 % for Type 4; 83 % vs. 17 % for Type 5),

433 although both types are infrequent. For Type 1 (canonical configuration) and Type 3 (*ascs_lat*

434 merged with *ascs_op*) no pronounced hemispheric differences were observed.

Figure 3: Morphological patterns. Schematic drawings illustrating the main morphological patterns of the subcentral region formed by the *ascs_lat* with neighbouring sulci. Type 1: Canonical configuration observed in the majority of hemispheres (also shown in Figure 2). Type 2: *ascs_lat* merged with the central sulcus. Type 3: *ascs_lat* merged with *ascs_op*. Type 4: *ascs_lat* merged with the inferior ramus of *iprs*. Type 5: *ascs_lat* anterior to *ascs_op*.

Table 1: Morphological patterns. Observed frequencies of morphological patterns based on 100 examined hemispheres (LH – left hemisphere; RH – right hemisphere).

	Type 1	Type 2	Type 3	Type 4	Type 5
LH	20	17	9	3	1

RH	24	3	9	9	5
Total	44	20	18	12	6

Spatial Probability Maps

The morphological variability of the central sulcus segments and the subcentral sulci was

quantified and visualized using spatial probability maps in 2D surface space and 3D volumetric

space (Figure 4, Figure 5). For both surface and volume probability maps, intensity values

decrease from the center of the map towards the edges, which is typical for overlap maps.

Maximal and median values of the probability maps are reported as Extended Data (Figure 4-

1). Values in the volumetric probability maps are overall lower given that they capture

variability in three spatial dimensions. The pattern of values across sulci and hemispheres,

however, is consistent across surface and volumetric probability maps. For all sulci,

hemispheric differences were observed with regard to the location in volume space (Figure 5).

All sulcal segments in the left hemisphere are located consistently further posterior compared

to the segments on the right hemisphere, which is in line with the Yakovlevian anticlockwise

torque of the two hemispheres (see Table 2 for coordinates of the voxel of maximal probability

and for the center of gravity for each label).

The first three segments of the central sulcus (*cs_1 - 3*) show high inter-individual spatial

consistency, which is characterized by high values, i.e. overlap, in the probability maps.

Consistency is lower for *cs_4* and *cs_5*. Consistency is also low for the subcentral sulci (*ascs_lat*,

ascs_op, *pscs*) with lowest consistency for *ascs_lat*, especially in the left hemisphere. The

hemispheric effect for the *ascs_lat* is evident in both surface and volumetric probability maps.

It was observed that the *ascs_lat* is larger in extent and deeper in the right hemisphere (see

for example Figure 5, slice *y* = -1). The low consistency in *ascs_lat* is in line with the variable

morphological subtypes that were described above. The two subcentral sulci, *ascs_op* and

pscs, are less consistent than the central sulcus segments, but more consistent than the

ascs_lat. *Ascs_op* and *cs_5* show a tendency for more variability in the right hemisphere, which

is the opposite pattern of lateralization in comparison with that for the *ascs_lat*.

Figure 4: Surface probability maps. Probability maps of central and subcentral sulcal labels shown on

an inflated average surface (*n* = 50). Labels for *cs_1* and *cs_2* are shown from a dorsal perspective.

Labels for *ascs_op* and *pscs* are shown from a ventral perspective with the temporal lobe removed for

better visibility of the opercular cortex. Intensity values show the amount of overlap with 100 %

indicating an overlap in all subjects.

Figure 5: Volumetric probability maps. Probability maps generated after linear registration to MNI space

overlaid onto the standard MNI template (*n* = 50). L – left hemisphere shown on right side of the image.

Intensity values show the amount of overlap across subjects. Color maps for the different sulci have

been matched in luminance so that darker colors indicate a lower overlap.

Table 2: Coordinates of probability maps. MNI-coordinates (*x, y, z*) of each sulcal segment's maximal

voxel and the center of gravity in the volumetric probability maps.

		maximum			center of gravity		
	sulcus	*x*	*y*	*z*	*x*	*y*	*z*
left	*cs_1*	-14	-30	69	-17	-30	-69

	cs_2	-36	-26	58	-36	-24	59
	cs_3	-38	-18	42	-47	-15	46
	cs_4	-54	-8	30	-55	-7	32
	cs_5	-58	-6	22	-60	-4	22
	ascs_lat	-60	2	12	-58	0	12
	ascs_op	-41	-4	14	-44	-3	10
	pscs	-39	-16	18	-48	-14	15
right	cs_1	16	-28	68	18	-28	69
	cs_2	38	-23	60	36	-21	58
	cs_3	44	-14	40	48	-12	46
	cs_4	58	-4	32	56	-5	32
	cs_5	62	-2	21	62	-2	22
	ascs_lat	58	1	10	58	2	12
	ascs_op	42	-3	14	44	-2	11
	pscs	41	-14	18	48	-13	16

Sulcal Registration and Structure-to-Function Relationship

We registered all individual surfaces based on the binary sulcal labels using MSM. We then

applied the registration to individual surface activation peaks from movement of different

effectors. By these means, we could visualize and assess the spatial relationship of functional

peaks to the sulcal labels at the group level. After registering the sulcal labels, we averaged all

transformed sulcal labels and thresholded ($t > 0.4$) the average labels to obtain the outlines of

group-level sulcal segments. Figure 6A shows the outlines of thresholded group-level sulci,

which demonstrates an orderly dorsal-ventral configuration of the sulcal segments.

Next, we applied the registration that was used to transform individual surface labels to the

group space to the functional activation peaks of different effectors in individual subjects.

Figure 6B-F shows the location of the resampled activation maxima in relation to the outlines

of the group-level sulcal segments. Given that the same registration was applied to individual

sulcal labels and the peaks, the location of resampled peaks with respect to the group-level

sulci reflects the original spatial relationship on the native surfaces of the individuals.

Overall, a strong correspondence between sulcal labels and functional peaks was observed.

The observed relationships between labels and peaks are reported in Table 3. The majority of

activation peaks from the hand localizer fall inside the center of *cs_2*. The dorsal peaks for

larynx activity are found on the anterior bank of *cs_3*. Peaks from the lip localizer also fall within

cs_3 and overlap with the location of the dorsal larynx peaks. Those for the lip, however, are

reliably located at the more ventral extent of *cs_3* whereas those for the dorsal larynx peaks

span the segment. Activation peaks from the tongue localizer fall inside *cs_4* with high

consistency. The ventral peaks for larynx activity are associated with *cs_5* and *ascs_lat* or with

the gyrus in between, but the relationship is less reliable than for the other functional peaks.

Figure 6: Structure-to-function relationships. **A**: Outlines of the averaged, normalized and thresholded

(> 0.4) sulcal maps after registration using MSM (i.e. the group-level sulci). **B-F**: Location of individual

activation peaks with respect to the sulcal maps after applying the sulcal registration (*n* = 20). Brighter

colors indicate overlapping peaks. Only the central strip of the brain is shown.

Table 3: Structure-to-function relationships. Observed relationships between functional activation

peaks and sulcal labels based on 40 examined hemispheres. The number of peaks for each effector are

given for each sulcal segment.

	cs_1	*cs_2*	*cs_3*	*cs_4*	*cs_5*	*ascs_lat*
hand (only LH)	0	16	4	0	0	0
dorsal larynx	0	2	37	1	0	0
lip	0	1	34	5	0	0
tongue	0	0	1	36	3	0

ventral larynx	0	0	0	2	24	14

To characterize the effect of the sulcal registration on the spatial variability of functional peaks,

we computed the median distance between all peaks for each effector before and after

applying the sulcal transformation, which is shown in the Extended Data (Figure 6-2). Spatial

variability decreased by 11 %, but not to the same extent for all effectors. Clusters in the left

hemisphere benefitted more from the registration, but an exception is the left lip cluster,

where values increased due to an outlier (see Figure 6D). A comparison with a registration

based on FreeSurfer's automatic labels and by sulcal depth demonstrated that all three

anatomical registrations decrease spatial variability when compared to baseline. Performance

of the three registrations is similar across all effectors, but spatial variability of the left ventral

larynx cluster decreased most using the sulcal label registration. Taken together, this

quantification indicates that a registration based on anatomical information decreases the

spatial variability of functional activation peaks. At the given sample size, however, the

reported differences do not reach statistical significance.

Discussion

We examined the morphological patterns and the inter-individual variability of the sulci in the

subcentral region of the cerebral cortex in the human brain and their relation to motor activity.

A robust relationship between morphological features of central and subcentral sulcal

segments and movement of different effectors at the individual subject level was

demonstrated. Cross-subject registration based on the individually drawn sulcal labels showed

that morphological variability explains functional variability in part.

Morphology of the Subcentral Cortex and the Central Operculum

The present study provided a systematic examination of the sulcal patterns of the subcentral

and adjacent opercular cortical region in the human brain, including inter-individual variability

of the sulcal morphology. In this region, one encounters an anterior and a posterior subcentral

sulcus (*ascs*, *pscs*) (Petrides, 2012, 2019). We showed that, in the majority of subjects, the *ascs*

consists of a distinct lateral (*ascs_lat*) and a distinct opercular segment (*ascs_op*).

The morphological patterns of the subcentral region were classified into five distinct subtypes,

based on the configuration of *ascs_lat*. In addition to the canonical configuration (Type 1,

Figure 2), the *ascs_lat* often merged with the central sulcus, so that the central sulcus complex

extended further ventral towards the lateral fissure (Type 2). In Type 3, the *ascs_lat* merges

with the inferior precentral sulcus (*iprs*). In Type 4, the position of the *ascs_lat* is notably more

opercular and it merges with the *ascs_op*, so that the *ascs_lat* and *ascs_op* form one

continuous sulcus. Type 5 is characterized by a more anterior position of the *ascs_lat*,

compared with its position in the other types. The variability in the configuration of the lateral

and opercular segments explains why the *ascs* appears as one continuous sulcus in a

volumetric average of structural brain images in MNI space (Petrides, 2019).

Sulcal Labelling in Surface Space

We labelled sulcal segments directly on the subject's native cortical surface, rather than in

volume space as traditionally done in several MRI-based labelling studies (Germann et al.,

2005, 2019; Zlatkina and Petrides, 2010; Amiez et al., 2013; Sprung-Much and Petrides, 2018,

2019). Surface space provides a parsimonious representation of the sheet-like geometry of the

cerebral cortex. We reproduced the segmentation of the central sulcus, as described in a

volumetric labeling study (Germann et al., 2019). The underlying morphological features are

easily visualized in surface space and the labelling is reproducible. Furthermore, using surface

labels allowed us to perform a surface-based registration, where distances are represented as

geodesic distances along the cortex rather than as Euclidean distances between voxels (Fischl

et al., 1999; Klein et al., 2010). In the current study, the labels were drawn manually, but future

research could develop automatic classification of sulcal segments based on supervised

learning algorithms (Clouchoux et al., 2006; Takerkart et al., 2015).

Spatial Probability Maps

We visualized the morphological variability and spatial extent of the central sulcus segments

and the subcentral sulci in surface and volumetric probability maps. The first three central

sulcus segments show the highest spatial consistency across subjects. Consistency decreases

for the fourth and even further for the most ventral (fifth) segment of the central sulcus, which

is in line with previous observations (Germann et al., 2019). The decrease in probability values

for *cs_4* and *cs_5* can also be attributed to their smaller spatial extent compared to the sulcal

labels for *cs_1-3*. Consistency for the subcentral sulci *ascs_lat*, *ascs_op* and *pscs* is also notably

reduced, which is in line with the inter-subject variability in the morphological patterns as

described above.

Hemispheric Differences

The classification of morphological subtypes and the spatial probability maps revealed

hemispheric differences in the subcentral cortex, indicating that the position of the *ascs_lat*

along the rostro-caudal axis differs between hemispheres, but the dorso-ventral position does

not. The *ascs_lat* in the left hemisphere tends to be located further posterior, which frequently

results in a merge with the central sulcus, while the *ascs_lat* in the right hemisphere is located

more anteriorly. The inter-individual variability as characterized in the spatial probability maps

showed that the *ascs_lat* is less variable and larger in extent in the right hemisphere.

These hemispheric differences can be interpreted in relation to sulcal variability of the ventro-

lateral cortex (Germann et al., 2005; Sprung-Much and Petrides, 2018, 2019). Language

processing is lateralized to the left hemisphere and the role of left inferior frontal cortex in

language function is widely established (Broca, 1861; Price, 2000; Vigneau et al., 2006; Hickok

et al., 2016). Functional language lateralization is associated with structural asymmetries

(Foundas et al., 1996; Josse and Tzourio-Mazoyer, 2004) and increased regional variability

(Croxson et al., 2018), but the structure-to-function relationships remain controversial

(Dorsaint-Pierre et al., 2006; Sprung-Much and Petrides, 2018). We presume that increased

surface area in the left inferior frontal cortex affects the neighboring sulci so that the *ascs_lat*

is 'pushed' to a relatively further posterior position in the left hemisphere. Despite the

hemispheric differences in morphology, the structure-to-function relationships for basic

movements of different effectors described below, did not exhibit notable differences

between hemispheres.

Functional Activation Peaks and Morphology

Here we demonstrated a tight link between individual morphological features of the cortex

and activation peaks for the different effectors examined. We replicated the relationships

between central sulcus segments and functional localizers for the hand, lip, tongue, and larynx

described in Germann et al. (2019). The functional localizer for larynx activity in the current

study differed substantially from the one used by Germann et al., where 'humming' was the

task instruction. Furthermore, we identified two larynx peaks in each subject, because more

recent studies suggest that two separate regions in the central motor strip correlate with larynx

activity (Belyk and Brown, 2017; Jarvis, 2019; Eichert et al., 2020a). The ventral 'humming' peak

in Germann et al. was associated with the fifth segment of the central sulcus. The current

analysis, however, showed that the ventral larynx peak was also localized in the *ascs_lat* in a

large portion of subjects. The difference across studies might be due to the significantly larger

sample size in the current study, which allowed a more robust assessment of the relationships.

It should be noted that the functional contribution of both larynx regions in motor control

during vocalization is still debated (Simonyan, 2014; Belyk and Brown, 2017). Quantifications

of cortical microstructure, such as myelin content, indicates that the ventral larynx region is

not located in primary motor cortex (Eichert et al., 2020a). The focus of the current study,

however, is not the interpretation of the functional activation peaks *per se* but the relationship

between their location and the underlying morphology. The individual variability in the cortical

location of the ventral larynx area could be one factor in explaining inconsistent reports in the

literature.

Relationship between Variability in Structure and Function

Cross-subject registration based on anatomical features decreased the spatial variability of the

functional peaks, indicating that anatomical variability explains some functional variability. The

somatomotor system, however, is only one example system, where a link between sulcal

morphology and function at the individual subject level was established (Boling et al., 1999;

Coulon et al., 2011; Zlatkina et al., 2016; Germann et al., 2019). Detailed anatomical and

functional studies have also revealed these relationships for other parts of the cortex (Amiez

et al., 2006; Bodin et al., 2018).

A registration based on individually drawn sulci showed a better registration for the left ventral

larynx area, when compared to a registration based on sulcal depth and based on FreeSurfer

labels, which rely on prominent and consistent landmarks, such as deep sulci. This observation

demonstrates that manual identification of anatomical features is particularly important to

examine spatial morphology in variable brain regions that have low gyrification. Further work

on structural and functional registrations may reveal a more nuanced picture of which

combination of anatomical features can better predict functional activations.

In the current study, we examined gross-anatomical features of the brain surface, an approach

that is directly applicable to neuronavigation and neurosurgical planning. Recent advances in

neuroimaging, however, allow us to acquire multiple image modalities in the same subjects,

some of which correlate with the underlying cellular microstructure (Fischl and Dale, 2000;

Zhang et al., 2012; Weiskopf et al., 2013). Although the histological analysis of post-mortem

brains remains the gold standard to define anatomical parcels of the brain, neuroimaging has

become a versatile tool for parcellating the cortex and to establish relationships between

different modalities (Glasser et al., 2016; Jakobsen et al., 2018; Smith et al., 2019; Eichert et

al., 2020b). Future work will show how detailed morphological labelling and segmentation

studies can benefit from other neuroimaging modalities.

In summary, the current study demonstrated a robust relationship between morphological

features of the central and subcentral sulcal segments and movement of different effectors at

the individual subject level. We described the morphological patterns of the sulci in the

subcentral and central opercular cortical region and their inter-individual variability. Laryngeal

activity during vocalization activates the subcentral region, which we found to be highly

variable across participants. This variability could explain inconsistencies in previous reports

about a ventral larynx area.

Extended Data

Figure 4-1: Quantification of probability maps. Bars show the maximal probability of overlap for sulcal labels in surface and volume space in the left (L) and right (R) hemisphere. Horizontal black lines indicate the median value of the probability map.

Figure 6-1: Individual sulcal labels and activation maxima. Data from the subset of subjects, where both anatomical and functional data were available.

Figure 6-2: Effect of anatomical registrations on functional variability. Distance of task activation peaks

prior to any anatomical registration (blank bars), after registration based on FreeSurfer labels

(precentral gyrus, central sulcus, postcentral gyrus, subcentral gyrus/sulcus) (striped bars), after

registration based on sulcal depth (dotted bars) and after registration based on manually drawn sulcal

labels (crossed bars). The median distance ± 95% confidence interval is shown across all peaks for each

task cluster (n = 20, 190 pairs of points for each cluster).

Author Contributions

712 <u>Author Contributions</u>

713 Conception and design: NE, MP. Acquisition of data: NE, KEW. Analysis and interpretation of

714 data: NE, KEW, RBM, MP. Original draft: NE, MP. Revising the article: NE, KEW, RBM, MP.

715 Contribution of analytic tools: NE, RBM.

Data Accessibility

<u>Data Accessibility</u>

Upon acceptance of the manuscript, all derived data supporting the findings of this study will

be made available from the Wellcome Centre for Integrative Neuroimaging's GitLab at

git.fmrib.ox.ac.uk/neichert/project_variability. Anatomical raw data of the subjects that

provided structural and functional data is publicly available at OpenNeuro under the accession

code ds002634 (version 1.0.1). The minimally pre-processed data of the HCP subjects is openly

available for download at https://db.humanconnectome.org.

Code and Software Accessibility

<u>Code and Software Accessibility</u>

Upon acceptance of the manuscript, all processing code will be made available from the

Wellcome Centre for Integrative Neuroimaging's GitLab at

git.fmrib.ox.ac.uk/neichert/project_variability. FSL tools, including MSM, are available from

fsl.fmrib.ox.ac.uk. Connectome Workbench is available at

www.humanconnectome.org/software/connectome-workbench.html.

References

<u>References</u>

Amiez C, Kostopoulos P, Champod AS, Petrides M (2006) Local morphology predicts functional

organization of the dorsal premotor region in the human brain. Journal of Neuroscience

26:2724–2731.

Amiez C, Neveu R, Warrot D, Petrides M, Knoblauch K, Procyk E (2013) The location of

feedback-related activity in the midcingulate cortex is predicted by local morphology.

Journal of Neuroscience 33:2217–2228.

Andersson JLR, Jenkinson M, Smith SM (2007) Non-linear optimisation FMRIB technial eport

TR07JA1.

Barch DM et al. (2013) Function in the human connectome: Task-fMRI and individual

differences in behavior. NeuroImage 80:169–189.

Belyk M, Brown S (2017) The origins of the vocal brain in humans. Neuroscience and

Biobehavioral Reviews 77:177–193.

Bennett CM, Miller MB (2010) How reliable are the results from functional magnetic resonance

imaging? Annals of the New York Academy of Sciences 1191:133–155.

Bodin C, Takerkart S, Belin P, Coulon O (2018) Anatomo-functional correspondence in the

superior temporal sulcus. Brain Structure and Function 223:221–232.

Boling W, Olivier A, Bittar RG, Reutens D (1999) Localization of hand motor activation in Broca's

pli de passage moyen. Journal of Neurosurgery 91:903–910.

Bouchard KE, Mesgarani N, Johnson K, Chang EF (2013) Functional organization of human

sensorimotor cortex for speech articulation. Nature 495:327–332.

Broca P (1861) Remarques sur le siège de la faculté du langage articulé suivies d'une

observation d'aphemie. Bull Soc Anthropol 6:330–357.

Carey D, Krishnan S, Callaghan MF, Sereno MI, Dick F (2017) Functional and quantitative MRI

mapping of somatomotor representations of human supralaryngeal vocal tract. Cerebral

Cortex 27:265–278.

Casey BJ et al. (2018) The Adolescent Brain Cognitive Development (ABCD) study: Imaging

acquisition across 21 sites. Developmental Cognitive Neuroscience 32:43–54.

Clouchoux C, Coulon O, Anton JL, Mangin JF, Régis J (2006) A new cortical surface parcellation

model and its automatic implementation. In: Lecture Notes in Computer Science

(including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in

Bioinformatics), pp 193–200. Springer, Berlin, Heidelberg.

Coulon O, Pizzagalli F, Operto G, Auzias G, Delon-Martin C, Dojat M (2011) Two new stable

anatomical landmarks on the Central Sulcus: Definition, automatic detection, and their

relationship with primary motor functions of the hand. In: Proceedings of the Annual

International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS,

pp 7795–7798. IEEE.

Croxson PL, Forkel SJ, Cerliani L, Thiebaut De Schotten M (2018) Structural variability across

the primate brain: A cross-species comparison. Cerebral Cortex 28:3829–3841.

Derrfuss J, Vogt VL, Fiebach CJ, Von Cramon DY, Tittgemeyer M (2012) Functional organization

of the left inferior precentral sulcus: Dissociating the inferior frontal eye field and the

inferior frontal junction. NeuroImage 59:3829–3837.

Dorsaint-Pierre R, Penhune VB, Watkins KE, Neelin P, Lerch JP, Bouffard M, Zatorre RJ (2006)

Asymmetries of the planum temporale and Heschl's gyrus: Relationship to language

lateralization. Brain 129:1164–1176.

Eichert N, Papp D, Mars RB, Watkins KE (2020a) Mapping human laryngeal motor cortex during

vocalization. bioRxiv:2020.02.20.958314.

Eichert N, Robinson EC, Bryant KL, Jbabdi S, Jenkinson M, Li L, Krug K, Watkins KE, Mars RB

(2020b) Cross-species cortical alignment identifies different types of neuroanatomical

reorganization in higher primates. bioRxiv (https://doi.org/101101/645234).

Fischl B, Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic

resonance images. Proceedings of the National Academy of Sciences of the United States

of America 97:11050–11055.

Fischl B, Sereno MI, Tootell RBH, Dale AM (1999) High-resolution intersubject averaging and a

coordinate system for the cortical surface. Human Brain Mapping 8:272–284.

Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL (2011) Unbiased average age-

appropriate atlases for pediatric studies. NeuroImage 54:313–327.

Foundas AL, Leonard CM, Gilmore RL, Fennell EB, Heilman KM (1996) Pars triangularis

asymmetry and language dominance. Proceedings of the National Academy of Sciences

of the United States of America 93:719–722.

Germann J, Chakravarty MM, Collins LD, Petrides M (2019) Tight coupling between

morphological features of the central sulcus and somatomotor body representations: A

combined anatomical and functional MRI study. Cerebral Cortex.

Germann J, Robbins S, Halsband U, Petrides M (2005) Precentral sulcal complex of the human

brain: Morphology and statistical probability maps. Journal of Comparative Neurology

493:334–356.

Glasser MF, Smith SM, Marcus DS, Andersson JLR, Auerbach EJ, Behrens TEJ, Coalson TS, Harms

MP, Jenkinson M, Moeller S, Robinson EC, Sotiropoulos SN, Xu J, Yacoub E, Ugurbil K, Van

Essen DC (2016) The Human Connectome Project's neuroimaging approach. Nature

Neuroscience 19:1175–1187.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S,

Webster M, Polimeni JR, Van Essen DC, Jenkinson M (2013) The minimal preprocessing

pipelines for the Human Connectome Project. NeuroImage 80:105–124.

Grabski K, Lamalle L, Vilain C, Schwartz JL, Vallée N, Tropres I, Baciu M, Le Bas JF, Sato M (2012)

Functional MRI assessment of orofacial articulators: Neural correlates of lip, jaw, larynx,

and tongue movements. Human Brain Mapping 33:2306–2321.

Hickok G, Lawrence Small S, Hillert DG (2016) Neurobiology of Language. Igarss 2014:1188.

Hitzig E, Fritsch GT (1870) Über die elektrische Erregbarkeit des Grosshirns. Arch Anat

Physiol:300–332.

Jackson J. (1863) Convulsive spasms of the right hand and arm preceding epileptic seizures.

Jakobsen E, Liem F, Klados MA, Bayrak Ş, Petrides M, Margulies DS (2018) Automated

individual-level parcellation of Broca's region based on functional connectivity.

NeuroImage 170:41–53.

Jarvis ED (2019) Evolution of vocal learning and spoken language. Science 366:50–54.

Jenkinson M (2003) Fast, automated, N-dimensional phase-unwrapping algorithm. Magnetic

Resonance in Medicine 49:193–197.

Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved optimization for the robust and

accurate linear registration and motion correction of brain images. NeuroImage 17:825–

841.

Jenkinson M, Smith S (2001) A global optimisation method for robust affine registration of

brain images. Medical Image Analysis 5:143–156.

Josse G, Tzourio-Mazoyer N (2004) Hemispheric specialization for language. Brain Research

Reviews 44:1–12.

Klein A, Ghosh SS, Avants B, Yeo BTT, Fischl B, Ardekani B, Gee JC, Mann JJ, Parsey R V. (2010)

Evaluation of volume-based and surface-based brain image registration methods.

NeuroImage 51:214–220.

Miller MB, Van Horn JD, Wolford GL, Handy TC, Valsangkar-smyth M, Inati S, Grafton S,

Gazzaniga MS (2002) Extensive individual differences in brain activations associated with

episodic retrieval are reliable over time. Journal of Cognitive Neuroscience 14:1200–1214.

Nieto-Castañón A, Fedorenko E (2012) Subject-specific functional localizers increase sensitivity

and functional resolution of multi-subject analyses. NeuroImage 63:1646–1669.

Olthoff A, Baudewig J, Kruse E, Dechent P (2008) Cortical Sensorimotor Control in Vocalization:

A Functional Magnetic Resonance Imaging Study. The Laryngoscope 118:2091–2096.

Penfield W, Boldrey E (1937) Somatic motor and sensory representation in the cerebral cortex

of man as studied by electrical stimulation. Brain 60:389–443.

Petrides M (2012) The human cerebral cortex. An MRI atlas of the sulci and gyri in MNI

stereotaxic space. :168.

Petrides M (2019) Atlas of the morphology of the human cerebral cortex on the average MNI

brain. New York: Academic Press.

Price CJ (2000) The anatomy of language: Contributions from functional neuroimaging. Journal

of Anatomy 197:335–359.

Ribas GC (2010) The cerebral sulci and gyri. Neurosurgical Focus 28.

Robinson EC, Jbabdi S, Glasser MF, Andersson J, Burgess GC, Harms MP, Smith SM, Van Essen

DC, Jenkinson M (2014) MSM: A new flexible framework for multimodal surface matching.

NeuroImage 100:414–426.

Simonyan K (2014) The laryngeal motor cortex: Its organization and connectivity. Current

Opinion in Neurobiology 28:15–21.

Smith S, Duff E, Groves A, Nichols TE, Jbabdi S, Westlye LT, Tamnes CK, Engvig A, Walhovd KB,

Fjell AM, Johansen-Berg H, Douaud G (2019) Structural variability in the human brain

reflects fine-grained functional architecture at the population level. The Journal of

Neuroscience 39:6136–6149.

Sprung-Much T, Petrides M (2018) Morphological patterns and spatial probability maps of two

defining sulci of the posterior ventrolateral frontal cortex of the human brain: the sulcus

diagonalis and the anterior ascending ramus of the lateral fissure. Brain Structure and

Function 223:4125–4152.

Sprung-Much T, Petrides M (2019) Morphology and spatial probability maps of the horizontal

ascending ramus of the lateral fissure. Cerebral Cortex 00:1–17.

Takerkart S, Auzias G, Brun L, Coulon O (2015) Mapping cortical shape differences using a

searchlight approach based on classification of sulcal pit graphs. In: Proceedings -

International Symposium on Biomedical Imaging, pp 1514–1517. IEEE.

Uğurbil K et al. (2013) Pushing spatial and temporal resolution for functional and diffusion MRI

in the Human Connectome Project. NeuroImage 80:80–104.

Van Essen DC, Smith SM, Barch DM, Behrens TEJJ, Yacoub E, Ugurbil K (2013) The WU-Minn

Human Connectome Project: An overview. NeuroImage 80:62–79.

Vigneau M, Beaucousin V, Hervé PY, Duffau H, Crivello F, Houdé O, Mazoyer B, Tzourio-

Mazoyer N (2006) Meta-analyzing left hemisphere language areas: Phonology, semantics,

and sentence processing. NeuroImage 30:1414–1432.

Vogt C, Vogt O (1919) Allgemeinere Ergebnisse unserer Hirnforschung. Journal für Psychologie

und Neurologie.

Weiskopf N, Suckling J, Williams G, Correia M. MM, Inkster B, Tait R, Ooi C, Bullmore T. ET, Lutti

A (2013) Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: A multi-

center validation. Frontiers in Neuroscience 7:95.

Woo CW, Krishnan A, Wager TD (2014) Cluster-extent based thresholding in fMRI analyses:

Pitfalls and recommendations. NeuroImage 91:412–419.

Woolrich MW, Ripley BD, Brady M, Smith SM (2001) Temporal autocorrelation in univariate

linear modeling of FMRI data. NeuroImage 14:1370–1386.

Yousry TA, U.D. Schmid HA, Schmidt D, Peraud A, Buettner A, Winkler P (1997) Localization of

the motor hand area to a knob on the precentral gyrus. Brain 120:141–157.

Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC (2012) NODDI: Practical in vivo

neurite orientation dispersion and density imaging of the human brain. NeuroImage

61:1000–1016.

Zlatkina V, Amiez C, Petrides M (2016) The postcentral sulcal complex and the transverse

postcentral sulcus and their relation to sensorimotor functional organization. European

Journal of Neuroscience 43:1268–1283.

Zlatkina V, Petrides M (2010) Morphological patterns of the postcentral sulcus in the human

brain. Journal of Comparative Neurology 518:3701–3724.