Can a population targeted by a CRISPR-based homing gene drive be rescued?

Nicolas O. Rode[*], Virginie Courtier-Orgogozo[†], Florence Débarre[‡]

[*] CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France
[†] Institut Jacques Monod, CNRS UMR7592, Univ. de Paris, 75013 Paris, France
[‡] Sorbonne Université, CNRS, IRD, INRA, Institute of Ecology and Environmental Sciences-Paris (IEES Paris), place Jussieu, 75005 Paris, France
ORCID: 0000-0002-1121-4202 (N.O.R.), 0000-0002-9297-9230 (V. C.-O.), 0000-0003-2497-833X (F.D.)
Corresponding authors: E-mails: florence.debarre@normalesup.org; virginie.courtier@ijm.fr; nicolas.rode@inrae.fr

Keywords: Genetic control, gene drive brake, theoretical model

Running title: Rescue of a population targeted by a CRISPR-based homing gene drive

Article type: Investigation

Abbreviations:

CRISPR: Clustered Regularly Interspaced Short Palindromic Repeats

Article summary for Issue Highlights (100 words)

Homing gene drive is a new genetic control technology that aims to spread a genetically engineered DNA construct within natural populations even when it impairs fitness. In case of unanticipated damages, it has been proposed to stop homing gene drives by releasing individuals carrying a gene-drive brake; however, the efficiency of such brakes has been little studied. The authors develop a model to investigate the dynamics of a population targeted by a homing drive in absence or in presence of brake. The model provides insights for the design of more efficient brakes and safer gene drives.

(96 words)

Abstract (250 words):

CRISPR-based homing gene drive is a genetic control technique aiming to modify or eradicate natural populations through the release of individuals carrying an engineered piece of DNA that can be inherited by all their progeny. Developing countermeasures is important to control the spread of gene drives, should they result in unanticipated damages. One proposed countermeasure is the introduction of individuals carrying a brake construct that targets and inactivates the drive allele but leaves the wild-type allele unaffected. Here we develop models to investigate the efficiency of such brakes. We consider a variable population size and use a combination of analytical and numerical methods to determine the conditions where a brake can prevent the extinction of a population targeted by an eradication drive. We find that a brake is not guaranteed to prevent eradication and that characteristics of both the brake and the drive affect the likelihood of recovering the wild-type population. In particular, brakes that restore fitness are more efficient than brakes that do not. Our model also suggests that threshold-dependent drives (drives that can spread only when introduced above a threshold) are more amenable to control with a brake than drives that can spread from an arbitrary low introduction frequency (threshold-independent drives). Based on our results, we provide practical recommendations and discuss safety issues.

Introduction

The use of engineered gene drives has been proposed as a technique for population control with potential applications in public health, agriculture and conservation (Burt 2003; Esvelt *et al.* 2014). This technique relies on the release of genetically engineered individuals that can rapidly propagate a transgene of interest into wild populations. Gene drive can be designed to modify, suppress or eradicate various target species (Scott *et al.* 2018; Rode *et al.* 2019). Potential target species include disease vectors (e.g. *Anopheles gambiae*, the main vector of malaria in Africa; Kyrou *et al.* 2018), agricultural pests (e.g. *Drosophila suzukii,* a major pest of soft fruits; Scott *et al.* 2018) or invasive rodents (e.g. invasive house mouse or black rats that threaten biodiversity on islands; Leitschuh *et al.* 2018).

Due to the universality of CRISPR genome editing, CRISPR-based gene drives can potentially be applied to a wide variety of organisms (Esvelt *et al.* 2014; Raban *et al.* 2020). Diverse CRISPR-based gene drive systems have already been developed in the laboratory as proofs-of-principle in a few model organisms (homing, split homing, translocation, X-shredder, killer-rescue, cleave-and-rescue and TARE gene drives; Webster *et al.* 2019; see Raban *et al.* 2020 for a review; Champer *et al.* 2020) or as theoretical possibilities (daisy chain drives; Noble *et al.* 2019). Gene drives have so far only been tested in the laboratory and no field trial has been conducted yet.

Among these systems, CRISPR-based homing gene drives are the most adaptable to new species and populations and the most advanced in terms of technological development (Raban *et al.* 2020). They involve a piece of DNA that includes a guide RNA (gRNA) gene and a *cas9* gene (encoding for the Cas9 endonuclease). The gRNA is designed to recognize a specific sequence in a wild-type chromosome, so that that in heterozygotes carrying a drive allele and a wild-type allele, the Cas9-gRNA molecular complex will cut the wild-type chromosome at the target site. The resulting double-strand DNA break can then be repaired through homology-directed repair (also known as "gene conversion"), using the drive allele as a template, which is designed to harbor sequences identical to the ones flanking the target site. Consequently, the drive allele is transmitted to the next generation at rates beyond those of regular Mendelian inheritance and, if its parameters allow it, will rapidly spread within the target population.

Homing gene drives are sometimes considered as "threshold-independent drives", i.e. as being able to spread in a population from an arbitrary low introduction frequency (e.g. Marshall and Akbari 2018). Mathematical models of homing gene drives (e.g. Deredec *et al.* 2008; Alphey and Bonsall 2014; Unckless *et al.* 2015; Tanaka *et al.* 2017) have however shown that depending on various parameters (the efficacy of gene conversion, its timing, the fitness cost incurred by the drive allele and its dominance over the wild-type allele), some of the homing gene drives can be threshold-dependent, i.e. only spread if they are introduced above a threshold frequency. Mathematically, when there is an equilibrium at an intermediate frequency of the drive allele ($0 < p_D < 1$) and when this equilibrium is unstable, then the drive is threshold-dependent; the value of the drive allele frequency at this equilibrium is the threshold above which the drive has to be introduced to spread (Deredec *et al.* 2008).

Given that gene drives can potentially impact biodiversity, national sovereignty and food security (Oye *et al.* 2014; Akbari *et al.* 2015; DiCarlo *et al.* 2015; NASEM 2016; Montenegro de Wit 2019), there is a crucial need to develop strategies to minimize the risks of unintentional spread (e.g. following the escape of gene drive individuals from a laboratory) and to mitigate unanticipated or premeditated and malevolent harm to humans or the environment. For example, a CRISPR-based eradication drive may spread into a non-target population or species (Noble *et al.* 2018; Courtier-Orgogozo *et al.* 2019a; Rode *et al.* 2019); a modification drive may alter the target population in an unexpected, detrimental manner;

or a gene drive could be used as bioweapon (Gurwitz 2014), for example to modify a locally important
pollinator insect species and cause heavy agricultural production losses. Decreasing the environmental
risks associated with the development of this technology can be achieved by designing safer gene drives
whose spread can be controlled spatially or temporally (Marshall and Akbari 2018; Raban *et al.* 2020)
and by developing countermeasures to stop the spread of an ongoing gene drive (Esvelt *et al.* 2014;
Gantz and Bier 2016; Vella *et al.* 2017).

Several countermeasure strategies for CRISPR-based homing gene drives have been proposed. One
strategy is to use gene drives whose non-Mendelian transmission is conditional on the presence of
synthetic molecules in the environment of the target species, so that the removal of the synthetic
molecule is expected to stop the spread of the gene drive, and natural selection to remove the drive from
the population (Esvelt *et al.* 2014; Del Amo *et al.* 2020). However, the development of such molecule-
dependent drives is still at its infancy and may have to be tailored for each ecosystem and target species.
Another strategy is to introduce resistant individuals carrying a modified target locus that prevents
homing ("synthetic resistant" (SR) allele; Burt 2003; Champer *et al.* 2016; Vella *et al.* 2017). Such
synthetic resistant alleles are however predicted to be rather ineffective against replacement drives with
small fitness costs (Vella *et al.* 2017), because of the limited selective advantage of synthetic resistant
alleles. Alternatively, it has been proposed to release suppressor individuals that carry a new piece of
DNA which will eventually lead to the knock-out of the initial gene drive (Esvelt *et al.* 2014; Marshall
and Akbari 2018). These alternative countermeasures rely on gene conversion and can be used against
virtually any type of CRISPR-based homing gene drive. Two types exist. The first type are
countermeasures that include the *cas9* gene and that can target the drive only (reversal drives sensu
Esvelt *et al.* 2014; overwriting drives; DiCarlo *et al.* 2015) or both the drive and wild-type alleles
(immunizing reversal drive (IRD); Esvelt *et al.* 2014; Vella *et al.* 2017). However, with these strategies,
a functional *cas9* gene will remain in the final population, which may increase the risk of subsequent
genetic modifications such as translocations, and possible negative environmental outcomes (Courtier-
Orgogozo *et al.* 2019b). The second type are countermeasures that do not encode for *cas9* and rely
instead on the *cas9* gene present in the initial gene drive construct. They can be contained in a single
locus (ERACR: element for reversing the autocatalytic chain reaction, Gantz and Bier 2016; CATCHA:
Cas9-triggered chain ablation, Wu *et al.* 2016), or be across two loci (CHACR: construct hitchhiking
on the autocatalytic chain reaction, Gantz and Bier 2016). These countermeasures might be safer for
the environment, due to the absence of a functional *cas9* gene. To our knowledge, neither ERACR nor
CHACR have been implemented in the lab; CHACR may be slow to spread due to its two-locus
structure, while ERACR may be sensitive to the evolution of resistance at its target sites (*cas9*-flanking
sequences whose mutation does not affect enzyme function).

We focus here on the in our opinion best gene-drive-based countermeasures proposed so far, the cas9-
devoid reversal drives (CATCHA, ERACR), which we call hereafter "brakes" for simplicity. In
drive/brake heterozygotes, the encoded guide RNA(s) target and inactivate the *cas9* gene of the initial
gene drive construct. Such brakes should be especially efficient, because even in absence of homology-
directed repair, the drive's *cas9* gene (targeted by the brake) is expected to be inactivated. However, for
simplicity, we will not model this additional scenario here.

Although mathematical modelling of the effects of brakes has been recommended (Wu *et al.* 2016), to
our knowledge only two such studies have been published (Vella *et al.* 2017; Girardin *et al.* 2019).
Vella et al. found that the introduction of a brake leads to a polymorphic equilibrium with transient
oscillatory dynamics (Figure 2d,e in Vella *et al.* 2017). They also showed that brakes with smaller
fitness costs increased the likelihood of long-term eradication of the homing gene drive (Figure 3 in
Vella *et al.* 2017). We note that because Vella et al. assumed 100% cleavage and germline conversion,

the drive they modeled was threshold-independent (Deredec *et al.* 2008). Girardin et *al.* (2019)
considered a spatial model, and found that a brake could stop a spatially spreading drive only if the
drive was threshold-dependent, and that threshold-independent drives led to an infinite spatial chase of
the drive by the brake. While both studies provided insights on our ability to control an ongoing gene
drive, they had limitations. First, both studies used classical population-genetic frameworks, and
focused on allele frequency dynamics, ignoring changes in population size. They thereby omitted
potential demographic feedbacks on allele frequency changes, which are likely to be important for
eradication drives. It thus remains unknown whether a brake can prevent the extinction of a population
targeted by an eradication drive. Second, both studies used deterministic models. Vella et al.
acknowledged that oscillations of the allele frequencies in their model could lead to the stochastic loss
of an allele. Similar oscillations were observed by Girardin et *al.* (2019), but their implications were not
explored.

To address some of the limitations of previous models and examine further the effectiveness of brakes,
we model here the dynamics of a population targeted by a drive, into which brake-carrying individuals
are released. We consider a variable population size and its potential feedback onto gene frequency
changes, and we also develop a stochastic version of the model. We compare two timings of gene
conversion for gene drive and brake alleles (in the germline or zygote, Figure S1) and explore the role
of parameters such as level of dominance, cleavage efficiency, brake-associated fitness costs (whether
or not it restores fitness), and the type of fitness component targeted by the gene drive (embryo survival,
fecundity or adult death rate). We contrast brakes that restore fitness with those that do not.
Implementing brakes that restore fitness (i.e. "specific brakes") require prior knowledge of the gene
disrupted by the homing drive in order to include in the brake a recoded version of this gene along with
a gRNA that targets the *cas9* sequence of the drive allele. Hence, drive-brake heterozygous individuals
have higher fitness than drive homozygotes, but may have lower fitness than wild-type homozygotes
(as they may incur a small fitness cost due to the expression of the gRNA). Implementing CATCHA
brakes that do not restore fitness (i.e. "universal brakes") does not require prior knowledge of the gene
disrupted by the homing drive as such brakes only include a gRNA that targets the *cas9* sequence of the
drive allele. Hence, drive-brake heterozygous individuals have the same fitness as drive homozygotes.

Eradication drives currently under development target genes involved in female development in various
human-disease vectors (Kyrou *et al.* 2018) or agricultural pests (Li and Scott 2016). Because they have
the strongest demographic consequences and pose the greatest risks of unwanted spread, we focus on
threshold-independent eradication drives in the numerical part of our study. We aim at finding the
characteristics of the brakes that can efficiently stop an ongoing gene drive and allow the recovery of a
wild-type population.

Methods

Analytical model

With three different alleles in the population (wild-type 0, drive D and brake B), we need to follow the
dynamics of six diploid genotypes. We denote by $G = \{00, 0D, DD, 0B, DB, BB\}$ the set of all possible
genotypes. To take into account gene drives that affect population size (e.g. eradication drives), we
consider the densities of individuals of each genotype and do not focus solely on genotype frequencies
as previous models did (Deredec *et al.* 2008; Unckless *et al.* 2015; Vella *et al.* 2017; Girardin *et al.*
2019). We denote the density of individuals of genotype g by N_g and the total population density by

N(omitting the time dependence (t) for concision; $N = \sum_g \; N_g$). We consider three traits affecting
fitness that can vary among genotypes: the survival of zygotes (ω_g), the death rate of adults (d_g), and
the fecundity of adults (β_g). We assume that reproduction is density-dependent: it depends on the total
population size N, following a classical logistic regulation with carrying capacity K. The death rate, on
the other hand, is density-independent. The change over time in the density of individuals of genotype
g is given by

$$\frac{d\,N_g}{d\,t} = \omega_g\, V_g\, N(1 - N/K) - d_g\, N_g, \tag{1}$$

where V_g corresponds to the production of new individuals of genotype g through sexual reproduction
and depends on the abundances of all genotypes, their fecundities β_g , but also on the timing of gene
conversion. The formulas of the V_g terms for each timing of gene conversion are given in the Appendix
(and also provided in the supplementary Mathematica file).

We consider that gene conversion in $0D$ or DB heterozygous individuals can either occur in the
germline or in the zygote (Fig. S1). When gene conversion occurs in the germline, $0D$ and DB
heterozygous individuals produce more than 50% of D and B gametes respectively. When gene
conversion occurs in newly formed zygotes (i.e. immediately after fertilization), $0D$ and
DB heterozygous individuals are converted into DD and BB homozygotes respectively and have the
corresponding traits. For both types of gene conversion, we denote the probabilities of successful gene
conversion by drive and by brake alleles by c_D and c_B respectively.

Numerical explorations

While our analytic results are obtained with generic parameters, numerical explorations require specific
parameter values. The number of parameter combinations to explore being very vast, we make a few
assumptions to reduce it. First, we consider that drive and brake affect either (i) zygote survival (ω),
(ii) adult survival (d) or (iii) adult fecundity (β), all other parameters remaining equal across genotypes.
To model an eradication drive, we chose ω_{DD}, d_{DD} or β_{DD} such that a 100% drive population is not
viable and standardised the parameters to yield the same negative equilibrium value of population size
(we set $\frac{d_{DD}}{\omega_{DD} b_{DD}^2} = 1.1$, see Mathematica Appendix for details). We consider that either the brake allele
does not restore the fitness loss due to the drive allele (i.e. it has the same fitness as the drive allele), or
that the brake allele restores partially the fitness loss and has a small fitness cost compared to the wild-
type allele (i.e. it contains a specific cargo that helps to restore fitness). We use the same dominance
parameter, h, for both drive and brake alleles. When the brake allele does not restore fitness, its
dominance is the same as that of the drive allele. When the brake allele does restore fitness, we consider
that h has little effect on the fitness of $0B$ heterozygous as the fitness BB homozygotes is much closer
to that of 00 than of DD homozygotes. For juvenile survival, the parameters of heterozygotes read:

$$\omega_{0D} = (1 - h)\omega_{00} + h\omega_{DD}$$

$$\omega_{0B} = (1 - h)\omega_{00} + h\omega_{BB} \tag{2}$$

$$\omega_{DB} = (1 - h)\omega_{BB} + h\omega_{DD},$$

and likewise for d and β parameters. In the numerical part of the study, we consider either complete
recessivity ($h = 0$) or codominance ($h = 0.5$).

We therefore have 24 combinations of parameters (2 timings of gene conversion x 3 traits affected x
dominance values x 2 types of brake). For each of them, we consider different timings of introduction
of the brake in the population; the timing is given in terms of the current frequency f_I of the drive allele
in the population at the time at which the brake is introduced. The $N^{(0)}{}_{0B}$ parameter represents the
number of released wild-type/brake heterozygous individuals. Unless stated, we assume that $N^{(0)}{}_{0B} =$
100. Other parameters are shown in tables S1-S3.

Reformulating the model

218   Our model is initially defined in terms of genotype densities (equation 1). To simplify the analyses, we
219   reparametrize the model in terms of total population size N, allele frequencies p_D and p_B (we have $p_0 =$
220   $1 - p_D - p_B$), and deviations from Hardy-Weinberg for each of the three heterozygotes
221   ($\delta_{0D}, \delta_{0B}, \delta_{DB}$). In particular,

$$N = N_{00} + N_{0D} + N_{DD} + N_{0B} + N_{DB} + N_{BB}, \tag{3a}$$

$$p_D = \frac{N_{DD} + \frac{1}{2}N_{0D} + \frac{1}{2}N_{DB}}{N}, \tag{3b}$$

$$\delta_{0D} = \frac{N_{0D}}{N} - 2p_D p_0, \tag{3c}$$

and likewise for p_B, δ_{0B} and δ_{DB} (the full equations are calculated in the supplementary Mathematica
file).

As usual with most continuous-time models (Nagylaki and Crow 1974), we cannot neglect deviations
from Hardy-Weinberg frequencies here (unlike models with discrete, non-overlapping generations).
The reformulated model (system (3)) also highlights interactions between total population size N and
changes in allele frequencies (eco-evolutionary feedbacks). The population growth rate depends on
population composition, since fecundity or survival parameters are genotype-dependent. Reciprocally,
changes in allele frequencies depend on the size of the population. This is because gene conversion,
which modifies allele frequencies, takes place upon reproduction (either in the germline, or in the newly
formed zygote). Given that reproduction is negatively density-dependent, changes in the frequencies of
drive and brake alleles slow down when population size is larger.

Stability analyses

234   We use the reformulated version of the model (system (3)) to find evolutionary equilibria and analyse
235   their stabilities.

Model without the brake

We first study the properties of our model when the brake is absent (setting all variables containing the
brake allele equal to zero). We determine the equilibrium states where only one allele is present (i.e.
boundary equilibria). At the wild-type-only equilibrium, we have $N = K(1 - \frac{d_{00}}{\omega_{00}b_{00}^2})$, $p_D = 0$, $\delta_{0D} =$
0 (see Mathematica Appendix for details). At the drive-only equilibrium, the size of the population
depends on the type of drive. As we only consider eradication drives (i.e. drives such that a drive-only

population is not viable), we have $N = 0$, $p_D = 1$, $\delta_{0D} = 0$ at the drive-only equilibrium (for
completeness though, we included in the Mathematica appendix a separate stability analysis of the
drive-only equilibrium for replacement drives). Generic formulas for interior equilibria (i.e. for which
$0 < p_D < 1$) could not be found analytically.

Model with the brake

For simplicity, in the full model with the three alleles, we only study the stability of the wild-type-only
equilibrium ($N = K(1 - \frac{d_{00}}{\omega_{00}b_{00}^2})$, $p_D = 0$, $p_B = 0$, $\delta_{0D} = 0$, $\delta_{0B} = 0$, $\delta_{DB} = 0$).

Numerical solutions and stochastic simulations

Deterministic solutions of the model

To test the robustness of the equilibrium states predicted by our analytical model, we solve the model
numerically for specific sets of parameters, using the original formulation in equation 1. We use
parameter values for a threshold-independent eradication drive (i.e. as explained in the result section
below, conditions where, according to the stability analysis of our model, the wild-type population
cannot be recovered after the introduction of the brake). Time is discretized; we consider small fixed
time steps $dt = 0.005$. When the system undergoes oscillations, genotype densities can go down to
extremely small values, possibly below computer precision. We therefore set a critical value $thr = $
0.01, below which the density of a genotype is considered to be zero.

Stochastic simulations

To explore the effect of stochasticity on our model, we implement a stochastic version of it using a
Gillespie algorithm. As for numeric simulations, we only consider parameter values for a threshold-
independent eradication drive. In short, within a time step we (i) compute the rates (or "propensities")
of all possible events (birth and death probabilities of each of the five genotypes); (ii) randomly pick
one event (the higher the event's rate, the more likely its occurence); (iii) update the population
according to the event that has taken place; (iv) draw the time interval that lasted the step (according to
an exponential distribution parameterized by the sum of all propensities). For each set of parameter
values, we run 10000 simulations, each of them until a maximum time value ($t_{max} = 25000$) or until
the population goes extinct. For each simulation, we list the different types of outcome (i.e., WT
recovery after introduction of the brake, coexistence between the wild type and either the brake or both
the initial gene drive and the brake, extinction before or after the introduction of the brake, drive loss
before brake introduction).

Data availability

273   Supplemental Material Files S1-S2 is available at Figshare:
274   https://doi.org/10.6084/m9.figshare.11982285.v1

275   File S1 contains supplemental script for the analytical model (Mathematica notebook). File S2 contains
276   scripts for numerical explorations and stochastic simulations.

Results

To assess the efficiency of various types of brakes to control gene drives, we use a combination of (i) analytical techniques (stability analysis of the deterministic model), (ii) numerical solutions of the deterministic model, and (iii) stochastic simulations. The stability analysis (i) is done with generic parameters. For the numerical steps of our exploration of the model ((ii) and (iii)), we use specific parameters corresponding to threshold-independent eradication drives, i.e. drives that spread from very low frequencies, and whose fixation leads to the extinction of the population.

There are four categories of homing drives

To better understand the dynamics of the full model with three alleles (wild-type, drive, brake), we first study the model in the absence of brake. This analysis is done using generic parameters, separately for each timing of gene conversion (zygote vs. germline conversion).

In this two-allele version of the model, there are two boundary equilibria: drive loss (the wild-type allele is fixed) and drive fixation. These two equilibria can be locally stable or unstable, so that there are up to four possible combinations of stabilities and therefore four possible outcomes: (i) drive loss, (ii) coexistence of the drive and wild-type alleles, (iii) drive fixation, (iv) bistability (Deredec *et al.* 2008; Alphey and Bonsall 2014; Unckless *et al.* 2015; Noble *et al.* 2017; Vella *et al.* 2017; Girardin *et al.* 2019). Drives in (ii) and (iii) will invade the wild-type population from an arbitrary low frequency and are "threshold-independent" (Marshall and Akbari 2018). Drives in (iv) can either spread and fix when the drive allele is introduced at a high enough frequency or will be lost when their introduction frequency is below a given threshold (i.e. there is a bistability). This type of drive is "threshold-dependent" (Akbari *et al.* 2013; Marshall and Akbari 2018). The parameter ranges corresponding to each outcome are illustrated in the supplementary Mathematica file, for replacement and eradication drives; they are consistent with the findings of previous studies (Deredec *et al.* 2008; Unckless *et al.* 2015; Vella *et al.* 2017; Girardin *et al.* 2019). The eradication drives used so far in laboratory studies (Kyrou *et al.* 2018) (large fitness cost, high conversion efficiency, recessivity and conversion in the germline) correspond to threshold-independent drives.

Stability analyses indicate that a brake can recover the wild-type population only if the drive is threshold-dependent

When the brake allele has lower fitness than the wild-type allele, the three alleles (wild-type, drive and brake), are involved in non-transitive interactions (rock-paper-scissors type; Vella *et al.* 2017): the wild-type is converted into a drive by the drive, the drive into a brake by the brake, and the brake is costly compared to the wild-type. A high frequency of the wild-type, drive or brake in the population favors the drive, brake or wild-type respectively. Such negative-frequency-dependent selection can result in the coexistence of the three alleles.

In the analytical model with the three alleles, we find that the conditions for the local stability of the wild-type-only equilibrium are the same as in the model without brake (details of the calculations are presented in the supplementary Mathematica file). In other words, our stability analysis indicates that the introduction of a brake can successfully restore a wild-type population only under two conditions. First, quite trivially, the wild-type population can be recovered when the population is targeted by a drive that would be lost in the absence of brake (drive loss equilibrium above; we ignore this case thereafter). Second, the wild-type population can be recovered when it is targeted by a threshold-

dependent drive (i.e. with parameters corresponding to a bistability in the model without brake, see above). In this case, introducing the brake allele can decrease the frequency of the drive allele below its invasion threshold; the drive is then lost. Once the drive is lost, if it is, the brake loses the competition against the wild-type allele because of its fitness cost, and the wild-type population is finally recovered.

Numerical explorations of the deterministic model and stochastic simulations show that brakes can stop threshold-independent drives under certain conditions

Numerical solutions of the deterministic model

The introduction of a brake in a population targeted by a threshold-independent drive may lead to oscillations of large amplitude. During these oscillations, the densities of some genotypes may reach extremely low values. Biologically, this is not realistic: however big a population, an extremely low density may correspond to less than one individual, and thus to the loss of an allele from the population. Computationally as well, these oscillations are challenging, because they may lead to values below the minimum number that a computer can represent, and therefore to the failure of numerical solutions. To solve both issues, we set a critical density below which a genotype is considered absent from the population and we numerically integrate our model to further explore the effect of the introduction of a brake in a population targeted by a threshold-independent eradication drive. Cutting large amplitude cycles means that alleles can be lost. The dynamics of the frequencies of the three alleles and of population size (scaled by the equilibrium density of the wild-type population) are shown in Figure 2. These dynamics depend on the trait that is affected by the drive and the brake (fecundity, adult mortality, or zygote survival; lines in Figure 2), the level of dominance (columns in Figure 2), and whether the brake restores fitness or not (Figures S3 vs. 2).

The addition of a critical density leads to outcomes that were not predicted by our stability analysis. Contrary to the predictions of the stability analysis for threshold-independent drives, in Figures 2(a) and 2(f), the drive is lost, allowing for population recovery. This is because the density of drive-carrying individuals reaches so small values at some point that the drive allele is considered extinct. Then, the brake allele being costly compared to the wild-type allele, it decreases in frequency and is lost as well. In Figure 2(b), the population goes extinct. This is because the overall population density goes down to very small values.

As expected, with our parameters, the wild-type population is more rarely recovered with a brake that does not restore fitness than with a brake that does (compare Figures 2 to S3, and S4 to S5).

We hypothesized that allele loss would happen when the amplitude of oscillations increases (i.e. when the interior equilibrium, where the three alleles coexist, is unstable). However, even when the amplitude of oscillations decreases (i.e. when the interior equilibrium is locally stable), the initial oscillations can be substantial, hindering our ability to predict the outcome. In addition, the outcome itself depends on non-biological contingencies such as the time interval at which the solutions are calculated and the critical density below which a genotype is considered extinct. As a consequence, a brake is not guaranteed to prevent the eradication of a population targeted by a threshold-independent drive.

We complemented our exploration with stochastic simulations. Notably, having integer numbers of
individuals of each genotype avoids the arbitrary choice of a critical density below which a genotype is
considered extinct. Importantly, the parameters that we chose in our simulations correspond to a large
wild-type population size (an expected density of $N^* = 10000$); the diversity of observed outcomes is
due to the large amplitude of oscillations in genotype densities triggered by the introduction of the
brake.

Among the different parameters investigated, whether or not the brake restored fitness has the highest
impact on the recovery of the wild type population (Figure 3 vs. 4 and 5 vs. 6). Our stochastic
simulations show that in many instances, the brake does not prevent population extinction when it does
not restore fitness (Figures 3 and 5). In contrast, the drive allele is always lost when the brake restores
fitness (Figures 4 and 6), resulting either in the full recovery of the wild-type population, or in a
coexistence between the wild type and the brake at the time at which the simulation ended (tmax =
2500). Noteworthily, as the fitness of the brake approaches that of the wild-type allele, the time
necessary to recover 100% wild-type individuals increases.

When the brake does not restore fitness, the recovery of the wild-type population is more frequent when
gene conversion occurs in the zygote than when it occurs in the germline, especially for recessive drives
and brakes ($h = 0$, Figure 3 vs. 5). When the brake restores fitness, the timing of conversion has little
effect on the final outcome (compare Figure 4 with Figure 6). The effects of other parameters such as
the type of trait targeted, the level of dominance or the drive frequency at brake introduction are more
difficult to predict. The most frequent outcome in stochastic simulations was often different from the
outcome predicted by deterministic models. For example, population extinction is the most frequent
outcome of some of the stochastic simulations, while the corresponding deterministic model predicts
the recovery of the wild-type population (e.g. Figures 3(a), 5(b)).

We cannot draw clear conclusions regarding the optimal timing of introduction of the brake. The
outcome strongly depends on the type of trait that is affected by the drive (and the brake), the level of
dominance and the timing of gene conversion. We conclude, in agreement with the results of Vella et
al. using infinite population size, that a brake is not guaranteed to prevent the eradication of a population
targeted by a threshold-independent eradication drive.

Discussion

We developed a model to investigate the consequences of introducing a brake allele in a population
targeted by a CRISPR-based homing gene drive. Our framework extends previous ones, which focused
on allele frequencies (ignoring fluctuations in population density) and assumed 100% cleavage
efficiency (Vella *et al.* 2017; Girardin *et al.* 2019). By accounting for the effects of both the initial gene
drive and the brake on population size, our model represents a first step towards the explicit integration
of changes in population size into the prediction of the dynamics of wild-type, gene drive and brake
alleles. While we concentrate here our numerical explorations on eradication drives and threshold-
independent drives, our model can also be used to study the dynamics of replacement drives and their
brakes, by adapting parameter values.Our model can form a basis for future studies investigating the
effect of CRISPR-based brakes against other types of gene drives (e.g. split gene drives; Li *et al.* 2020),
to check whether these alternatives might be easier to control.

Our model does not account for the potential evolution of resistance against gene drives. Such resistance
can be due to cleavage repair by non-homologous end joining or to natural variation at the target locus,
and can occur frequently after the release of gene drive individuals (Drury *et al.* 2017; Unckless *et al.*
2017; Bull *et al.* 2019). However, several strategies are under way to prevent the evolution of gene drive
resistance, such as the use of multiple gRNAs (Champer *et al.* 2018; Oberhofer *et al.* 2018; Edgington
et al. 2020) or the targeting of a functionally constrained locus whose mutations are highly deleterious
and cannot increase in frequency (e.g. Kyrou *et al.* 2018). Given these efforts to limit the evolution of
resistance against gene drives, we chose not to include this feature in our model. In addition, Vella et
al. (2017) investigated the evolution of resistance at the target locus in addition to the introduction of a
countermeasure and found that the qualitative behavior of the brake remains unchanged (polymorphic
equilibrium of all alleles).

Furthermore, we did not model the evolution of resistance against brakes either. If such resistant alleles
were to form, for the types of brakes we investigated, the consequences would differ between ERACR
and CATCHA brakes. For ERACR brakes, mutations arising in flanking sequences targeted by the
brake could prevent cleavage and conversion of the drive into a brake. If these mutations do not alter
the rate of conversion of the wild-type allele into a drive allele, a drive resistant to the ERACR brake
could continue spreading. Thus, ERACR brake could fail to prevent a population from extinction. For
CATCHA brakes, mutations in the target *cas9* sequence would result in non-functional Cas9 enzymes.
These brake-resistant alleles would have the same fitness cost as the drive allele, but without the gene-
conversion advantage of the drive. Should they appear, they would be expected to remain at a low
frequency in the population. Overall, we thus expect CATCHA brakes to overcome the evolution of
resistance against brake while ERACR brakes would not, so we recommend using the former.

Overall, our model shows that the success of recovering the wild-type population using a brake depends
both on the type of brake introduced and the type of gene drive targeted. More specifically, our
conclusions depend on the method chosen to explore the model. Our stability analysis indicates that the
wild-type population can only be recovered after the introduction of a brake if the drive is threshold-
dependent. Nevertheless, our numerical integration of the model -- including a critical population
density to avoid unrealistically low genotype densities -- and stochastic simulations show that the wild-
type population can also be recovered in certain cases when a threshold-independent drive is used. In
these cases, brakes that restore fitness can better control a gene drive than universal brakes that do not.
However, we could not draw general conclusions on the effect of other parameters (e.g. fitness trait
affected by the drive, dominance level, timing of conversion, and frequency of the drive for introducing
the brake) on the final outcome.

Our model shows that, even when the brake is introduced when the eradication drive is still at a low
frequency, the frequency of the eradication drive continues to increase and results in a strong population
bottleneck (e.g. Figure 1a). Such a strong bottleneck could result in a long term alteration of the
recovered wild-type population (e.g. due to inbreeding depression). This point is important to keep in
mind even though it is not explicitly incorporated in our model.

Our study has practical implications. First, we advise against using universal brakes as the sole
countermeasure because they are not guaranteed to succeed and stop a drive. In contrast, we recommend
using specific brakes which restore fitness, as they are more likely to be effective. They spread at a
faster rate and increase the chances of recovering a population of wild-type individuals when they
include a recoded version of the gene disrupted by the initial gene drive. We recommend that the
development of homing gene drives goes in pair with the co-development of such specific brakes.
Although they are not guaranteed to be successful, specific brakes currently represent the best

countermeasure against the spread of homing drives following an escape from a laboratory. We also
recommend laboratory studies to assess the efficacy of brakes using experimental evolution under
controlled conditions. Second, because they are easier to control with brake, we believe that threshold-
dependent homing gene drives are a safer alternative to threshold-independent homing drives, that are
currently being developed in laboratories. These threshold-independent homing drives are
characterized by large and recessive large fitness costs, high conversion efficiency and germline
conversion (e.g. Kyrou *et al.* 2018). Several studies (Tanaka *et al.* 2017; Min *et al.* 2018) have
recommended the use of spatially and/or temporally limited threshold-dependent homing drives,
because they are less likely to spread into non-target populations. However, we emphasize that it might
be difficult in practice to implement a threshold-dependent drive whose threshold remains as expected
for several reasons. First, theoretical models show that the range of parameter values for threshold-
dependent gene drives is larger when conversion occurs in the zygote than when it occurs in the
germline (compare Figures 1 and 4 in Deredec *et al.* 2008; Figure S1-S2). So ideally, it might be better
to use drives with conversion in the zygote. Nevertheless, such drives are more difficult to create and
so far all homing drives have been engineered with germline promoters (Table 2 in Courtier-Orgogozo
et al. 2019b). A few conserved genes are expressed in the germline of all animals (*nanos*, *vasa*, *piwi*;
Extavour and Akam 2003; Juliano *et al.* 2010) and their promoters constitute preferred tools for
engineering gene drive constructs in various animal species, in contrast to zygotically expressed genes,
which tend to be less conserved across taxa (Heyn *et al.* 2014). Second, "real life" ecological conditions
are likely to alter the genetic parameters of any gene drive, in particular its fitness cost. Fitness costs
are difficult to estimate in the field and can vary either across genomic backgrounds, spatially or
temporally (Marshall and Hay 2012; Backus and Delborne 2019). Hence, depending on ecological
conditions, the threshold-value for the invasion of a threshold-dependent homing drive could change,
or even decrease to 0. Thus, a homing drive that is threshold-dependent in the laboratory might turn
into a threshold-independent drive in the wild.

Conclusion

Our model is a step towards the development of more complex analytical models of gene drive that
account for the feedback between population demography and evolution. Our results suggest that the
recessive eradication drives with germline conversion currently developed in mosquitoes (e.g. Kyrou *et*
al. 2018) are likely to be threshold-independent and could be particularly difficult to control using
brakes. In addition, our results show that a brake that carries a version of the gene disrupted by the
initial gene drive, and therefore restores fitness, can prevent the extinction of the target population under
certain conditions. We think that the development of countermeasures should go in par with the
development of drives. Given the diversity of outcomes that we find and the difficulty to precisely
estimate the relevant parameters determining each outcome, specific experimental studies will be
necessary to confirm modelling outcomes that a given brake can indeed stop the spread of drives. A
brake should not be considered reliable before population experiments are carried out.

Acknowledgements:

We thank Arnaud Martin for discussions. Funding for this project was provided by European Research
Council (FP7/2007-2013 Grant Agreement no. 337579) to VCO, ANR-14-ACHN-0003 and ANR-19-

CE45-0009-01 to FD and the CeMEB LabEx/University of Montpellier (ANR-10-LABX-04-01) and
an INRAE-SPE starting grant to NOR.

Authors contributions: VCO brought the research topic, all authors developed the model, FD did the
analysis, implemented numerical solutions, ran stochastic simulations and prepared the figures. All
authors analysed data and wrote the manuscript.

Literature cited

Akbari O. S., K. D. Matzen, J. M. Marshall, H. Huang, C. M. Ward, *et al.*, 2013 A synthetic gene

drive system for local, reversible modification and suppression of insect populations. Curr.

Biol. 23: 671–677.

Akbari O. S., H. J. Bellen, E. Bier, S. L. Bullock, A. Burt, *et al.*, 2015 Safeguarding gene drive

experiments in the laboratory. Science 349: 927–929.

Alphey N., and M. B. Bonsall, 2014 Interplay of population genetics and dynamics in the genetic

control of mosquitoes. J. R. Soc. Interface 11: 20131071.

Backus G. A., and J. A. Delborne, 2019 Threshold-dependent gene drives in the wild: spread,

controllability, and ecological uncertainty. BioScience 69: 900–907.

Bull J. J., C. H. Remien, R. Gomulkiewicz, and S. M. Krone, 2019 Spatial structure undermines

parasite suppression by gene drive cargo, (D. Lahr, Ed.). PeerJ 7: e7921.

https://doi.org/10.7717/peerj.7921

Burt A., 2003 Site-specific selfish genes as tools for the control and genetic engineering of natural

populations. Proc. R. Soc. B Biol. Sci. 270: 921–928. https://doi.org/10.1098/rspb.2002.2319

Champer J., A. Buchman, and O. S. Akbari, 2016 Cheating evolution: engineering gene drives to

manipulate the fate of wild populations. Nat. Rev. Genet. 17: 146–159.

https://doi.org/10.1038/nrg.2015.34

Champer J., J. Liu, S. Y. Oh, R. Reeves, A. Luthra, *et al.*, 2018 Reducing resistance allele formation

in CRISPR gene drive. Proc. Natl. Acad. Sci. 115: 5522–5527.

https://doi.org/10.1073/pnas.1720354115

Champer J., E. Lee, E. Yang, C. Liu, A. G. Clark, *et al.*, 2020 A toxin-antidote CRISPR gene drive

system for regional population modification. Nat. Commun. 11: 1082.

https://doi.org/10.1038/s41467-020-14960-3

Courtier-Orgogozo V., A. Danchin, P.-H. Gouyon, and C. Boëte, 2019a Evaluating the Probability of

CRISPR-based Gene Drive Contaminating Another Species. bioRxiv 776609.

Courtier-Orgogozo V., A. Danchin, P.-H. Gouyon, and C. Boëte, 2019b Evaluating the Probability of

CRISPR-based Gene Drive Contaminating Another Species. BioRxiv 776609.

Del Amo V. L., A. L. Bishop, J. B. Bennett, X. Feng, J. M. Marshall, *et al.*, 2020 A

transcomplementing gene drive provides a flexible platform for laboratory investigation and

potential field deployment. Nat. Commun. 11: 1–12.

Deredec A., A. Burt, and H. C. J. Godfray, 2008 The Population Genetics of Using Homing

Endonuclease Genes in Vector and Pest Management. Genetics 179: 2013–2026.

https://doi.org/10.1534/genetics.108.089037

DiCarlo J. E., A. Chavez, S. L. Dietz, K. M. Esvelt, and G. M. Church, 2015 Safeguarding CRISPR-

Cas9 gene drives in yeast. Nat. Biotechnol. 33: 1250–1255. https://doi.org/10.1038/nbt.3412

Drury D. W., A. L. Dapper, D. J. Siniard, G. E. Zentner, and M. J. Wade, 2017 CRISPR/Cas9 gene

drives in genetically variable and nonrandomly mating wild populations. Sci. Adv. 8.

Edgington M. P., T. Harvey-Samuel, and L. Alphey, 2020 Population-level multiplexing, a promising

strategy to manage the evolution of resistance against gene drives targeting a neutral locus.

Evol. Appl. n/a. https://doi.org/10.1111/eva.12945

Esvelt K. M., A. L. Smidler, F. Catteruccia, and G. M. Church, 2014 Emerging technology:

concerning RNA-guided gene drives for the alteration of wild populations. Elife 3: e03401.

Extavour C. G., and M. Akam, 2003 Mechanisms of germ cell specification across the metazoans:

epigenesis and preformation. Development 130: 5869–5884.

Gantz V. M., and E. Bier, 2016 The dawn of active genetics. BioEssays 38: 50–63.

https://doi.org/10.1002/bies.201500102

Girardin L., V. Calvez, and F. Débarre, 2019 Catch me if you can: a spatial model for a brake-driven

gene drive reversal. Bull. Math. Biol. 81: 5054–5088.

Gurwitz D., 2014 Gene drives raise dual-use concerns. Science 345: 1010–1010.

Heyn P., M. Kircher, A. Dahl, J. Kelso, P. Tomancak, *et al.*, 2014 The earliest transcribed zygotic

genes are short, newly evolved, and different across species. Cell Rep. 6: 285–292.

Juliano C. E., S. Z. Swartz, and G. M. Wessel, 2010 A conserved germline multipotency program.

Development 137: 4113–4126.

Kyrou K., A. M. Hammond, R. Galizi, N. Kranjc, A. Burt, *et al.*, 2018 A CRISPR–Cas9 gene drive

targeting doublesex causes complete population suppression in caged Anopheles gambiae

mosquitoes. Nat. Biotechnol. https://doi.org/10.1038/nbt.4245

Leitschuh C. M., D. Kanavy, G. A. Backus, R. X. Valdez, M. Serr, *et al.*, 2018 Developing gene drive

technologies to eradicate invasive rodents from islands. J. Responsible Innov. 5: S121–S138.

https://doi.org/10.1080/23299460.2017.1365232

Li F., and M. J. Scott, 2016 CRISPR/Cas9-mediated mutagenesis of the white and Sex lethal loci in

the invasive pest, Drosophila suzukii. Biochem. Biophys. Res. Commun. 469: 911–916.

Marshall J. M., and B. A. Hay, 2012 Confinement of gene drive systems to local populations: a

comparative analysis. J. Theor. Biol. 294: 153–171.

Marshall J. M., and O. S. Akbari, 2018 Can CRISPR-Based Gene Drive Be Confined in the Wild? A

Question for Molecular and Population Biology. ACS Chem. Biol. 13: 424–430.

https://doi.org/10.1021/acschembio.7b00923

Montenegro de Wit M., 2019 Gene driving the farm: who decides, who owns, and who benefits?

Agroecol. Sustain. Food Syst. 43: 1054–1074.

Nagylaki T., and J. F. Crow, 1974 Continuous selective models. Theor. Popul. Biol. 5: 257–283.

NASEM, 2016 *Gene Drives on the Horizon: Advancing Science, Navigating Uncertainty, and*

Aligning Research with Public Values. National Academies Press, Washington, D.C.

Noble C., J. Olejarz, K. M. Esvelt, G. M. Church, and M. A. Nowak, 2017 Evolutionary dynamics of

CRISPR gene drives. Sci. Adv. 8.

Noble C., B. Adlam, G. M. Church, K. M. Esvelt, and M. A. Nowak, 2018 Current CRISPR gene

drive systems are likely to be highly invasive in wild populations. eLife 7: e33423.

Noble C., J. Min, J. Olejarz, J. Buchthal, A. Chavez, *et al.*, 2019 Daisy-chain gene drives for the

alteration of local populations. Proc. Natl. Acad. Sci. 116: 8275–8282.

Oberhofer G., T. Ivy, and B. A. Hay, 2018 Behavior of homing endonuclease gene drives targeting

genes required for viability or female fertility with multiplexed guide RNAs. Proc. Natl.

Acad. Sci. 115: E9343. https://doi.org/10.1073/pnas.1805278115

Oye K. A., K. Esvelt, E. Appleton, F. Catteruccia, G. Church, *et al.*, 2014 Regulating gene drives.

Science 345: 626–628. https://doi.org/10.1126/science.1254287

Raban R. R., J. M. Marshall, and O. S. Akbari, 2020 Progress towards engineering gene drives for

population control. J. Exp. Biol. 223.

Rode N. O., A. Estoup, D. Bourguet, V. Courtier-Orgogozo, and F. Débarre, 2019 Population

management using gene drive: molecular design, models of spread dynamics and assessment

of ecological risks. Conserv. Genet. 20: 671–690. https://doi.org/10.1007/s10592-019-01165-

5

Scott M. J., F. Gould, M. Lorenzen, N. Grubbs, O. Edwards, *et al.*, 2018 Agricultural production:

assessment of the potential use of Cas9-mediated gene drive systems for agricultural pest

control. J. Responsible Innov. 5: S98–S120. https://doi.org/10.1080/23299460.2017.1410343

Tanaka H., H. A. Stone, and D. R. Nelson, 2017 Spatial gene drives and pushed genetic waves. Proc.

Natl. Acad. Sci. 114: 8452–8457.

Unckless R. L., P. W. Messer, T. Connallon, and A. G. Clark, 2015 Modeling the Manipulation of

Natural Populations by the Mutagenic Chain Reaction. Genetics 201: 425–431.

https://doi.org/10.1534/genetics.115.177592

Unckless R. L., A. G. Clark, and P. W. Messer, 2017 Evolution of Resistance Against CRISPR/Cas9

Gene Drive. Genetics 205: 827–841. https://doi.org/10.1534/genetics.116.197285

Vella M. R., C. E. Gunning, A. L. Lloyd, and F. Gould, 2017 Evaluating strategies for reversing

CRISPR-Cas9 gene drives. Sci. Rep. 7. https://doi.org/10.1038/s41598-017-10633-2

Webster S. H., M. R. Vella, and M. J. Scott, 2019 Development and testing of a novel Killer-Rescue

self-limiting gene drive system in *Drosophila melanogaster*. bioRxiv 680629.

https://doi.org/10.1101/680629

Wu B., L. Luo, and X. J. Gao, 2016 Cas9-triggered chain ablation of cas9 as a gene drive brake. Nat.

Biotechnol. 34: 137–138. https://doi.org/10.1038/nbt.3444

Appendix

In the main text, the change over time in the density of individuals of genotype g is given by

$$\frac{dN_g}{dt} = \omega_g V_g N(1 - N/K) - d_g N_g.$$

We provide below the expressions for V_g for the two timings of gene conversion that we consider in the article.

Germline conversion

When gene conversion takes place in the germline, individuals born heterozygous remain heterozygous as adults, their life-history parameters are those of heterozygotes, but then gene conversion takes place in the germline, and if successful, predominantly one type of gamete is produced by the individual. We have

$$V_{00} = \frac{\gamma_0^2}{N^2}, V_{0D} = \frac{2\gamma_0\gamma_D}{N^2}, V_{DD} = \frac{\gamma_D^2}{N^2}, V_{0B} = \frac{2\gamma_0\gamma_B}{N^2}, V_{DB} = \frac{2\gamma_D\gamma_B}{N^2}, V_{BB} = \frac{\gamma_B^2}{N^2},$$

where

$$\gamma_0 = \beta_{00}N_{00} + \frac{1}{2}\beta_{0D}N_{0D}(1 - c_D) + \frac{1}{2}\beta_{0B}N_{0B},$$

$$\gamma_D = \beta_{DD}N_{DD} + \frac{1}{2}\beta_{0D}N_{0D}(1 + c_D) + \frac{1}{2}\beta_{DB}N_{DB}(1 - c_B),$$

$$\gamma_B = \beta_{BB}N_{BB} + \frac{1}{2}\beta_{0B}N_{0B} + \frac{1}{2}\beta_{DB}N_{DB}(1 + c_B).$$

Zygote conversion

When conversion takes place in zygotes, and when gene conversion is successful, an initially heterozygous zygote becomes homozygous, and develops into a homozygous adult. We have

$$V_{00} = \frac{\gamma_0^2}{N^2}, V_{0D} = (1 - c_D)\frac{2\gamma_0\gamma_D}{N^2}, V_{DD} = c_D\frac{2\gamma_0\gamma_D}{N^2} + \frac{\gamma_D^2}{N^2},$$

$$V_{0B} = \frac{2\gamma_0\gamma_B}{N^2}, V_{DB} = (1 - c_B)\frac{2\gamma_D\gamma_B}{N^2}, V_{BB} = c_B\frac{2\gamma_D\gamma_B}{N^2} + \frac{\gamma_B^2}{N^2},$$

where

$$\gamma_0 = \beta_{00}N_{00} + \frac{1}{2}\beta_{0D}N_{0D} + \frac{1}{2}\beta_{0B}N_{0B},$$

$$\gamma_D = \beta_{DD}N_{DD} + \frac{1}{2}\beta_{0D}N_{0D} + \frac{1}{2}\beta_{DB}N_{DB},$$

$$\gamma_B = \beta_{BB}N_{BB} + \frac{1}{2}\beta_{0B}N_{0B} + \frac{1}{2}\beta_{DB}N_{DB}.$$

Figures

Figure 1: Life-cycles with the two timings of gene conversion, germline (a) and zygote (b). The blue color corresponds to the wild-type allele, the red color to the drive allele and drive-homozygous individuals; the drive/wild-type heterozygous individual is represented in purple. The tombstone represents death. Notation: 0: WT, D: drive; c probability of gene conversion; ω: zygote survival; d: adult mortality; β: adult fecundity.

Conversion in the germline, brake does not restore fitness

Figure 2: Deterministic dynamics of the frequencies of each allele in the population, and scaled total population size (black curve). Conversion takes place in the germline, and the brake does not restore fitness. Population size is scaled relative to the equilibrium size of a 100% wild-type population $\left(K\left(1-d_{00}/(\beta_{00}^2\omega_{00})\right)\right)$. The arrow indicates the timing of drive introduction, here chosen to be when the drive allele is at 50% ($f_I = 0.5$). A cross indicates population extinction.

Figure 3: Frequency of each type of outcome in the simulations (color-coded), depending on the frequency of drive f_I at the time at which the drive is introduced (horizontal axis), on the dominance coefficient h (columns) and on the trait that is affected by the drive and the brake (rows). The dots show, with the same color code, the output of the deterministic model.

Figure 4: Frequency of each type of outcome in the simulations (color-coded), depending on the frequency of drive f_I at the time at which the drive is introduced (horizontal axis), on the dominance coefficient h (columns) and on the trait that is affected by the drive and the brake (rows). The dots show, with the same color code, the output of the deterministic model.

Figure 5: Frequency of each type of outcome in the simulations (color-coded), depending on the frequency of drive f_I at the time at which the drive is introduced (horizontal axis), on the dominance coefficient h (columns) and on the trait that is affected by the drive and the brake (rows). The dots show, with the same color code, the output of the deterministic model.

Figure 6: Frequency of each type of outcome in the simulations (color-coded), depending on the frequency of drive f_I at the time at which the drive is introduced (horizontal axis), on the dominance coefficient h (columns) and on the trait that is affected by the drive and the brake (rows). The dots show, with the same color code, the output of the deterministic model.

$$
\begin{array}{rl}
K & 25000 \\
c_D & 0.9 \\
c_B & 0.8 \\
N_{0D}^{(0)} & 1000 \\
N_{0B}^{(0)} & 100 \\
t_{\max} & 2500
\end{array}
$$

Table S1: Fixed parameters

$$
\begin{array}{rl}
f_I & \{0.025, 0.1375, 0.25, 0.375, 0.5, 0.625, 0.75, 0.8625, 0.975\} \\
h_{D0} = h_{B0} = h_{DB} = h & \{0, 0.5\} \\
convType & \{Z, G\}
\end{array}
$$

Table S2: Varying parameters

Scenario #	(1)	(2)	(3)	(4)	(5)	(6)
Brake…	does not restore fitness			restores fitness		
Effects on	d	ω	β	d	ω	β
d_{00}	0.6	0.6	0.6	0.6	0.6	0.6
d_{DD}	1.1	0.6	0.6	1.1	0.6	0.6
d_{BB}	1.1	0.6	0.6	0.64	0.6	0.6
ω_{00}	1.0	1.0	1.0	1.0	1.0	1.0
ω_{DD}	1.0	0.545	1.0	1.0	0.545	1.0
ω_{BB}	1.0	0.545	1.0	1.0	0.938	1.0
β_{00}	1.0	1.0	1.0	1.0	1.0	1.0
β_{DD}	1.0	1.0	0.738	1.0	1.0	0.738
β_{BB}	1.0	1.0	0.738	1.0	1.0	0.968

Table S3: Parameters for the different scenarios, depending on whether the brake restores fitness (modulo a small cost) or not, and on which life-history parameter is affected (adult survival d, zygote survival ω, adult fecundity β).

Eradication drive

Conversion in the germline   Conversion in the zygote

Figure S1: Local stabilities of the drive-only and the wild-type only equilibria in the absence of brake, for an eradication drive. The wild-type only equilibrium is locally stable in the blue-shaded region left of the blue curve; the drive-only equilibrium is locally stable in the red-shaded region right of the red curve. Neither equilibrium is locally stable in the white area, in which the two alleles coexist. Both equilibria are locally stable in the purple area; the final outcome depends on the initial conditions (bistability). Drives whose parameters put them in the purple area are threshold-dependent.

Figure S2: Local stabilities of the drive-only and the wild-type only equilibria in the absence of brake, for a replacement drive. The legend is the same as figure S1.

Conversion in the germline, brake restores fitness

Figure S3: Same legend as figure 2.

Conversion in the zygote, brake does not restore fitness

Figure S4: Same as figure 2

Conversion in the zygote, brake restores fitness

Figure S5: Same as figure 2

Conversion in the germline

Brake does not restore fitness　**Brake restores fitness**

Figure S6: Deterministic dynamics when the drive is threshold-dependent; conversion takes place in the germline. Parameters are the same as in the other figures, except for the dominance parameter ($h = 1$) and for conversion efficiencies ($c_D = 0.3$, $c_B = 0.25$ in panels (a)–(b); $c_D = 0.6$, $c_B = 0.55$ in panels (c)–(d); $c_D = 0.5$, $c_B = 0.45$ in panels (e)–(f)). Introduction densities are $N_{0D} = 10^5$ and $N_{0B} = 10^4$.

Figure S7: Deterministic dynamics when the drive is threshold-dependent; conversion takes place in the zygote. See figure S6 for parameter values.