Technical Report: Targeted Proteomic Analysis Reveals Enrichment of Atypical Ubiquitin Chains in
Contractile Murine Tissues

Tiaan Heunis[1,2], Frederic Lamoliatte[1,2], José Luis Marín-Rubio[1], Abeer Dannora[1] and Matthias
Trost[1]*

[1]Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4HH, United Kingdom.

[2]These authors contributed equally to this work

*Corresponding author:

Matthias Trost, Tel: +44 191 2087009, E-mail: matthias.trost@ncl.ac.uk

Summary

Ubiquitylation is an elaborate post-translational modification involved in all biological processes. Its
pleotropic effect is driven by the ability to form complex polyubiquitin chain architectures that can
influence biological functions. In this study, we optimised sample preparation and chromatographic
separation of Ubiquitin peptides for Absolute Quantification by Parallel Reaction Monitoring (Ub-
AQUA-PRM). Using this refined Ub-AQUA-PRM assay, we were able to quantify all ubiquitin chain
types in 10-minute LC-MS/MS runs. We used this method to determine the ubiquitin chain-linkage
composition in murine bone marrow-derived macrophages and different mouse tissues. We could
show tissue-specific differences in ubiquitin levels in murine tissues, with polyubiquitin chain types
contributing a small proportion to the total pool of ubiquitin. Interestingly, we observed enrichment
of atypical (K33) ubiquitin chains in heart and muscle. Our approach enabled high-throughput
screening of ubiquitin chain-linkage composition in different murine tissues and highlighted a
possible role for atypical ubiquitylation in contractile tissues.

Main Text

Ubiquitylation is a post-translational modification that results from the addition of ubiquitin, a 76
amino acid protein, to generally the ε-amino group of a lysine residue. Ubiquitylation plays a role in
practically all eukaryotic biological processes. It is therefore not surprising that aberrant

ubiquitylation is linked to several human diseases, including cancer, infectious, neurodegenerative,

and developmental diseases [1]. Protein substrates can be modified by monoubiquitin or can be

polyubiquitylated, which is formed by further ubiquitylation of any lysine (K6, K11, K27, K29, K33,

K48, and K63) or the N-terminal methionine (M1) of ubiquitin. This allows formation of complex

chain architectures (homotypic, heterotypic, or branched chains), which can ultimately influence a

substrates fate [2]. However, the biological functions of atypical and mixed ubiquitin chains are only

beginning to emerge [1,3]. Therefore, methods are needed to rapidly screen the ubiquitin chain-

linkage landscape in cells and tissues to advance our understanding of ubiquitin signalling, especially

for atypical linkage types as there are few tools available that facilitate their analysis [4]. Recent

technological advances in mass spectrometry-based proteomics have enabled quantification of

endogenous levels of ubiquitin chain types by selected reaction monitoring (Ub-AQUA-SRM) [5]. This

approach has subsequently been adapted to quadrupole-Orbitrap instruments using parallel

reaction monitoring (PRM) [6] to determine chain-linkage composition of purified substrates [7],

whole-cell lysates [8], to quantify K48-K63 branched chains [9,10], to gain information on higher-

order chain-linkage architecture of ubiquitylated substrates [7] and to quantify phosphorylated and

acetylated ubiquitin [11,12]. Here we further refined the Ub-AQUA-PRM method for absolute

quantification of ubiquitin chain types. We optimised peptide oxidation, ion-pairing agents, and

implemented microflow chromatographic separation for increased assay sensitivity and throughput.

Our refined Ub-AQUA-PRM assay is ideal for high-throughput screening of ubiquitin chain types in

diverse samples, as demonstrated here using murine primary cells and tissues.

Methionine-containing peptides can be challenging to analyse by PRM-based methods, as

they can occur in different oxidation states [13]. This can become problematic for quantification

when the peptide signal is divided across multiple oxidation states or, importantly, when there are

differences between the endogenous peptide and the spike-in standard. Current Ub-AQUA-PRM

strategies analyse the methionine sulfoxide version of methionine-containing peptides, induced by

using low concentrations of H_2O_2 at 4°C [13,14]. However, these peptides are stable for only 21.5

hours, before oxidation state shifts are observed [13]. We therefore fully oxidised methionine

residues into the sulfone derivative using H_2O_2. No difference in methionine sulfone-containing

peptide (M1 or K6) intensity was observed without H_2O_2 at room temperature or 60°C for 2 hours

(Supplementary Figure 1a, b). However, a methionine sulfone was generated when using 0.1% H_2O_2

at 60°C for 2 hours, but the sulfoxide version still contributed substantially to M1 and K6 peptide

intensity. Both methionine-containing peptides (M1 and K6) were converted into a stable

methionine sulfone by more than 99.9%, based on peptide intensity, when using 1% H_2O_2 for 2 hours

at 60°C (Supplementary Figure S1a, b). The other peptides analysed were unaffected under the same

conditions (Supplementary Figure S1c). This indicates that our oxidation procedure does not

introduce undesirable artifacts. We, therefore, used 1% H_2O_2 for 2 hours at 60°C for peptide

oxidation in all further analyses. Next, we determined the optimal normalised collision energy (NCE)

required for fragmenting each peptide. Detailed information on NCE used for each peptide is

available in Supplementary Table S1. Additionally, we also tested the effect of two commonly used

ion-pairing agents in reversed-phase chromatography for their effect on peptide retention and ion

intensity, namely formic acid (FA) and trifluoroacetic acid (TFA). We compared FA and TFA at

concentrations ranging from 0.2 to 5.0% (v:v) in the loading buffer (Supplementary Figure S2). For all

these injections, buffer A and B remained the same. The K29 peptide is the most hydrophilic of the

ubiquitin peptides, but we did not observe a major difference in peak intensity or peak shape

between 5.0% FA and 0.2% TFA for this peptide (Supplementary Figure S2a). However, we observed

a marked decrease in intensity for other ubiquitin peptides when using TFA, even as low as 0.2%

(Supplementary Figure S2b). We, therefore, used 5.0% FA in the sample buffer for all further

analyses. Finally, we determined the lower limit of detection (LLOD) and lower limit of quantification

(LLOQ) of each peptide in a simple (ubiquitin peptides only) and complex (*E. coli* tryptic digest)

matrix. We opted for an *E. coli* tryptic digested as spike-in matrix as no endogenous ubiquitin is

present in this organism that would interfere with the assay. LLOD for peptides were as low as 0.5

amol on column. In the simple matrix we could achieve LLOQ ranging from 50 amol for the M1 and

K29 peptides to 1.5 fmol for the K11, K63 and TITTITLEVEPSDTIENVK ubiquitin peptides. Similar

results were obtained for ubiquitin peptides in the complex matrix (Supplementary Figure S3),

reaching LLOQ as low as 0.1 fmol/μg protein injected. These results indicate that our refined Ub-

AQUA-PRM assay could accurately quantify low levels of endogenous ubiquitin chain types in

complex matrixes.

We used our Ub-AQUA-PRM assay to screen the ubiquitin chain-linkage composition in

primary murine macrophages (BMDMs), given the importance of ubiquitin in innate immunity [15–

18]. Additionally, we validated our Ub-AQUA-PRM assay using MG132, a proteasome inhibitor that

results in ubiquitin accumulation, especially K48 chains [19,20]. An increase in the percentage of

polyubiquitin chains was observed in BMDMs after treatment with MG132, as would be expected

(Figure 1a, b). All linkage types could be detected in murine BMDMs with K48 (63.2%), K63 (24.2%)

and K29 (8.0%) contributing 95.4% of the total ubiquitin chain-linkage composition of whole-cell

lysates (Figure 1c). These results are in agreement with previous reports in cell lines [7,21]. Similarly,

K48 (71.8%), K63 (15.5%) and K29 (5.2%) contributed 92.5% of the total ubiquitin chain-linkage

composition of BMDMs during MG132 treatment (Figure 1c). Next, we determined whether there

was any difference in ubiquitin chain types during treatment with MG132 for 6 hours. We observed

a statistically significant increase in the majority of ubiquitin chain types, including total ubiquitin,

during MG132 treatment (Supplementary Figure S4), which is in agreement with previous reports

[7,19,22]. However, K11 and K27 chain types were below the limit of quantification during our

analysis. These results indicate that K48, K63 and K29 linkage types dominate the ubiquitin chain-

linkage compositions in murine BMDM whole-cell lysates and that up to ~29.2% of total ubiquitin is

in the polyubiquitylated form in resting BMDMs.

Lastly, we used our Ub-AQUA-PRM assay to screen the ubiquitin chain-linkage composition

of six different murine tissues (brain, heart, kidney, lung, muscle, and spleen) from three different

mice. We were able to detect ubiquitin in all samples, with the brain, kidney, and spleen having the

highest levels of total ubiquitin (Supplementary Figure S5). A statistically significant difference in

total ubiquitin was observed in the brain, which had higher levels of total ubiquitin than the heart,

lung, and muscle (Supplementary Figure S5). Interestingly, only a small percentage of total ubiquitin

seems to be polyubiquitylated in murine tissues, with the contribution of chain types to total

ubiquitin being ~1% in most tissues (Figure 2a). The lung had ~1.4% of total ubiquitin in the

polyubiquitylated form, whereas muscle had the highest with ~8.7%. In most tissues, K48 was the

most dominant chain type (Figure 2b), similar to what we observed in BMDMs. However, we

observed differences in ubiquitin chain-linkage composition in heart and muscle. Interestingly, K33

was enriched in heart and muscle, compared to the other tissues (Figure 2b). In order to determine

the percentage contribution of the K33 linkage type to the total pool of ubiquitin, we normalised this

chain type to total ubiquitin in each tissue. K33 contributed only a small proportion to the total pool

of ubiquitin, with 5.3 ± 1.9% in muscle and 3.0 ± 1.3% in heart. We verified that these results were

not an artefact of our experimental procedure by using a different extraction buffer before Ub-

AQUA-PRM analysis. We observed similar results in muscle when using TCA extraction buffer in an

independent experiment, with K33 contributing to 5.7 ± 0.9% of the total pool of ubiquitin.

Noteworthy, we observed a slightly lower contribution of K33 to the total ubiquitin chain-linkage

composition when using TCA extraction, but K33 still contributed >56% to the total ubiquitin chain-

linkage composition in muscle (Supplementary Figure S6). This discrepancy is likely due to the

different extraction buffer used, which could affect the protein composition extracted from the

tissues.

These results indicate that there are tissue-specific differences in total ubiquitin and

ubiquitin chain types in murine tissues, with atypical linkages increased in heart and muscle.

Furthermore, our data indicate that only a small proportion of total ubiquitin in murine tissues are in

the polyubiquitin form. It is, therefore, possible that many proteins in murine tissues are either

monoubiquitylated, or that the majority of ubiquitin is in its unconjugated, free form. To test this,

we performed immunoblots for total ubiquitin (conjugated and unconjugated) using the same

tissues analysed by Ub-AQUA-PRM. We observed high levels of unconjugated ubiquitin in the

majority of tissues (Supplementary Figure S7). However, we did observe conjugated ubiquitin in all

tissues. More unconjugated ubiquitin was observed in the brain, kidney, lung, and spleen, whereas

heart and muscle had the least amount of unconjugated ubiquitin. In our Ub-AQUA-PRM assay, we

also observed lower levels of total ubiquitin in the heart and muscle, which corresponds to the data

obtained with the western blots. This data indicates that there are major tissue-specific differences

in ubiquitylation and that only a small proportion of total ubiquitin in murine tissues are in the

conjugated form. Further analysis of the ubiquitylome and ubiquitin chain-linkage landscape, in

combination with recently described methods such as Ub-clipping [21] and Ubiquitin Chain

Enrichment Middle-Down Mass Spectrometry (UbiChEM-MS) [23], could reveal detailed information

on atypical ubiquitylation and higher-order chain-linkage architecture in different tissues and

pathological states.

Methods – online methods

Preparation of AQUA peptides

Synthetic ubiquitin peptides were obtained from Cambridge Research Biochemicals (Cleveland,

United Kingdom) (Supplementary Table S1). Methionine-containing peptides were synthesised under

their reduced form, and were subsequently oxidised during sample preparation, as described below.

Peptides were dissolved at a concentration of 1 nmol/µl in 50:50 acetonitrile:water (v:v) and peptide

purity was monitored by LC-MS/MS. Peptides lacking net peptide content were injected together

with their isotope counterpart and net peptide content was determined based on the ratio between

the two isotopes. Two sets of master mixes were prepared, with the 10 light and 10 heavy peptides

mixed, separately, in equimolar concentrations. The light and heavy peptide master mixes were

diluted, separately, to 10 pmol/µl in HPLC-grade water (Thermo Fisher Scientific) and stored at -

80°C.

Optimization of methionine oxidation and ion-pairing agents for targeted proteomic analysis of

Ubiquitin AQUA peptides

Methionine oxidation was optimised to generate a stable methionine sulfone for targeted proteomic

analysis. Heavy and light peptides were mixed in a 1:1 ratio and incubated with different

concentrations of H_2O_2 at either room temperature (RT) or 60°C. The conversion of reduced

methionine to sulfoxide and sulfone derivatives were monitored by LC-MS/MS. The effect of ion-

pairing agents on peptide retention and intensity was assessed by using increasing concentrations of

formic acid (FA) and trifluoracetic acid (TFA) in the sample and loading buffers, as indicated in the

text.

Preparation of standard curves

The heavy master mix was diluted to 100 fmol/µl and the light master mix was serially diluted from

10 pmol/µl to 1 amol/µl using a dilution factor of $\sqrt{10}$. The standard curve was prepared on ice by

combining reaction components in the following order: 20 µl of HPLC-grade water, 10 µl of heavy

mix at 100 fmol/µl, 10 µl of light mix at varying concentrations, 50 µl of 10% FA, and 10 µl of 10%

H_2O_2. A second standard curve was prepared by substituting 10 µl of HPLC-grade water with 10 µl of

Escherichia coli tryptic digest at 1 µg/µl. The *E. coli* tryptic digest was generated as described below.

Samples were incubated for 2h at 60°C for oxidation, before targeted proteomic analysis.

Bacterial cell culture for proteomic analysis

E. coli DH5-α was cultured overnight in Luria-Bertani (Thermo Fisher Scientific) broth at 37°C with

shaking at 180 rpm. The next day, cells were harvested (4,000 xg, 15 min, 4°C), washed twice with

ice-cold PBS pH 7.4 and stored at -80°C until further analysis. Cell pellets were lysed in 5% SDS in 50

mM triethylammonium bicarbonate (TEAB) pH 7.5, by sonication. Whole-cell lysates were

subsequently centrifuged (10,000 xg, 5 min, RT) and the supernatants used for proteomic sample

preparation.

Murine cell culture and tissue collection

Animal work was approved by the Newcastle University ethical committee and performed under UK

Home Office project licence (PCI 23A338). Bone marrow-derived macrophages (BMDMs) were

derived from femurs of 8 to 12-week-old C57BL/6 mice. Bone marrow was extracted and

differentiated into macrophages using Iscove's Modified Dulbecco's Media (IMDM; Sigma-Aldrich)

supplemented with 10% heat-inactivated foetal calf serum (Thermo Fisher Scientific), 1% L-

glutamine (Lonza), 1% penicillin/streptomycin (Lonza) and 20% L929-conditioned media for 7 days at

37 °C in 5% CO_2 humidified atmosphere. After differentiation, BMDMs were stimulated with 10 µM

MG132 (Sigma-Aldrich) for 6 hours or mock treated. After stimulation, BMDMs were washed twice

with ice-cold PBS pH 7.4. Cells were subsequently lysed by sonication in 5% SDS in 50 mM TEAB pH

7.5, supplemented with 100 mM N-ethylmaleimide (NEM, Sigma-Aldrich) to inhibit deubiquitylases.

Whole-cell lysates were used for proteomic sample preparation.

198   Tissues (brain, heart, lung, kidney, muscle and spleen) were collected from 8 to 12-week-old

C57BL/6J mice, washed once with ice-cold PBS pH 7.4, flash frozen in liquid nitrogen and stored at -

80°C until further analysis. Tissues were ground to a powder in liquid nitrogen using a mortar and

pestle, and lysed by sonication in lysis buffer (5% SDS in 50 mM TEAB pH 7.5) supplemented with

100 mM NEM. Tissue lysates were subsequently centrifuged (10,000 xg, 5 min, RT) and the

supernatants used for methanol chloroform precipitation. Protein pellets were resuspended in lysis

buffer and used for proteomic sample preparation. In a separate experiment, muscle samples were

ground to a powder in liquid nitrogen using a mortar and pestle, and proteins extracted in 20%

trichloroacetic acid (TCA) at 4°C. Samples were centrifuged (10,000 xg, 10 min, RT) and washed twice

with ice-cold acetone to remove remaining TCA. Protein precipitates were subsequently

resuspended in lysis buffer, containing 100 mM NEM, centrifuged to remove debris, and processed

as described below.

Proteomics sample preparation

Protein concentration for all samples was determined using the bicinchoninic acid assay (BCA) assay

(Thermo Fisher Scientific). Fifty µg of total protein was used for proteomics sample preparation by

suspension trapping (S-Trap) [24,25], with minor modifications to that recommended by the supplier

(ProtiFi, Huntington NY, USA). Samples were reduced with 5 mM tris(2-carboxyethyl)phosphine

(Pierce) for 15 min at 37°C, and subsequently alkylated with 10 mM NEM for 15 min at room

temperature in the dark. Five µl of 12% phosphoric acid was added to each sample after reduction

and alkylation, followed by the addition of 330 µl S-Trap binding buffer (90% methanol in 100 mM

TEAB pH 7.1). Acidified samples were added, separately, to S-Trap mini-spin columns and

centrifuged at 4,000 xg for 1 minute. Each S-Trap mini-spin column was washed with 400 µl S-Trap

binding buffer and centrifuged at 4,000 xg for 1 minute. This process was repeated for five washes.

Twenty-five µl of 50 mM TEAB pH 8.0 containing sequencing-grade trypsin (1:10 ratio of

trypsin:protein) was added to each sample, followed by proteolysis for 2 hours at 47°C using a

thermomixer (Eppendorf). Peptides were eluted with 50 mM TEAB pH 8.0 and centrifugation at

1,000 xg for 1 minute. Elution steps were repeated using 0.2% FA and 0.2% FA in 50% acetonitrile,

respectively. The three eluates from each sample were combined and dried using a speed-vac before

storage at -80°C. Peptides were subsequently reconstituted to 1 µg/µl using 5% FA in water and

prepared for PRM, on ice, by combining reagents in the following order: 30 µl of HPLC-grade water,
10 µl peptide digest at 1 µg/µl, 10 µl of heavy master mix at 100 fmol/µl, 40 µl of 10% FA and 10 µl
of 10% H_2O_2. Samples were incubated for 2h at 60°C for oxidation before targeted mass
spectrometry analysis.

Targeted LC-MS/MS analysis - Parallel Reaction Monitoring (PRM)

Peptide samples were injected on a Dionex Ultimate 3000 RSLC (Thermo Fisher Scientific) connected
to a Q-Exactive HF mass spectrometer (Thermo Fisher Scientific). Fifty fmol of each ubiquitin peptide
was injected during analysis, along with 500 ng of tryptic peptide mixture from either *E. coli* or *Mus*
musculus as indicated in the text. In order to minimise dead volume and shorten the analysis to only
10 min, we used a direct injection LC setup. During the loading phase, samples are loaded using a 5
µl loop and full loop injection, with a 2.4 µl needle and an overfill ratio of 1.5. Samples were injected
onto a 150 µm x 150 mm EASY-Spray LC column (Thermo Fisher Scientific) using the nano-pump and
100% buffer A (96.9% HPLC-grade water, 3% DMSO, 0.1% FA) at 1.5 µl/min. After 4 min (6 µl), the
column oven valve was switched to bypass the loop and peptides were eluted by increased
concentration of buffer B (80% HPLC-grade acetonitrile, 16.9% HPLC-grade water, 3% DMSO, 0.1%
FA) using a multistep gradient (Supplementary Table S2). The source was heated to 320°C and 2.5 kV
was applied between the column emitter and the MS. The Q-Exactive HF was operated in PRM mode
using a resolution of 15,000, AGC target of 2×10^5 and maximum injection time of 25 ms. Peptides
were selected for MS/MS data acquisition using pre-determined isolation windows and fragmented
using collision energies described in Supplementary Table S1.

Targeted proteomic data analysis

All Ub-AQUA-PRM RAW mass spectrometry files were analysed using Skyline-daily (version
4.1.1.11871) [26]. Fragment ion intensities were extracted and light to heavy (L/H) ratios of peptide
pairs were automatically determined in Skyline. All extracted ion chromatograms were manually
inspected. Standard curves were generated from log transformed L/H ratios and peptide
concentrations, to determine the lower limit of detection (LLOD) and lower limit of quantification
(LLOQ) for each peptide. The R^2 was determined based on the LLOQ of each peptide. Only curves
with a R^2 value > 0.99 were used for determining the absolute concentration of total ubiquitin and
the different ubiquitin chain types. Group comparisons were performed using a Student's t-test, and
multiple group comparisons were performed using an analysis of variance (ANOVA), followed by a

Tukey's HSD test for determining significance for pairwise comparisons, in GraphPad Prism (version

8.0.1). The mass spectrometry proteomics data have been deposited to the ProteomeXchange [27]

Consortium via the PRIDE [28] partner repository with the dataset identifier PXD017754. Reviewers

can access it through the Username: reviewer39102@ebi.ac.uk and the password vJ56s8uk.

Western blots

Total protein from brain, heart, kidney, lung, muscle and spleen were extracted as described above.

Ten-microgram protein from each sample were separated on 12% SDS-PAGE gels before being

electrotransferred to Immobilon-P transfer membranes (Merck Millipore). Membranes were probed

with monoclonal antibody for mono-and polyubiquitylated proteins (FK2)(BML-PW8810; Enzo Life

Sciences), anti-α-Tubulin (T9026; Sigma-Aldrich) and anti-Histone H3 (D1H2) (4499; Cell Signaling)

antibodies. Corresponding secondary antibodies used included anti-mouse IgM, HRP-conjugated

(AP307P; Sigma-Aldrich) and anti-rabbit IgG, HRP-conjugated (7074; Cell Signaling Technology)

antibodies. Peroxidase activity was developed using the Amersham™ ECL™ Prime Western Blotting

Detection Reagent (GE Healthcare) and imaged on an Amersham Imager 600 digital imaging system

(GE Healthcare). To assess equal loading of protein, Ponceau staining (Sigma-Aldrich) of the PVDF

membranes was performed.

Acknowledgements

This work was funded by a Wellcome Trust Investigator Award (215542/Z/19/Z) and the generous

start-up funding of Newcastle University to MT. FL was funded by a Marie Skłodowska-Curie

Fellowship from the Horizon 2020 programme of the EU (project number 795670).

Conflict of interest

The authors declare no conflict of interest.

Author contributions

TH, FL and MT designed the study. TH and FL performed mass spectrometry experiments and data

analysis. JLM-R performed western blots. AD was responsible for animal husbandry and animal

experimental procedures. TH and MT wrote the paper with contribution of all authors.

References

[1]   H.J. Meyer, M. Rape, Enhanced protein degradation by branched ubiquitin chains, Cell. 157 (2014) 910–921. https://doi.org/10.1016/j.cell.2014.03.037.

[2]   K.N. Swatek, D. Komander, Ubiquitin modifications, Cell Res. 26 (2016) 399–422. https://doi.org/10.1038/cr.2016.39.

[3]   C.N. Cunningham, J.M. Baughman, L. Phu, J.S. Tea, C. Yu, M. Coons, D.S. Kirkpatrick, B. Bingol, J.E. Corn, USP30 and parkin homeostatically regulate atypical ubiquitin chains on mitochondria, Nat. Cell Biol. 17 (2015) 160–169. https://doi.org/10.1038/ncb3097.

[4]   R.E. Heap, M.S. Gant, F. Lamoliatte, J. Peltier, M. Trost, Mass spectrometry techniques for studying the ubiquitin system, Biochem. Soc. Trans. 45 (2017) 1137–1148. https://doi.org/10.1042/BST20170091.

[5]   D.S. Kirkpatrick, N.A. Hathaway, J. Hanna, S. Elsasser, J. Rush, D. Finley, R.W. King, S.P. Gygi, Quantitative analysis of in vitro ubiquitinated cyclin B1 reveals complex chain topology, Nat. Cell Biol. 8 (2006) 700–710. https://doi.org/10.1038/ncb1436.

[6]   H. Tsuchiya, K. Tanaka, Y. Saeki, The parallel reaction monitoring method contributes to a highly sensitive polyubiquitin chain quantification, Biochem. Biophys. Res. Commun. 436 (2013) 223–229. https://doi.org/10.1016/j.bbrc.2013.05.080.

[7]   H. Tsuchiya, D. Burana, F. Ohtake, N. Arai, A. Kaiho, M. Komada, K. Tanaka, Y. Saeki, Ub-ProT reveals global length and composition of protein ubiquitylation in cells, Nat. Commun. 9 (2018). https://doi.org/10.1038/s41467-018-02869-x.

[8]   T.P. Miettinen, J. Peltier, A. Härtlova, M. Gierliński, V.M. Jansen, M. Trost, M. Björklund, Thermal proteome profiling of breast cancer cells reveals proteasomal activation by CDK 4/6 inhibitor palbociclib, EMBO J. 37 (2018) 1–19. https://doi.org/10.15252/embj.201798359.

[9]   F. Ohtake, Y. Saeki, S. Ishido, J. Kanno, K. Tanaka, The K48-K63 branched ubiquitin chain regulates NF-κB signaling, Mol. Cell. 64 (2016) 251–266. https://doi.org/10.1016/j.molcel.2016.09.014.

[10]   F. Ohtake, H. Tsuchiya, Y. Saeki, K. Tanaka, K63 ubiquitylation triggers proteasomal degradation by seeding branched ubiquitin chains, Proc. Natl. Acad. Sci. 115 (2018) E1401–E1408. https://doi.org/10.1073/PNAS.1716673115.

320   [11]   F. Koyano, K. Okatsu, H. Kosako, Y. Tamura, E. Go, M. Kimura, Y. Kimura, H. Tsuchiya, H.
321   Yoshihara, T. Hirokawa, T. Endo, E.A. Fon, J.F. Trempe, Y. Saeki, K. Tanaka, N. Matsuda,
322   Ubiquitin is phosphorylated by PINK1 to activate parkin, Nature. 510 (2014) 162–166.
323   https://doi.org/10.1038/nature13392.

324   [12]   F. Ohtake, Y. Saeki, K. Sakamoto, K. Ohtake, H. Nishikawa, H. Tsuchiya, T. Ohta, K. Tanaka, J.
325   Kanno, Ubiquitin acetylation inhibits polyubiquitin chain elongation, EMBO Rep. 16 (2015)
326   192–201. https://doi.org/10.15252/embr.201439152.

327   [13]   L. Phu, A. Izrael-Tomasevic, M.L. Matsumoto, D. Bustos, J.N. Dynek, A. V. Fedorova, C.E.
328   Bakalarski, D. Arnott, K. Deshayes, V.M. Dixit, R.F. Kelley, D. Vucic, D.S. Kirkpatrick, Improved
329   quantitative mass spectrometry methods for characterizing complex ubiquitin signals, Mol.
330   Cell. Proteomics. 10 (2011) 1–19. https://doi.org/10.1074/mcp.M110.003756.

331   [14]   H. Tsuchiya, K. Tanaka, Y. Saeki, The parallel reaction monitoring method contributes to a
332   highly sensitive polyubiquitin chain quantification, Biochem. Biophys. Res. Commun. (2013).
333   https://doi.org/10.1016/j.bbrc.2013.05.080.

334   [15]   L. Deng, C. Wang, E. Spencer, L. Yang, A. Braun, J. You, C. Slaughter, C. Pickart, Z.J. Chen,
335   Activation of the Iκb kinase complex by TRAF6 requires a dimeric ubiquitin-conjugating
336   enzyme complex and a unique polyubiquitin chain, Cell. (2000).
337   https://doi.org/10.1016/S0092-8674(00)00126-4.

338   [16]   F. Tokunaga, S.I. Sakata, Y. Saeki, Y. Satomi, T. Kirisako, K. Kamei, T. Nakagawa, M. Kato, S.
339   Murata, S. Yamaoka, M. Yamamoto, S. Akira, T. Takao, K. Tanaka, K. Iwai, Involvement of
340   linear polyubiquitylation of NEMO in NF-κB activation, Nat. Cell Biol. 11 (2009) 123–132.
341   https://doi.org/10.1038/ncb1821.

342   [17]   B. Gerlach, S.M. Cordier, A.C. Schmukle, C.H. Emmerich, E. Rieser, T.L. Haas, A.I. Webb, J.A.
343   Rickard, H. Anderton, W.W.L. Wong, U. Nachbur, L. Gangoda, U. Warnken, A.W. Purcell, J.
344   Silke, H. Walczak, Linear ubiquitination prevents inflammation and regulates immune
345   signalling, Nature. 471 (2011) 591–596. https://doi.org/10.1038/nature09816.

346   [18]   M.U. Gack, Y.C. Shin, C.H. Joo, T. Urano, C. Liang, L. Sun, O. Takeuchi, S. Akira, Z. Chen, S.
347   Inoue, J.U. Jung, TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated
348   antiviral activity, Nature. 446 (2007) 916–920. https://doi.org/10.1038/nature05732.

349   [19]   F. Ohtake, H. Tsuchiya, Y. Saeki, K. Tanaka, K63 ubiquitylation triggers proteasomal
350   degradation by seeding branched ubiquitin chains, Proc. Natl. Acad. Sci. U. S. A. 115 (2018)

E1401–E1408. https://doi.org/10.1073/pnas.1716673115.

[20] K.L. Rock, C. Gramm, L. Rothstein, K. Clark, R. Stein, L. Dick, D. Hwang, A.L. Goldberg,

Inhibitors of the proteasome block the degradation of most cell proteins and the generation

of peptides presented on MHC class I molecules, Cell. (1994). https://doi.org/10.1016/S0092-

8674(94)90462-6.

[21] K.N. Swatek, J.L. Usher, A.F. Kueck, C. Gladkova, T.E.T. Mevissen, J.N. Pruneda, T. Skern, D.

Komander, Insights into ubiquitin chain architecture using Ub-clipping, Nature. (2019).

https://doi.org/10.1038/s41586-019-1482-y.

[22] H. Tsuchiya, F. Ohtake, N. Arai, A. Kaiho, S. Yasuda, K. Tanaka, Y. Saeki, In vivo ubiquitin

linkage-type analysis reveals that the Cdc48-Rad23/Dsk2 axis contributes to K48-linked chain

Ssecificity of the proteasome, Mol. Cell. 66 (2017) 488-502.e7.

https://doi.org/10.1016/j.molcel.2017.04.024.

[23] S.O. Crowe, A.S.J.B. Rana, K.K. Deol, Y. Ge, E.R. Strieter, Ubiquitin chain enrichment middle-

down mass spectrometry enables characterization of branched ubiquitin chains in cellulo,

Anal. Chem. 89 (2017) 4428–4434. https://doi.org/10.1021/acs.analchem.6b03675.

[24] A. Zougman, P.J. Selby, R.E. Banks, Suspension trapping (STrap) sample preparation method

for bottom-up proteomics analysis, Proteomics. 14 (2014) 1006–1010.

https://doi.org/10.1002/pmic.201300553.

[25] M. HaileMariam, R.V. Eguez, H. Singh, S. Bekele, G. Ameni, R. Pieper, Y. Yu, S-Trap, an

ultrafast sample-preparation approach for shotgun proteomics, J. Proteome Res. 17 (2018)

2917–2924. https://doi.org/10.1021/acs.jproteome.8b00505.

[26] B. MacLean, D.M. Tomazela, N. Shulman, M. Chambers, G.L. Finney, B. Frewen, R. Kern, D.L.

Tabb, D.C. Liebler, M.J. MacCoss, Skyline: An open source document editor for creating and

analyzing targeted proteomics experiments, Bioinformatics. 26 (2010) 966–968.

https://doi.org/10.1093/bioinformatics/btq054.

[27] E.W. Deutsch, A. Csordas, Z. Sun, A. Jarnuczak, Y. Perez-Riverol, T. Ternent, D.S. Campbell, M.

Bernal-Llinares, S. Okuda, S. Kawano, R.L. Moritz, J.J. Carver, M. Wang, Y. Ishihama, N.

Bandeira, H. Hermjakob, J.A. Vizcaíno, The ProteomeXchange consortium in 2017: Supporting

the cultural change in proteomics public data deposition, Nucleic Acids Res. 45 (2017) D1100–

D1106. https://doi.org/10.1093/nar/gkw936.

[28] Y. Perez-Riverol, A. Csordas, J. Bai, M. Bernal-Llinares, S. Hewapathirana, D.J. Kundu, A.

Inuganti, J. Griss, G. Mayer, M. Eisenacher, E. Pérez, J. Uszkoreit, J. Pfeuffer, T. Sachsenberg,

Ş. Yilmaz, S. Tiwary, J. Cox, E. Audain, M. Walzer, A.F. Jarnuczak, T. Ternent, A. Brazma, J.A.

Vizcaíno, The PRIDE database and related tools and resources in 2019: Improving support for

quantification data, Nucleic Acids Res. 47 (2019) D442–D450.

https://doi.org/10.1093/nar/gky1106.

Figure legends

Figure 1. All eight polyubiquitin chain types are present in murine bone marrow-derived

macrophages. Percentage of polyubiquitin chain types present in resting, unstimulated (a) and

MG132-treated (b) bone marrow-derived macrophages, compared to the total pool of ubiquitin.

Contribution of each of the eight different ubiquitin chain types to the total pool of polyubiquitin in

resting, unstimulated (c) and MG132-treated bone marrow-derived macrophages (d). Error bars

indicate standard deviation.

396
397 **Figure 2.** Tissue-specific differences in ubiquitin chain types and compositions exist in murine tissues.

398 (a) Percentage of polyubiquitin chain types present in different murine tissues, compared to the

399 total pool of ubiquitin present in these tissues. (b) Contribution of each of the eight different

     ubiquitin chain types to the total pool of polyubiquitin in different murine tissues. Error bars indicate

     standard deviation.

     Supporting documents

     Supplementary Figure S1. Generation of a stable methionine sulfone by oxidation. Methionine-

     containing ubiquitin peptides were oxidised using hydrogen peroxide (H_2O_2) and analysed by LC-

     MS/MS to determine the generation of a methionine sulfone. A stable methionine sulfone was

     generated using 1.0% H_2O_2 at 60°C for 2 hours in M1 (a) and K6 (b) peptides. (c) Peptide intensities

     observed for non-methionine-containing ubiquitin peptides when using H_2O_2 for oxidation. L = light

     version of the peptide, H = heavy version of the peptide, LoD = limit of detection. Error bars indicate

     standard deviation.

412
Supplementary Figure S2. Formic acid is a superior ion-pairing agent for ubiquitin peptides. (a)
Peptide intensities observed for M1, K6 and K29 ubiquitin peptides when using formic acid (FA) and
trifluoroacetic acid (TFA) as ion-pairing agent in the loading buffer. (b) Peptide intensities observed
for the other ubiquitin peptides when using FA and TFA in the loading buffer. L = light version of the
peptide, H = heavy version of the peptide, LoD = limit of detection. Error bars indicate standard
deviation.

419
Supplementary Figure S3. Limits of detection and quantification of spiked-in, heavy and light

ubiquitin peptides in an *E. coli* proteome digest. Standard curves were generated using 10 synthetic

ubiquitin peptides in their light and heavy counterparts. Heavy peptides were used at 10 fmol/µl,

while light peptides were serially diluted from 1 pmol/µl to 1 amol/µl. Peptides were oxidised using

H_2O_2 before injected on a Q-Exactive HF mass spectrometer for PRM analysis. The peptide master

mix contained each of the eight different ubiquitin chain-type peptides (a-h) and two peptides that

served as proxy for total ubiquitin (i, j). Error bars indicate standard deviation.

Supplementary Figure S4. MG132 treatment causes an increase in ubiquitin chain types. Differences observed in the quantified ubiquitin chain types at resting state and during MG132 treatment. Statistical significance is denoted with an * and indicates a p-value < 0.05 determined using a Student's t-test. Error bars indicate standard deviation.

Supplementary Figure S5. Murine tissues contain different levels of ubiquitin. Absolute concentration of ubiquitin in different murine tissues when using non-modified peptides as proxy for total ubiquitin. Total ubiquitin was derived from the sum of the absolute concentrations of TITLEVPSDTIENVK and K11 ubiquitin peptides. Statistical significance is denoted with an * and

indicates an adjusted p-value < 0.05 after a Tukey's HSD test and ANOVA. Error bars indicate

standard deviation.

Supplementary Figure S6. Mouse muscle is enriched in K33

ubiquitin chain types. (a) Contribution of each of the eight different ubiquitin chain types to the total

pool of polyubiquitin in mouse muscle when using 20% TCA for protein extraction to inhibit

deubiquitylase activity.

Supplementary Figure S7. Murine tissues contain a large amount of unconjugated ubiquitin. Western blot for total ubiquitin (conjugated and unconjugated) in murine tissues. The same tissues were used as those analysed by Ub-AQUA-PRM.

Supplementary Table S1. Peptides used for absolute quantification of ubiquitin chain types by Ub-AQUA-PRM analysis.

Supplementary Table S2. Multi-step gradient used for separating ubiquitin peptides during Ub-AQUA-PRM analysis.

459 **Supplementary Table S1.** Peptides used for absolute quantification of ubiquitin chain types by Ub-
460 AQUA-PRM analysis.

Sequence	m/z	Z	RT (min)	NCE (%)	Transitions
GGM(ox^2)QIFVK[$^{13}C_6^{15}N_2$]	460.2435	2+	6.7	20	y3$^+$, y5$^+$, y6$^+$, y4$^+$, y2$^+$
GGM(ox^2)QIFVK	456.2364	2+	6.7	20	y3$^+$, y5$^+$, y6$^+$, y4$^+$, y2$^+$
M(ox^2)QIFVK(**GG**)TLTGK[$^{13}C_6^{15}N_2$]	710.3914	2+	7.2	24	y8$^+$, y6$^+$, y9$^+$, y7$^+$, y5$^+$
M(ox^2)QIFVK(**GG**)TLTGK	706.3843	2+	7.2	24	y8$^+$, y9$^+$, y6$^+$, y7$^+$, y5$^+$
TLTGK(**GG**)TITLEVEPSDTIENVK[$^{13}C_6^{15}N_2$]	804.0983	3+	9.3	20	y9$^+$, y9^{2+}, y10$^+$, y19^{2+}, y11$^+$
TLTGK(**GG**)TITLEVEPSDTIENVK	801.4269	3+	9.3	20	y9$^+$, y9^{2+}, y10$^+$, y19^{2+}, y11$^+$
TITLEVEPSDTIENVK(**GG**)AK[$^{13}C_6^{15}N_2$]	1055.061	2+	9	22	y11$^+$, y12$^+$, y13$^+$, y3$^+$, y14$^+$
TITLEVEPSDTIENVK(**GG**)AK	1051.054	2+	9	22	y11$^+$, y12$^+$, y13$^+$, y3$^+$, y14$^+$
AK(**GG**)IQDK[$^{13}C_6^{15}N_2$]	412.7394	2+	5.7	26	y4$^+$, y2$^+$, y3$^+$, y5$^+$, y5^{2+}
AK(**GG**)IQDK	408.7323	2+	5.7	26	y4$^+$, y2$^+$, y3$^+$, y5$^+$, y5^{2+}
IQDK(**GG**)EGIPP[$^{13}C_5^{15}N$]DQQR[$^{13}C_6^{15}N_4$]	551.9536	3+	6.3	22	y6^{2+}, y6$^+$, y5$^+$, y5^{2+}, y3$^+$
IQDK(**GG**)EGIPPDQQR	546.6129	3+	6.3	22	y6^{2+}, y6$^+$, y5$^+$, y5^{2+}, y3$^+$
LIFAGK(**GG**)QLEDGR[$^{13}C_6^{15}N_4$]	735.9006	2+	7.3	24	y8$^+$, y10$^+$, y10^{2+}, y9$^+$, y2$^+$
LIFAGK(**GG**)QLEDGR	730.8964	2+	7.3	24	y8$^+$, y10$^+$, y10^{2+}, y9$^+$, y2$^+$
TLSDYNIQK(**GG**)ESTLHLVLR[$^{13}C_6^{15}N_4$]	752.0737	3+	9.2	20	y16^{2+}, y13^{2+}, y15^{2+}, y14^{2+}, y11^{2+}
TLSDYNIQK(**GG**)ESTLHLVLR	748.7376	3+	9.2	20	y16^{2+}, y13^{2+}, y15^{2+}, y14^{2+}, y11^{2+}
ESTESTLHLVLR[$^{13}C_6^{15}N_4$]	539.3182	2+	7.2	28	y5$^+$, y4$^+$, y3$^+$, y5^{2+}, y6$^+$
ESTESTLHLVLR	534.314	2+	7.2	28	y5$^+$, y4$^+$, y3$^+$, y5^{2+}, y6$^+$
TITTITLEVEPSDTIENVK[$^{13}C_6^{15}N_2$]	898.4744	2+	9.5	20	y9$^+$, y10$^+$, y11$^+$, y12$^+$, y14$^+$

TITTITLEVEPSDTIENVK	894.4673	2+	9.5	20	y9$^+$, y10$^+$, y11$^+$, y12$^+$, y14$^+$

461   *Ox^2 = sulfone version of methionine

464   **Supplementary Table S2.** Multi-step gradient used for separating ubiquitin peptides during Ub-

465   AQUA-PRM analysis.

Time (min)	%B
0	0
4	0
4.25	7
5	12
6	15
7	16
8	24
8.25	35
8.5	90
8.75	90
9	0
10	0