1 ***Trpm8* modulates thermogenesis in a sex-dependent manner without influencing cold-**
2 **induced bone loss**
3
4
5

6 Adriana Lelis Carvalho, PhD[1], Annika Treyball, BA[1], Daniel J. Brooks, MS[2], Samantha Costa,
7 BS[1], Ryan J. Neilson, BS[1], Michaela R. Reagan, PhD[1,3], Mary L. Bouxsein, PhD[2], Katherine J.
8 Motyl, PhD[1,3]*
9
10
11 [1]Center for Molecular Medicine, Maine Medical Center Research Institute, Scarborough, ME
12 [2]Center for Advanced Orthopaedic Studies, Beth Israel Deaconess Medical Center, Boston, MA
13 [3]Tufts University School of Medicine, Tufts University, Boston, MA

14
15
16
17

Grant Funding:
K01AR067858 to Katherine Motyl, PhD, Principal Investigator
P30GM106391 to Robert Friesel, PhD, Program Director
P30GM103392 to Don St. Germain, MD, Program Director
P20GM121301 to Lucy Liaw, PhD, Program Director
U54GM115516 to Clifford Rosen, MD, Program Director

24
25
26

Address correspondence to:
Katherine J. Motyl
81 Research Drive
Scarborough, ME 04074
(207) 396-8004
motylk@mmc.org

33
34
35 Figures: 7
36 Tables: 3
37 Supplementary Figures: 1
38 Supplementary Tables: 1
39
40
41
42
43 Running Head: Sex and *Trpm8*-mediated effects of cold on fat and bone
44
45
46

Abstract

Trpm8 (transient receptor potential cation channel, subfamily M, member 8) is expressed by sensory neurons, and it is involved in the detection of cold temperatures. Increased TRPM8 activity triggers an increase in uncoupling protein 1 (*Ucp1*)-dependent brown adipose tissue (BAT) thermogenesis. Bone density and marrow adipose tissue are regulated by rodent housing temperature and brown adipose tissue, suggesting one mediator of this effect may be TRPM8. In order to test whether TRPM8 was involved in the co-regulation of thermogenesis and bone homeostasis, we examined the bone phenotypes of one-year-old *Trpm8* knockout mice (*Trpm8-KO*) and then exposed them to a 4-week cold temperature challenge. Male *Trpm8-KO* mice had lower bone mineral density than WT, with smaller bone size (femur length and cross-sectional area) being the most striking finding, and exhibited a delayed cold acclimation with increased BAT expression of *Ucp1, Dio2* and *Cidea* at the end of the study. In contrast to males, female *Trpm8-KO* mice had low vertebral bone microarchitectural parameters, but no genotype-specific alterations in body temperature. Interestingly, *Trpm8* was not required for cold-induced trabecular bone loss in either sex, but may be required for cold-induced changes in marrow adiposity in males. Furthermore, bone marrow adipose tissue accumulation in females was significantly blunted by *Trpm8* deletion. In summary, we identified sex differences in the role of TRPM8 in maintaining body temperature, bone density and marrow adipose tissue. Identifying mechanisms through which cold temperature and brown adipose tissue influence bone could help to ameliorate potential bone side effects of obesity treatments designed to stimulate thermogenesis.

Introduction

Maintaining body temperature homeostasis involves both the regulation of heat loss and generation of heat (i.e. thermogenesis). Mammals require energy to generate and/or conserve heat when in temperatures below their thermoneutral zone, which is room temperature in humans, and warmer (28-32°C) in mice [1]. During adaptation to cooler temperatures, sympathetic nervous system (SNS) output increases, elevating energy expenditure, fat oxidation, and heat production by brown adipose tissue (BAT). Uncoupling protein 1 (*Ucp1*) in the inner mitochondrial membrane of BAT generates heat by uncoupling oxidative phosphorylation from ATP production [2,3]. In addition to this, cold exposure can also trigger white adipose tissue (WAT) beiging to enhance heat production [4]. Previous studies have demonstrated a relationship between SNS activity and the skeleton, revealing that sympathetic outflow is not only involved in the activation of BAT thermogenesis for thermoregulation purposes, but that sympathetic tone can also suppress bone formation by signaling through the β2-adrenergic receptor (β2AR) expressed by osteoblasts [11][12]. Marrow adipocytes and osteoblasts have a common mesenchymal precursor and are known to be elevated in many conditions of low bone density. However, the SNS suppresses both bone formation and bone marrow adipose tissue (BMAT), and a four-week cold exposure in on-year-old C3H/HeJ mice leads to reduced BMAT at the most active sites of bone remodeling [13]. In that study, however, bone parameters were largely unchanged, and there was no indication of whether C57BL/6J mice (a commonly used model for skeletal research) would have bone loss from cold exposure. A more recent study, however, showed that thermoneutral (32°C) housing in C57BL/6J mice exerted a protective effect on the rodent skeleton, and this was associated with reduced expression of the *Ucp1* in BAT [1]. These findings suggest that long-term cold-stimulus may be detrimental to bone

density, but it is unclear what particular thermoregulatory pathways may promote bone and

marrow adipose tissue loss.

Transient receptor potential melastatin 8 (TRPM8) is a Ca^{2+} permeable, non-selective

cation channel expressed by temperature-sensing sensory neurons and is known to be the

principal detector of innocuous cold [14]. In mice, TRPM8 is activated by a wide temperature

range, approximately between 28°C to 8°C, and also by cooling agents (e.g. menthol and icilin)

[15]. Previously, TRPM8 was primarily described to work as a peripheral thermal sensor;

however, there is evidence that this cold-sensing channel is also involved in thermoregulation by

promoting heat conservation and energy homeostasis [16]. Furthermore, TRPM8 is a potential

target for obesity treatment due to its function in regulating energy metabolism [16–21]. TRPM8

has been found to trigger non-shivering thermogenesis by BAT and the beiging of WAT in mice

[21][22] and humans [19]. Moreover, *Trpm8* knockout (*Trpm8*-KO) mice are hyperphagic and

have reduced fat oxidation when housed at 21°C, promoting the development of obesity [16].

However, it is unknown whether TRPM8 has a role in regulating bone density *in vivo* and by

inference, whether modulation of TRPM8 by obesity therapeutics might influence bone.

Thus, we conducted this study to determine whether the cold-sensing TRPM8 channel

would be required for any cold-induced changes in the skeleton and bone marrow adipose tissue.

We hypothesized that *Trpm8* deletion would impair body temperature regulation but may protect

mice from cold-induced bone loss and marrow adipose tissue changes. We exposed one-year-old

male and female WT and *Trpm8* knockout (KO) mice to a cold housing (4°C) treatment and

examined body temperature, peripheral and marrow adipose tissue, bone microarchitecture and

remodeling. We found that male *Tprm8-KO* mice had impaired temperature regulation while

females did not. Male *Trpm8-KO* mice had a more overt low bone density phenotype, but

females also had reduced microarchitectural parameters in cortical bone of the femur and

trabecular bone of the vertebrae. Overall, cold temperature reduced trabecular bone volume

fraction in males, while reducing cortical area fraction in females, but these effects were not

dependent upon genotype. Surprisingly, bone marrow adiposity was lower in *Trpm8-KO* females

compared to WT, but there was no striking effect of cold on either genotype or sex of mice, with

the exception of an interaction suggesting TRPM8 may be require for BMAT loss after cold in

males. There were, however, sex and genotype specific effects on the utilization of peripheral

adipose tissue stores, suggesting TRPM8 regulates the energy demands of thermogenesis in a

sex-dependent manner.

Materials and Methods

Wild-type (WT) and *Trpm8-knockout* (*Trpm8-KO*) mice were purchased from The

Jackson Laboratory (stock #008198). Mice were bred and housed in a barrier animal facility at

Maine Medical Center Research Institute (MMCRI) on a 14-hr light and 10-hr dark cycle. Mice

were aged to 52 weeks while being housed up to four animals per cage at 22°C. Mice were given

water and regular chow *ad libitum*. All animal protocols in this study were approved by the

Institutional Animal Care and Use Committee (IACUC) of MMCRI, an Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC)-accredited facility.

Euthanasia was performed using isoflurane followed by cervical dislocation or decapitation.

Dual-energy X-ray absorptiometry (DXA)

All mice had body weight measured prior to DXA. Areal bone mineral content (aBMC),

bone mineral density (aBMD), lean mass and fat mass measurements were performed on each

mouse by a PIXImus dual energy X-ray densitometer (GE Lunar, GE Healthcare). The PIXImus

was calibrated daily with a mouse phantom provided by the manufacturer. Mice were ventral

side down with each limb and tail positioned away from the body. Full-body scans were

obtained, and the head was excluded from analyses because of concentrated mineral content in

skull and teeth. DXA data were processed and analyzed with Lunar PIXImus 2 (version 2.1)

software [23]. Adiposity index was defined as fat mass divided by fat-free mass.

Cold exposure Study

Fifty-two week-old WT and *Trpm8-KO* mice were separated into individual cages. Half

of the mice were maintained at room temperature, while the other half were moved to a cold

room with automatic lights set to a 14 hr light, 10 hr dark cycle, equivalent to the light schedule

of room temperature mice. Light intensity was not measured, but was comparable to that in the

barrier animal facility and the source of the light was at the ceiling. Cold-treated mice were

acclimatized to 18°C for 1 week, then 4°C for 3 weeks. This method of cold-temperature

exposure has been previously published to produce changes in marrow adipose tissue C3H/HeJ

mice and in C57BL/6J mice as well as used to determine dependence of cold-acclimation on

energy expenditure in mice [13][24]. Rectal temperature was measured daily in cold room mice

and every 2 hours on the day of change to 4°C with a Type T thermocouple rectal probe (RET-3,

Physitemp Instruments, Inc., Clifton, NJ, USA) with a MicroTherma 2T hand-held thermometer

(ThermoWorks, Inc., Lindon, UT; cat. no. THS-227-193). Cages contained standard bedding.

Mice were fed regular chow and water *ad libitum.* Environmental temperature and humidity was

measured and recorded daily. Mice were monitored for signs of pain and distress twice per day

and every 2 hours on the day of change to 4°C. We monitored and recorded any of the following

signs: shivering, hunched posture, reduced mobility, ruffled fur and vocalization. Signs that were

found included occasional shivering and ruffled fur, and these generally resolved immediately if

not associated with low core temperature. If core temperature fell below 30°C, mice were

immediately removed to room temperature and euthanized. No mice exhibited temperatures

below this threshold or exhibited signs of distress during the transition from 18°C to 4°C. During

the 4°C period, two mice (one male +/+ and one male -/-) exhibited a low body temperature and

were immediately euthanized. No mice died spontaneously.

Histology

Femur and all adipose tissue samples were placed in 10% neutral buffered formalin

(Sigma-Aldrich) for 48 hr and then stored in 70% ethanol. Bones were decalcified with EDTA

and bone and adipose tissue samples were paraffin embedded and tissues sections of 5 µm were

deparaffinized and dehydrated in a graded series (100-70%) of ethanol washes and stained with

hematoxylin and eosin (H&E). Photomicrographs of H&E stained tissues were taken at 4x

magnification.

Bone Marrow Adipocyte Quantification and Analysis

A blinded analysis of bone marrow (BM) adipocytes was performed on each histology

slide using ImageJ software (https://imagej.nih.gov/ij/) [25,26]. A global scale was set at 1000

pixels/mm based on the microscope imaging objective and settings. The tissue area (T.Ar) was

selected in the distal femoral metaphysis (0.1 mm below the epiphyseal growth plate) using the

polygon tool to contain the BM, excluding cortical bone. One histological slide per distal femur

was measured. The T.Ar was measured using ImageJ to find the total tissue area, which was used

to normalize adipocyte numbers per T.Ar for each image. Color images were converted to 32-bit

and then thresholded using the *Huang* filter at the automatic level determined by ImageJ [25,26].

[Note: thresholding an image converts it to black (255) and white (0) values [25,26]. Images

were then processed with the *Despeckle* and *Adjustable Watershed* plugins. [*Despeckle* was used

to automatically remove salt-and-pepper background noise that could cause inaccuracies with

particle analysis, while the *Adjustable Watershed* ImageJ segmentation algorithm was used to

systematically correct any adjacent adipocytes that had been merged into 1, back into 2 distinct

objects. ImageJ typically applies a watershed segmentation threshold at a level of 0.5, but with

the *Adjustable Watershed* the amount of segmentation can be changed to correct for closely

packed adipocyte, tissue tears, or staining inconsistencies [25,26]. A blinded investigator set the

Adjustable Watershed within the range of 0.3-0.8 to select the most appropriate segmentation

threshold to apply to each image based solely on the quality of the individual image. Next,

particle analysis for BM adipocytes was performed within the T.Ar using the *Analyze Particles*

plugin. This plugin scans a selected area and measures objects that meet parameters based on

area and circularity [25,26]. The particle circumference range was set to 50-2500 pixels (area

results displayed in mm^2 due to the set global scale) and the circularity was set at 0.60-1.00 (1.00

being a perfect circle). These parameters were determined through an iterative, trial and error

analysis of different images until a universal measurement system was decided upon and then

used for every image. After particle analysis, the *Result* and *Summary* pop-up windows provided

the individual and average area (mm^2) for each counted adipocyte within the T.Ar; these data

were used to quantify adiposity in each sample (based on adipocyte number and average

adipocyte area).

Micro-computed tomography (μCT)

A high-resolution desktop micro-tomographic imaging system (μCT40, Scanco Medical

AG, Brüttisellen, Switzerland) was used to assess trabecular bone architecture in the distal

femoral metaphysis and L5 vertebral body and cortical bone morphology of the femoral mid-

diaphysis. Scans were acquired using a 10 μm^3 isotropic voxel size, 70 kVP, 114 μA, 200 ms

integration time, and were subjected to Gaussian filtration and segmentation. Image acquisition

and analysis protocols adhered to guidelines for the assessment of rodent bones by μCT [27]. In

the femur, trabecular bone microarchitecture was evaluated in a 1500 μm (150 transverse slices)

region beginning 200 μm superior to the peak of the growth plate and extending proximally. In

the L5 vertebral body, trabecular bone was evaluated in a region beginning 100 μm inferior to

the cranial end-plate and extending to 100 μm superior to the caudal end-plate. The trabecular

bone regions were identified by manually contouring the endocortical region of the bone.

Thresholds of 335 mgHA/cm^3 and 385 mgHA/cm^3 were used to segment bone from soft tissue in

the femur and L5 vertebrae, respectively. The following architectural parameters were measured

using the Scanco Trabecular Bone Morphometry evaluation script: trabecular bone volume

fraction (Tb.BV/TV, %), t rabecular bone mineral density (Tb. BMD, mgHA/cm^3), specific bone

surface (BS/BV, mm^2/mm^3), trabecular thickness (Tb.Th, mm), trabecular number (Tb.N, mm^{-1}),

and trabecular separation (Tb.Sp, mm), connectivity density (Conn.D, 1/mm^3). Cortical bone was

assessed in 50 transverse μCT slices (500 μm long region) at the femoral mid-diaphysis and the

region included the entire outer most edge of the cortex. Cortical bone was segmented using a

fixed threshold of 708 mgHA/cm^3. The following variables were computed: total cross-sectional

area (bone + medullary area) (Tt.Ar, mm^2), cortical bone area (Ct.Ar, mm^2), medullary area

(Ma.Ar, mm^2), bone area fraction (Ct.Ar/Tt.Ar, %), cortical tissue mineral density (Ct.TMD,

mgHA/cm^3), cortical thickness (Ct.Th, mm), cortical porosity (%), as well as bone strength

indicators such as maximum, minimum and polar moments of inertia (I_{max}, I_{min}, and pMOI,

mm^4), which describe the shape/distribution of cortical bone (larger values indicate a higher

bending strength). Cortical bone images were taken of the mouse using the median total area

value within each group. The scale for all of the images is 1 mm equals 315 pixels.

Real-time PCR

Adipose tissues (brown and gonadal adipose tissue) were collected from WT and *Trpm8-*

KO mice for RNA extraction under liquid nitrogen conditions. Total RNA was prepared using

the standard TRIzol (Sigma-Aldrich) method for tissues. cDNA was generated using the High

Capacity cDNA Reverse Transcriptase Kit (Applied Biosystems) according to the

manufacturer's instructions. mRNA expression analyses was carried out using an iQ SYBR

Green Supermix with a BioRad® Laboratories CFX Connect Real-time System thermal cycler

and detection system. TATA binding protein 1 (Tbp1) was used as an internal standard control

gene for all quantification [28]. Primers used were from Integrated DNA Tecnologies (IDT)

(Coralvile, IA) [29–31] and Primer Design (Southampton, UK). All primer sequences are listed

in Supplementary Table I.

Supplemental Table I. qPCR primer information.

Gene	Source	Sequence	Reference
Acadl	Primer Design	For: 5'-GGA ATG AAA GCT CAG GAC ACA -3' Rev: 5'-AGC CTT TAT TCT CTT CTC CAA GT -3'	N/A
Cidea	IDT	For: 5'-TGC TCT TCT GTA TCG CCC AGT -3' Rev: 5'-GCC GTG TTA AGG AAT CTG CTG -3'	Seale et al. (2008)
Dio2	IDT	For: 5'-CAG TGT GGT GCA CGT CTC CAA TC -3' Rev: 5'-TGA ACC AAA GTT GAC CAC CAG -3'	Cooper et al. (2008)

Pdk4	Primer Design	For: 5'-AAA GTG GGT CTG TGG CAT TG -3' Rev: 5'-AAG GTT TGT ACT CGT GTT TGT G -3'	N/A
Pparc1α	IDT	For: 5'- TGA TGT GAA TGA CTT GGA TAC AGA CA -3' Rev: 5'- GCT CAT TGT TGT ACT GGT TGG ATA TG -3'	Cooper et al. (2008)
Pparγ2	IDT	For: 5'-GCA TGG TGC CTT CGC TGA -3' Rev: 5'-TGG CAT CTC TGT GTC AAC CAT G -3'	Cooper et al. (2008)
Prdm16	IDT	For: 5'-CAG CAC GGT GAA GCC ATT C -3' Rev: 5'-GCG TGC ATC CGC TTG TG -3'	Cooper et al. (2008)
Tbp1	IDT	For: 5'-GAA GCT GCG GTA CAA TTC CAG -3' Rev: 5'-CCC CTT GTA CCC TTC ACC AAT -3'	Sanchez-Gurmaches and Guertin (2014)
Ucp1	IDT	For: 5'-ACT GCC ACA CCT CCA GTC ATT-3' Rev: 5'- CTT TGC CTC ACT CAG GAT TGG -3'	Cooper et al. (2008)

249   N/A: not applicable.

Statistical analysis

GraphPad Prism 8® software was used to visualize data and perform statistical analyses. Data are presented as mean ± standard error. Student's t test or two-way ANOVA was performed with Holm-Sidak *post hoc* multiple comparison test as needed. *p*<0.05 was considered statistically significant.

Results

Body composition and two-dimensional areal bone density (n=14-19/group) were assessed at 52 weeks of age in mice housed at standard room temperature (22°C) (Figure 1). Both male and female *Trpm8-KO* mice had reduced body mass compared to WT mice, which was in part due to a significant reduction in lean mass in males (Figure 1A,B). In both sexes, two-way ANOVA indicated that *Trpm8* deletion significantly affected fat mass and % body fat, but in pairwise comparisons, only female % body fat was lower after *Trpm8* deletion (Figure 1C,D). Significant sex and genotype effects were found in bone mineral density (BMD) and bone mineral content (BMC) parameters (Figure 1E,F). In the latter, there was a significant interaction between sex and genotype such that *Trpm8* deletion caused significantly lower bone mineral content in males, but not in females (Figure 1F).

Figure 1. *Trpm8* deletion causes reduced body mass, lean mass and bone density in male mice. (A-D) Body composition in 52-week old male (blue) and female (purple), wildype (closed circles) and *Trpm8-KO* (open squares) mice was analyzed by DXA. (E,F) Areal total body bone mineral density (aBMD) and areal bone mineral content (BMC) were measured excluding the head. Data are expressed as individual points with bars representing mean ± SEM. Tables below each panel indicate results of the two-way ANOVA, with *p*-values for sex, genotype and their interaction. *p*-values above the bars represent results from Holm-Sidak *post hoc* test for pairwise significance.

Because significant changes in two-dimensional DXA data (aBMD and aBMC) are most

likely to indicate alterations in cortical microarchitecture, we next examined whether *Trpm8*

deletion led to changes in cortical bone by micro Computed Tomography (µCT) (Figure 2). Male

Trpm8-KO mice had significantly lower femur total cross-sectional area (total area) and lower

marrow area compared to male WT mice (Figure 2A-E). These differences were not significant

in females. Polar moment of inertia (an indicator of strength), was dependent upon genotype

(p=0.0479) but within sex pairwise comparisons did not reach statistical significance (Figure

2G). Other parameters, such as cortical tissue mineral density, cortical thickness and porosity

were not dependent upon genotype (Figure 2F,H-J). However, femur length was shorter in male

Trpm8-KO mice compared to WT (Figure 2K). Finally, to test whether changes in bone density

were a result of altered trabecular microarchitecture, we performed µCT on the distal femur

trabecular bone (Figure 3). Deletion of *Trpm8* did not influence volumetric trabecular

architecture in male or female mice (Figure 3). Taken together, these results indicate smaller

bone size in the male *Trpm8-KO* mice may have led to the reduced total body bone mineral

content and density.

Figure 2. *Trpm8* deletion reduced femur size in male mice. Femurs from room temperature
housed male and female *Trpm8-KO* and WT mice were scanned by µCT. (A,B) Representative
images of the femur cross section from the midshaft, where cortical microarchitectural
measurements were performed (C-J). Shaded areas of the pictograms in C-D represent the region
of interest for each measurement. Data are expressed as individual points with bars representing
mean ± SEM. Tables below each panel indicate results of the two-way ANOVA, with *p*-values
for sex, genotype and their interaction. *p*-values above the bars represent results from Holm-
Sidak *post hoc* test for pairwise significance.
300

Figure 3. *Trpm8* deletion did not overtly impact trabecular microarchitecture of the distal
femur. Bone volume fraction (BV/TV) of the trabecular bone of the distal femur was measured
in male and female *Trpm8-KO* and WT mice. Data are expressed as individual points with bars
representing mean ± SEM. Table below the panel indicate results of the two-way ANOVA, with
p-values for sex, genotype and their interaction. *p*-values above the bars represent results from
Holm-Sidak *post hoc* test for pairwise significance.

Previous studies have demonstrated that TRPM8 regulates brown adipose tissue function,

and thermogenesis has been associated with changes in trabecular bone remodeling [12,22,23].

We next wanted to test, 1) whether a 4-week cold exposure would lead to trabecular bone loss

and 2) whether TRPM8 was necessary for the adipose tissue and bone responses to cold (Figure

4A). Mice were separated into individual cages and then left at room temperature or moved to

18ºC (Figure 4A). After one week, the temperature was changed from 18ºC to 4ºC. Male *Trpm8-*

KO mice had a poorer response to the cold challenge than WT, with core temperature being

lower in the KO (Figure 4B). Interestingly, the impact of housing temperature change on female

mouse core temperature was not dependent upon genotype. By the end of week 2, male *Trpm8-*

KO mice had acclimated to the cold exposure and reached core temperatures similar to those of

male WT mice (Figure 4C) which was maintained until the endpoint of the study.

319
Figure 4. Male *Trpm8-KO* mice had lower core temperature during acclimatization to cold.
(A) At 52 weeks of age, mice were separated into individual cages and half were kept at room
temperature while the other half were moved to a cold room set at 18ºC. After one week, the
temperature was changed to 4ºC and mice were maintained for 3 additional weeks. (B) During
the temperature change from 18ºC to 4ºC, core temperature was measured every 2 hours. (C)
Weekly core temperature, starting from time zero before mice were moved into the cold, and
including the data from the 8 hour time point of the day the temperature changed from 18ºC to
4ºC. (D) Percent change in body weight. Temperature data points represent mean ± SEM. Body
weight data are expressed as individual points with bars representing mean ± SEM. Table below
panels B and D indicate results of the two-way ANOVA (performed using data from 8 hr time
point in B), with *p*-values for sex, genotype and their interaction.
331

Previous studies had shown that impaired BAT function in response to cold was

associated with bone loss. Therefore, we next wanted to identify changes in brown adipose tissue

with cold treatment. Cold exposure induced BAT expansion in male WT and *Trpm8-KO* mice

(p<0.05) (Figure 5A). In contrast, the expansion of BAT mass in female mice was observed only

in the WT group (p=0.002) in response to cold, but not in female *Trpm8-KO* mice. In H&E

stained sections of BAT, we observed larger lipid droplets in BAT from males of all genotypes

and treatments, as well as in female *Trpm8-KO* mice at room temperature. Other female groups

had less lipid accumulation, including in the female *Trpm8-KO* mice after cold. We also

observed that cold treatment increased *Ucp1* gene expression in BAT from male WT mice, but

the level of *Ucp1* expression was not different between cold treated WT and *Trpm8-KO* at either

temperature. Overall, KO cold-treated mice had elevated expression of other fatty acid

metabolism markers (*Acadl, Prdm16,* and *Pparc1α*) and thermogenic markers (*Cidea* and *Dio2*)

(Figure 5C). Changes in BAT gene expression were less striking in females compared to males,

with the notable exceptions of cold treatment inducing *Ucp1* to a greater degree in the KO

females than in the WT females and suppressing *PParg2* in both genotypes (Figure 5C).

Together, these results indicate *Trpm8* regulates the BAT response to temperature change in

males and females.

Figure 5. TRPM8 regulates the response of BAT to cold temperature. (A) BAT weights from
male and female WT and *Trpm8-KO* housed at 22°C and 4°C. (B) Representative H&E stained
image of BAT. N=5-8. (C) Gene expression of fatty acid metabolism and thermogenic markers
in male (top) and female (bottom) BAT. Data presented as mean ± standard error. *p<0.05 and
**p<0.01 by Holm-Sidak *post doc* test after a significant two-way ANOVA.
355

Although we did not find any significant differences in inguinal WAT weight (Figure

S1), we did identify that gonadal adipose tissue (gWAT) mass was significantly lowered by cold

treatment in male WT mice and in female *Trpm8-KO* mice (Figure 6A). H&E staining was

suggestive of the presence of beige adipocytes after cold treatment in all sexes and genotypes

except the *Trpm8-KO* cold treated males. The development of beige adipocytes was most

striking in female *Trpm8-KO* mice, however, indicating a clear sex-difference in the way

TRPM8 regulates the acclimatization to cold. These findings were supported by gene expression

in gWAT in female mice, which showed, among other things, a 5 to 8-fold increase in *Ucp1*

expression after cold, while the male mice only had a 2- and 4-fold increase in WT and KO,

respectively, after cold treatment (Figure 6C). Similarly, female mice also had significant

increases in *Acadl, Pparc1a, Pdk4, Cidea* and *Dio2* in the cold, whereas males did not have any

other changes (Figure 6C). Taken together, these data indicate TRPM8 may be required for

beiging of gWAT tissue in males but not females. Further, females may be more prone to

generate beige adipocytes in gWAT than males, which could explain their advantage over males

in adjusting to cold temperature.

Figure 6. Loss of *Trpm8* prevented beige adipocyte formation in gonadal adipose tissue
from male but not female mice. (A) gWAT weights from male and female WT and *Trpm8-KO*
housed at 22°C and 4°C. (B) Representative H&E stained image of gWAT. (C) Gene expression
of fatty acid metabolism and thermogenic markers in male (top) and female (bottom) gWAT.
Data presented as mean ± standard error. *p<0.05 and **p<0.01 by Holm-Sidak *post doc* test
after a significant two-way ANOVA.
378

Because altered thermogenesis has previously been associated with changes in bone

homeostasis, we next examined bone microarchitectural phenotypes of WT and *Trpm8-KO* mice

at room temperature and with cold exposure. In addition to cortical bone changes due to

genotype and sex noted above in mice at room temperature (Figure 2), we determined that cold

temperature caused trabecular bone loss (decreased BV/TV) in male mice of both gentoypes, in

both the femur and the vertebrae (Table I). However, cold temperature treatment did not

influence cortical bone in either genotype of male mice. In contrast to males, females had only

had increased L5 trabecular BS/BV (indicative of trabecular thinning) in response to cold, but no

changes in BV/TV in either the femur or the vertebral trabecular bone (Table II). Medullary area

was increased with cold, leading to a reduction in cortical area fraction in females (Table II).

However, there were no significant temperature by genotype interactions in trabecular or cortical

bone parameters in either sex, suggesting bone loss from cold was not dependent upon TRPM8

in males or females (Tables I and II). Of note, female, but not male, mice also had reduced

vertebral trabecular microarchitectural parameters after *Trpm8* deletion (Table II), but this was

also not dependent upon housing temperature.

Table I. Femoral and Vertebral Bone outcomes assessed by μCT of Male wildtype (WT) and *Trpm8-Knockout* (*KO*) mice housed at room temperature or cold.

	Room temperature (22°C)		Cold (4°C)		*p*-value		
	WT	KO	WT	KO	genotype	temp	interaction
Distal Femur Trabecular Bone							
BV/TV (%)	11.1 ± 0.8	11.9 ± 0.9	8.9 ± 1.0	9.9 ± 1.1	0.36	**0.04**	0.91
Conn.D (1/mm^3)	43 ± 7	58.5 ± 9.0	31.2 ± 5.8	43.1 ± 7.3	0.07	0.08	0.79
Tb.N (1/mm)	3.0 ± 0.1	3.2 ± 0.1	2.8 ± 0.1	3.0 ± 0.1	0.07	0.06	0.67
Tb.Th (mm)	0.059 ± 0.001	0.061 ± 0.003	0.058 ± 0.002	0.056 ± 0.002	0.95	0.15	0.19
Tb.Sp (mm)	0.323 ± 0.017	0.304 ± 0.007	0.347 ± 0.007	0.321 ± 0.011	0.07	0.09	0.78
Femur Midshaft Cortical Bone							
Ct.Th (mm)	0.159 ± 0.007	0.161 ± 0.008	0.160 ± 0.006	0.154 ± 0.004	0.20	0.67	0.52
Ct.Ar (mm^2)	0.88 ± 0.04	0.86 ± 0.04	0.88 ± 0.03	0.82 ± 0.03	0.26	0.49	0.52
Ct.TMD (mgHA/cm^3)	1195 ± 7	1197 ± 5	1198 ± 8	1192 ± 5	0.73	0.79	0.50
Ma.Ar (mm^2)	1.77 ± 0.10	1.52 ± 0.05	1.74 ± 0.11	1.54 ± 0.06	**0.01**	0.97	0.81
Tt.Ar (mm^2)	2.64 ± 0.09	2.38 ± 0.06	2.62 ± 0.12	2.36 ± 0.07	**0.005**	0.77	0.99
Ct.Ar/Tt.Ar (%)	33.5 ± 1.9	36.2 ± 1.5	33.7 ± 1.4	34.8 ± 0.9	0.21	0.67	0.58
Ct. Porosity (%)	0.23 ± 0.03	0.26 ± 0.03	0.20 ± 0.01	0.23 ± 0.02	0.27	0.14	0.87
pMOI (mm^4)	0.64 ± 0.04	0.58 ± 0.04	0.64 ± 0.04	0.54 ± 0.03	**0.04**	0.63	0.72
Imax (mm^4)	0.42 ± 0.03	0.40 ± 0.03	0.42 ± 0.03	0.37 ± 0.03	0.19	0.53	0.67
Imin (mm^4)	0.22 ± 0.01	$0.18 \pm 0.01*$	0.22 ± 0.02	$0.18 \pm 0.01*$	**0.0009**	0.89	0.89
L5 Vertebral Trabecular Bone							
BV/TV (%)	21.6 ± 1.1	20.7 ± 1.8	18.7 ± 0.5	18.4 ± 0.7	0.59	**0.04**	0.78
BS/BV (mm^2/mm^3)	43.9 ± 1.5	44.5 ± 1.9	46.2 ± 0.4	46.9 ± 0.3	0.62	0.08	0.95
BMD (mgHA/cm^3)	226.4 ± 10.0	217.3 ± 16.1	198.2 ± 4.7	197.6 ± 5.9	0.65	**0.03**	0.69

Conn.D (1/mm^3)	127 ± 11	115 ± 8	113 ± 5	119 ± 8	0.73	0.53	0.32
Tb.N (1/mm)	4.4 ± 0.1	4.3 ± 0.1	4.1 ± 0.1	4.2 ± 0.1	0.94	0.11	0.43
Tb.Th (mm)	0.050 ± 0.001	0.053 ± 0.002	0.050 ± 0.003	0.053 ± 0.002	0.10	0.96	0.96
Tb.Sp (mm)	0.215 ± 0.007	0.223 ± 0.009	0.236 ± 0.005	0.230 ± 0.009	0.89	0.08	0.37

*p<0.05 compared to WT mice group. N=6–7. *Abbreviations:* BV/TV, bone volume fraction; Conn.D, connectivity density; Tb.N, trabecular number; Tb.Th, trabecular thickness; Tb.Sp, trabecular separation; Ct.Th, average cortical thickness; Ct.Ar, cortical bone area; Ct.TMD, cortical tissue mineral density; M.Ar, marrow area; T.Ar, total cross-sectional area; Ct.Ar/T.Ar, cortical area fraction; Ct.Porosity, cortical porosity; pMOI, polar moment of inertia; Imax, maximum moment of inertia; Imin, minimum moment of inertia; BMD, bone mineral density; BS/BV, specific bone surface.

Table II. Femoral and Vertebral Bone outcomes assessed by μCT of Female wildtype (WT) and *Trpm8-Knockout* (*KO*) mice housed at room temperature or cold.

	Room temperature (22°C)		Cold (4°C)		*p*-value		
	WT	KO	WT	KO	genotype	temp	interaction
Distal Femur Trabecular Bone							
BV/TV (%)	2.13 ± 0.38	2.78 ± 0.90	1.87 ± 0.20	2.58 ± 0.78	0.32	0.73	0.97
Conn.D (1/mm^3)	4.3 ± 1.4	14.4 ± 7.9	6.6 ± 1.7	15.9 ± 10.7	0.21	0.81	0.96
Tb.N (1/mm)	1.7 ± 0.1	1.8 ± 0.1	1.7 ± 0.1	1.8 ± 0.1	0.46	0.74	0.62
Tb.Th (mm)	0.060 ± 0.003	0.056 ± 0.003	0.052 ± 0.001	0.054 ± 0.003	0.56	0.07	0.27
Tb.Sp (mm)	0.611 ± 0.039	0.584 ± 0.037	0.588 ± 0.027	0.585 ± 0.32	0.67	0.74	0.72
Femur Midshaft Cortical Bone							
Ct.Th (mm)	0.188 ± 0.007	0.177 ± 0.004	0.177 ± 0.004	0.174 ± 0.004	0.16	0.16	0.42
Ct.Ar (mm^2)	0.85 ± 0.04	0.78 ± 0.02	0.82 ± 0.02	0.78 ± 0.02	**0.04**	0.71	0.62
Ct.TMD (mgHA/cm^3)	1251 ± 6	1244 ± 5	1248 ± 5	1241 ± 4	0.16	0.55	0.93
Ma.Ar (mm^2)	1.12 ± 0.04	1.05 ± 0.03	1.21 ± 0.05	1.14 ± 0.03	0.08	**0.02**	0.89

Tt.Ar (mm^2)	1.97 ± 0.07	1.83 ± 0.04	2.03 ± 0.06	1.92 ± 0.04	**0.02**	0.14	0.81
Ct.Ar/Tt.Ar (%)	43.1 ± 1.2	42.6 ± 1.0	40.7 ± 0.9	40.7 ± 0.8	0.80	**0.03**	0.76
Ct. Porosity (%)	0.19 ± 0.01	0.21 ± 0.02	0.21 ± 0.01	0.24 ± 0.03	0.28	0.26	0.86
pMOI (mm^4)	0.44 ± 0.04	0.37 ± 0.02	0.44 ± 0.02	0.37 ± 0.03	**0.02**	0.91	0.98
Imax (mm^4)	0.29 ± 0.03	0.24 ± 0.01	0.28 ± 0.02	0.25 ± 0.01	**0.02**	0.80	0.78
Imin (mm^4)	0.15 ± 0.01	0.13 ± 0.01	0.16 ± 0.01	0.15 ± 0.01	0.05	0.33	0.65
L5 Vertebral Trabecular Bone							
BV/TV (%)	15.8 ± 1.2	13.1 ± 0.5	14.2 ± 1.2	12.0 ± 0.5	**0.01**	0.16	0.81
BS/BV (mm^2/mm^3)	38.2 ± 1.3	40.9 ± 1.1	41.5 ± 1.2	43.4 ± 1.7	0.10	**0.04**	0.77
BMD (mgHA/cm^3)	174 ± 12	148 ± 4	158 ± 12	138 ± 4	**0.02**	0.15	0.80
Conn.D (1/mm^3)	51 ± 9	48 ± 7	60. ± 7	48 ± 8	0.36	0.55	0.57
Tb.N (1/mm)	2.67 ± 0.07	2.43 ± 0.12	2.68 ± 0.11	2.44 ± 0.09	**0.02**	0.90	0.99
Tb.Th (mm)	0.061 ± 0.002	0.057 ± 0.002	0.056 ± 0.001	0.054 ± 0.002	0.15	0.06	0.64
Tb.Sp (mm)	0.382 ± 0.009	0.428 ± 0.024	0.384 ± 0.014	0.423 ± 0.016	**0.02**	0.90	0.83

*p<0.05 compared to WT mice group. N=8-11. *Abbreviations:* BV/TV, bone volume fraction; Conn.D, connectivity density; Tb.N, trabecular number; Tb.Th, trabecular thickness; Tb.Sp, trabecular separation; Ct.Th, average cortical thickness; Ct.Ar, cortical bone area; Ct.TMD, cortical tissue mineral density; M.Ar, marrow area; T.Ar, total cross-sectional area; Ct.Ar/T.Ar, cortical area fraction; Ct.Porosity, cortical porosity; pMOI, polar moment of inertia; Imax, maximum moment of inertia; Imin, minimum moment of inertia; BMD, bone mineral density; BS/BV, specific bone surface.

Finally, we also evaluated whether cold exposure and/or loss of *Trpm8* would affect the

adiposity of the bone marrow (Figure 7). In general, bone marrow adipose tissue (BMAT) is

limited in male mice compared to females (Figure 7A-G). In male mice, we only found a

significant interaction between genotype and housing temperature (p=0.045) in the % total

adiposity, suggesting *Trpm8-KO* males do not lose marrow adipose tissue at cold temperatures,

while WT mice do (Figure 7D). In female mice, there was a significant effect of *Trpm8* deletion

on adipocyte number (Figure 7E) and % adiposity (Figure 7G), but not adipocyte size. In

contrast to males, there was no clear impact of housing temperature on bone marrow adipose

tissue females.

Figure 7. *Trpm8* deletion suppressed marrow fat in female mice. (A and B) Representative
μCT images of cortical bone from male and female WT and *Trpm8-KO* mice housed at standard
room temperature (RT) (22°C) and cold temperature (4°C). (C and D) Quantification of
adipocyte numbers (Ad.N/T.Ar), average adipocyte size (Ad.A), and % adiposity (T.Ar
Adiposity) within the bone marrow of male and female WT and *Trpm8-KO* mice housed at 22°C
and 4°C. (E and F) Representative H&E stained image of distal femur area from male and female
WT and *Trpm8-KO* mice housed at 22°C and 4°C (scale bar: 500 μm; magnification: 4x). Data
presented as mean ± standard error.
431

Discussion

In this study, we determined that the impact of *Trpm8* deletion on bone, body temperature

and adipose tissue was dependent upon sex and housing temperature (Table IV summary). Our

original hypothesis was that *Trpm8* deletion would impair body temperature regulation, but

would also protect against cold-induced changes in bone and marrow adipose tissue. Males, but

not females, had impaired temperature regulation after *Trpm8* deletion. However, cold housing

led to trabecular bone loss in males and cortical bone loss in females, independent of genotype,

which is consistent with previous studies showing that cold exposure can negatively impact the

skeleton [1,12]. It has also previously been shown that cold exposure is associated with reduced

BMAT [32], and our findings indicate that in males this may be dependent upon *Trpm8* (Figure

7). On the other hand, female BMAT was highly dependent upon *Trpm8* but not dependent upon

housing temperature, pointing toward potentially diverse function of TRPM8 in regulating

BMAT levels and response to stimuli. Despite the lack of any temperature by genotype

interaction effects on bone parameters, it is worth noting that male KO mice, with impaired

temperature regulation, also had smaller bones, while female KO mice had normal temperature

regulation had normal bone size.

Table IV. Summary of results.

	Males		Females	
	Trpm8 **Deletion**	**Cold Housing**	*Trpm8* **Deletion**	**Cold Housing**
Body Temperature	Sex by genotype interaction – temperature reduced in male KOs but not female KOs in cold			
Bone Size	decreased	uc	uc	uc
Cortical Bone	uc	uc	uc	decreased
Trabecular Bone	uc	decreased	decreased (L5 only)	uc
Marrow Adiposity	Genotype by temp interaction		decreased	uc

Abbreviation: uc, unchanged.

A bone phenotype in *Trpm8-KO* mice has not been previously described. Bone size was

smaller in male *Trpm8*-KO mice compared to WT, but this difference did not occur in females.

Furthermore, female mice had significant trabecular bone loss in the vertebrae but not the femur,

indicating potential site-specific effects of *Trpm8* deletion on bone. Recently, Iwaniec et al.

(2016) highlighted that the current standard room temperature housing (22°C) can already be

considered a cold environment to mice and that it was associated with bone loss in growing

females compared to thermoneutral housing [1]. A *Trpm8* functional cold circuit is established

after birth around postnatal day 14 [33]. Others have found that *Trpm8* is a regulator of energy

metabolism and thermogenesis [16,19,21], and our findings are consistent: showing that its

deletion can cause an increase in the expression of thermogenic markers, gWAT fat pad

reduction and weight loss. This suggests that the absence of *Trpm8* during the developmental

phase might be associated with defective cold circuits which partially fail to sustain optimal body

temperature, as previously described [16]. Thus, it is plausible that the lack of *Trpm8* during

growth causes a shift towards energy utilization for heat generation, decreasing ATP availability

for bone cells, leading to altered bone density. However, the mechanism of low bone mass in

Trpm8-KO mice, and the relative contributions of bone forming osteoblasts and bone resorbing

osteoclasts, are yet to be determined. Some *in vitro* evidence suggests human osteoblast-like

cells and murine osteoblast cells express *Trpm8,* but the functional impact on bone is unclear

[34]. Although the deletion of other *Trp* channels, such as *Trpv1* [35] and *Trpv5* [36], has been

reported to induce bone loss in animal models, there is a lack of data exploring the physiological

role of TRPM8 in skeletal homeostasis.

One interesting finding was that *Trpm8* deletion in females caused decreased BMAT

content. Aging generally leads to accumulation of bone marrow adipose tissue in the distal femur

and proximal tibia in mice, but the mechanisms and consequences of this accumulation remain

an area of active investigation. Both osteoblasts and adipocytes arise from a shared mesenchymal

precursor, thus one hypothesis is that situations of bone loss may lead to marrow adiposity by

shunting cells away from the osteoblast lineage to the adipocyte lineage (as in calorie restriction,

PPARg2 inhibition, and aging). However, the alternate, shifting from marrow adipose lineage to

osteoblast lineage, does not seem likely, since female *Trpm8-KO* mice had indices of reduced

bone density in concert with low marrow adiposity. In other situations, like bone loss from the

atypical antipsychotic drug risperidone, elevated SNS activity leads to a reduction in both

osteoblast-mediated bone formation and marrow adiposity as well as increased expression of

Ucp1 in BAT [23]. Recently, Wee et al., (2019) elegantly demonstrated that iWAT and BMAT

shared sympathetic innervation using retroviral nerve tracing [37]. They revealed that leptin-

responsive regions such as paraventricular hypothalamus, arcuate nucleus, dorsomedial

hypothalamus, ventromedial hypothalamus, and area postrema were labelled after viral tracing

from iWAT, bone marrow, and BMAT [37]. However, it remains unclear if sex-differences exist

in neural regulation of these tissues, and if gWAT would be regulated by similar common

sympathetic pathways. Taken together, these findings suggest elevated SNS activity to stimulate

thermogenesis could be responsible for bone loss and reduced marrow adiposity with *Trpm8*

deletion.

Investigations into sex differences in the thermogenic response to *Trpm8* deletion have

been limited. Clearly, *Trpm8* is important for the detection of environmental cooling [14,15,38]

and it has also been implicated in BAT thermogenesis and thermoregulatory processes

[16,21][22]. Early publications on *Trpm8* had reported no differences in core temperature

associated with its global deletion in mice [38]. Ma et al. (2012) demonstrated that menthol

activates Ucp-1 dependent thermogenesis and increases core temperature in WT mice, but that

when TRPM8 is deleted, this activation is absent [22]. In contrast, our studies use cold-

temperature to induce thermogenesis, and show a partial dependence on TRPM8. However, our

findings also indicate that other pathways may compensate for the loss of TRPM8, especially in

the females. Specifically, *Trpm8-KO* male mice struggled to increase body temperature to the

level of WT in a cold environment (of which, menthol treatment is partly mimicking). However,

the study by Ma, et al. (2012) did not directly compare WT to KO body temperature, so it is

difficult to align perfectly with our findings. Recently, Reimúndez et al. (2018) demonstrated

that *Trpm8* is required for thermoregulation in male mice when housed in colder environments,

which corroborates our findings [16]. Interestingly, Reimúndez et al. (2018) also demonstrated

that this difficulty in maintaining core temperature was associated with greater tail vasodilation

and heat loss in male mice with global deletion of *Trpm8*, indicating that this cold-sensing

channel has an important role in regulating both thermogenesis and heat dissipation [16].

However, female core temperature status was not investigated in any of these studies.

In contrast, one strength of our study is the comprehensive analyses of mice of both

sexes. Cooling sensitivity is the first step to activate the cold response neuronal mechanism that

modulates lipolysis and energy expenditure [39]. In our study, we observed that cold-induced

weight loss occurred in both sexes, irrespective of the genotype. However, there were *Trpm8*-

and sex-differences among the various adipose tissues' response to housing temperature. Male

mice housed at 4°C exhibited BAT expansion with fewer lipid droplets and increased *Ucp1*

expression in WT, which is indicative of thermogenesis activity. However, this difference did

appear to be blunted in KO mice (Figure 5). Moraes et al. (2017) found reduced *Ucp1* and *Pparg*

expression at ZT2 and ZT14 in male TRPM8-KO mice ranging from 3-6 months of age,

compared to ours that were 12 months [20]. Although we did not see lower *Ucp1* or *Pparg* in our

room temperature mice, we expect that this difference might be due to the age of the mice or the

time of the day that our tissues were collected ZT6-ZT10. However, our overall finding that

thermogenesis is impaired in males is consistent with Moraes et al. finding of reduced *Ucp1* [20].

Further, we found that cold housing in male KO, but not WT, mice induced BAT expression of

other fatty acid metabolism markers (*Acadl, Prdm16,* and *Pgc1α*) as well as a robust expression

of other thermogenic markers such as *Cidea* and *Dio2* in BAT, which may have been

compensatory for a lack of *Ucp1* induction compared to room temperature.

In females, we found that *Trpm8* deletion led to an increase in *Ucp1* expression and

reduction in lipid droplets in BAT during cold exposure, indicating TRPM8 was not required for

the response. Furthermore, cold reduced gWAT mass in female mice from both genotypes, in

addition to increasing expression of fatty acid metabolism (*Acadl, Pgc1α, Pdk4*) and

thermogenic (*Ucp1, Cidea, Dio2*) markers. It should be considered that estrogen status may play

a role in sex-differences in thermogenesis and WAT beiging found in the present study. Several

studies observed that the beiging process does take place in gWAT in females [40,41]. Increased

expression of beiging markers such as *Ucp1*, *Pgc1α*, *Elovl3*, *Cidea* was observed in gWAT after

either 10 days of cold exposure [41] or β3-adrenergic receptor agonist treatment [40] in female

mice. Kim et al. (2016) demonstrated that gWAT in females is more susceptible to the beiging

process than in males, due to more extensive sympathetic innervation inducing brown

adipogenesis [40]. Also, the authors showed that sympathetic innervation of gWAT is sex-

hormone dependent, as reduced tyrosine hydroxylase protein levels and lack of beiging of gWAT

was found after chemical ovarian ablation [40]. Indeed, lipolysis can be directly modulated by

estrogen through estrogen receptor alpha (ERα) signaling in BAT and WAT adipocytes [42].

Estrogen also has been described to play a critical role in cold sensitivity and hyperalgesia

[43,44], which are also functions thought to be regulated by *Trpm8* [45,46]. However,

interestingly, a previous preclinical study observed that *Trpm8* expression was upregulated in the

skin of ovariectomized rats compared to sham control rats [47] and β-estradiol treatment might

reduce *Trpm8* overexpression in the skin of ovariectomized rats [48]. These findings support our

data that suggest regulation of energy metabolism in female mice is independent of *Trpm8* cold-

sensing function and estrogen might be a more relevant factor to normal thermoregulation in

females. It is unclear however, whether ovariectomized mice, or mice aged to natural

amenorrhea, would have a response to cold that is modified by *Trpm8* deletion.

The global deletion of *Trpm8* may have led to compensatory changes in other

thermogenic mechanisms, which may have been sex-dependent. In addition to TRPM8, there is

another cold-activated *Trp* channel known as transient receptor potential ankyrin 1 (TRPA1)

which is considered to be a noxious cold receptor [49]. Interestingly, TRPM8 and TRPA1 appear

to function in both complementary and synergistic fashion to detect environmental innocuous

and noxious cold [49]. Winter et al. (2017) found that the lack of *Trpm8* and *Trpa1* led to a

striking reduction in cold avoidance behavior in mice [49]. However, an acute cold avoidance

response was still observed in *Trpm8-KO* mice with functional TRPA1 when exposed to a

temperature range between 28°C and 15°C [49]. This suggests that TRPA1 can only partially

compensate for the absence of TRPM8 [49]. When cold-defense mechanisms, such as

thermogenesis and vascular tone, fail to sustain an optimal body temperature, a shivering

response is triggered to increase heat generation [50]. In our study, only occasional shivering-

like behaviors were observed in some mice. These resolved quickly and we did not note any

differences in the genotypes of the mice that shivered. Previous studies found that *Trpm8* is not

required to induce shivering and tachycardia during cold challenge [50]. Instead, shivering is

found to be regulated by TRPV1 [50], which once more indicates TRPM8 is a major regulator of

non-shivering thermogenesis.

In summary, our findings support that *Trpm8* is important for body temperature

maintenance in male mice and its absence triggers a compensatory response to counteract

reduction in core temperature. Moreover, *Trpm8* does not appear to regulate the bone response to

colder environments, where it is presumed that the sympathetic nervous system-mediated stress

response and/or diversion of energy away from bone contributes to bone loss. Interestingly,

deletion of *Trpm8* was associated with bone size and shape differences in male mice as well.

Therefore, although any bone loss associated with acute cold housing may be independent of

TRPM8, low bone mass found in *Trpm8*-KO mice compared to WT, more striking in males,

indicates that the functions of TRPM8 could contribute to bone modeling and remodeling

throughout lifespan.

Limitations

Some limitations of this study do exist. First, addition of a thermoneutral group would potentially

exacerbate differences observed, and could provide additional insight into whether *Trpm8*

regulates bone and marrow adiposity independent of cold temperature sensation and subsequent

physiologic response. We regret that we did not have the capability to age mice to 52 weeks of

age in a thermoneutral (~28-30ºC) rodent incubator when this study was first designed. Next,

because the room temperature mice were housed in a separate room from the cold-treated, we

cannot exclude the possibility that the additional noise of the cold room contributed to the

phenotype. There is no known association between noise stress and changes in brown adipose

tissue, but there are some studies examining the impact of chronic mild stress (CMS), of which

one of the stressors is noise, on bone density. One study found that the CMS caused bone loss,

and this could be blocked by propranolol, which suggests the SNS is involved [54]. However,

other stressors were involved in that study, including water deprivation, continuous overnight

illumination, stroboscopic illumination, cage tilt, and time in a soiled cage. Thus, it is unclear

whether noise alone would have influenced bone microarchitecture. Although the sound was

greater in the cold room in our study, it was consistent over time, and the fact remains that

compared to WT males, and females of both genotypes all treated in the cold, the body

temperature in the male KO mice had a delayed response to cold. Other key findings, like

Trpm8-KO mediated bone loss, were not dependent upon housing conditions. Light intensity was

not measured, but was similar and from the ceiling in both rooms. Furthermore, additional N

would have helped to reduce biological variability, which was high in some of the secondary

outcomes of interest, such as inguinal WAT mass and expression of *Ucp1* in both BAT and

gWAT. Although differences were not detected in iWAT mass, it is possible that histology and

mRNA analyses of this tissue would have provided additional valuable information. However,

iWAT samples were not saved to analyze at a later time point. Additionally, examining the

thermogenic response at the time point where we observed major differences in body

temperature in male mice exposed to cold would have provided better insight into mechanism.

However, analysis of bone defects after cold were the primary outcomes of interest, and these

would not have been apparent at such an early time point.

Despite these limitations, the current data provide new insight into our understanding of

the role of *Trpm8* in the regulation of thermogenesis and skeletal homeostasis and demonstrate

important and unexpected sex-differences in these processes. This work suggests that in males in

particular, TRPM8 is necessary for normal thermoregulation and bone density, while in females,

TRPM8 regulates marrow adiposity.

Acknowledgments

The authors thank Phuong Le and Casey Doucette for the assistance during the cold housing

experiment, and Audrey Bergeron for critical review of the manuscript. This work was supported

by the National Institute of Arthritis And Musculoskeletal And Skin Diseases (NIAMS) and the

National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health

(NIH) under award numbers K01AR067858 and P20GM121301 to KJM. This work utilized

services of the Maine Medical Center Research Institute (MMCRI) Molecular Phenotyping Core,

which is supported by NIH/NIGMS P30GM106391, the Physiology Core, which is supported by

NIH/NIGMS P30GM106391 and P20GM121301, the Histopathology and Histomorphometry

Core, which is supported by NIH/NIGMS P30GM106391, P20GM121301, and P30103392, and

the Mouse Transgenic and In Vivo Imaging Core which is supported by NIH/NIGMS

P30GM103392. All cores also received

support from the Norther New England Clinical and Translational Research Network

NIH/NIGMS U54GM115516. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

Authors Contributions

ALC was responsible for data acquisition, data analysis, interpretation, drafting and critical

review of the manuscript. AT was responsible for data acquisition, data analysis, interpretation,

drafting and critical review of the manuscript. DJB was responsible for data acquisition, data

analysis, interpretation, drafting and critical review of the manuscript. SC was responsible for

data acquisition, data analysis, interpretation, drafting and critical review of the manuscript. RN

was responsible for data acquisition, data analysis and critical review of the manuscript. MRR

was responsible for data interpretation and critical review of the manuscript. MLB was

responsible for data interpretation and critical review of the manuscript. KJM was responsible for

experimental design, data acquisition, data analysis, interpretation, drafting, and critical review

of the manuscript. All authors had final approval of the manuscript.

Declaration of interests

The authors have declared that no conflict of interest exists.

References

1. Iwaniec UT, Philbrick KA, Wong CP, Gordon JL, Kahler-Quesada AM, Olson DA, et al. Room temperature housing results in premature cancellous bone loss in growing female mice: implications for the mouse as a preclinical model for age-related bone loss. Osteoporos Int. 2016;27: 3091–3101. doi:10.1007/s00198-016-3634-3

2. Fedorenko A, Lishko P V., Kirichok Y. Mechanism of fatty-acid-dependent UCP1 uncoupling in brown fat mitochondria. Cell. 2012;151: 400–413. doi:10.1016/j.cell.2012.09.010

3. Tan CL, Knight ZA. Regulation of Body Temperature by the Nervous System. Neuron. 2018;98: 31–48. doi:10.1016/j.neuron.2018.02.022

4. Contreras GA, Lee YH, Mottillo EP, Granneman JG. Inducible brown adipocytes in subcutaneous inguinal white fat: The role of continuous sympathetic stimulation. Am J Physiol - Endocrinol Metab. 2014;307: E793–E799. doi:10.1152/ajpendo.00033.2014

5. Bahler L, Verberne HJ, Admiraal WM, Stok WJ, Soeters MR, Hoekstra JB, et al. Differences in sympathetic nervous stimulation of brown adipose tissue between the young and old, and the lean and obese. J Nucl Med. 2016;57: 372–377. doi:10.2967/jnumed.115.165829

6. Liu CK. the Impact of Sarcopenia on a Physical Activity Intervention □: the Lifestyle Interventions and. J Nutr Heal Aging. 2014;18: 59–64. doi:10.1007/s12603-013-0369-

0.THE

7.   Zoico E, Rubele S, De Caro A, Nori N, Mazzali G, Fantin F, et al. Brown and beige

adipose tissue and aging. Front Endocrinol (Lausanne). 2019;10: 1–10.

doi:10.3389/fendo.2019.00368

8.   Wang W, Ishibashi J, Trefely S, Shao M, Cowan AJ, Sakers A, et al. A PRDM16-Driven

Metabolic Signal from Adipocytes Regulates Precursor Cell Fate. Cell Metab. 2019;30:

174-189.e5. doi:10.1016/j.cmet.2019.05.005

9.   Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated

with osteoporotic fractures. Osteoporos Int. 2006;17: 1726–1733. doi:10.1007/s00198-

006-0172-4

10.   Odén A, McCloskey E V., Kanis JA, Harvey NC, Johansson H. Burden of high fracture

probability worldwide: secular increases 2010–2040. Osteoporos Int. 2015;26: 2243–

2248. doi:10.1007/s00198-015-3154-6

11.   Elefteriou F. Neuronal signaling and the regulation of bone remodeling. Cell Mol Life Sci.

2005;62: 2339–2349. doi:10.1007/s00018-005-5175-3

12.   Motyl KJ, Bishop KA, Demambro VE, Bornstein SA, Le P, Kawai M, et al. Altered

thermogenesis and impaired bone remodeling in Misty mice. J Bone Miner Res. 2013;28:

1885–1897. doi:10.1002/jbmr.1943

13.   Scheller EL, Doucette CR, Learman BS, Cawthorn WP, Khandaker S, Schell B, et al.

Region-specific variation in the properties of skeletal adipocytes reveals regulated and

constitutive marrow adipose tissues. Nat Commun. 2015;6: 7808.

doi:10.1038/ncomms8808

14.   Dhaka A, Murray AN, Mathur J, Earley TJ, Petrus MJ, Patapoutian A. TRPM8 Is

Required for Cold Sensation in Mice. Neuron. 2007;54: 371–378.

doi:10.1016/j.neuron.2007.02.024

15.   McKemy DD, Neuhausser WM, Julius D. Identification of a cold receptor reveals a

general role for TRP channels in thermosensation. Nature. 2002;416: 52–58.

doi:10.1038/nature719

16.   Reimúndez A, Fernández-Peña C, García G, Fernández R, Ordás P, Gallego R, et al.

Deletion of the cold thermoreceptor TRPM8 increases heat loss and food intake leading to

reduced body temperature and obesity in mice. J Neurosci. 2018;38: 3643–3656.

doi:10.1523/JNEUROSCI.3002-17.2018

17.   Zhu Z, Ma S, Yu H, Zhao Z, Luo Z, Chen J, et al. Activation of the cold-sensing TRPM8

channel triggers UCP1-dependent thermogenesis and prevents obesity. J Mol Cell Biol.

2012;4: 88–96. doi:10.1093/jmcb/mjs001

18.   McCoy DD, Zhou L, Nguyen AK, Watts AG, Donovan CM, McKemy DD. Enhanced

insulin clearance in mice lacking TRPM8 channels. Am J Physiol - Endocrinol Metab.

2013;305: 78–88. doi:10.1152/ajpendo.00542.2012

19.   Rossato M, Granzotto M, Macchi V, Porzionato A, Petrelli L, Calcagno A, et al. Human

white adipocytes express the cold receptor TRPM8 which activation induces UCP1

expression, mitochondrial activation and heat production. Mol Cell Endocrinol. 2014;383:

137–146. doi:10.1016/j.mce.2013.12.005

20.   Moraes MN, de Assis LVM, Henriques F dos S, Batista ML, Güler AD, Castrucci AM de

L. Cold-sensing TRPM8 channel participates in circadian control of the brown adipose

tissue. Biochim Biophys Acta - Mol Cell Res. 2017;1864: 2415–2427.

doi:10.1016/j.bbamcr.2017.09.011

21.   Jiang C, Zhai M, Yan D, Li D, Li C, Zhang Y, et al. Dietary menthol-induced TRPM8

714   activation enhances WAT 1C "browning 1D" and ameliorates diet-induced obesity.

715   Oncotarget. 2017;8: 75114–75126. doi:10.18632/oncotarget.20540

716   22.   Ma S, Yu H, Zhao Z, Luo Z, Chen J, Ni Y, et al. Activation of the cold-sensing TRPM8

717   channel triggers UCP1-dependent thermogenesis and prevents obesity. J Mol Cell Biol.

718   2012;4: 88–96. doi:10.1093/jmcb/mjs001

719   23.   Motyl KJ, DeMambro VE, Barlow D, Olshan D, Nagano K, Baron R, et al. Propranolol

720   attenuates risperidone-induced trabecular bone loss in female mice. Endocrinology.

721   2015;156: 2374–2383. doi:10.1210/en.2015-1099

722   24.   Fischer AW, Cannon B, Nedergaard J. Optimal housing temperatures for mice to mimic

723   the thermal environment of humans: An experimental study. Mol Metab. 2018.

724   doi:10.1016/j.molmet.2017.10.009

725   25.   Eliceiri K, Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ□: 25 years of

726   image analysis HISTORICAL commentary NIH Image to ImageJ□: 25 years of image

727   analysis. Nat Methods. 2012;9: 671–675. doi:10.1038/nmeth.2089

728   26.   T. Ferreira WR. ImageJ User Guide IJ 1.46r. IJ 146r. 2012; 185. doi:10.1038/nmeth.2019

729   27.   Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Müller R. Guidelines

730   for assessment of bone microstructure in rodents using micro-computed tomography. J

731   Bone Miner Res. 2010;25: 1468–1486. doi:10.1002/jbmr.141

732   28.   Vengellur A, LaPres JJ. The role of hypoxia inducible factor 1α in cobalt chloride induced

733   cell death in mouse embryonic fibroblasts. Toxicol Sci. 2004. doi:10.1093/toxsci/kfh278

734   29.   Seale et al. PRDM16 Controls a Brown Fat/Skeletal Muscle Switch. Bone. 2008;23: 1–7.

735   doi:10.1038/nature07182

30. Sanchez-Gurmaches□; Guertin D. Adipocytes arise from multiple lineages that are

heterogeneously and dynamically distributed Access. Nat Commun. 2014;5: 4099.

doi:10.1038/ncomms5099

31. Cooper MP, Uldry M, Kajimura S, Arany Z, Spiegelman BM. Modulation of PGC-1

coactivator pathways in brown fat differentiation through LRP130. J Biol Chem. 2008.

doi:10.1074/jbc.M805431200

32. Scheller EL, Doucette CR, Learman BS, Cawthorn WP, Khandaker S, Schell B, et al.

Region-specific variation in the properties of skeletal adipocytes reveals regulated and

constitutive marrow adipose tissues. Nat Commun. 2015;6: 1–13.

doi:10.1038/ncomms8808

33. Takashima Y, Ma L, McKemy DD. The development of peripheral cold neural circuits

based on TRPM8 expression. Neuroscience. 2010;169: 828–842.

doi:10.1016/j.neuroscience.2010.05.039

34. Abed E, Labelle D, Martineau C, Loghin A, Moreau R. Expression of transient receptor

potential (TRP) channels in human and murine osteoblast-like cells. Mol Membr Biol.

2009;26: 146–158. doi:10.1080/09687680802612721

35. Takahashi N, Matsuda Y, Sato K, De Jong PR, Bertin S, Tabeta K, et al. Neuronal TRPV1

activation regulates alveolar bone resorption by suppressing osteoclastogenesis via CGRP.

Sci Rep. 2016;6: 1–11. doi:10.1038/srep29294

36. Nilius B, Voets T, Peters J. TRP channels in disease. Sci STKE. 2005;2005: 1–10.

37. Wee NKY, Lorenz MR, Bekirov Y, Jacquin MF, Scheller EL. Shared autonomic

pathways connect bone marrow and peripheral adipose tissues across the central neuraxis.

Front Endocrinol (Lausanne). 2019;10: 1–16. doi:10.3389/fendo.2019.00668

38.   Bautista DM, Siemens J, Glazer JM, Tsuruda PR, Basbaum AI, Stucky CL, et al. The

760   menthol receptor TRPM8 is the principal detector of environmental cold. Nature.

761   2007;448: 204–208. doi:10.1038/nature05910

762   39.   Shin H, Ma Y, Chanturiya T, Cao Q, Wang Y, Kadegowda AKG, et al. Lipolysis in

763   Brown Adipocytes Is Not Essential for Cold-Induced Thermogenesis in Mice. Cell Metab.

764   2017;26: 764-777.e5. doi:10.1016/j.cmet.2017.09.002

765   40.   Kim SN, Jung YS, Kwon HJ, Seong JK, Granneman JG, Lee YH. Sex differences in

766   sympathetic innervation and browning of white adipose tissue of mice. Biol Sex Differ.

767   2016;7: 1–13. doi:10.1186/s13293-016-0121-7

768   41.   Rosell M, Kaforou M, Frontini A, Okolo A, Chan YW, Nikolopoulou E, et al. Brown and

769   white adipose tissues: Intrinsic differences in gene expression and response to cold

770   exposure in mice. Am J Physiol - Endocrinol Metab. 2014;306.

771   doi:10.1152/ajpendo.00473.2013

772   42.   Clookey SL, Welly RJ, Shay D, Woodford ML, Fritsche KL, Rector RS, et al. Beta 3

773   adrenergic receptor activation rescues metabolic dysfunction in female estrogen receptor

774   alpha-null mice. Front Physiol. 2019;10: 1–15. doi:10.3389/fphys.2019.00009

775   43.   Cankar K, Music M, Finderle Z. Cutaneous microvascular response during local cold

776   exposure - the effect of female sex hormones and cold perception. Microvasc Res.

777   2016;108: 34–40. doi:10.1016/j.mvr.2016.07.006

778   44.   Smeester BA, O'Brien EE, Michlitsch KS, Lee JH, Beitz AJ. The relationship of bone-

779   tumor-induced spinal cord astrocyte activation and aromatase expression to mechanical

780   hyperalgesia and cold hypersensitivity in intact female and ovariectomized mice.

781   Neuroscience. 2016;324: 344–354. doi:10.1016/j.neuroscience.2016.03.030

782   45.   Weyer AD, Lehto SG. Development of TRPM8 antagonists to treat chronic pain and

783   migraine. Pharmaceuticals. 2017;10: 1–9. doi:10.3390/ph10020037

784   46.   Caudle RM, Caudle SL, Jenkins AC, Ahn AH, Neubert JK. Sex differences in mouse

785   Transient Receptor Potential Cation Channel, Subfamily M, Member 8 expressing

786   trigeminal ganglion neurons. PLoS One. 2017;12: 1–20.

787   doi:10.1371/journal.pone.0176753

788   47.   Noguchi W, Ishizuka O, Imamura T, Kurizaki Y, Yamagishi T, Yokoyama H, et al. The

789   relationship between α1-adrenergic receptors and TRPM8 channels in detrusor

790   overactivity induced by cold stress in ovariectomized rats. J Urol. 2013;189: 1975–1981.

791   doi:10.1016/j.juro.2012.10.014

792   48.   Kubo T, Tsuji S, Amano T, Yoshino F, Niwa Y, Kasahara K, et al. Effects of β-estradiol

793   on cold-sensitive receptor channel TRPM8 in ovariectomized rats. Exp Anim. 2017;66:

794   337–343. doi:10.1538/expanim.17-0028

795   49.   Winter Z, Gruschwitz P, Eger S, Touska F, Zimmermann K. Cold temperature encoding

796   by cutaneous TRPA1 and TRPM8-carrying fibers in the mouse. Front Mol Neurosci.

797   2017;10. doi:10.3389/fnmol.2017.00209

798   50.   Feketa V V., Balasubramanian A, Flores CM, Player MR, Marrelli SP. Shivering and

799   tachycardic responses to external cooling in mice are substantially suppressed by TRPV1

800   activation but not by TRPM8 inhibition. Am J Physiol - Regul Integr Comp Physiol.

801   2013;305: 1040–1050. doi:10.1152/ajpregu.00296.2013

802   51.   Ukropec J, Anunciado RP, Ravussin Y, Hulver MW, Kozak LP. UCP1-independent

803   thermogenesis in white adipose tissue of cold-acclimated Ucp1-/- mice. J Biol Chem.

804   2006;281: 31894–31908. doi:10.1074/jbc.M606114200

805   52.   Gao Y, Qimuge NR, Qin J, Cai R, Li X, Chu GY, et al. Acute and chronic cold exposure

806   differentially affects the browning of porcine white adipose tissue. Animal. 2018;12:

807   1435–1441. doi:10.1017/S1751731117002981

808   53.   Liang X, Pan J, Cao C, Zhang L, Zhao Y, Fan Y, et al. Transcriptional response of

809   subcutaneous white adipose tissue to acute cold exposure in mice. Int J Mol Sci. 2019;20.

810   doi:10.3390/ijms20163968

811   54.   Yirmiya R, Goshen I, Bajayo A, Kreisel T, Feldman S, Tam J, et al. Depression induces

812   bone loss through stimulation of the sympathetic nervous system. Proc Natl Acad Sci U S

813   A. 2006;103: 16876–16881. doi:10.1073/pnas.0604234103

Supplemental References

Cooper, M.P., Uldry, M., Kajimura, S., Arany, Z., Spiegelman, B.M., 2008. Modulation of PGC-

1 coactivator pathways in brown fat differentiation through LRP130. J. Biol. Chem. 283,

31960–31967. https://doi.org/10.1074/jbc.M805431200

Sanchez-Gurmaches☐; Guertin, D., 2014. Adipocytes arise from multiple lineages that are

heterogeneously and dynamically distributed Access. Nat Commun. 5, 4099.

https://doi.org/10.1038/ncomms5099

Seale et al., 2008. PRDM16 Controls a Brown Fat/Skeletal Muscle Switch. Bone 23, 1–7.

https://doi.org/10.1038/nature07182

Figure 1

Figure 2

Figure 3

Source of Variation	% of total variation	P value
Interaction	0.2518	0.4360
Sex	89.23	<0.0001
Genotype	0.08133	0.6567

Figure 4

A

Experimental Design

B

2-way ANOVA at 8 hrs:

Source of Variation	% of total variation	P value
Interaction	5.216	0.0187
Genotype	9.561	0.0021
Sex	58.15	<0.0001

C

D

Source of Variation	% of total variation	P value
Interaction	0.1715	0.8168
Sex	1.081	0.5618
Genotype	7.356	0.1367

Figure 5

Figure 7

A

Male

Wild-type (WT)   *Trpm8-KO*

Female

Wild-type (WT)   *Trpm8-KO*

Distal Femur

RT (22°C)

Cold (4°C)

B

Ad.N/T.Ar (#/mm²)

Source of Variation	% of total variation	P value
Interaction	5.126	0.3004
Genotype	1.227	0.6087
Temperature	2.939	0.4303

C

Average Ad.A (µm²)

Source of Variation	% of total variation	P value
Interaction	0.5785	0.7178
Genotype	3.338	0.3888
Temperature	5.915	0.2545

D

T.Ar Adiposity (%)

Source of Variation	% of total variation	P value
Interaction	18.14	0.0452
Genotype	0.0004265	0.9918
Temperature	2.331	0.4529

E

Ad.N/T.Ar (#/mm²)

Source of Variation	% of total variation	P value
Interaction	0.1225	0.8434
Genotype	29.20	0.0052
Temperature	0.05022	0.8993

F

Average Ad.A (µm²)

Source of Variation	% of total variation	P value
Interaction	1.655	0.5242
Genotype	5.539	0.2489
Temperature	2.432	0.4411

G

T.Ar Adiposity (%)

Source of Variation	% of total variation	P value
Interaction	0.4325	0.7234
Genotype	21.97	0.0177
Temperature	0.6035	0.6760