Title Page

Plasmidome analysis of carbapenem-resistant Enterobacteriaceae isolated in Vietnam

Running title. Plasmidome analysis of CRE in Vietnam

Aki Hirabayashi[a], Koji Yahara[a], Satomi Mitsuhashi[b#], So Nakagawa[b], Tadashi Imanishi[b],

Van Thi Thu Ha[c], An Van Nguyen[c##], Son Thai Nguyen[c], Keigo Shibayama[d*], and Masato

Suzuki[a*]

[a]AMR Research Center, National Institute of Infectious Diseases, Tokyo, Japan

[b]Department of Molecular Life Science, Tokai University School of Medicine, Kanagawa,

Japan

[c]Microbiology Department, Hospital 103, Military Medical University, Hanoi, Vietnam

[d]Department of Bacteriology II, National Institute of Infectious Diseases, Tokyo, Japan

*Corresponding authors. Phone: +81-3-5285-1111, Fax: +81-3-561-1150, E-mail: suzuki-

m@nih.go.jp for MS, and Phone: +81-3-5285-1111, Fax: +81-3-561-1150, E-mail:

keigo@nih.go.jp for KS.

[#]Present address. Department of Human Genetics, Yokohama City University

Graduate School of Medicine, Kanagawa, Japan

[##]Present address. Department of Viral Infection and International Health, Graduate

School of Medical Sciences, Kanazawa University, Ishikawa, Japan

Abstract

Carbapenem-resistant Enterobacteriaceae (CRE) represent a serious threat to public health due to limited management of severe infections and high mortality. The rate of resistance of Enterobacteriaceae isolates to major antimicrobials, including carbapenems, is much higher in Vietnam than in Western countries, but the reasons remain unknown due to the lack of genomic epidemiology research. A previous study suggested that carbapenem resistance genes, such as the carbapenemase gene bla_{NDM-1}, spread via plasmids among Enterobacteriaceae in Vietnam. In this study, we performed detection and molecular characterization of bla_{NDM-1}-carrying plasmids in CRE isolated in Vietnam, and identified several possible cases of horizontal transfer of plasmids both within and among species of bacteria. Twenty-five carbapenem-resistant isolates from Enterobacteriaceae clinically isolated in a reference medical institution in Hanoi were sequenced on Illumina short-read sequencers, and 12 isolates harboring bla_{NDM-1} were sequenced on an Oxford Nanopore Technologies long-read sequencer to obtain complete plasmid sequences. Most of the plasmids co-carried genes conferring resistance to clinically relevant antimicrobials, including third-generation cephalosporins, aminoglycosides, and fluoroquinolones, in addition to bla_{NDM-1}, leading to multidrug resistance of their bacterial hosts. These results provide insight into the genetic basis of CRE in Vietnam, and could help control nosocomial infections.

Introduction

Carbapenem-resistant Enterobacteriaceae (CRE) represent a serious threat to public health due to limited management of severe infections and high mortality. Carbapenem-hydrolyzing β-lactamase (carbapenemase) genes, such as bla_{NDM}, bla_{KPC}, and bla_{OXA-48}, that confer resistance to a broad range of β-lactams including third-generation cephalosporins and carbapenems are predominantly encoded on conjugative plasmids and have been transferred among Enterobacteriaceae around the world [1]. Therefore, it is very important to detect carbapenemase-carrying plasmids and to aggregate information on the plasmids spreading in each country. According to a report from China [2], CRE isolates simultaneously harbored other important antimicrobial resistance (AMR) genes, including those encoding extended-spectrum β-lactamase (ESBL) that hydrolyze third-generation cephalosporins, plasmid-mediated quinolone resistance, and aminoglycoside resistance genes, suggesting that the isolates harbored plasmids associated with multidrug resistance.

Several studies have revealed the co-existence of clinically relevant AMR genes on the same plasmid. The acquired fluoroquinolone resistance gene *qnr*, which encodes an efflux pump that excretes quinolones, was co-carried with carbapenemase genes in the same plasmids in Taiwan [3], and with bla_{KPC} in Israel [4]. Aminoglycoside acetyltransferases such as AAC(6')-Ib and 16S ribosomal RNA (rRNA) methyltransferases such as ArmA and RmtB are involved in well-known mechanisms of high-level resistance to aminoglycosides. In Switzerland, 16S rRNA methyltransferase genes have been reported to be co-carried with bla_{NDM-1}, in the same plasmids [5]. Another acquired fluoroquinolone resistance gene, *qep* [6], which also encodes a

quinolone efflux pump, co-existed with aminoglycoside resistance and ESBL genes in

the same plasmids in China [7]. The spread of plasmids co-carrying fluoroquinolone,

aminoglycoside, and carbapenem resistance genes both within and among species of

bacteria poses a threat to public health. In this study, we report the detailed genetic

structures of the plasmids that co-carried bla_{NDM-1} with other important AMR genes; this

knowledge provides insight into characteristics and transmission of the plasmids among

Enterobacteriaceae in Vietnam.

The Vietnamese Ministry of Health developed a national action plan on AMR in 2013

[8]. The Viet Nam Resistance (VINARES) project implements measures in the following

areas: infection control and healthcare-associated infections (HAI), antimicrobial

consumption, and microbiological analysis and reporting capacity [9]. Through VINARES,

a surveillance network consisting of 16 central and provincial-level medical institutions

was established in 2013. Vietnam has implemented AMR surveillance and stewardship

according to the national action plan, and research on AMR using whole-genome

sequencing has begun. According to VINARES data, the rate of resistance of *Klebsiella*

pneumoniae to third-generation cephalosporins, carbapenems, aminoglycoside, and

fluoroquinolones was 66.4%, 17.1%, 29.5%, and 53.0%, respectively, in Vietnam [10],

whereas the rate was 11.4%, 0.6%, 7.9%, and 9.3%, respectively, in UK [11], and 17.2,

4.3%, 4.5%, and 16.8%, respectively, in U.S.A [12]. The rate of resistance to major

antimicrobials, including carbapenems, is much higher in Vietnam than in Western

countries, but the reasons remain unknown due to the lack of genomic epidemiology

research. A point prevalence survey with screening for colonization by CRE in 12 medical

institutions in Vietnam revealed that 52% of patients were colonized with CRE [13].

Previously, we frequently detected bla_{NDM} genes in CRE isolated in Vietnam [14], and

suspected that bla_{NDM} genes were spreading via plasmids among Enterobacteriaceae in

medical institutions and communities. To examine the prevalence of important AMR

genes in plasmids, whole-genome sequencing of CRE isolates is required. Long-read

sequencing is useful for constructing complete sequences of whole plasmids and

tracking horizontal transfer of AMR plasmids during nosocomial infections [15]. Recently,

long-read nanopore sequencing using MinION from Oxford Nanopore Technologies

(ONT) has been applied to such investigations [16,17].

In this study, we performed nanopore sequencing to achieve comprehensive

characterization of plasmids carrying bla_{NDM-1} among clinical isolates of

Enterobacteriaceae from Vietnam.

Results

A total of 122 isolates of ESBL-producing Enterobacteriaceae were collected in daily

diagnosis in a reference medical institution in Hanoi, Vietnam between 2013 and 2017.

Twenty-five CRE isolates, including 19 *K. pneumoniae*, two *Escherichia coli*, two

Citrobacter freundii, one *Morganella morganii*, and one *Proteus mirabilis* isolates, were

sequenced on Illumina short-read sequencers, and confirmed actually carbapenemase-

positive and carrying carbapenemase genes (12 bla_{NDM-1}, five bla_{NDM-4}, eight bla_{KPC-2}, one

bla_{OXA-48}, and one $bla_{OXA-181}$). As shown in Fig. 1, many isolates had the same types of

carbapenemase genes and plasmid replicons. For example, *M. morganii* MH16-367M,

P. mirabilis MH13-009N, *C. freundii* MH17-012N, and *E. coli* MH13-051M had the IncN

replicon and bla_{NDM-1}, suggesting that they shared a common plasmid carrying bla_{NDM-1}

that had been transferred via conjugation. Multilocus sequence typing (MLST) classified

three *K. pneumoniae* isolates (MH15-289M, MH16-398D, and MH15-258M) as

sequence type 11 (ST11), and another three *K. pneumoniae* isolates (MH15-208H,

MH15-191M, and MH13-055M) as sequence type 395 (ST395). The isolates that

belonged to the same STs had the same replicons and bla_{NDM-1}, suggesting that they

shared a common plasmid carrying bla_{NDM-1} and had been transferred via conjugation or

suggesting that their common origin clonally disseminated. Enterobacteriaceae

harboring bla_{NDM-1} were first reported in Vietnam in 2013 [18], suggesting that bla_{NDM-1} is

both important and prevalent in this country; hence, there is an urgent need to elucidate

the mechanism of plasmid transmission. Twelve isolates harboring bla_{NDM-1} and one

isolate harboring bla_{NDM-4} were sequenced on a MinION nanopore sequencer, yielding

complete sequences of the plasmids carrying the gene. The results of BLAST search

revealed that some plasmids had high sequence identities with each other and with

plasmids in the GenBank database (Figs. 2, 3, 4, and 5). However, bla_{NDM-4}-carrying

plasmid pMH16-335M_1 was not identical to any other plasmids (data not shown).

P. mirabilis MH13-009N and *M. morganii* MH16-367M had only one plasmid, pMH13-

009N_1 (72.6 kb) and pMH16-367M_1 (73.1 kb), respectively; the sequences of these

plasmids were almost identical, including the bla_{NDM-1} and IncFII(pSE11)::IncN hybrid

replicon regions (Figs. 2A and 2B). Both of the plasmids carried bla_{NDM-1} and an integron

cassette that contains the class 1 integrase gene (*intI1*) and the AMR genes (bla_{TEM-1b},

rmtB, and *qepA1*). The same integron cassette was observed in previous reports [6][19].

In another case, *E. coli* MH13-051M had five plasmids, and *C. freundii* MH17-012N had

four plasmids. *E. coli* MH13-051M harbored an IncFII(pRSB107)::IncN hybrid plasmid

carrying bla_{NDM-1}, pMH13-051M_1 (111.5 kb), which shared common structures with two

plasmids, pMH17-012N_1 (39.2 kb) and pMH17-012N_2 (87.6 kb), both found in *C.*

freundii MH17-012N (Figs. 2C and 2D). The plasmid pMH13-051M_1 and the smaller

IncR plasmid pMH17-012N_1 carried AMR genes (bla_{NDM-1} and bla_{TEM-1b}), and the only

pMH13-051M_1 carried *rmtB* and ESBL gene $bla_{CTX-M-55}$. The larger

IncFII(pHN7A8)::IncN hybrid plasmid pMH17-012N_2 carried no known AMR genes, but

did have a set of conjugation-associated type IV secretion system (T4SS) genes, which

shared common structures with plasmid pMH13-051M_1.

BLAST search showed that *P. mirabilis* pMH13-009N_1 was highly identical with

Salmonella enterica FDAARGOS_70 plasmid unnamed1 from human in U.S.A. (100.0%

identity in 61% region of pMH13-009N_1), which carried bla_{TEM-1B} (Fig. 3A). Also, *C.*

freundii pMH17-012N_2 was highly identical with *E. coli* p103-2-4 in China (99.9%

identity in 94% region of pMH17-012N_2) and with *Salmonella enterica* serovar

Enteritidis p12367A from human in China (99.9% identity in 73% region of pMH17-

012N_2) (Fig. 3B). Though the plasmid pMH17-012N_2 had no AMR gene, the plasmids

p103-2-4 and p12367A carried several important AMR genes, such as *rmtB*, *bla*TEM-1b,

and *bla*CTX-M-55 (Fig. 3B).

Furthermore, as shown in Fig. 4, three *K. pneumoniae* isolates belonging to ST11

(MH15-289M, MH16-398D, and MH15-258M) shared identical IncFIA(HI1)::IncA/C2

hybrid plasmids carrying *bla*NDM-1 (147–149 kb), and all of these plasmids carried other

important AMR genes, such as *qnrB9*, *rmtC*, *bla*OXA-1, and *bla*CTX-M-15. BLAST search

showed that these three plasmids were highly identical with *K. pneumoniae* plasmid

tig00000169_pilon in U.S.A. (99.8% identity in 82% region of pMH15-289M_1) and with

E. coli pK71-77-1-NDM from human in Norway (99.9% identity in 62% region of pMH15-

289M_1). The plasmids tig00000169_pilon and pK71-77-1-NDM carried *rmtC*, *bla*NDM-1,

and *bla*CMY-6, and the plasmid tig00000169_pilon further   carried *qnrB58* and *bla*SHV-11.

K. pneumoniae MH16-398D had 235 sequence variants (214 single-nucleotide variants

[SNVs], 14 multiple-nucleotide variants [MNVs], four deletions, two insertions, and one

replacement), and *K. pneumoniae* MH15-258M had six sequence variants (five SNVs

and one replacement) relative to *K. pneumoniae* MH15-289M.

As shown in Fig. 5, another three *K. pneumoniae* isolates belonging to ST395 (MH15-

208H, MH15-191M, and MH13-055M) shared identical IncFII(Yp) plasmids carrying

*bla*NDM-1 (75–76 kb), and all plasmids carried another important AMR gene, *rmtB*. BLAST

search showed that these three plasmids were highly identical with *Enterobacter*

hormaechei pNDM1_045001 from human in China (99.9% identity in 98% region of

pMH15-208H_1) and with *K. pneumoniae* pSECR18-2374C from human in South Korea

(99.9% identity in 95% region of pMH15-208H_1). The plasmid pNDM1_045001 carried

rmtB, *bla*TEM-1b, and *bla*NDM-1, and the plasmid pSECR18-2374C carried *rmtB*, *bla*TEM-1b,

and bla_{NDM-4}. *K. pneumoniae* MH15-191M had three sequence variants (two SNVs and

one MNV) and *K. pneumoniae* MH13-055M had only one SNV relative to *K. pneumoniae*

MH15-208H.

The bla_{NDM-1}-carrying plasmids, *P. mirabilis* pMH13-009N_1 and *M. morganii* pMH16-

367M_1 were successfully transferred to recipient *E. coli* at a frequency of 10^{-2}–10^{-3} *in*

vitro, however, *C. freundii* pMH17-012N_1 and pMH17-012N_2, *E. coli* pMH13-051M_1,

K. pneumoniae pMH15-258M_1, and *K. pneumoniae* pMH15-208H_1 were not

transferred to recipient *E. coli* under our experimental conditions (data not shown).

Discussion

In this study, we performed plasmidome analysis (i.e., comprehensive genetic analysis of plasmids) on plasmids carrying carbapenemase genes in CRE isolated from patients in Vietnam. We focused on CRE isolates harboring bla_{NDM}, one of the most important carbapenemase genes in the world, and completely sequenced 12 plasmids carrying bla_{NDM-1} from *K. pneumoniae*, *E. coli*, *C. freundii*, *M. morganii*, and *P. mirabilis* isolates and one plasmid carrying bla_{NDM-4} from *K. pneumoniae* isolate. Some plasmids were nearly identical to each other, suggesting that they represent common and important AMR plasmids disseminated in this country. Moreover, the co-existence of clinically relevant AMR genes, such as ESBL, aminoglycoside resistance, and fluoroquinolone resistance genes, with bla_{NDM-1} was observed in plasmids pMH13-009N_1, pMH16-367M_1, pMH15-289M_1, pMH16-398D_1, and pMH15-258M_1. Hence, the spread of the plasmids both within and among species of bacteria will pose a threat to public health.

Almost identical IncFII(pSE11)::IncN hybrid plasmids, pMH13-009N_1 and pMH16-367M_1, were found in *P. mirabilis* and *M. morganii* isolates, respectively (Figs. 2A and 2B). These plasmid shared common structures with the IncN plasmid from *S. enterica* isolate in U.S.A (Fig. 3A). IncN plasmids are prevalent in the microbiota of animals, and disseminate AMR genes such as $bla_{CTX-M-1}$ and *qnr* [20], suggesting that their origin plasmid could be disseminated in communities, including humans, animals, and the environment, in Vietnam and other countries.

The IncFII(pRSB107)::IncN hybrid plasmid carrying bla_{NDM-1}, pMH13-051M_1 in *E. coli* MH13-051M, had partial sequence identity with an IncR plasmid carrying bla_{NDM-1}, pMH17-012N_1, and an IncFII(pHN7A8)::IncN hybrid plasmid carrying conjugation genes but no known AMR gene, pMH17-012N_2, both in *C. freundii* MH17-012N (Figs.

2C and 2D). The plasmid pMH17-012N_2 shared common structures with the plasmids

p103-2-4 from *E. coli* in China and p12367A from *Salmonella enterica* in China, carrying

many AMR genes (Fig. 3B). Rearrangement of conjugation genes plays an important

role in the dissemination of AMR genes between bacteria [21]. Because known IncR

plasmids are non-transferable [22], an IncR plasmid carrying AMR genes, such as

pMH17-012N_1, and another plasmid carrying conjugation genes, such as pMH17-

012N_2, could be assumed to have been fused into a single plasmid with both AMR and

conjugation genes, such as pMH13-051M_1, in a bacterium, and then been propagated

by conjugation to another bacterium. Reports of IncR plasmids carrying various AMR

genes have been increasing. The pool of AMR genes on IncR plasmids is thought to

spread to transmissible plasmids via recombination, contributing to the high evolutionary

plasticity of bacterial genomes [23]. IncFII(pRSB107)::IncN hybrid plasmids such as

pMH13-051M_1 are partially homologous to the pHN7A8 plasmid, which carries several

AMR genes, such as $bla_{CTX-M-65}$, *fosA3*, and *rmtB*, and is widespread in *E. coli* from

animals in China [24]. The plasmid pMH13-051M_1 also carries $bla_{CTX-M-55}$, *fosA3*, and

rmtB in addition to bla_{NDM-1} and bla_{TEM-1B}.

The IncFII(pSE11)::IncN, IncFII(pRSB107)::IncN, and IncFII(pHN7A8)::IncN hybrid

plasmids shown in Figs. 2A and 2C consisted of two replicons belonging to different

incompatibility groups, IncFII and IncN. Similarly, IncFIA(HI1)::IncA/C2 hybrid plasmids

from three *K. pneumoniae* ST11 isolates consisted of two replicons belonging to different

incompatibility groups, IncFIA and IncA/C2 (Fig. 4). Hybrids of multiple replicons

belonging to different incompatibility groups represent a plasmid strategy for expansion

of host range and dissemination of important acquired genes, including AMR genes,

among bacteria [25]. The plasmid pMH15-289M_1 was similar in structure to those

reported in U.S.A and Europe, and the plasmid pMH15-208H_1 was similar in structure

to those reported in Asia (Figs. 4 and 5). These *K. pneumoniae* and plasmids would have

also been spreading in medical institutions and community in Vietnam.

It is important to estimate whether *K. pneumoniae* isolates belonging to the same STs

were originated from the same clones and spread within the medical institution.

According to a report [26], SNVs accumulated in *K. pneumoniae* (ST258) at a rate of 3.9

SNVs per year. Our analysis of SNVs of *K. pneumoniae* (ST11) revealed 214 SNVs in

MH16-398D relative to MH15-289M; based on the rate of accumulation, it could take

many years for these many SNVs to arise. This suggests that many ST11 clones would

be prevalent in the medical institutions and communities, and that they have intermixed

for a long time. In support of this theory, a report [27] showed that *K. pneumoniae* isolates

from medical institutions, sewage, canals, and agricultural waste water were intermixed

in the phylogenetic classification. Because ST11 is one of major international epidemic

clones of *K. pneumoniae* [28], an ST11 clone carrying this AMR plasmid could have

disseminated in a medical institution or local community in Hanoi, Vietnam. Another three

isolates of *K. pneumoniae* (ST395) are more clonal because only a few SNVs were

present in MH15-191M and MH13-055M relative to MH15-208H.

Conclusion

Plasmidome analysis in this study enabled estimation of the origins and diversity of

plasmids carrying bla_{NDM-1} in Vietnam. Hybrid analysis with both Illumina short-read and

ONT long-read sequencing is a promising method for detecting important AMR plasmids

in CRE isolates and controlling nosocomial infections.

Materials and Methods

Bacterial isolates

A total of 122 ESBL-producing Enterobacteriaceae isolates were collected from blood,

sputum, urine, and pus from patients in daily diagnosis in a reference medical institution

in Hanoi, Vietnam between 2013 and 2017. Bacterial species identification and

antimicrobial susceptibility testing were performed by Vitek 2 (BioMèrieux) and E-test

(BioMèrieux). Carbapenemase production was examined by CarbaNP test according to

the Clinical and Laboratory Standards Institute (CLSI) guidelines. Major carbapenemase

genes (bla_{NDM}, bla_{KPC}, bla_{OXA-48}, bla_{IMP}, and bla_{VIM}) were detected by a multiplex PCR

method [29]. Twenty-five carbapenem-resistant and carbapenemase-positive isolates

(19 *K. pneumoniae*, two *E. coli*, two *C. freundii*, one *M. morganii*, and one *P. mirabilis*)

were subjected to the whole-genome sequencing analysis described below.

Whole-genome sequencing and bioinformatics analysis

Whole genome sequencing using MiniSeq, MiSeq, and/or HiSeq systems (Illumina)

was performed for phylogenetic and MLST analysis, and detection of acquired AMR

genes and plasmid replicon types. A maximum-likelihood phylogenetic tree was

generated by PhyML from a concatenated core-gene alignment consisting of 13,305

SNVs constructed using the Roary pipeline (https://sanger-pathogens.github.io/Roary/).

Sequence types, carbapenemase genes, and plasmid replicon types were analyzed

using MLST v2.0, ResFinder v3.2 with minimum threshold of 90% identity and 60%

coverage, and PlasmidFinder v2.1 with minimum threshold of 90% identity and 60%

coverage, respectively, on the Center for Genomic Epidemiology (CGE) server at

Technical University of Denmark (http://www.genomicepidemiology.org).

Twelve Enterobacteriaceae isolates with bla_{NDM-1} (seven *K. pneumoniae*, one *E. coli*,

two *C. freundii*, one *M. morganii*, and one *P. mirabilis*) and one *K. pneumoniae* isolate

with bla_{NDM-4} were further sequenced on a MinION nanopore sequencer (Oxford

Nanopore Technologies) using the SQK-RAD002 or SQK-RBK001 kits and R9.4

flowcells to obtain complete sequences of plasmids carrying carbapenem resistance

genes. *De novo* assembly was performed using Canu v1.5 [30] and Miniasm [31], error

correction using Illumina reads with CLC Genomics Workbench v12.0 (QIAGEN), and

coding sequences (CDS) annotation using the PATRIC server (https://www.patricbrc.org).

Sequence variant detection was performed using CLC Genomics Workbench. Linear

comparison of sequences was performed using BLAST and visualized by Easyfig

(http://mjsull.github.io/Easyfig/). bla_{NDM-1}, other important AMR genes (ARG), type IV

secretion system (T4SS)-associated genes involved in conjugation that were detected

by SecReT4 program [32], and mobile gene elements (MGE) were identified manually

from CDS annotations.

Draft genome and complete plasmid sequences of CRE isolated in Vietnam have been

deposited at GenBank/EMBL/DDBJ under BioProject number PRJDB6655.

Bacterial conjugation

Bacterial conjugation was performed according to the following protocol. The same

amount of Luria-Bertani (LB) broth cultures of each donor bacteria and the recipient

azide–resistant *E. coli* J53 (*F- met pro Azi^r*), were mixed and spotted onto Mueller-Hinton

agar and then incubated at 37°C overnight. The mixed cells were recovered and

suspended into PBS buffer, plated onto LB agar after 10-fold serial dilution, and

incubated at 37°C overnight. Transconjugants were selected on LB agar containing 2

µg/mL of meropenem and 100 µg/mL of sodium azide.

Acknowledgements

We thank Meiko Takeshita for technical assistance and are grateful to the participating

medical institution for providing the bacterial isolates and clinical information. This work

was supported by Grants-in-Aid for the Research Program on Emerging and Re-

emerging Infectious Diseases (JP18fk0108020, JP19fk0108061, JP19fk0108093,

JP19fk0108052, and JP19fk0108049) and the Japan Initiative for Global Research

Network on Infectious Diseases (J-GRID) (JP17fm0108023, JP18fm0108003, and

JP19fm0108006) from the Japan Agency for Medical Research and Development

(AMED). This work was also supported by the Cooperative Research Grant from

Research Center on Tropical Disease, Institute of Tropical Medicine, Nagasaki University

(2019-Ippan-20).

Conflicts of Interest

None to declare.

Figure Legends

Figure 1.

CRE isolates sequenced on Illumina systems. Red nodes, but not black nodes, show

isolates subsequently sequenced with ONT MinION. Bar lengths represent the number

of substitutions per site in the core genome. Minimum inhibitory concentrations (MICs)

of meropenem (MEM) and imipenem (IPM) of isolates, years in which the bacteria were

isolated, plasmid replicon types, and sizes of bla_{NDM}-carrying plasmids are shown.

Figure 2.

bla_{NDM-1}-carrying plasmids in CRE isolates sequenced using ONT MinION and Illumina

systems. Sets of plasmids from *P. mirabilis* MH13-009N and *M. morganii* MH16-367M,

and from *E. coli* MH13-051M and *C. freundii* MH17-012N, are shown. (A) and (C) Linear

comparison of bla_{NDM-1}-carrying plasmid sequences. Sets of plasmids from *P. mirabilis*

MH13-009N and *M. morganii* MH16-367M (A and B), and from *E. coli* MH13-051M and

C. freundii MH17-012N (C and D), are shown. Red, yellow, green, blue, and gray arrows

indicate bla_{NDM-1}, other important AMR genes (ARG), type IV secretion system (T4SS)-

associated genes involved in conjugation, mobile gene elements (MGE), and other

genes, respectively.

Figure 3.

(A) and (B) Linear comparison of plasmid sequences from Vietnam and other countries.

P. mirabilis pMH13-009N_1 and *Salmonella enterica* FDAARGOS_70 plasmid

unnamed1 (accession: CP026053) (A), and *C. freundii* pMH17-012N_2, *E. coli* p103-2-

4 (accession: CP034846), and *Salmonella enterica* serovar Enteritidis p12367A

(accession: CP041177) (B) are shown.

Figure 4.

Linear comparison of bla_{NDM-1}-carrying plasmid sequences from Vietnam and other

countries. *K. pneumoniae* pMH15-289M_1, *K. pneumoniae* pMH16-398D_1, *K.*

pneumoniae pMH15-258M_1, *K. pneumoniae* tig00000169_pilon (accession:

CP021952), and *E. coli* pK71-77-1-NDM (accession: CP040884) are shown.

Figure 5.

Linear comparison of bla_{NDM}-carrying plasmid sequences from Vietnam and other

countries. *K. pneumoniae* pMH15-208H_1, *K. pneumoniae* pMH15-191M_1, *K.*

pneumoniae pMH13-055M_1, *E. hormaechei* pNDM1_045001 (accession: CP043383),

and *K. pneumoniae* pSECR18-2374C (accession: CP041930) are shown.

References

1.   Tzouvelekis LS, Markogiannakis A, Psichogiou M, Tassios PT, Daikos GL.

Carbapenemases in Klebsiella pneumoniae and other Enterobacteriaceae: an

evolving crisis of global dimensions. Clin Microbiol Rev. 2012;25: 682–707.

2.   Huang S, Dai W, Sun S, Zhang X, Zhang L. Prevalence of plasmid-mediated

quinolone resistance and aminoglycoside resistance determinants among

carbapeneme non-susceptible Enterobacter cloacae. PLoS One. 2012;7:

e47636.

3.   Wu JJ, Ko WC, Tsai SH, Yan JJ. Prevalence of plasmid-mediated quinolone

resistance determinants QnrA, QnrB, and QnrS among clinical isolates of

Enterobacter cloacae in a Taiwanese hospital. Antimicrob Agents Chemother.

2007.

4.   Chmelnitsky I, Navon-Venezia S, Strahilevitz J, Carmeli Y. Plasmid-mediated

qnrB2 and carbapenemase gene blaKPC-2 carried on the same plasmid in

carbapenem-resistant ciprofloxacin-susceptible Enterobacter cloacae isolates.

Antimicrob Agents Chemother. 2008.

5.   Poirel L, Dortet L, Bernabeu S, Nordmann P. Genetic features of blaNDM-1-

positive Enterobacteriaceae. Antimicrob Agents Chemother. 2011;55: 5403–7.

6.   Yamane K, Wachino JI, Suzuki S, Kimura K, Shibata N, Kato H, et al. New

plasmid-mediated fluoroquinolone efflux pump, QepA, found in an Escherichia

coli clinical isolate. Antimicrob Agents Chemother. 2007.

7.   Deng Y, He L, Chen S, Zheng H, Zeng Z, Liu Y, et al. F33:A-:B- and F2:A-:B-

plasmids mediate dissemination of rmtB-blaCTX-M-9 group genes and rmtB-

qepA in Enterobacteriaceae isolates from pets in China. Antimicrob Agents

Chemother. 2011;55: 4926–9.

8.   Kinh N Van, Wertheim HFL, Thwaite GE, Khue LN, Thai CH, Khoa NT, et al. Developing an antimicrobial resistance reference laboratory and surveillance programme in Vietnam. The Lancet Global Health. 2017.

9.   Wertheim HFL, Chandna A, Vu PD, Pham C Van, Nguyen PDT, Lam YM, et al. Providing impetus, tools, and guidance to strengthen national capacity for antimicrobial stewardship in Viet Nam. PLoS Med. 2013;10: e1001429.

10.   Vu TVD, Do TTN, Rydell U, Nilsson LE, Olson L, Larsson M, et al. Antimicrobial susceptibility testing and antibiotic consumption results from 16 hospitals in Viet Nam: The VINARES project 2012–2013. J Glob Antimicrob Resist. 2019;18: 269–278.

11.   Monnet DL, Eemborg HD, Andersen SR, Schöller C, Sørensen TL, Bager F. Surveillance of antimicrobial resistance in Denmark. Eurosurveillance. 2000.

12.   Sanchez G V., Master RN, Clark RB, Fyyaz M, Duvvuri P, Ekta G, et al. Klebsiella pneumoniae antimicrobial drug resistance, United States, 1998-2010. Emerg Infect Dis. 2013;19: 133–136.

13.   Tran DM, Larsson M, Olson L, Hoang NTB, Le NK, Khu DTK, et al. High prevalence of colonisation with carbapenem-resistant Enterobacteriaceae among patients admitted to Vietnamese hospitals: Risk factors and burden of disease. J Infect. 2019;79: 115–122.

14.   Tran HH, Ehsani S, Shibayama K, Matsui M, Suzuki S, Nguyen MB, et al. Common isolation of New Delhi metallo-beta-lactamase 1-producing Enterobacteriaceae in a large surgical hospital in Vietnam. Eur J Clin Microbiol Infect Dis. 2015;34: 1247–1254.

15.   Conlan S, Thomas PJ, Deming C, Park M, Lau AF, Dekker JP, et al. Single-

molecule sequencing to track plasmid diversity of hospital-associated

carbapenemase-producing Enterobacteriaceae. Sci Transl Med. 2014;6:

254ra126.

16.   Lemon JK, Khil PP, Frank KM, Dekker JP. Rapid Nanopore Sequencing of

Plasmids and Resistance Gene Detection in Clinical Isolates. Mellmann A,

editor. J Clin Microbiol. 2017;55: 3530–3543.

17.   Gorrie CL, Mirceta M, Wick RR, Judd LM, Wyres KL, Thomson NR, et al.

Antimicrobial-Resistant Klebsiella pneumoniae Carriage and Infection in

Specialized Geriatric Care Wards Linked to Acquisition in the Referring Hospital.

Clin Infect Dis. 2018;67: 161–170.

18.   Hoang TH, Wertheim H, Minh NB, Duong TN, Anh DD, Phuong TTL, et al.

Carbapenem-Resistant Escherichia coli and Klebsiella pneumoniae Strains

Containing New Delhi Metallo-Beta-Lactamase Isolated from Two Patients in

Vietnam: Table 1. J Clin Microbiol. 2013;51: 373–374.

19.   Berçot B, Poirel L, Nordmann P. Plasmid-mediated 16S rRNA methylases

among extended-spectrum beta-lactamase-producing Enterobacteriaceae

isolates. Antimicrob Agents Chemother. 2008;52: 4526–7.

20.   Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. Characterization of IncN

plasmids carrying blaCTX-M-1 and qnr genes in Escherichia coli and Salmonella

from animals, the environment and humans. J Antimicrob Chemother. 2013;68:

333–339.

21.   He S, Chandler M, Varani AM, Hickman AB, Dekker JP, Dyda F. Mechanisms of

Evolution in High-Consequence Drug Resistance Plasmids. MBio. 2016;7:

e01987-16.

22.   Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B,

Guerra B, et al. Plasmids carrying antimicrobial resistance genes in

Enterobacteriaceae. J Antimicrob Chemother. 2018;73: 1121–1137.

23.   Compain F, Frangeul L, Drieux L, Verdet C, Brisse S, Arlet G, et al. Complete

Nucleotide Sequence of Two Multidrug-Resistant IncR Plasmids from Klebsiella

pneumoniae. Antimicrob Agents Chemother. 2014;58: 4207–4210.

24.   He L, Partridge SR, Yang X, Hou J, Deng Y, Yao Q, et al. Complete nucleotide

sequence of pHN7A8, an F33:A-:B-type epidemic plasmid carrying blaCTX-M-

65, fosA3 and rmtB from China. J Antimicrob Chemother. 2013;68: 46–50.

25.   Garza-Ramos U, Barrios H, Hernandez-Vargas MJ, Rojas-Moreno T, Reyna-

Flores F, Tinoco P, et al. Transfer of quinolone resistance gene qnrA1 to

Escherichia coli through a 50 kb conjugative plasmid resulting from the splitting

of a 300 kb plasmid. J Antimicrob Chemother. 2012;67: 1627–1634.

26.   Bowers JR, Kitchel B, Driebe EM, MacCannell DR, Roe C, Lemmer D, et al.

Genomic Analysis of the Emergence and Rapid Global Dissemination of the

Clonal Group 258 Klebsiella pneumoniae Pandemic. Bengoechea JA, editor.

PLoS One. 2015;10: e0133727.

27.   Runcharoen C, Moradigaravand D, Blane B, Paksanont S, Thammachote J,

Anun S, et al. Whole genome sequencing reveals high-resolution

epidemiological links between clinical and environmental Klebsiella pneumoniae.

Genome Med. 2017;9: 6.

28.   Mathers AJ, Peirano G, Pitout JDD. The role of epidemic resistance plasmids

and international high-risk clones in the spread of multidrug-resistant

Enterobacteriaceae. Clin Microbiol Rev. 2015;28: 565–91.

29. Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of

acquired carbapenemase genes. Diagn Microbiol Infect Dis. 2011;70: 119–123.

30. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu:

scalable and accurate long-read assembly via adaptive k-mer weighting and

repeat separation. Genome Res. 2017;27: 722–736.

31. Li H. Minimap and miniasm: Fast mapping and de novo assembly for noisy long

sequences. Bioinformatics. 2016;32: 2103–2110.

32. Bi D, Liu L, Tai C, Deng Z, Rajakumar K, Ou HY. SecReT4: A web-based

bacterial type IV secretion system resource. Nucleic Acids Res. 2013;41.

Fig. 1

Fig. 2

(A)

P. mirabilis
pMH13-009N_1
[IncFII(pSE11)::IncN]
72.6 Kb

M. morganii
pMH16-367M_1
[IncFII(pSE11)::IncN]
73.1 Kb

bla_{NDM-1} bla_{TEM-1b} $rmtB$ $qepA1$

(B)

P. mirabilis
MH13-009N
(1 plasmid)
73 Kb bla_{NDM-1}

M. morganii
MH16-367M
(1 plasmid)
73 Kb bla_{NDM-1}

87%   100%

(C)

C. freundii
pMH17-012N_1
[IncR]
39.2 Kb

E. coli
pMH13-051M_1
[IncFII(pRSB107)::IncN]
111.5 Kb

C. freundii
pMH17-012N_2
[IncFII(pHN7A8)::IncN]
87.6 Kb

bla_{NDM-1} bla_{TEM-1b}

bla_{NDM-1} bla_{TEM-1b} $rmtB$ $bla_{CTX-M-55}$

(D)

E. coli
MH13-051M
(5 plasmids)
112 Kb bla_{NDM-1}

C. freundii
MH17-012N
(4 plasmids)
89 Kb 88 Kb bla_{NDM-1}

95%   100%

| bla_{NDM-1} | ARG | T4SS | MGE | Others |

Fig. 3

(A)

P. mirabilis
pMH13-009N_1
[IncFII(pSE11)::IncN]
72.6 Kb, Vietnam

S. enterica FDAARGOS_70
unnamed1
[IncN]
47.8 Kb, USA

(B)

C. freundii
pMH17-012N_2
[IncFII(pHN7A8)::IncN]
87.6 Kb, Vietnam

E. coli
p103-2-4
[IncFII(pHN7A8)]
100.8 Kb, China

S. enterica Enteritidis
p12367A
[IncFII(pHN7A8)::IncX1]
111.8 Kb, China

Fig. 4

K. pneumoniae
pMH15-289M_1
[IncFIA(HI1)::IncA/C2]
147.8 Kb, Vietnam

K. pneumoniae
pMH16-398D_1
[IncFIA(HI1)::IncA/C2]
147.8 Kb, Vietnam

K. pneumoniae
pMH15-258M_1
[IncFIA(HI1)::IncA/C2]
148.8 Kb, Vietnam

K. pneumoniae
tig00000169_pilon
[IncFIA(HI1)::IncA/C2]
176.3 Kb, USA

E. coli
pK71-77-1-NDM
[IncA/C2]
145.3 Kb, Norway

94%   100%

bla_{NDM-1} ARG   T4SS   MGE   Others

Fig. 5

K. pneumoniae pMH15-208H_1 [IncFII(Yp)] 75.4 Kb, Vietnam

K. pneumoniae pMH15-191M_1 [IncFII(Yp)] 75.6 Kb, Vietnam

K. pneumoniae pMH13-055M_1 [IncFII(Yp)] 75.3 Kb, Vietnam

E. hormaechei pNDM1_045001 [IncFII(Yp)] 85.7 Kb, China

K. pneumoniae pSECR18-2374C [IncFII(Yp)] 86.0 Kb, South Korea

bla_{NDM-1} ARG   T4SS   MGE   Others

99%   100%