Title Page

Plasmid analysis of NDM metallo-β-lactamase-producing Enterobacterales isolated in

Vietnam

Running title. Plasmid analysis of NDM-producing Enterobacterales in Vietnam

Aki Hirabayashi[a], Koji Yahara[a], Satomi Mitsuhashi[b#], So Nakagawa[b], Tadashi Imanishi[b],

Van Thi Thu Ha[c], An Van Nguyen[c##], Son Thai Nguyen[c], Keigo Shibayama[d*], and Masato

Suzuki[a*]

[a]AMR Research Center, National Institute of Infectious Diseases, Tokyo, Japan

[b]Department of Molecular Life Science, Tokai University School of Medicine, Kanagawa,

Japan

[c]Microbiology Department, Hospital 103, Military Medical University, Hanoi, Vietnam

[d]Department of Bacteriology II, National Institute of Infectious Diseases, Tokyo, Japan

*Corresponding authors. Phone: +81-3-5285-1111, Fax: +81-3-561-1150, E-mail: suzuki-

m@nih.go.jp for MS, and Phone: +81-3-5285-1111, Fax: +81-3-561-1150, E-mail:

keigo@nih.go.jp for KS.

[#]Present address. Medical Genomics, Medical Research Institute, Tokyo Medical and

Dental University, Tokyo, Japan

[##]Present address. Department of Viral Infection and International Health, Graduate

School of Medical Sciences, Kanazawa University, Ishikawa, Japan

Abstract

Carbapenem-resistant Enterobacterales (CRE) represent a serious threat to public health due to the lack of treatment and high mortality. The rate of antimicrobial resistance of Enterobacterales isolates to major antimicrobials, including carbapenems, is much higher in Vietnam than in Western countries, but the reasons remain unknown due to the lack of genomic epidemiology research. A previous study suggested that carbapenem resistance genes, such as the carbapenemase gene bla_{NDM}, spread via plasmids among Enterobacterales in Vietnam. In this study, we characterized bla_{NDM}-carrying plasmids in Enterobacterales isolated in Vietnam, and identified several possible cases of horizontal transfer of plasmids both within and among species of bacteria. Twenty-five carbapenem-nonsusceptible isolates from a medical institution in Hanoi were sequenced on Illumina short-read sequencers, and 13 bla_{NDM}-positive isolates, including isolates of *Klebsiella pneumoniae*, *Escherichia coli*, *Citrobacter freundii*, *Morganella morganii*, and *Proteus mirabilis*, were further sequenced on an Oxford Nanopore Technologies long-read sequencer to obtain complete plasmid sequences. Almost identical 73 kb IncFII(pSE11)::IncN hybrid plasmids carrying bla_{NDM-1} were found in a *P. mirabilis* isolate and an *M. morganii* isolate. A 112 kb IncFII(pRSB107)::IncN hybrid plasmid carrying bla_{NDM-1} in an *E. coli* isolate had partially identical sequences with a 39 kb IncR plasmid carrying bla_{NDM-1} and an 88 kb IncFII(pHN7A8)::IncN hybrid plasmid in a *C. freundii* isolate. 148–149 kb IncFIA(HI1)::IncA/C2 plasmids and 75–76 kb IncFII(Yp) plasmids, both carrying bla_{NDM-1} were shared among three sequence type 11 (ST11) isolates and three ST395 isolates of *K. pneumoniae*, respectively. Most of the plasmids co-carried genes conferring resistance to clinically relevant antimicrobials, including third-generation cephalosporins, aminoglycosides, and fluoroquinolones, in addition to bla_{NDM}-

49 $_1$. These results provide insight into the genetic basis of CRE in Vietnam, and could help

50 control nosocomial infections.

Introduction

Carbapenems have a broad spectrum of antimicrobial activity and are reserved for the treatment of infections caused by multidrug-resistant Gram-negative bacteria including Enterobacterales. The increase of cases infected by carbapenem-resistant Enterobacterales (CRE) is a serious threat to public health due to limited management of severe infections and risk of increased mortality [1,2]. Carbapenem-hydrolyzing β-lactamase (carbapenemase) genes, such as bla_{NDM}, bla_{KPC}, and bla_{OXA-48}, that confer resistance to a broad range of β-lactams including third-generation cephalosporins and carbapenems are predominantly encoded on conjugative plasmids and have been transferred among Enterobacterales around the world [3]. There was a report that out of 27 carbapenem-resistant *Klebsiella pneumoniae* isolated from patients in a medical institution in Hanoi, Vietnam from 2014 to 2015, two isolates belonging to sequence type 395 (ST395) harbored bla_{NDM-1}, and five isolates belonging to ST15, ST16, and ST2353 harbored bla_{NDM-4} [4]. There was another report that six *K. pneumoniae* isolates with bla_{NDM-1} were isolated from water samples in Hanoi, Vietnam in 2011 and belonging to ST283 [5]. Furthermore, we previously detected bla_{NDM} in 68.1% (47/69) of CRE isolates from 45 patients in a medical institution in Hanoi, Vietnam from 2010 to 2012, and some of isolates could be shown to transfer bla_{NDM} to *Escherichia coli* [6], suggesting that bla_{NDM} had spread via plasmids among Enterobacterales in medical institutions and communities in Vietnam.

The common replicon types of bla_{NDM}-carrying plasmids associated with Enterobacterales includes IncA/C, IncFIA, IncFIB, IncFII, IncH, IncL/M, IncN, and IncX [7,8]. The plasmids occasionally co-carried other clinically relevant antimicrobial resistance (AMR) genes, such as extended-spectrum β-lactamase (ESBL) genes

conferring resistance to third-generation cephalosporins, such as bla_{CTX-M}, 16S ribosomal

RNA (rRNA) methyltransferase genes conferring high-level resistance to

aminoglycosides, such as *armA* and *rmt*, and fluoroquinolone resistance genes, such as

qnr and *qep* [9]. Metallo-β-lactamase gene bla_{NDM-1} is often observed in a Tn*125*

transposon in carbapenem-resistant *Acinetobacter* spp. Tn*125* is bracketed by two

copies of insertion sequences, IS*Aba125*, belonging to the IS*30* family [10,11]. On

plasmids of CRE, bla_{NDM-1} is also observed in the Tn*125*-like mobile gene element (MGE)

which contains IS*Aba125* [11–14].

In this study, we report the detailed structures of bla_{NDM-1}-carrying plasmids, and co-

existing AMR genes and MGEs on the plasmids in Enterobacterales isolated in Vietnam.

This knowledge provides insight into genetic characteristics and potential transmissions

of the plasmids among Enterobacterales in Vietnam for the first time.

Results

Molecular characterization of carbapenem-nonsusceptible Enterobacterales isolated in Vietnam

A total of 122 isolates of ESBL-producing Enterobacterales were collected in daily diagnosis from individual patients in a medical institution in Hanoi, Vietnam between 2013 and 2017. Twenty-five Enterobacterales isolates, including *K. pneumoniae* (19/25, 76.0%), *E. coli* (2/25, 8.0%), *Citrobacter freundii* (2/25, 8.0%), *Morganella morganii* (1/25, 4.0%), and *Proteus mirabilis* (1/25, 4.0%) isolates, from clinical specimens, including blood (12/25, 48.0%), pus (10/25, 40.0%), urine (2/25, 8.0%), and sputum (1/25, 4.0%), were nonsusceptible to meropenem or imipenem and tested positive with CarbaNP test. The minimum inhibitory concentrations (MICs) of meropenem ranged from 1.5 to >32 μg/mL (median at 6 μg/mL) and those of imipenem ranged from 2 to >32 μg/mL (median at 12 μg/mL) (Fig. 1 and Table S1).

Whole-genome sequencing of all 25 carbapenem-nonsusceptible Enterobacterales isolates with short-read sequencers resulted in the draft genome sequences ranged from 4.1 Mb to 5.9 Mb sizes with 132 to 868 contigs covered by at least x60 sequencing data (Table S1). Carbapenemase gene bla_{NDM-1} was detected in 12 Enterobacterales isolates (12/25, 48.0%), including *K. pneumoniae* (7/25, 28.0%), *C. freundii* (2/25, 8.0%), *E. coli* (1/25, 4.0%), *M. morganii* (1/25, 4.0%), and *P. mirabilis* (1/25, 4.0%) isolates. Carbapenemase gene bla_{NDM-4} was detected in five *K. pneumoniae* isolates (5/25, 20.0%), in which two isolates co-harbored other carbapenemase gene $bla_{OXA-48-like}$ (bla_{OXA-48} and $bla_{OXA-181}$, respectively) (2/25, 8.0%). Carbapenemase gene bla_{KPC-2} was detected in eight Enterobacterales isolates (8/25, 32.0%), including *K. pneumoniae* (7/25, 28.0%) and *E. coli* (1/25, 4.0%) isolates (Fig. 1 and Table S1).

Multilocus sequence typing (MLST) analysis revealed that the 19 carbapenemase-producing *K. pneumoniae* isolates belonged to five sequence types (STs), including ST15 (9/19, 47.4%), ST11 (4/19, 21.1%), ST395 (3/19, 15.8%), ST656 (2/19, 10.5%), and ST16 (1/19, 5.3%). The two *E. coli* isolates with bla_{NDM-1} and bla_{KPC-2} belonged to ST405 and ST457, respectively (Fig. 1). The presence of bla_{NDM-1} and replicons IncFII(pSE11) and IncN were detected in *M. morganii* MH16-367M and *P. mirabilis* MH13-009N, suggesting that these isolates might harbor bla_{NDM-1} on the identical IncFII(pSE11), IncN, or IncFII(pSE11)::IncN plasmid (Fig. 1).

MLST classified three *K. pneumoniae* isolates (MH15-289M, MH16-398D, and MH15-258M) as ST11 and another three *K. pneumoniae* isolates (MH15-208H, MH15-191M, and MH13-055M) as ST395. Interestingly, the presence of bla_{NDM-1} and replicons IncFIA(HI1), IncFIB(K), and IncFII(K) were detected in all six isolates (Fig. 1). In other examples, in the nine *K. pneumoniae*-ST15 isolates, bla_{NDM-4}, IncFIB(pKPHS1), and IncFII(K) replicons were detected in four isolates (MH17-556D, MH16-335M, MH17-016D, and MH17-017D), and bla_{KPC-2}, IncFIB(K), and IncN replicons were detected in five isolates (MH16-444M, MH17-553M, MH17-536D, MH16-519D, and MH17-011M) (Fig. 1).

Putative transfer of bla_{NDM-1}-carrying plasmids between Enterobacterales isolated in Vietnam

To investigate the similarities of the bla_{NDM-1}-carrying plasmids in more detail, long-read sequencing was performed for 12 bla_{NDM-1}-positive Enterobacterales isolates, including seven *K. pneumoniae* belonging to ST11 (three isolates), ST395 (three isolates), and ST656 (one isolate), two *C. freundii*, one *E. coli* belonging to ST405, one *M. morganii*,

and one *P. mirabilis* isolates, and one bla_{NDM-4}-positive *K. pneumoniae* isolate belonging

to ST15 were further sequenced with the long-read sequencer. Hybrid analysis using

short-read and long-read sequencing data resulted in complete sequences of 13 bla_{NDM}-

carrying plasmids that ranged from 39.2 kb to 345.7 kb sizes (covered by at least x125

long-read sequencing data; Table S2). Interestingly, many bla_{NDM}-carrying plasmids co-

carried other clinically relevant AMR genes, such as 16S rRNA methyltransferases genes

(9/13, 69.2%) and quinolone resistance genes (6/13, 46.2%) on the same plasmid (Table

S2).

BLAST search revealed that several bla_{NDM-1}-carrying plasmids had high sequence

identities with each other and with plasmids in the GenBank database (Figs. 2, 3, 4, and

5). However, bla_{NDM-1}-carrying plasmid pMH16-522D_1 (52.7 kb plasmid IncX3 replicon,

accession: AP018571) in *C. freundii* MH16-522D, bla_{NDM-1}-carrying plasmid pMH16-

390M_1 [345.7 kb plasmid with IncFIA(HI1)::IncR replicons, accession: AP018583] in *K.*

pneumoniae MH16-390M, and bla_{NDM-4}-carrying plasmid pMH16-335M_1 [105.6 kb

plasmid with IncFII(K) replicon, accession: AP018584] in *K. pneumoniae* MH16-335M

were not homologous to any other plasmids in this study (Table S2). BLAST search

revealed that numerous plasmids in Enterobacterales isolated around the world had 98-

100% identity in more than 70% of the regions of pMH16-522D_1, pMH16-390M_1, and

pMH16-335M_1, respectively. All three plasmids harbored additional β-lactamase

and/or *qnr* genes: *C. freundii* pMH16-522D_1 carried bla_{SHV-12}, *K. pneumoniae* pMH16-

390M_1 carried $bla_{CTX-M-3}$, bla_{TEM-1b}, and *qnrB6*, and *K. pneumoniae* pMH16-335M_1

carried bla_{OXA-9}, $bla_{SHV-106}$, and bla_{SHV-28} (Table S2). Furthermore, all three plasmids

contained the Tn*125*-like MGEs around bla_{NDM}, and the sequences were highly identical

to the reference sequence of *Tn*125 (accession: HQ857107) (96.9 to 100% identity with

87-89% region) [14].

P. mirabilis MH13-009N (isolated in 2013) and *M. morganii* MH16-367M (isolated in

2016) had only one plasmid, pMH13-009N_1 [72.6 kb plasmid with IncFII(pSE11)::IncN

replicons, accession: AP018566] and pMH16-367M_1 [73.1 kb plasmid with

IncFII(pSE11)::IncN replicons, accession: AP018565], respectively; the sequences of

these plasmids were almost identical (100% identity in 99% regions), including the

bla_{NDM-1} and IncFII(pSE11)::IncN hybrid replicon regions (Figs. 2A and 2B). Both of the

plasmids carried bla_{NDM-1} and an integron cassette that contains the class 1 integrase

gene (*intI1*) and the AMR genes, such as bla_{TEM-1b}, *rmtB*, and *qepA1*. The same integron

cassette was observed on *E. coli* pHPA (10.5 kb plasmid with IncFII replicon, accession:

AB263754) isolated from a human in Japan in 2002 [15]. Carbapenemase gene bla_{NDM-}

$_1$ was surrounded by several MGEs, including IS*Aba125*, IS*CR21*, and two IS*26*, on *P.*

mirabilis pMH13-009N_1 and *M. morganii* pMH16-367M_1, and shared identical

sequences with Tn*125* (Fig. 2A): 100% identity with 89% region of the reference

sequence (accession: HQ857107) [14,16,17]. The nearly identical bla_{NDM-1}-carrying

plasmids, *P. mirabilis* pMH13-009N_1 and *M. morganii* pMH16-367M_1 were

successfully transferred to recipient *E. coli* at a frequency of 10^{-2}–10^{-3} *in vitro*.

E. coli MH13-051M (isolated in 2013) harbored bla_{NDM-1} on plasmid pMH13-051M_1

[111.5 kb plasmid with IncFII(pRSB107)::IncN replicons, accession: AP018572], which

had partially identical sequences with two plasmids, pMH17-012N_1 (39.2 kb plasmid

with IncR replicon, accession: AP018567, 100% identity in 24% region of pMH13-

051M_1) and pMH17-012N_2 [87.6 kb plasmid with IncFII(pHN7A8)::IncN replicons,

accession: AP018568, 98.8% identity in 78% region of pMH13-051M_1], both found in

C. freundii MH17-012N (isolated in 2017) (Figs. 2C and 2D). *E. coli* pMH13-051M_1

[IncFII(pRSB107)::IncN plasmid] and *C. freundii* pMH17-012N_1 (smaller IncR plasmid)

commonly carried bla_{NDM-1} and bla_{TEM-1b}, and pMH13-051M_1 further carried $bla_{CTX-M-55}$

and *rmtB*. *C. freundii* pMH17-012N_2 [larger IncFII(pHN7A8)::IncN plasmid] carried no

known AMR genes, but did have a set of conjugation-associated type IV secretion

system (T4SS) genes, which shared identical sequences with *E. coli* pMH13-051M_1.

The genetic structures surrounding bla_{NDM-1} in *E. coli* pMH13-051M_1 and *C. freundii*

pMH17-012N_1 contained MGEs, including IS*Aba125*, IS*CR21*, and two IS*26*, and

shared identical sequences with Tn*125* (Fig. 2C): 98.6% and 100% identity with 89%

region of the reference sequence (accession: HQ857107) [14].

Comparison of bla_{NDM-1}-carrying plasmids in Vietnam with those in other countries

Next, we compared bla_{NDM-1}–carrying plasmids identified in Vietnam with those

reported in other countries. A set of conjugation-associated T4SS genes in *P. mirabilis*

pMH13-009N_1 [72.6 kb plasmid with IncFII(pSE11)::IncN replicons] was partially

identical with *Salmonella enterica* FDAARGOS_70 plasmid unnamed1 (47.8 kb plasmid

with IncN replicon, accession: CP026053, 100% identity in 61% region of pMH13-

009N_1) isolated from a human in the United States in 2013, which carried bla_{TEM-1b} (Fig.

3A). However, *S. enterica* FDAARGOS_70 plasmid unnamed1 did not carry bla_{NDM-1} as

P. mirabilis pMH13-009N_1 did. *C. freundii* pMH17-012N_2 [87.6 kb plasmid with

IncFII(pHN7A8)::IncN replicons and no AMR genes] was highly identical with *E. coli*

p103-2-4 [100.8 kb plasmid with IncFII(pHN7A8) replicons, accession: CP034846,

99.9% identity in 94% region of pMH17-012N_2] from a goose farm in China in 2018 and

with *S. enterica* serovar Enteritidis p12367A [111.8 kb plasmid with

IncFII(pHN7A8)::IncX1 replicons, accession: CP041177, 99.9% identity in 73% region of

pMH17-012N_2] isolated from a human in China in 2013 (Fig. 3B). *E. coli* p103-2-4 and

S. enterica serovar Enteritidis p12367A carried several AMR genes, such as $bla_{CTX-M-55}$

and bla_{TEM-1b}.

Furthermore, three *K. pneumoniae* isolates belonging to ST11 (MH15-289M, MH16-

398D, and MH15-258M isolated in 2015, 2016, and 2015, respectively) shared nearly

identical bla_{NDM-1}-carrying plasmids, pMH15-289M_1, pMH16-398D_1, and pMH15-

258M_1 [147–149 kb plasmids with IncFIA(HI1)::IncA/C2 replicons, accessions:

AP018577, AP018578, and AP018579, respectively, 99.6% identity in 99% regions], and

all of these plasmids carried other AMR genes, such as $bla_{CTX-M-15}$, bla_{OXA-1}, *rmtC*, and

qnrB9 (Fig. 4). bla_{NDM-1} was surrounded by several MGEs, including IS*3000*, IS*1*, and

IS*CR1*. These three plasmids were highly identical with *K. pneumoniae* plasmid

tig00000169_pilon [176.3 kb plasmid with IncFIA(HI1)::IncA/C2 replicons, accession:

CP021952, 99.8% identity in 82% region of pMH15-289M_1] in the United States and

with *E. coli* pK71-77-1-NDM (145.3 kb plasmid with IncA/C2 replicon, accession:

CP040884, 99.9% identity in 62% region of pMH15-289M_1) isolated from a human in

Norway in 2010. The plasmids tig00000169_pilon and pK71-77-1-NDM had identical

sequences in the T4SS region of pMH15-289M_1, and carried bla_{NDM-1}, bla_{CMY-6}, and

rmtC, and the plasmid tig00000169_pilon further carried bla_{SHV-11} and *qnrB58*. *K.*

pneumoniae MH16-398D had 235 sequence variants (214 single-nucleotide variants

[SNVs], 14 multiple-nucleotide variants [MNVs], four deletions, two insertions, and one

replacement), and *K. pneumoniae* MH15-258M had six sequence variants (five SNVs

and one replacement) relative to *K. pneumoniae* MH15-289M.

Another three *K. pneumoniae* isolates belonging to ST395 (MH15-208H, MH15-191M,

and MH13-055M isolated in 2015, 2015, and 2013, respectively) shared nearly identical

bla_{NDM-1}-carrying plasmids, pMH15-208H_1, pMH15-191M_1, and pMH13-055M_1 [75–

76 kb plasmids with IncFII(Yp) replicon, accessions: AP018580, AP018581, and

AP018582, respectively, 99.8–99.9% identity in 99% regions], and all plasmids carried

another AMR gene, $rmtB$ (Fig. 5). bla_{NDM-1} in these plasmids was surrounded by several

MGEs, including IS$Aba125$, IS1, and IS91, and shared partially identical sequences with

Tn125 (Fig. 5): 99.5-100% identity with 88-89% region of the reference sequence

(accession: HQ857107) [14]. These three plasmids were highly identical with

Enterobacter hormaechei pNDM1_045001 [85.7 kb plasmid with IncFII(Yp) replicon,

accession: CP043383, 99.9% identity in 98% region of pMH15-208H_1] isolated from a

human in China in 2018 and with *K. pneumoniae* pSECR18-2374C [86.0 kb plasmid with

IncFII(Yp) replicon, accession: CP041930, 99.9% identity in 95% region of pMH15-

208H_1] isolated from a human in South Korea in 2018. *E. hormaechei* pNDM1_045001

and *K. pneumoniae* pSECR18-2374C had a partially identical sequence of conjugation-

associated T4SS genes with pMH15-208H_1. *E. hormaechei* pNDM1_045001 carried

bla_{NDM-1}, bla_{TEM-1b}, and $rmtB$, and *K. pneumoniae* pSECR18-2374C carried bla_{NDM-4},

bla_{TEM-1b} and $rmtB$. *K. pneumoniae* MH15-191M had three sequence variants (two SNVs

and one MNV) and *K. pneumoniae* MH13-055M had only one SNV relative to *K.*

pneumoniae MH15-208H.

In bacterial conjugation experiments with *E. coli*, *C. freundii* pMH17-012N_1 (39.2 kb

plasmid with IncR replicon), pMH17-012N_2 [87.6 kb plasmid with IncFII(pHN7A8)::IncN

replicons], *E. coli* pMH13-051M_1 [111.5 kb plasmid with IncFII(pRSB107)::IncN

replicons], *K. pneumoniae* pMH15-258M_1 [148.8 kb plasmid with IncFIA(HI1)::IncA/C2

replicons], and *K. pneumoniae* pMH15-208H_1 [75.4 kb plasmid with IncFII(Yp) replicon]

were not transferred to recipient *E. coli* under our experimental conditions though a set

of T4SS-associated genes were detected in *C. freundii* pMH17-012N_2, *E. coli* pMH13-

051M_1, *K. pneumoniae* pMH15-258M_1, and *K. pneumoniae* pMH15-208H_1.

Discussion

The Viet Nam Resistance (VINARES), a nationwide surveillance network consisting of 16 central and provincial-level medical institutions was established in Vietnam in 2013 [18]. According to VINARES data, the rates of resistance of $K.$ $pneumoniae$ to third-generation cephalosporins, carbapenems, aminoglycosides, and fluoroquinolones were 66.4%, 17.1%, 29.5%, and 53.0%, respectively, in Vietnam [19], whereas the rates were 13.6%, 0.5%, 6.5%, and 8.7%, respectively, in the United Kingdom in 2013 [20], and 17.2%, 4.3%, 4.5%, and 16.8%, respectively, in the United States in 2010 [21]. Though the rates of resistance of Enterobacterales to major antimicrobials are much higher in Vietnam than in Western countries, the genetic causes remain unknown.

In this study, we performed comprehensive genetic analysis on plasmids carrying bla_{NDM} carbapenemase genes in Enterobacterales isolated from patients in Vietnam. We focused on carbapenem-nonsusceptible Enterobacterales isolates harboring bla_{NDM}, one of the most prevalent carbapenemase genes in the world, and completely sequenced 12 plasmids carrying bla_{NDM-1} from $K.$ $pneumoniae$ (ST11, ST395, and ST656), $E.$ $coli$ (ST405), $C.$ $freundii$, $M.$ $morganii$, and $P.$ $mirabilis$ isolates and one plasmid carrying bla_{NDM-4} from $K.$ $pneumoniae$ isolate (ST15). Nearly identical plasmids carrying bla_{NDM-1} were detected in different bacterial species of this study, suggesting that they represent common and important AMR plasmids disseminated in Vietnam (Figs. 2, 4, and 5). Moreover, the co-existence of clinically relevant AMR genes, such as ESBL genes (e.g., bla_{CTX-M}, bla_{OXA}, and bla_{TEM}), aminoglycoside resistance genes (e.g., $armA$ and rmt), and fluoroquinolone resistance genes (e.g., qep and qnr), with bla_{NDM-1} was observed in several plasmids, such as $P.$ $mirabilis$ pMH13-009N_1 (Fig. 2) and $K.$ $pneumoniae$ pMH15-289M_1 (Fig. 4). Hence, the spread of the plasmids both within and among

species of bacteria would have caused an increase in multidrug-resistant bacteria and

the infected hosts in Vietnam, and will pose a threat to public health.

Almost identical IncFII(pSE11)::IncN hybrid plasmids, pMH13-009N_1 and pMH16-

367M_1, were found in *P. mirabilis* and *M. morganii* isolates, respectively (Figs. 2A and

2B). These plasmids had identical sequences including a set of conjugation-associated

T4SS genes with the IncN plasmid from *S. enterica* isolated in the United States (Fig.

3A). IncN plasmids are prevalent in the microbiota of animals, and disseminate AMR

genes such as $bla_{CTX-M-1}$ and *qnr* [22], suggesting that IncN plasmids with the same origin

of *P. mirabilis* pMH13-009N_1 and *M. morganii* pMH16-367M_1 could be disseminated

in communities, including humans, animals, and the environment, in Vietnam and other

countries.

The IncFII(pRSB107)::IncN hybrid plasmid pMH13-051M_1 in *E. coli*, had partial

sequence identity with the IncR plasmid pMH17-012N_1 and the IncFII(pHN7A8)::IncN

hybrid plasmid pMH17-012N_2 both in *C. freundii* (Figs. 2C and 2D). Rearrangement of

conjugation genes plays an important role in the dissemination of AMR genes between

bacteria [23]. Because known IncR plasmids are non-transferable [24], an IncR plasmid

carrying AMR genes (e.g. *C. freundii* pMH17-012N_1) and another plasmid carrying

conjugation genes (e.g. *C. freundii* pMH17-012N_2) could be assumed to have been

fused into a single plasmid with both AMR and conjugation genes (e.g. *E. coli* pMH13-

051M_1) in a bacterium. Subsequently the plasmid was propagated by conjugation to

another bacterium. There is an increasing number of reports of IncR plasmids carrying

various AMR genes and the pool of AMR genes on IncR plasmids is thought to spread

to transmissible plasmids via recombination, contributing to the high evolutionary

plasticity of bacterial genomes [25]. Further analysis on the possible recombination

events are needed to understand the ways of transmission of these plasmids between

bacterial isolates and species.

The IncFII(pHN7A8)::IncN hybrid plasmid pMH17-012N_2 had highly identical

sequences including a set of conjugation-associated T4SS genes with the plasmids *E.*

coli p103-2-4 and *S. enterica* Enteritidis p12367A, that carry several AMR genes, such

as $bla_{CTX-M-55}$ and bla_{TEM-1b}, but no bla_{NDM} (Fig. 3B). The IncFII plasmid pHN7A8

(accession: JN232517), which also carries several AMR genes, such as $bla_{CTX-M-65}$, *rmtB*,

and *fosA3*, and is widespread in *E. coli* from farm and companion animals in China [26],

suggesting that plasmids with IncFII(pHN7A8) replicon are associated with AMR.

bla_{NDM-1} was found on several hybrid plasmids, such as IncFII(pSE11)::IncN,

IncFII(pRSB107)::IncN hybrid plasmids (Figs. 2A and 2C), and IncFIA(HI1)::IncA/C2

hybrid plasmids from three *K. pneumoniae* ST11 isolates (Fig. 4). Hybrids of multiple

replicons belonging to different incompatibility groups represent a plasmid strategy for

expansion of host range and dissemination of acquired AMR genes among bacteria [27].

The IncFIA(HI1)::IncA/C2 hybrid plasmids or IncFII(Yp) plasmids were reported from

other regions, including the United States, Europe, and Asia (Figs. 4 and 5).

IncFIA(HI1)::IncA/C2 plasmids and IncFII(Yp) plasmids were detected in *K. pneumoniae*

in this study, suggesting that these plasmids and/or bacterial isolates would have also

been spreading in medical institutions and community in Vietnam.

The bla_{NDM-1}-surrounding regions in plasmids, *P. mirabilis* pMH13-009N_1, *M. morganii*

pMH16-367M_1, *C. freundii* pMH17-012N_1, and *E. coli* pMH13-051M_1 (Fig. 2), *K.*

pneumoniae pMH15-208H_1, pMH15-191M_1, and pMH13-055M_1 (Fig. 5), and *K.*

pneumoniae pMH16-390M_1, pMH16-335M_1, and *C. freundii* pMH16-522D_1 had

almost identical sequence including Tn*125* which was discovered in *A. baumannii*

[10,14,16]. Tn*125* includes IS*Aba125* and IS*CR21*, but sequences of the Tn*125*-like

region in *P. mirabilis* pMH13-009N_1, *M. morganii* pMH16-367M_1, and *E. coli* pMH13-

051M_1 (Fig. 2) include two IS*26* in addition to IS*Aba125* and IS*CR21*. This Tn*125*-like

structure has also been detected in other plasmids from Enterobacterales, such as *C.*

freundii pCF104a-T3 (accession: MN657241) isolated from a human in Germany in 2015

[17]. This report showed that IS*26*-mediated transfer of bla_{NDM-1}-containing Tn*125*-like

regions had occurred in different Enterobacterales species within one medical institution.

The MGEs surrounding bla_{NDM-1} on plasmids, *K. pneumoniae* pMH15-289M_1,

pMH16-398D_1, and pMH15-258M_1 (Fig. 4), were highly identical with the

corresponding sequence of *E. coli* pEC2-NDM-3 (accession: KC999035) isolated from a

human in Australia in 2010 though pEC2-NDM-3 carried bla_{NDM-3} with a part of the same

set of MGEs, including IS*CR1*. IS*CR1* was suggested to provide a vehicle for bla_{NDM-1}

dissemination [8, 28].

MLST analysis is helpful to estimate whether bacterial isolates belonging to the same

STs were originated from the same clones and spread within the medical institution, and

whole-genome SNV-based analysis enables us to estimate more accurate genetic

relationship. According to a report [29], SNVs accumulated in *K. pneumoniae* ST258

isolates at a rate of 3.9 SNVs per year. Our analysis of SNVs of *K. pneumoniae* ST11

isolates (Fig. 4) revealed 214 SNVs in MH16-398D relative to MH15-289M. This

suggests that different ST11 clones are present in the medical institutions and

communities. In support of this theory, a report [30] showed that *K. pneumoniae* isolates

from medical institutions, sewage, canals, and agricultural waste water were intermixed

in the phylogenetic classification. Because ST11 is one of major international epidemic

clones of *K. pneumoniae* [31], an ST11 clone carrying AMR plasmid could have

disseminated in a medical institution or local community in Hanoi, Vietnam. Another three

ST395 isolates of *K. pneumoniae* (Fig. 5) are very clonal based on the rate of SNV

accumulation [29] and could be disseminated from the same clone within the medical

institution because only a few SNVs were present in MH15-191M and MH13-055M

relative to MH15-208H.

Conclusion

Plasmid analysis in this study shows the complex structures and diversity of plasmids

carrying $bla_{\text{NDM-1}}$ in Vietnam. Hybrid analysis with both Illumina short-read and ONT long-

read sequencing is a promising method for detecting important AMR plasmids in CRE

isolates and the real-time analysis is useful for controlling nosocomial infections.

Materials and Methods

Bacterial isolates

A total of 122 ESBL-producing Enterobacterales isolates were collected from selected specimens, including blood, sputum, urine, and pus from both in-patients and out-patients with infections in daily diagnosis in a reference medical institution in Hanoi, Vietnam between 2013 and 2017. All bacterial isolates were excluded from duplicates from the same patient. The medical institution has 700 beds, and includes 13 surgical departments and 10 internal medicine departments. The medical services are provided for both military and civil patients. Bacterial species identification, prediction of ESBL activity, and antimicrobial susceptibility testing were performed by Vitek 2 (BioMèrieux) and E-test (BioMèrieux). ESBL gene(s)-harboring isolates could also harbor other important AMR genes, including carbapenemase gene(s). According to the Clinical and Laboratory Standards Institute (CLSI) 2020 guidelines, breakpoints of meropenem and imipenem are ≤1 μg/mL (susceptible), 2 μg/mL (intermediate), ≥4 μg/mL (resistant), respectively. Carbapenemase activity was examined by CarbaNP test according to the CLSI 2020 guidelines. Major carbapenemase genes (bla_{NDM}, bla_{KPC}, bla_{OXA-48}, bla_{IMP}, and bla_{VIM}) were detected by a multiplex PCR method [32]. Twenty-five carbapenem-nonsusceptible and carbapenemase-positive isolates (19 *K. pneumoniae*, two *E. coli*, two *C. freundii*, one *M. morganii*, and one *P. mirabilis*) were subjected to the whole-genome sequencing analysis described below.

Preparation of genomic DNA

For short-read sequencing, genomic DNAs of bacterial isolates were extracted with the phenol-chloroform method, and purified with QIAquick PCR Purification Kit

(QIAGEN) according to the manufacturer's instructions. For long-read sequencing, high-

molecular-weight genomic DNAs of bacterial isolates were extracted with MagAttract

HMW DNA Kit (QIAGEN) according to the manufacturer's instructions. The extracted

genomic DNAs were quantified with a Qubit 2.0 fluorometer (Thermo Fisher Scientific).

Whole-genome sequencing and bioinformatics analysis

Whole-genome sequencing was performed to examine the prevalence of important

AMR genes on plasmids. Long-read sequencing is useful for constructing complete

sequences of whole plasmids and tracking horizontal transfer of AMR plasmids during

nosocomial infections [33]. Recently, long-read nanopore sequencing using MinION

nanopore sequencer from Oxford Nanopore Technologies (ONT) has been applied to

such investigations [34,35].

Whole genome sequencing using MiniSeq or HiSeq systems (Illumina) was performed

for phylogenetic and MLST analysis, and screening of acquired AMR genes and plasmid

replicon types. Library for Illumina sequencing (insert size of 500-900 bp) was prepared

using Nextera XT DNA Library Prep Kit (Illumina). Paired-end sequencing was performed

using MiniSeq (2 x 150 bp) or HiSeq 4000 systems (2 x 150 bp). Trimming and *de novo*

assembly of paired-end reads was performed using CLC Genomics Workbench v12.0

(QIAGEN) with default parameters. A maximum-likelihood phylogenetic tree was

generated by PhyML from a concatenated core-gene alignment consisting of 13,305

SNVs constructed using the Roary pipeline (https://sanger-pathogens.github.io/Roary/).

For PhyML, we used the following parameters that indicate the GTR +G4 model of DNA

substitution with estimation of the shape parameter of the gamma distribution by

maximizing the likelihood: -m GTR -c 4 -a e.

Twelve Enterobacterales isolates with bla_{NDM-1} (seven *K. pneumoniae*, one *E. coli*, two

C. freundii, one *M. morganii*, and one *P. mirabilis*) and one *K. pneumoniae* isolate with

bla_{NDM-4} were further sequenced on MinION (ONT) using the SQK-RAD002 or SQK-

RBK001 kits and R9.4 flowcells according to the manufacturer's instructions to obtain

complete sequences of plasmids carrying carbapenem resistance genes. *De novo*

assembly was performed using Canu v1.5 [36] and Miniasm [37] with default parameters.

The overlap region in the assembled contig was detected by genome-scale sequence

comparison using LAST (http://last.cbrc.jp) and was trimmed manually. Illumina paired-

end reads were mapped onto the resulted circular sequences, and error correction was

performed by extracting consensus of mapped reads using CLC Genomics Workbench

v12.0 with default parameters.

Sequence types, carbapenemase genes, and plasmid replicon types were analyzed

using MLST v2.0, ResFinder v3.2 with minimum threshold of 90% identity and 60%

coverage, and PlasmidFinder v2.1 with minimum threshold of 90% identity and 60%

coverage, respectively, on the Center for Genomic Epidemiology (CGE) server at

Technical University of Denmark (http://www.genomicepidemiology.org).

Coding sequences (CDS) annotation using RASTtk on the PATRIC v3.6.3 server

(https://www.patricbrc.org) with default parameters. Sequence variant detection was

performed using CLC Genomics Workbench v12.0 with default parameters of Fixed

Ploidy Variant Detection (Ploidy=1). Linear comparison of sequences was performed

using BLAST with the default settings (the nucleotide collection database and the

megablast program), and visualized by Easyfig (http://mjsull.github.io/Easyfig/). bla_{NDM-1},

other important AMR genes (ARG), type IV secretion system (T4SS)-associated genes

involved in conjugation that were detected by SecReT4 program [38], and mobile gene

elements (MGEs) were identified manually from CDS annotations and basically analyzed

by comparing the sequences analyzed in previous studies.

Draft genome and complete plasmid sequences of carbapenem-nonsusceptible

Enterobacterales isolated in Vietnam have been deposited at GenBank/EMBL/DDBJ

under BioProject number PRJDB6655.

Bacterial conjugation

Bacterial conjugation was performed using six bla_{NDM-1}-carrying plasmid-positive

Enterobacterales isolates [*P. mirabilis* MH13-009N, *M. morganii* MH16-367M, *C. freundii*

MH17-012N, *E. coli* MH13-051M (ST405), *K. pneumoniae* MH15-258M (ST11), and *K.*

pneumoniae MH15-208H (ST395)] according to the following protocol. The same amount

of Luria-Bertani (LB) broth cultures of each donor bacteria and the recipient azide–

resistant *E. coli* J53 (ATCC BAA-2731, *F- met pro Azi^r*), were mixed and spotted onto

Mueller-Hinton agar and then incubated at 37°C overnight. The mixed cells were

recovered and suspended into PBS buffer, plated onto LB agar after 10-fold serial dilution,

and incubated at 37°C overnight. Transconjugants were selected on LB agar containing

2 μg/mL of meropenem and 100 μg/mL of sodium azide.

Acknowledgements

We thank Meiko Takeshita for technical assistance and are grateful to the participating

medical institution for providing the bacterial isolates and clinical information.

462 **Conflicts of Interest**

463   None to declare.

 Figure Legends

 Figure 1

   Carbapenem-nonsusceptible Enterobacterales isolates sequenced on Illumina

 systems. Red nodes, but not black nodes, show isolates subsequently sequenced with

 ONT MinION. Bar lengths represent the number of substitutions per site in the core

 genome constructed using the Roary pipeline (see Materials and Methods). Sequence

 types of multilocus sequence typing (MLST) analysis determined from genomes,

 minimum inhibitory concentrations (MICs) of meropenem (MEM) and imipenem (IPM) of

 isolates, carbapenemase genes detected by ResFinder and replicon types detected by

 PlasmidFinder in genomes, years in which the bacteria were isolated, and replicon types

 and sizes of bla_{NDM}-carrying plasmids are shown.

 Figure 2

   bla_{NDM-1}-carrying plasmids and the detailed genetic structures around bla_{NDM-1} in

 carbapenem-nonsusceptible Enterobacterales isolates sequenced using ONT MinION

 and Illumina systems. Sets of plasmids from *P. mirabilis* MH13-009N (pMH13-009N_1,

 accession: AP018566) and *M. morganii* MH16-367M (pMH16-367M_1, accession:

 AP018565), and from *E. coli* MH13-051M (pMH13-051M_1, accession: AP018572) and

 C. freundii MH17-012N (pMH17-012N_1, accession: AP018567 and pMH17-012N_2,

 accession: AP018568), are shown. (A) and (C) Linear comparison of bla_{NDM-1}-carrying

 plasmid sequences. (B) and (D) Schematic representation of possible transfer of

 plasmids, including bla_{NDM-1}-carrying plasmids. Circles within bacterial cells show

 plasmids, and red squares on plasmids show bla_{NDM-1}. Sets of plasmids from *P. mirabilis*

 MH13-009N and *M. morganii* MH16-367M (A and B), and from *E. coli* MH13-051M and

C. freundii MH17-012N (C and D), are shown. Red, yellow, green, blue, and gray arrows

indicate *bla*_{NDM}, other important AMR genes (ARGs), type IV secretion system (T4SS)-

associated genes involved in conjugation, mobile gene elements (MGEs), and other

genes, respectively. The colors in comparison of plasmids show percent identity and

sequence direction as indicated. Blue for matches in the same direction and red for

matches in the inverted direction.

Figure 3

(A) and (B) Linear comparison of plasmid sequences from Vietnam and other countries.

P. mirabilis pMH13-009N_1 (this study, accession: AP018566) and *Salmonella enterica*

FDAARGOS_70 plasmid unnamed1 (accession: CP026053) (A), and *E. coli* pMH13-

051M_1 (this study, accession: AP018572), *C. freundii* pMH17-012N_2 (this study,

accession: AP018568), *E. coli* p103-2-4 (accession: CP034846), and *Salmonella*

enterica serovar Enteritidis p12367A (accession: CP041177) (B) are shown. Red, yellow,

green, blue, and gray arrows indicate *bla*_{NDM}, other important AMR genes (ARGs), type

IV secretion system (T4SS)-associated genes involved in conjugation, mobile gene

elements (MGEs), and other genes, respectively. The colors in comparison of plasmids

show percent identity and sequence direction as indicated. Blue for matches in the same

direction and red for matches in the inverted direction.

Figure 4

Linear comparison of *bla*_{NDM-1}-carrying plasmid sequences and the detailed genetic

structures around *bla*_{NDM-1} from Vietnam and other countries. *K. pneumoniae* pMH15-

289M_1 (this study, accession: AP018577), *K. pneumoniae* pMH16-398D_1 (this study,

accession: AP018578), *K. pneumoniae* pMH15-258M_1 (this study, accession:

AP018579), *K. pneumoniae* tig00000169_pilon (accession: CP021952), and *E. coli*

pK71-77-1-NDM (accession: CP040884) are shown. Red, yellow, green, blue, and gray

arrows indicate bla_{NDM}, other important AMR genes (ARGs), type IV secretion system

(T4SS)-associated genes involved in conjugation, mobile gene elements (MGEs), and

other genes, respectively. The colors in comparison of plasmids show percent identity

and sequence direction as indicated. Blue for matches in the same direction and red for

matches in the inverted direction.

Figure 5

Linear comparison of bla_{NDM}-carrying plasmid sequences and the detailed genetic

structures around bla_{NDM-1} from Vietnam and other countries. *K. pneumoniae* pMH15-

208H_1 (this study, accession: AP018580), *K. pneumoniae* pMH15-191M_1 (this study,

accession: AP018581), *K. pneumoniae* pMH13-055M_1 (this study, accession:

AP018582), *E. hormaechei* pNDM1_045001 (accession: CP043383), and *K.*

pneumoniae pSECR18-2374C (accession: CP041930) are shown. Red, yellow, green,

blue, and gray arrows indicate bla_{NDM}, other important AMR genes (ARGs), type IV

secretion system (T4SS)-associated genes involved in conjugation, mobile gene

elements (MGEs), and other genes, respectively. The colors in comparison of plasmids

show percent identity and sequence direction as indicated. Blue for matches in the same

direction and red for matches in the inverted direction.

Table S1

Bacterial species and isolates, specimen types for the bacterial isolation, years in

which the bacteria were isolated, and minimum inhibitory concentrations (MICs) of

meropenem (MEM) and imipenem (IPM) of isolates are shown. Also, sequence types of

multilocus sequence typing (MLST) analysis determined from genomes, carbapenemase

genes detected by ResFinder in genomes, sizes and contigs of genomes, coverages in

short-read sequencing, Illumina sequencing platforms, and accession numbers of

genomes are shown. According to the CLSI 2020 guidelines, Breakpoints of meropenem

and imipenem are as follows: ≤1 μg/mL, susceptible; 2 μg/mL, intermediate; ≥4 μg/mL,

resistant (R).

Table S2

Bacterial species and isolates, and bla_{NDM}-carrying plasmids are shown. Also, replicon

types detected by PlasmidFinder and antimicrobial resistance genes (ARGs) detected

by ResFinder in plasmids, sizes of plasmids, and coverages in long-read sequencing,

and accession numbers of plasmids are shown.

References

1. Martin A, Fahrbach K, Zhao Q, Lodise T. Association between carbapenem resistance and mortality among adult, hospitalized patients with serious infections due to enterobacteriaceae: Results of a systematic literature review and meta-analysis. Open Forum Infectious Diseases. Oxford University Press; 2018. p. ofy150.

2. Kohler PP, Volling C, Green K, Uleryk EM, Shah PS, McGeer A. Carbapenem Resistance, Initial Antibiotic Therapy, and Mortality in Klebsiella pneumoniae Bacteremia: A Systematic Review and Meta-Analysis. Infection Control and Hospital Epidemiology. Cambridge University Press; 2017. pp. 1319–1328.

3. Tzouvelekis LS, Markogiannakis A, Psichogiou M, Tassios PT, Daikos GL. Carbapenemases in Klebsiella pneumoniae and other Enterobacteriaceae: an evolving crisis of global dimensions. Clin Microbiol Rev. 2012;25: 682–707.

4. Tada T, Tsuchiya M, Shimada K, Nga TTT, Thu LTA, Phu TT, et al. Dissemination of Carbapenem-resistant Klebsiella pneumoniae clinical isolates with various combinations of Carbapenemases (KPC-2, NDM-1, NDM-4, and OXA-48) and 16S rRNA Methylases (RmtB and RmtC) in Vietnam. BMC Infect Dis. 2017;17: 467.

5. Isozumi R, Yoshimatsu K, Yamashiro T, Hasebe F, Nguyen BM, Ngo TC, et al. BlaNDM-1-positive Klebsiella pneumoniae from environment, Vietnam. Emerging Infectious Diseases. 2012. pp. 1383–1385.

6. Tran HH, Ehsani S, Shibayama K, Matsui M, Suzuki S, Nguyen MB, et al. Common isolation of New Delhi metallo-beta-lactamase 1-producing Enterobacteriaceae in a large surgical hospital in Vietnam. Eur J Clin Microbiol

Infect Dis. 2015;34: 1247–1254.

7.   Wu W, Feng Y, Tang G, Qiao F, McNally A, Zong Z. NDM metallo-β-lactamases

and their bacterial producers in health care settings. Clinical Microbiology

Reviews. American Society for Microbiology; 2019. pp. e00115-18.

8.   Wailan AM, Sidjabat HE, Yam WK, Alikhan NF, Petty NK, Sartor AL, et al.

Mechanisms involved in acquisition of blaNDM genes by IncA/C2 and IncFIIY

plasmids. Antimicrob Agents Chemother. 2016;60: 4082–4088.

9.   Nordmann P, Dortet L, Poirel L. Carbapenem resistance in Enterobacteriaceae:

Here is the storm! Trends in Molecular Medicine. Elsevier Current Trends; 2012.

pp. 263–272.

10.   Poirel L, Bonnin RA, Boulanger A, Schrenzel J, Kaase M, Nordmann P. Tn125-

related acquisition of blaNDM-like genes in Acinetobacter baumannii. Antimicrob

Agents Chemother. 2012;56: 1087–1089.

11.   Partridge SR, Iredell JR. Genetic Contexts of bla NDM-1. Antimicrobial Agents

and Chemotherapy. American Society for Microbiology Journals; 2012. pp.

6065–6067.

12.   Wailan AM, Paterson DL, Kennedy K, Ingram PR, Bursle E, Sidjabat HE.

Genomic characteristics of NDM-producing Enterobacteriaceae isolates in

Australia and their blaNDM genetic contexts. Antimicrob Agents Chemother.

2016;60: 136–141.

13.   Baraniak A, Machulska M, Literacka E, Izdebski R, Urbanowicz P, Bojarska K, et

al. Towards endemicity: large-scale expansion of the NDM-1-producing

Klebsiella pneumoniae ST11 lineage in Poland, 2015-16. J Antimicrob

Chemother. 2019;74: 3199–3204.

14.   Pfeifer Y, Wilharm G, Zander E, Wichelhaus TA, Göttig SG, Hunfeld KP, et al.

601   Molecular characterization of blaNDM-1 in an Acinetobacter baumannii strain

602   isolated in Germany in 2007. J Antimicrob Chemother. 2011;66: 1998–2001.

603   15.   Yamane K, Wachino JI, Suzuki S, Kimura K, Shibata N, Kato H, et al. New

604   plasmid-mediated fluoroquinolone efflux pump, QepA, found in an Escherichia

605   coli clinical isolate. Antimicrob Agents Chemother. 2007;51: 3354–3360.

606   16.   Bontron S, Nordmann P, Poirel L. Transposition of Tn125 encoding the NDM-1

607   carbapenemase in Acinetobacter baumannii. Antimicrob Agents Chemother.

608   2016;60: 7245–7251.

609   17.   Weber RE, Pietsch M, Frühauf A, Pfeifer Y, Martin M, Luft D, et al. IS26-

610   Mediated Transfer of blaNDM–1 as the Main Route of Resistance Transmission

611   During a Polyclonal, Multispecies Outbreak in a German Hospital. Front

612   Microbiol. 2019;10: 2817.

613   18.   Wertheim HFL, Chandna A, Vu PD, Pham C Van, Nguyen PDT, Lam YM, et al.

614   Providing impetus, tools, and guidance to strengthen national capacity for

615   antimicrobial stewardship in Viet Nam. PLoS Med. 2013;10: e1001429.

616   19.   Vu TVD, Do TTN, Rydell U, Nilsson LE, Olson L, Larsson M, et al. Antimicrobial

617   susceptibility testing and antibiotic consumption results from 16 hospitals in Viet

618   Nam: The VINARES project 2012–2013. J Glob Antimicrob Resist. 2019;18:

619   269–278.

620   20.   Ecdc. SURVEILLANCE REPORT. Antimicrobial resistance surveillance in

621   Europe 2013. 2013.

622   21.   Sanchez G V., Master RN, Clark RB, Fyyaz M, Duvvuri P, Ekta G, et al.

623   Klebsiella pneumoniae antimicrobial drug resistance, United States, 1998-2010.

624   Emerg Infect Dis. 2013;19: 133–136.

625 22. Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. Characterization of IncN

626   plasmids carrying blaCTX-M-1 and qnr genes in Escherichia coli and Salmonella

627   from animals, the environment and humans. J Antimicrob Chemother. 2013;68:

628   333–339.

629 23. He S, Chandler M, Varani AM, Hickman AB, Dekker JP, Dyda F. Mechanisms of

630   Evolution in High-Consequence Drug Resistance Plasmids. MBio. 2016;7:

631   e01987-16.

632 24. Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B,

633   Guerra B, et al. Plasmids carrying antimicrobial resistance genes in

634   Enterobacteriaceae. J Antimicrob Chemother. 2018;73: 1121–1137.

635 25. Compain F, Frangeul L, Drieux L, Verdet C, Brisse S, Arlet G, et al. Complete

636   Nucleotide Sequence of Two Multidrug-Resistant IncR Plasmids from Klebsiella

637   pneumoniae. Antimicrob Agents Chemother. 2014;58: 4207–4210.

638 26. He L, Partridge SR, Yang X, Hou J, Deng Y, Yao Q, et al. Complete nucleotide

639   sequence of pHN7A8, an F33:A-:B-type epidemic plasmid carrying blaCTX-M-

640   65, fosA3 and rmtB from China. J Antimicrob Chemother. 2013;68: 46–50.

641 27. Garza-Ramos U, Barrios H, Hernandez-Vargas MJ, Rojas-Moreno T, Reyna-

642   Flores F, Tinoco P, et al. Transfer of quinolone resistance gene qnrA1 to

643   Escherichia coli through a 50 kb conjugative plasmid resulting from the splitting

644   of a 300 kb plasmid. J Antimicrob Chemother. 2012;67: 1627–1634.

645 28. Li J, Lan R, Xiong Y, Ye C, Yuan M, Liu X, et al. Sequential isolation in a patient

646   of Raoultella planticola and Escherichia coli bearing a novel IS CR1 element

647   carrying blaNDM-1. PLoS One. 2014;9: 1–9.

29.   Bowers JR, Kitchel B, Driebe EM, MacCannell DR, Roe C, Lemmer D, et al.

Genomic Analysis of the Emergence and Rapid Global Dissemination of the

Clonal Group 258 Klebsiella pneumoniae Pandemic. Bengoechea JA, editor.

PLoS One. 2015;10: e0133727.

30.   Runcharoen C, Moradigaravand D, Blane B, Paksanont S, Thammachote J,

Anun S, et al. Whole genome sequencing reveals high-resolution

epidemiological links between clinical and environmental Klebsiella pneumoniae.

Genome Med. 2017;9: 6.

31.   Mathers AJ, Peirano G, Pitout JDD. The role of epidemic resistance plasmids

and international high-risk clones in the spread of multidrug-resistant

Enterobacteriaceae. Clin Microbiol Rev. 2015;28: 565–91.

32.   Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of

acquired carbapenemase genes. Diagn Microbiol Infect Dis. 2011;70: 119–123.

33.   Conlan S, Thomas PJ, Deming C, Park M, Lau AF, Dekker JP, et al. Single-

molecule sequencing to track plasmid diversity of hospital-associated

carbapenemase-producing Enterobacteriaceae. Sci Transl Med. 2014;6:

254ra126.

34.   Lemon JK, Khil PP, Frank KM, Dekker JP. Rapid Nanopore Sequencing of

Plasmids and Resistance Gene Detection in Clinical Isolates. Mellmann A,

editor. J Clin Microbiol. 2017;55: 3530–3543.

35.   Gorrie CL, Mirceta M, Wick RR, Judd LM, Wyres KL, Thomson NR, et al.

Antimicrobial-Resistant Klebsiella pneumoniae Carriage and Infection in

Specialized Geriatric Care Wards Linked to Acquisition in the Referring Hospital.

Clin Infect Dis. 2018;67: 161–170.

36.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu:

scalable and accurate long-read assembly via adaptive k-mer weighting and

repeat separation. Genome Res. 2017;27: 722–736.

37.   Li H. Minimap and miniasm: Fast mapping and de novo assembly for noisy long

sequences. Bioinformatics. 2016;32: 2103–2110.

38.   Bi D, Liu L, Tai C, Deng Z, Rajakumar K, Ou HY. SecReT4: A web-based

bacterial type IV secretion system resource. Nucleic Acids Res. 2013;41: D660–

D665.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Species	Isolate	Specimen	Year	MEM MIC	IPM MIC	MLST	Carbapenemase gene	Size	Contig	Coverage	Platform	Accession no.
M. morganii	MH16-367M	blood	2016	1.5	8 (R)	N/A	bla_{NDM-1}	4,132,245 bp	297	88.2x	MiniSeq	BFCJ00000000
P. mirabilis	MH13-009N	urine	2013	>32 (R)	>32 (R)	N/A	bla_{NDM-1}	4,116,783 bp	868	296.9x	MiniSeq	BFCK01000000
C. freundii	MH17-012N	urine	2017	2	3	N/A	bla_{NDM-1}	5,555,817 bp	156	235.1x	HiSeq 4000	BFCL01000000
C. freundii	MH16-522D	pus	2016	>32 (R)	12 (R)	N/A	bla_{NDM-1}	5,311,537 bp	182	119.4x	MiniSeq	BFCM01000000
E. coli	MH17-008M	blood	2017	4 (R)	6 (R)	457	bla_{KPC-2}	5,406,936 bp	606	129.6x	HiSeq 4000	BFCN01000000
E. coli	MH13-051M	blood	2013	6 (R)	3	405	bla_{NDM-1}	5,656,255 bp	259	107.9x	MiniSeq	BFCO00000000
K. pneumoniae	MH17-004D	pus	2017	4 (R)	>32 (R)	11	bla_{KPC-2}	5,694,352 bp	280	108.3x	HiSeq 4000	BFCP00000000
K. pneumoniae	MH15-289M	blood	2015	>32 (R)	>32 (R)	11	bla_{NDM-1}	5,771,413 bp	159	158.8x	HiSeq 4000	BFCQ00000000
K. pneumoniae	MH16-398D	pus	2016	8 (R)	2	11	bla_{NDM-1}	5,640,361 bp	263	87.3x	MiniSeq	BFCR00000000
K. pneumoniae	MH15-258M	blood	2015	6 (R)	2	11	bla_{NDM-1}	5,868,091 bp	377	444.1x	MiniSeq	BFCS00000000
K. pneumoniae	MH15-208H	sputum	2015	>32 (R)	>32 (R)	395	bla_{NDM-1}	5,685,569 bp	202	216.5x	HiSeq 4000	BFCT00000000
K. pneumoniae	MH15-191M	blood	2015	>32 (R)	>32 (R)	395	bla_{NDM-1}	5,787,568 bp	375	350.8x	MiniSeq	BFCU01000000
K. pneumoniae	MH13-055M	blood	2013	>32 (R)	>32 (R)	395	bla_{NDM-1}	5,718,215 bp	248	255.0x	HiSeq 4000	BFCV00000000
K. pneumoniae	MH17-010D	pus	2017	>32 (R)	>32 (R)	16	bla_{NDM-4}, $bla_{OXA-181}$	5,608,215 bp	160	261.8x	HiSeq 4000	BFCW01000000
K. pneumoniae	MH17-007D	pus	2017	2	3	656	bla_{KPC-2}	5,637,531 bp	149	131.1x	HiSeq 4000	BFCX00000000
K. pneumoniae	MH16-390M	blood	2016	8 (R)	8 (R)	656	bla_{NDM-1}	5,638,714 bp	219	97.1x	MiniSeq	BFCY00000000
K. pneumoniae	MH17-556D	pus	2017	>32 (R)	>32 (R)	15	bla_{NDM-4}	5,533,931 bp	132	126.3x	HiSeq 4000	BFCZ01000000
K. pneumoniae	MH16-335M	blood	2016	16 (R)	16 (R)	15	bla_{NDM-4}	5,840,408 bp	274	140.2x	MiniSeq	BFDA00000000
K. pneumoniae	MH17-016D	pus	2017	>32 (R)	8 (R)	15	bla_{NDM-4}, bla_{OXA-48}	5,871,712 bp	429	64.8x	MiniSeq	BFAI01000000
K. pneumoniae	MH17-017D	pus	2017	6 (R)	8 (R)	15	bla_{NDM-4}	5,847,449 bp	250	203.0x	HiSeq 4000	BFDB00000000
K. pneumoniae	MH16-444M	blood	2016	6 (R)	4 (R)	15	bla_{KPC-2}	5,640,233 bp	288	122.7x	MiniSeq	BFDC00000000
K. pneumoniae	MH17-553M	blood	2017	>32 (R)	>32 (R)	15	bla_{KPC-2}	5,673,369 bp	177	138.2x	HiSeq 4000	BFDD01000000
K. pneumoniae	MH17-536D	pus	2017	3	8 (R)	15	bla_{KPC-2}	5,652,240 bp	292	74.8x	MiniSeq	BFDE00000000
K. pneumoniae	MH16-519D	pus	2016	6 (R)	12 (R)	15	bla_{KPC-2}	5,682,307 bp	287	78.7x	MiniSeq	BFDF00000000
K. pneumoniae	MH17-011M	blood	2017	>32 (R)	>32 (R)	15	bla_{KPC-2}	5,652,498 bp	202	182.1x	HiSeq 4000	BFDG00000000

Table S1

Species	Isolate	Plasmid	Replicon type	ARG (β-lactam)	ARG (aminoglycoside)	ARG (quinolone)	ARG (others)	Size	Coverage	Accession no.
M. morganii	MH16-367M	pMH16-367M_1	IncFII(pSE11)::IncN	bla_{NDM-1}, bla_{TEM-1B}	*rmtB*	*qepA1*	*tet(A)*	73.1 kb	1860.7x	AP018565
P. mirabilis	MH13-009N	pMH13-009N_1	IncFII(pSE11)::IncN	bla_{NDM-1}, bla_{TEM-1B}	*rmtB*	*qepA1*	*tet(A)*	72.6 kb	758.8x	AP018566
C. freundii	MH17-012N	pMH17-012N_1	IncR	bla_{NDM-1}, bla_{TEM-1B}	N.D.	N.D.	N.D.	39.2 kb	930.6x	AP018567
C. freundii	MH17-012N	pMH17-012N_2	IncFII(pHN7A8)::IncN	N.D.	N.D.	N.D.	N.D.	87.6 kb	930.6x	AP018568
C. freundii	MH16-522D	pMH16-522D_1	IncX3	bla_{NDM-1}, bla_{SHV-12}	N.D.	N.D.	N.D.	52.7 kb	924.5x	AP018571
E. coli	MH13-051M	pMH13-051M_1	IncFII(pRSB107)::IncN	bla_{NDM-1}, $bla_{CTX-M-55}$, bla_{TEM-1B}	*rmtB*	N.D.	*fosA3*	111.5 kb	388.7x	AP018572
K. pneumoniae	MH15-289M	pMH15-289M_1	IncFIA(HI1)::IncA/C2	bla_{NDM-1}, $bla_{CTX-M-15}$, bla_{OXA-1}	*rmtC, aac(6')-Ib-cr, aac(3)-IIa, aac(3)-IId*	*qnrB9*	*catA2, catB3, sul1, dfrA14*	147.8 kb	125.0x	AP018577
K. pneumoniae	MH16-398D	pMH16-398D_1	IncFIA(HI1)::IncA/C2	bla_{NDM-1}, $bla_{CTX-M-15}$, bla_{OXA-1}	*rmtC, aac(6')-Ib-cr, aac(3)-IIa, aac(3)-IId*	*qnrB9*	*catA2, catB3, sul1, dfrA14*	147.7 kb	224.1x	AP018578
K. pneumoniae	MH15-258M	pMH15-258M_1	IncFIA(HI1)::IncA/C2	bla_{NDM-1}, $bla_{CTX-M-15}$, bla_{OXA-1}	*rmtC, aac(6')-Ib-cr, aac(3)-IIa, aac(3)-IId*	*qnrB9*	*catA2, catB3, sul1, dfrA14*	148.8 kb	584.9x	AP018579
K. pneumoniae	MH15-208H	pMH15-208H_1	IncFII(Yp)	bla_{NDM-1}	*rmtB*	N.D.	N.D.	75.4 kb	252.3x	AP018580
K. pneumoniae	MH15-191M	pMH15-191M_1	IncFII(Yp)	bla_{NDM-1}	*rmtB*	N.D.	N.D.	75.6 kb	788.4x	AP018581
K. pneumoniae	MH13-055M	pMH13-055M_1	IncFII(Yp)	bla_{NDM-1}	*rmtB*	N.D.	N.D.	75.3 kb	243.8x	AP018582
K. pneumoniae	MH16-390M	pMH16-390M_1	IncFIA(HI1)::IncR	bla_{NDM-1}, $bla_{CTX-M-3}$, bla_{TEM-1B}	*aac(6')-Ib-cr, aadA16*	*qnrB6*	*ARR-3, sul1, dfrA27*	345.7 kb	431.1x	AP018583
K. pneumoniae	MH16-335M	pMH16-335M_1	IncFII(K)	bla_{NDM-4}, bla_{OXA-9}, $bla_{SHV-106}$, bla_{SHV-28}	*aac(6')-Ib-cr, aadA1*	N.D.	N.D.	105.6 kb	180.6x	AP018584

Table S2