BK polyomavirus evades innate immune sensing by disrupting the mitochondrial network and membrane potential and promotes mitophagy

Julia Manzetti[1]†, Fabian H. Weissbach[1], Fabrice E. Graf[1], Gunhild Unterstab[1]‡, Marion Wernli[1], Helmut Hopfer[2], Cinthia B. Drachenberg[3], Christine Hanssen Rinaldo[4,5], Hans H. Hirsch[1,6,7,*]

Affiliations

[1] Transplantation & Clinical Virology, Department Biomedicine (Haus Petersplatz), University of Basel, Basel, Switzerland.

[2] Institute for Pathology, University Hospital Basel, Basel, Switzerland.

[3] Departments of Pathology, University of Maryland School of Medicine, Baltimore, MD, USA.

[4] Department of Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway.

[5] Metabolic and Renal Research Group, UiT The Arctic University of Norway, Tromsø, Norway.

[6] Clinical Virology, Laboratory Medicine, University Hospital Basel, Basel, Switzerland.

[7] Infectious Diseases & Hospital Epidemiology, University Hospital Basel, Basel, Switzerland.

†present address: Oncology, Laboratory Medicine, University Hospital Basel, Basel, Switzerland

‡present address: Immunobiology, Department Biomedicine, University of Basel, Basel, Switzerland

*Correspondence

Hans H. Hirsch, MD, MSc

Transplantation & Clinical Virology,

Department Biomedicine,

University of Basel, Petersplatz 10, CH-4009 Basel, Switzerland

Phone: +41 61 207 32 62

Fax: +41 61 207 3283

E-mail: hans.hirsch@unibas.ch

Running Title

BKPyV disrupts mitochondria and promotes mitophagy

Key words

Innate immunity, virus, polyoma, mitophagy, interferon

Authors emails

julia.manzetti@usb.ch, fabian.weissbach@unibas.ch, fabrice.graf@unibas.ch;

gunhild.unterstab@unibas.ch, m.wernli@unibas.ch, helmut.hopfer@usb.ch,

CDrachenberg@som.umaryland.edu, christine.rinaldo@unn.no, hans.hirsch@unibas.ch

Graphical Abstract

Highlights

- **BK Polyomavirus agnoprotein disrupts mitochondrial network and membrane potential**
- **Agnoprotein impairs nuclear IRF3-translocation, interferon-β expression**
- **Agnoprotein facilitates innate immune evasion during the late viral replication phase**
- **Damaged mitochondria are targeted for p62/SQMTS autophagy**

In Brief

BKPyV infects 90% of humans and asymptomatically persists in the kidneys, but causes premature allograft failure in kidney transplant patients. Manzetti and colleagues show that BKPyV-agnoprotein facilitates viral replication by impairing cytosolic innate immune sensing in late replication by disrupting the mitochondrial network and membrane potential, and promotes p62/SQSTM1 mitophagy.

Abstract

Immune escape contributes to viral persistence, yet little is known about human polyomaviruses. BK-polyomavirus (BKPyV) asymptomatically infects 90% of humans, but causes premature allograft failure in kidney transplant patients. Despite potent virus-specific T-cells and neutralizing antibodies, BKPyV persists in kidneys and regularly evades immune control as evidenced by urinary shedding in immunocompetent individuals. Here, we report that BKPyV disrupts the mitochondrial network and membrane potential when expressing the 66aa-long agnoprotein during late replication. Agnoprotein is necessary and sufficient, using its amino-terminal and central domain for mitochondrial targeting and network disruption, respectively. Agnoprotein impairs nuclear IRF3-translocation, interferon-β expression, and promotes p62-mitophagy. Agnoprotein-mutant viruses unable to disrupt mitochondria show reduced replication, which can be rescued by type-I-interferon-blockade, TBK1-inhibition, or CoCl2-treatment. Mitochondrial fragmentation and p62-autophagy occur in allograft biopsies of kidney transplant patients with BKPyV-nephropathy. JCPyV- and SV40-infection similarly disrupt mitochondrial networks indicating a conserved mechanism facilitating polyomavirus persistence relevant to post-transplant disease.

Introduction

Viruses infect all forms of life, and while metagenomics are revealing an increasing complexity of viromes (Hirsch, 2019; Simmonds et al., 2017; Virgin, 2014), the underlying principle remains the same: viral genetic information present as DNA or RNA is decoded by host cells, thereby enabling a programmed take-over of cell metabolism to accomplish the essential steps of viral genome replication, packaging, and progeny release to infect new susceptible cells and hosts (Lynn W. Enquist, 2013). With progeny rates of ten- to one hundred-thousand per cell, virus replication represents a severe burden compromising host cell function and viability, and ultimately organ and host integrity. Host defense mechanisms include antiviral restriction factors (Kluge et al., 2015) as well as innate and adaptive immune responses (McNab et al., 2015) intercepting early and late steps of the viral life cycle. Given the obligatory intracellular step of virus replication, the innate immune response is faced with the challenge of discriminating virus and its regulatory and structural units as "non-self" amidst physiological host cell constituents. This task is partly accomplished by identifying pathogen-associated molecular patterns (PAMPs) such as repetitive protein, lipid, and sugar structures through pattern recognition receptors (PRRs). In the last decade, cytoplasmic sensing of nucleic acids has emerged as a key mechanism of intracellular innate immune sensing (Takeuchi and Akira, 2010). Evolutionarily linked to detecting DNA damage and the failing integrity of nuclei or mitochondria resulting from metabolic stress, toxicity, or cancer (Hartlova et al., 2015; Li and Chen, 2018; West and Shadel, 2017), viral RNA and DNA were found to be similarly sensed through PRRs such as RIG-I/MDA-5/MAVS and cGAS/STING (Goubau et al., 2013; Hartlova et al., 2015; McFadden et al., 2017). MAVS and STING have been located on membranous platforms of mitochondria and the endoplasmic reticulum (ER) in the cytosol. Besides direct cross-talk (Zevini et al., 2017) and proximity via mitochondria-associated ER-membranes (MEM), both pathways converge in inducing type-1 interferon expression following the activation of the downstream kinases TBK-1 and IKK$_\varepsilon$, and the phosphorylation and nuclear translocation of transcription factors such as IRF3, IRF7, and NFkB (Liu et al., 2015; McFadden et al., 2017). Interferons are key

mediators of the antiviral state in infected and neighbouring cells and help to activate the adaptive antigen-specific immune responses (Iwasaki and Medzhitov, 2010; Schneider et al., 2014). Different molecular mechanisms of innate immune activation are induced in acute and chronic infections with human RNA or DNA viruses, and a variety of strategies has been described which may permit transient or persistent viral evasion (Garcia-Sastre, 2017). However, such aspects are incompletely understood for small non-enveloped DNA viruses such as human polyomaviruses.

BK polyomavirus (BKPyV) is one of at least 10 human polyomaviruses and infects >90% of the general population typically during childhood without specific illness (Greenlee and Hirsch, 2017; James A. DeCaprio, 2013). Although BKPyV induces potent virus-specific CD4 and CD8 T-cells (Binggeli et al., 2007; Chen et al., 2008; Cioni et al., 2016; Leboeuf et al., 2017) and neutralizing antibodies (Kaur et al., 2019; Pastrana et al., 2012; Shah et al., 1980; Solis et al., 2018), the virus latently persists in the kidneys and regularly escapes from immune control as evidenced by asymptomatic urinary virus shedding in immunocompetent healthy individuals (Egli et al., 2009; Imperiale and Jiang, 2016). In immunosuppressed patients, BKPyV-replication increases in rate and magnitude, progressing to nephropathy and haemorrhagic cystitis in 1%-15% and 5%-25% of kidney transplant and allogeneic bone marrow transplant recipients, respectively (Cesaro et al., 2018; Hirsch et al., 2019). Since specific antiviral agents and vaccines are not available, reducing immunosuppression is the current mainstay of therapy in order to regain control over BKPyV-replication (Cesaro et al., 2018; Hirsch et al., 2019). However, this manoeuvre increases the risk of immunological injury such as allograft rejection and graft-versus-host disease. In our ongoing study to identify functionally and diagnostically relevant targets of BKPyV-specific antibody and T-cell responses (Binggeli et al., 2007; Cioni et al., 2016; Leboeuf et al., 2017), we noted that the small BKPyV-agnoprotein of 66 amino acids (aa) is abundantly located in the cytoplasm during the late viral life cycle *in vivo*, but largely ignored by the adaptive immunity (Leuenberger et al., 2007; Rinaldo et al., 1998). BKPyV-agnoprotein co-localizes with lipid droplets (LD) (Unterstab et al., 2010) and membranous structures of the ER (Unterstab et

al., 2013). We now report that the BKPyV-agnoprotein also targets mitochondria subverting interferon-β induction by disrupting the mitochondrial network and its membrane potential, and promotes p62-mitophagy in cell culture and in kidney allograft biopsies.

Results

BKPyV-Agnoprotein colocalizes with mitochondria and induces mitochondrial fragmentation. To investigate the function of BKPyV agnoprotein in the absence of LD, we noted that the N-terminal amino acid (aa) sequence predicted similarity with mitochondrial targeting sequences (MTS) like the cytochrome c oxidase cox8 (**Fig. S1**). To address the potential mitochondrial localisation, we infected primary human renal proximal tubular epithelial cells (RPTECs) with the agnoprotein wildtype BKPyV-Dunlop (Dun-*AGN*), a well-characterized model of renal allograft nephropathy (Bernhoff et al., 2008; Hirsch et al., 2016; Low et al., 2004). The results were compared with the isogenic derivative Dun-*agn25D39E*, of which the encoded point mutant agnoprotein is known to no longer bind to LD after replacing the hydrophobic A25 and F39 with D and E, respectively, which abrogate the amphipathic character of the central domain without affecting the α-helix prediction (Unterstab et al., 2010).

At 48 h post-infection (hpi) with wildtype Dun-*AGN* and mutant Dun-*agn25D39E* viruses, immunofluorescent staining detected both the early viral protein large T-antigen (LTag) and the late viral protein Vp1 capsid in the nucleus and agnoprotein in the cytoplasm (**Fig. S2**). Using the mitochondrial outer membrane protein Tom20 as a marker, specific colocalization of both the *AGN*-wildtype and *agn25D39E*-mutant protein was found, demonstrating agnoprotein colocalization to mitochondria (**Fig. 1A**). Of note, the mitochondria of Dun-*AGN*-replicating RPTECs had lost the network-like pattern typical of uninfected cells and appeared in short, fragmented units (**Fig. 1A, supplementary Movie1**). In contrast, Dun-*agn25D39E*-replicating RPTECs exhibited a regular mitochondrial network similar to neighbouring uninfected cells (**Fig. 1A; supplementary Movie2**). Quantification of the mitochondrial morphology indicated a large excess of short mitochondrial fragments in Dun-*AGN*-infected cells, whereas mostly elongated mitochondria in a network-like pattern were seen in the Dun-*agn25D39E*-infected cells (**Fig. 1B**). Given these striking differences in mitochondrial phenotype, we examined whether or not the mutant *agn25D39E*- agnoprotein

was still able to target the ER as reported for the wildtype agnoprotein (Unterstab et al., 2013). Confocal microscopy revealed that the *agn25D39E*-protein colocalized with the ER-marker calreticulin (**Fig. 1C**). However, whereas the mitochondrial colocalization of the *agn25D39E*-protein appeared in network strings, the ER-colocalization with calreticulin was patchy and reminiscent of the contact sites with the mitochondria-associated membranes (MEMs) (**Fig. 1C**). The patchy ER-colocalization pattern was independently confirmed using protein disulphide isomerase (PDI), another ER-marker protein (**Fig. S2C**). The results indicated that targeting of ER and mitochondria remained intact and implicated the amphipathic character of the central α-helix in the disruption of the mitochondrial network. To correlate the severely altered mitochondrial morphology with the viral life cycle, we examined a time course of Dun-*AGN* infection demonstrating that expression of the early viral LTag at 24 h post-infection (hpi) had no effect on the mitochondrial network (**Fig. 1D**). After 36 hpi, agnoprotein expression started to appear in the cytoplasm, but mitochondrial fragmentation and perinuclear condensation became apparent only from 48 hpi onwards. At 72 hpi, Dun-*AGN*-replicating cells could be readily identified by the dramatically fragmented mitochondrial network using Tom20 staining alone (**Fig. 1D**). Thus, wildtype and *agn25D39E*-agnoprotein were both able to localize to mitochondria and to the ER, which in case of the wildtype led to mitochondrial fragmentation, while this was not observed for the Dun-*agn25D39E* mutant.

To further characterize the role of agnoprotein and its amphipathic helix in BKPyV replication, two additional isogenic derivatives were generated: Dun-*agn25L39L* encoding a mutant agnoprotein retaining the amphipathic character of the central α-helix, and Dun-*ATCagn*, in which the *ATG* start codon had been changed to *ATC* to create a "null"-agnoprotein virus. Ribbon models predicted that the overall secondary structures of *agn25D39E* and *agn25L39L* were similar to the *AGN*-encoded wildtype protein in having a short N-terminal and a central helix, whereas the C-terminal structure was not predictable

except for a short terminal α-helical tail (**Fig. 2A**). All BKPyV variants were found to proceed to the late viral replication phase, as evidenced by nuclear Vp1 staining (**Fig. 2B**). Similar to Dun-*AGN* wildtype infection, the Dun-*agn25L39L*-infected RPTECs exhibited a cytoplasmic agnoprotein distribution and mitochondrial fragmentation (**Fig. 2B**). In contrast, Dun-*ATCagn*-infected RPTECs retained intact mitochondrial networks similar to Dun-*agn25D39E,* but lacked agnoprotein expression as expected (**Fig. 2B**). Since LD-binding had been shown to require the amphipathic character of the central α-helix (**Fig. 2A**) now implicated in mediating mitochondrial fragmentation (Unterstab et al., 2010), the effect of LD-formation on the mitochondrial network was investigated. To this end, RPTECs were infected with Dun-*AGN* or with Dun-*agn25D39E* and 300μM oleate was added at 24 hpi. Confocal microscopy of Dun-*AGN* at 48hpi revealed that agnoprotein was sequestered around LD as described previously, and that mitochondrial fragmentation of Dun-*AGN* infected cells was abrogated (**Fig. 2C**). In contrast, Dun-*agn25D39E*-infected RPTECs retained the mitochondrial colocalization of the mutant agnoprotein without sequestering to LD (**Fig. 2C, magnification panel bottom right**) which in the absence of LD-staining are known to appear as punched-out holes (Unterstab et al., 2010). These results linked the amphipathic helix of agnoprotein to LD-binding and to mitochondrial fragmentation in a competitive manner. To investigate functional consequences of agnoprotein expression, we examined the mitochondrial membrane potential (MMP) using the ($Ψ_m$)-dependent accumulation of JC-1 dye, whereby mitochondrial depolarization is indicated by a decrease in the red/green fluorescence intensity. At 48 hpi, Dun-*AGN*-infected cells exhibited fragmented mitochondria (green channel) and a significant decrease in MMP (red channel), whereas MMP changed little in Dun-*agn25D39E*- or Dun-*ATCagn*-infected RPTECs, or in mock-treated cells (**Fig. 2D**). Similarly, automated measurements of overall JC-1 red and green signals in cell culture revealed significant MMP decreases of about 40% in Dun-*AGN*-infected RPTECs at 48 hpi compared to controls. Together, the results indicated that

infection with BKPyV expressing an agnoprotein with a central amphipathic helix was necessary for MMP-breakdown and network fragmentation in the late viral replication phase.

Agnoprotein is sufficient for mitochondrial fragmentation and breakdown of the mitochondrial membrane potential and confers impaired innate immune signalling. To investigate whether or not agnoprotein expression alone is sufficient for mitochondrial fragmentation, expression vectors containing the full-length gene of wildtype agnoprotein or agnoprotein subdomains fused to monomeric enhanced green fluorescent protein (mEGFP) (Unterstab et al., 2010) were transfected into UTA6 cells (**Fig. 3A**). At 24 hours post-transfection (hpt) of the agno(1-66)mEGFP construct, GFP and agnoprotein overlapped in colocalization to fragmented mitochondria (**Fig. 3A, top row**). Similarly, mitochondrial colocalization and fragmentation was seen for agno(1-53)mEGFP lacking the C-terminal 13 aa of agnoprotein, but not for any of the other truncated agnoprotein constructs shown. Since the truncated agno(20-66)mEGFP had been previously demonstrated to be able to still colocalize with LD (Unterstab et al., 2010), we hypothesized that the N-terminal domain was necessary for mitochondrial targeting in order to mediate the breakdown of the mitochondrial network. Therefore, the N-terminal mitochondrial targeting sequence of cytochrome c oxidase (MTScox8) was fused yielding MTScox8-Agno(20-66)mEGFP lacking its natural N-terminus (**Fig. S1**). The results showed that agno(20-66) was sufficient to induce mitochondrial fragmentation if mitochondrial targeting was provided by the MTScox8 sequence (**Fig. 3A, bottom row and extra panels; supplementary movie3**). This was not observed for MTScox8-mEGFP targeting mitochondria, but lacking agnoprotein sequences (**Fig. S3A**). To the ability of binding to LD upon addition of oleate, MTScox8-mEGFP and MTScox8-Agno(20-66)mEGFP were transfected and stained for LD using 4,4′-Difluoro-4-bora-(3a,4a)-diaza-s-indacene (bodipy) and Tom70, and analysed by confocal microscopy (**Fig. S3B and C**). The results showed that MTScox8-Agno(20-66)mEGFP were co-localizing to LD while MTScox8-mEGFP was not. The data indicated that the amphipathic helix present in the aa20 – aa66 of truncated MTScox8-Agno(20-66)mEGFP fusion protein

was available for interaction with LD on the surface of mitochondria. As an independent approach, we examined UTA6-2C9 cells harbouring a wildtype full-length agnoprotein under the control of an inducible tetracycline (tet)-off promoter (Cioni et al., 2013). At 24 h in the absence of tet (-tet), agnoprotein was expressed in the cytoplasm and colocalized with fragmented mitochondria, whereas in the presence of tet concentrations suppressing agnoprotein expression (+tet), an intact mitochondrial network was seen (**Fig. 3B**). Importantly, inducing agnoprotein was associated with MMP disruption when examining aggregate (red) and monomeric (green) signals or when using automated JC-1 dye measurements of the cell cultures (**Fig. 3C**). Since mitochondria have a key role in innate immunity (Koshiba et al., 2011), UTA6-2C9 cells were cultured for 24 h in the presence or absence of agnoprotein expression, and transfected with either poly(I:C)RNA or poly(dA:dT)DNA, both of which are potent inducers of the type-1-interferon expression via cytoplasmic phosphorylation and nuclear translocation of the interferon regulatory factor 3 (IRF3). At 4 hpt, the effect of agnoprotein expression on the nuclear localization of the IRF3 was examined. The results showed that in the presence of agnoprotein, nuclear IRF3 translocation was significantly reduced after transfecting poly(I:C) (**Fig. 3D**) or poly(dA:dT) (**Fig. 3E**). To functionally relate the agnoprotein-dependent differences in nuclear IRF-3 translocation, interferon (IFN)-β expression was quantified after transfection of increasing amounts of poly(dA:dT). The results indicated that agnoprotein expression resulted in a significant reduction of IFN-β transcripts and secreted protein levels (**Fig. 3F**). Together, the data indicated that the expression of BKPyV-agnoprotein was necessary and sufficient to induce mitochondrial fragmentation, breakdown of MMP, and impairment of the innate immune sensing of transfected cytosolic DNA and RNA.

BKPyV replication is inhibited by IFN-β. To determine whether or not BKPyV replication is sensitive to IFN-β, RPTECs were pre-treated, which resulted in an IFN-β dose-dependent reduction of LTag-positive cells and supernatant BKPyV loads at 72 hpi (**Fig. 4A**).

SDS/PAGE Immunoblot analysis demonstrated an increase of IFN-induced protein with tetratricopeptide repeats (IFIT) and a reduction of the viral capsid protein Vp1 and agnoprotein (**Fig. 4B**). A time course of IFN-β addition revealed maximal inhibitory effects before or at 2 hpi, but addition at 36 hpi still reduced the supernatant BKPyV loads by 50% (**Fig. 4B**). The reduction in supernatant BKPyV loads could be restored by type-I interferon blockade consisting of a cocktail of blocking antibodies against IFN-α, IFN-β and IFN-alpha/beta receptor (**Fig. 4B**). Thus, BKPyV replication in primary human RPTECs was inhibited by IFN-β, but could be prevented by anti-IFN-blockade. To examine the role of agnoprotein in BKPyV replication, RPTECs were infected with equivalent infectious doses of Dun-*AGN* or the mutant derivatives Dun-*agn25D39E* and Dun-*ATCagn*. Although the mutant variants replicated, both had lower supernatant BKPyV loads at 72 hpi and a reduced number of LTag- and Vp1-positive cells compared to wildtype virus (**Fig. 4C**). To investigate whether or not the reduced replication of Dun-*agn25D39E* involved innate immune activation*,* the TBK-1-inhibitor (Bx795) to prevent IRF3-phosphorylation, or the type-1 interferon-blocking cocktail was added at 36 hpi. The results demonstrated that TBK-1 inhibition or type-1 interferon blockade were able to partially rescue BKPyV Dun-*agn25D39E* replication (**Fig. 4C**). Under these conditions, there was not effect of TBK-1 inhibition on BKPyV Dun-*AGN* replication (**Fig. S4**). Since $CoCl_2$-treatment has been described to induce functional hypoxia disrupting the MMP (Jung and Kim, 2004), RPTECs were infected with Dun-*agn25D39E* and treated with 150 μM or 300 μM $CoCl_2$ at 24 hpi. Partial mitochondrial fragmentation of BKPyV Dun-*agn25D39E* infected cells was seen together with increased supernatant viral loads at 72 hpi (**Fig. 4D**). Together, the results indicated that the failure of the *agn25D39E* mutant protein to disrupt the MMP and the mitochondrial network was associated with reduced replication, which could be partially rescued by interfering with innate immune activation, type-1-interferon expression and MMP-dependent mitochondrial signalling relays.

Since MAVS is an important sensor of cytosolic nucleic acids and located on mitochondria, we examined its distribution in Dun-*AGN*- and mock-infected RPTECs by confocal microscopy. As shown, MAVS colocalized with both, intact and agnoprotein-fragmented mitochondria in RPTECs (**Fig. 5A**). Similarly, MAVS also colocalized with fragmented mitochondria in UTA6-2C9 cells following tet-off induced agnoprotein expression (data not shown). To compare the levels of MAVS in Dun-*AGN*-infected and mock-treated RPTECs, cell extracts were analysed by SDS/PAGE and immunoblotting showing reduced MAVS levels following BKPyV Dun-*AGN* infection (**Fig. 5A, top row**). Sequential immunoblotting for GAPDH as loading control and for Tom20 revealed a 60%-70% decrease in intensity of the major MAVS band and Tom20 in Dun-*AGN*-infected RPTECs compared to non-infected control cells (**Fig. 5A, bottom panels**).

Fragmentation of the mitochondrial network with intact MMP occurs physiologically prior to mitosis and mitochondria partitioning into daughter cells, and involves phosphorylation of the dynamin-related protein (Drp)1 at S616 by the cell division kinase CDK1/cyclin B (**Fig. 5A, mock**). In BKPyV Dun-*AGN*-infected RPTECs, increased Drp1-S616 phosphorylation was detected Vp1-positive cells, but unrelated to mitosis (**Fig. 5A, *AGN***). By SDS/PAGE and immunoblotting, overall Drp1 levels were similar (**Fig. S5A**), but Drp1-S616 phosphorylation was increased in Dun-*AGN*-infected cells compared to mock (**Fig. 5A, right panel**). Comparing isogenic mutant viruses by confocal microscopy revealed that Drp1-S616 phosphorylation was increased in Dun-*agn25D39E* and Dun-*ATCagn*-replicating RPTECs, in which the mitochondrial network remained intact (**Fig. S5B**). The results indicated that increased Drp1-S616 phosphorylation was related to BKPyV infection occurring independent of cell division or expression or functionality of agnoprotein. However, agnoprotein-mediated fragmentation of the mitochondrial network was associated with declining protein levels of mitochondrial MAVS and Tom20.

Agnoprotein-disrupted mitochondria are targeted for p62/SQSTM1-mitophagy. Given the striking differences in mitochondrial structure by confocal microscopy, we applied by

transmission electron microscopy to examine BKPyV-Dun-*AGN* or Dun-*agn25D39E*-infected RPTECs. At 72 hpi, viral particles in the nucleus served as a marker of the late viral life cycle equivalent to nuclear Vp1 staining in confocal microscopy. In the cytoplasm of Dun-*AGN*-infected RPTECs, small disrupted mitochondrial vesicles were seen, matching the results obtained by confocal microscopy, as well as several smaller dense multilaminar structures. In contrast, longer filamentous mitochondrial structures were seen in BKPyV Dun-*agn25D39E* (**Fig. 6A**) in line with tangential cuts of the intact three-dimensional mitochondrial network seen by confocal microscopy (**Fig. S6**). Since damaged mitochondria with MMP breakdown are known to be targeted for autophagy, we investigated p62/SQSTM1 as a marker of mitochondrial autophagosomes (Johansen and Lamark, 2011). At 72 hpi, a significant increase in large confluent p62/SQSTM1-positive signals was observed in Dun-*AGN* infected RPTECs compared to the disperse cytoplasmic distribution p62/SQSTM1 in mutant Dun-*agn25D39E*-infected cells (**Fig. 6A**). Moreover, cytoplasmic aggregates of p62/SQSTM1 colocalizing with Tom20 were prominent in Dun-*AGN*-infected RPTECs with mitochondrial fragmentation, but rare in Dun-*agn25D39E*-infected cells with intact mitochondrial networks (**Fig. S6**). Similarly, agnoprotein-dependent p62/SQSTM1-aggregates were seen in UTA6-2C9 cells at 24 h post-induction, which condensed at 48 h indicating that agnoprotein-induced breakdown of the mitochondrial membrane potential and network was followed by p62/SQSTM1-positive autophagosome formation (**Fig. 6B**).

To investigate whether or not similar changes could be observed in BKPyV-associated nephropathy in kidney transplant patients, allograft biopsy samples were analysed. Indeed, transmission electron microscopy revealed small disrupted mitochondria in the cytoplasm of tubular epithelial cells having viral particles in the nuclei, whereas noninfected tubular epithelial cells showed prominent long mitochondria (**Fig. 6C**). Immunofluorescent staining of kidney allograft biopsies and confocal microscopy for agnoprotein, Tom20 and p62/SQSTM1 revealed fragmented mitochondria and p62/SQSTM1-aggregates in agnoprotein-positive cells in BKPyV-infected parts, which was not seen in non-infected tubular epithelial cells of the same biopsy core (**Fig. 6C**). Together, the data extended the

cell culture results to renal allograft nephropathy showing that BKPyV replication was associated with mitochondrial fragmentation and p62/SQSTM1-positive autophagosome formation in kidney transplant patients.

To investigate autophagic flux as a dynamic marker of p62/SQSTM1-mitophagy, the expression levels of LC3-I and its activated lipid-derivative LC3-II were examined in UTA6-2C9 cells by immunoblotting. In untreated cells, mostly LC3-I was detected, but in the presence of the lysosomal protease inhibitors pepstatin-A1/E64d, LC3-I increased and LC3-II became apparent as expected for inhibition of the steady-state autophagic flux (**Fig. 7A**). Following agnoprotein expression, LC3-I and the derivative LC3-II levels were also increased, and increased further in the presence of pepstatin-A1/E64d indicating that agnoprotein expression increased the autophagic flux, which could be partly blocked by lysosomal protease inhibitors (**Fig. 7A**). Treatment with carbonyl cyanide m-chlorophenylhydrazone (CCCP) is known to chemically disrupt the MMP and to induce PINK-Parkin-dependent mitophagy (Narendra et al., 2008). Indeed, CCCP treatment caused an increase in LC3-I and -II, which further increased in the presence of pepstatin-A1/E64d (**Fig. 7A**). However, agnoprotein expression followed by CCCP treatment did not result in a further increase of LC3-I/II levels, but rather increased the relative LC3-II proportion suggesting that the autophagic flux induced by agnoprotein was further maximized by CCCP treatment. To investigate the role of parkin in this process, UTA6-2C9 cells were transfected with a yellow fluorescent protein (YFP)-Parkin expression construct and analysed by confocal microscopy. In YFP-transfected UTA6-2C9 cells not expressing agnoprotein (+tet), little co-localization with fragmented mitochondria was observed. Following the addition of CCCP, extensive cytoplasmic YFP-parkin positive aggregates overlaid Tom20-positive structures in the cytoplasm (**Fig. S7, +tet, CCCP, magnification**). In UTA6-2C9 cells expressing agnoprotein (-tet), the YFP-Parkin signals appeared displaced by agnoprotein from Tom20-colocalizing structures accumulating in the perinuclear cytoplasm (**Fig. S7, tet-, magnification**). Using isosurface rendering to analyze the agnoprotein-induced

mitochondrial fragmentation supported the notion that Tom20-positive mitochondrial structures were surrounded agnoprotein-positive layer, which displaced another layer of YFP-parking (**Fig. S7, tet-, bottom panels, dashed circle**). Together, the data suggested that despite a shared breakdown of the MMP, CCCP- and agnoprotein-induced mitophagy differed in associating directly and indirectly with Parkin, respectively.

To independently investigate the role of p62/SQSTM1 in autophagic flux following BKPyV infection of RPTECs, LC3 was analysed in RPTEC with or without siRNA-p62 knockdown. Immunoblotting of the siRNA-p62 knockdown RPTECs revealed a reduction of p62/SQSTM1 levels to approximately 30 percent of the control RPTECs (**Fig. 7B, left panel**). Upon pepstatin-A1/E64d treatment, p62/SQSTM1 levels as well as the levels of LC3-I and -II increased in the control cells, but not to the same extent in the siRNA-p62 knockdown cells, indicating that the decrease in p62/SQSTM1 levels was associated with lower LC3-II formation and reduced steady-state autophagic flux (**Fig. 7B, left panel**). Following Dun-*AGN*-infection and pepstatin-A1/E64 treatment, LC3-I and -II levels increased, but the overall levels were lower in the siRNA-p62 knockdown cells, and the LC3-II/-I ratio was decreased as well as the Vp1 levels (**Fig. 7B, left panel**). To investigate the impact of the siRNA-p62 knockdown, whole cell extracts were prepared and analysed by SDS/PAGE-immunoblotting for MAVS levels in infected and uninfected RPTECs. The results after normalization to actin revealed that MAVS levels were not affected by siRNA-p62-knockdown, but decreased upon BKPyV-*AGN*-infection, but which could be partly inhibited in the presence of the lysosomal protease inhibitors pepstatin-A1/E64d (**Fig. 7B, right panel**). Together, the data suggested that BKPyV Dun-*AGN* infection of RPTECs was associated with an increased p62/SQSTM1-dependent autophagic flux, which involved MAVS degradation, and which could be reduced by siRNA-p62 knockdown or pepstatin-A1/E64d treatment.

To further investigate mitophagy following BKPyV infection, the tandem tag mitophagy reporter mCherry-mEGFP-OMP25TM carrying the transmembrane domain (TM of OMP25) for targeting to the mitochondrial outer membrane (Bhujabal et al., 2017) was transfected

into RPTECs infected with Dun-*AGN* or Dun-*agn25D39E*. Quantifying the red and green signals in z-stacks of Vp1-expressing cells following confocal microscopy revealed that the number of mitochondrial signals with mCherry red signals exceeding GFP green signals were higher in Dun-*AGN*-infected RPTECs and increased as the number of residual mitochondrial fragments progressively decreased in advanced replication phase as compared to Dun-*agn25D39E*-infected cells or mock-treated controls (**Fig. 7C**). The data indicated that the mitochondrial tandem tag reporter was associated with the mitochondrial network in RPTECs infected with Dun-*AGN* or Dun-*agn25D39E*, but was progressively disrupted and targeted to the acidic environment of autophagosomes in the former. Together, the data supported the notion that BKPyV Dun-*AGN* infection of RPTECs was associated with increased mitophagy, which was not observed to this extent for infection with the mutant Dun-*agn25D39E*.

Mitochondrial colocalization and fragmentation of agnoproteins is conserved among BKPyV, JCPyV, and SV40. To investigate the impact of agnoprotein expression in another BKPyV strain, infection of RPTECs was studied using a well-characterized, yet slowly replicating BKPyV-WW(1.4) strain carrying an archetype NCCR (Bethge et al., 2015; Gosert et al., 2008). Similar to Dun-*AGN*, the BKPyV-WW(1.4) strain showed that agnoprotein expression was associated with mitochondrial fragmentation (**Fig. S8A**). Since agnoprotein homologues have been identified in the human polyomavirus JCPyV, we examined SVG-A cells infected with JCPyV-Mad4 strain, for which cytoplasmic agnoprotein expression has been reported previously (Gosert et al., 2010). Colocalization of JCPyV-agnoprotein with Tom20 and mitochondrial fragmentation was observed (**Fig. S8B**). Moreover, treatment with oleate and staining with LipidTox[TM] revealed colocalization of the JCPyV-agnoprotein with LD as has been reported for the BKPyV-encoded agnoprotein (Unterstab et al., 2010). SV40 infection was studied in CV-1 cells using Vp2/3-expression in the nucleus as a marker of the late viral replication phase (**Fig. 8C**). Since SV40-agnoprotein antibodies were not available, we used the partly cross-reacting antibody raised against the JCPyV-agnoprotein

together with Tom20 demonstrating fragmentation of the mitochondrial network and co-localisation with SV40 agnoprotein (**Fig. 8C**). Together, the data support the view that mitochondrial fragmentation is conserved a conserved feature among different yet related renotropic polyomaviruses found in human and animal species.

Discussion

In the last decade, significant knowledge has been accumulated on how cytoplasmic sensing of viral infections is achieved by the innate immune system (Takeuchi and Akira, 2010; Zevini et al., 2017) and how this crucial first line of defense is subverted to allow for transient or persistent immune escape of acute and chronic viral infections, respectively (Garcia-Sastre, 2017; Goubau et al., 2013). Despite a high infection rate (Virgin et al., 2009) and evidence of immune evasion in the general population (Egli et al., 2009; Kaur et al., 2019), comparatively little is known about relevant mechanisms operating in human polyomaviruses. In this study, we report that the small BKPyV-encoded agnoprotein of 66aa facilitates polyomavirus replication by disrupting the mitochondrial network and its membrane potential during the late phase of the viral replication cycle. Thereby, BKPyV replication is able to evade cytosolic innate immune sensing in this critical phase of viral progeny accumulation, during which DNA damage (Hein et al., 2009) as well as abundant viral DNA genomes and RNA transcripts cumulate in the infected host cell (Funk, 2008; Funk et al., 2006). The precise timing to the critical late viral replication phase (Bernhoff et al., 2008; Low et al., 2004) permits this window of immune evasion to be equally open during *de novo* infection or reactivation, permitting viral cell-to-cell spread below the radar of the immune system.

Transfection experiments using the entire agnoprotein or different subdomains fused to the reporter mEGFP as well as the tet-off inducible agnoprotein expression indicate that

agnoprotein is necessary and sufficient, using its amino-terminal and central helix for mitochondrial targeting and mitochondrial disruption, respectively. As expected from the key role of mitochondria in relaying innate immune sensing, the agnoprotein-mediated disruption of the MMP was associated with significantly reduced IRF3 translocation into the cell nucleus as well as IFN-β transcript and protein expression following poly(*I:C*)RNA or poly(*dA:dT*)DNA stimulation. Importantly, BKPyV viral variants either lacking agnoprotein expression due to a start codon mutation (Dun-*ATCagn*) or carrying mutations abrogating the amphipathic character of the central helix (Dun-*agn25D39E*) showed intact mitochondrial networks and little mitophagy, but significantly impaired replication compared to the wildtype strain Dun-*AGN*. The lower replication rate of the mutant Dun-*agn25D39E* could be partially reversed on three levels of the mitochondrial innate immune relay, namely by CoCl$_2$ treatment affecting the respiratory chain and disrupting the MMP (Jung and Kim, 2004), by BX795 inhibiting the downstream signalling kinase TBK-1, or by type-1 interferon blockade. Although BKPyV as a DNA-virus would be expected to preferentially be sensed via cGAS/STING, increasing data indicate a close interaction between DNA- and RNA-sensing pathway coupling MAVS- and STING-activation on mitochondria and endoplasmic reticulum to TBK-1 activation downstream including through RNA-polymerase III transcription of cytoplasmic DNA-fragments (Chiu et al., 2009; Zevini et al., 2017). Although we can presently not exclude a more direct role of agnoprotein on STING signalling through MMP-breakdown or through its ER-co-localization, our results clearly demonstrate that agnoprotein expression inhibits innate immune responses to both, poly(I:C)-RNA and poly(dA:dT)-DNA. Notably, experimental MMP breakdown by CCCP has been shown to also suppress STING signalling, further merging the cytosolic RNA and DNA-triggered responses (Kwon et al., 2017).

Our study also reveals for the first time a critical role of mitophagy in polyomavirus biology. Mitophagy is known to assist in the disposal of irreversibly damaged mitochondrial fragments. The pepstatin-A1/E64d protease inhibition and siRNA-knockdown experiments

indicate that agnoprotein increases the p62/SQSTM1-dependent steady-state autophagic flux via LC3-II lipidation in a fashion similar to the one observed following the CCCP-induced mitochondrial membrane potential breakdown (Johansen and Lamark, 2011). However, whereas CCCP-induced mitophagy has been reported to directly involve the PINK-Parkin pathway (Bhujabal et al., 2017), our results indicate that mitophagy by agnoprotein involves shifting Parkin by an agnoprotein layer away from the fragmented mitochondria. Confocal microscopy and comparison of LCI/II levels indicate that mitophagy by agnoprotein can be further increased CCCP. Together, this suggests that the agnoprotein-mediated mitophagy differs from the direct CCCP-induced Parkin-involving process. Further evidence for increased autophagic flux was obtained using the tandem tag mitophagy reporter mCherry-mEGFP-OMP25TM revealing acidification in mitophagosomes of BKPyV-*AGN*-infected RPTECs, which was not observed to the same extent in *agn25D39E*-mutant infected RPTECs. Notably, pepstatin-A1/E64d protease inhibition and p62-siRNA knockdown, both of which act downstream of the agnoprotein-mediated impaired innate immune sensing and reduced interferon production, were associated with the reduced Vp1-protein levels in BKPyV Dun-*AGN*-infected RPTECs. This observation suggests that the autophagic flux may be relevant for the effective biosynthesis in the exhaustive late viral replication phase following the functional and structural loss of the mitochondrial power plants (Forbes, 2016). However, the precise adaptor proteins and pathways need further study (Farre and Subramani, 2016).

Innate immune sensors have been characterized in primary human RPTECs and in kidney biopsies by transcriptional profiling identifying antiviral, proinflammatory and proapoptotic responses (Heutinck et al., 2012; Ribeiro et al., 2012). Other studies reported that BKPyV infection of RPTECs can occur without inducing innate immune sensors by as yet unknown mechanisms (Abend et al., 2010; An et al., 2019; de Kort et al., 2017). However, these studies failed to reveal inhibition of BKPyV replication by type-1 interferons, which is in contrast to our results for BKPyV shown here and those recently reported for JCPyV

(Assetta et al., 2016). Importantly, confocal and transmission electron microscopy studies of biopsies obtained from kidney transplant patients provide independent evidence of mitochondrial fragmentation and p62/SQSTM1-mitophagy *in vivo*. These data suggest that the combined effects of immune escape, mitophagy, and viral spread are operating not only in a relevant primary human cell culture model of RPTECs, but also in one of the currently most challenging pathologies affecting kidney transplantation (Ramos et al., 2009). Intriguingly, the BKPyV agnoprotein-induced immune subversion and p62/SQSTM1-mitophagy may explain the earlier reported paradox of abundant detection of agnoprotein *in vivo* and the low agnoprotein-specific antibody and T-cell response (Leuenberger et al., 2007), and provide a new twist to BKPyV-promoting hypoxic mechanisms following renal ischemia/reperfusion injury (Atencio et al., 1993; Fishman, 2002; Hirsch et al., 2006).

Our results also shed new light on previous reports on BKPyV agnoprotein and the closely related JCPyV and SV40 homologues (Gerits and Moens, 2012; Saribas et al., 2018): These include facilitating polyomavirus replication (Ng et al., 1985), increasing viral late phase production (Carswell et al., 1986) and plaques size (Hou-Jong et al., 1987), acting as viroporin (Suzuki et al., 2010) or egress factor (Panou et al., 2018) from the nucleus, by disrupting the tight surrounding mitochondrial network. We demonstrate that mitochondrial fragmentation occurs not only for BKPyV strains carrying an archetype NCCR, but also for the human JCPyV or the monkey SV40 suggesting that this dramatic agnoprotein-mediated function is evolutionary conserved and active across different species, cell types and hosts. In line with this notion, we were unable to identify relevant point mutations in the critical amphipathic wheel of agnoprotein among more than 300 whole genome sequences available in the GenBank (unpublished observation)(Leuzinger et al., 2019). Thus, we conclude that the small agnoprotein of less than 66 aa facilitates BKPyV replication by an effective mechanism also seen in other renotropic human and animal polyomaviruses, which is unmatched in other DNA viruses through its dramatic simplicity (Koshiba et al., 2011).

Indeed, rather complex viral mechanisms has been described targeting the cytosolic sensing platforms at steps up- and down-stream of mitochondria and associated ER-membranes (Khan et al., 2015). Thus, the matrix protein (M-protein) of parainfluenza-3 has been reported to translocate to mitochondria, where it induces mitophagy via LC3-II in a PINK/Parkin-independent manner (Ding et al., 2017). However, the M-protein is a structural protein, and according to the current model acts early following entry into host cells and requires piggy-back import via the mitochondrial elongation factor TUFM (Ding et al., 2017), while our study indicates that the non-structural BKPyV agnoprotein targets mitochondria directly during the late replication phase. Unlike BKPyV, HCV and other (+)-ssRNA viruses replicating in the cytoplasm appear to induce mitochondrial fragmentation and mitophagy in a PINK-Parkin-dependent fashion (Gou et al., 2017). Although structurally unrelated to agnoprotein, the 247aa-long cytomegalovirus (CMV) US9 glycoprotein appears to suppress both MAVS and STING pathways, TBK-1 activation and IFN-β expression by disrupting the mitochondrial membrane potential in the late CMV replication phase (Choi et al., 2018; Mandic et al., 2009). Given the association of agnoprotein with LD, we were intrigued by the similarity to the antiviral host cell protein viperin, which is expressed following RNA- and DNA-virus infections (Gizzi et al., 2018; Hee and Cresswell, 2017). The antiviral effects of viperin have been attributed to a perplexing variety of functions including interference with lipid metabolism and formation of detergent-resistant lipid rafts at the sites of influenza budding (Wang et al., 2007). Similar to agnoprotein, viperin contains an amphipathic helix required for adsorbing to the cytosolic face of LD- and ER-membranes (Hinson and Cresswell, 2009a, b). However, viperin may facilitate viral replication (Hee and Cresswell, 2017), when re-directed to mitochondria by the CMV-encoded Bax-specific inhibitor viral mitochondria-localized inhibitor of apoptosis (vMIA) carrying a mitochondrial targeting domain (Cam et al., 2010; Seo et al., 2011). Thus, the hijacked viperin-CMV-vMIA complex shares properties accommodated in only 66aa of BKPyV-agnoprotein.

Although our study indicates that agnoprotein is both necessary and sufficient, it is presently unclear, whether or not the agnoprotein-mediated MMP breakdown and mitochondrial fragmentation require the direct interaction with cellular proteins other than the mitochondrial import machinery. Given the abundance of BKPyV agnoprotein in the cytoplasm and the clearly distinct colocalization patterns to ER in addition to mitochondria, further studies need to be carefully conducted in order to avoid misleading artefacts. For the Bcl-2 family of proteins, a hierarchy of interactome complexes has emerged over the last decade (Bleicken et al., 2017; Edlich et al., 2011). Moreover, for the related JCPyV-agnoprotein, a strong tendency to form multimeric aggregates *in vitro* has been reported (Sami Saribas et al., 2013), which undoubtedly reduces the specificity of otherwise straight-forward pull-down approaches and complicates potential functional attributions. This notion has been recently confirmed by a detailed study of potential proteomic interaction partners for the homologous JCPyV-agnoprotein. Presently, we favour a minimal model in which mitochondrial targeting of N-terminal domain of agnoprotein allows for embedding the central amphipathic helix into the outer leaflet of the outer mitochondrial membrane similar to other amphipathic proteins, where it remains available for LD-binding (**Fig.9**). This is indirectly supported by the recruitment of LD to the MTScox8-agno(20-66)mEGFP not observed for MTScox8-mEGFP following transfection and oleate treatment. However, further studies are needed to investigate whether or not this simple plug-and-play suffices to progressively leak protons causing loss of membrane potential, preventing TBK-1 activation, and targeting for p62/SQSTM1 mitophagy (Jarsch et al., 2016; McMahon and Gallop, 2005; Shen et al., 2012).

In summary, our study provides important novel insights into how BKPyV-replication is effectively facilitated in the critical late phase of the viral life cycle by the small accessory agnoprotein in at least three complementary ways: i) inactivating immune sensing and the inhibitory effects of interferon-β expression by breakdown of the mitochondrial membrane potential; ii) enhancing the supply of biosynthetic building blocks via increased autophagic

flux, and iii) facilitating viral release of nuclear virions by fragmenting the nucleus-surrounding mitochondrial network. Finally, our observations suggest that BKPyV agnoprotein is not only important for immune escape and urinary shedding in healthy immunocompetent hosts, but may also facilitate the rapid cell-to-cell spread inside the renal tubules, and hence the progression to BKPyV-nephropathy in kidney transplant patients. These insights may help to design novel antiviral and immunization strategies and permit identifying novel markers readily distinguishing BKPyV nephropathy from allograft rejection (Hirsch et al., 2019).

Materials and Methods

<u>Reagents</u>

The following reagents were used (Table 1). Nuclei were stained with Hoechst 33342 (Invitrogen, H21492). To measure mitochondrial membrane potential, cells were stained with 5 μM JC-1 dye (T3168, Molecular Probes) for 30 minutes at 37°C, washed twice with D-PBS for 5 min, followed by addition of live cell imaging solution (A14291DJ, Thermo Fisher). The cells were illuminated at 488 nm and the emission was measured between 515/545 nm and 575/625 nm with Mithras[2] (Berthold Technologies GmbH & Co. KG, Bad Wildbad Germany) or the Safire II plate reader (Tecan, Maennedorf, Switzerland). Mock infected- or no agnoprotein-expressing cells were set to 100%. To induce LDs, culture medium was supplemented with 300 μM oleate (O1383, Sigma-Aldrich) bound to essentially fatty acid free bovine albumin (A6003; Sigma-Aldrich) as described (Unterstab et al., 2010). Poly(I:C) (HMW) Rhodamine (tlrl-picr, InvivoGen) or Poly(dA:dT) Rhodamine (tlrl-patrh, InvivoGen) was transfected with ViaFect™ Transfection Reagent (E4982, Promega) at a reagent: DNA ratio of 3:1 according to manufacturer's instructions. As TBK-1 inhibitor BX-795, (S1274, Selleck Chemicals) was used. BX-795 stock was diluted in DMSO and a stock concentration of 10 mM was generated. CCCP powder (C2759, Sigma-Aldrich) was diluted in DMSO to generate a stock concentration of 100 mM. The protease inhibitors pepstatin A1 (ALX-260-085, Enzo Life Science) and E64d (BML-PI107, Enzo Life Science) were diluted to generate a 10 mg/mL stock concentration in DMSO and Ethanol, respectively.

<u>Antibodies</u>

Goat anti-mouse IgG2a Alexa Fluor 568 (A21134, Invitrogen), donkey anti-rabbit Alexa Fluor 647 (Abcam, ab150075), goat anti-mouse IgG1 Alexa Fluor 488 (A-21121, Molecular Probe), goat anti-mouse IgG1 Alexa Fluor 647 (A-21240, Invitrogen), goat anti-mouse IgG2b Alexa Fluor 647 (A21242, Molecular Probes), goat anti-chicken Alexa Fluor 488 (A11039, Molecular Probes), goat anti-rabbit Alexa Fluor 488 (ab150077, Abcam), donkey anti-guinea pig IRDye 680RD (926-68077, LI-COR) donkey anti-mouse Alexa Fluor 680 (A10038,

Invitrogen), goat anti-rabbit IRDye 800CW (926-32211, LI-COR), donkey anti-goat IRDye 800CW (926-32214, LI-COR), polyclonal rabbit anti-aa40-59 JCPyV agno-sera (generated on request by Eurogentec, Belgium), polyconal anti-BKPyV agnosera (generous gift from C. Hanssen-Rinaldo, clone 81038), polyclonal rabbit anti-aa52-66 BKPyV agno-sera (generated on request by Eurogentec, Belgium, clone 753), polyclonal rabbit anti-aa40-53 BKPyV agno-sera (generated on request by Eurogentec, clone 1163), polyclonal rabbit anti-BKPyV LTag sera (generous gift from C. Hanssen-Rinaldo, clone81048), mouse monoclonal IgG2a anti-SV40-LTag cross-reacting with BKPyV LTag (PAb416, Calbiochem), mouse monoclonal IgG1 anti-BKPyV Vp1 (MAB3204-M19, Abnova), polyclonal rabbit anti-SV40 Vp2/Vp3 (ab53983, Abcam), mouse monoclonal IgG2a anti-Tom20 (sc-17764, Santa Cruz), mouse monoclonal IgG2a anti-MAVS (sc-166583, Santa Cruz), mouse monoclonal IgG1 anti-IRF3 (NBP1-04308, Novus Biologicals), mouse monoclonal IgG1 anti-Interferon α (21116-1, pbl assay science), mouse monoclonal IgG1 anti-Interferon β (21400-1, pbl assay science), mouse monoclonal IgG2a anti-Interferon α / β receptor chain 2 (21385-1, pbl assay science), mouse monoclonal IgG1 anti-p62/SQSTM1 (sc-28359, Santa Cruz), polyclonal guinea pig anti-p62/SQSTM1 (GP62-C, Progen), mouse monoclonal IgG1 anti-PDI (SPA-891, Enzo Life Science), polyclonal chicken anti-Calreticulin (ab14234, Abcam), polyclonal rabbit anti-Phospho-Drp1-(Ser616) (3455, Cell Signalling), mouse monoclonal IgG1 anti-Drp1 (NBP2-23489, Novus Biologicals), mouse monoclonal IgG2b anti-STING (MAB7169, R&D Systems), polyclonal rabbit anti-LC3A/B (4108S, Cell Signalling), mouse monoclonal IgG1 anti-beta Actin (ab6276, Abcam), mouse monoclonal IgG1 and anti-Tubulin (A-11126, Molecular Probes), mouse monoclonal IgG1 anti-GAPDH (sc-47724, Santa Cruz).

Cell lines and viruses

Primary human renal proximal tubular epithelial cells (RPTEC Lot:5111; 4100, ScienCell) were maintained in epithelial cell medium (EpiCM; 4101, ScienCell) and passaged with Trypsin (T3924, Sigma-Aldrich) and Defined Trypsin Inhibitor (DTI; R007100, Invitrogen).

UTA6-2C9 cells stably transfected with the *AGN* gene under a tetracyclin-dependent suppressor (tet-off) have been described previously (Cioni et al., 2013).The African green monkey kidney cell line CV-1 (ATCC CCL-70) was cultured in Dulbecco's modified Eagle's medium (D5671, Sigma-Aldrich) supplemented with 5% FBS (S0115, Biochrom) and 2 mM Stable Glutamine (5-10K50-H, Amimed). The SVG-A cells (generous gift from C. Hanssen-Rinaldo) were kept in Minimum Essential Medium (M2279, Sigma-Aldrich) including 10% FBS. One day after seeding, RPTEC were infected with BKPyV Dunlop at a MOI of 1.0 determined by nuclear LTag staining on RPTECs. The infection was carried out for 2 h before surplus infectious units were removed and EpiCM containing 0.5% FCS was added. One day after seeding, CV-1 cells were infected with SV40 (776 strain) at a MOI of 0.5-1.0 pfu/cell using a supernatant from SV40-infected BS-C-1 cells. The infection was carried out for 2 h before surplus infectious units were removed and complete medium added. One day after seeding, SVG-A cells were infected with JCPyV Mad-4 using a supernatant from JCPyV Mad-4 transfected COS-7 cells. The infection was carried out for 2 h before surplus infectious units were removed and complete medium added.

Quantitative nucleic acid testing (QNAT)

BKPyV loads were quantified form cell culture supernatant as described previously (Hirsch et al., 2016). To quantify interferon-β transcripts, cellular RNA was extracted using the QIAshredder kit and the RNeasy Mini kit (79654 and 74106, Qiagen). The cDNA was generated using High Capacity cDNA reverse transcript kit (4368814, Applied Biosystems) and RT-QNAT was performed on a Veriti 96 Well Thermal Cycler (Applied Biosystems, Lincoln, USA) using an human interferon b1 assay (4331182, Thermo Fisher Scientific) and human HPRT1 endogenous control (4333768F, Applied Biosystems).

Plasmids

The different agnoprotein constructs have been described in (Unterstab et al., 2010). Mitochondrial membrane localization signal (MTS) of cytochrome c oxidase cox8 was N-

terminally fused to agno(20-66)-EGFP and named MTScox8-Agno(20-66). The mCherry-mEGFP-OMP25TM tandem tag mitophagy reporter was a generous gift of Professor Terje Johansen, UiT The Arctic University of Norway, Tromsø, Norway. The YFP-Parkin plasmid was a gift from Richard Youle (Addgene plasmid # 23955; http://n2t.net/addgene:23955 ; RRID:Addgene_23955)(Narendra et al., 2008).

Transfection

Transfection of BKPyV genomic DNA or plasmids containing the agnoprotein into RPTECs or UTA6 cells was performed at 90-95% confluency using ViaFect™ Transfection Reagent (E4982, Promega) at a reagent: DNA ratio of 3:1 according to manufacturer's instructions. At 24 h post-transfection, medium was replaced with EpiCM with 0.5% FBS and DMEM high Glucose containing 5% FBS (S0115, Biochrom AG), respectively. For p62/SQSTM1 knock-down experiments the SMARTpool siGENOME SQSTM1 siRNA (M-010230-00-0005, Dharmacon) and siGENOME Non-Targeting siRNA #2 (D-001210-02-05, Dharmacon) were transfected with Lipofectamine 3000 Reagent (L3000001, Thermo Fisher) according to manufacturer's instructions, using 100 nM siRNA.

Electron microscopy

Samples were rinsed with ice-cold cacodylate 0.1 M pH 7.4 and fixed with 2% paraformaldehyde, 2.5% glutaraldehyde (01909; Polysciences) in 0.1 M cacodylate pH 7.4. Samples were washed again with cacodylate buffer (0.1 M, pH 7.0 and postfixed in 1% osmium tetroxide and 1.5% potassium ferrocyanide in cacodylate buffer (0.1 M pH 7.4), followed by a 1% osmium in cacolylate buffer treatment. Sections were washed in distilled water. Samples were stained with 1% uracil acetate in water and rinsed once with H_2O. Samples were de-hydrated in graded alcohol series and embedded in Epon. The images were taken with a TecnaiTm Spirit TEM (Fei, Thermo Fisher). Biopsy specimens were processed and analyzed as described previously (Drachenberg et al., 1999).

Immunofluorescence microscopy

Microscopy was performed using an epifluorescence microscope (model eclipseE800; Nikon, Tokio, Japan) equipped with suitable filters and a digital camera (Hamamatsu, Tokio, Japan). The pictures were analyzed by Fiji (U.S. National Institutes of Health, Bethesda, MD) as described previously (Hirsch et al., 2016).

Confocal Laser Scanning Microscopy (CLSM)

Confocal pictures were taken with a LeicaSP5 (Leica, Wetzlar, Germany) with a 63x Plan Apochromat/NA1.4 oil objective and pictures were further processed by Fiji. Z-stacks were acquired with a 63x Plan Apochromat/NA1.4 oil objective and the requirements of the Nyquist theorem were fulfilled, with voxel *xyz* size of 45, 45, 150 nm, respectively. To exclude the possibility of channel crosstalk, images were acquired sequentially using the multi-track mode. Deconvolution and visualization were done essentially as described in (Unterstab et al., 2010). For deconvolution, the Huygens professional software (Scientific Volume Imaging, Hilversum, the Netherlands) was used, applying the Classic Maximum Likelihood Estimation Mode (CMLE) using a theoretical point spread function (PSF) and IMARIS (Bitplane AG, Zurich, Switzerland) was used for visualization, colocalization analyzes, and quantification.

SDS/PAGE and immunoblotting

Cells were lysed with RIPA buffer: 10 mM Tris/HCl pH 7.5; 150 mM NaCl; 0.5 mM EDTA. 1.0% Nonidet P-40 and proteinase inhibitor (04693132001, Roche). Cell lysates were separated by Mini Protean TGX Gradient Gel 4-20% (4561095, Bio-Rad) and electrotransferred onto 0.45-µm Immobilon-FL polyvinylidene difluoride (PVDF) membrane (IPFL00010, Millipore/Merck). Membranes were blocked with Odyssey blocking buffer (927-40000, LI-COR) diluted 1:2 in Tris-buffered saline (TBS). Primary and secondary antibodies were diluted in Odyssey blocking buffer diluted 1:2 in TBS-0.1% Tween 20 and incubated at RT for 1 h (or o/n at 4°C) or 45 min, respectively. Washing in between was performed with

TBS–0.1% Tween 20. For the detection of p62/SQSTM1, cells were trypsinized and 2.0×10^4 cells were resuspended and lysed per 10µL 1x Laemmli Sample Buffer (161-0747, Bio-Rad). 10µL cell lysate was separated by SDS-PAGE, electrotransfered onto 0.2-µm Immobilon®-P^{SQ} PVDF membrane (ISEQ00010, Millipore/Merck) and immunoblotting was done essentially as described above, but 3.0% milk (T145.2, Roth) in TBS-0.1% Tween 20 was used instead of the Odyseey blocking buffer. Detection and quantification were done with the Odyssey CLx system (LI-COR, Lincoln, USA).

Enzyme-linked immunosorbent assays (ELISA)

The ELISA was performed using the VeriKine-HsTM Human IFN Beta Serum ELISA KIT (41415, pbl assay science). The recommended enhanced protocol form improved performance in serum evolution was used. Samples were measured by Safire II plate reader in triplicates and the standard curve was generated using GraphPad Prism software (v8.1.0).

Immunohistochemistry

After deparaffinization samples were heated for 30 min in the microwave at 98°C in citrate-buffer (pH 6.0) and cooled down for 30 min. Samples were washed 1×5min with D-PBS and 1×5 min in 10 mM D-PBS + 0.1% Tween20 (93773, Fluka). Samples were blocked with 5% normal goat serum (50197Z, Thermo Fisher), for 1 h at RT. Primary antibodies were diluted in 3% BSA/PBS (A9647, Sigma-Aldrich) and samples were incubated over night at 4°C followed by 2×5min washes in 10 mM PBS. Secondary antibodies were diluted in 3% BSA/PBS containing 1 µg/ml Hoechst 33342 (B2261, Sigma-Aldrich), followed by 2×5 min washes in 10 mM PBS. Samples were mounted with Prolong Gold (P36935, Thermo Fisher).

Statistics

All results were analyzed by GraphPad Prism software (version 8.1.0). Level of significant variability between groups was tested as indicated in figure legends, using paired t-test or two-way analyses of variance (ANOVA), respectively. Significant differences were assumed for P values of <0.05. Unless indicated differently, groups were plotted as mean values and standard deviation (SD).

Acknowledgments:

We thank Pascal Lorentz of the confocal imaging core facility of the Department Biomedicine, University of Basel, and Mohamed Chami at the electron microscopy core facility of the Biocenter, University of Basel for advice and support with the imaging studies. We are grateful to Professor Terje Johansen, UiT The Arctic University of Norway, Tromsø, Norway, for helpful discussions and for providing the tandem tag mitophagy reporter construct mCherry-mEGFP-OMP25TM. We are indebted to Ms Erika Hofmann for the timely updates of the reference library.

The data presented in this manuscript are tabulated in the main paper and in the supplementary materials.

Funding: This study was funded by an appointment grant of the University of Basel to H.H.H.

Author contributions: J Manzetti, F Weissbach, FE Graf, G Unterstab, M Wernli, C Hanssen-Rinaldo, C Drachenberg, H Hopfer conducted experiments, analyzed data, and contributed to writing the manuscript; HHH designed the study, analyzed data, and wrote the manuscript.

Declaration of Interests: The authors declare no competing interests

Table 1: KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-BKPyV agno (clone 81038)	(Rinaldo et al., 1998)	N/A
Rabbit polyclonal anti-aa52-66 BKPyV agno (clone 753)	This paper (unpublished)	N/A
Rabbit polyclonal anti-aa40-53 BKPyV agno (clone 1163)	This paper (unpublished)	N/A
Rabbit polyclonal anti-aa40-59 JCPyV agno (clone 1172)	This paper (unpublished)	N/A
Rabbit polyclonal anti-BKPyV LTag (clone 81048)	(Rinaldo et al., 1998)	N/A
Mouse monoclonal IgG2a anti-SV40 LTag	Calbiochem	Cat#PAb416
Mouse monoclonal IgG1 anti-BKPyV Vp1	Abnova	Cat#MAB3204-M19
Rabbit polyclonal anti-SV40 Vp2/Vp3	Abcam	Cat#ab53983
Mouse monoclonal IgG2a anti-Tom20	Santa Cruz	Cat#sc-17764
Mouse monoclonal IgG2a anti-MAVS	Santa Cruz	Cat#sc-166583
Mouse monoclonal IgG1 anti-IRF3	Novus Biological	Cat#NBP1-04308
Mouse monoclonal IgG1 anti-Interferon α	pbl assay science	Cat#21116-1
Mouse monoclonal IgG1 anti-Interferon β	pbl assay science	Cat#21400-1
Mouse monoclonal IgG1 anti-Interferon α/β receptor chain 2	pbl assay science	Cat#21385-1
Mouse monoclonal IgG1 anti-p62/SQSTM1	Santa Cruz	Cat#sc-28359
Guinea pig polyclonal anti-p62/SQSTM1	Progen	Cat#GP62-C
Mouse monoclonal IgG1 anti-PDI	Enzo Life	Cat#SPA-891
Chicken polyclonal anti-Calreticulin	Abcam	Cat#ab14234
Rabbit polyclonal phosphor-Drp1(Ser616)	Cell Signalling	Cat#3455
Mouse monoclonal IgG1 anti-Drp1	Novus Biologicals	Cat#23489
Mouse monoclonal IgG2b anti-STING	R&D Systems	Cat#MAB7169
Polyclonal rabbit anti-LC3A/B	Cell Signalling	Cat#4108S
Mouse monoclonal IgG1 anti-beta Actin	Abcam	Cat#ab6276
Mouse monoclonal IgG1 anti-Tubulin	Invitrogen	Cat#A-11126
Mouse monoclonal IgG anti-GAPDH	Santa Cruz	Cat#sc-47724
Goat anti-mouse IgG2a Alexa Fluor 568	Invitrogen	Cat#A-21134
Donkey anti-rabbit Alexa Fluor 647	Abcam	Cat#ab150075
Goat anti-mouse IgG1 Alexa Fluor 488	Invitrogen	Cat#A-21121
Goat anti-mouse IgG1 Alexa Fluor 647	Invitrogen	Cat#A-21240
Goat anti-mouse IgG2b Alexa Fluor 647	Invitrogen	Cat#A-21242
Goat anti-chicken Alexa Fluor 488	Invitrogen	Cat#A-11039
Goat anti-rabbit Alexa Fluor 488	Abcam	Cat#ab150077

Donkey anti-guinea pig IRDye 680RD	LI-COR	Cat#926-68077
Donkey anti-mouse Alexa Fluor 680	Invitrogen	Cat#A-10038
Goat anti-rabbit IRDye 800CW	LI-COR	Cat#926-32211
Donkey anti-goat IRDye 800CW	LI-COR	Cat#926-32214
Bacterial and Virus Strains		
BKPyV Dunlop-AGN, GENBANK: V01108.1	(Henriksen et al., 2015)	N/A
SV40 (776 strain), GENBANK: AF316139.1	(Henriksen et al., 2016)	N/A
JCPyV Mad-4	ATCC	Cat#ATCC VR-1583
BKPyV-WW(1.4), GENBANK: AB211371.1	(Gosert et al., 2008)	N/A
Chemicals, Peptides, and Recombinant Proteins		
Oleate	Sigma-Aldrich	Cat#O1383
JC-1 dye	Invitrogen	Cat#T3168
Hoechst 33342	Invitrogen	Cat#H21492
LipidTox Red	Molecular Probes	H34476
Poly(I:C) (HMW) Rhodamine	InvivoGen	Cat#tlrl-picr
Poly(dA:dT) Rhodamine	InvivoGen	Cat#tlrl-patrh
ViaFect	Promega	Cat#E4982
Lipofectamine 3000	Thermo Fisher	Cat#L3000001
IFN-beta 1a	pbl assay science	Cat#11410-2
BX795	Selleck Chemicals	Cat#S1274
$CoCl_2$	Sigma-Aldrich	Cat#C8661
Tetracycline	Sigma-Aldrich	Cat#T7660
CCCP	Sigma-Aldrich	Cat#C2759
Pepstatin A1	Enzo Life Science	Cat#ALX-260-085
E64d	Enzo Life Science	Cat#BML-PI107
Critical Commercial Assays		
VeriKine-HsTM Human IFN Beta Serum ELISA KIT	pbl assay science	Cat#41415
High Capacity cDNA reverse transcript kit	Applied Biosystems	Cat#4368814
QIAshredder kit	Qiagen	Cat#79654
RNeasy Mini kit	Qiagen	Cat#74106
Human Interferon b1 assay	Thermo Fisher Scientific	Cat#4331182
Human-HPRT1 endogenous control	Applied Biosystems	Cat#4333768F
Experimental Models: Cell Lines		
RPTEC (Lot: 5111)	ScienCell	4100
CV-1	ATCC	ATCC CCL-70
SVG-A	(Henriksen et al., 2014)	N/A

Vero E6	ATCC	ATCC No. CRL-1586
UTA6-2C9	(Cioni et al., 2013)	N/A
Oligonucleotides		
SMARTpool siGENOME SQSTM1 siRNA	Dharmacon	Cat#M-010230-00-0005
siGENOME Non-Targeting siRNA #2	Dharmacon	Cat#D-001210-02-05
BKPyV qPCR primer F: AGCAGGCAAGGGTTCTATTACTAAAT	(Hirsch et al., 2016)	N/A
BKPyV qPCR primer R: GAAGCAACAGCAGATTCTCAACA	(Hirsch et al., 2016)	N/A
BKPyV qPCR probe: FAM-AAGACCCTAAAGACTTTCCCTCTGATCTACACCAGTTT-TAMRA	(Hirsch et al., 2016)	N/A
Recombinant DNA		
Agno(1-66)-mEGFP	(Unterstab et al., 2010)	N/A
Agno(1-53)-mEGFP	(Unterstab et al., 2010)	N/A
Agno(1-20)-mEGFP	(Unterstab et al., 2010)	N/A
Agno(20-66)-mEGFP	(Unterstab et al., 2010)	N/A
Agno(42-66)-mEGFP	(Unterstab et al., 2010)	N/A
MTScox8-Agno(20-66)-mEGFP	This paper (unpublished)	N/A
MTScox8-mEGFP	This paper (unpublished)	N/A
mCherry-mEGFP-OMP25TM reporter plasmid	(Bhujabal et al., 2017)	N/A
YFP-Parkin	(Narendra et al., 2010)	Addgene Plasmid #23955
Software and Algorithms		
GraphPad Prism software (version 8.1.0)	GraphPad	https://www.graphpad.com
Fiji	Fiji	https://fiji.sc
Huygens professional software (version 19.10)	Scientific Volume Imaging	https://svi.nl/Huygens-Professional
IMARIS (version 9.3.1)	Bitplane AG	https://imaris.oxinst.com/

Figure legends

Fig. 1. Agnoprotein colocalizes with mitochondria and induces mitochondrial fragmentation in the late replication phase of BKPyV infection.

(**A**) Z-stacks of RPTECs infected with BKPyV Dun-*AGN* (top row) or with Dun-*agn25D39E* (bottom row, large replicating cell next to small non-replicating cell) at 48 hpi, stained for Tom20 (red), agnoprotein (green), and DNA (blue). Colocalizing voxels are shown in yellow.

(**B**) Quantification of mitochondrial morphology in six fields of two independent experiments using Fiji software (mean ± SD, 2-way ANOVA).

(**C**) Z-stacks of BKPyV Dun-*agn25D39E* infected cells at 48 hpi, stained for mitochondrial marker Tom20 (red), calreticulin as marker for the endoplasmic reticulum (magenta), agnoprotein (green), and DNA (blue). Colocalizing voxels are shown in yellow (Scale bar 5 µm).

(**D**) Confocal images of RPTECs infected with BKPyV-Dun-*AGN* at indicated times post infection. Cells were stained for LTag (red), agnoprotein (green), mitochondrial marker Tom20 (cyan), and DNA (blue). White arrows indicate cells magnified (Scale bar 20 µm).

Fig. 2. The amphipathic character of the central agnoprotein helix is required for mitochondrial fragmentation.

(**A**) Three-dimensional ribbon model of each agnoprotein derivate as predicted with the Quark online algorithm (https://zhanglab.ccmb.med.umich.edu/I-TASSER/) and predicted amphipathic helical wheel of the agnoprotein amino acids 22 to 39.

(http://cti.itc.virginia.edu/~cmg/Demo/wheel/wheelApp.html).

(**B**) Confocal images of RPTEC-infected with BKPyV Dun-*AGN* and isogenic derivatives Dun-*agn25D39E*, Dun-*agn25L39L*, and Dun-*ATCagn*, respectively. Cells were fixed at 48 hpi and stained for mitochondrial marker Tom20 (red), agnoprotein (green), Vp1 (cyan), and DNA (blue)(Scale bar 20 µm).

(**C**) Confocal images of RPTECs infected with BKPyV-Dun-*AGN* and BKPyV-Dun-*agn25D39E*, respectively, at 48 hpi. Cells were mock-treated or treated with 300 μM oleate at 24 hpi. Cells were stained for Tom20 (red), agnoprotein (green), and DNA (blue) (Scale bar 20 μm).

(**D**) Live cell imaging using JC-1 dye (5μM) of mock-infected or infected with BKPyV Dun-*AGN* and isogenic derivatives Dun-*agn25D39E*, Dun-*agn25L39L*, and Dun-*ATCagn*, respectively at 48 hpi. Quantification of mitochondrial membrane potential (Ψ_m) by measuring red fluorescent signal (JC-1 aggregates) with the Safire II plate reader of three independent experiments (mean ± SD, Kruskal-Wallis test).

Fig. 3. Agnoprotein is sufficient to mediate structural and functional alterations of mitochondrial network

(**A**) Schematic presentation of agnoprotein-mEGFP fusion constructs transfected into UTA-6 cells. Retained aa indicated in parenthesis and presented as solid line, deleted parts presented as dotted line. The central amphipathic helix (aa 22–39) is shown as blue bar. Confocal images of transfected UTA-6 cells, transiently expressing the indicated agnoprotein-mEGFP fusion constructs were taken at 24 hpt. Immunofluorescent staining for Tom20 (red) and agnoprotein (magenta), GFP (green); DNA (blue)(Scale bar 20 μm). For MTScox8-agno(20-66), Z-stacks were obtained and colocalizing voxels are shown in yellow, white dashed-lined rectangle indicating enlarged section.

(**B**) UTA6-2C9 cells stably transfected with tetracycline (tet)-off inducible BKPyV agnoprotein were cultured for 24 h in the presence (+tet) or absence (-tet) of tetracycline to suppress or induce BKPyV agnoprotein expression, respectively. Confocal images of cells stained for DNA (blue), agnoprotein (green), and Tom20 (red). White rectangle indicating enlarged section. Graph representing corresponding mitochondrial morphology, quantification of six fields using Fiji software of two independent experiments (mean ± SD; 2-way ANOVA)

(**C**) Ψ_m was assessed by JC-1 dye and imaging of live cells using the signal ratio of aggregate (red) / monomeric (green) normalized to UTA6-2C9 cells not expressing agnoprotein (+tet)

versus cells expressing agnoprotein (-tet) using Mithras[2] (mean ± SD, unpaired parametric t-test).

(**D**) Nuclear IRF3 translocation following poly(I:C) transfection was compared in UTA6-2C9 cells cultured for 24 h in the presence or absence of tetracycline to suppress or induce BKPyV agnoprotein expression, respectively. Increasing amounts of rhodamine-labeled poly(I:C) was delivered to the cells via lipofection, cells were fixed at 4 hpt, stained for IRF3 (magenta), agnoprotein (green) and DNA (blue) (left images, 1000 ng/mL poly(I:C), right panel quantification of nuclear IRF3 of six fields using Fiji software (mean ± SD, Mann-Whitney).

(**E**) Nuclear IRF3 translocation following poly(dA:dT) transfection was compared in UTA6-2C9 cells as described in D (left images, 1000 ng/mL poly(dA:dT), right panel quantification of nuclear IRF3 (mean ± SD, Mann-Whitney).

(**F**) Quantification of IFN-β mRNA and IFN-β secretion into cell culture supernatants following poly(dA:dT) stimulation of UTA6-2C9 cells, three experiments (mean ± SD, Mann-Whitney t-test).

Fig. 4. BKPyV replication in primary human RPTECs is sensitive to type-1 interferon

(**A**) RPTECs were treated overnight with the indicated concentrations of IFN-β or solvent. LTag positive cells (left panel; triplicates, mean ± SD, 2-Way ANOVA) and supernatant BKPyV loads (middle panel; triplicates, mean ± SD, 2-Way ANOVA) were quantified at the indicated times, and a representative LTag staining of RPTECs at 72 hpi is shown (right panel).

(**B**) RPTECs were pre-treated with the indicated concentrations of IFN-β and expression of IFIT1 (*ISG56*) and BKPyV late viral proteins Vp1 and agnoprotein were analyzed at the indicated times post-infection by immunoblot analysis (left panel). RPTECs were treated before or at the indicated times post-infection with 200 U/mL IFN-β, and supernatant BKPyV loads were measured at 72 hpi (middle panel; triplicates, mean ± SD, Kruskal-Wallis test). BKPyV-infected RPTECs were treated with IFN-β in the presence or absence of anti-IFN consisting of antibodies blocking IFN-α, IFN-β and interferon α/β receptor (right panel), and

supernatant BKPyV loads were measured at 72 hpi (right panel; duplicates, mean ± SD, unpaired parametric t-test).

(**C**) RPTECs were infected with the indicated BKPyV variants (MOI = 1 by nuclear LTag staining of RPTCs) and the number of infected cells were quantified by immunofluorescence at 72 hpi (left panel; triplicates, mean ± SD, unpaired parametric t-test) and supernatant BKPyV loads were measure at the indicated time post-infection (middle panel; triplicates, mean ± SD, 2-Way ANOVA). RPTECS were infected with BKPyV Dun-*agn25D39E*, the TBK-1 inhibitor BX795, antibodies blocking IFN-α, IFN-β and interferon α/β receptor or solvent were added at 36 hpi, and supernatant BKPyV loads were measured at 72 hpi (right panel; triplicates of two independent experiments, mean ± SD, unpaired parametric t-test).

(**D**) RPTECs were infected with BKPyV-Dun-*agn25D39E* and treated with the indicated concentrations of $CoCl_2$ or solvent at 24 hpi, and at 72 hpi, confocal microscopy was performed for Tom20 (red), agnoprotein (green) and DNA (blue), and supernatant BKPyV loads quantified (right panel; triplicates, mean ± SD, unpaired parametric t-test).

Fig. 5. MAVS co-localizes to fragmented mitochondria and is decreased in BKPyV-infected RPTECs

(**A**) RPTECs were infected with BKPyV-Dun-*AGN* or mock-treated and were either fixed after 72 h for confocal microscopy staining for DNA (blue), agnoprotein (green), and MAVS (red) (top left panel), or were harvested lysing $2.0x10^4$ cells per 10 μL Laemmli Sample Buffer and analysed by SDS/PAGE and immunoblotting for MAVS and agnoprotein (top panels). The immunoblot (lower panel) stained for MAVS and agnoprotein (1st) was subsequently stained for GAPDH and Tom20 (2nd). MAVS and Tom20 levels were normalized to GAPDH levels as indicated (right panel). GAPDH signal was overlapping with MAVS specific band (indicated by asterisk) which was subtracted prior normalization.

(**B**) RPTECs were infected with BKPyV-Dun-*AGN* or mock-treated and were either fixed after 72 h for confocal microscopy for DNA (blue), Vp1 (cyan), MAVS (red), phosphorylated Drp1-S616 (magenta) (bottom left panel). Cell lysates were analysed by SDS/PAGE and

immunoblotting using antibodies to total Drp1, phosphorylated Drp1-S616, agnoprotein, and actin (right panel).

Fig. 6. Agnoprotein-mediated mitochondrial fragmentation and p62/SQSTM1-autophagosomes in cell culture and kidney transplant biopsy tissue.

(**A**) RPTECs were infected with BKPyV-Dun-*AGN* or BKPyV-Dun-*agn25D39E*. At 72 hpi, cells were fixed and processed for TEM (top panels, scale bar 2μm) or for confocal microscopy (bottom left panels) staining for Tom20 (red), agnoprotein (green), p62/SQSTM1 (cyan) and DNA (blue). p62/SQSTM1-positive autophagosomes of six fields were quantified using Fiji at 48 hpi and 72 hpi (bottom right panels; mean ± SD, 2-way ANOVA).

(**b**) UTA6-2C9 cells were cultured for 24 h or 48 h in the presence or absence of tetracycline to suppress or induce BKPyV agnoprotein expression, respectively. At the indicated times, confocal microscopy (left panels) was performed after fixing and staining for Tom20 (red), agnoprotein (green), p62/SQSTM1 (cyan) and DNA (blue). p62/SQSTM1-positive autophagosomes were quantified as described in A at 24 h and 48 h (right panels; mean ± SD, 2-way ANOVA).

(**c**) Tissue biopsies from kidney transplant patients with (infected) or without (non-infected) BKPyV-associated nephropathy were studied by transmission electron microscopy (left panels, scale bar 1μm) or confocal microscopy (right panels) staining for Tom20 (red), agnoprotein (green), p62/SQSTM1 (cyan), and DNA (blue).

Fig. 7. Agnoprotein mediates p62/SQSTM1-dependent autophagic flux and mitophagy.

(**A**) UTA6-2C9 cells were cultured for 48 h in the presence or absence of tetracycline to suppress or induce BKPyV agnoprotein expression, respectively. At 24 h post-induction, pepstatin-A1/E64d (10 μg/mL) and CCCP (10 μM) was added as indicated and cell extracts were prepared and analysed by immunoblotting as described in Materials & Methods, using RIPA buffer, for LC3-I and II, agnoprotein expression and actin. LC3-II/I ratio were normalized using untreated cells without agnoprotein (+tet) expression as reference.

(**B**) RPTECs transfected with siRNA-p62 (p62 knockdown) or control were infected with BKPyV Dun-*AGN* (indicated as *AGN*). Pepstatin-A1/E64d (10 µg/mL) was added at 48 hpi, when indicated, 2.0x10^4 cells were harvested and lysed per 10 µL Laemmli Sample Buffer and analysed by immunoblotting as described in Material & Methods for p62/SQSTM1, BKPyV Vp1, and LC3-I and -II expression (on 0.22 µm PVDF membrane, left panel) or MAVS, actin, BKPyV LTag, and agnoprotein (on 0.45 µm PVDF-FL membrane, right panel).

(**C**) RPTECs transfected with mCherry-GFP-OMP25TM tandem tag mitophagy reporter were infected with BKPyV Dun-*AGN* or BKPyV-Dun-*agn25D39E*. At 72 hpi, cells were fixed and stained for Vp1 and DNA. Colocalization of mCherry and GFP is shown (yellow), Vp1 (cyan), and DNA (blue). Z-stacks were acquired and analyzed in IMARIS, the mCherry signal was transformed into countable spots (according voxel intensity). The GFP and mCherry mean intensity within the spots were quantified (bars represent mean ± 95% CI, Wilson/Brown method).

Supplementary Material 1: Figure S1.

Comparing the prediction of mitochondrial presequence BKPyV agnoprotein, point mutant derivatives and cytochrome c oxidase cox8.

A mitochondrial presequence was predicted within the blue bar of the first 25 amino acids with a precision of 0.79 and a recall of 0.80 using the MitoFate prediction tool (http://mitf.cbrc.jp/MitoFates/cgi-bin/top.cgi, accessed 26.02.2020) for the amino acids encoded in the *AGN, agn25D39E, agn25L39L* sequences and for the mitochondrial targeting sequence (MTS) of cytochrome c oxidase cox-8.

Supplementary Material 2: Figure S2.

BKPyV early and late viral protein expression in primary human RPTECs

Cells were infected with the indicated viral strains, fixed at 48 hpi, stained for confocal microscopy and z-stack acquisition as described in Materials & Methods.

(**A**) BKPyV Dun-*AGN*: DNA (blue), early viral LTag (red), late viral Vp1 (cyan) and agnoprotein (green).

(**B**) BKPyV Dun-*agn25D39E:* DNA (blue), early viral LTag (red), late viral Vp1 (cyan) and agnoprotein (green).

(**C**) BKPyV Dun-*agn25D39E*: DNA (blue), Tom20 (red), agnoprotein (green), PDI (magenta). Colocalizing voxels are shown in yellow.

Supplementary Material 3: Figure S3.

MTScox8-mEGFP and MTScox8-Agno(20-66)mEGFP co-localize to mitochondria, but only MTScox8- Agno(20-66)mEGFP disrupts mitochondrial network and binds to lipid droplets (LD).

(**A**) Confocal images of transfected UTA-6 cells, transiently expressing MTScox8-mEGFP fusion protein at 24 hpt. DNA (blue), Tom20 (red), and mEGFP (green).

(**B**) Confocal images of transfected Vero cells, transiently expressing MTScox8-mEGFP or MTScox8-Agno(20-66)mEGFP fusion proteins at 24 hpt. Left panels in the absence (- oleate): DNA (blue), Tom70 (red), and mEGFP (green). Right panels in the presence of 300 µM oleate (+ oleate). DNA (blue), Tom70 (magenta), and mEGFP (green), LD (red).

(**C**) Confocal images of transfected Vero cells, transiently expressing MTScox8- Agno(20-66)mEGFP fusion proteins at 24 hpt in the presence of 300 µM oleate (+ oleate). DNA (blue), Tom70 (magenta), and mEGFP (green), LD (red).

Supplementary Material 4: Figure S4.

Inhibition of TBK-1 under conditions mediating partial rescue of BKPyV Dun-*agn25E39D* replication is not affecting BKPyV-Dun-*AGN* replication.

RPTECS were infected with BKPyV Dun-*AGN*, the TBK-1 inhibitor BX795 or solvent were added at 36 hpi, and supernatant BKPyV loads were measured at 72 hpi, input virus load is indicated by the 24 hpi time-point (triplicates of two independent experiments, mean ± SD, unpaired parametric t-test).

Supplementary Material 5: Figure S5.

Phosphorylation of Drp1 at serine 626 (Drp1-S626) is increased in BKPyV infected RPTECs and occurs independent in isogenic BKPyV variants expressing the amphipathic helix mutant (BKPyV Dun-*agn25D39E*) or no agnoprotein (BKpyV Dun-*ATCagn*).

(**A**) RPTECs were infected with BKPyV-Dun-*AGN* or mock-treated and at 72 hpi, cell lysates were prepared using RIPA buffer, and analysed by SDS/PAGE and immunoblotting using antibodies to Drp1-total protein (red triangle), BKPyV LTag (green triangle), Vp1, and agnoprotein.

(**B**) RPTECs were infected with Dun-*agn25D39E* , Dun-*ATCagn,* or mock-treated and confocal microscopy was performed for Vp1 (cyan), Drp1-S616 (magenta), MAVS (red), and DNA (blue).

Supplementary Material 6: Figure S6.

BKPyV Dun-*AGN*-replicating RPTECs show mitochondrial fragmentation and increased Tom20-p62/SQSTM1 colocalization compared to BKPyV Dun-*agn25D39E*-replicating RPTECs

RPTECs were infected with BKPyV Dun-*AGN* or Dun-*agn25D39E* and at 72 hpi confocal microscopy was performed for Tom20 (red) agnoprotein (green), p62/SQSTM1 (cyan), and DNA (blue). Colocalizing voxels shown in yellow.

Supplementary Material 7: Figure S7.

UTA6-2C9 cells transfected with YFP-Parkin show parkin-positive mitophagy following CCCP treatment, but displacement by intermediate agnoprotein layer following agnoprotein expression.

UTA6-2C9 cells harboring the tetracycline (tet)-off inducible BKPyV agnoprotein were transfected with yellow-fluorescence protein-Parkin construct, and confocal microscopy was performed at 48 hpt.

BKPyV agnoprotein suppression (tet+) and expression (tet-) is indicated, respectively. CCCP treatment (10 µM) for 3 h served as positive control for parkin-positive mitophagy. Confocal images of cells stained for DNA (blue), Tom20 (red), YFP-parkin (yellow), and agnoprotein (green). Magnification and 3D-isosurface rendering showing agnoprotein layer (green) displacing YFP-parkin layer (yellow).

Supplementary Material 8: Figure S8.

Agnoprotein of archetype BKPyV, JC Polyomavirus and the simian SV40 colocalize to mitochondria and disrupt the mitochondrial network.

(**A**) RPTECs were infected with BKPyV-WW(1.4) carrying an archetype non-coding control region, and confocal microscopy was performed at 6 dpi after staining for Tom20 (red), agnoprotein (green), and DNA (blue).

(**B**) SVG-A cells were infected with JCPyV-*Mad4* and confocal microscopy was performed at 72 hpi after staining for Tom20 (red), anti-JCPyV agnoprotein (green), and DNA (blue) (top panels). Z-stacks of JCPyV-replicating SVG-A cells were acquired, deconvolved and processed with IMARIS. 3D isosurface-renderings of the mitochondrial marker Tom20 (red), agnoprotein (green), and DNA (blue) are shown. Colocalizing voxels are shown in yellow (middle panels). Confocal Image of SVG-A cells infected with JCPyV-Mad4, 48 hpi, treated with 300 µM oleate at 24 hpi. Cells were stained for lipid droplets (red), agnoprotein (green), Vp1 (cyan), and DNA (blue). Lipid droplets indicated by white arrow (bottom panels).

(**C**) CV-1 cells were infected with SV40 and confocal microscopy was performed at 48 hpi after staining for Tom20 (red), Vp2/3 (blue) (top panels) or cross-reacting antisera raised against JCPyV agnoprotein (green)(lower panels).

Supplementary Material 9: Figure S9

Working model of BKPyV agnoprotein mediating innate immune evasion by mitochondrial membrane breakdown, network fragmentation and mitophagy.

Supplementary Material 10: Video S1. Dun-*AGN* Agnoprotein colocalizes with mitochondria and induces mitochondrial fragmentation. Cells were infected with the indicated viral strains and fixed at 48 hpi as described in Materials & Methods. Z-stacks of BKPyV Dun-*AGN*, stained for Tom20 (red), agnoprotein (green), and DNA (blue). Colocalizing voxels are shown in yellow.

Supplementary Material 11: Video S2. Dun-*agn25D39E* mutant agnoprotein colocalizes with mitochondria, but does not induces fragmentation of mitochondrial network.
 Z-stacks of BKPyV Dun-*agn25D39E* infected cells, stained for mitochondrial marker Tom20 (red), agnoprotein (green), and DNA (blue). Colocalizing voxels are shown in yellow.

Supplementary Material 12: Video S3. MTScox8-agno-(20-66)-mEGFP is targeted to mitochondria and induced mitochondrial fragmentation. UTA6 cells transfected with MTScox8-agno(20-66) at 24 hpt. Immunofluorescent staining for Tom20 (red) and agnoprotein-mEGFP (green); DNA (blue). Colocalizing voxels are shown in yellow.

References

Abend, J.R., Low, J.A., and Imperiale, M.J. (2010). Global effects of BKV infection on gene expression in human primary kidney epithelial cells. Virology *397*, 73-79.

An, P., Saenz Robles, M.T., Duray, A.M., Cantalupo, P.G., and Pipas, J.M. (2019). Human polyomavirus BKV infection of endothelial cells results in interferon pathway induction and persistence. PLoS pathogens *15*, e1007505.

Assetta, B., De Cecco, M., O'Hara, B., and Atwood, W.J. (2016). JC Polyomavirus Infection of Primary Human Renal Epithelial Cells Is Controlled by a Type I IFN-Induced Response. mBio *7*, e00903-00916.

Atencio, I.A., Shadan, F.F., Zhou, X.J., Vaziri, N.D., and Villarreal, L.P. (1993). Adult mouse kidneys become permissive to acute polyomavirus infection and reactivate persistent infections in response to cellular damage and regeneration. Journal of Virology *67*, 1424-1432.

Bernhoff, E., Gutteberg, T.J., Sandvik, K., Hirsch, H.H., and Rinaldo, C.H. (2008). Cidofovir Inhibits Polyomavirus BK Replication in Human Renal Tubular Cells Downstream of Viral Early Gene Expression. American Journal of tTansplantation *8*, 1413-1422.

Bethge, T., Hachemi, H.A., Manzetti, J., Gosert, R., Schaffner, W., and Hirsch, H.H. (2015). Sp1 sites in the noncoding control region of BK polyomavirus are key regulators of bidirectional viral early and late gene expression. Journal of Virology *89*, 3396-3411.

Bhujabal, Z., Birgisdottir, A.B., Sjottem, E., Brenne, H.B., Overvatn, A., Habisov, S., Kirkin, V., Lamark, T., and Johansen, T. (2017). FKBP8 recruits LC3A to mediate Parkin-independent mitophagy. EMBO Rep *18*, 947-961.

Binggeli, S., Egli, A., Schaub, S., Binet, I., Mayr, M., Steiger, J., and Hirsch, H.H. (2007). Polyomavirus BK-Specific Cellular Immune Response to VP1 and Large T-Antigen in Kidney Transplant Recipients. American Journal of Transplantation *7*, 1131-1139.

Bleicken, S., Hantusch, A., Das, K.K., Frickey, T., and Garcia-Saez, A.J. (2017). Quantitative interactome of a membrane Bcl-2 network identifies a hierarchy of complexes for apoptosis regulation. Nat Commun *8*, 73.

Cam, M., Handke, W., Picard-Maureau, M., and Brune, W. (2010). Cytomegaloviruses inhibit Bak- and Bax-mediated apoptosis with two separate viral proteins. Cell Death Differ *17*, 655-665.

Carswell, S., Resnick, J., and Alwine, J.C. (1986). Construction and characterization of CV-1P cell lines which constitutively express the simian virus 40 agnoprotein: alteration of plaquing phenotype of viral agnogene mutants. Journal of Virology *60*, 415-422.

Cesaro, S., Dalianis, T., Hanssen Rinaldo, C., Koskenvuo, M., Pegoraro, A., Einsele, H., Cordonnier, C., Hirsch, H.H., and Group, E.-. (2018). ECIL guidelines for the prevention, diagnosis and treatment of BK polyomavirus-associated haemorrhagic cystitis in haematopoietic stem cell transplant recipients. The Journal of Antimicrobial Chemotherapy *73*, 12-21.

Chen, Y., Trofe, J., Gordon, J., Autissier, P., Woodle, E.S., and Koralnik, I.J. (2008). BKV and JCV large T antigen-specific CD8(+) T cell response in HLA A*0201(+) kidney transplant recipients with polyomavirus nephropathy and patients with progressive multifocal leukoencephalopathy. Journal of Clinical Virology *42*, 198-202.

Chiu, Y.H., Macmillan, J.B., and Chen, Z.J. (2009). RNA polymerase III detects cytosolic DNA and induces type I interferons through the RIG-I pathway. Cell *138*, 576-591.

Choi, H.J., Park, A., Kang, S., Lee, E., Lee, T.A., Ra, E.A., Lee, J., Lee, S., and Park, B. (2018). Human cytomegalovirus-encoded US9 targets MAVS and STING signaling to evade type I interferon immune responses. Nat Commun *9*, 125.

Cioni, M., Leboeuf, C., Comoli, P., Ginevri, F., and Hirsch, H.H. (2016). Characterization of Immunodominant BK Polyomavirus 9mer Epitope T Cell Responses. American Journal of Transplantation *16*, 1193-1206.

Cioni, M., Mittelholzer, C., Wernli, M., and Hirsch, H.H. (2013). Comparing effects of BK virus agnoprotein and herpes simplex-1 ICP47 on MHC-I and MHC-II expression. Clin Dev Immunol *2013*, 626823.

de Kort, H., Heutinck, K.M., Ruben, J.M., Ede, V.S.A., Wolthers, K.C., Hamann, J., and Ten Berge, I.J.M. (2017). Primary Human Renal-Derived Tubular Epithelial Cells Fail to Recognize and Suppress BK Virus Infection. Transplantation *101*, 1820-1829.

Ding, B., Zhang, L., Li, Z., Zhong, Y., Tang, Q., Qin, Y., and Chen, M. (2017). The Matrix Protein of Human Parainfluenza Virus Type 3 Induces Mitophagy that Suppresses Interferon Responses. Cell Host Microbe *21*, 538-547 e534.

Drachenberg, C.B., Beskow, C.O., Cangro, C.B., Bourquin, P.M., Simsir, A., Fink, J., Weir, M.R., Klassen, D.K., Bartlett, S.T., and Papadimitriou, J.C. (1999). Human polyoma virus in renal allograft biopsies: morphological findings and correlation with urine cytology. Hum Pathol *30*, 970-977.

Edlich, F., Banerjee, S., Suzuki, M., Cleland, M.M., Arnoult, D., Wang, C., Neutzner, A., Tjandra, N., and Youle, R.J. (2011). Bcl-x(L) retrotranslocates Bax from the mitochondria into the cytosol. Cell *145*, 104-116.

Egli, A., Infanti, L., Dumoulin, A., Buser, A., Samaridis, J., Stebler, C., Gosert, R., and Hirsch, H.H. (2009). Prevalence of Polyomavirus BK and JC Infection and Replication in 400 Healthy Blood Donors. The Journal of Infectious Diseases *199*, 837-846.

Farre, J.C., and Subramani, S. (2016). Mechanistic insights into selective autophagy pathways: lessons from yeast. Nat Rev Mol Cell Biol *17*, 537-552.

Fishman, J.A. (2002). BK virus nephropathy--polyomavirus adding insult to injury. The New England Journal of Medicine *347*, 527-530.

Forbes, J.M. (2016). Mitochondria-Power Players in Kidney Function? Trends Endocrinol Metab *27*, 441-442.

Funk, G.A., Gosert, R., Comoli, P., Ginevri, F., Hirsch, H.H. (2008). Polyomavirus BK replication dynamics in vivo and in silico to predict cytopathology and viral clearance in kidney transplants. American Journal of Transplantation *8*, 2368-2377.

Funk, G.A., Steiger, J., and Hirsch, H.H. (2006). Rapid dynamics of polyomavirus type BK in renal transplant recipients. The Journal of Infectious Diseases *193*, 80-87.

Garcia-Sastre, A. (2017). Ten Strategies of Interferon Evasion by Viruses. Cell Host Microbe *22*, 176-184.

Gerits, N., and Moens, U. (2012). Agnoprotein of mammalian polyomaviruses. Virology *432*, 316-326.

Gizzi, A.S., Grove, T.L., Arnold, J.J., Jose, J., Jangra, R.K., Garforth, S.J., Du, Q., Cahill, S.M., Dulyaninova, N.G., Love, J.D.*, et al.* (2018). A naturally occurring antiviral ribonucleotide encoded by the human genome. Nature *558*, 610-614.

Gosert, R., Kardas, P., Major, E.O., and Hirsch, H.H. (2010). Rearranged JC virus noncoding control regions found in progressive multifocal leukoencephalopathy patient samples increase virus early gene expression and replication rate. Journal of Virology *84*, 10448-10456.

Gosert, R., Rinaldo, C.H., Funk, G.A., Egli, A., Ramos, E., Drachenberg, C.B., and Hirsch, H.H. (2008). Polyomavirus BK with rearranged noncoding control region emerge in vivo in renal transplant patients and increase viral replication and cytopathology. J Exp Med *205*, 841-852.

Gou, H., Zhao, M., Xu, H., Yuan, J., He, W., Zhu, M., Ding, H., Yi, L., and Chen, J. (2017). CSFV induced mitochondrial fission and mitophagy to inhibit apoptosis. Oncotarget *8*, 39382-39400.

Goubau, D., Deddouche, S., and Reis e Sousa, C. (2013). Cytosolic sensing of viruses. Immunity *38*, 855-869.

Greenlee, J.E., and Hirsch, H.H. (2017). Polyomaviruses. In Clinical Virology, 4th Edition, A.A.S.f. Microbiology, ed. (asmscience.org), pp. 599-623.

Hartlova, A., Erttmann, S.F., Raffi, F.A., Schmalz, A.M., Resch, U., Anugula, S., Lienenklaus, S., Nilsson, L.M., Kroger, A., Nilsson, J.A.*, et al.* (2015). DNA damage primes the type I interferon system via the cytosolic DNA sensor STING to promote anti-microbial innate immunity. Immunity *42*, 332-343.

Hee, J.S., and Cresswell, P. (2017). Viperin interaction with mitochondrial antiviral signaling protein (MAVS) limits viperin-mediated inhibition of the interferon response in macrophages. PloS one *12*, e0172236.

Hein, J., Boichuk, S., Wu, J., Cheng, Y., Freire, R., Jat, P.S., Roberts, T.M., and Gjoerup, O.V. (2009). Simian virus 40 large T antigen disrupts genome integrity and activates a DNA damage response via Bub1 binding. Journal of Virology *83*, 117-127.

Henriksen, S., Hansen, T., Bruun, J.A., and Rinaldo, C.H. (2016). The Presumed Polyomavirus Viroporin VP4 of Simian Virus 40 or Human BK Polyomavirus Is Not Required for Viral Progeny Release. Journal of Virology *90*, 10398-10413.

Henriksen, S., Mittelholzer, C., Gosert, R., Hirsch, H.H., and Rinaldo, C.H. (2015). Human BK Polyomavirus Plasmid pBKV (34-2) (Dunlop) Contains Mutations Not Found in the Originally Published Sequences. Genome Announcements *3*, 1-2.

Henriksen, S., Tylden, G.D., Dumoulin, A., Sharma, B.N., Hirsch, H.H., and Rinaldo, C.H. (2014). The human fetal glial cell line SVG p12 contains infectious BK polyomavirus. Journal of Virology *88*, 7556-7568.

Heutinck, K.M., Rowshani, A.T., Kassies, J., Claessen, N., van Donselaar-van der Pant, K.A., Bemelman, F.J., Eldering, E., van Lier, R.A., Florquin, S., Ten Berge, I.J.*, et al.* (2012). Viral double-stranded RNA sensors induce antiviral, pro-inflammatory, and pro-apoptotic responses in human renal tubular epithelial cells. Kidney international *82*, 664-675.

Hinson, E.R., and Cresswell, P. (2009a). The antiviral protein, viperin, localizes to lipid droplets via its N-terminal amphipathic alpha-helix. Proc Natl Acad Sci U S A *106*, 20452-20457.

Hinson, E.R., and Cresswell, P. (2009b). The N-terminal amphipathic alpha-helix of viperin mediates localization to the cytosolic face of the endoplasmic reticulum and inhibits protein secretion. J Biol Chem *284*, 4705-4712.

Hirsch, H.H. (2019). Spatio-temporal Virus Surveillance for Severe Acute Respiratory Infections in Resource-limited Settings: How Deep Need We Go? Clinical Infectious Diseases *68*, 1126-1128.

Hirsch, H.H., Drachenberg, C.B., Steiger, J., and Ramos, E. (2006). Polyomavirus-associated nephropathy in renal transplantation: critical issues of screening and management. Advances in Experimental Medicine and biology *577*, 160-173.

Hirsch, H.H., Randhawa, P.S. (2019). BK polyomavirus in solid organ transplantation-Guidelines from the American Society of Transplantation Infectious Diseases Community of Practice. Clinical Transplantation, e13528.

Hirsch, H.H., Yakhontova, K., Lu, M., and Manzetti, J. (2016). BK Polyomavirus Replication in Renal Tubular Epithelial Cells Is Inhibited by Sirolimus, but Activated by Tacrolimus Through a Pathway Involving FKBP-12. American Journal of Transplantation *16*, 821-832.

Hou-Jong, M.H., Larsen, S.H., and Roman, A. (1987). Role of the agnoprotein in regulation of simian virus 40 replication and maturation pathways. Journal of Virology *61*, 937-939.

Imperiale, M.J., and Jiang, M. (2016). Polyomavirus Persistence. Annu Rev Virol *3*, 517-532.

Iwasaki, A., and Medzhitov, R. (2010). Regulation of adaptive immunity by the innate immune system. Science *327*, 291-295.

James A. DeCaprio, M.J.I., Eugene O .Major (2013). Polyomaviruses. In Fields Virology 6th Edition, P.H. David M. Knipe ed. (Wolters Kluwer, Lippincott Williams & Wilkins), pp. 1633-1661.

Jarsch, I.K., Daste, F., and Gallop, J.L. (2016). Membrane curvature in cell biology: An integration of molecular mechanisms. J Cell Biol *214*, 375-387.

Johansen, T., and Lamark, T. (2011). Selective autophagy mediated by autophagic adapter proteins. Autophagy *7*, 279-296.

Jung, J.Y., and Kim, W.J. (2004). Involvement of mitochondrial- and Fas-mediated dual mechanism in CoCl2-induced apoptosis of rat PC12 cells. Neurosci Lett *371*, 85-90.

Kaur, A., Wilhelm, M., Wilk, S., and Hirsch, H.H. (2019). BK polyomavirus-specific antibody and T-cell responses in kidney transplantation: update. Curr Opin Infect Dis *32*, 575-583.

Khan, M., Syed, G.H., Kim, S.J., and Siddiqui, A. (2015). Mitochondrial dynamics and viral infections: A close nexus. Biochim Biophys Acta *1853*, 2822-2833.

Kluge, S.F., Sauter, D., and Kirchhoff, F. (2015). SnapShot: antiviral restriction factors. Cell *163*, 774-774 e771.

Koshiba, T., Yasukawa, K., Yanagi, Y., and Kawabata, S. (2011). Mitochondrial membrane potential is required for MAVS-mediated antiviral signaling. Science signaling *4*, ra7.

Kwon, D., Park, E., Sesaki, H., and Kang, S.J. (2017). Carbonyl cyanide 3-chlorophenylhydrazone (CCCP) suppresses STING-mediated DNA sensing pathway through inducing mitochondrial fission. Biochem Biophys Res Commun *493*, 737-743.

Leboeuf, C., Wilk, S., Achermann, R., Binet, I., Golshayan, D., Hadaya, K., Hirzel, C., Hoffmann, M., Huynh-Do, U., Koller, M.T.*, et al.* (2017). BK Polyomavirus-Specific 9mer CD8 T Cell Responses Correlate With Clearance of BK Viremia in Kidney Transplant Recipients: First Report From the Swiss Transplant Cohort Study. American Journal of Transplantation *17*, 2591-2600.

Leuenberger, D., Andresen, P.A., Gosert, R., Binggeli, S., Strom, E.H., Bodaghi, S., Rinaldo, C.H., and Hirsch, H.H. (2007). Human polyomavirus type 1 (BK virus) agnoprotein is abundantly expressed but immunologically ignored. Clinical and Vaccine Immunology *14*, 959-968.

Leuzinger, K., Naegele, K., Schaub, S., and Hirsch, H.H. (2019). Quantification of plasma BK polyomavirus loads is affected by sequence variability, amplicon length, and non-encapsidated viral DNA genome fragments. Journal of Clinical Virology *121*, 104210.

Li, T., and Chen, J.Z. (2018). The cGAS-cGAMP-STING pathway connects DNA damage to inflammation, senescence, and cancer. J Exp Med *215*, 1287-1299.

Liu, S., Cai, X., Wu, J., Cong, Q., Chen, X., Li, T., Du, F., Ren, J., Wu, Y.T., Grishin, N.V.*, et al.* (2015). Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation. Science *347*, aaa2630-2631.

Low, J., Humes, H.D., Szczypka, M., and Imperiale, M. (2004). BKV and SV40 infection of human kidney tubular epithelial cells in vitro. Virology *323*, 182-188.

Lynn W. Enquist, V.R.R. (2013). Virology: From Contagium Fluidum to Virome. In Fields Virology 6th Edition, P.H. David M. Knipe ed. (Wolters Kluwer, Lippincott Williams & Wilkins), pp. 1-20.

Mandic, L., Miller, M.S., Coulter, C., Munshaw, B., and Hertel, L. (2009). Human cytomegalovirus US9 protein contains an N-terminal signal sequence and a C-terminal mitochondrial localization domain, and does not alter cellular sensitivity to apoptosis. The Journal of General Virology *90*, 1172-1182.

McFadden, M.J., Gokhale, N.S., and Horner, S.M. (2017). Protect this house: cytosolic sensing of viruses. Current Opinion in Virology *22*, 36-43.

McMahon, H.T., and Gallop, J.L. (2005). Membrane curvature and mechanisms of dynamic cell membrane remodelling. Nature *438*, 590-596.

McNab, F., Mayer-Barber, K., Sher, A., Wack, A., and O'Garra, A. (2015). Type I interferons in infectious disease. Nat Rev Immunol *15*, 87-103.

Narendra, D., Kane, L.A., Hauser, D.N., Fearnley, I.M., and Youle, R.J. (2010). p62/SQSTM1 is required for Parkin-induced mitochondrial clustering but not mitophagy; VDAC1 is dispensable for both. Autophagy *6*, 1090-1106.

Narendra, D., Tanaka, A., Suen, D.F., and Youle, R.J. (2008). Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. J Cell Biol *183*, 795-803.

Ng, S.C., Mertz, J.E., Sanden-Will, S., and Bina, M. (1985). Simian virus 40 maturation in cells harboring mutants deleted in the agnogene. J Biol Chem *260*, 1127-1132.

Panou, M.M., Prescott, E.L., Hurdiss, D.L., Swinscoe, G., Hollinshead, M., Caller, L.G., Morgan, E.L., Carlisle, L., Muller, M., Antoni, M.*, et al.* (2018). Agnoprotein Is an Essential Egress Factor during BK Polyomavirus Infection. International journal of molecular sciences *19*, 1-16.

Pastrana, D.V., Brennan, D.C., Cuburu, N., Storch, G.A., Viscidi, R.P., Randhawa, P.S., and Buck, C.B. (2012). Neutralization serotyping of BK polyomavirus infection in kidney transplant recipients. PLoS pathogens *8*, e1002650.

Ramos, E., Drachenberg, C.B., Wali, R., and Hirsch, H.H. (2009). The decade of polyomavirus BK-associated nephropathy: state of affairs. Transplantation *87*, 621-630.

Ribeiro, A., Wornle, M., Motamedi, N., Anders, H.J., Grone, E.F., Nitschko, H., Kurktschiev, P., Debiec, H., Kretzler, M., Cohen, C.D.*, et al.* (2012). Activation of innate immune defense mechanisms contributes to polyomavirus BK-associated nephropathy. Kidney International *81*, 100-111.

Rinaldo, C.H., Traavik, T., and Hey, A. (1998). The agnogene of the human polyomavirus BK is expressed. Journal of Virology *72*, 6233-6236.

Sami Saribas, A., Abou-Gharbia, M., Childers, W., Sariyer, I.K., White, M.K., and Safak, M. (2013). Essential roles of Leu/Ile/Phe-rich domain of JC virus agnoprotein in dimer/oligomer formation, protein stability and splicing of viral transcripts. Virology *443*, 161-176.

Saribas, A.S., Coric, P., Bouaziz, S., and Safak, M. (2018). Expression of novel proteins by polyomaviruses and recent advances in the structural and functional features of agnoprotein of JC virus, BK virus, and simian virus 40. J Cell Physiol, 1-21.

Schneider, W.M., Chevillotte, M.D., and Rice, C.M. (2014). Interferon-stimulated genes: a complex web of host defenses. Annu Rev Immunol *32*, 513-545.

Seo, J.Y., Yaneva, R., Hinson, E.R., and Cresswell, P. (2011). Human cytomegalovirus directly induces the antiviral protein viperin to enhance infectivity. Science *332*, 1093-1097.

Shah, K., Daniel, R., Madden, D., and Stagno, S. (1980). Serological investigation of BK papovavirus infection in pregnant women and their offspring. Infect Immun *30*, 29-35.

Shen, H., Pirruccello, M., and De Camilli, P. (2012). SnapShot: membrane curvature sensors and generators. Cell *150*, 1300, 1300 e1301-1302.

Simmonds, P., Adams, M.J., Benko, M., Breitbart, M., Brister, J.R., Carstens, E.B., Davison, A.J., Delwart, E., Gorbalenya, A.E., Harrach, B.*, et al.* (2017). Consensus statement: Virus taxonomy in the age of metagenomics. Nature Reviews Microbiology *15*, 161-168.

Solis, M., Velay, A., Porcher, R., Domingo-Calap, P., Soulier, E., Joly, M., Meddeb, M., Kack-Kack, W., Moulin, B., Bahram, S.*, et al.* (2018). Neutralizing Antibody-Mediated

Response and Risk of BK Virus-Associated Nephropathy. Journal of the American Society of Nephrology *29*, 326-334.

Suzuki, T., Orba, Y., Okada, Y., Sunden, Y., Kimura, T., Tanaka, S., Nagashima, K., Hall, W.W., and Sawa, H. (2010). The human polyoma JC virus agnoprotein acts as a viroporin. PLoS pathogens *6*, e1000801.

Takeuchi, O., and Akira, S. (2010). Pattern recognition receptors and inflammation. Cell *140*, 805-820.

Unterstab, G., Gosert, R., Leuenberger, D., Lorentz, P., Rinaldo, C.H., and Hirsch, H.H. (2010). The polyomavirus BK agnoprotein co-localizes with lipid droplets. Virology *399*, 322-331.

Unterstab, G., Manzetti, J., and Hirsch, H.H. (2013). Corrigendum to 'The polyomavirus BK agnoprotein co-localizes with lipid droplets' [Virology 399 (2) (2010) 322-331]. Virology *441*, 197-199.

Virgin, H.W. (2014). The virome in mammalian physiology and disease. Cell *157*, 142-150.

Virgin, H.W., Wherry, E.J., and Ahmed, R. (2009). Redefining chronic viral infection. Cell *138*, 30-50.

Wang, X., Hinson, E.R., and Cresswell, P. (2007). The interferon-inducible protein viperin inhibits influenza virus release by perturbing lipid rafts. Cell Host Microbe *2*, 96-105.

West, A.P., and Shadel, G.S. (2017). Mitochondrial DNA in innate immune responses and inflammatory pathology. Nat Rev Immunol *17*, 363-375.

Zevini, A., Olagnier, D., and Hiscott, J. (2017). Crosstalk between Cytoplasmic RIG-I and STING Sensing Pathways. Trends Immunol *38*, 194-205.

A

AGN

Merge | Colocalization | Colocalization, Tom20

agn25D39E

B

Number of cells (%)

Mitochondria at 48hpi

AGN | agn25D39E

<0.001 | <0.001

Elongated | Intermediate | Fragmented

C

DNA | Calr | Agno | Tom20

Colocalization

Calr + Tom20 | Calr + Agno | Tom20 + Agno | Calr + Tom20 + Agno

Colocalization + Agno

D

AGN

24hpi | 36hpi | 48hpi | 72hpi

LTag

Agno

Tom20

DNA

Merge

Tom20

A

B

A

AGN agn25D39E

Tom20 Agno p62 Merge

AGN

agn25D39E

p62 autophagosomes per cell

<0.001

AGN
agn25D39E

48 72

Hours post infection

B

Tom20 Agno p62 Merge

24h + tet

24h − tet

48h + tet

48h − tet

p62 autophagosomes per cell

<0.001

+ tet
− tet

24 48

Hours post induction

C

Infected Non-Infected

Infected Non-Infected

Agno

Tom20

p62

Merge