Calculation of fungal and bacterial inorganic nitrogen immobilization rates in soil

Xiaobo Li[1,2], Hongbo He[3], Xudong Zhang[3], Caner Kazanci[4,5], Zhian Li[1,2], Magdalena Necpalova[6], Qianqian Ma[1,2]*

[1] *Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China.*

[2] *Center of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Guangzhou 510650, China.*

[3] *Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China.*

[4] *Department of Mathematics, University of Georgia, Athens, GA 30602, USA.*

[5] *College of Engineering, University of Georgia, Athens, GA 30602, USA*

[6] *School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Ireland.*

*Corresponding author:

Q, Ma, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China. *E-mail*: maqianqian@scbg.ac.cn

Abstract

Microbial inorganic nitrogen (N) immobilization is an important mechanism in the retention of N in soils. However, as a result of the high diversity and complexity of soil microorganisms, there is still no effective approach to measuring the respective immobilization rates of inorganic N by fungi and bacteria, which are the two dominant microbial communities in soils. We propose a mathematical framework, combining the experimentally measurable gross inorganic N immobilization rate and proxies for fungal and bacterial inorganic N immobilization rates, to quantify the respective immobilization rates of inorganic N by fungal and bacterial communities in soil. Our approach will help to unravel the mechanisms of N retention in soils.

Keywords:

Amino sugars; fungi; bacteria; microbial inorganic N immobilization; ^{15}N; stable isotope probing

The microbial immobilization of inorganic nitrogen (N) has a vital role in controlling the size of the soil inorganic N pool and is therefore an important mechanism for the retention of N in ecosystems (Davidson et al 1992, Stark and Hart 1997, Zhang et al 2013, Zogg et al 2000). Through this immobilization process, inorganic N in soil is converted to microbial biomass N and subsequently re-mineralized or converted to stable organic N, eventually reducing the risk of N losses from soil (Recous et al 1990, Tahovská et al 2013, Zhang et al 2019). As the dominant microorganisms in soil,

fungi and bacteria are probably the main participants in inorganic N immobilization

(Bottomley et al 2012, Boyle et al 2008, Myrold and Posavatz 2007). Given the

distinct physiologies, morphologies, lifestyles and quantities of these two microbial

groups in soil (Lauber et al 2008, Rousk and Bååth 2011, Six et al 2006, Waring et al

2013), the relative importance of fungi and bacteria in soil inorganic N

immobilization is likely to be unequal (Bottomley et al 2012, Li et al 2019, Myrold

and Posavatz 2007). However, as a result of the high diversity and complexity of soil

microorganisms, quantifying the respective rates of immobilization of inorganic N by

fungal and bacterial communities in soil is challenging (Fierer 2017, Li et al 2019, Li

et al 2020), although the gross inorganic N immobilization rate can be measured using

well-established ^{15}N isotope techniques (e.g.,   the ^{15}N pool dilution method) (Cheng

et al 2017, Murphy et al 2003).

Amino sugars, which are important constituents of microbial cell walls, have different

origins in microorganisms. Among the amino sugars identified in microorganisms,

muramic acid (MurN) originates exclusively from bacterial peptidoglycan, whereas

glucosamine (GlcN) is mainly in the form of chitin in fungal cell walls (Amelung

2001, Parsons 1981, Zhang and Amelung 1996). Based on their microbial source

specificity, stable isotope probing based on amino sugars (^{15}N-AS-SIP) has been

developed to disentangle the immobilization processes of inorganic N by fungi and

bacteria in soils (He et al 2006, He et al 2011a, He et al 2011b, Liang and Balser 2010,

Reay et al 2019a, Reay et al 2019b).

This approach has now been extended to indicate the inorganic N immobilization

rates of fungal and bacterial communities in soils (Li et al 2019, Li et al 2020). More

specifically, given the relatively long persistence of amino sugars in soils (mean

turnover time >2 years, much longer than that of the living microorganisms) (Derrien

and Amelung 2011, Glaser et al 2006, Liu et al 2016), the newly formed ^{15}N-labeled

amino sugars are considered to be stable in soil even after cell death (Glaser et al 2004,

Gunina et al 2017). The fungal-derived ^{15}N-GlcN and bacterial-derived ^{15}N-MurN

synthesis rates within a short period of incubation after ^{15}N tracer addition have

therefore been used as proxies for the rates of immobilization of inorganic N by fungi

and bacteria, respectively (Li et al 2019, Li et al 2020). However, mainly as a result of

the variation in the composition of tissues of massive microbial species, but also

within each species under different growth conditions, the actual contents of GlcN

and MurN in the respective biomasses of fungi and bacteria in soil are almost

unobtainable (Appuhn and Joergensen 2006, Engelking et al 2007, Glaser et al 2004,

Joergensen 2018). It is also still unclear how fast do the cell N-containing components

turn over intracellularly and extracellularly in soil (Dippold et al 2019, Engelking et al

2007, Gunina et al 2017). As a consequence, converting the synthesis rates of

^{15}N-labeled amino sugars specific for fungi and bacteria to the actual inorganic N

immobilization rates in soil is challenging.

To bypass this intractable problem, we propose a mathematical framework to estimate

the conversion coefficients between fungal and bacterial inorganic N immobilization

rates and their respective proxies by combining the gross inorganic N immobilization

rate with proxies for the respective inorganic N immobilization rates of fungi and

bacteria. In this way, we can obtain the respective immobilization rates of inorganic N

by fungal and bacterial communities in soil.

Calculation of fungal and bacterial inorganic N immobilization rates

Our proposed calculation is based on the assumption that fungi and bacteria are the

dominant participants in soil microbial inorganic N immobilization. If both the gross

inorganic N immobilization rate and the proxies for inorganic N immobilization rates

of fungi and bacteria have been measured on n soil samples ($n \geq 2$), then the respective

immobilization rates of inorganic N by fungal and bacterial communities can be

calculated.

The measured variables are:

$G = \begin{bmatrix} G_1 \\ G_2 \\ \vdots \\ G_n \end{bmatrix}$: gross microbial inorganic N immobilization rates for n samples (mg N

$\text{kg}^{-1}\ \text{day}^{-1}$);

$F = \begin{bmatrix} F_1 \\ F_2 \\ \vdots \\ F_n \end{bmatrix}$: fungal-derived ^{15}N-GlcN synthesis rates for n samples (mg N $\text{kg}^{-1}\ \text{day}^{-1}$);

$B = \begin{bmatrix} B_1 \\ B_2 \\ \vdots \\ B_n \end{bmatrix}$: bacterial-derived ^{15}N-MurN synthesis rates for n samples (mg N kg^{-1}

day^{-1}).

The two parameters to be estimated are:

K_F: the conversion coefficient from the fungal-derived ^{15}N-GlcN synthesis rate to the

fungal inorganic N immobilization rate;

K_B: the conversion coefficient from the bacterial-derived [15]N-MurN synthesis rate to

the bacterial inorganic N immobilization rate.

Using the [15]N-labeled amino sugars synthesis rates and conversion coefficients, the

estimated fungal and bacterial inorganic N immobilization rates (mg N kg^{-1} day^{-1}) are,

respectively, calculated as:

$$R_F = K_F \times F \qquad (1)$$

111   and

$$R_B = K_B \times B \qquad (2)$$

113   Their sum is therefore the estimated gross microbial inorganic N immobilization rate

114   (mg N kg^{-1} day^{-1}):

$$\hat{G} = R_F + R_B = K_F \times F + K_B \times B$$

115   The measured gross microbial inorganic N immobilization rate results are included in

116   the equation:

$$G = \hat{G} + e = K_F \times F + K_B \times B + e$$

117   where e is the estimation error. This equation can be rewritten in a matrix format:

$$G = \begin{bmatrix} F & B \end{bmatrix} \begin{bmatrix} K_F \\ K_B \end{bmatrix} + e$$

118   Alternatively,

$$\begin{bmatrix} G_1 \\ G_2 \\ \vdots \\ G_n \end{bmatrix} = \begin{bmatrix} F_1 & B_1 \\ F_2 & B_2 \\ \vdots & \vdots \\ F_n & B_n \end{bmatrix} \begin{bmatrix} K_F \\ K_B \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_n \end{bmatrix}$$

120   If we let $K = \begin{bmatrix} K_F \\ K_B \end{bmatrix}$ and $X = \begin{bmatrix} F_1 & B_1 \\ F_2 & B_2 \\ \vdots & \vdots \\ F_n & B_n \end{bmatrix}$, we obtain:

$$G = X K + e$$

121   The least-squares estimators that minimize the sum of the squared residuals are given

122   in the following (see Appendix for the detailed derivation) (Wackerly et al 2014):

123                           $\hat{K} = (X^T X)^{-1} X^T G$ (3)

To illustrate how this approach works, we calculated the soil nitrate (NO_3^-)

immobilization rates of fungi and bacteria using the gross NO_3^- immobilization rates

reported by Zhang et al (2013) and the [15]N-labeled amino sugars synthesis rates

reported by Li et al (2019). Both studies studied the effect of land conversion from

forest to agriculture on the soil NO_3^- immobilization in subtropical zones of China.

Ideally, the gross NO_3^- immobilization rates and the [15]N-labeled amino sugars

synthesis rates should be measured under the same experimental conditions such as

sampling sites. Due to the unavailability of such data, we roughly treat selected

studies as being conducted at the same sites. Therefore, the results in Table 1 are

presented as an illustrative example, rather than as reliable estimates. For simplicity,

only the mean rates for forest and agricultural lands were used in this example ($n = 2$).

The conversion coefficients were obtained by substituting the measured gross NO_3^-

immobilization rates and the [15]N-labeled amino sugars synthesis rates into Equation

(3). The fungal and bacterial NO_3^- immobilization rates were then calculated using

Equations (1) and (2). A summary of measured data and estimated values is provided

in Table 1.

The results showed that the NO_3^- immobilization rates of fungi in woodland and

agricultural soils were about 8.4 and four times those of bacteria, indicating that fungi

dominated the microbial NO_3^- immobilization in the studied soil (Table 1).

Conversion to agricultural use led to decreases in the fungal and bacterial NO_3^-

immobilization rates of 0.34 and 0.03 mg N kg^{-1} day^{-1}, respectively, which suggests

that the decrease in the fungal NO_3^- immobilization rate dominates the decrease in the

gross soil microbial NO_3^- immobilization caused by the land use change.

Table 1. An illustration of the method of calculating soil fungal and bacterial NO_3^-

immobilization rates under different land use scenarios. The gross NO_3^-

immobilization rates (G) were obtained from Zhang et al (2013). The synthesis rates

of fungal-derived ^{15}N-GlcN (F) and bacterial-derived ^{15}N-MurN (B) were obtained

from Li et al (2019) (see Table S1 for detailed calculations). These values are

presented as an illustrative example, rather than as reliable estimates.

Land use	G	F	B	K_F	K_B	R_F	R_B
	mg N kg^{-1} day^{-1}					mg N kg^{-1} day^{-1}	
Woodland	0.47	0.0303	0.0022	13.78	23.83	0.42	0.05
Agriculture	0.10	0.0057	0.0009	13.78	23.83	0.08	0.02

Note: K_F and K_B are the conversion coefficients between F, B and the NO_3^-

immobilization rates of fungi (R_F) and bacteria (R_B), respectively.

Advantages and limitations of this approach

Understanding the microbially mediated N cycling processes in soil is central to

unraveling soil N retention mechanisms and has ramifications for reducing N losses

and managing ecosystem productivity. As a result of the high diversity and

complexity of microbial communities, quantifying the process rates of different

microbial groups has been a great challenge, especially in soil (Bardgett and Van Der

Putten 2014, Fierer 2017, Stres and Tiedje 2006). Our approach provides an effective

way to mathematically, rather than mechanically, quantify the relative importance of

fungal and bacterial communities in soil inorganic N immobilization. It circumvents

the bottleneck of directly measuring or estimating the inorganic N immobilization

rates of fungi and bacteria in soil. The experimentally accessible gross inorganic N

immobilization rate and proxies of fungal and bacterial inorganic N immobilization

rates are used to estimate the conversion coefficients between fungal and bacterial

inorganic N immobilization rates and their respective proxies. The conversion

coefficients obtained inherently take into account both the actual contents of GlcN

and MurN in the respective biomasses of fungi and bacteria and the turnover of cell

N-containing components in the studied soil. Because the rationale and mathematical

derivation are universal, our method may also be applicable to other environmental

systems, such as freshly colonized organic substrates (Appuhn and Joergensen 2006).

This approach relies on the simplifying assumption that only fungi and bacteria are

involved in soil microbial inorganic N immobilization. This assumption may not quite

hold true, because Archaea may also contribute to inorganic N immobilization

(Laughlin et al 2009). Considering that Archaea contain GlcN, but not MurN

(Joergensen 2018), the contribution of Archaea, if any, is included in the fungal

inorganic N immobilization rates by adopting our approach. Nevertheless, considering

that Archaea account for less than <1% of the soil microbial biomass (Fierer 2017),

the errors caused by this assumption are probably trivial.

183 Conclusions

We propose a mathematical approach that combines the mechanically accessible gross

inorganic N immobilization rate and proxies for fungal and bacterial inorganic N

immobilization rates (measured by ^{15}N-AS-SIP) to quantify the inorganic N

immobilization rates of fungal and bacterial communities in soil. This approach,

although not without its limitations, allows us for the first time to disentangle the

actual contribution of fungi and bacteria to the immobilization of N-containing

substrates in soil. Promisingly, integrating both fungal and bacterial inorganic N

immobilization rates into terrestrial ecosystem models (e.g., microbial models) will

improve our ability to understand, predict and manage the N retention capacity in

soils under different scenarios (Waring et al 2013).

194 Acknowledgments

This study was supported by the National Natural Science Foundation of China

(41977097, 41401279, 31600392 and 41630862), the Natural Science Foundation of

Guangdong Province (2019A1515012067 and 2016A030310013), R & D program of

Guangdong Provincial Department of Science and Technology (2018B030324003),

China Soil Microbiome Initiative of the Chinese Academy of Sciences

(XDB15040200), the Research Fund of State Key Laboratory of Soil and Sustainable

Agriculture, Nanjing Institute of Soil Science, Chinese Academy of Sciences

(Y412201422), and National Key Research and Development Program of China

(2017YFC0506305). H. He and X. Zhang acknowledge the CAS Interdisciplinary

Innovation Team. X. Li appreciates China Scholarship Council for providing funds to

support his study in Switzerland.

References

Amelung W (2001). Methods using amino sugars as markers for microbial residues in soil.

Assessment methods for soil carbon. Lewis Publishers Boca Raton. pp 233-272.

Appuhn A, Joergensen RG (2006). Microbial colonisation of roots as a function of plant

species. *Soil Biology and Biochemistry* **38**: 1040-1051.

Bardgett RD, Van Der Putten WH (2014). Belowground biodiversity and ecosystem functioning.

Nature **515**: 505-511.

Bottomley PJ, Taylor AE, Myrold DD (2012). A consideration of the relative contributions of

different microbial subpopulations to the soil N cycle. *Frontiers in Microbiology* **3**: 373.

Boyle SA, Yarwood RR, Bottomley PJ, Myrold DD (2008). Bacterial and fungal contributions to

soil nitrogen cycling under Douglas fir and red alder at two sites in Oregon. *Soil Biology and*

Biochemistry **40**: 443-451.

Cheng Y, Wang J, Wang J, Chang SX, Wang S (2017). The quality and quantity of exogenous

organic carbon input control microbial NO3− immobilization: A meta-analysis. *Soil Biology and*

Biochemistry **115**: 357-363.

Davidson EA, Hart SC, Firestone MK (1992). Internal cycling of nitrate in soils of a mature

coniferous forest. *Ecology* **73**: 1148-1156.

Derrien D, Amelung W (2011). Computing the mean residence time of soil carbon fractions

using stable isotopes: impacts of the model framework. *Eur J Soil Sci* **62**: 237-252.

Dippold MA, Gunina A, Apostel C, Boesel S, Glaser B, Kuzyakov Y (2019). Metabolic tracing

unravels pathways of fungal and bacterial amino sugar formation in soil. *Eur J Soil Sci* **70**:

421-430.

Engelking B, Flessa H, Joergensen RG (2007). Shifts in amino sugar and ergosterol contents

after addition of sucrose and cellulose to soil. *Soil Biology and Biochemistry* **39**: 2111-2118.

Fierer N (2017). Embracing the unknown: disentangling the complexities of the soil

microbiome. *Nature Reviews Microbiology* **15**: 579.

Glaser B, Turrión Ma-B, Alef K (2004). Amino sugars and muramic acid—biomarkers for soil

microbial community structure analysis. *Soil Biology and Biochemistry* **36**: 399-407.

Glaser B, Millar N, Blum H (2006). Sequestration and turnover of bacterial- and fungal-derived

carbon in a temperate grassland soil under long-term elevated atmospheric pCO2. *Global*

Change Biology **12**: 1521-1531.

Gunina A, Dippold M, Glaser B, Kuzyakov Y (2017). Turnover of microbial groups and cell

components in soil: 13C analysis of cellular biomarkers. *Biogeosciences* **14**: 271-283.

He H, Xie H, Zhang X (2006). A novel GC/MS technique to assess 15N and 13C incorporation

into soil amino sugars. *Soil Biology and Biochemistry* **38**: 1083-1091.

He H, Li X, Zhang W, Zhang X (2011a). Differentiating the dynamics of native and newly

immobilized amino sugars in soil frequently amended with inorganic nitrogen and glucose. *Eur*

J Soil Sci **62**: 144-151.

He H, Zhang W, Zhang X, Xie H, Zhuang J (2011b). Temporal responses of soil

microorganisms to substrate addition as indicated by amino sugar differentiation. *Soil Biology*

and Biochemistry **43**: 1155-1161.

Joergensen RG (2018). Amino sugars as specific indices for fungal and bacterial residues in

soil. *Biol Fertil Soils* **54**: 559-568.

Lauber CL, Strickland MS, Bradford MA, Fierer N (2008). The influence of soil properties on

the structure of bacterial and fungal communities across land-use types. *Soil Biology and*

Biochemistry **40**: 2407-2415.

Laughlin R, Rütting T, Müller C, Watson C, Stevens R (2009). Effect of acetate on soil

respiration, N2O emissions and gross N transformations related to fungi and bacteria in a

grassland soil. *Applied Soil Ecology* **42**: 25-30.

Li X, He H, Zhang X, Yan X, Six J, Cai Z *et al* (2019). Distinct responses of soil fungal and

bacterial nitrate immobilization to land conversion from forest to agriculture. *Soil Biology and*

Biochemistry **134**: 81-89.

Li X, Li Z, Zhang X, Xia L, Zhang W, Ma Q *et al* (2020). Disentangling immobilization of nitrate

by fungi and bacteria in soil to plant residue amendment. *Geoderma* **374**: 114450.

Liang C, Balser TC (2010). Mass spectrometric characterization of amino sugar aldononitrile

acetate derivatives used for isotope enrichment assessment of microbial residues. *Soil Biology*

and Biochemistry **42**: 904-909.

Liu X, Hu G, He H, Liang C, Zhang W, Bai Z *et al* (2016). Linking microbial immobilization of

fertilizer nitrogen to in situ turnover of soil microbial residues in an agro-ecosystem. *Agriculture,*

Ecosystems and Environment **229**: 40-47.

Murphy DV, Recous S, Stockdale EA, Fillery IRP, Jensen LS, Hatch DJ *et al* (2003). Gross

nitrogen fluxes in soil : theory, measurement and application of 15N pool dilution techniques.

Advances in Agronomy. Academic Press. pp 69-118.

Myrold DD, Posavatz NR (2007). Potential importance of bacteria and fungi in nitrate

assimilation in soil. *Soil Biology and Biochemistry* **39**: 1737-1743.

Parsons JW (1981). Chemistry and distribution of amino sugars in soils and soil organisms. In:

E.A. Paul JNL (ed). *Soil Biochemistry.* Marcel Dekker: New York. pp 197-227.

Reay MK, Charteris AF, Jones DL, Evershed RP (2019a). 15N-amino sugar stable isotope

probing (15N-SIP) to trace the assimilation of fertiliser-N by soil bacterial and fungal

communities. *Soil Biology and Biochemistry* **138**: 107599.

Reay MK, Knowles TDJ, Jones DL, Evershed RP (2019b). Development of Alditol Acetate

Derivatives for the Determination of 15N-Enriched Amino Sugars by Gas

Chromatography–Combustion–Isotope Ratio Mass Spectrometry. *Analytical Chemistry* **91**:

3397-3404.

Recous S, Mary B, Faurie G (1990). Microbial immobilization of ammonium and nitrate in

cultivated soils. *Soil Biology and Biochemistry* **22**: 913-922.

Rousk J, Bååth E (2011). Growth of saprotrophic fungi and bacteria in soil. *FEMS Microbiol*

Ecol **78**: 17-30.

Six J, Frey SD, Thiet RK, Batten KM (2006). Bacterial and fungal contributions to carbon

sequestration in agroecosystems. *Soil Science Society of America Journal* **70**: 555-569.

Stark JM, Hart SC (1997). High rates of nitrification and nitrate turnover in undisturbed

coniferous forests. *Nature* **385**: 61-64.

Stres B, Tiedje JM (2006). New frontiers in soil microbiology: how to link structure and function

of microbial communities? *Nucleic Acids and Proteins in Soil.* Springer. pp 1-22.

Tahovská K, Kaňa J, Bárta J, Oulehle F, Richter A, Šantrůčková H (2013). Microbial N

immobilization is of great importance in acidified mountain spruce forest soils. *Soil Biology and*

Biochemistry **59**: 58-71.

Wackerly D, Mendenhall W, Scheaffer RL (2014). *Mathematical statistics with applications.*

Cengage Learning.

Waring BG, Averill C, Hawkes CV (2013). Differences in fungal and bacterial physiology alter

soil carbon and nitrogen cycling: insights from meta-analysis and theoretical models. *Ecology*

Letters **16**: 887-894.

Zhang J, Zhu T, Meng T, Zhang Y, Yang J, Yang W *et al* (2013). Agricultural land use affects

nitrate production and conservation in humid subtropical soils in China. *Soil Biology and*

Biochemistry **62**: 107-114.

Zhang W, Zhang X, Bai E, Cui Y, He H, Zhang X (2019). The strategy of microbial utilization of

the deposited N in a temperate forest soil. *Biol Fertil Soils*: 1-9.

Zhang X, Amelung W (1996). Gas chromatographic determination of muramic acid,

glucosamine, mannosamine, and galactosamine in soils. *Soil Biology and Biochemistry* **28**:

1201-1206.

Zogg GP, Zak DR, Pregitzer KS, Burton AJ (2000). Microbial immobilization and the retention

of anthropogenic nitrate in a northern hardwood forest. *Ecology* **81**: 1858-1866.

348