ARTICLE

Clinal and seasonal change are correlated in *Drosophila melanogaster*

natural populations

Running title: Correlation between clinal and seasonal change

Murillo F. Rodrigues[1,2*], Maria D. Vibranovski[1], Rodrigo Cogni[3]

[1] Department of Genetics and Evolutionary Biology, Institute of Biosciences, University of
Sao Paulo, Sao Paulo, SP, Brazil
[2] Present address: Institute of Ecology and Evolution, University of Oregon, Eugene, OR,
USA
[3] Department of Ecology, Institute of Biosciences, University of Sao Paulo, Sao Paulo, SP,
Brazil
12

* Corresponding author:

Murillo F. Rodrigues

Institute of Ecology and Evolution

University of Oregon

272 Onyx Bridge

5289 University of Oregon

Eugene, OR 97403-5289.

Author contributions

M.F.R. and R.C. designed the research. M.F.R. performed the research and analyzed the data.

M.F.R., M.D.V. and R.C. discussed the results and conclusions. M.F.R. wrote the manuscript

with input from M.D.V and R.C.

Acknowledgements

We would like to thank all the members of the Cogni Lab for input and support throughout

the development of this work. We thank Diogo Meyer, Paul Schmidt, Reinaldo Azevedo and

Peter Ralph for discussions and three anonymous reviewers for comments on the manuscript.

Funding for this work was provided by São Paulo Research Foundation (FAPESP)

(13/25991-0 and 17/02206-6 to RC, 15/20844-4 to MDV, 16/01354-9 and 17/06374-0 to

MFR), and CNPq (307015/2015-7 and 307447/2018-9 to RC), and a Newton Advanced

Fellowship from the Royal Society to RC.

34   Data accessibility

All the data used in this project are available on the NCBI Short Read Archive (BioProject

accession numbers PRJNA256231 and PRJNA308584, and NCBI Sequence Read Archive

SRA012285.16).

Abstract

Spatial and seasonal variation in the environment are ubiquitous. Environmental

heterogeneity can affect natural populations and lead to covariation between environment and

allele frequencies. *Drosophila melanogaster* is known to harbor polymorphisms that change

both with latitude and seasons. Identifying the role of selection in driving these changes is not

trivial, because non-adaptive processes can cause similar patterns. Given the environment

changes in similar ways across seasons and along the latitudinal gradient, one promising

approach may be to look for parallelism between clinal and seasonal change. Here, we test

whether there is a genome-wide correlation between clinal and seasonal change, and whether

the pattern is consistent with selection. Allele frequency estimates were obtained from pooled

samples from seven different locations along the east coast of the US, and across seasons

within Pennsylvania. We show that there is a genome-wide correlation between clinal and

seasonal variation, which cannot be explained by linked selection alone. This pattern is

stronger for coding than intergenic regions, consistent with natural selection. We derive a

way to biologically interpret these correlations, and we show that around 3.7% of the

common, autosomal variants could be under parallel seasonal and spatial selection. Our

results highlight the contribution of natural selection in driving fluctuations in allele

frequencies in *D. melanogaster*.

Introduction

Species occur in environments that vary both in space and time (Ewing 1979; Cardini et al. 2007; Dionne et al. 2007; Hancock et al. 2008; Zuther et al. 2012; Campitelli and Stinchcombe 2013; Kooyers et al. 2015). Populations may adapt to the local conditions of the environment in which they occur, resulting in covariation between traits and space (Endler 1977; Barton 1983; Barton 1999; Kawecki and Ebert 2004). Similarly, predictable changes in the environment through time can lead covariation between relevant traits and time (Levene, 1953; Ewing 1979). Although correlations between environment and traits (either in time or space) are indicative of selection, these patterns can be produced by non-adaptive processes such as migration, isolation by distance and range expansion (Wright 1943; Vasemägi 2006; Excoffier et al. 2009; Duchen et al. 2013; Bergland et al. 2016). It is not trivial to identify the role of selection in diversifying traits, but a promising approach might be to jointly model changes across space and time.

Drosophila melanogaster is a uniquely suited to study both spatial and temporal adaptation. These sub-Saharan flies recently invaded most of the world (David and Capy 1988), and adaptations at the phenotypic and genotypic levels evolved in response to the colonization of new habitats (Mettler et al. 1977; Knibb 1982; Oakeshott et al. 1982; David and Capy 1988; Schmidt et al. 2000; de Jong and Bochdanovits 2003; Sezgin et al. 2004). Many traits, polymorphisms and inversions were observed to covary with latitude (also called clinal) in natural fly populations (Hoffmann et al. 2002; Hoffmann and Weeks 2007; Turner et al. 2008; Paaby et al. 2010; Yukilevich et al. 2010; Reinhardt et al. 2014; Schrider et al. 2016). For instance, flies from colder environments are darker (David et al. 1985), bigger (Arthur et al. 2008) and show higher incidence of reproductive diapause than flies from lower latitudes (Schmidt et al. 2005).

In higher latitudes, fly populations started to experience dramatic cyclical changes in

the environment through seasons. Given these *D. melanogaster* have multiple generations per

year, differential fitness across seasons could theoretically lead to temporal adaptations

(Levene 1953; Ewing 1979). Traits that favor rapid reproduction in the summer can be

particularly different than those which favor endurance in the winter (Behrman et al. 2015).

Concordant with this hypothesis, chromosomal inversions in *D. pseudoobscura* were

observed to cycle with seasons (Dobzhansky 1943). In *D. melanogaster*, flies collected in the

spring are more tolerant to stress (Behrman et al. 2015), show higher diapause inducibility

(Schmidt and Conde 2006), have increased immune function (Behrman et al. 2018) and have

different cuticular hydrocarbon profiles than those collected in the fall (Rajpurohit et al.

2017). Genome-wide analyses have identified polymorphisms and inversions that oscillate in

seasonal timescales in several localities in the United States and Europe (Bergland et al.

2014; Kapun et al. 2016; Machado et al. 2018). However, a recent analysis suggested

seasonal fluctuations in allele frequencies seems small and temporal structure independent of

seasons may be more important in this system (Buffalo and Coop 2019).

It is challenging to characterize the role of selection in producing spatial or seasonal

change in allele frequencies. At the spatial scale, the axis of demography and environmental

heterogeneity are confounded in this system (i.e., migration and environment are structured

along the south-north axis) (Caracristi and Schlotterer 2003; Yukilevich and True 2008;

Duchen et al. 2013; Kao et al. 2015). At the seasonal scale, the magnitude of allele frequency

change with seasons is expected to be rather small, and stochastic environmental events not

aligned with seasons complicate inferences even further (Machado et al. 2018). However, we

can gain power by jointly modelling latitudinal and seasonal changes (Cogni et al. 2015).

Two adaptive mechanisms are expected to induce correlations between clinal and

seasonal fluctuations in allele frequencies. First, the environment changes similarly with

latitude and through seasons (at least with respect to temperature). Second, the onset of

spring changes with latitude, and so seasonal changes in polymorphisms alone could produce

clinal variation, a mechanism termed seasonal phase clines (Roff 1980; Rhomberg and Singh

1986). Concordant with this interpretation, the prevalence of reproductive diapause and the

frequency of a variant in the gene that mediates diapause inducibility vary both latitudinally

and seasonally in *D. melanogaster* (Schmidt et al. 2005; Cogni et al. 2014). The frequency of

this diapause-inducing variant drops in the summer and is positively correlated with latitude.

Importantly, demographic processes that can generate clinal patterns are independent to

variation across seasons, given the short time scale at which the seasonal processes happen

(Bergland et al. 2014; Cogni et al. 2015). Thus, parallel latitudinal and seasonal variation in a

trait is strong evidence in favor of natural selection.

A few other studies have looked for signatures of parallel clinal and seasonal variation

in *D. melanogaster* (Bergland et al. 2014; Cogni et al. 2015; Kapun et al. 2016; Behrman et

al. 2018; Machado et al. 2018). Cogni et al. (2015) found an association between clinal and

seasonal change in central metabolic genes, which are likely important drivers of climatic

adaptation. Kapun et al. (2016) dissected the role of selection in shaping clinal variation in *D.*

melanogaster cosmopolitan inversions. A more recent study also investigated the association

between clinal and seasonal variation and showed the direction of allele frequency change

with latitude is concordant with the change with seasons in outlier polymorphisms across the

D. melanogaster genome (Machado et al. 2018).

Here, we aim to quantify the relationship clinal and seasonal change and to evaluate

whether the pattern observed is consistent with natural selection. Our work is fundamentally

different from previous studies because (i) we use seasonal samples collected over six years,

as opposed to at most three years that were used in previous studies, increasing power to find

seasonal effects; (ii) our samples are all from the same location in Pennsylvania, where

seasonality is strong and phenotypes are known to cycle seasonally, and where there is little

evidence of population substructure or large scale migrations events (Schmidt and Conde

2006; Behrman et al. 2015; Rajpurohit et al. 2017; Behrman et al. 2018); (iii) we analyze how

the correlation between clinal and seasonal variation changes across genomic regions that

differ in density of functional sites, what allows us to better disentangle demography and

selection; and (iv) we dissect the role of linkage disequilibrium in driving these patterns.

Material and Methods

Population samples

We analyzed 20 samples from seven locations along the United States east coast, collected by

(Bergland et al. 2014) (10 samples), and (Machado et al. 2018) (10 samples) (see Table S1).

The samples were based on pools of wild-caught individuals. We decided to not include

previously collected samples from Maine because they were collected in the fall, whereas all

of our other samples were collected in the spring, and we also did not include the DGRP

sample from North Carolina, as it is hard to ascertain when they were obtained (Fabian et al.

2012; Mackay et al. 2012; Bergland et al. 2014). The Linvilla (Pennsylvania) population was

sampled extensively from 2009 to 2015 (six spring, seven fall samples), and was therefore

used in our analysis of seasonal variation. One of the Pennsylvania samples was exclusively

used for the clinal analysis. We also replicated our clinal analysis using data from four

Australian samples (Anderson et al. 2005; Kolaczkowski et al. 2011). All the data used in this

project are available on the NCBI Short Read Archive (BioProject accession numbers

PRJNA256231, PRJNA308584 and NCBI Sequence Read Archive SRA012285.16).

Mapping and processing of sequencing data

Raw, paired-end reads were mapped against the FlyBase *D. melanogaster* (r6.15) and *D.*

simulans (r2.02) reference genomes (Gramates et al. 2017) using BBSplit from the BBMap

suite (https://sourceforge.net/projects/bbmap/; version from February 11, 2019). We removed

any reads that preferentially mapped to *D. simulans* to mitigate effects of contamination (the

proportion of reads preferentially mapping to *D. simulans* was minimal, never exceeding

3%). Then, reads were remapped to *D. melanogaster* reference genome using bwa (MEM

algorithm) version 0.7.15 (Li and Durbin 2010). Files were converted from SAM to BAM

format using Picard Tools (http://broadinstitute.github.io/picard). PCR duplicates were

marked and removed using Picard Tools and local realignment around indels was performed

using GATK version 3.7 (McKenna et al. 2010). Single nucleotide polymorphisms (SNPs)

and indels were called using CRISP with default parameters (Bansal et al. 2016).

We applied several filters to ensure that the identified SNPs were not artifacts. SNPs

in repetitive regions, identified using the RepeatMasker library for *D. melanogaster* (obtained

from http://www.repeatmasker.org), and SNPs within 5bp of polymorphic indels were

removed from our analyses. SNPs with mean minor allele frequency in the clinal or seasonal

samples less than 5%, with minimum per-population coverage less than 10x (or 4x for the

Australian samples) or maximum per-population coverage greater than the 99[th] quantile were

excluded from our analyses. Also, SNPs with multi-allelic states were removed from the

analysis. We did not include SNPs in any of the sex chromosomes. SNPs were annotated with

their genic classes using SNPeff version 4.3o (Cingolani et al. 2012).

Clinal and seasonal changes in allele frequency

The allele frequencies were calculated by diving the number of reads supporting each allele,

divided by the total number of reads. Because pool-seq data contain an additional component

of error due to sampling, we did not weight the allele frequencies by total depth at each site;

instead we used the effective sample size, or effective number of chromosomes (N_E), as the

denominator. This metric can be computed as follows:

$$N_E = \left(\frac{1}{D} + \frac{1}{N_C}\right)^{-1}$$

where N_C is the number of chromosomes in the pool and D is the read depth at that site

(Kolaczkowski et al. 2011; Feder et al. 2012; Bergland et al. 2014).

To assess latitudinal variation, we fitted a binomial linear model of allele frequency

against latitude for each site. Similarly, we regressed allele frequency at each site against a

season dummy variable (June and July were encoded as Spring, and September, October and

November as Fall) and included the year of sampling as a covariate. For either regression, we

required the variant to be polymorphic in at least two samples. Further, we computed

pairwise F_{ST} for all our samples using the R package poolfstat (Hivert et al., 2018).

We defined clinal and seasonal SNPs using an outlier approach, because we do not

have an adequate genome-wide null distribution to compare our estimates. We considered

that SNPs were outliers if their regression P-value fell in the bottom 1% (or 5%) of the

distribution.

Correlation between clinal and seasonal variation

Our main goal was to evaluate whether clinal and seasonal change are correlated, pooling

information across the thousands of polymorphisms that segregate in natural populations. To

do so, we regressed the slopes of the clinal regressions and the slope of the seasonal

regressions. The regression line was fit using Huber's M estimator to improve robustness to

outliers. Before fitting the regression, we z-normalized the clinal and seasonal slopes, so the

slope of the regression of clinal and seasonal change is actually the same as the correlation.

We also investigated how the correlation between clinal and seasonal change differed

across genomic regions. For that, we used a dummy variable with annotations as a covariate.

The regions analyzed were exon, intron, 5' UTR, 3' UTR, upstream, downstream intergenic

and splice. There are some chromosomal inversions segregating in the populations we

studied, and they are known to contribute to adaptation (Wright and Dobzhansky 1946;

García-Vázquez and Sánchez-Refusta 1988; Kapun et al. 2014). We annotated SNPs

surrounding (2Mb) common inversion breakpoints and added inversion status as a covariate

in the linear model (Corbett-Detig and Hartl 2012).

206   To confirm our results are robust to potential model misspecifications, we

implemented a permutation test in which we rerun the regressions for each SNP using

shuffled season and latitude labels 2,000 times. The same procedure was implemented for

most of the statistical tests, except where indicated otherwise.

Enrichment tests

We tested for enrichment of genic classes using our sets of clinal and seasonal SNPs using

Fisher's exact test for each genic region and statistic. To control for confounders, such as

read depth and allele frequency variation, we shuffled the season and latitude labels and reran

the generalized regressions. Using the P-values obtained from regressions in which season

and latitude labels were shuffled, we defined, for each iteration, lists of top clinal and

seasonal SNPs. Then, we calculated the enrichment of each genic class using Fisher's exact

test. To obtain a P-value for an enrichment of a given genic class, we compared the observed

odds ratio in the actual dataset to the distribution of odds ratios observed for datasets in

which season and latitude labels were shuffled.

Mitigating the impact of linkage disequilibrium

Selection at one site affects genetic variation at nearby, linked neutral sites (Smith and Haigh

1974). Because we assume that sites are independent in our models, the indirect effects of

selection can inflate the magnitude of the patterns we investigated. To test the effect of

linkage disequilibrium (LD) in our outlier analyses, we plotted P-values against distance to a

top SNP. We then smoothed the scatterplot using cubic splines as implemented in ggplot2

(Wickham 2016). To test the effect of linkage on the relationship between clinal and seasonal

variation, we implemented a thinning approach. Sampling one SNP per *L* base pairs one

thousand times, we constructed sets of SNPs with minimized dependency, where *L* ranged

from 1 to 20kb. For each of these sets for a given *L*, we computed the correlation between

clinal and seasonal slopes, and compared the distribution of the thinned regression

coefficients to the coefficients we obtained using all SNPs.

All statistical analyses were performed in R 3.5.0 (R Core Team 2018) and can be

found at gitlab.com/mufernando/clinal_sea.git.

Results

We assembled 20 *D. melanogaster* population samples collected from seven localities across

multiple years in the east coast of the United States. All of these samples are the result of a

collaborative effort of many researchers from a consortium, the DrosRTEC (Bergland et al.

2014; Machado et al. 2018). Seven of our samples span from Florida to Massachusetts and

together comprise our clinal set. The seasonal samples were collected in Pennsylvania in the

spring (6 samples collected in June or July) and in the fall (6 samples collected in September,

October or November). For each sample, a median of 55 individuals (with a range of 33 to

116) was pooled and resequenced to an average 75x coverage (ranging from 17 to 215). We

also used four clinal samples from the Australia (Anderson et al. 2005; Kolaczkowski et al.

2011). More details about the samples can be found on Table S1 (also see Machado et al.

2018; Bergland et al. 2014). After all the filtering steps, we identified 798,176 common

autosomal SNPs, which were used in our downstream analyses.

Allele frequency changes with latitude, seasons and years

Latitude explains much of allele frequency variation along the surveyed populations, as there

is an excess of low GLM P-value SNPs (Fig. 1A). The mean absolute difference in allele

frequency between the ends of the clines is 9.2%. Seasons, on the other hand, explain less of

the variation in allele frequency. There is only a minor excess of low GLM P-value SNPs

(Fig. 1B) and the mean absolute difference in allele frequency between seasons is 2.6%. We

also found that year of sampling is a good predictor of allele frequency change (in

Pennsylvania), more so than seasons, given there is a huge excess of low GLM P-values (Fig.

1C).

Figure 1. Distribution of P-values from the generalized linear models of allele frequency and
latitude, and allele frequency and seasons/years.
259
Our generalized linear models do not account for dependency between samples,

which can be a problem when regressing allele frequency on seasons. To investigate whether

this could be an issue, we performed Durbin-Watson tests for autocorrelation in the residuals

of the seasonal regressions using Julian days as the time variable. There is no excess of low

P-values (Fig. S1), and the season P-values are not correlated with Durbin-Watson test P-

value ($P = 0.77$). This indicates that the assumption of independency is being met for most

variants, and that autocorrelation is not artificially creating patterns of seasonality in allele

frequency.

Given we do not have enough information to build an appropriate null distribution to

calibrate our P-values, we sought to demonstrate that top significant clinal and seasonal SNPs

are enriched for functional variants, which are more likely to contribute to adaptation.

Latitudinal SNPs are more likely to be in exonic and UTR 3' regions (Fig. 2A), whereas

seasonal SNPs are enriched for exonic, UTR 3' and UTR 5' regions (Fig. 2B). Further, top

latitudinal and seasonal SNPs seem to be underrepresented within upstream and downstream

regions. Similar enrichment patterns have been observed for both top clinal and seasonal

(Kolaczowski et al. 2011; Fabian et al. 2012,; Bergland et al. 2014; Machado et al. 2016,

2018). Using a 5% cutoff, our enrichment results are largely replicated (Fig. S2).

Figure 2. Top SNPs are enriched for functionally relevant classes. Enrichment of top 1%
SNPs in each genic class for A) latitudinal P-value and B) seasonal P-value. The histograms
show the distribution of odds ratios when latitude and season labels were permuted, and the
vertical bars show the observed odds ratios.

Clinal variation is related to seasonal variation

A clinal pattern can arise solely as a result of demographic processes, such as isolation by

distance and admixture (Duchen et al. 2013; Kao et al. 2015; Bergland et al. 2016). Although

seasonality is less affected by such processes, seasonal change is less pronounced and more

subject to stochastic changes in the environment, making it harder to detect seasonal change

with precision. Here, we integrate both clinal and seasonal change estimates across a large

number of SNPs in the genome. We expect the overall pattern that emerges to be informative

of the relative role of natural selection, because selection is a plausible process to produce a

pattern of clinal variation mirroring seasonal variation (Cogni et al. 2015).

We found a significant negative correlation between clinal and seasonal regression

coefficients (Fig. 3A, Table S2). The correlation is strongest for SNPs within exons, and the

weakest for unclassified SNPs and those within intergenic regions (Fig. 3A; Table S3).

Nonetheless, the correlation is different than zero for all classes (except for the unclassified),

what would be consistent either with pervasive linked selection or widespread distribution of

variants that are important for adaptation, even within non-coding regions. Qualitatively

similar results were replicated using a different minor allele frequency cutoff and using

samples from Maine, which were obtained the summer – in contrast to all the other clinal

populations that were sampled in the fall (Fig. S3B-C).

Figure 2. Clinal change parallels seasonal change. Correlation between clinal change and
seasonal change for each genic class (A) and by inversion breakpoint (B). (C) Association
between top latitudinal and seasonal variants for different P-value cutoffs. Gray histograms

are the null distribution of the correlation (after permuting the latitude and season labels) and
vertical bars represent the observed correlation.
306
Given that previous studies have demonstrated the importance of cosmopolitan

inversions in climatic adaptation (e.g., Kapun et al. 2016), we looked at the correlation

between clinal and seasonal change near common cosmopolitan inversions breakpoints. We

found that the correlation between clinal and seasonal change is strongest near the

breakpoints of inversions In(2R)NS, In(3R)P and In(3L)P (Fig. 2B, Table S4). Nevertheless,

the pattern is still strong outside these regions, indicating our main results are not purely

driven by frequency changes of inversions.

Another way of testing for parallelism between clinal and seasonal change is by

testing if clinal SNPs are more likely to be seasonal (and vice-versa). We observed that clinal

SNPs are enriched for seasonal SNPs (Fig. 2C). The enrichment increases with more

stringent lower P-value quantile cut-offs, as we would expect if even strictly non-significant

variants were informative of the role of selection.

We also confirmed our main finding, that clinal and seasonal change are correlated

using clinal samples from Australia. To measure clinal change in Australia, we only used

four low coverage samples in Australia, two for each of low and high latitude locations.

There is a negative correlation between clinal variation in Australia and seasonal variation in

Pennsylvania that, although minor, is significant (Fig. S3C).

The effects of linkage disequilibrium on clinal and seasonal variation

Variation at one site is linked to variation at other sites, and selection will increase this

dependency (Smith and Haigh 1974). First, we assessed if latitudinal and seasonal P-values,

were dependent on how distant a SNP was from our top 1% SNPs. We show that both

statistics are dependent on distance from the outlier SNPs (Fig. 3A, B), but the effect

virtually disappears after 5kb.

330    We assessed the impact of linkage on the correlation between clinal and seasonal

331 change by implementing a thinning approach. First, we tested how the genome-wide

332 regression estimate varied with changing window sizes. The effect of non-independency of

333 variants on the correlation is rather small (Fig. 3C) and the genome-wide correlation remains

334 significantly different from zero (P= 0.001; Fig. 3D). The thinning did not significantly

335 reduce the signal for many regions, but the strength of the signal within splicing, UTR 5',

336 upstream, downstream and intergenic regions decreased and is not significantly different

337 from zero (Fig. 3D). It may be that most of the signal coming from those regions are due to

338 the linked effects of selection.

340 **Figure 3.** Effects of linkage disequilibrium. The mean (A) latitudinal P-value and (B)
341 seasonal P-value depend on distance to the respective top 1% outliers. The correlation
342 between clinal and seasonal variation is affected by dependency among SNPs. C) the
343 correlation between clinal and seasonal variation changes with the size of the thinning
344 window. D) comparison among original estimates (arrows) and values obtained after thinning
345 using a window size of 5kb (histogram). Histograms show the distributions across sampled
346 thinned datasets, and the black arrows point to the original estimates.

Biological interpretation of the correlation between clinal and seasonal change

Although the negative correlation between clinal and seasonal change indicates a role for

selection, it is unclear how strongly parallel selection would need to be to generate this

correlation. Intuitively, we expect the correlation to be rather small, as the majority of the

variants are likely not under parallel selection. Below, we derive how to get a rough estimate

of the number of SNPs under parallel selection from the observed correlation.

Suppose that for a proportion p of the SNPs the clinal and seasonal regression

coefficients are correlated due to parallel selection, whereas for the remainder of the SNPs

they are independent. What genome-wide correlation would we expect? To find this, we can

write (Z_1, Z_2) for the two z-normalized regression coefficients of a randomly chosen SNP.

SNPs can be either under parallel selection or not, so we define (X_1, X_2) and (Y_1, Y_2) as

random draws from the z-normalized regression coefficients for SNPs under parallel

selection and not, respectively. Then, $(Z_1, Z_2) = (X_1, X_2)$ with probability p, and $(Z_1, Z_2) =$

(Y_1, Y_2) otherwise. To find the $\mathrm{cor}(Z_1, Z_2)$, note that since Z_1 and Z_2 have mean 0 and

standard deviation 1, $\mathrm{cor}(Z_1, Z_2) = E[Z_1 Z_2]$ (and similarly for X and Y). So,

$\mathrm{cor}(Z_1, Z_2) = p\,\mathrm{cor}(X_1, X_2) + (1 - p)\,\mathrm{cor}(Y_1, Y_2).$

For the subset of SNPs under parallel selection, we suppose $\mathrm{cor}(X_1, X_2) = \rho$ and for

the remainder of the SNPs we suppose no correlation, or $\mathrm{cor}(Y_1, Y_2) = 0$. Then the genome-

wide correlation is $\mathrm{cor}(Z_1, Z_2) = p\,\rho$. Here, p is the proportion of SNPs under parallel

selection and it can be estimated as $p = \mathrm{cor}(Z_1, Z_2) \div \rho$. We do not know what the

correlation ρ between clinal and seasonal change should be for the variants under selection,

but since it is expected to be $-1 < \rho < 0$, estimating p as $-\mathrm{cor}(Z_1, Z_2)$ is conservative.

Recall we assume ρ is negative because the climate becomes colder the greater the latitude,

but it gets warmer from spring to fall.

Note our model has few assumptions: (i) our measures of clinal and seasonal change

have mean zero and variance one, which is met given we are dealing with the z-normalized

regression coefficients and (ii) all polymorphisms are independent from one another.

Accounting for linkage disequilibrium is notoriously complicated, especially because we

cannot accurately measure LD from pooled sequencing (Feder et al. 2012). In the previous

section we show that we are able to mitigate the effects of LD on the correlation between

clinal and seasonal change using a thinning approach.

We can now readily interpret our observed correlations as proportion of SNPs under

parallel selection (ignoring the negative sign). Using our thinned estimates, the patterns

uncovered here are consistent with 3.78% of the common, autosomal variants being under

parallel selection. It is curious our estimated proportion is close to previous estimates for the

proportion of clinal (3.7% in Machado et al. 2015) and seasonal SNPs (~4% in Machado et

al. 2018).

Discussion

Clinal patterns have been observed in both phenotypic and genotypic traits in many different

species (Hancock et al. 2008; Baxter et al. 2010; Adrion et al. 2015). Especially in systems in

which there is collinearity between the axis of gene flow and environmental heterogeneity,

disentangling the contribution of selection and demography in producing clines is not trivial.

Detecting seasonal cycling in allele frequencies is also challenging, mostly because the effect

size is likely to be small and the environment may change unpredictably within seasons. We

relied on integrating patterns of clinal and seasonal variation, since variation over short

periods is independent of most confounding demographic processes. Parallel environmental

change and the differences in the onset of seasons with latitude can create parallelism

between clinal and seasonal variation. Natural selection would be the process driving the

coupling between latitudinal and seasonal change in both scenarios, thus investigating it can

help discern the role of natural selection.

Here, using *D. melanogaster* samples from the east coast of the United States and

Australia, we show there is a genome-wide correlation between clinal and seasonal change.

The correlation varies across genomic regions, being stronger for genic when compared to

intergenic regions (Fig. 2A,C). Demographic processes are expected to impact the genome as

a whole, but the effects of selection are stronger in regions with higher density of functional

(Andolfatto 2005). We derived a way to get a rough estimate of the number of SNPs under

parallel selection from the observed genome-wide correlation and found that 3.78% of

common, autosomal SNPs could be driven by selection both in space and time.

Segregating inversions are known to underlie much of climatic adaptation in *D.*

melanogaster (Fabian et al. 2012, Kapun et al. 2014, 2016), therefore we investigated how

much of our signal depended on inversion status. We found the correlation between clinal

and seasonal change to be stronger surrounding common inversions, highlighting the role of

selection in driving frequency changes in common, cosmopolitan inversions in *D.*

melanogaster. The correlation between clinal and seasonal change is particularly high for

SNPs near In(3R)Payne breakpoints, an inversion known to be associated to phenotypes

relevant to adaptation to cooler climates (reviewed in Kapun et al. 2019). Importantly, clinal

and seasonal change are significantly coupled for SNPs far from inversion breakpoints,

suggesting loci involved in adaptation at the spatial and seasonal scales are not restricted to

inversions.

Because we expect selection to intensify linkage disequilibrium, the correlation

between clinal and seasonal variation could be mostly driven by a few large effect loci.

However, we show that our results cannot be fully explained by linkage (Fig. 3). When

accounting for autocorrelation along the chromosomes, the correlation disappears for variants

within intergenic regions, but not for exons (Fig. 3D). The effects of linkage disequilibrium

are important in this short time scale, but it decays rapidly, and seems to return to background

levels after 5kb (Fig. 3A-C).

Population substructure and migration could be causing seasonal variation in allele

frequency in *D. melanogaster*. One possibility is that rural populations of *D. melanogaster* in

temperate regions collapse during the winter and recover from spring to fall. However,

reproductive diapause cycles in orchards and reaches high frequencies early in the spring,

whereas its frequency in urban fruit markets in Philadelphia is much lower (Schmidt and

Conde 2006). Another possibility is that seasonal variation is produced by migration of flies

from the south in the summer, and from the north in the winter. There is little evidence of

long-range migration in *D. melanogaster*, though this process seems important in *D. simulans*

(Bergland et al. 2014; Machado et al. 2016). *D. melanogaster* have been shown to survive

and reproduce during winter season in temperate regions, so rural flies can withstand a harsh

winter season and be subject to selection (Mitrovski and Hoffmann 2001; Hoffmann et al.

2003, Rudman et al. 2019). These seasonal patterns have been replicated in many populations

across North America and Europe (Machado et al. 2018), bolstering the argument for

seasonal selection. Given the patterns we uncovered here are the result of subtle, but

repeatable changes across multiple seasons, it is hard to imagine that selection is not the main

causing force, even if it is acting to maintain cryptic population structure within each

location.

Importantly, our results should be robust to differential admixture from Europe and

Africa to the ends of the clines (Duchen et al. 2013; Kao et al. 2015; Bergland et al. 2016).

Given variation over seasonal time scales are less affected by broader scale migration

patterns, it is not plausible that secondary contact can create parallel clinal and seasonal

variation. Furthermore, the evidence for secondary contact in Australia is quite weak (see

Bergland et al. 2016), but we show that clinal variation in Australia is related to seasonal

variation in Pennsylvania (Fig. S3). Secondary contact may have contributed ancestral

variation, that have since been selectively sorted along the cline (Flatt 2016). Consistent with

this interpretation, the proportion of African ancestry is lower in low recombination regions,

suggesting selection is mediating admixture in *D. melanogaster* (Pool 2015).

An important mechanism that can cause clinal patterns has been neglected from

recent discussions of clinal variation in *Drosophila*. The latitudinal variation on the onset of

seasons can produce clines, a phenomenon termed "seasonal phase clines" (Roff 1980;

Rhomberg and Singh 1986). Under this model, a correlation between clinal and seasonal

change is expected. Our latitudinal samples were all collected within one month of difference

(during the spring), and so our observations could be partially explained by differences in the

seasonal phase. Our data does not allow for proper disentangling of seasonal phase clines

from parallel environmental change but change on the onset of seasons alone cannot explain

our results. We found that latitude is usually a much better predictor of allele frequency

differences (Fig. 1A-B), and the magnitude of change along the cline is much greater than

what we found within a population across seasons (9.2% vs. 2.64%). We show that including

Maine samples (which were obtained in the fall, in contrast to all other samples that were

sampled in the spring) in our analyses does not meaningfully change our main results (Fig.

S3). However, the rate of change (per degree of latitude) in allele frequency from Florida to

Maine is smaller than the rate from Florida to Massachusetts. This could be due to

differences in sampling year, but we also found the rate of change in frequency between

Virginia and Maine to be smaller than the rate between Virginia and Massachusetts (FL and

ME were sampled in 2009, whereas VA and MA were sampled in 2012; Fig. S5). We

believe these differences are due to the shift in seasonal phase, as the samples from Maine

were collected in the fall, but all other samples were collected in the spring.

A recent study suggested the temporal changes in allele frequency reported in

Bergland et al. (2014) is only weakly consistent with seasonal selection (Buffalo et al. 2019).

Consecutive spring-fall pairs showed some signal of adaptation, but the effect was small and

disappeared at larger timescales (same season but across different years). Similarly, Machado

et al. 2018 found that when they flipped the season labels of some samples the seasonal

model fit was greatly improved. Consistent with these observations, we found there is strong

temporal structure across years (Fig. 1C) and the matrix of pairwise F_{ST} shows some strong

and seemingly haphazard temporal events (e.g., consider the entries for PA_07_2010 and

PA_07_2015 in Fig. S5). Here with an expanded seasonal set of samples, we show that the

distribution of seasonal P-values is only slightly enriched for low P-values (Fig. 1B), but top

seasonal SNPs are enriched for functional genic classes, when compared to datasets in which

the season labels were permuted (Fig. 2B). These results highlight how difficult it is to find

truly seasonal SNPs with current datasets. Once more comprehensive time series data is

available, environmental heterogeneity could be explored without the need for an imperfect

proxy (such as seasons).

Many species occur along spatially structured environments and show clinal variation

in traits, so a question that remains open is: what is the role of selection in producing and

maintaining these patterns? Seasonal variation is also ubiquitous, especially in temperate

environments, so seasonal change could be an important feature of organisms that have

multiple generations each year (Behrman et al. 2015). Here, we demonstrate that by

integrating clinal and seasonal variation, we can discern the contributions of selection in

driving allele frequency changes with the environment. This approach could potentially be

applied to other multivoltine species that occur along environmental gradients, including

invasive species, which are known to often have short generation times and to reproduce

quickly (Sakai et al. 2001).

References

Adrion JR, Hahn MW, Cooper BS. 2015. Revisiting classic clines in Drosophila melanogaster in the age of genomics. Trends Genet. 31:434–444.

Agis M, Schlötterer C. 2001. Microsatellite variation in natural Drosophila melanogaster populations from New South Wales (Australia) and Tasmania. Mol. Ecol. 10:1197–1205.

Anderson, A. R., Hoffmann, A. A., McKechnie, S. W., Umina, P. A., & Weeks, A. R. 2005. The latitudinal cline in the In (3R) Payne inversion polymorphism has shifted in the last 20 years in Australian Drosophila melanogaster populations. Molecular Ecology. 14(3), 851-858.

Andolfatto, P. 2005. Adaptive evolution of non-coding DNA in Drosophila. Nature. 437(7062), 1149.

Arthur AL, Weeks AR, Sgrò CM. 2008. Investigating latitudinal clines for life history and stress resistance traits in Drosophila simulans from eastern Australia. J. Evol. Biol. 21:1470–1479.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat. Genet. 25:25–29.

Bansal V, Bansal V, Libiger O. 2016. A statistical method for the detection of variants from next-generation resequencing of DNA pools. Bioinformatics 32:3213.

Barton NH. 1983. Multilocus clines. Evolution 37:454–471.

Barton NH. 1999. Clines in polygenic traits. Genet. Res. 74:223–236.

Baxter I, Brazelton JN, Yu D, Huang YS, Lahner B, Yakubova E, Li Y, Bergelson J, Borevitz JO, Nordborg M, et al. 2010. A Coastal Cline in Sodium Accumulation in Arabidopsis thaliana Is Driven by Natural Variation of the Sodium Transporter AtHKT1;1. PLoS Genet. 6:e1001193.

Behrman EL, Howick VM, Kapun M, Staubach F, Bergland AO, Petrov DA, Lazzaro BP, Schmidt PS. 2018. Rapid seasonal evolution in innate immunity of wild Drosophila melanogaster. Proc. Biol. Sci. [Internet] 285. Available from: http://dx.doi.org/10.1098/rspb.2017.2599

Behrman EL, Watson SS, O'Brien KR, Heschel MS, Schmidt PS. 2015. Seasonal variation in life history traits in two Drosophila species. J. Evol. Biol. 28:1691–1704.

Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. 2014. Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in Drosophila. PLoS Genet. 10:e1004775.

Bergland AO, Tobler R, González J, Schmidt P, Petrov D. 2016. Secondary contact and local adaptation contribute to genome-wide patterns of clinal variation in Drosophila melanogaster. Mol. Ecol. 25:1157–1174.

Berry A, Kreitman M. 1993. Molecular analysis of an allozyme cline: alcohol dehydrogenase in Drosophila melanogaster on the east coast of North America. Genetics 134(3):869-93.

Bock IR, Parsons PA. 1981. Species of Australia and New Zealand. Genetics and biology of Drosophila 3:291–308.

Božičević V, Hutter S, Stephan W, Wollstein A. 2016. Population genetic evidence for cold adaptation in European Drosophila melanogaster populations. Molecular ecology 25(5):1175-91.

Buffalo V, Coop G. 2019. Estimating the genome-wide contribution of selection to temporal allele frequency change. BioRxiv, 798595.

Campitelli BE, Stinchcombe JR. 2013. Natural selection maintains a single-locus leaf shape cline in Ivyleaf morning glory, Ipomoea hederacea. Mol. Ecol. 22:552–564.

Caracristi, G., and C. Schlötterer. 2003. "Genetic Differentiation between American and European Drosophila Melanogaster Populations Could Be Attributed to Admixture of African Alleles." Molecular Biology and Evolution 20 (5): 792–99.

Cardini A, Jansson A-U, Elton S. 2007. A geometric morphometric approach to the study of ecogeographical and clinal variation in vervet monkeys. J. Biogeogr. 34:1663–1678.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6:80–92.

Cogni R, Kuczynski C, Koury S, Lavington E, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2014. The intensity of selection acting on the couch potato gene--spatial-temporal variation in a diapause cline. Evolution 68:538–548.

Cogni R, Kuczynski K, Lavington E, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2015. Variation in Drosophila melanogaster central metabolic genes appears driven by natural selection both within and between populations. Proceedings of the Royal Society of London B: Biological Sciences 282:20142688.

Cogni R, Kuczynski K, Koury S, Lavington E, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2017. On the long-term stability of clines in some metabolic genes in Drosophila melanogaster. Scientific reports 7:42766.

Comeron JM, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in Drosophila melanogaster. PLoS Genet. 8:e1002905.

Corbett-Detig, Russell B., and Daniel L. Hartl. Population Genomics of Inversion Polymorphisms in Drosophila Melanogaster. PLOS Genet. 8, no. 12 (December 20, 2012): e1003056.

556
Coyne JA, Beecham E. 1987. Heritability of two morphological characters within and among natural
populations of Drosophila melanogaster. Genetics 117:727–737.
David J, Capy P, Payant V, Tsakas S. 1985. Thoracic trident pigmentation in Drosophila melanogaster:
Differentiation of geographical populations. Genet. Sel. Evol. 17:211–224.
David JR, Bocquet C. 1975a. Evolution in a cosmopolitan species: Genetic latitudinal clines in Drosophila
melanogaster wild populations. Experientia 31:164–166.
David JR, Bocquet C. 1975b. Similarities and differences in latitudinal adaptation of two Drosophila sibling
species. Nature 257:588–590.
David JR, Capy P. 1988. Genetic variation of Drosophila melanogaster natural populations. Trends Genet.
4:106–111.
Dionne M, Miller KM, Dodson JJ, Caron F, Bernatchez L. 2007. Clinal Variation In Mhc Diversity With
Temperature: Evidence For The Role Of Host? Pathogen Interaction On Local Adaptation In Atlantic Salmon.
Evolution 61:2154-2164.
Dobzhansky T. 1943. Genetics of Natural Populations IX. Temporal Changes in the Composition of Populations
of Drosophila Pseudoobscura. Genetics 28:162–186.
Duchen P, Zivkovic D, Hutter S, Stephan W, Laurent S. 2013. Demographic inference reveals African and
European admixture in the North American Drosophila melanogaster population. Genetics 193:291–301.
Duvernell DD, Schmidt PS, Eanes WF. 2003. Clines and adaptive evolution in the methuselah gene region in
Drosophila melanogaster. Molecular ecology 12(5):1277-85.
Endler JA. 1977. Geographic variation, speciation, and clines. Monogr. Popul. Biol. 10:1–246.
Ewing EP. 1979. Genetic Variation in a Heterogeneous Environment VII. Temporal and Spatial Heterogeneity
in Infinite Populations. Am. Nat. 114:197–212.
Excoffier L, Foll M, Petit RJ. 2009. Genetic Consequences of Range Expansions. Annu. Rev. Ecol. Evol. Syst.
40:481–501.
Fabian DK, Kapun M, Nolte V, Kofler R, Schmidt PS, Schlötterer C, Flatt T. 2012. Genome-wide patterns of
latitudinal differentiation among populations of Drosophila melanogaster from North America. Mol. Ecol.
21:4748–4769.
Feder AF, Petrov DA, Bergland AO. 2012. LDx: estimation of linkage disequilibrium from high-throughput
pooled resequencing data. PLoS One 7:e48588.
Fisher RA. 1992. Statistical Methods for Research Workers. In: Kotz S, Johnson NL, editors. Breakthroughs in
Statistics: Methodology and Distribution. New York, NY: Springer New York. p. 66–70.
Flatt T. 2016. Genomics of clinal variation in Drosophila: disentangling the interactions of selection and
demography. Molecular ecology 25(5):1023-6.
García-Vázquez E, Sánchez-Refusta F. 1988. Chromosomal polymorphism and extra bristles of Drosophila
melanogaster: joint variation under selection in isofemale lines. Genetica 78:91–96.
Gillespie J. 1973. Polymorphism in random environments. Theor. Popul. Biol. 4:193–195.
Gramates LS, Marygold SJ, Santos GD, Urbano J-M, Antonazzo G, Matthews BB, Rey AJ, Tabone CJ, Crosby
MA, Emmert DB, et al. 2017. FlyBase at 25: looking to the future. Nucleic Acids Res. 45:D663–D671.
Hancock AM, Witonsky DB, Gordon AS, Eshel G, Pritchard JK, Coop G, Di Rienzo A. 2008. Adaptations to
climate in candidate genes for common metabolic disorders. PLoS Genet. 4:e32.
Hivert, Valentin, Raphaël Leblois, Eric J. Petit, Mathieu Gautier, and Renaud Vitalis. 2018. Measuring Genetic
Differentiation from Pool-Seq Data. Genetics. 210, no. 1: 315–30.
Hoffmann AA, Anderson A, Hallas R. 2002. Opposing clines for high and low temperature resistance in
Drosophila melanogaster. Ecol. Lett. 5:614–618.
Hoffmann AA, Scott M, Partridge L, Hallas R. 2003. Overwintering in Drosophila melanogaster: outdoor field
cage experiments on clinal and laboratory selected populations help to elucidate traits under selection. J. Evol.
Biol. 16:614–623.
Hoffmann AA, Weeks AR. 2007. Climatic selection on genes and traits after a 100 year-old invasion: a critical
look at the temperate-tropical clines in Drosophila melanogaster from eastern Australia. Genetica 129:133–147.
de Jong G, Bochdanovits Z. 2003. Latitudinal clines inDrosophila melanogaster: Body size, allozyme
frequencies, inversion frequencies, and the insulin-signalling pathway. J. Genet. 82:207–223.
Kao JY, Zubair A, Salomon MP, Nuzhdin SV, Campo D. 2015. Population genomic analysis uncovers African
and European admixture in Drosophila melanogaster populations from the south-eastern United States and
Caribbean Islands. Mol. Ecol. 24:1499–1509.
Kapun M, Fabian DK, Goudet J, Flatt T. 2016. Genomic Evidence for Adaptive Inversion Clines in Drosophila
melanogaster. Mol. Biol. Evol. 33:1317–1336.
Kapun M, van Schalkwyk H, McAllister B, Flatt T, Schlötterer C. 2014. Inference of chromosomal inversion
dynamics from Pool-Seq data in natural and laboratory populations of Drosophila melanogaster. Mol. Ecol.
23:1813–1827.

Kapun, Martin, and Thomas Flatt. 2019. The Adaptive Significance of Chromosomal Inversion Polymorphisms
in Drosophila Melanogaster. Molecular Ecology 28, no. 6: 1263–82.
Kawecki, Tadeusz J., and Dieter Ebert. 2004. Conceptual Issues in Local Adaptation. Ecology Letters 7, no. 12:
1225–41.
Keller A. 2007. Drosophila melanogaster's history as a human commensal. Curr. Biol. 17:R77–R81.
Kennington WJ, Killeen JR, Goldstein DB, Partridge L. 2003. Rapid laboratory evolution of adult wing area in
Drosophila melanogaster in response to humidity. Evolution 57:932–936.
Knibb WR. 1982. Chromosome inversion polymorphisms in Drosophila melanogaster II. Geographic clines and
climatic associations in Australasia, North America and Asia. Genetica 58:213–221.
Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C, Schlötterer C. 2011.
PoPoolation: A Toolbox for Population Genetic Analysis of Next Generation Sequencing Data from Pooled
Individuals. PLoS One 6:e15925.
Kofler R, Schlötterer C. 2012. Gowinda: unbiased analysis of gene set enrichment for genome-wide association
studies. Bioinformatics 28:2084–2085.
Kolaczkowski B, Kern AD, Holloway AK, Begun DJ. 2011. Genomic differentiation between temperate and
tropical Australian populations of Drosophila melanogaster. Genetics 187:245–260.
Kooyers NJ, Greenlee AB, Colicchio JM, Oh M, Blackman BK. 2015. Replicate altitudinal clines reveal that
evolutionary flexibility underlies adaptation to drought stress in annual Mimulus guttatus. New Phytol.
206:152–165.
Krimbas CB, Powell JR. 1992. Drosophila inversion polymorphism. CRC press.
Lavington E, Cogni R, Kuczynski C, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF. 2014. A
small system--high-resolution study of metabolic adaptation in the central metabolic pathway to temperate
climates in Drosophila melanogaster. Mol. Biol. Evol. 31:2032–2041.
Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics
26:589–595.
Li H, Stephan W. 2006. Inferring the demographic history and rate of adaptive substitution in Drosophila. PLoS
Genet. 2:e166.
Lynch M, Bost D, Wilson S, Maruki T, Harrison S. 2014. Population-genetic inference from pooled-sequencing
data. Genome Biol. Evol. 6:1210–1218.
Machado HE, Bergland AO, O'Brien KR, Behrman EL, Schmidt PS, Petrov DA. 2016. Comparative population
genomics of latitudinal variation in Drosophila simulans and Drosophila melanogaster. Mol. Ecol. 25:723–740.
Machado HE, Bergland AO, Taylor R, Tilk S, Behrman E, Dyer K, Fabian DK, Flatt T, González J, Karasov
TL, et al. 2018. Broad geographic sampling reveals predictable and pervasive seasonal adaptation in Drosophila.
Available from: http://dx.doi.org/10.1101/337543
Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM,
Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference Panel. Nature 482:173–178.
McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel
S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation
DNA sequencing data. Genome Res. 20:1297–1303.
Mettler LE, Voelker RA, Mukai T. 1977. Inversion Clines in Populations of DROSOPHILA
MELANOGASTER. Genetics 87:169–176.
Mitrovski P, Hoffmann AA. 2001. Postponed reproduction as an adaptation to winter conditions in Drosophila
melanogaster: evidence for clinal variation under semi-natural conditions. Proc. Biol. Sci. 268:2163–2168.
Oakeshott JG, Gibson JB, Anderson PR, Knibb WR, Anderson DG, Chambers GK. 1982. Alcohol
dehydrogenase and glycerol-3-phosphate dehydrogenase clines in Drosophila melanogaster on different
continents. Evolution 36:86–96.
Paaby AB, Bergland AO, Behrman EL, Schmidt PS. 2014. A highly pleiotropic amino acid polymorphism in
the Drosophila insulin receptor contributes to life-history adaptation. Evolution 68:3395–3409.
Paaby AB, Blacket MJ, Hoffmann AA, Schmidt PS. 2010. Identification of a candidate adaptive polymorphism
for Drosophila life history by parallel independent clines on two continents. Mol. Ecol. 19:760–774.
Pavlidis P, Jensen JD, Stephan W, Stamatakis A. 2012. A critical assessment of storytelling: gene ontology
categories and the importance of validating genomic scans. Mol. Biol. Evol. 29:3237–3248.
Pool JE. The mosaic ancestry of the Drosophila genetic reference panel and the D. melanogaster reference
genome reveals a network of epistatic fitness interactions. 2015. Molecular biology and evolution. 32(12):3236-
51.
Rajpurohit S, Hanus R, Vrkoslav V, Behrman EL, Bergland AO, Petrov D, Cvačka J, Schmidt PS. 2017.
Adaptive dynamics of cuticular hydrocarbons in Drosophila. J. Evol. Biol. 30:66–80.
R Core Team. 2018. R: A Language and Environment for Statistical Computing. Available from:
https://www.R-project.org/

Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD. 2014. Parallel Geographic Variation in
Drosophila melanogaster. Genetics 197:361–373.
Rhomberg LR, Singh RS. 1986. Evidence for a link between local and seasonal cycles in gene frequencies and
latitudinal gene clines in a cyclic parthenogen. Genetica78(1):73-9.
Roff D. 1980. Optimizing development time in a seasonal environment: the 'ups and downs' of clinal variation.
Oecologia 45(2):202-8.
Rudman, Seth M., Sharon Greenblum, Rachel C. Hughes, Subhash Rajpurohit, Ozan Kiratli, Dallin B. Lowder,
Skyler G. Lemmon, Dmitri A. Petrov, John M. Chaston, and Paul Schmidt. 2019. "Microbiome Composition
Shapes Rapid Genomic Adaptation of Drosophila Melanogaster." BioRxiv, 632257.
Sakai AK, Allendorf FW, Holt JS, Lodge DM, Molofsky J, With KA, Baughman S, Cabin RJ, Cohen JE,
Ellstrand NC, et al. 2001. The Population Biology of Invasive Species. Annu. Rev. Ecol. Syst. 32:305–332.
Schmidt PS, Conde DR. 2006. Environmental heterogeneity and the maintenance of genetic variation for
reproductive diapause in Drosophila melanogaster. Evolution 60:1602–1611.
Schmidt PS, Duvernell DD, Eanes WF. 2000. Adaptive evolution of a candidate gene for aging in Drosophila.
Proc. Natl. Acad. Sci. U. S. A. 97:10861–10865.
Schmidt PS, Matzkin L, Ippolito M, Eanes WF, Hey J. 2005. Geographic variation in diapause incidence, life-
history traits, and climatic adaptation in Drosophila melanogaster. Evolution 59:1721–1732.
Schmidt PS, Paaby AB. 2008. Reproductive diapause and life-history clines in North American populations of
Drosophila melanogaster. Evolution 62:1204–1215.
Schrider DR, Hahn MW, Begun DJ. 2016. Parallel evolution of copy-number variation across continents in
Drosophila melanogaster. Mol. Biol. Evol. 33:1308–1316.
Sezgin E, Duvernell DD, Matzkin LM, Duan Y, Zhu C-T, Verrelli BC, Eanes WF. 2004. Single-locus
latitudinal clines and their relationship to temperate adaptation in metabolic genes and derived alleles in
Drosophila melanogaster. Genetics 168:923–931.
Singh RS, Rhomberg LR. 1987a. A Comprehensive Study of Genic Variation in Natural Populations of
Drosophila melanogaster. II. Estimates of Heterozygosity and Patterns of Geographic Differentiation. Genetics
117:255–271.
Singh RS, Rhomberg LR. 1987b. A Comprehensive Study of Genic Variation in Natural Populations of
Drosophila melanogaster. I. Estimates of Gene Flow from Rare Alleles. Genetics 115:313–322.
Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genet. Res. 23:23–35.
Turner TL, Levine MT, Eckert ML, Begun DJ. 2008. Genomic analysis of adaptive differentiation in Drosophila
melanogaster. Genetics 179:455–473.
Vasemägi A. 2006. The adaptive hypothesis of clinal variation revisited: single-locus clines as a result of
spatially restricted gene flow. Genetics 173:2411–2414.
Verrelli BC, Eanes WF. 2001. Clinal variation for amino acid polymorphisms at the Pgm locus in Drosophila
melanogaster. Genetics 157(4):1649-63.
Vigue CL, Johnson FM. 1973. Isozyme variability in species of the genus Drosophila. VI. Frequency-property-
environment relationships of allelic alcohol dehydrogenases in D. melanogaster. Biochemical genetics 9(3):213-
27.
H. Wickham. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
Wittmann MJ, Bergland AO, Feldman MW, Schmidt PS, Petrov DA. 2017. Seasonally fluctuating selection can
maintain polymorphism at many loci via segregation lift. Proceedings of the National Academy of Sciences
114(46): E9932-41.
Wright S. 1943. Isolation by distance. Genetics 28:114–138.
Wright S, Dobzhansky T. 1946. Genetics of natural populations; experimental reproduction of some of the
changes caused by natural selection in certain populations of Drosophila pseudoobscura. Genetics 31:125–156.
Yukilevich, Roman, and John R. True. 2008. "African Morphology, Behavior and Phermones Underlie Incipient
Sexual Isolation between Us and Caribbean Drosophila Melanogaster." Evolution; International Journal of
Organic Evolution 62 (11): 2807–28.
Yukilevich R, Turner TL, Aoki F, Nuzhdin SV, True JR. 2010. Patterns and processes of genome-wide
divergence between North American and African Drosophila melanogaster. Genetics 186:219–239.
Zuther E, Schulz E, Childs LH, Hincha DK. 2012. Clinal variation in the non-acclimated and cold-acclimated
freezing tolerance of Arabidopsis thaliana accessions. Plant Cell Environ. 35:1860–1878.
728

Supplementary information

Figure S1. Distribution of Durbin-Watson P-values. We tested whether there is
autocorrelation in the residuals of the seasonal generalized linear models.

Figure S2. Enrichment of top 5% SNPs in each genic class for A) latitudinal P-value, B)
seasonal P-value. Histograms show the distribution of odds ratios when season labels were
permuted, and the vertical bars indicated the observed odds ratio.

Figure S3. Correlation between clinal and seasonal change using A) 5% allele frequency
cutoff (main results), B) latitudinal samples from Maine, which were collected in the summer
and C) samples from Australia to calculate latitudinal change.

Figure S4. Relationship between proportion of points linearly related (in comparison to
points that are not related) and the correlation between two z-normalized variables. Each
color represents a different assumed degree of linear relatedness (ranging from -1 to -0.1).
Note how for a given estimated relationship between two variables, the slope we assume (-1)
results in the smallest possible proportion of points linearly related, demonstrating that our
test is conservative.

Figure S5. Rate of allele frequency differences between two populations (normalized by
difference in latitude). Note the rates are smaller for comparisons involving the samples from
Maine. FL: Florida (July 2008 and 2010), ME: Maine (October 2009), MA: Massachusetts
(July 2012), VA: Virginia (July 2012).

Figure S6. Mean pairwise F_{ST} values for all US samples included in our main analyses. Refer
to Table S1 for more information on each population.

759 **Table S1.** Information of the samples used in this study.

Accession #	Sample name	Location	Latitude	Collection date	# Flies	Median depth	Month	Season	Seasonal set	Clinal set
SRR1177951	AU_LO1	Queensland, Australia	16.90	2004	17	10				
SRR1177952	AU_LO2	Queensland, Australia	16.90	2004	17	12				
SRR1177953	AU_HI1	Tasmania, Australia	42.77	2004	15	10				
SRR1177955	AU_HI2	Tasmania, Australia	42.77	2004	15	14				
SRR1525685	FL1	Homestead, FL	25.47	Jul-08	39	59	7	Spring	0	1
SRR1525694	FL2	Homestead, FL	25.47	Jul-10	48	37	7	Spring	0	1
SRR1525698	ME1	Bowdoinham, ME	44.02	Oct-09	75	86	9	Fall	0	0
SRR2006283	ME2	Bowdoinham, ME	44.02	Oct-09	75	22	9	Fall	0	0
SRR1525695	GA	Hahira, GA	30.99	Jul-10	51	101	7	Spring	0	1
SRR1525696	SC	Euatwville, SC	33.40	Jul-10	48	83	7	Spring	0	1
SRR3590551	VA_07_2012	Charlotttesville, VA	38.03	Jul-12	69	70	7	Spring	0	1
SRR3939095	PA_06_2013	Linvilla, PA	39.88	Jun-13	54	37	6	Spring	0	1
SRR3590557	MA_07_2012	Lancaster, MA	42.46	Jul-12	90	51	7	Spring	0	1
SRR1525768	PA_07_2009	Linvilla, PA	39.53	Jul-09	55	186	7	Spring	1	0
SRR1525769	PA_11_2009	Linvilla, PA	39.53	Nov-09	74	66	11	Fall	1	0
SRR1525770	PA_07_2010	Linvilla, PA	39.53	Jul-10	116	17	7	Spring	1	0
SRR1525771	PA_11_2010	Linvilla, PA	39.53	Nov-10	33	76	11	Fall	1	0
SRR1525772	PA_07_2011	Linvilla, PA	39.53	Jul-11	75	53	7	Spring	1	0
SRR1525773	PA_10_2011	Linvilla, PA	39.53	Oct-10	47	74	10	Fall	1	0
SRR3590560	PA_10_2012	Linvilla, PA	39.53	Oct-12	100	25	10	Fall	1	0
SRR3590561	PA_07_2012	Linvilla, PA	39.53	Jul-12	115	59	7	Spring	1	0
SRR3590563	PA_9_2012	Linvilla, PA	39.53	Sep-12	50	55	9	Fall	1	0
SRR39	PA_10_2	Linvilla, PA	39.88	Oct-14	50	103	10	Fall	1	0

39096	014									
SRR39 39097	PA_06_2 014	Linvilla, PA	39.88	Jun-14	92	68	6	Spri ng	1	0
SRR39 39098	PA_10_2 015	Linvilla, PA	39.88	Oct-15	52	97	10	Fall	1	0
SRR39 39099	PA_07_2 015	Linvilla, PA	39.88	Jul-15	74	215	7	Spri ng	1	0

761 **Table S2.** Summary of a regression of z-normalized latitude regression coefficients against z-

762 normalized season regression coefficients genome-wide. CI stands for 95% confidence

763 interval.

	Latitudinal slope		
Predictors	*Estimates*	*CI*	*p*
Intercept	0.014	0.012 – 0.016	**<0.001**
Seasonal slope	-0.039	-0.042 – -0.037	**<0.001**
Observations	798196		

765 **Table S3.** Summary of a regression of z-normalized latitude regression coefficients against z-

766 normalized season regression coefficients for each genic class. CI stands for 95% confidence

767 interval.

	Latitudinal slope		
Predictors	*Estimates*	*CI*	*p*
Intercept	0.014	0.011 – 0.016	**<0.001**
Seasonal slope:Intergenic	-0.029	-0.038 – -0.021	**<0.001**
Seasonal slope:Exon	-0.051	-0.058 – -0.044	**<0.001**
Seasonal slope:Intron	-0.046	-0.051 – -0.042	**<0.001**
Seasonal slope:Downstream	-0.037	-0.043 – -0.031	**<0.001**
Seasonal slope:Upstream	-0.036	-0.040 – -0.033	**<0.001**
Seasonal slope:UTR_3	-0.046	-0.056 – -0.036	**<0.001**
Seasonal slope:UTR_5	-0.045	-0.058 – -0.032	**<0.001**
Seasonal slope:Splice	-0.049	-0.071 – -0.026	**<0.001**

Seasonal slope:Other	-0.004	-0.016 – 0.009	0.584
Observations	798196		

769 **Table S4.** Summary of a regression of z-normalized latitude regression coefficients against z-

770 normalized season regression coefficients for each SNPs surrounding inversion breakpoints.

771 CI stands for 95% confidence interval.

	Latitudinal slope		
Predictors	*Estimates*	*CI*	*p*
Intercept	0.014	0.012 – 0.016	**<0.001**
Seasonal slope:In(2L)t	0.004	-0.003 – 0.010	0.235
Seasonal slope:In(2R)NS	-0.020	-0.026 – -0.013	**<0.001**
Seasonal slope:In(3L)P	-0.069	-0.076 – -0.063	**<0.001**
Seasonal slope:In(3R)P	-0.173	-0.181 – -0.166	**<0.001**
Seasonal slope:Outside	-0.031	-0.033 – -0.028	**<0.001**
Observations	798196		

772