α-Synuclein strains that cause distinct pathologies differentially inhibit proteasome

Genjiro Suzuki[1*], Sei Imura[1,2], Masato Hosokawa[1], Ryu Katsumata[1], Takashi Nonaka[1], Shin-Ichi

Hisanaga[2], Yasushi Saeki[3] and Masato Hasegawa[1*]

[1] Department of Dementia and Higher Brain Function, Tokyo Metropolitan Institute of Medical

Science, Tokyo, Japan. [2] Laboratory of Molecular Neuroscience, Department of Biological

Sciences, Tokyo Metropolitan University, Tokyo, Japan. [3] Laboratory of Protein Metabolism,

Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan.

E-mail: suzuki-gj@igakuken.or.jp, hasegawa-ms@igakuken.or.jp

TEL: +81-3-6834-2349

Abstract

Abnormal α-synuclein aggregation has been implicated in several diseases and is known to

spread in a prion-like manner. There is a relationship between protein aggregate structure

(strain) and clinical phenotype in prion diseases, however, whether differences in the strains of

α-synuclein aggregates account for the different pathologies remained unclear. Here, we generated

two types of α-synuclein fibrils from identical monomer and investigated their seeding and

propagation ability in mice and primary-cultured neurons. One α-synuclein fibril induced

marked accumulation of phosphorylated α-synuclein and ubiquitinated protein aggregates, while

the other did not, indicating the formation of α-synuclein two strains. Notably, the former

α-synuclein strain inhibited proteasome activity and co-precipitated with 26S proteasome

complex. Further examination indicated that structural differences in the C-terminal region of

α-synuclein strains lead to different effects on proteasome activity. These results provide a

possible molecular mechanism to account for the different pathologies induced by different

α-synuclein strains.

Introduction

Misfolding and aggregation of normally soluble proteins are common pathological features of

many neurodegenerative diseases, including Alzheimer's, Parkinson's, Creutzfeldt–Jacob and

Huntington's diseases (Ross & Poirier, 2004). For example, Parkinson's disease (PD), dementia

with Lewy bodies (DLB) and multiple system atrophy (MSA) are characterized by

accumulation of misfolded α-synuclein in neuronal and/or glial cells, and therefore these

diseases are termed α-synucleinopathies. In PD and DLB, α-synuclein pathologies are mainly

observed in neurons in the form of Lewy bodies (LBs) and Lewy neurites (LNs) (Baba et al.,

1998), while glial cytoplasmic inclusions (GCIs) are seen in oligodendrocytes in MSA

(Wakabayashi, Yoshimoto, Tsuji, & Takahashi, 1998). The abnormal α-synuclein observed in

brains of patients is accumulated as fibrous or filamentous forms with cross-β structures (Araki

et al., 2019; Spillantini et al., 1997), existing in phosphorylated and partially ubiquitinated states

(Fujiwara et al., 2002; Hasegawa et al., 2002). These abnormal α-synuclein species exhibit

seeding activity for prion-like conversion, being similar in this respect to the infectious forms of

prion protein (PrP) causing Creutzfeldt-Jakob disease (CJD) and bovine spongiform

encephalopathy (Goedert, 2015). Various other neurodegenerative disease-related proteins,

including amyloid-β, tau and TDP-43, can also propagate through neural networks in a similar

manner.

α-Synuclein is a natively unfolded protein of 140 amino acid residues, normally found in both

soluble and membrane-associated fractions and localized in synaptic termini. Although its

physiological function has not been fully clarified, it appears to be involved in the regulation of

SNARE complex and in dopamine production. Disease-linked missense mutations and

multiplication of the *SNCA* gene encoding α-synuclein have been reported in familial forms of

α-synucleinopathies, indicating that structural changes and overexpression of α-synuclein

protein are involved in the development of α-synucleinopathies (Wong & Krainc, 2017).

Recombinant soluble α-synuclein proteins purified from bacterial cells form amyloid-like fibrils

that are morphologically and physicochemically similar to those observed in patients' brains

(Araki et al., 2019; Goedert, 2015). These synthetic α-synuclein fibrils can act as seeds and

induce seeded aggregation of α-synuclein in cultured cells or primary cultured neurons, as well

as in animal brains. Intracerebral inoculation of synthetic α-synuclein fibrils induces

phosphorylated and ubiquitinated α-synuclein pathologies even in wild-type (WT) mice (Luk et

al., 2012; Masuda-Suzukake et al., 2013).   It has also been reported that extracts from brains of

patients with α-synucleinopathies induce α-synuclein pathologies in cellular and animal models

(Bernis et al., 2015; Watts et al., 2013). In addition, recent studies have suggested that

α-synuclein strains with distinct conformations exist, which is a characteristic of prions

(Bousset et al., 2013; Gribaudo et al., 2019; Guerrero-Ferreira et al., 2019; Peelaerts &

Baekelandt, 2016; Peelaerts et al., 2015; Shahnawaz et al., 2020; Woerman et al., 2019).

Synthetic α-synuclein fibrils formed under different physiological conditions *in vitro* show

distinct seeding activities and cytotoxicity in cultured cells and rat brains. Furthermore, MSA

brain extracts exhibit distinct infectivity compared to PD or control brain extracts in cultured

cells or mice expressing mutant A53T or WT α-synuclein (Lau et al., 2020; Peng et al., 2018;

Prusiner et al., 2015; Woerman et al., 2019; Woerman et al., 2015).

These observations support the idea that α-synuclein shows prion-like behavior, because they

can be accounted for by a typical hallmark of the prion phenomenon, i.e., the presence of strains.

In prion diseases, the variety of strains that can be differentiated in terms of the clinical signs,

incubation period after inoculation, and the vacuolation lesion profiles in the brain of affected

animals is due to structural differences of PrP aggregates, as identified by biochemical analyses

including glycosylation profile, electrophoretic mobility, protease resistance, and

sedimentation.   These PrP strains are thought to correspond to different conformations of PrP

aggregates, as demonstrated for the yeast prion [*PSI*[+]], which induces aggregates of Sup35p

(Ohhashi, Ito, Toyama, Weissman, & Tanaka, 2010). Thus, as the case of prion disease,

differences of lesions among α-synucleinopathies are thought to be caused by conformational

heterology of α-synuclein assemblies, probably amyloid-like fibrils (Lau et al., 2020;

Schweighauser et al., 2020; Shahnawaz et al., 2020). However, little is known about how

conformational differences of protein aggregates induce a variety of lesions, not only in prion

disease, but also in other neurodegenerative diseases, such as α-synucleinopathies.

In this study, we prepared two α-synuclein assemblies from identical wild-type α-synuclein

monomer under different conditions, and established that they have distinct conformations, i.e.,

we succeeded to generating two α-synuclein fibrils from the same monomer. We examined their

seeding abilities to convert endogenous soluble α-synuclein monomers into phosphorylated

aggregates in mice and primary-cultured neurons, indicating the formation of two α-synuclein

strains. Notably, only one strain induced the accumulation of ubiquitinated proteins, as well as

phosphorylated α-synuclein aggregates, indicating the ability of this strain to inhibit proteasome

activity. Moreover, only this strain strongly inhibited proteasome activity and co-precipitated

with purified 26S proteasome complex *in vitro*. Structural studies suggested that the C-terminal

region plays a key role in the different properties of the two strains. Taken together, these results

provide a possible molecular mechanism to account for the different lesions induced by distinct

α-synuclein strains.

Results

Formation of Amyloid-like Fibrils from α-Synuclein Proteins in vitro

We generated two distinct α-synuclein assemblies from the identical purified α-synuclein

monomer using the method described in the previous report, with minor modifications (*Figure*

1A) (Bousset et al., 2013; Lau et al., 2020). Specifically, we prepared   recombinant α-synuclein

monomer and agitated it in the presence or absence of salt at a physiological concentration. In

the presence of physiological salt (30 mM Tris, pH 7.5, 150 mM KCl), the monomer formed a

cloudy solution of assemblies with higher turbidity, while in the absence of salt (30 mM Tris,

pH 7.5), a clear solution containing assemblies with lower turbidity was formed (*Figure 1A and*

B). Almost all of the α-synuclein was present as aggregates rather than soluble oligomers under

both conditions (*Figure 1-figure supplement 1*) (Thibaudeau, Anderson, & Smith, 2018). Both

assemblies showed fibrillar morphology, but the previously reported ribbon-like morphology

was not observed (*Figure 1C*) (Bousset et al., 2013). Therefore, we will refer to the former

assemblies as α-synuclein fibrils (+) and the later assemblies as α-synuclein fibrils (-). Both

fibrils were stained with Thioflavin T and Congo red, indicating their amyloid-like nature

(*Figure 1D and 1E*). The differences of Thioflavin T fluorescence and Congo red binding of

these α-synuclein fibrils suggested that these fibrils have a different structures each other. Next,

we examined the in vitro seeding activity of these fibrils and found that α-synuclein fibrils (+)

showed higher seeding activity than α-synuclein fibrils (-) *in vitro* (*Figure 1F*). These results

indicated that we had successfully prepared two distinct types of α-synuclein fibrils from the

same monomer.

Formation of Phosphorylated α-Synuclein Pathology by Injection of α-Synuclein Strains

into Mouse Brain

We next investigated whether there was a strain-dependent difference of prion-like propagation

in mouse brain. We injected α-synuclein fibrils (+) and α-synuclein fibrils (-) into striatum of

wild-type mice, and after one month, we examined the accumulation of phosphorylated

α-synuclein deposits resembling those observed in patients' brains (Luk et al., 2012;

Masuda-Suzukake et al., 2013). In contrast to the *in vitro* seeding activities, α-synuclein fibrils

(-) induced Lewy body/Lewy neurite-like abnormal phosphorylated α-synuclein deposits

through the mouse brain, including cortex, striatum and corpus callosum, whereas few

phosphorylated α-synuclein deposits were induced by α-synuclein fibrils (+) (*Figure 2A* and

Figure 2-figure supplement 1). We used three mice each for this experiment and confirmed the

reproducibility (*Figure 2-figure supplement 2*). The deposits induced by α-synuclein fibrils (-)

were also positive for ubiquitin staining in cortex, like Lewy bodies and Lewy neurites (*Figure*

2B). Compared with phosphorylated α-synuclein, which is the most representative marker in the

α-synucleinopathies, ubiquitin staining is weaker, thus we could find only few ubiquitin positive

deposits in striatum and corpus callosum (data not shown). Taken together, there was a

strain-dependent difference of α-synuclein fibril formation from the identical α-synuclein

monomer when the two strains were inoculated into the mouse brain, and α-synuclein fibrils (-)

had higher prion-like seeding activity than α-synuclein fibrils (+) in mouse brain, contrary to the

in vitro seeding results (*Figure 1F*).

Formation of Phosphorylated α-Synuclein Pathology Induced by the Two α-Synuclein

Strains in Cultured Neurons

To further study the difference in the formation of pathological α-synuclein aggregates in

neurons, we compared the ability of the two α-synuclein strains to induce seed-dependent

aggregation of α-synuclein in primary mouse cortical cells. When primary mouse cortical cells

from non-transgenic mice were treated with α-synuclein fibrils for two weeks, we observed a

dramatic increase of phosphorylated α-synuclein accumulation, which is also positive for

ubiquitin, only in the case of α-synuclein fibrils (-), while little accumulation was seen with

α-synuclein fibrils (+) (*Figure 3A*). These results are consistent with those observed in the

mouse brain. To investigate whether these accumulations occur in neurons, we performed

double staining for phosphorylated α-synuclein and neuronal markers or an astrocyte marker

using cells treated with α-synuclein fibrils (-) (*Figure 3-figure supplement 1*). We found that

phosphorylated α-synuclein accumulations were seen in neurons, not in astrocytes. Thus,

α-synuclein fibrils (-) could induce phosphorylated α-synuclein accumulations in primary mouse

cortical neurons.

Next, we performed biochemical analysis of detergent-insoluble α-synuclein prepared from

these cells. Phosphorylated α-synuclein were accumulated in cells treated with both α-synuclein

fibrils (-) and α-synuclein fibrils (+) (*Figure 3B and C*). However, α-synuclein fibrils (-) induced

a greater accumulation of phosphorylated α-synuclein than did α-synuclein fibrils (+) (*Figure

3D*). The phosphorylated and aggregated α-synuclein in these cells was found to be endogenous

mouse α-synuclein, indicating that the introduced human α-synuclein fibrils worked as seeds

(*Figure 3B*). We also examined the accumulation of detergent-insoluble ubiquitinated proteins

and found that not only ubiquitinated α-synuclein, but also other ubiquitinated proteins were

accumulated in cells treated with α-synuclein fibrils (-). There was no significant increase of

ubiquitinated protein accumulation in cells treated with α-synuclein fibrils (+) (*Figure 3B and

D*). These results indicated that only α-synuclein fibrils (-) induced much accumulation of

phosphorylated α-synuclein and ubiquitinated proteins in primary mouse cortical neurons, in

accordance with the findings in mouse brain. Seed dependent aggregation of α-synuclein must

depend on the efficiency of fibril uptake by cells. We confirmed whether the two α-synuclein

fibrils were taken up into primary mouse cortical cells with similar efficiency. Cells were

treated with the two α-synuclein fibrils or buffer. A day after treatment, we treated cells with

trypsin for digestion of extracellular fibrils, collected them and examined the α-synuclein fibrils

taken up into the cells by western blotting of sarkosyl insoluble fractions. We found no

α-synuclein signal in pellet fraction of cells with buffer control indicating that detected

α-synuclein signals were originated from added fibrils. We found α-synuclein fibrils (-), which

induced much α-synuclein accumulation, were taken up into the cells lesser than α-synuclein

fibrils (+), demonstrating that α-synuclein fibrils (-) efficiently caused the accumulation of

α-synuclein and ubiquitinated proteins in spite of their inefficient uptake into the cells (*Figure*

3-figure supplement 2).

Different Interactions of α-Synuclein Strains with 26S Proteasome

The above results motivated us to examine proteasome activity in the presence of these two

types of α-synuclein fibrils. We purified 26S proteasome complex from budding yeast

expressing FLAG-tagged Rpn11p, a subunit of 19S regulatory complex (*Figure 4-figure*

supplement 1)(Saeki, Isono, & Toh-E, 2005). The activity of the purified 26S proteasome was

examined in the presence or absence of α-synuclein fibrils. The eukaryotic proteasome has three

active subunits, β1, β 2 and β 5, each displaying a specific catalytic activity, trypsin-like,

chymotrypsin-like and caspase-like activity, respectively. Thus, we examined the

chymotrypsin-like (*Figure 4A*), caspase-like (*Figure 4B*) and trypsin-like (*Figure 4C*) activities

of 26S proteasome in the presence of these α-synuclein fibrils using each fluorogenic peptide

substrate in vitro. The catalytic activity of 26S proteasome was drastically impaired in the

presence of α-synuclein fibrils (-), whereas α-synuclein fibrils (+) were ineffective (*Figure*

4A-C). These results accord with our mouse and primary-cultured neuron data, i.e., only

α-synuclein fibrils (-) could induce much accumulation of phosphorylated α-synuclein and

ubiquitinated proteins. Co-aggregation of functional proteins with misfolded protein aggregates

may cause the impairment of their function. Therefore, we investigated the interaction of 26S

proteasome with the fibrils. α-Synuclein fibrils were mixed with purified 26S proteasome, and

the mixture was centrifuged. The supernatants and precipitates were analyzed by western

blotting. We found that both α-synuclein fibrils (-) and α-synuclein fibrils (+) were fractionated

to the pellet fractions (*Figure 4D*). However, only α-synuclein fibrils (-) co-precipitated with

26S proteasome, while α-synuclein fibrils (+) did not (*Figure 4E*). Taken together, these results

indicate that only α-synuclein fibrils (-) interact with 26S proteasome and impair the proteasome

activity.

If α-synuclein fibrils (-) bind other proteins, such as 26S proteasome, while α-synuclein fibrils

(+) do not, the structures of the two types of fibrils should be distinguishable. Indeed, we did

observe slight differences between these two types of fibrils (*Figure 1*). Amyloid-like fibrils are

composed of core regions consisting of β-sheet-rich rigid structure, and exposed regions, which

might interact with other molecules. The core regions tend to be resistant to protease attack,

whereas exposed regions tend to be easily digested by protease. To identify the core regions and

exposed regions of these α-synuclein fibrils, we carried out limited proteolysis with proteinase

K, followed by mass spectrometric analysis (*Figure 4F*) (Suzuki, Shimazu, & Tanaka, 2012).

We found that α-synuclein fibrils (-) have a smaller core region (amino acid residues 39-96 and

40-94; m/z 5656 and 5268, respectively) corresponding to the NAC region (amino acid residues

61-95), which had previously been reported as the core region of α-synuclein fibrils

(Guerrero-Ferreira et al., 2018; Li et al., 2018). In contrast, α-synuclein fibrils (+) had a larger

core region, extending to the C-terminal regions (residues 31-109, 28-118 and 25-132; m/z 7860,

9115 and 10907, respectively), indicating that α-synuclein fibrils (-) have amyloid structure with

a more exposed C-terminal region than α-synuclein fibrils (+). To confirm these results, we

performed dot-blot analysis using various antibodies against different regions of α-synuclein

protein. Native α-synuclein fibrils were spotted on nitrocellulose membranes and detected by

applying various α-synuclein antibodies. Antibodies raised against the N-terminal region of

α-synuclein (residues 1-10) and the NAC region (91-99) almost equally recognized both types

of α-synuclein fibrils, indicating that they have similar structure in the N-terminal and NAC

regions (*Figure 4G*). However, as expected from the above data, the antibodies raised against

C-terminal regions (amino acid 115-122 and 131-140) bound more strongly to α-synuclein

fibrils (-) than to α-synuclein fibrils (+), supporting the idea that the C-terminal region of

α-synuclein fibrils (-) is more exposed than that of α-synuclein fibrils (+). Thus, we considered

that the C-terminal region of α-synuclein fibrils (-) might interact with 26S proteasome and

impair its activities. Indeed, we previously reported that C-terminally truncated α-synuclein

fibrils induced pathology in mouse brain less potently than did full-length α-synuclein fibrils,

even though the C-terminally truncated α-synuclein fibrils had higher seeding activity in vitro

(Terada et al., 2018). To confirm this, we next examined the seeding activity of C-terminally

truncated fibrils formed in the absence of salt in primary-cultured neurons. We prepared

C-terminally truncated α-synuclein monomer (residues 1-120) and agitated it in the absence of

salt. The resulting assemblies showed fibrillar morphology and thioflavin T binding (data not

shown), and we refer to them as αSΔC20 fibrils (-). We treated primary mouse cortical cells

with αSΔC20 fibrils (-) and examined the accumulation of phosphorylated α-synuclein and

ubiquitinated proteins. As expected, we observed little accumulation of these proteins (*Figure*

5A and B). Next, we examined the activities of 26S proteasome in the presence of αSΔC20

fibrils (-) as shown in Figure 4A-C. The catalytic activity of 26S proteasome was not impaired

in the presence of αSΔC20 fibrils (-) (*Figure 5C*). We also investigated the interaction of 26S

proteasome with αSΔC20 fibrils (-) as shown in Figure 4D and E and found that αSΔC20 fibrils

(-) did not co-precipitated with 26S proteasome (*Figure 5D and E)*. Considering all these results,

we can conclude that the C-terminal region of α-synuclein is exposed only in α-synuclein fibrils

(-) and this region interacts with 26S proteasome and impairs its activity.

Discussion

According to the prion hypothesis, differences in disease symptoms and lesions are caused by

differences in the conformation of strains. Therefore, if α-synuclein is prion-like, differences in

the structure of α-synuclein aggregates should cause the differences in the lesions observed in

various α-synucleopathies. Recent analyses have shown that introduction of extracts from brains

of patients with PD and MSA into mice and cells induces different pathologies (Prusiner et al.,

2015; Tarutani, Arai, Murayama, Hisanaga, & Hasegawa, 2018; Woerman et al., 2019;

Woerman et al., 2015). Similar results were obtained with brain extracts of patients with

tauopathies, in which tau is abnormally accumulated (Narasimhan et al., 2017; Saito et al.,

2019). Moreover, it has been clarified that TDP-43 aggregates have a different structure

depending on the disease, and different lesions appear when the patients' brain extracts are

introduced into cells or animals (Laferriere et al., 2019; Nonaka et al., 2013; Porta et al., 2018;

Tsuji et al., 2012). As for α-synuclein, it has been reported that aggregates having different

structures are formed *in vitro* (Bousset et al., 2013; Shahnawaz et al., 2020), and the lesions

caused by introduction of aggregates having different structures into cells and rodents are

different (Gribaudo et al., 2019; Lau et al., 2020; Peelaerts et al., 2015). However, it has

remained unclear what molecular mechanism might lead to the differences in phenotypes

induced by different protein aggregates.

In this study, we confirmed that different pathologies were caused by two α-synuclein strains

with different structures, and we also found that these α-synuclein strains differ in their ability

to inhibit 26S proteasome activity. It has already been reported that protein aggregates that can

cause neurodegenerative diseases may interact with proteasome and inhibit its activity (Bence,

Sampat, & Kopito, 2001; Thibaudeau et al., 2018; Zondler et al., 2017). However, it is a

noteworthy finding in this work that one of the two differently structured fibrils formed from

identical monomer under different conditions inhibited proteasome, while the other did not.

This clearly raises the possibility that inhibition of proteasome by abnormal α-synuclein plays a

role in the pathology. These results provide a possible molecular mechanism to account for the

different lesions induced by distinct α-synuclein strains. Considering that PD and MSA have

different structures of accumulated α-synuclein (Klingstedt et al., 2019; Lau et al., 2020;

Prusiner et al., 2015; Schweighauser et al., 2020; Shahnawaz et al., 2020; Woerman et al., 2019;

Woerman et al., 2015), it seems highly likely that the structures of α-synuclein fibrils determine the lesions observed in these diseases.

Protein inclusions found in many neurodegenerative diseases are often ubiquitinated, and this is consistent with reports that ubiquitin proteasome systems are impaired in neurons of patients with neurodegenerative diseases (Bence et al., 2001). Recently, it was reported that protein aggregates co-aggregate with proteasome at the molecular level (Guo et al., 2018). Furthermore, α-synuclein oligomers, Aβ oligomers and polyglutamine inhibit 20S proteasome (Thibaudeau et al., 2018). In our α-synuclein fibril formation experiments, almost all the proteins formed insoluble aggregates, and soluble oligomers could not be isolated (*Figure 1-figure supplement 1*). Thus, our results indicate that protein aggregates may also show proteasome-inhibitory activity, and the extent of the inhibitory activity depends on the structure of the aggregates. These facts indicate that pathogenic protein oligomers and aggregates have different effects on cells depending on their structure, and both may be toxic.

Our results suggest that the difference in the structure of the two strains is mainly at the C-terminal region. The C-terminal region (residues 96-140) of α-synuclein is acidic and contains negatively charged residues, including aspartate and glutamate, as well as proline residues. When intermolecular repulsion at the C-terminal region is weakened by changes in ionic strength, exposure of the C-terminal region decreases, and more tightly packed α-synuclein fibrils are formed in the presence of salt. On the other hand, in the absence of salt, intermolecular repulsion at the C-terminal region causes the formation of α-synuclein fibrils in which the C-terminal region is exposed. Only this latter type of α-synuclein fibrils can interact with 26S proteasome complex in vitro, causing inhibition of 26S proteasome activity. If this also occurs in cells, abnormal α-synuclein aggregates, which might be partially degraded by proteasome, would be accumulated (*Figure 6*). More analysis is desired to show that α-synuclein fibrils (-), but not fibrils (+), interact with the mammalian proteasome and

inhibit its activity in mouse primary neurons or brains of injected mice. It may also be

necessary to examine whether protein aggregates co-aggregate with the proteasome in

the patient's brain. Consequently, phosphorylated and ubiquitinated α-synuclein aggregates

would be accumulated in neurons and then presumably would propagate throughout the brain in

a prion-like manner. Our results imply that inhibition of proteasome activity by protein

aggregates is critical for prion-like propagation of these protein aggregates. Thus, inhibiting the

interaction between aggregates and the proteasome may be a promising therapeutic strategy for

neurodegenerative diseases.

Forming fibrils from purified monomeric protein and introducing them into cells or animals

might be a useful approach for analyzing the mechanisms of propagation and toxicity of

diseases in which protein aggregates accumulate. However, based on our results and previous

reports, it seems possible that the results obtained might vary markedly in response to even

slight differences in the structure of the introduced fibrils (Bousset et al., 2013; Gribaudo et al.,

2019; Peelaerts et al., 2015). Introducing brain extract from a patient with a neurodegenerative

disease into cells or animals would also be one approach for identifying the cause of the disease

and developing a treatment, but even in patients with the same disease, the results might vary

due to slight differences in the structure of the accumulated protein aggregates. Therefore, when

conducting such experiments, it is extremely important to analyze the structure of the protein

aggregates.

In this study, we found that the degree of interaction with proteasome and the inhibition of

proteasome activity vary depending on the strain of α-synuclein aggregates. The relationship

between proteasome inhibition and the induction of the pathology supports the hypothesis that

prion-like activity of α-synuclein aggregates contributes to disease progression.

2 Materials and methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*Mus musculus*)	C57BL/6J	CLEA, Japan, Inc.		WT mouse
Genetic reagent (*Saccharomyces cerevisiae*)	*RPN11-FLAG*x3	This paper		
strain, strain background (*Escherichia coli*)	BL21(DE3)	Sigma-Aldrich	Cat. #: 69450	
recombinant DNA reagent	pRK172-α-synuclein	PMID: 8194594		
recombinant DNA reagent	pRK172-α-synucleinΔC20	PMID: 30030380		
antibody	anti-α-Synuclein-pSer129 #64 (mouse monoclonal)	FUJIFILM Wako Chemicals	Cat. #: 015-25191	
antibody	anti-α-Synuclein 1-10 (rabbit polyclonal)	Cosmo Bio	Cat. #: TIP-SN-P01	
antibody	anti-α-Synuclein 131-140 (rabbit polyclonal)	Cosmo Bio	Cat. #: TIP-SN-P09	

antibody	anti-α-Synuclein 91-99 (mouse monoclonal)	BD Bioscience	Cat. #: 610787	
antibody	anti-α-Synuclein-pSer129 (rabbit monoclonal)	Abcam	Cat. #: 51253	
antibody	anti-α-Synuclein 115-122 (mouse monoclonal)	Santa Cruz Biotechnology	Cat. #: 58480	
antibody	anti-α-Synuclein D37A6 (rabbit monoclonal)	Cell Signaling Technology	Cat. #: 4179	
antibody	anti-α-Synuclein-pSer129 (rabbit polyclonal)	PMID: 23466394		
antibody	anti-α-Synuclein (Syn205) (mouse monoclonal)	Cell Signaling Technology	Cat. #: 2644	
antibody	anti-ubiquitin (rabbit polyclonal)	Proteintech	Cat. #: 10201-2-AP	
antibody	anti-ubiquitin (rabbit polyclonal)	Dako	Cat. #: Z0458	
antibody	anti-GAPDH (mouse monoclonal)	Sigma-Aldrich	Cat. #: MAB374	
antibody	anti-FLAG (mouse monoclonal)	Sigma-Aldrich	Cat. #: F1804	

antibody	anti-GFAP (rabbit monoclonal)	Cell Signaling Technology	Cat #: 12389	
antibody	anti-Neurofilament-L (rabbit monoclonal)	Cell Signaling Technology	Cat #: 2837	
antibody	anti-NeuN (rabbit monoclonal)	abcam	Cat. #: ab177487	
antibody	anti-tau (tauC) (rabbit polyclonal)	PMID: 26374846		
antibody	Goat anti-rabbit IgG, Alexa Fluor 488	Thermo Fisher	Cat. #: A11008	
antibody	Goat anti-mouse IgG, Alexa Fluor 568	Thermo Fisher	Cat. #: A11004	
antibody	Goat anti-rabbit IgG, HRP	Thermo Fisher	Cat. #: A27036	
antibody	Goat anti-mouse IgG, HRP	Thermo Fisher	Cat. #: A28177	
antibody	Goat anti-rabbit IgG, biotin	Vector laboratories	Cat. #: BA-1000	
software, algorithm	EZR	PMID: 23208313		
other	hematoxylin	Muto Pure Chemicals	Cat. #: 30002	
other	DAPI stain	Vector laboratories	Cat #: H-1500	

Expression and purification of Recombinant Wild-type and C-Terminally Truncated

Human α-Synuclein

Full-length and C-terminally truncated α-synuclein encoded in pRK172 plasmids were

transformed into *Escherichia coli* BL21 (DE3).   Recombinant proteins were purified as

described previously (Nonaka, Watanabe, Iwatsubo, & Hasegawa, 2010). Protein concentration

was determined by HPLC.

Preparation of α-Synuclein Fibrils

α-Synuclein fibrils were prepared as follows. Purified recombinant α-synuclein proteins were

dissolved in 30 mM Tris-HCl, pH 7.5, containing 150 mM KCl and 0.1% NaN_3, to a final

concentration of 6 mg/ml. The samples were incubated at 37 °C under rotation at 20 rpm for 7

days. The assembled α-synuclein was sonicated with an ultrasonic homogenizer (VP-5S,

TAITEC) in 30 mM Tris-HCl, pH 7.5. For the measurement of turbidity, the resultant

α-synuclein assemblies were diluted to 1 mg/ml and the absorbance at 440 nm was measured.

Transmission Electron Microscopy

α-Synuclein fibrils were diluted to 15 μM in 30 mM Tris-HCl, pH 7.5, plated on carbon-coated

300-mesh copper grids (Nissin EM), and stained with 2% [v/v] phosphotungstate. Micrographs

were recorded on a JEM-1400 electron microscope (JEOL).

Thioflavin T-binding Assay

The degree of fibrillation was measured in terms of Thioflavin T (ThT) fluorescence intensity,

which increases when ThT binds to amyloid-like fibrils. The samples (7.5 μM) were incubated

with 20 μM ThT in 30 mM Tris-HCl buffer (pH 7.5) for 30 min at 37 °C. Fluorometry was

performed using a microplate reader (Infinite 200, TECAN, excitation 442 nm, emission 485

nm).

Congo Red Binding Assay

The binding of Congo red was measured as described previously (Suzuki et al., 2012).

α-Synuclein monomer and fibrils (37.5 μM) were mixed with Congo red (1 μM) and incubated

for 1 hour at 37 °C. Absorbance between 400 and 700 nm was measured with a plate reader

(Infinite 200, TECAN). The binding of Congo red was calculated as $A_{540}/25296-A_{477}/46306$.

α-Synuclein Aggregation Assay

Full-length α-synuclein aggregation experiments were performed using a microplate reader

(Infinite 200, TECAN, excitation 442 nm, emission 485 nm) and monitored by measuring ThT

fluorescence in the absence or presence of 5% α-syn fibril seeds. All experiments were

performed at 37 °C, under quiescent conditions in flat-bottomed 96-well black plates

(Sumitomo Bakelite) sealed with MicroAmp Optical Adhesive Film (Applied Biosystems). The

reaction mixture consisted of PBS containing 1 mg/ml α-synuclein monomer. During

experiments under quiescent conditions, ThT fluorescence was read every 2 min.

Primary-cultured cells and Introduction of α-Synuclein Proteins into Cells

Dissociated cultures of embryonic (E15) mouse cortical cells were prepared from pregnant

C57BL/6 mice using Neuron Dissociation Solutions (FUJI FILM Wako) according to the

manufacturer's protocol. Briefly, dissected brain was digested with enzyme solution for 30 min

at 37 °C, then centrifuged. Dispersion solution was added and tissues were suspended, then

isolation solution was added. Cells were collected by centrifugation, resuspended and plated on

poly-L-lysine-coated cover glass or plates. Cells were maintained at 37 °C in 5% CO_2 in

Neurobasal Medium (Gibco) supplemented with 1x B27 and 1x Glutamax. Cells were cultured

for 7 days in vitro (DIV) in 6-well plates, and then treated with sonicated α-synuclein fibrils

diluted in culture medium. Cells were collected or fixed at 14 days post treatment (21 DIV).

Purification of 26S Proteasome from Budding Yeast

Yeast 26S proteasome was purified as described previously (Saeki et al., 2005). Briefly, the

yeast strain (BY4741, *Rpn11-FLAGx3::KanMX*) was cultured in YPD for 2 days, harvested,

washed and stocked at -80°C. Buffer A'' (50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 10%

glycerol, 4 mM ATP, 10 mM $MgCl_2$) and glass beads were added and the cells were lysed in a

Beads Shocker (Yasui Kikai). Cell debris was removed by centrifugation and then anti-FLAG

M2 antibody-conjugated agarose beads (Sigma) were added. The mixture was incubated for 2

hours at 4°C. Agarose beads were washed with Buffer A'' and Buffer A'' containing 0.1% Triton,

and proteasome was eluted by adding 3x FLAG peptide (400 μg/ml) (Sigma) in Buffer A''.

26S Proteasome Activity Assays

Inhibition of α-synuclein fibrils on proteasome activity was measured using fluorogenic

peptides in 96-well black flat-bottomed plates. 26S proteasome (10 μg/ml) was added to

α-synuclein fibrils (35 μM, calculated based on the monomer protein concentration), and the

mixture was incubated in buffer A'' containing 100 μM fluorogenic substrate (suc-LLVY-mca,

Z-LLE-mca, Peptide Institute) for 60 minutes at 37°C or 50 μM fluorogenic substrate

(boc-LRR-mca, Peptide Institute) for 10 minutes at 37°C. Fluorescence was measured before

and after incubation (Infinite 200, TECAN, excitation 360 nm, emission 440 nm). The rate of

increase in fluorescence intensity is regarded as representing proteasome activity.

Binding Assay of Proteasome with α-Synuclein Fibrils

α-Synuclein fibrils were mixed with purified 26S proteasome in Buffer A'', and the mixture was

centrifuged at 21,500 x g for 20 minutes. The supernatant and pellet fractions were analyzed by

the western blotting.

Immunocytochemistry

Introduction of α-synuclein fibrils were conducted as described above, using mouse primary

cultured cells grown on coverslips. At 2 weeks after introduction of fibrils, the cells were fixed

with 4% paraformaldehyde and treated with the indicated primary antibodies. After incubation

overnight, the cells were washed and treated with secondary antibodies conjugated with Alexa

Fluor for 1 hour. The cells were mounted with DAPI to counterstain nuclear DNA and analyzed

with a BZ-X710 (Keyence) and BZ-X analyzer (Keyence).

α-Synuclein Inoculation into Mice and Immunohistochemistry of Mouse Brains

Ten-week-old, male C57BL/6J mice were purchased from CLEA Japan, Inc. All experimental

protocols were performed according to the recommendations of the Animal Care and Use

Committee of Tokyo Metropolitan Institute of Medical Science. α-Synuclein samples (150 μM,

5 μl) were injected into the striatum (anterior-posterior, 0.2 mm; medial-lateral, -2.0 mm;

dorsal-ventral, 2.6 mm). Inoculation into mouse brain was performed as described previously

(Masuda-Suzukake et al., 2013). Non-injected wild- type mice were used as negative controls.

One month after inoculation, mice were deeply anesthetized with isoflurane (Pfizer) and

sacrificed, and the brain was perfused with 0.1 M phosphate buffer. Sections were fixed in 4%

paraformaldehyde and preserved in 20% sucrose in 0.01 M phosphate buffered saline, pH 7.4.

were cut serially on a freezing microtome at 30 μm thickness. The sections were then mounted

on glass slides. Sections were incubated with 1% H_2O_2 for 30 min to eliminate endogenous

peroxidase activity and were treated with 100% formic acid (Wako) for 10 min for antigen

retrieval and washed under running tap water. Immunohistochemistry with polyclonal antibody

1175 (1:1,000) directed against α-synuclein phosphorylated at Ser129 or anti-ubiquitin (DAKO,

1:10,000) were performed as described previously (Masuda-Suzukake et al., 2013). Antibody

labeling was performed by incubation with biotinylated goat anti-rabbit IgG for 3 hours. The

antibody labeling was visualized by incubation with avidin-biotinylated horseradish peroxidase

complex (ABC Elite, Vector Laboratories, 1:1,000) for 3 hours, followed by incubation with a

solution containing 0.01% 3,3'-diaminobenzidine, 0.05 M imidazole and 0.00015% H_2O_2 in

0.05 M Tris-HCl buffer, pH 7.6. Counter nuclear staining was performed with hematoxylin

(Muto Pure Chemicals). The sections were then rinsed with distilled water, treated with xylene,

and coverslipped with Entellan (Merck). Images were analyzed with a BZ-X710 (Keyence) and

BZ-X analyzer (Keyence). We examined three mice for each sample and each mouse yielded

highly similar results. We examined at least 6 slices for each mouse and confirmed the similar

results. Animal experiments were done by the experimenter who is blind.

Sedimentation Analysis and Western blotting

Cells were harvested, collected by centrifugation (2,000 $\times$ g, 5 min) and washed with PBS. The

cellular proteins were extracted by sonication in 200 μl of buffer A68 (10 mM Tris-HCl, pH 7.5,

1 mM EGTA, 10% sucrose, 0.8 M NaCl containing sarkosyl (final 1%, w/v) and protease

inhibitor (Roche). After ultracentrifugation at 135,000 $\times$ g for 20 min at 25 °C, the supernatant

was collected as sarkosyl-soluble fraction, and the protein concentration was determined by

Bradford assay. The pellet was solubilized in 50 μl of SDS-sample buffer. Both sarkosyl-soluble

and insoluble fractions were analyzed by immunoblotting with appropriate antibodies as

indicated.

Fibril uptake assay

14 Mouse cortical cells were prepared and treated with the α-synuclein fibrils as described above. A

15 day after treatment, cells were washed with PBS and treated with 0.25% trypsin at 37 °C for 10

16 minutes, then cells were collected. Sedimentation analysis and western blotting were performed

17 as described above.

Statistical analysis

19 Student's t-test was performed when comparing 2 groups. One-way ANOVA and Tukey's post

20 hoc test were performed with EZR when comparing 3 groups (Kanda, 2013). P values below

21 0.05 were considered to be statistically significant.

Acknowledgments

We would like to thank all members of the laboratory for helpful discussion.

Competing Interests

The authors declare no competing financial interests.

References

Araki, K., Yagi, N., Aoyama, K., Choong, C. J., Hayakawa, H., Fujimura, H., . . . Mochizuki,
H. (2019). Parkinson's disease is a type of amyloidosis featuring accumulation of
amyloid fibrils of alpha-synuclein. *Proc Natl Acad Sci U S A*.
doi:10.1073/pnas.1906124116

Baba, M., Nakajo, S., Tu, P. H., Tomita, T., Nakaya, K., Lee, V. M., . . . Iwatsubo, T. (1998).
Aggregation of alpha-synuclein in Lewy bodies of sporadic Parkinson's disease and
dementia with Lewy bodies. *Am J Pathol, 152*(4), 879-884.

Bence, N. F., Sampat, R. M., & Kopito, R. R. (2001). Impairment of the ubiquitin-proteasome
system by protein aggregation. *Science, 292*(5521), 1552-1555.
doi:10.1126/science.292.5521.1552

Bernis, M. E., Babila, J. T., Breid, S., Wusten, K. A., Wullner, U., & Tamguney, G. (2015).
Prion-like propagation of human brain-derived alpha-synuclein in transgenic mice
expressing human wild-type alpha-synuclein. *Acta Neuropathol Commun, 3*, 75.
doi:10.1186/s40478-015-0254-7

Bousset, L., Pieri, L., Ruiz-Arlandis, G., Gath, J., Jensen, P. H., Habenstein, B., . . . Melki, R.
(2013). Structural and functional characterization of two alpha-synuclein strains.
Nat Commun, 4, 2575. doi:10.1038/ncomms3575

Fujiwara, H., Hasegawa, M., Dohmae, N., Kawashima, A., Masliah, E., Goldberg, M. S., . . .
Iwatsubo, T. (2002). alpha-Synuclein is phosphorylated in synucleinopathy lesions.
Nat Cell Biol, 4(2), 160-164. doi:10.1038/ncb748

Goedert, M. (2015). NEURODEGENERATION. Alzheimer's and Parkinson's diseases: The
prion concept in relation to assembled Abeta, tau, and alpha-synuclein. *Science,*
349(6248), 1255555. doi:10.1126/science.1255555

Gribaudo, S., Tixador, P., Bousset, L., Fenyi, A., Lino, P., Melki, R., . . . Perrier, A. L. (2019).
Propagation of alpha-Synuclein Strains within Human Reconstructed Neuronal
Network. *Stem Cell Reports, 12*(2), 230-244. doi:10.1016/j.stemcr.2018.12.007

Guerrero-Ferreira, R., Taylor, N. M., Arteni, A. A., Kumari, P., Mona, D., Ringler, P., . . .
Stahlberg, H. (2019). Two new polymorphic structures of human full-length
alpha-synuclein fibrils solved by cryo-electron microscopy. *Elife, 8*.
doi:10.7554/eLife.48907

Guerrero-Ferreira, R., Taylor, N. M., Mona, D., Ringler, P., Lauer, M. E., Riek, R., . . . Stahlberg, H. (2018). Cryo-EM structure of alpha-synuclein fibrils. *Elife, 7*. doi:10.7554/eLife.36402

Guo, Q., Lehmer, C., Martinez-Sanchez, A., Rudack, T., Beck, F., Hartmann, H., . . . Fernandez-Busnadiego, R. (2018). In Situ Structure of Neuronal C9orf72 Poly-GA Aggregates Reveals Proteasome Recruitment. *Cell, 172*(4), 696-705 e612. doi:10.1016/j.cell.2017.12.030

Hasegawa, M., Fujiwara, H., Nonaka, T., Wakabayashi, K., Takahashi, H., Lee, V. M., . . . Iwatsubo, T. (2002). Phosphorylated alpha-synuclein is ubiquitinated in alpha-synucleinopathy lesions. *J Biol Chem, 277*(50), 49071-49076. doi:10.1074/jbc.M208046200

Kanda, Y. (2013). Investigation of the freely available easy-to-use software 'EZR' for medical statistics. *Bone Marrow Transplant, 48*(3), 452-458. doi:10.1038/bmt.2012.244

Klingstedt, T., Ghetti, B., Holton, J. L., Ling, H., Nilsson, K. P. R., & Goedert, M. (2019). Luminescent conjugated oligothiophenes distinguish between alpha-synuclein assemblies of Parkinson's disease and multiple system atrophy. *Acta Neuropathol Commun, 7*(1), 193. doi:10.1186/s40478-019-0840-1

Laferriere, F., Maniecka, Z., Perez-Berlanga, M., Hruska-Plochan, M., Gilhespy, L., Hock, E. M., . . . Polymenidou, M. (2019). TDP-43 extracted from frontotemporal lobar degeneration subject brains displays distinct aggregate assemblies and neurotoxic effects reflecting disease progression rates. *Nat Neurosci, 22*(1), 65-77. doi:10.1038/s41593-018-0294-y

Lau, A., So, R. W. L., Lau, H. H. C., Sang, J. C., Ruiz-Riquelme, A., Fleck, S. C., . . . Watts, J. C. (2020). alpha-Synuclein strains target distinct brain regions and cell types. *Nat Neurosci, 23*(1), 21-31. doi:10.1038/s41593-019-0541-x

Li, B., Ge, P., Murray, K. A., Sheth, P., Zhang, M., Nair, G., . . . Jiang, L. (2018). Cryo-EM of full-length alpha-synuclein reveals fibril polymorphs with a common structural kernel. *Nat Commun, 9*(1), 3609. doi:10.1038/s41467-018-05971-2

Luk, K. C., Kehm, V., Carroll, J., Zhang, B., O'Brien, P., Trojanowski, J. Q., & Lee, V. M. (2012). Pathological alpha-synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice. *Science, 338*(6109), 949-953. doi:10.1126/science.1227157

Masuda-Suzukake, M., Nonaka, T., Hosokawa, M., Oikawa, T., Arai, T., Akiyama, H., . . . Hasegawa, M. (2013). Prion-like spreading of pathological alpha-synuclein in brain. *Brain, 136*(Pt 4), 1128-1138. doi:10.1093/brain/awt037

Narasimhan, S., Guo, J. L., Changolkar, L., Stieber, A., McBride, J. D., Silva, L. V., . . . Lee, V. M. Y. (2017). Pathological Tau Strains from Human Brains Recapitulate the Diversity of Tauopathies in Nontransgenic Mouse Brain. *J Neurosci, 37*(47), 11406-11423. doi:10.1523/JNEUROSCI.1230-17.2017

Nonaka, T., Masuda-Suzukake, M., Arai, T., Hasegawa, Y., Akatsu, H., Obi, T., . . . Hasegawa, M. (2013). Prion-like properties of pathological TDP-43 aggregates from diseased brains. *Cell Rep, 4*(1), 124-134. doi:10.1016/j.celrep.2013.06.007

Nonaka, T., Watanabe, S. T., Iwatsubo, T., & Hasegawa, M. (2010). Seeded aggregation and toxicity of {alpha}-synuclein and tau: cellular models of neurodegenerative diseases. *J Biol Chem, 285*(45), 34885-34898. doi:10.1074/jbc.M110.148460

Ohhashi, Y., Ito, K., Toyama, B. H., Weissman, J. S., & Tanaka, M. (2010). Differences in prion strain conformations result from non-native interactions in a nucleus. *Nat Chem Biol, 6*(3), 225-230. doi:10.1038/nchembio.306

Peelaerts, W., & Baekelandt, V. (2016). a-Synuclein strains and the variable pathologies of synucleinopathies. *J Neurochem, 139 Suppl 1*, 256-274. doi:10.1111/jnc.13595

Peelaerts, W., Bousset, L., Van der Perren, A., Moskalyuk, A., Pulizzi, R., Giugliano, M., . . . Baekelandt, V. (2015). alpha-Synuclein strains cause distinct synucleinopathies after local and systemic administration. *Nature, 522*(7556), 340-344. doi:10.1038/nature14547

Peng, C., Gathagan, R. J., Covell, D. J., Medellin, C., Stieber, A., Robinson, J. L., . . . Lee, V. M. (2018). Cellular milieu imparts distinct pathological alpha-synuclein strains in alpha-synucleinopathies. *Nature, 557*(7706), 558-563. doi:10.1038/s41586-018-0104-4

Porta, S., Xu, Y., Restrepo, C. R., Kwong, L. K., Zhang, B., Brown, H. J., . . . Lee, V. M. (2018). Patient-derived frontotemporal lobar degeneration brain extracts induce formation and spreading of TDP-43 pathology in vivo. *Nat Commun, 9*(1), 4220. doi:10.1038/s41467-018-06548-9

Prusiner, S. B., Woerman, A. L., Mordes, D. A., Watts, J. C., Rampersaud, R., Berry, D. B., . . . Giles, K. (2015). Evidence for alpha-synuclein prions causing multiple system atrophy in humans with parkinsonism. *Proc Natl Acad Sci U S A, 112*(38), E5308-5317. doi:10.1073/pnas.1514475112

Ross, C. A., & Poirier, M. A. (2004). Protein aggregation and neurodegenerative disease. *Nat Med, 10 Suppl*, S10-17. doi:10.1038/nm1066

Saeki, Y., Isono, E., & Toh-E. A. (2005). Preparation of ubiquitinated substrates by the PY motif-insertion method for monitoring 26S proteasome activity. *Methods Enzymol, 399*, 215-227. doi:10.1016/S0076-6879(05)99014-9

Saito, T., Mihira, N., Matsuba, Y., Sasaguri, H., Hashimoto, S., Narasimhan, S., . . . Saido, T. C. (2019). Humanization of the entire murine Mapt gene provides a murine model of pathological human tau propagation. *J Biol Chem, 294*(34), 12754-12765. doi:10.1074/jbc.RA119.009487

Schweighauser, M., Shi, Y., Tarutani, A., Kametani, F., Murzin, A. G., Ghetti, B., . . . Goedert, M. (2020). Structures of alpha-synuclein filaments from multiple system atrophy. *Nature*. doi:10.1038/s41586-020-2317-6

Shahnawaz, M., Mukherjee, A., Pritzkow, S., Mendez, N., Rabadia, P., Liu, X., . . . Soto, C. (2020). Discriminating alpha-synuclein strains in Parkinson's disease and multiple system atrophy. *Nature*. doi:10.1038/s41586-020-1984-7

Spillantini, M. G., Schmidt, M. L., Lee, V. M., Trojanowski, J. Q., Jakes, R., & Goedert, M.
(1997). Alpha-synuclein in Lewy bodies. *Nature, 388*(6645), 839-840.
doi:10.1038/42166

Suzuki, G., Shimazu, N., & Tanaka, M. (2012). A yeast prion, Mod5, promotes acquired drug
resistance and cell survival under environmental stress. *Science, 336*(6079), 355-359.
doi:10.1126/science.1219491

Tarutani, A., Arai, T., Murayama, S., Hisanaga, S. I., & Hasegawa, M. (2018). Potent
prion-like behaviors of pathogenic alpha-synuclein and evaluation of inactivation
methods. *Acta Neuropathol Commun, 6*(1), 29. doi:10.1186/s40478-018-0532-2

Terada, M., Suzuki, G., Nonaka, T., Kametani, F., Tamaoka, A., & Hasegawa, M. (2018). The
effect of truncation on prion-like properties of alpha-synuclein. *J Biol Chem, 293*(36),
13910-13920. doi:10.1074/jbc.RA118.001862

Thibaudeau, T. A., Anderson, R. T., & Smith, D. M. (2018). A common mechanism of
proteasome impairment by neurodegenerative disease-associated oligomers. *Nat
Commun, 9*(1), 1097. doi:10.1038/s41467-018-03509-0

Tsuji, H., Arai, T., Kametani, F., Nonaka, T., Yamashita, M., Suzukake, M., . . . Tamaoka, A.
(2012). Molecular analysis and biochemical classification of TDP-43 proteinopathy.
Brain, 135(Pt 11), 3380-3391. doi:10.1093/brain/aws230

Wakabayashi, K., Yoshimoto, M., Tsuji, S., & Takahashi, H. (1998). Alpha-synuclein
immunoreactivity in glial cytoplasmic inclusions in multiple system atrophy.
Neurosci Lett, 249(2-3), 180-182. doi:10.1016/s0304-3940(98)00407-8

Watts, J. C., Giles, K., Oehler, A., Middleton, L., Dexter, D. T., Gentleman, S. M., . . .
Prusiner, S. B. (2013). Transmission of multiple system atrophy prions to transgenic
mice. *Proc Natl Acad Sci U S A, 110*(48), 19555-19560. doi:10.1073/pnas.1318268110

Woerman, A. L., Oehler, A., Kazmi, S. A., Lee, J., Halliday, G. M., Middleton, L. T., . . .
Prusiner, S. B. (2019). Multiple system atrophy prions retain strain specificity after
serial propagation in two different Tg(SNCA*A53T) mouse lines. *Acta Neuropathol,
137*(3), 437-454. doi:10.1007/s00401-019-01959-4

Woerman, A. L., Stohr, J., Aoyagi, A., Rampersaud, R., Krejciova, Z., Watts, J. C., . . .
Prusiner, S. B. (2015). Propagation of prions causing synucleinopathies in cultured
cells. *Proc Natl Acad Sci U S A, 112*(35), E4949-4958. doi:10.1073/pnas.1513426112

Wong, Y. C., & Krainc, D. (2017). alpha-synuclein toxicity in neurodegeneration: mechanism
and therapeutic strategies. *Nat Med, 23*(2), 1-13. doi:10.1038/nm.4269

Zondler, L., Kostka, M., Garidel, P., Heinzelmann, U., Hengerer, B., Mayer, B., . . . Danzer, K.
M. (2017). Proteasome impairment by alpha-synuclein. *PLoS One, 12*(9), e0184040.
doi:10.1371/journal.pone.0184040

Figure legends.

Figure 1. Preparation and characterization of two α-synuclein strains.

(A) Schematic representation of two α-synuclein (αS) strains (left) and the resulted two

α-synuclein assemblies (right). (B) Turbidity of these assemblies. Analysis was performed using

student t test. (mean±S.E.M; n=3) Formation of aggregates rather than soluble oligomer is

shown in Figure 1-figure supplement 1. (C) Transfer electron microscopy (TEM) images of

these assemblies. Scale bar, 200nm. (D) Thioflavin T fluorescence of these assemblies and αS

monomer. Analysis was performed using one-way ANOVA and Tukey post hoc test.

(mean±S.E.M; n=3) (E) Congo red bindings of these strains and αS monomer. (mean±S.E.M;

n=3) (F) In vitro seeding activity of these strains. α-synuclein monomers were incubated with

α-synuclein fibril (-) (orange) or α-synuclein fibril (+) (gray). Kinetics of Thioflavin T

fluorescence were shown. Analysis was performed using one-way ANOVA and Tukey post hoc

test. **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Source Data 1. Quantification for graph in Figure 1B, D and E.

Source Data 2. Three independent data in Figure 1F.

Figure 2. Comparison of pathologies in WT mouse brains by inoculation of two α-synuclein

strains.

(A) Distribution of phosphorylated α-synuclein pathology in mouse brain. α-synuclein fibrils (-)

or α-synuclein fibrils (+) were injected into the striatum of WT mouse brain and stained with

phosphorylated α-synuclein antibody 1 month after injection. Regions surrounded by white

rectangles in left panels are magnified and shown in right panels. Typical pathological

α-synuclein deposits are indicated by the arrowheads. Sections were counterstained with

hematoxylin. Representative images of cortex of mice injected with α-synuclein fibrils (-)

 (upper), α-synuclein fibrils (+) (middle) or not injected mice as negative controls (lower) are

 shown. Representative images of corpus callosum and striatum are shown in Figure 2-figure

 supplement 1. Representative images of cortex of other two mice injected with α-synuclein

 fibrils are also shown in Figure 2-figure supplement 2. (B) Distribution of ubiquitin pathology

 in mouse brain. α-synuclein fibrils (-) or α-synuclein fibrils (+) were injected into the striatum

 of WT mouse brain and stained with ubiquitin antibody 1 month after injection. Representative

 images of cortex of mice injected with α-synuclein fibrils (-) (upper), α-synuclein fibrils (+)

 (middle) or not injected mice as negative controls (lower) are shown. Regions surrounded by

 white rectangles in left panels are magnified and shown in right panels. Typical pathological

 ubiquitin positive deposits are indicated by the arrowheads. Sections were counterstained with

 hematoxylin. Scale bars, 50 µm.

 Figure 3. Seeding activities α-synuclein strains in primary mouse cortical neurons.

 The two α-synuclein fibrils were transduced into primary mouse cortical cells. 14 days after

 fibril transduction, the accumulation of abnormal phosphorylated α-synuclein and ubiquitinated

 aggregates are detected by immunofluorescence microscopy and western blotting. (A)

 Phosphorylated α-synuclein (Phos-αS), ubiquitin (Ub) and nuclei (DAPI) were stained. Scale

 bar, 20 µm. Images of cells double stained with antibody against phosphorylated α-synuclein

 and neuronal or astrocyte markers are shown in Figure 3-figure supplement 1. Uptake of these

 α-synuclein fibrils into cells are shown in Figure 3-figure supplement 2. (B) Detection of

 sarkosyl insoluble phosphorylated α-synuclein (left), endogenous mouse α-synuclein (center)

 and ubiquitinated proteins (right) by western blotting. (C) Detection of sarkosyl soluble

 GAPDH as a loading control. (D) The quantification data of sarkosyl insoluble phosphorylated

α-synuclein (left) and ubiquitinated proteins (right) shown in (B) (mean±S.E.M; n=3). Analysis

was performed using one-way ANOVA and Tukey post hoc test. ****$P<0.0001$.

Source Data 1. Quantification for graph in Figure 3D and Figure 3-figure supplement 2C.

Figure 4. Different interaction of α-synuclein strains with 26S proteasome

(A-C) Effects of the α-synuclein strains on the 26S proteasome activity. 26S proteasome was

purified from budding yeast as shown in Figure 4-figure supplement 1. 26S proteasome

activities in the presence of the two α-synuclein fibrils were measured. Chymotrypsin like

activity was measured by LLVY-MCA hydrolysis(A). Caspase like activity was measured by

LLE-MCA hydrolysis (B). Trypsin like activity was measured by LRR-MCA hydrolysis

(mean±S.E.M; n=3).　(D and E) Co-precipitation of 26S proteasome with the α-synuclein

strains. 26S proteasome was mixed with the two α-synuclein fibrils and centrifuged. Resulted

supernatants (sup) and precipitates (ppt) were analyzed by western blotting against α-synuclein

(D, upper) and Rpn11-HA (E, upper). Quantification of the α-synuclein (D, lower) and

Rpn11-HA (E, lower) in ppt fractions were shown. (mean±S.E.M; n=3) (F) The core regions of

the α-synuclein strains. These α-synuclein fibrils were mildly digested by proteinase K and

centrifuged. The resulted pellet fractions were denatured by guanidine and analyzed by

MALDI-TOF-MS. Peptide peaks identified by mass analysis and the predicted peptide regions

corresponding to each peak are shown. (G) Dot blot analysis of the α-synuclein strains. These

α-synuclein fibrils were spotted on nitrocellulose membranes and detected by various antibodies

against α-synuclein (upper).　Reactivity against the antibody raised against 115-122 region of

α-synuclein was quantified (mean±S.E.M; n=3) (lower). Analysis was performed using one-way

ANOVA and Tukey post hoc test. *$P<0.05$, ***$P<0.001$, ****$P<0.0001$ (A, B, C and E).

Analysis was performed using student t test. *$P<0.05$ (D and G).

Source Data 1. Quantification for graph in Figure 4A, B and C.

Source Data 2. Quantification for graph in Figure 4D, E and G.

Figure 5. Effects of C-terminal truncated α-synuclein fibrils on proteasome in neurons and in

vitro.

(A-C) Seeding activities of C-terminal truncated α-synuclein fibrils in primary mouse cortical

neurons. C-terminal truncated α-synuclein fibrils (αSΔC20 (KCl-)) were transduced into

primary mouse cortical cells. 14 days after fibril transduction, the accumulation of abnormal

phosphorylated α-synuclein and ubiquitinated proteins are detected by immunofluorescence

microscopy and western blotting. (A) Phosphorylated α-synuclein (Phos-αS), ubiquitin (Ub) and

nuclei (DAPI) were stained. Scale bar, 20 µm. (B) Detection of sarkosyl insoluble

phosphorylated α-synuclein (left) and endogenous mouse α-synuclein (right) by western blotting.

(C) Detection of sarkosyl soluble GAPDH as a loading control. (D) Effects of the αSΔC20

(KCl-) fibrils on the 26S proteasome activity. 26S proteasome activities in the presence of the

αSΔC20 (KCl-) fibrils were measured. Chymotrypsin like activity (left), caspase like activity

(center) and trypsin like activity (right) are shown. (mean±S.E.M; n=3). (E and F)

Co-precipitation of 26S proteasome with the αSΔC20 (KCl-) fibrils. 26S proteasomes were

mixed with the αSΔC20 (KCl-) fibrils and centrifuged. Resulted supernatants (sup) and

precipitates (ppt) were analyzed by western blotting against α-synuclein (E) and Rpn11-HA (F,

left). Quantification of the Rpn11-HA in ppt fractions were shown (F, right). (mean±S.E.M;

n=3) Analysis was performed using student t test.

Source Data 1. Quantification for graph in Figure 5D and F.

Figure 6

Schematic representation of α-synuclein strain formation and strain dependent interaction with

26S proteasome.

In the absence of salt, α-synuclein monomers have exposed C-terminal region with high electric

repulsion. These form the α-synuclein strain with exposed C-terminal region and this type of the

α-synuclein strain can interact with 26S proteasomes and inhibit their activities resulting the

accumulation of phosphorylated α-synuclein aggregates and ubiquitinated proteins (left). In the

presence of salt, α-synuclein monomers have packed C-terminal region with low electric

repulsion. These form the α-synuclein strain with masked C-terminal region and this type of the

α-synuclein strain can not interact with 26S proteasomes.

Figure Supplement

Figure 1-figure supplement 1

Formation of α-synuclein aggregates.

α-synuclein monomers are incubated for 7days and then centrifuged at 10,000 x g for 10 min to

separate soluble oligomers and insoluble aggregates. CBB staining of soluble fractions (sup)

and insoluble fractions (ppt) are shown. Soluble oligomers are hardly detected in both

α-synuclein strains.

Figure 2-figure supplement 1

Comparison of pathologies in WT mouse brains by inoculation of two α-synuclein strains.

Distribution of phosphorylated α-synuclein pathology in mouse brain. α-synuclein fibrils (-) or

α-synuclein fibrils (+) were injected into the striatum of WT mouse brain and stained with

phosphorylated α-synuclein antibodies 1 month after injection. Representative images of corpus

callosum (A) and striatum (B) are shown. Typical pathological α-synuclein deposits are

indicated by the arrowheads. Sections were counterstained with hematoxylin. Scale bar, 50 µm.

Figure 2-figure supplement 2

Images of cortex of other two mice injected with the α-synuclein fibrils.

Distribution of phosphorylated α-synuclein pathology in mouse brain. α-synuclein fibrils (-) or

α-synuclein fibrils (+) were injected into three mice each and stained with phosphorylated

α-synuclein antibodies 1 month after injection. Representative images of cortex of other two

mice (#2 and #3), not shown in Figure 2, are shown. Typical pathological α-synuclein deposits

are indicated by the arrowheads. Sections were counterstained with hematoxylin. Scale bar, 50

µm.

Figure 3-figure supplement 1

Double stained Images of phosphorylated α-synuclein and neuronal or astrocyte markers.

α-synuclein fibrils (-), which induced the accumulation of phosphorylated α-synuclein, were

transduced into primary mouse cortical cells. 14 days after fibril transduction, phosphorylated

α-synuclein and neuronal or astrocyte markers were doubly stained. (A) Phosphorylated

α-synuclein (Phos-αS), GFAP (an astrocyte marker) and nuclei (DAPI) were stained. (B)

Phosphorylated α-synuclein (Phos-αS), NeuN (a mature neuronal cell marker) and nuclei

(DAPI) were stained. (C) Phosphorylated α-synuclein (Phos-αS), tau (a mature neuronal cell

marker) and nuclei (DAPI) were stained. (D) Phosphorylated α-synuclein (Phos-αS),

Neurofilament-L (a mature neuronal cell marker) and nuclei (DAPI) were stained. Scale bars,

50 µm.

Figure 3-figure supplement 2

Uptake of the α-synuclein fibrils into the cells.

The two α-synuclein fibrils were transduced into primary mouse cortical cells. A day after fibril

transduction, cells were washed and treated with trypsin to digest extracellular α-synuclein

fibrils. Cells were collected and lysed in the presence of sarkosyl. The cell lysates were

centrifuged at 135,000 × g for 20 min, then the supernatant (sup) was collected as

sarkosyl-soluble fraction, and the protein concentration was determined by Bradford assay. The

precipitate (ppt) was solubilized in SDS-sample buffer and used sarkosyl-insoluble fractions.

(A) α-synuclein detected in sarkosyl- insoluble fractions were regarded as the α-synuclein fibrils

taken up by the cells. (B) GAPDH detected in sarkosyl- soluble fractions as loading controls.

(C) The quantification data of sarkosyl insoluble α-synuclein shown in (A) (mean±S.E.M; n=3).

Analysis was performed using student t test. **P<0.01

Figure 4-figure supplement 1

Purification of 26S proteasome.

CBB staining of 26S proteasome complex used in this study.

Suzuki et al., Figure 1.

Suzuki et al., Figure 2.

Suzuki et al., Figure 3.

Suzuki et al., Figure 4.

Suzuki et al., Figure 5

α-synuclein

Suzuki et al., Figure 6