Unclusterable, underdispersed arrangement of plant species in pollinator niche space

Carlos M. Herrera [1]

Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Avenida

Americo Vespucio 26, E-41092 Sevilla, Spain

Running head: Plant species in pollinator niche space

[1] E-mail: herrera@ebd.csic.es

Abstract. Pollinators can mediate facilitative or competitive relationships between plant

species, but the comparative importance of these two conflicting phenomena in shaping

pollinator resource use in plant communities remains unexplored. This paper provides a case

example and proof of concept of the idea that the arrangement in pollinator niche space of large

species samples comprising complete or nearly complete plant communities can be instrumental

to evaluate the importance of facilitation and competition as drivers of pollinator resource at the

plant community level. Pollinator composition data for 221 plant species from the Sierra de

Cazorla mountains (southeastern Spain), comprising 85% of families and ~95% of widely

distributed species of entomophilous plants, were used to address the following questions: (1)

Do objectively identifiable species clusters occur in pollinator niche space ? Three different

pollinator niche spaces were considered whose axes were defined by insect orders ($N = 7$),

families ($N = 94$) and genera ($N = 346$); and (2) If all plant species form a single, indivisible

cluster in pollinator niche space, Are species overdispersed or underdispersed relative to a

random arrangement ? "Clusterability" tests failed to reject the null hypothesis that there was

only one pollinator-defined species cluster in pollinator niche space, irrespective of the space

axes (insect orders, families or genera) or pollinator importance measurement (proportions of

pollinator individuals or flowers visited by each pollinator type) considered. Randomly

simulated species arrangements in each pollinator niche space showed that observed means of

pairwise interspecific distances in pollinator composition were smaller than values from

simulations, thus revealing significantly non-random, underdispersed arrangement of plant

species within the single cluster existing in pollinator niche space. Arrangement of plant species

in niche space in the montane plant community studied did not support a major role for

interspecific competition as a force shaping pollinator resource use by plants, but rather a

situation closer to the facilitation-dominated extreme in a hypothetical competition-facilitation

gradient. Results also illustrate the potential of investigations on complete or nearly complete

entomophilous plant communities for addressing novel hypotheses and contributing insights to

the understanding of the ecology and evolution of plant-pollinator systems.

Key words:clusterability; Mediterranean mountain habitats; plant community; pollinator

composition; pollinator niche space; pollinator syndromes.

42   INTRODUCTION

The maturation of a full-fledged niche theory was a significant landmark in the trajectory of

twentieth-century ecology (Vandermeer 1972, Chase and Leibold 2003, Ricklefs 2012).

Originally aimed at investigating how many and how similar coexisting species could be in a

community, niche theory was predominantly concerned with animal communities and rested on

the overarching assumption that interspecific competition for limiting resources is a major

driving force in shaping community composition and resource use by individual species (Chase

and Leibold 2003). Notions derived from niche theory sometimes also made their way into plant

community studies. In particular, the niche theory framework is behind investigations of plant-

pollinator interactions that have envisaged animal pollinators as "resources" exploited by

entomophilous plants to ensure sexual reproduction. In this context, pollinator partitioning over

daily or seasonal gradients (Heinrich 1975, Herrera 1990, Stone et al. 1998, de Souza et al.

2017), generalization *vs.* specialization on pollinators as alternative strategies of resource use

(Herrera 1996, 2005, Waser et al. 1996, Gómez and Zamora 1999, Armbruster 2017), or

divergence in pollinator composition in response to interspecific competition (Moldenke 1975,

Kephart 1983, Johnson 2010, Bouman et al. 2016, Johnson and Bronstein 2019), are all notions

rooted in niche theory and share its canonical assumption of competition as major driving force.

One insufficiently acknowledged limitation of the application of competition-laden niche

theory to plant-pollinator systems is that, under certain circumstances, pollinator sharing by

different plant species can result in facilitation rather than competition (Thomson 1978, Rathcke

1983, Callaway 1995). This possibility has been verified experimentally many times (see Braun

and Lortie 2019 for review). Proximate mechanisms leading to pollinator-mediated interspecific

facilitation include the "magnet species effect", whereby species highly attractive to pollinators

can benefit co-flowering ones in the neighborhood that are visited less often (Feinsinger et al.

1986, Laverty 1992, Johnson et al. 2003, Hansen et al. 2007, Molina-Montenegro et al. 2008,

Pellegrino et al. 2016); sequential facilitation, whereby earlier-flowering species benefit later-

flowering ones at the same location by sustaining populations of site-faithful, shared pollinators

(Waser and Real 1979, Ogilvie and Thomson 2016); floral mimicry, in which species can derive

reproductive success when the colour, shape or scent of their flowers resemble those of a co-

blooming plant (Gumbert and Kunze 2001, Johnson et al. 2003, Pellegrino et al. 2008); and

diffuse beneficial effects of floral diversity at the plant community level on the pollination

success of individual species (Ghazoul 2006, Seifan et al. 2014, Norfolk et al. 2016).

While the reality of both facilitative and competitive phenomena in plant-pollinator systems

is well established, their comparative importance as drivers of pollinator resource use in natural

communities remain essentially unexplored to date (but see, e.g., Tur et al. 2016). The

deleterious effects of pollinator sharing and interspecific pollen transfer have been traditionally

interpreted as supporting the role of competition in organizing multispecies plant-pollinator

interactions by favoring pollinator partitioning (Stebbins 1970, Waser 1978, Wilson and

Thomson 1996, Morales and Traveset 2008, Mitchell et al. 2009). The hitherto little-explored

possibility remains, however, that facilitative interactions among animal-pollinated plants could

also be significant drivers of plant-pollinator interactions at the community level, since benefits

of pollinator sharing can sometimes outweigh its costs (Tur et al. 2016) and the net outcome of

opposing facilitation and competition might depend on environmental context (Moragues and

Traveset 2005, Ghazoul 2006, Ye et al. 2014, Fitch 2017). I propose here that the comparative

importance of competition *vs.* facilitation in shaping community-wide plant-pollinator systems

could be evaluated by examining the arrangement of plant species in the pollinator niche space,

as outlined in the next section.

89   *A hypothesis on species arrangement in pollinator niche space*

Pollinator-mediated interspecific facilitation and competition can be envisaged as exerting

opposite forces on pollinator resource use, tending to favor pollinator sharing and divergence,

respectively. Insights into the relative importance of these two conflicting forces could thus be

gained by examining how *the whole set* of animal-pollinated species in a plant community are

arranged in the multidimensional niche space whose axes are biologically distinct pollinator

"types". Such types should include all objectively recognizable pollinators types that are

sufficiently different in size, morphology, behavior and/or pollinating features as to represent

different resources from the plants' perspective. A simple sketch of this idea referred to just two

pollinator niche axes is shown in Fig. 1. Divergence in pollinator resource use, denoted by two

or more distinct species clusters in pollinator space (Fig. 1A), would be consistent with

competition playing a prevailing role in the organization of plant-pollinator interactions. On the

opposite extreme, a single cluster of densely packed (underdispersed relative to a random

distribution) species in the pollinator niche space (Fig. 1C) would be suggestive of predominant

facilitation having promoted convergence in pollinator resource use by the different plant

species. The intermediate situation, in which species form a single cluster of uniformly

distributed species (overdispersed relative to a random distribution) in pollinator niche space

(Fig. 1B) would point to a sort of balance between facilitation (favoring a single cluster) and

competition (driving species repulsion in niche space via limiting similarity effects).

The main objective of this paper is to provide a case example and proof of concept for the

preceding framework by examining the arrangement in the pollinator niche space of a very large

sample of entomophilous plant species from a hyperdiverse community. A secondary objective

is to highlight the importance of adopting "zooming out" approaches in the study of the ecology

of plant-pollinator interactions (Herrera 2020). Current knowledge of pollination system niches

has been mostly obtained from studies dealing with small species samples (usually < 25

species), or focusing on specific pollination guilds associated with particular pollinators. While

irreplaceable for other purposes, these investigations lack the biological amplitude needed to

address broad-scale questions and warrant generalizations on the overall organization of plant-

pollinator systems (Johnson 2010, Herrera 2019, 2020). Only a comprehensive coverage of the

ecological and phylogenetic range of the insect-pollinated plants in a region and their whole set

of pollinators could help to discriminate among the scenarios hypothesized in Fig. 1. The

species sample examined here virtually encompasses the whole community of entomophilous

plants in a southeastern Spanish biodiversity hotspot (see *Study area and plant species sample*

below) and the whole spectrum of their insect pollinators.

The following two specific questions will be addressed: (1) Are there distinct, objectively

identifiable plant species clusters in the pollinator niche space ? Three different pollinator niche

spaces with broadly different dimensionalities will be considered, with axes defined by insect

orders, families and genera; and (2) If plant species form a single indivisible cluster in the

pollinator niche space, Is there evidence for either overdispersed or underdispersed arrangement

of species within the cluster ? Two simplifications in the scheme of Fig. 1 must be

acknowledged. First, the sign and magnitude of the hypothesized competition-facilitation

balance mediated by pollinators may depend on the spatial scale considered (Tur et al. 2016,

Braun and Lortie 2019). And second, the scheme does not incorporate the possibility that even if

divergence in pollinator composition can not be detected, pollinator partitioning associated with

differential blooming time or habitat segregation could still reflect the effect of competition. The

possible relationships between interspecific differences in habitat type and flowering time, and

interspecific divergence in pollinator composition, will also be addressed.

136   MATERIALS AND METHODS

137   *Study area and plant species sample*

Pollinator composition data considered in this paper were collected during February-

December 1997-2019 (with a gap in 2000-2002) in the Sierras de Cazorla-Segura-Las Villas

Natural Park (Jaén Province, southeastern Spain), a region characterized by outstanding

biological diversity and well-preserved mountain habitats. Sampling sites were the same as in

Herrera (2019, 2020), where details on locations can be found. Quantitative data on pollinator

composition were obtained for 221 plant species in 48 families (Appendix S1: Table S1). This

sample comprises 85% of families and ~95% of widely distributed species of entomophilous

plants in the study area (Gómez Mercado 2011, C. M. Herrera, *unpublished data*), and is a

superset of the 191 species in 43 families studied by Herrera (2020: Table 1). Asteraceae (41

species), Lamiaceae (24), Fabaceae (14), Brassicaceae (12), Rosaceae (11) and Ranunculaceae

(9) contributed most species to the sample. About two thirds of species ($N = 156$) were sampled

for pollinators only one year, and the rest were sampled ≥ 2 years. In the vast majority of species

($N = 211$) pollinator sampling was conducted on a single site. Data for the same species

obtained in more than one site or year were combined into a single sample for the analyses.

Pollinators of some of the species considered here vary among sites or years, but such

intraspecific variation is always small in comparison to the broad interspecific range occurring

in the large species sample considered here (e.g., Herrera 1988, 2019, C. M. Herrera,

unpublished data), which justifies characterizing each plant species by the combined sample of

pollinator data from different sites and years.

Pollinator sampling dates for each species roughly matched its peak flowering date. The

mean date of pollinator censuses for each species, expressed as days from 1 January, was used

in analyses relating differences in pollinator composition to variation in flowering phenology.

Mean census dates for the species studied ranged between 39–364 days from 1 January, and

most species flowered during May-July (interquartile range = 145-190 days from 1 January)

(Appendix S1: Table S1). For analyses relating differences in pollinator composition to

dissimilarity in habitat type, each plant species was classed into one of the following nine major

habitat categories (total species per habitat in parentheses; see also Herrera 2020): banks of

permanent streams or flooded/damp areas around springs (18); dwarf mountain scrub dominated

by cushion plants (27); forest edges and large clearings (34); forest interior (20); patches of

grasslands and meadows on deep soils in relatively flat terrain (40); local disturbances caused by

humans, large mammals or natural abiotic processes (24); sandy or rocky dolomitic outcrops

(29); tall, dense Mediterranean sclerophyllous forest and scrub (17); vertical rock cliffs (12

species). Individual species' assignments to habitat types are shown in Appendix S1: Table S1.

171   *Pollinator composition*

Data on pollinator composition were obtained for every species as described in Herrera

(2019, 2020). The elemental sampling unit was the "pollinator census", consisting of a 3-min

watch of a flowering patch whose total number of open flowers was also counted. All

pollinators visiting some flower in the focal patch during the 3-min period were identified, and

total number of flowers probed by each individual was recorded. Detailed information on

sampling effort is shown in Appendix S1: Table S1. For all plant species combined, a total of

28,128 censuses were conducted on 575 different dates. Assessing the number of flowers visited

by pollinators was impractical in the case of species with tiny flowers densely packed into

compact inflorescences (e.g., Apiaceae, Asteraceae). In these species the number of

inflorescences available per patch and visited per census were counted rather than individual

flowers, although for simplicity I will always refer to "flower visitation". Close-up photographs

of pollinators visiting flowers were taken routinely during censuses, which were used for

identification, keeping photographic vouchers, and ascertaining the pollinating status of

different insect taxa. Only taxa whose individuals contacted anthers or stigmas, or with

discernible pollen grains on their bodies, were considered as pollinators.

187   *Data analysis*

Two measurements of pollinator importance and three different pollinator niche spaces will

be considered throughout the analyses. Pollinator composition for each plant species will be

assessed using the proportion of pollinator individuals recorded and the proportion of flowers

visited for each pollinator type, which will be termed "proportion of visitors" and "proportion of

visits", respectively. Three distinct, taxonomically-defined classifications of pollinators will be

considered in the analyses, based respectively on their affiliation to insect orders, families and

genera. These classifications defined three pollinator niche spaces that differed in their

dimensionality and the biological meaning of their axes ($n = 7$, 94 and 346 dimensions for the

spaces defined by orders, families and genera, respectively). Out of a total of 33,835 individual

pollinators recorded in censuses, 100% were identified to order and family, and 98.3% to genus,

but only 80.3% to species. Analyses on a possible fourth pollinator niche space defined by

species were not undertaken because excluding data with missing species-level identifications

would have led to severe taxonomic biases and distortion of species arrangements in pollinator

niche space. For instance, about half (46%) of all individual pollinators that remained

unidentified at the species level belonged to just four bee genera (*Andrena*, *Ceratina*,

Lasioglossum, Panurgus).

The question of whether species clusters could be recognized in each of the three pollinator

niche spaces considered (defined by insect orders, families and genera) was first addressed

informally by comparing the distribution of species in a reduced bidimensional space with the

various scenarios envisaged in Fig. 1. For each pollinator niche space, nonmetric

multidimensional scaling (NMDS) was performed on the matrix of interspecific pairwise

distances in relative importance of pollinator types (proportion of visitors and proportion of

visits). Computations were performed with the function metaMDS in the package vegan

(Oksanen et al. 2019) for the R computing environment (R Core Team 2018; all statistical

analyses in this paper were carried out using R). In metaMDS the ordination is followed by a

rotation via principal components analysis, which consistently ensures that the first axis reflects

the principal source of variation in the data and thus enhances the possibility of revealing

clusters. Unless otherwise stated the Manhattan distance metric (also known as L1 norm, city

block or taxicab metric; Deza and Deza 2013) was used to compute interspecific distances

matrices for NMDS and rest of analyses involving distance matrices. The Manhattan metric is

better suited than the widely used Euclidean metric for analyses of high dimensionality data as

those done here (Hinneburg et al. 2000, Aggarwal et al. 2001, Houle et al. 2010).

Formal statistical analyses of "clusterability", or whether a meaningful clustered structure

can be objectively established in a multidimensional data set (Ackerman and Ben-David 2009,

Nowakowska et al. 2015, Simovici and Hua 2019), were conducted on the set of species

coordinates in each of the three pollinator niche spaces considered (each defined by either

proportion of visitors or proportion of visits). Dip multimodality tests (Hartigan and Hartigan

1985) were applied to distributions of pairwise distances between species, a method particularly

robust to outliers, anomalous topologies and high dimensionality (Adolfsson et al. 2019). These

tests evaluated the null hypothesis that the distribution of interspecific distances in pollinator

space was unimodal, denoting a single species cluster, against the alternative hypothesis that the

distribution had two or more modes, denoting the existence of at least two clusters (Adolfsson et

al. 2019; see Brownstein et al. 2019, Simovici and Hua 2019, for examples of applications). The

clusterabilitytest function in the clusterability package (Neville and Brownstein 2019) was used

for computations, and statistical significance was determined by Monte Carlo simulations with

10^5 repetitions. Since results of clusterability tests did not support the Fig. 1A scenario, it was

important to assess whether the failure to identify species clusters in the pollinator niche spaces

considered was an spurious outcome of pooling species from different habitats and with

different flowering phenologies, which could be partitioning pollinators in time or space. This

possibility was examined by simultaneously regressing each matrix of interspecific distance in

pollinator composition against corresponding pairwise matrices of distances in blooming time

(absolute difference between mean pollinator census dates) and habitat type dissimilarity

(simple mismatching coefficient), using the multi.mantel function in the phytools package

(Revell 2012). Statistical significance was evaluated using permutations with 10^5 repetitions.

Simulations were performed to elucidate whether the arrangement of species in pollinator

niche space matched any of the scenarios depicted in Figs. 1B and 1C. The observed mean

pairwise distance between species in each pollinator space was compared with a distribution of

means obtained from simulated pollinator composition data. Each simulation consisted of

permuting the values in each column of the raw plant species x pollinator type data matrix

(number of pollinator individuals or number of flower visits contributed by each pollinator

type); computing for each plant species the proportional importance of each pollinator type in

the simulated data; obtaining the pairwise distances matrix; and finally computing the mean

pairwise distance between species. This simulation procedure created random configurations of

proportional pollinator compositions for individual plant species, and thus random arrangements

of plants in the corresponding pollinator niche space, but preserved exactly both the observed

distributional properties of visitors or visits for every pollinator type, and the absolute and

relative numerical importances of the different pollinator types for all plant species combined. A

total of 10^5 simulated species arrangements were computed in each of the three pollinator niche

spaces, using proportion of visitors and proportion of visits as measurements of pollinator type

importance. To test whether the observed arrangement of species in the pollinator niche space

departed significantly from the randomly simulated ones, the mean interspecific distance

observed was compared with the 0.025 and 0.975 quantiles of the distribution of simulated

means.

261   RESULTS

Most plant species in the sample studied ($N = 221$) had very taxonomically diverse arrays of

insect pollinators. On average ($\pm$ SE), individual plant species were pollinated by 3.3 ± 0.1

different insect orders (interquartile range, IQR = 3–4), 10.7 ± 0.4 insect families (IQR = 6–14)

and 16.7 ± 0.8 insect genera (IQR = 7–22). The number of orders, families and genera of

pollinators recorded for each plant species are summarized in Appendix S2: Table S1, and the

complete list of taxonomic categories recorded is presented in Appendix S2: Table S2.

Neither the informal analysis based on nonmetric multidimensional scaling (NMDS) nor the

formal one testing clusterability by means of dip tests supported the existence of objectively

definable plant species clusters in any of the three pollinator niche spaces considered.

Irrespective of whether pollinator composition was assessed using proportion of visitors or

proportion of visits, the cloud of points in the reduced two-dimensional space obtained with

NMDS was roughly centered on the bivariate origin (0, 0), and there was not any indication of

clusters or discontinuities suggestive of species groups with contrasting, well-defined pollinator

types (Fig. 2). Clusterability analyses using dip tests likewise failed to detect clustered structures

underlying the data in any of the three pollinator niche spaces considered (Table 1).

Multiple Mantel regressions revealed no statistically significant effects of interspecific

dissimilarity in flowering time or habitat type on interspecific distance in pollinator composition

when the axes of the pollinator niche space were insect orders (Table 2). When niche spaces

were defined by families and genera, in contrast, statistically significant relationships did exist

between distance in pollinator composition and dissimilarities in flowering time and habitat

type. In all these instances the relationships were positive, denoting that species pairs sharing the

same habitat type or flowering at similar times of year tended to be significantly more similar in

pollinator composition at the family and genus levels of taxonomic resolution than species from

different habitats or flowering at different times of year.

Simulations showed that plant species were significantly underdispersed in the pollinator

niche space. Regardless of the niche space considered or the measurement of pollinator

importance used, the observed mean of pairwise interspecific distances in pollinator

composition was always significantly smaller than simulated values, falling well below the

lower limit of the range defined by the 0.025–0975 quantiles (Fig. 3). The extent of

underdispersion, as intuitively assessed by the disparity between the observed mean and the

range of simulated ones, increased from order- through family- to genus-defined pollinator

niche spaces.

294   DISCUSSION

Considerable effort has been invested to develop reliable methods for grouping sets of

objects in multivariate space in such a way that objects falling in the same group are more

similar among themselves than to objects in other groups, a technique known as "cluster

analysis" (Hennig et al. 2016) widely used in ecological applications. Clustering techniques

often focus on the problem of optimally splitting a multivariate data set into $k \geq 2$ clusters

(Halkidi et al. 2016), yet the problem of elucidating whether cohesive and well-differentiated

groups of objects actually exist in the data, i.e., testing the null hypothesis that $k = 1$ against the

alternative $k \geq 2$, has been rarely investigated due to computational difficulties (the so-called

"clusterability problem"; Ackerman and Ben-David 2009, Nowakowska et al. 2015, Simovici

and Hua 2019). Following recent technical developments, it has been advised that a prior study

of clusterability should become an integral part of any cluster analysis (Adolfsson 2016,

Adolfsson et al. 2019, Simovici and Hua 2019). Clusterability tests could be profitably

incorporated as a first step in analyses of ecological questions traditionally addressed by means

of cluster analysis, so that the existence in multivariate spaces of objectively recognizable

clusters of objects (e.g., species, plots, communities) can be rigorously evaluated before

proceeding to clustering. This applies also to pollination studies, which have sometimes used

clustering to identify plant groupings in multivariate spaces defined by floral traits or pollinator

composition (Ollerton et al. 2009, de Jager et al. 2011, Abrahamczyk et al. 2017, Schurr et al.

2019, Vandelook et al. 2019). As exemplified by this study, clusterability analysis allows tests

of hypotheses in pollination ecology by providing an objective means of ascertaining whether

groups of plants defined by pollinators actually exist in a given species sample.

The present study failed to identify distinct clusters of plant species in the pollinator niche

space. Groups with species more similar to each other in pollinator composition than to species

in other groups were not statistically recognizable in the very large species sample of

entomophilous plants considered. This conclusion emerges as a particularly robust one, since it

was consistently attained regardless of the particular taxonomic resolution adopted to define the

axes of the pollinator niche space (insect orders, families or genera) or the measurement of

pollinator importance (proportion of visitors or proportion of visits). Variation in pollinator

composition across the large species sample was essentially continuous. This was largely a

consequence of the generalized pollination systems exhibited by the vast majority of plant

species considered, each of which had pollinators belonging to a substantial number of different

insect orders and families. Similar predominance of generalized pollination systems has been

reported by other community studies (McCall and Primack 1992, Waser et al. 1996, Bosch et al.

1997), but I am not aware of any previous investigation evaluating the reality of species clusters

in multidimensional pollinator spaces using objective statistical methods. Studies of plant-

pollinator networks have often looked for, and sometimes found, "modules" consisting of

subsets of plant and animal species that are closely linked internally but weakly linked to other

modules (Olesen et al. 2007, Dormann and Strauss 2014), which would suggest some parallels

with the concept of species clusters in pollinator space considered in this study. The concepts

are not strictly comparable, however, because plant-pollinator network data usually contain a

blend of information on plant species abundances, pollinator visitation rates and pollinator

composition (C. M. Herrera, *unpublished information*). As a matter of fact, "modular" plant-

pollinator networks can fail to pass the clusterability test after information on plant species

abundance and pollinator visitation rates is purged from the data by computing for each plant

species the proportions of different pollinator types. For instance, the set of $N = 25$ plant species

in Memmott's (1999) plant-pollinator network, which was presented as an example of modular

structure by Dormann and Strauss (2014; see also Olesen et al. 2007), is unclusterable in the

pollinator niche space defined by the 79 insect taxa involved using the methods applied in the

present study (C. M. Herrera, *unpublished information*). This contrast between plant-pollinator

network modularity and plant clusterability in pollinator niche space seems another factor

contributing to the disconnection between community ecology and network approaches pointed

out by Blüthgen (2010).

Pollinators from different insect orders, families or genera generally have contrasting

morphological and behavioral attributes, and these differences can eventually lead to variation

in important pollinating features such as, e.g., ability to access floral rewards, daily or seasonal

activity rhythm, flower visitation rate, frequency of pollen removal or deposition, or flight

distance between flowers. On this basis, taxonomically-defined groupings of insects have been

often treated as distinct "functional pollinator groups" and related to distinct plant "pollination

syndromes", or suites of floral features that are associated to particular pollinator groups (Fægri

and van der Pijl 1979, Fenster et al. 2004, Ollerton et al. 2007, 2009, Gómez et al. 2008). The

relationship between insect taxonomic affiliation and pollination attributes lends biological

justification to using taxonomic categories to define the axes of multidimensional pollinator

niche spaces. This study has shown that the arrangement of plant species in pollinator niche

space did not match any of the scenarios postulated on the assumption that interspecific

competition is an important force shaping pollinator resource use by plants. Not only there were

no discernible species clusters that could reflect a prevalence of pollinator niche partitioning, but

also (1) the arrangement of species was significantly more dense than expected by chance given

the observed abundances of the different pollinator types considered; and (2) Mantel multiple

regressions revealed that species sharing the same habitats or blooming at the same time of year

were more similar to each other in pollinator composition than species that segregated

temporally or spatially.

Results of this study suggest that concepts quintessential to competition-laden niche theory,

such as limiting similarity, character displacement and competitive exclusion (Chase and

Leibold 2003, Johnson and Bronstein 2019), do not generally apply to entomophilous plants in

my study region in relation to their exploitation of insect pollinator resources. In addition, the

absence in my plant species sample of objectively recognizable clusters linked to particular

pollinator groups does not support the generality of plant pollination syndromes associated to

specialization on particular insect functional groups (see also Ollerton et al. 2009, 2015).

Observed patterns, consisting of plants not showing objectively-definable partitioning of

pollinators and being underdispersed in pollinator niche space, would fall close to the

facilitation-dominated extreme in a hypothetical competition-facilitation gradient. The

competition-facilitation balance in plant-pollinator systems can depend on particularities of the

ecological scenario under consideration (Moragues and Traveset 2005, Ghazoul 2006, Ye et al.

2014, Fitch 2017), and results of the present study can be tentatively related to features of plant-

pollinator interactions in the study region. On one side, insect pollinators probably are not a

limiting resource for the majority of entomophilous plants in the region, as shown by the high

flower visitation rates in most habitat types (Herrera 2020), generally weak or inexistent pollen

limitation even in plants blooming at the most adverse periods of year (Herrera 1995, 2002,

Alonso 2005, Alonso et al. 2012, 2013), and increased pollinator abundance in recent times in

parallel to the long-term warming trend exhibited by the regional climate (Herrera 2019). And

on the other, sharing of individual pollinators by different plant species seems a widespread

phenomenon, as denoted by the frequent occurrence of heterospecific pollen grains in the

stigmas of most species (Arceo-Gómez et al. 2018), common field observations of insect

individuals interspersing visits to flowers of different species in the same foraging bout, and

close-up photographs often revealing conspicuous mixed pollen loads on pollinator bodies (C.

M. Herrera, *unpublished observations*). Taken together, all these observation are in accordance

with the interpretation that the unclusterable, underdispersed arrangement of plant species in

pollinator niche space found in this study reflects a prevailing role of interspecific facilitation as

the organizing driver of plant-pollinator systems in the montane habitats studied.

394   CONCLUDING REMARKS

Excepting a few classical monographs considering complete or nearly complete regional

plant–pollinator assemblages (Müller 1883, Robertson 1928, Moldenke 1976), the vast majority

of pollination community studies have examined only small numbers of plant species at a time

(e.g., Ollerton 2017:Appendix 2, and datasets included in the bipartite package for R, Dormann

et al. 2008). Small species sets drawn from much larger pools of entomophilous species provide

suitable systems for investigating whether a particular phenomenon occurs or not (e.g., testing

for pollinator-mediated competition or facilitation in a well-chosen group of species; see

references in *Introduction*), but are intrisically unable to address more ambitious, broadly-

encompassing questions bearing on the strength or frequency with which the phenomenon of

interest operates in a particular ecological context. In addition, unacknowledged biases in the

selection of pollination study systems can lead to small plant species samples rarely representing

truly scaled-down, taxonomically- and ecologically-unbiased versions of whole plant

communities, possibly leading to distorted representations of natural patterns (Ollerton et al.

2015). Place-based ecological research pursuing "general understanding through the sort of

detailed understanding of a particular place" (Price and Billick 2010, p. 4) has been only

infrequently adopted by pollination studies. As exemplified by the present investigation (see

also Herrera 2020), comprehensive studies of complete or nearly complete entomophilous plant

communities possess an unexplored potential for testing novel hypotheses and contributing

realistic insights to the understanding of the ecology and evolution of plant-pollinator systems.

ACKNOWLEDGMENTS

I am deeply indebted to all insect taxonomists that helped over the years with insect

identifications, a complete list of which was presented in Herrera (2020: *Acknowledgments*).

Consejería de Medio Ambiente, Junta de Andalucía, granted permission to work in the Sierra de

Cazorla and provided invaluable facilities there. Mónica Medrano provided cogent criticisms at

an early stage of the analyses which were instrumental in refining my thinking and improving

the analytical approach. The research reported in this paper received no specific grant from any

funding agency.

LITERATURE CITED

Abrahamczyk, S., S. Lozada-Gobilard, M. Ackermann, E. Fischer, V. Krieger, A. Redling, and

M. Weigend. 2017. A question of data quality – Testing pollination syndromes in

Balsaminaceae. PLoS ONE 12:e0186125.

Ackerman, M., and S. Ben-David. 2009. Clusterability: a theoretical study. Proceedings of the

12th International Conference on Artificial Intelligence and Statistics 5:1–8.

Adolfsson, A. 2016. An effective and efficient approach for clusterability evaluation. Master of

Science Thesis, Department of Computer Science, Florida State University, Tallahassee,

Florida, USA.

Adolfsson, A., M. Ackerman, and N. C. Brownstein. 2019. To cluster, or not to cluster: an

analysis of clusterability methods. Pattern Recognition 88:13–26.

Aggarwal, C. C., A. Hinneburg, and D. A. Keim. 2001. On the surprising behavior of distance

metrics in high dimensional spaces. Pages 420–434 in J. VanDenBussche and V. Vianu,

editors. Proceedings 8th International Conference on Database Theory. Springer, Berlin,

Germany.

Alonso, C. 2005. Pollination success across an elevation and sex ratio gradient in gynodioecious

Daphne laureola. American Journal of Botany 92:1264–1269.

Alonso, C., C. M. Herrera, and T.-L. Ashman. 2012. A piece of the puzzle: a method for

comparing pollination quality and quantity across multiple species and reproductive events.

New Phytologist 193:532–542.

Alonso, C., C. M. Navarro-Fernández, G. Arceo-Gómez, G. A. Meindl, V. Parra-Tabla, and T.

L. Ashman. 2013. Among-species differences in pollen quality and quantity limitation:

implications for endemics in biodiverse hotspots. Annals of Botany 112:1461–1469.

Arceo-Gómez, G., C. Alonso, T.-L. Ashman, and V. Parra-Tabla. 2018. Variation in sampling

effort affects the observed richness of plant-plant interactions via heterospecific pollen

transfer: implications for interpretation of pollen transfer networks. American Journal of

Botany 105:1601–1608.

Armbruster, W. S. 2017. The specialization continuum in pollination systems: diversity of

concepts and implications for ecology, evolution and conservation. Functional Ecology

31:88–100.

Price, M. V., and I. Billick. 2010. The ecology of place. Pages 1–10 *in* I. Billick and M. V.

Price, editors. The ecology of place. Contributions of place-based research to ecological

understanding. University of Chicago Press, Chicago, Illinois, USA.

Blüthgen, N. 2010. Why network analysis is often disconnected from community ecology: A

critique and an ecologist's guide. Basic and Applied Ecology 11:185–195.

Bosch, J., J. Retana, and X. Cerdá. 1997. Flowering phenology, floral traits and pollinator

composition in a herbaceous Mediterranean plant community. Oecologia 109:583–591.

Bouman, R. W., S. L. Steenhuisen, and T. van der Niet. 2016. The role of the pollination niche

in community assembly of *Erica* species in a biodiversity hotspot. Journal of Plant Ecology

10:634–648.

Braun, J., and C. J. Lortie. 2019. Finding the bees knees: a conceptual framework and

systematic review of the mechanisms of pollinator-mediated facilitation. Perspectives in

Plant Ecology, Evolution and Systematics 36:33–40.

Brownstein, N. C., A. Adolfsson, and M. Ackerman. 2019. Descriptive statistics and

visualization of data from the R datasets package with implications for clusterability. Data

in Brief 25:104004.

Callaway, R. M. 1995. Positive interactions among plants. Botanical Review 61:306–349.

Chase, J. M., and M. A. Leibold. 2003. Ecological niches. Linking classical and contemporary

approaches. University of Chicago Press, Chicago, Illinois, USA.

de Jager, M. L., L. L. Dreyer, and A. G. Ellis. 2011. Do pollinators influence the assembly of

flower colours within plant communities? Oecologia 166:543–553.

de Souza, J. M. T., C. Snak, and I. G. Varassin. 2017. Floral divergence and temporal pollinator

partitioning in two synchronopatric species of *Vigna* (Leguminosae-Papilionoideae).

Arthropod-Plant Interactions 11:285–297.

Deza, M. M., and E. Deza. 2013. Encyclopedia of distances. 2nd edition. Springer, Berlin,

Germany.

Dormann, C. F., B. Gruber, and J. Fründ. 2008. Introducing the bipartite package: analysing

ecological networks. R news 8:8–11.

Dormann, C. F., and R. Strauss. 2014. A method for detecting modules in quantitative bipartite

networks. Methods in Ecology and Evolution 5:90–98.

Fægri, K., and L. van der Pijl. 1979. The principles of pollination ecology. Third edition.

Pergamon Press, Oxford, United Kingdom.

Feinsinger, P., K. G. Murray, S. Kinsman, and W. H. Busby. 1986. Floral neighborhood and

pollination success in four hummingbird-pollinated cloud forest plant species. Ecology

67:449–464.

Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004.

Pollination syndromes and floral specialization. Annual Review of Ecology, Evolution and

Systematics 35:375–403.

Fitch, G. M. 2017. Urbanization-mediated context dependence in the effect of floral

neighborhood on pollinator visitation. Oecologia 185:713–723.

Ghazoul, J. 2006. Floral diversity and the facilitation of pollination. Journal of Ecology 94:295–

304.

Gómez Mercado, F. 2011. Vegetación y flora de la Sierra de Cazorla. Guineana 17:1–481.

Gómez, J. M., and R. Zamora. 1999. Generalization vs. specialization in the pollination system

of *Hormathophylla spinosa* (Cruciferae). Ecology 80:796–805.

Gómez, J. M., J. Bosch, F. Perfectti, J. D. Fernández, M. Abdelaziz, and J. P. M. Camacho.

2008. Association between floral traits and rewards in *Erysimum mediohispanicum*

(Brassicaceae). Annals of Botany 101:1413–1420.

Gumbert, A., and J. Kunze. 2001. Colour similarity to rewarding model plants affects

pollination in a food deceptive orchid, *Orchis boryi*. Biological Journal of the Linnean

Society 72:419–433.

Halkidi, M., M. Vazirgiannis, and C. Hennig. 2016. Method-independent indices for cluster

validation and estimating the number of clusters. Pages 595–618 in Hennig, C., M. Meila,

F. Murtagh, and R. Rocci, editors. Handbook of cluster analysis. CRC Press, Boca Raton,

Florida, USA.

Hansen, D. M., H. C. Kiesbuy, C. G. Jones, and C. B. Muller. 2007. Positive indirect

interactions between neighboring plant species via a lizard pollinator. American Naturalist

169:534–542.

Hartigan, J. A., and P. M. Hartigan. 1985. The dip test of unimodality. Annals of Statistics

13:70–84.

Heinrich, B. 1975. Bee flowers: a hypothesis on flower variety and blooming times. Evolution
29:325–334.

Hennig, C., M. Meila, F. Murtagh, and R. Rocci, editors. 2016. Handbook of cluster analysis.
CRC Press, Boca Raton, Florida, USA.

Herrera, C. M. 1988. Variation in mutualisms: the spatio-temporal mosaic of a pollinator
assemblage. Biological Journal of the Linnean Society 35:95–125.

Herrera, C. M. 1990. Daily patterns of pollinator activity, differential pollinating effectiveness,
and floral resource availability, in a summer-flowering Mediterranean shrub. Oikos 58:277–
288.

Herrera, C. M. 1995. Floral biology, microclimate, and pollination by ectothermic bees in an
early-blooming herb. Ecology 76:218–228.

Herrera, C. M. 1996. Floral traits and plant adaptation to insect pollinators: a devil's advocate
approach. Pages 65–87 *in* D. G. Lloyd, and S. C. H. Barrett, editors. Floral Biology. Studies
on floral evolution in animal-pollinated plants. Chapman and Hall, New York, New York,
USA.

Herrera, C. M. 2002. Censusing natural microgametophyte populations: variable spatial mosaics
and extreme fine-graininess in winter-flowering *Helleborus foetidus* (Ranunculaceae).
American Journal of Botany 89:1570–1578.

Herrera, C. M. 2005. Plant generalization on pollinators: species property or local phenomenon?
American Journal of Botany 92:13–20.

Herrera, C. M. 2019. Complex long-term dynamics of pollinator abundance in undisturbed
Mediterranean montane habitats over two decades. Ecological Monographs 89:e01338.

Herrera, C. M. 2020. Flower traits, habitat and phylogeny as predictors of pollinator service: a

plant community perspective. Ecological Monographs 90. doi:10.1002/ecm.1402

Hinneburg, A., Aggarwal, C. C., and Keim, D. A. 2000. What is the nearest neighbor in high

dimensional spaces? Pages 506–515 in El Abbadi, A., Brodie, M. L., Chakravarthy, S.,

Dayal, U., Kamel, N., Schlageter, G., and Whang, K. Y., editors. Proceedings of the 26th

International Conference on Very Large Databases, Cairo, Egypt.

Houle, M. E., Kriegel, H. P., Kröger, P., Schubert, E., and Zimek, A. 2010. Can shared-neighbor

distances defeat the curse of dimensionality ? Pages 482–500 in Gertz, M., and Ludäscher,

B., editors. Proceedings 22th International Conference on Scientific and Statistical Database

Management. Springer, Berlin, Germany.

Johnson, C. A., and J. L. Bronstein. 2019. Coexistence and competitive exclusion in mutualism.

Ecology 100:e02708.

Johnson, S. D. 2010. The pollination niche and its role in the diversification and maintenance of

the southern African flora. Philosophical Transactions of the Royal Society B 365:499–516.

Johnson, S. D., R. Alexandersson, and H. P. Linder. 2003. Experimental and phylogenetic

evidence for floral mimicry in a guild of fly-pollinated plants. Biological Journal of the

Linnean Society 80:289–304.

Johnson, S. D., C. I. Peter, L. A. Nilsson, and J. Ågren. 2003. Pollination success in a deceptive

orchid is enhanced by co-occurring rewarding magnet plants. Ecology 84:2919–2927.

Kephart S. R. 1983. The partitioning of pollinators among three species of *Asclepias*. Ecology

64:120–133.

Laverty, T. M. 1992. Plant interactions for pollinator visits: a test of the magnet species effect.

Oecologia 89:502–508.

Legendre, P., and L. Legendre. 2012. Numerical ecology. Elsevier, Amsterdam, The

Netherlands.

McCall, C., and R. B. Primack. 1992. Influence of flower characteristics, weather, time of day,

and season on insect visitation rates in three plant communities. American Journal of

Botany 79:434–442.

Memmott, J. 1999. The structure of a plant-pollinator food web. Ecology Letters 2:276–280.

Mitchell, R. J., R. J. Flanagan, B. J. Brown, N. M. Waser, and J. D. Karron. 2009. New frontiers

in competition for pollination. Annals of Botany 103:1403–1413.

Moldenke, A. R. 1975. Niche specialization and species diversity along a California transect.

Oecologia 21:219–242.

Moldenke, A. R. 1976. California pollination ecology and vegetation types. Phytologia 34:305–

361.

Molina-Montenegro, M. A., E. I. Badano, and L. A. Cavieres. 2008. Positive interactions among

plant species for pollinator service: assessing the 'magnet species' concept with invasive

species. Oikos 117:1833–1839.

Moragues, E., and A. Traveset. 2005. Effect of *Carpobrotus* spp. on the pollination success of

native plant species of the Balearic Islands. Biological Conservation 122:611–619.

Morales, C. L., and A. Traveset. 2008. Interspecific pollen transfer: magnitude, prevalence and

consequences for plant fitness. Critical Reviews in Plant Sciences 27:221–238.

Müller, H. 1883. The fertilisation of flowers. Translated by D. W. Thompson. MacMillan,

London, UK.

Neville, Z., and Brownstein, N. 2019. clusterability: Performs tests for cluster tendency of a data

set. R package version 0.1.0.0. https://CRAN.R-project.org/package=clusterability

Norfolk, O., M. P. Eichhorn, and F. Gilbert. 2016. Flowering ground vegetation benefits wild

pollinators and fruit set of almond within arid smallholder orchards. Insect Conservation

and Diversity 9:236–243.

Nowakowska, E., J. Koronacki, and S. Lipovetsky. 2015. Clusterability assessment for Gaussian

mixture models. Applied Mathematics and Computation 256:591–601.

Ogilvie, J. E., and J. D. Thomson. 2016. Site fidelity by bees drives pollination facilitation in

sequentially blooming plant species. Ecology 97:1442–1451.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.

B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2019.

vegan: Community ecology package. R package version 2.5-4. https://CRAN.R-

project.org/package=vegan

Olesen, J. M., J. Bascompte, Y. L. Dupont, and P. Jordano. 2007. The modularity of pollination

networks. Proceedings of the National Academy of Sciences USA 104:19891–19896.

Ollerton, J. 2017. Pollinator diversity: distribution, ecological function, and conservation.

Annual Review of Ecology, Evolution, and Systematics 48:353–376.

Ollerton, J., A. Killick, E. Lamborn, S. Watts, and M. Whiston. 2007. Multiple meanings and

modes: on the many ways to be a generalist flower. Taxon 56:717–728.

Ollerton, J., R. Alarcón, N. M. Waser, M. V. Price, S. Watts, L. Cranmer, A. Hingston, C. I.

Peter, and J. Rotenberry. 2009. A global test of the pollination syndrome hypothesis. Annals

of Botany 103:1471–1480.

Ollerton, J., A. Rodrigo Rech, N. M. Waser, and M. V. Price. 2015. Using the literature to test

pollination syndromes – some methodological cautions. Journal of Pollination Ecology

16:119–125.

Pellegrino, G., F. Bellusci, and A. Musacchio. 2008. Double floral mimicry and the magnet

species effect in dimorphic co-flowering species, the deceptive orchid *Dactylorhiza*

sambucina and rewarding *Viola aethnensis*. Preslia 80:411–422.

Pellegrino, G., F. Bellusci, and A. M. Palermo. 2016. Who helps whom? Pollination strategy of

Iris tuberosa and its relationship with a sexually deceptive orchid. Journal of Plant Research

129:1051–1059.

R Core team. 2018. R: a language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rathcke, B. 1983. Competition and facilitation among plants for pollination. Pages 305–329 *in*

L. Real, editor. Pollination biology. Academic Press, Orlando, Florida, USA.

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223.

Ricklefs, R. E. 2012. Naturalists, natural history, and the nature of biological diversity.

American Naturalist 179:423–435.

Robertson, C. 1928. Flowers and insects. Lists of visitors of four hundred and fifty-three

flowers. Science Press, Lancaster, Pennsylvania, USA.

Schurr, L., L. Affre, F. Flacher, T. Tatoni, L. Le Mire Pecheux, and B. Geslin. 2019. Pollination

insights for the conservation of a rare threatened plant species, *Astragalus tragacantha*

(Fabaceae). Biodiversity and Conservation 28:1389–1409.

Seifan, M., E.-M. Hoch, S. Hanoteaux, and K. Tielbörger. 2014. The outcome of shared

pollination services is affected by the density and spatial pattern of an attractive neighbour.

Journal of Ecology 102:953–962.

Simovici, D., and K. Hua. 2019. Data ultrametricity and clusterability. Journal of Physics

Conference Series 1334:012002.

Stebbins, G. L. 1970. Adaptive radiation of reproductive characteristics in angiosperms, I:

Pollination mechanisms. Annual Review of Ecology and Systematics 1:307–326.

Stone, G. N., P. Willmer, and J. A. Rowe. 1998. Partitioning of pollinators during flowering in

an African Acacia community. Ecology 79:2808–2827.

Thomson, J. D. 1978. Effects of stand composition on insect visitation in two-species mixtures

of *Hieracium*. American Midland Naturalist 100:431–440.

Tur, C., A. Sáez, A. Traveset, and M. A. Aizen. 2016. Evaluating the effects of pollinator-

mediated interactions using pollen transfer networks: evidence of widespread facilitation in

south Andean plant communities. Ecology Letters 19:576–586.

Vandelook, F., S. B. Janssens, P. Gijbels, E. Fischer, W. Van den Ende, O. Honnay, and S.

Abrahamczyk. 2019. Nectar traits differ between pollination syndromes in Balsaminaceae.

Annals of Botany 124:269–279.

Vandermeer, J. H. 1972. Niche theory. Annual Review of Ecology and Systematics 3:107–132.

Waser, N. M. 1978. Interspecific pollen transfer and competition between co-occurring plant

species. Oecologia 36:223–236.

Waser, N. M., and L. A. Real. 1979. Effective mutualism between sequentially flowering plant

species. Nature 281:670–672.

Waser, N. M., L. Chittka, M. V. Price, N. M. Williams, and J. Ollerton. 1996. Generalization in

pollination systems, and why it matters. Ecology 77:1043–1060.

Wilson, P., and J. D. Thomson. 1996. How do flowers diverge? Pages 88–111 *in* D. G. Lloyd,

and S. C. H. Barrett, editors. Floral biology. Studies on floral evolution in animal-pollinated

plants. Chapman and Hall, New York, New York, USA.

Ye, Z.-M., W.-K. Dai, X.-F. Jin, R. W. Gituru, Q.-F. Wang, and C.-F. Yang. 2014. Competition

and facilitation among plants for pollination: can pollinator abundance shift the plant–plant

interactions? Plant Ecology 215:3–13.

653   SUPPORTING INFORMATION

654   **Appendix S1**

655 Plant species studied, habitat types, flowering time and pollinator sampling effort.

656   **Appendix S2**

657 Orders, families and genera of insect pollinators recorded.

659   TABLE 1. Tests for multimodality of pairwise distances between plant species ($N = 221$) in three

660   different pollinator niche spaces whose axes are defined by insect orders, families and

661   genera.

Axes of pollinator niche space *	Pollinator importance measurement			
	Proportion of visitors		Proportion of visits	
	Dip statistic	*P* value	Dip statistic	*P* value
Orders ($n = 7$)	0.0067	0.96	0.0088	0.64
Families ($n = 94$)	0.0020	0.92	0.0016	0.99
Genera ($n = 346$)	0.0012	0.69	0.00089	0.98

Notes: Each test evaluates the null hypothesis that the distribution of interspecific pairwise

distances in the given pollinator space is unimodal, denoting a single cluster of plant species,

against the alternative hypothesis that the distribution has two or more modes, denoting the

existence of at least two clusters. *P* values determined by Monte Carlo simulation with 10^5

repetitions. Computations were performed with the clusterabilitytest function of the

clusterability package (Neville and Brownstein 2019), called with the following options:

test="dip", distance_metric = "manhattan", d_reps=1e5, d_simulatepvalue = TRUE,

reduction="none", distance_standardize="none".

* n = dimensionality of the pollinator niche space.

TABLE 2. Results of multiple Mantel regressions simultaneously assessing the significance of dissimilarity between plant species in

blooming date and habitat type as predictors of distance in pollinator composition, using three different pollinator niche spaces

whose axes are defined by insect orders, families and genera.

	Pollinator importance measurement							
	Proportion of visitors				Proportion of visits			
	Predictor:				Predictor:			
	Blooming date		Habitat type		Blooming date		Habitat type	
Axes of pollinator niche space *	Parameter estimate	P value	Parameter estimate	P value	Parameter estimate	P value	Parameter estimate	P value
Orders ($n = 7$)	-0.0178	0.65	1.5353	0.21	-0.0456	0.32	0.9529	0.47
Families ($n = 94$)	0.1039	0.0003	1.5937	0.062	0.0912	0.0025	1.7140	0.076
Genera ($n = 346$)	0.1175	0.0001	1.9740	0.001	0.1152	0.0001	2.3612	0.0009

* n = dimensionality of the pollinator niche space.

Captions to figures

FIG. 1. Schematic depiction of three possible arrangements of 30 animal-pollinated plant

species (black dots) in an imaginary pollinator niche space defined by two pollinator types

(horizontal and vertical axes). The coordinates of each species on the axes are the relative

contributions of pollinator types to its pollinator assemblage. It is proposed that the importance

of interspecific competition for pollinators as a major driving force shaping the arrangement of

plants in the pollinator niche space should decline from A to C (see text).

FIG. 2. Distribution of the $N = 221$ plant species studied on the plane defined by axes from

Nonmetric Multidimensional Scaling applied to the matrices of pairwise distances in pollinator

composition (MDS1, MDS2). Original pairwise distances were computed using two different

measurements of relative pollinator importance (proportions of visitors recorded and proportion

of flowers visited), and referred to three different pollinator niche spaces whose axes were

defined by insect orders ($n = 7$ dimensions), families ($n = 94$) and genera ($n = 346$). Stress

values, an inverse measurement of goodness of fit, are shown in graphs. Note that meanings of

MDS1 and MDS2 axes are not comparable across graphs, since axis pairs were obtained from

independent ordination analysis on different data.

FIG. 3. Observed (thick solid lines) and randomly simulated (grey histograms; $N = 10^5$

simulations) values for the mean pairwise distance between plant species in pollinator niche

space. Distances were computed for two different measurements of relative pollinator

importance (proportions of visitors recorded or proportions of flowers visited), and referred to

three different pollinator niche spaces whose axes are defined by insect orders ($n = 7$

dimensions), families ($n = 94$) and genera ($n = 346$). The dashed lines denote the 0.025 and

0.975 quantiles of the distributions of simulated mean values.

Fig. 1.

Fig. 2

Fig. 3