From iron to antibiotics: Identification of conserved bacterial-fungal interactions across diverse partners

Emily C. Pierce[1], Manon Morin[1], Jessica C. Little[2], Roland Liu[1], Joanna Tannous[3], Nancy P. Keller[3,4,5], Benjamin E. Wolfe[6], Laura M. Sanchez[2], and Rachel J. Dutton[1,7]

1. Division of Biological Sciences, University of California San Diego, La Jolla, California, USA
2. Department of Medicinal Chemistry and Pharmacognosy, University of Illinois at Chicago, Chicago, Illinois, USA
3. Department of Medical Microbiology & Immunology, University of Wisconsin-Madison, Madison, Wisconsin, USA
4. Department of Bacteriology, University of Wisconsin – Madison, Madison, Wisconsin, USA
5. Food Research Institute, University of Wisconsin – Madison, Madison, Wisconsin, USA
6. Department of Biology, Tufts University, Medford, MA, USA
7. Center for Microbiome Innovation, Jacobs School of Engineering, University of California San Diego, La Jolla, United States

ABSTRACT

Microbial interactions are major determinants in shaping microbiome structure and function. Although fungi are found across diverse microbiomes, the mechanisms through which fungi interact with other species remain largely uncharacterized. In this work, we explore the diversity of ways in which fungi can impact bacteria by characterizing interaction mechanisms across 16 different bacterial-fungal pairs, involving 8 different fungi and 2 bacteria (*Escherichia coli* and *Pseudomonas psychrophila*). Using random barcode transposon-site sequencing (RB-TnSeq), we identified a large number of bacterial genes and pathways important in fungal interaction contexts. Within each interaction, fungal partners elicit both antagonistic and beneficial effects. Using a panel of phylogenetically diverse fungi allowed us to identify interactions that were conserved across all species. Our data show that all fungi modulate the availability of iron and biotin, suggesting that these may represent conserved bacterial-fungal interactions. Several fungi also appear to produce previously uncharacterized antibiotic compounds. Generating a mutant in a master regulator of fungal secondary metabolite production showed that fungal metabolites are key shapers of bacterial fitness profiles during interactions. This work demonstrates a diversity of mechanisms through which fungi are able to interact with bacterial species. In addition to many species-specific effects, there appear to be conserved interaction mechanisms which may be important across microbiomes.

INTRODUCTION

Despite awareness that fungi have an immense capacity to produce biologically active metabolites and to reshape ecosystems, fungi are frequently overlooked in microbiome studies in favor of a bacterial-centric focus[1–3]. Recently, fungi and other microeukaryotes have received increased attention in sequencing-based studies[4–9], and there is growing interest in exploring the role fungi and bacterial-fungal interactions play in environmental and host-associated microbiomes[10–17]. However, the broader patterns and diversity of bacterial-fungal interaction mechanisms have been challenging to characterize given the abiotic and biotic complexity of many microbiomes. While specific interaction mechanisms have been elucidated for particular pairwise bacterial-fungal associations, little is known about the conservation of these

mechanisms for other bacterial-fungal pairs and thus about general and widespread interaction
mechanisms that could be a major factor in shaping microbiomes.
Fermented foods have been developed as experimentally tractable systems to study multi-
kingdom microbial communities[18–20]. Cheese rind biofilms, in particular, represent an ideal
system to investigate the diversity and conservation of bacterial-fungal interactions due to the
presence of culturable bacterial and fungal species representing diverse phyla. Previous work has
shown that biofilms that form on the surface of cheese are composed of an average of six
bacterial and three fungal genera[18]. These genera are frequently found in natural and human
environments, suggesting that interactions in this system may resemble interspecies interactions
in other microbiomes. Communities of isolated fungal and bacterial species from cheese can be
reassembled in order to investigate how these members interact in coculture[21]. Prior work in this
system has demonstrated that fungi in rind biofilms can have both strong positive and negative
impacts on bacterial growth and that specialized metabolites can impact these
interactions[18,19,22,23].
To systematically characterize the potential effects of fungal species on bacteria within
this system, we have employed an interdisciplinary approach. Specifically, we used the high-
throughput genetic screen RB-TnSeq[24], RNASeq, bacterial cytological profiling, and
metabolomics to investigate the diversity of mechanisms through which eight diverse fungal
species impact two bacteria (*Escherichia coli* or a cheese-associated *Pseudomonas*
psychrophila). A previous study used RB-TnSeq in the cheese rind microbiome system to
identify interspecies interactions and to characterize the occurrence of higher-order interactions
in a microbial community[21]. RB-TnSeq has been used in other systems to identify genes
important for host colonization, to determine essential gene sets, and to characterize genes of
unknown function[25–27].
Here, to infer potential mechanisms of bacterial-fungal interactions, we have developed
customized experimental and computational RB-TnSeq pipelines to identify genes impacting
bacterial fitness in a mixed biofilm with a fungal partner. We observed a diversity of gene
functions that affect bacterial fitness in the presence of a fungal partner. Additionally, we found
conservation among the impacts of yeasts and filamentous molds on bacteria; a key example of
this is the widespread effect on iron metabolism in bacteria, which is mediated by the acquisition
of fungal siderophores. Furthermore, we observed similar effects when we expanded our analysis
to include soil and skin fungi, suggesting that this mechanism is relevant not only within cheese
rind biofilms, but also in other systems. Consistent with the understanding that fungal species
have diverse metabolic repertoires, we find several examples of species-specific effects including
the apparent production of antimicrobial compounds by filamentous fungal species. Analysis of a
fungal mutant defective in secondary metabolite production revealed changes in the bacterial
response compared to the wild type (WT) strain, including genes involved in siderophore uptake
and antibiotic resistance. This work provides new perspectives on the biology and mechanisms
of bacterial-fungal interactions and highlights the key roles of fungi in microbiomes.

RESULTS

Selection of Bacterial-Fungal Interaction Partners

To represent fungal diversity within the cheese environment, we selected a panel of 8
species of commonly found yeasts and filamentous fungi (Figure 1). The selected fungal species

from cheese rind microbiomes, which represent five different genera and three fungal classes,
include two yeasts, *Candida* sp. str. 135E and *Debaryomyces* sp. str. 135B, and filamentous
fungi *Penicillium* sp. str. #12*, Penicillium* sp. str. SAM3*, Penicillium* sp. str. RS17*, Fusarium* sp.
str. 554A*, Scopulariopsis* sp. str. JB370, and *Scopulariopsis* sp. str. 165-5. These fungal genera
were all present at >1% average abundance in 137 geographically diverse cheese microbiomes
analyzed in a previous study of rind diversity[18]. These genera are also found in the human gut
mycobiome[28], in the soil[29] , and in marine environments[30].
The bacterial interaction partners selected were two species of Gammaproteobacteria,
Pseudomonas psychrophila str. JB418 and *Escherichia coli*. We decided to focus on
Proteobacteria for this work, as they are common inhabitants of cheese rind communities from
diverse geographic locations and have been shown to be responsive to the presence of fungi in
experimental community conditions[18] (Supplementary Figure 1 and Supplementary Figure 2). *P.*
psychrophila is a native and relatively uncharacterized cheese community member isolated from
a Robiola due latti cheese rind. *Pseudomonas* species are of interest not only in cheese, but also
in human and soil environments[27,31–33]. *E. coli* was also included as a bacterial partner in this
study in order to take advantage of the vast genetic resources available for this organism. While
E. coli is not a common member of cheese rind communities, it is relevant both as a causative
agent of foodborne illness in cheese and other foods, and as a commensal member of the human
gut microbiome that encounters species from consumed fermented foods[34,35].
107
Characterization of bacterial genes with differential fitness in the presence of fungal
partners
Using a pooled library of barcoded transposon-insertion mutants, RB-TnSeq[24]
experiments and analyses generate a fitness value for each gene, reflecting the importance of a
gene for survival in the experimental condition. Here, we used RB-TnSeq to identify bacterial
mutants that have a differential fitness in the presence of a fungal partner compared to growth
alone. To do this, we created a modified experimental and computational pipeline that allowed us
to measure and quantitatively compare fitness values across multiple conditions. Previous
applications of RB-TnSeq have been designed for intra-condition fitness comparisons, allowing
the quantitative comparison of fitness values of different genes within a given condition.
However, they were not developed for quantitative inter-condition comparisons, which allow the
quantitative comparison of fitness values of the same genes between two different conditions.
Specifically, additional normalization steps were needed for inter-condition comparisons. Our
updated pipeline includes custom R scripts (available at https://github.com/DuttonLab/RB-
TnSeq-Microbial-interactions) which provide step-by-step data visualization to follow the
progress of data transformation from raw number of reads to normalized fitness values.
Pooled *P. psychrophila*[21] or *E. coli*[24] RB-TnSeq mutant libraries were grown as a lawn
for seven days on solid cheese curd agar (CCA) plates[38] either alone or mixed evenly with one of
the eight fungal species (Supplementary Figure 3). Mutant abundances at T0 (inoculation) and
day 7 were measured via barcode sequencing and differences in barcode abundance were used to
calculate gene fitness values within each condition (Supplementary Data 1 and Supplementary
Data 2). To identify genes related to interactions, we looked for mutants whose fitness values
were significantly different in a "with fungus" condition versus alone ($p<0.05$) (Figure 2). We
hereafter refer to these differences as an interaction fitness. In some cases, the presence of a
fungus increases the fitness of a mutant (positive interaction fitness), whereas in others the
fitness of a mutant is decreased (negative interaction fitness).

In total, we found 453 *E. coli* and 692 *P. psychrophila* genes whose disruption leads to
fitness alteration in the presence of at least one of the fungal partners used in this study
(Supplementary Data 3 and Supplementary Data 4). This represents an average of 163 ± 24 *E.*
coli genes per fungal condition and 290 ± 32 *P. psychrophila* genes per fungal condition that
have interaction fitness (Figure 3a). These gene sets represent around ten percent of the genes of
each bacterium. For *E. coli*, interaction fitness values range from -5.66 to 5.72, and for *P.*
psychrophila, -6.18 to 6.02. These values are consistent with previously reported fitness values
in other experimental systems, where fitness values greater than 2 and less than -2 are generally
thought to represent strong fitness effects[24].

Comparison of interaction fitness across fungal partners
To assess the specificity of interactions, we evaluated the intersections of gene sets across
the entire set of fungal interaction conditions as well as all possible subsets (Figure 3b,
Supplementary Data 5 and Supplementary Data 6). A large number of genes show differential
fitness only in the presence of specific fungal partners. For *E. coli*, 40 percent of the genes with
interaction fitness are specific to a single fungus (n=171), and for *P. psychrophila,* 32 percent
(n=219). We identified a number of genes that had interaction fitness across the entire set of
partners (n=30 for *E. coli*, and n=65 for *P. psychrophila*). In addition, around 40 percent (n=269)
of the interaction-related genes for *P. psychrophila* and 30 percent (n=137) for *E. coli* were
common to at least four of the eight fungal interaction conditions. For both *E. coli* and *P.*
psychrophila, growth with *Penicillium* sp. str. #12 and *Penicillium* sp. str. SAM3 results in a
large number of the same genes with significant interaction fitness (Figure 3a and Figure 3b).
These species also do not cluster with the other fungi in Principal Component Analysis (PCA) of
the raw fitness values for all *E. coli* (left) or *P. psychrophila* (right) genes having interaction
fitness in at least one fungal condition (Fig 3c).

Mechanisms of fungal impacts on bacterial gene fitness
To identify potential mechanisms underlying bacterial-fungal interactions, we examined
the functional distribution of genes associated with interaction fitness. Three functional themes
common to multiple fungi used in this study were identified using Clusters of Orthologous Genes
(COG) categorization and functional enrichment analysis (Figure 4, Supplementary Data 7 and
Supplementary Data 8, Supplementary Figure 4), ranking of strength of fitness effect, and
analysis of conservation of the effect across diverse fungal species. These interactions include
antimicrobial stress on the bacterial cell envelope, competition for biotin, and provision of
bioavailable iron.

Penicillium sp. str. #12 and *Penicillium* sp. str. SAM3 induce bacterial envelope stress

Penicillium sp. str. #12 and *Penicillium* sp. str. SAM3 consistently shared impacts on
bacterial mutant fitness, as seen by their large number of network connections for both bacteria
(Figure 3a and Figure 3b- For *E. coli,* of 116 total genes shared, 22 of these genes are shared
only between these two fungi; for *P. psychrophila,* of 187 total genes shared, 19 of these genes
are shared only between these two fungi). The genes within this set suggest that these two fungal
species are producing antibiotic molecules. For *E. coli*, this overlapping gene set includes the
genes encoding for the stress protein CspC, beta-lactam antibiotic resistance membrane protein
Blr, and the MdtK multidrug efflux pump. Mutants of these genes displayed decreased fitness in
the presence of these fungi. This is also supported by the gene set specific to *Penicillium* sp. str.

#12/*E.coli,* which includes a regulator of the EmrA multidrug efflux pump and three gene
components of the Rcs regulator of capsule synthesis system (*rcsA, rcsC, rcsD*). The Rcs system
can respond to damage to the cell envelope and is induced, for example, by membrane-active
antimicrobial peptides[40], peptidoglycan damage caused by inhibition of penicillin-binding
proteins[41], and mutations in lipopolysaccharide biosynthesis genes[42]. Additionally, three genes in
the *waa* operon (*waaY, waaP, waaQ*) affecting the modification of the heptose region of the core
LPS, which plays a crucial role in membrane stability[43], have increased fitness in the presence of
Penicillium sp. str. #12. WaaY activity is dependent on WaaP and WaaQ[43]. Inactivation of *waaY*
has previously been shown to increase *E. coli*'s resistance to antimicrobial peptide LL-37, due to
reduced binding of the peptide to the cell surface[44], and it has also been shown to increase
resistance of *Salmonella typhimurium* to antimicrobial peptides[45].
When grown in coculture with *P. psychrophila,* these two fungi induce an interaction
fitness for universal stress protein A, whose mRNA is stabilized by the protein encoded by *cspC,*
one of the genes with mutants impacted in *E. coli.* With *Penicillium* sp. str. #12*,* three *P.*
psychrophila genes involved in LPS/lipid A/outer membrane biogenesis (*msbA, arnB, waaG*)
also had interaction fitness. While the *P. psychrophila* genome is not as well documented as *E.*
coli's, these data are consistent with the *E. coli* data and with the production of antibiotics by
these fungi.
To investigate fungal antibiotic activity that could be related to the observed changes in
bacterial gene fitness, we modified bacterial cytological profiling (BCP) protocols for use on *in*
vitro cheese biofilms[46]. This microscopy technique was previously developed to rapidly
determine the bacterial cellular pathway targeted by antibiotic compounds. We grew wild type
(WT) or Δ*mdtK E. coli* alone or in a mixed biofilm with *Penicillium* sp. str. #12 or *Penicillium*
sp. str. SAM3 on CCA plates for seven days. Δ*mdtK E. coli* was chosen due to the RB-TnSeq
fitness defect of this mutant specifically in the presence of these *Penicillium.* We performed BCP
on bacterial cells either co-cultured in the mixed biofilms or after 2 days of growth on CCA
plates with known antibiotic compounds as reference controls (Supp Figure 5, Figure 5).
Microscopy showed a strong change in cell morphology for both WT and Δ*mdtK E. coli* when
grown with *Penicillium* compared to growth alone (Figure 5). When cultured with these fungi, *E*
coli cells exhibit a rounded phenotype, consistent with a reduction in cell wall integrity and
reminiscent of cells treated with antibiotics that target cell wall biosynthesis such as mecillinam
and amoxicillin. Δ*mdtK* cells are strongly affected and have spheroplasted, indicative of the
complete loss of structural integrity. Neither of these two fungal strains are known producers of
penicillin, and analysis of the draft genomes failed to detect penicillin biosynthesis gene
clusters[47]. However, these BCP results are consistent with our RB-TnSeq data and suggest that
these two fungal strains are inducing stress on the bacterial cell envelope through an
undetermined mechanism that may involve novel antimicrobials.

Fungi increase bacterial need for biotin biosynthesis

Pseudomonas interaction fitness genes suggested competition for environmental biotin
between *P. psychrophila* and all of the studied fungal partners. Biotin, found in both milk and
cheese,[48] is present in our cheese curd agar medium at 73 nmol/mg and represents an essential
cofactor for enzymes involved in key cellular functions like amino acid metabolism and lipid
synthesis[49]. Three *P. psychrophila* genes associated with biotin biosynthesis mutants (*bioB,*
bioD, bioF) have a negative interaction fitness in the presence of all eight fungi. An additional
three genes are associated with a negative interaction fitness in the presence of seven fungi

(*bioA, bioC, bioH*) (Figure 4). The biotin biosynthesis pathway was significantly enriched in
functional enrichment analysis for all fungal partners except *Penicillium* sp. str. SAM3
(Supplementary Data 8). Although some species of fungi are capable of biotin biosynthesis,
others are capable of only partial synthesis or are incapable [50]. Additionally, differential
expression analysis from RNA-Seq of *E. coli* grown either alone or in the presence of
Penicillium sp. str. #12 showed that *bioA, bioB, bioC, bioD,* and *bioF* were all significantly
upregulated in the presence of the fungus with an average fold change of 4.4 (Supplementary
Data 9). This highlights an increased need for bacterial biotin synthesis, which again strongly
supports that fungi and bacteria are consistently competing for available biotin in the medium.
Notably, this interaction appears to be conserved across both bacteria and all fungi.

Fungi increase iron availability for bacterial community members

Because iron is essential for bacterial growth and cheese is an iron-limited environment,
with free iron levels measured to be approximately 3 ppm, microbial species growing on cheese
require iron chelators with specific transport systems to sustain their growth[21,51–53]. Our RB-
TNseq fitness data revealed that *E. coli* mutants defective in enterobactin transport grow better in
the presence of all fungal partners than they do alone (Figure 4, Figure 6a). These genes include
members of the *fep* operon (*fepC, fepG, fepA*), which encodes enterobactin transport functions,
and *exbD,* which encodes a component of an iron-siderophore transport complex. In the majority
of fungal conditions, mutants in an enterobactin esterase encoded by *fes* in addition to iron-
siderophore related transport genes *fepB, fepD, tonB* and *exbB* also grow better in the presence of
a fungal partner. When grown with any fungal partner, ferric-enterobactin transport is
significantly enriched among all the functions associated with genes that have an interaction
fitness (Supplementary Data 7).

For *P. psychrophila,* we noticed two genes that have a strong fitness defect alone, but
positive interaction fitness when any fungus is present. These genes, Ga0212129_114259 (avg.
interaction fitness 3.3) and Ga0212129_114260 (avg. interaction fitness 5.3) are annotated as
uncharacterized protein DUF3649 and as an uncharacterized iron-regulated membrane protein.
The protein encoded by Ga0212129_114259 is 99% identical to an iron transporter from
Pseudomonas fragi (Acc. WP_133145017.1). Moreover, immediately upstream of these two
genes we find a ferric enterobactin receptor (fepA) and the PfeR-PfeS two-component regulatory
system required for production of the ferric enterobactin receptor, suggesting that this region
may be involved in siderophore uptake. Again, positive interaction fitness for these likely
siderophore-uptake associated genes suggests that possibly more iron is available for *P.*
psychrophila when fungi are present.

To validate the RB-TnSeq results related to the Fep system genes, we performed
competitive mutant fitness assays with a 1:1 ratio of WT *E. coli* and Δ*fepC,* Δ*fepA,* and Δ*fepG*
mutants either alone or in the presence of *Scopulariopsis* sp. str. JB370 or *Penicillium* sp. str.
#12. When no fungus is present, there is a clear growth defect for Δ*fep* mutants. After 7 days,
both Δ*fepA* and Δ*fepG* mutants grew significantly better with both fungi, whereas there was no
difference in wild-type growth with or without fungus. The Δ*fepC* mutant grew significantly
better with *Penicillium* sp. str. #12 (Figure 6b).

Additionally, differential expression analysis from RNA-Seq of *E. coli* grown on CCA
for three days either alone or in the presence of *Penicillium* sp. str. #12 revealed 34 genes (out of
a total of 348 significantly upregulated genes) involved in iron acquisition that are specifically
upregulated in the presence of the fungus (Supplementary Data 10). Notably, we find genes

associated with the uptake machinery for hydroxamate siderophores (including *fhuA* and *fhuE*),
which are commonly produced by fungi. We also observe upregulation of enterobactin
biosynthesis and uptake, suggesting that *E. coli*, even in the presence of fungi, still produces and
utilizes its native siderophore, enterobactin (Figure 6c).
All filamentous molds in this study, but not yeast, produce siderophores that were
detectable by liquid Chrome Azurol S (CAS) assay (Supplementary Figure 6). Given the positive
CAS assays, we next sought to identify the siderophores produced by these fungal species. We
performed liquid chromatography mass spectrometry (LC-MS and LC-MS/MS) for all fungal
species. This metabolomics data showed evidence of the hydroxamate fungal siderophores
coprogen and ferrichrome in *Fusarium* and *Penicillium* species (Figure 6d). Although not
detected in these extracts, *Scopulariopsis* sp. str. JB370 is predicted to make dimethylcoprogen
based on antiSMASH secondary metabolite biosynthesis gene cluster prediction analysis of the
draft genome[54].
We hypothesized that the increased fitness of enterobactin *fep* mutants was due to the
uptake of fungal siderophores through an alternate pathway. Indeed, we found that coprogen and
ferrichrome can rescue the growth defect of Δ*fepA* and Δ*fepC* on CCA (Figure 6e). The *E. coli*
Fhu system has previously been shown to allow uptake of hydroxamate-type siderophores[55–58].
In *E. coli,* ferrichrome uptake is known to be mediated by the outer membrane receptor FhuA,
and coprogen uptake is mediated by the outer membrane receptor FhuE. Mutants of *fhuA* or *fhuE*
alone do not show a growth defect on CCA, likely due to the fact that the enterobactin system is
intact. Thus, to specifically examine fungal siderophore uptake, we constructed mutants of *fhuA*
or *fhuE* in an enterobactin-uptake defective background. Δ*fepA*Δ*fhuA*, Δ*fepA*Δ*fhuE,* and
Δ*fepC*Δ*fhuE* were grown on CCA in the presence or absence of fungal siderophores (Figure 6e).
Combined loss of enterobactin uptake and *fhuA* eliminates the alleviation seen with ferrichrome,
whereas loss of either *fhuA* or *fhuE* in the Δ*fepA* background seems to eliminate the alleviation
seen with coprogen. This suggests that *E. coli* requires *fhuA* for ferrichrome uptake, and both
fhuA and *fhuE* for coprogen uptake.
To validate that the fungi used in this study are alleviating *E. coli's* need for enterobactin
through the same mechanism, equal volumes of WT or Δ*fep*/Δ*fhu* mutants were spotted on CCA
at varying distances from a pre-cultured fungal front (Figure 6f, Supplementary Figure 7).
Growth of Δ*fepA* and Δ*fepC* is restored closest to the fungal fronts of all filamentous molds, but
not yeast species. For *Penicillium* sp. str. #12, *Scopulariopsis* sp. str. JB370, and *Scopulariopsis*
sp. str. 165-5, this effect is lost in the Δ*fepA*Δ*fhuE* and Δ*fepC*Δ*fhuE* double mutant, suggesting it
is *fhuE*-dependent. For *Fusarium* sp. str. 554A and *Penicillium* sp. str. RS17*,* it is both *fhuE* and
fhuA-dependent. For *Penicillium* sp. str. SAM3, loss of *fhuA* decreases but does not totally
eliminate alleviation. Thus, we can conclude that in the vicinity of a fungal partner, *E. coli* is
likely to use and benefit from fungal hydroxamate siderophores that are taken up by the FhuA
and FhuE uptake systems independently of the enterobactin uptake system.
Given that iron limitation is a common challenge across many environments, we wanted
to examine whether fungal species from other ecosystems could also be producing siderophores
accessible to neighboring bacterial species. We performed similar assays with *Aspergillus*
fumigatus, a soil-dwelling filamentous ascomycete that was originally isolated from the lung
tissue of a patient who had aspergillosis[59]. This assay was also performed for *Malassezia*
pachydermatis, a yeast commensal resident on animal skin. *M. pachydermatis* is also sometimes
found on human skin and can act as an opportunistic pathogen; this species has caused
bloodstream infections in hospitalized neonates[60,61]. Interestingly, Δ*fepA*Δ*fhuE* and Δ*fepC*Δ*fhuE*

mutants are rescued next to *A. fumigatus*, whereas the Δ*fepA*Δ*fhuA* mutant lacking the
ferrichrome receptor was not, suggesting that *A. fumigatus* produces a siderophore capable of
being imported through FhuA (Figure 6f). *A. fumigatus* is known to produce the extracellular
siderophores fusarinine C and triacetylfusarinine C, which are not known to be imported via the
Fhu system, and the intracellular siderophore ferricrocin, which is similar to ferrichrome but not
expected to be excreted[62]. Ultimately, it is unclear what siderophore is responsible for the effects
by *A. fumigatus*. We see a similar effect using *M. pachydermatis,* suggesting that bacteria are
able to utilize siderophores from a yeast species using the Fhu system (Figure 6f). *Malassezia*
restricta and *Malassezia globosa* have previously been found to possess genes for siderophore
biosynthesis[63,64]. We performed AntiSMASH[54] analysis on a previously published genome of
this *Malassezia pachydermatis* strain, and were able to identify a NRPS biosynthetic gene cluster
containing a ferrichrome peptide synthetase[54,65,66]. In sum, our results suggest that diverse fungi
can reduce bacterial dependence on their own siderophores by secreting xenosiderophores.

Loss of a fungal secondary metabolite regulator alters the profile of interaction fitness

The cases above show that bacterial gene fitness can be impacted by the production of
fungal secondary metabolites, including antimicrobial compounds and siderophores. Previous
studies have shown that fungal secondary metabolite production is regulated by the master
regulator, LaeA[67]. Loss of LaeA in *Aspergillus* spp., *Fusarium oxysporum,* and *Penicillium*
chrysogenum is associated with loss of secondary metabolite production[68,69]. To test the impact
of alterations in fungal metabolite production on bacterial gene fitness, we generated a Δ*laeA*
mutant in *Penicillium* sp. str. #12.
To assess the impact of the *laeA* knockout on fungal metabolites in *Penicillium* sp. str.
#12, we performed RNA-Seq and liquid chromatography with mass spectrometry (LC-MS) on
the WT and *ΔlaeA* mutant. Fourteen percent of the genome (1925 out of 13261 genes) was
differentially expressed between WT and *ΔlaeA* (Figure 7a, Supplementary Data 11). This is
consistent with previous findings in *A. fumigatus* that LaeA influenced expression of around 10
percent of the fungal genome[67]. GO term enrichment analysis identified melanin, organic
hydroxy compound, phenol-containing compound, pigment, sterigmatocystin, depsipeptide,
lactone, mycotoxin, and organic heteropentacyclic compound biosynthesis as pathways
overrepresented in in the set of 1070 genes more expressed in WT (Supplementary Data 12).
Among the genes more expressed in WT, we find four genes in a nonribosomal peptide
synthetase cluster region predicted by AntiSMASH[54]; these four genes have homology to *sid2,*
sidF, sidH, and *sitT,* genes associated with siderophore biosynthesis and transport in
Aspergillus[62]. This gene cluster also includes genes with homology to *sidJ* and *mirB,* providing
further evidence that this region has a siderophore-related function. Five genes in a predicted
gene cluster encoding production of serinocyclin A and four genes in a predicted cluster
encoding production of nidulanin A are also more highly expressed in WT than in *ΔlaeA.* All
together, this confirms that deletion of *laeA* is likely to reduce production of fungal secondary
metabolites including siderophores. Among the 855 genes more expressed in *ΔlaeA,* GO term
enrichment identified amino acid and nucleoside metabolism, fungal cell wall, and ion transport
(Supplementary Data 12), highlighting a possible reorganization of *Penicillium* sp. str. #12
metabolism in the absence of *laeA*.
LC-MS comparison of the two extracts showed differential production of many
metabolites, 94 of which have a >10-fold change between the two (Supplementary Data 13). Of
these, 93 are less abundant in the *ΔlaeA* mutant, which is consistent with the loss of secondary

metabolite production in the *ΔlaeA* mutant (Figure 7b). Some of these metabolites matched with
known secondary metabolites and their analogs based on mass and fragmentation pattern.
Namely, atlantinone A and cyclopenol were found to be produced by the WT *Penicillium* sp. str.
#12 in >10-fold higher quantity than the *ΔlaeA* mutant, and pyripyropene O was putatively
identified and produced in >2-fold higher quantity. Cyclopenol is an alkaloid that is described as
a mycotoxin in the benzodiazepine class[70] and serves as an biosynthetic intermediate for
viridicatol[71], which has been described as having antibacterial activity against *Staphylococcus*
aureus[72]. Small amounts of viridicatol were indeed found in extracts with high quantities of
cyclopenol. Atlantinone A is a meroterpenoid that is derived from the same biosynthetic pathway
as other mixed polyketide-terpenoids such as andrastins and citrehybridones produced by various
Penicillium and *Aspergillus* species[73]. Both pyripyropene O and atlantinone A have tested
negative for antimicrobial activity when screened against a panel of bacteria, including an *E. coli*
strain[74,75]. The ecological role of these metabolites remains undetermined. In combination with
our RNASeq analysis, these data highlight an important diminution of secondary metabolite
production in the *ΔlaeA* strain.
We next performed RB-TnSeq experiments with the Penicillium sp. str. #12 ΔlaeA
mutant to determine whether this single fungal mutation leads to changes in the genes needed for
bacterial fitness. While some interaction fitnesses are preserved between ΔlaeA and WT, loss of
LaeA appears to significantly change the profile of interactions (Figure 7c). As LaeA is predicted
to impact siderophore production, we would expect that E. coli enterobactin uptake mutants
would not be as strongly impacted by the presence of the ΔlaeA mutant. Although we see
positive interaction fitness of fepA, fepB, fepC and fepG genes when E. coli is grown with WT
Penicillium sp. str. #12 relative to E. coli growth alone, we do not see positive interaction fitness
of fep genes with ΔlaeA Penicillium sp. str. #12 (Supplementary Data 14). Indeed, liquid CAS
assays demonstrate that this mutant produces less siderophores than WT on cheese media (Figure
7d). Together, these data support the hypothesis that loss of a fungal secondary metabolite
regulator corresponds to changes in bacterial interaction fitness effects.
Our RB-TnSeq results, combined with the antibiotic assay BCP, suggested that
antibiotic activity by Penicillium sp. str. #12 causes damage to E. coli's cell envelope. As the
laeA mutant should have a decreased ability to produce antimicrobial compounds, we examined
whether there were changes in the fitness of genes related to responses to antibiotics. The mdtK
efflux pump gene, which was previously seen to have negative interaction fitness when grown
with WT Penicillium sp. str. #12, no longer has an interaction fitness with ΔlaeA, suggesting that
the ΔlaeA mutant has decreased antibiotic production. Although the exact nature of the
antimicrobials produced by this strain is unknown, these results are consistent with previous
studies, which show that LaeA regulates antibiotic production in other fungal species, and with
our metabolite analysis, which showed a decrease in production of many metabolites in ΔlaeA.
Overall, these findings suggest that a single fungal gene can strongly impact the bacterial genes
needed for fitness and in particular point out that fungal specialized metabolite production may
play a large role in determination of bacterial fitness in bacterial-fungal interactions.

DISCUSSION

Fungi have been shown to strongly impact bacterial neighbors in diverse systems, from
soil to polymicrobial infections[77–85]. To characterize the effects of fungi from a simple
microbiome on bacteria, we used a high-throughput genetic screen, RB-TnSeq, to identify

bacterial genes relevant to fungal interactions. We observed a diversity of interactions, in terms of direction (+/-), strength, and mechanism. These mechanisms include effects on important cellular pathways such as biotin synthesis and antibiotic resistance. Our work demonstrates that there is both a large diversity of bacterial-fungal interactions as well as conservation of key interaction mechanisms across different fungi.

One of the strongest and most widespread bacterial-fungal interactions that we observed suggests that fungal species can dramatically modulate the access to iron through the provision of fungal siderophores, such as ferrichrome and coprogen. Although it has long been known that bacteria grown in isolation are able to uptake purified fungal siderophores[55,56], our results demonstrate that this exchange takes place between bacteria and fungi growing in a biofilm and that this exchange can have impacts on the competitive fitness of bacteria.

Hydroxamate siderophore receptors homologous to FhuE and FhuA are widespread in Proteobacteria, suggesting that this interkingdom siderophore transfer may play an important role in altering metal availability in diverse systems. *Pseudomonas aeruginosa* has previously been shown to produce phenazine metabolites that led to upregulation of extracellular siderophore production by *A. fumigatus*[86]. Fhu uptake systems have also been identified in the Gram-positive bacterial pathogens *Streptococcus agalactiae* and *Listeria monocytogenes*, and growth of non-siderophore producing mutants of *Streptomyces coelicolor* was restored by the presence of siderophores from airborne contaminant fungi[87–90]. Additionally, hydroxamate siderophore uptake systems have been shown to impact *Staphylococcus aureus* fitness in a murine infection model[91]. Another study has shown that the presence of a siderophore-producing, cheese-associated *Scopulariopsis* produced siderophores and caused downregulation of siderophore production by *Staphylococcus equorum*[52]. *Bacteroides fragilis,* a human gut symbiont, is also able to use ferrichrome to grow in iron-limiting conditions, and *fhu* genes are expressed by *E. coli* in colonic mucus[92,93]. Some fungi consumed as a part of fermented foods have been shown to survive digestive system transit, and fermented foods are known to contain fungal siderophores[94,95], which could be a source of fungal siderophores in the gut in addition to potential siderophore production by gut-resident species[94–96]. All together, these results suggest that bacterial use of fungal siderophores may be a widespread mechanism of interkingdom interaction. Considering the crucial role of iron acquisition and metabolism in microbial survival, that this conserved interaction mechanism could be a determinant in shaping environmental and host-associated microbiomes.

This study provides new insight into the wide range of fungal impacts on bacteria that can occur even in a simple microbiome. Our data also suggest that even among Proteobacteria, fungal impacts may be diverse. Of the 692 *P. psychrophila* genes and 453 *E. coli* genes that had an interaction fitness with at least one fungal partner, only 58 were homologous based on BLAST (Supplementary Data 15). Further, many genes with interaction fitness do not have a known function. We anticipated that by looking for fungal impacts on *E. coli,* we could leverage the vastly superior genetic information available for this species. However, even in this well-characterized organism, 43% of genes identified as having interaction fitness are annotated as hypothetical or putative. For *P. psychrophila,* 29% of genes with interaction fitness are hypothetical proteins. This highlights that many genes involved in interspecies interactions might yet to be characterized, and that studying microbes in the context of their interactions with other species, and not just in monoculture, provides an avenue for uncovering new areas of biology.

MATERIALS AND METHODS

Fungal Ribosomal RNA Sequencing

Genomic DNA was extracted with phenol-chloroform (pH 8) from cultures of the eight cheese fungal species used in this study. For each extraction: 125 µL of 425–600 µm acid-washed beads and 125 µL of 150–212 µm acid-washed beads were poured in a screw-capped 2 mL tube. 500 µL of 2X buffer B (200 mM NaCl, 20 mM EDTA) and 210 µL of SDS 20% were added to the tube as well as the pellet and 500 µL of Phenol:Chloroform (pH 8). Cells were lysed by vortexing the tubes for 2 min at maximum speed. Aqueous and organic phases were separated by centrifugation at 4°C, 8,000 RPM for 3 min and 450 µL of the aqueous phase (upper phase) were recovered in a 1.5 mL eppendorf tube. 45 µL of sodium acetate 3M and 450 µL of ice cold isopropanol were added before incubating the tubes at −80°C for 10 min. The tubes were then centrifuged for 5 min at 4°C at 13,000 RPM. The pellet was then washed in 750 µL of 70% ice cold ethanol and re-suspended in 50 µL of DNAse/RNAse free water. Following DNA extraction, LROR (ACCCGCTGAACTTAAGC) and LR6 (CGCCAGTTCTGCTTACC)[97] primers were used to amplify the large subunit of the ribosomal RNA and for *Penicillium* species, Bt2a (GGTAACCAAATCGGTGCTGCTTTC) Bt2b (ACCCTCAGTGTAGTGACCCTTGGC)[98] primers were used to amplify the beta-tubulin gene. PCR was performed in a final volume of 50 µL: 25 µL of Q5 polymerase master mix (New England Biolab), 2.5 µL of the forward primer at 10 µM, 2.5 µL of the reverse primer at 10 µM, 100 ng of genomic DNA, and water using the following PCR programs: LSU-(i) 98°C - 30 s, (ii) 35 cycles of: 98°C – 10 s; 52°C – 30 s; 72°C – 1.5 min, (iii) 72°C – 5 min; Beta-tubulin-(i) 98°C - 30 s, (ii) 35 cycles of: 98°C – 10 s; 57°C – 30 s; 72°C – 1 min, (iii) 72°C – 5 min. PCR products were purified with the QIAquick PCR purification kit (Qiagen) and sequenced using the forward and reverse primer by Eton Bioscience Inc (San Diego, USA). Consensus sequences from forward and reverse sequencing reactions were aligned using Geneious version R9 9.1.8 (http://www.geneious.com).he MrBayes[36] plugin for Geneious was used to build the phylogenetic tree with the following parameters: Substitution model- JC69; Rate variation- gamma; Outgroup- *Fusarium sp. str. 554A*; Gamma categories-4; Chain Length- 1100000; Subsampling freq- 200; Heated chains-4; Burn-in length- 100000; Heated chain temp- 0.2; Random seed-1160; Unconstrained branch lengths- 1, 0.1, 1, 1.

Bacterial-Fungal Growth Assays

We aimed to inoculate 60000 bacterial cells alone or with 6000 fungal spores per well on 10% cheese curd agar solid medium (CCA medium)[38] adjusted to pH 7 in a 96-well plate. Each bacterial or bacterial-fungal assay was done in triplicate. After 7 days of growth, the entire well was harvested and homogenized in PBS1X-Tween0.05% prior to dilution and plating on LB with 20 µg/ml cycloheximide (for bacterial counts) or milk plate count agar with 50 µg/ml chloramphenicol (for fungal spore counts). Counts were done at inoculation and after harvest. Final growth counts were then compared in co-culture condition relative to growth alone to identify interaction impacts. Significant growth impacts were determined based on Dunnett's test[99], p-value < 0.05. Plots were made with R package ggplot2 3.2.1[100].

Microbial Culturing for LC-MS/MS extraction

All fungal cultures were grown on plate count agar milk salt (PCAMS; 1 g/L whole milk
powder, 1 g/L dextrose, 2.5 g/L yeast extract, 5 g/L tryptone, 10 g/L sodium chloride, 15 g/L
agar). Plates were kept at room temperature and spores were harvested at 7 days (or until
sporulation was observed) of growth for subsequent experiments. Spores harvested from fungi
were put into PBS and normalized to an OD_{600} of 0.1 for working stock.

Extraction of cultures

Three biological replicates of each condition were plated and extracted from solid agar. For
extraction from solid agar plates, 5 µL of fungal working stock were spotted onto 10% CCA
medium adjusted to pH 7. Following 7 days of growth, agar was removed from the Petri dish and
placed into 50 mL falcon tubes. Acetonitrile (10 mL) was added to each tube and all were
sonicated for 30 minutes. All falcon tubes were centrifuged and liquid was removed from the
solid agar pieces and transferred to 15 mL falcon tubes. The 15 mL falcon tubes containing
liquid were then centrifuged and liquid was again removed from any residual solid debris and
transferred to glass scintillation vials. These liquid extractions were then dried *in vacuo*. Dried
extracts were weighed and diluted with MeOH to obtain 1 mg/mL solutions which were stored at
-20°C until analysis via LC-MS/MS.

LC-MS/MS data collection

High resolution LC-MS and LC-MS/MS data were collected on a Bruker impact II qTOF in
positive mode with the detection window set from 50 to 1500 Da, on a 2.1x150mm C18 cortecs
UPLC column with a flow rate of 0.5mL/min for a gradient of 10-100% ACN with 0.1% formic
acid over 16 minutes. For each sample, 10 μL of a 1mg/mL solution was injected. The ESI
conditions were set with the capillary voltage at 4.5 kV. For MS^2, dynamic exclusion was used,
and the top nine precursor ions from each MS^1 scan were subjected to collision energies scaled
according to mass and charge state for a total of nine data dependent MS^2 events per MS^1. MS^2
data for pooled biological replicates is deposited under MassIVE accession number
MSV000085070. MS^1 and MS^2 data for *ΔlaeA* and WT *Penicillium sp. str. #12* is deposited
under MassIVE accession number MSV000085054 and was collected under identical conditions
on a Bruker compact qTOF.

Molecular Networking

For all extractions, all precursor *m/z*'s that were found in solvent and agar controls (based on
both retention time and mass tolerance) were removed prior to input into GNPS using the
BLANKA algorithm.[101] A molecular network (
https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=464b331ef9d54de9957d23b4f9b9db14) was
created using the online workflow at GNPS. The data was filtered by removing all MS/MS peaks
within +/- 17 Da of the precursor m/z. MS/MS spectra were window filtered by choosing only
the top 6 peaks in the +/- 50Da window throughout the spectrum. The data was then clustered
with MS-Cluster with a parent mass tolerance of .02 Da and a MS/MS fragment ion tolerance of
.02 Da to create consensus spectra. Further, consensus spectra that contained less than 2 spectra
were discarded. A network was then created where edges were filtered to have a cosine score
above 0.7 and more than 6 matched peaks. Further edges between two nodes were kept in the

network if and only if each of the nodes appeared in each other's respective top 10 most similar
nodes. The spectra in the network were then searched against GNPS' spectral libraries. The
library spectra were filtered in the same manner as the input data. All matches kept between
network spectra and library spectra were required to have a score above 0.7 and at least 6
matched peaks. Solvent and agar control files were also loaded into the networks in order to
perform removal based also on fragmentation patterns. All nodes with precursor masses less than
200Da were also removed. The extensive background and low *m/z* Da removal was done to more
accurately reflect the metabolomic profiles of fungal genera in an attempt to represent only true
metabolites. The DEREPLICATOR was used to annotate MS/MS spectra[102,103]. The molecular
networks were visualized using Cytoscape software[104].

RB-TnSeq Assays

All RB-TnSeq assays were performed on 10% CCA medium adjusted to pH 7. Prior to
inoculation, one aliquot of each library was thawed and inoculated into 25 mL of liquid LB-
kanamycin (50 µg/mL). Once the culture reached mid-log phase (OD = 0.6–0.8), 5 mL of that
pre-culture was pelleted and stored at −80°C for the T0 reference in the fitness analysis. The
remaining cells were used to inoculate the fitness assay conditions. For each RB-TnSeq fitness
assay, we aimed to inoculate seven million cells of the bacterial library (on average 50 cells per
insertion mutant). For fitness assays including a fungal partner, 700,000 fungal cells were
inoculated based on spore counts. Pre-cultured cells were washed in PBS1X-Tween0.05%,
mixed with appropriate volumes of quantified fungal spore stocks, and then inoculated by
spreading evenly on a 100 mm petri dish containing 10% CCA medium, pH 7. For each
condition, assays were performed in triplicate. After seven days, the plate was flooded with 1.5
mL of PBS1X-Tween0.05% and cells were scraped off, taking care to not disturb the CCA. The
liquid was then transferred into a 1.5 mL microfuge tube and cells were pelleted by
centrifugation. After removing the supernatant, the cells were washed in 1 mL of RNA-protect
solution (Qiagen, Hilden, Germany), pelleted and stored at −80°C until gDNA extraction.
Genomic DNA was extracted with phenol-chloroform (pH 8) using the same protocol used for
fungal gDNA extraction described above. Samples were stored at −80°C until further analysis.
After gDNA extraction, extracts containing *Penicillium sp. str. #12* DNA were purified
using the OneStep PCR Inhibitor Removal Kit (Zymo Research, CA, USA). Then, the 98°C
BarSeq PCR as described in Wetmore et al., 2015 was used to amplify only the barcoded region
of the transposons. PCR was performed in a final volume of 50 µL: 25 µL of Q5 polymerase
master mix (New England Biolab, MA, USA), 10 µL of GC enhancer buffer (New England
Biolab), 2.5 µL of the common reverse primer (BarSeq_P1 – Wetmore et al., 2015) at 10 µM,
2.5 µL of a forward primer from the 96 forward primers (BarSeq_P2_ITXXX) at 10 µM and
either 200 ng of gDNA for alone conditions, or 2 µg of gDNA for fungal interaction conditions.
For *E. coli* analysis, we performed 84 PCRs (T0 sample and 28 harvest samples in triplicate)
involving 28 different multiplexing indexes. For *P. psychrophila* JB418 analysis, we performed
84 PCRs (T0 sample and 28 harvest samples in triplicate) involving 28 different multiplexing
indexes. We used the following PCR program: (i) 98°C - 4 min, (ii) 30 cycles of: 98°C – 30 s;
55°C – 30 s; 72°C – 30 s, (iii) 72°C – 5 min. After the PCR, for both *E. coli* and *P. psychrophila,*
10 µL of each of the PCR products were pooled together to create the BarSeq sequencing library
and 200 µL of the pooled library were purified using the MinElute purification kit (Qiagen,
Germany). The final elution of the BarSeq library was performed in 30 µL in DNAse and RNAse
free water. The BarSeq libraries were then quantified using Qubit dsDNA HS assay kit

(Invitrogen, CA, USA) and sequenced on HiSeq4000 (75 bp, single-end reads), by the IGM
Genomics Center at the University of California San Diego. The sequencing depth for each
condition varied between 6.1 and 11.7 million reads for *E. coli* and 5.8 and 13.3 million reads for
P. psychrophila.

RB-TnSeq Data Processing

Custom R scripts were used to determine the average fitness scores for each gene across three
RB-TnSeq assay replicates. These scripts are available at https://github.com/DuttonLab/RB-
TnSeq-Microbial-interactions. The Readme document provides an in-depth explanation of all
data processing steps performed in these scripts. In brief, insertion mutants that did not have a
sufficient T0 count (3) in each condition or that are not centrally inserted (10-90% of gene) were
removed from analysis. Counts determined by Wetmore *et al.* scripts were then normalized to a
reference gene (*allB* in *E. coli* and closest protein BLAST match Ga0212129_11710 in *P.*
psychrophila) to be able to compare across conditions and to account for differences in
sequencing depth. These genes do not have a strong fitness effect in any condition based on
former fitness determination developed by Wetmore *et al.*, 2015. Un-normalized strain fitness
was then calculated per insertion mutant as the log2 of the ratio of the corrected counts in the
condition and the corrected counts in the T0 sample. Un-normalized gene fitness and variance
was next calculated by averaging insertion mutants within a gene. These values were then
normalized based on the position of the gene along the chromosome, as well as by the mean of
the data distribution. These steps were all done on individual replicates. Next, the average gene
fitness and associated variance were calculated using the weighted average of the fitness values
across the three different replicates. Then, for each condition, fitness values are corrected by the
mean of the replicate means (the replicate means being the mean values used to center fitness
value within a replicate). Finally, based on the assumption that most genes should have no fitness
effect, we corrected gene fitness values in each condition by the mode (peak of the density
distribution). Final fitness values were then compared for fungal interaction conditions compared
to bacterial alone conditions and comparisons associated with a p-value lower than 5% were
considered significant interaction fitness (alpha parameter=0.05 in
2conditions_FitnessComparison.R code). Networks of fitness values were visualized in
Cytoscape v. 3.5.1[104] and PCA plots were made with R package ggplot2 3.2.1[100] and ggfortify
0.4.7[105]. COG category mapping of *E. coli* and *P. psychrophila* protein sequences was done by
eggNOG-mapper[106] and visualized with R package ggplot2 3.2.1[100].

Bacterial Cytological Profiling

Approximately seven million WT *E. coli* K-12 strain BW25113 or Keio collection *mdtK* mutant
cells [107] were inoculated alone or co-inoculated with 700,000 *Penicillium* sp. str. #12 or
Penicillium sp. str. SAM3 spores on 10% CCA pH 7. After 7 days of growth, 1 mL of T-Base
buffer was added to the surface of the biofilms, and biofilms were scraped into the buffer. For
co-culture conditions, the sample was filtered through a 0.5 µm filter to specifically remove
fungal material. 2 µL of concentrated dye mix (1 µL 1 mg/mL FM4-64, 1 µL 2 mg/mL DAPI in
48 µL T-Base) were added to 20 µL of filtrate. Dye-filtrate mix was spotted onto agarose-LB
pads (1% Agarose, 20% LB liquid medium, 80% ddH$_2$O) and imaged by fluorescence and phase
contrast. Control compound references on CCA medium were obtained by spotting 30 µL of 5x,
10x, 25x, and 100x MIC dilutions of antibiotics onto quadrants on CCA medium pH 7 plates,

allowing to dry, and spread-plating 200 µL of log-phase (OD 0.1) *E. coli* cultures. After two
days of growth, cells near the edge of the zone of inhibition on appropriate dilution spots were
resuspended in 10 µL of prediluted dye mix (1 µL 1 mg/mL FM4-64, 1 µL 2 mg/mL DAPI in
998 µL T-Base) and spotted onto agarose-LB pads and imaged as described above. Resulting
images were deconvolved using Deltavision SoftWorx software (Applied Precision, Inc., WA,
USA), analyzed using Fiji[108], and assembled in Adobe Photoshop (Adobe, CA, USA).

CCA medium biotin quantification

Biotin quantification of CCA medium was performed on three replicate samples by Creative
Proteomics (NY, USA) as follows: 100 mg of each sample was homogenized in water (10
µL/mg) for 1 min three times with the aid of 5-mm metal balls on a MM400 mill mixer.
Methanol at 10 µL/mg was then added. Water-soluble vitamins were extracted by vortex-mixing
for 2 min and sonication in a water batch for 5 min. After centrifugation, the clear supernatants
were cleaned up by solid-phase extraction on a Strata-X (60 mg/mL) cartridge. The eluted
fractions containing water-soluble vitamins were collected, pooled and then dried under a gentle
nitrogen gas flow in a nitrogen evaporator. The residues were dissolved in 200 µL of 10%
methanol. Twenty microliter aliquots were injected to run UPLC-MRM/MS with the use of a
C18 UPLC column and with (+) ion detection and (-) ion detection, respectively. Calibration
curves were prepared by injection of serially-diluted mixed standard solutions of water-soluble
vitamins. Concentrations of detected vitamins were calculated by interpolating the linear
calibration curves.

Δf*ep* mutant competitive growth assays

Approximately 600000 WT and Δ*fepA,* Δ*fepC,* or Δ*fepG* Keio collection[107] mutant cells were
inoculated at a 1:1 ratio either alone or co-inoculated with approximately 60,000 *Penicillium* sp.
str. #12 or *Scopulariopsis* sp. str. JB370 spores on 10% CCA pH 7 in a 96-well plate in four
replicates each. After seven days of growth, the entire well was harvested and homogenized in
PBS1X-Tween0.05% prior to dilution and plating on LB with 20 µg/ml cycloheximide (total
bacterial counts) or with 20 µg/ml cycloheximide and 50 µg/ml kanamycin (bacterial mutant
counts). Final growth counts were then compared in co-culture condition relative to growth alone
to identify interaction impacts. Significant growth impacts were determined by significantly
different growth in the presence of a fungus relative to growth alone based on a two-sample t-test
p-value < 0.05. Plots made with R package ggplot2 3.2.1[100].

RNA-Seq and differential expression analysis of *E. coli* with *Penicillium* sp. str. #12

Approximately seven million *E. coli* cells were inoculated in triplicate either alone or with
approximately 700,000 *Penicillium* sp. str. #12 spores on 10% CCA pH 7. After 3 days, the
biofilms were harvested for RNA extraction and washed with 1 mL of RNAprotect. RNA was
extracted by a phenol-chloroform extraction (pH 8) using the same extraction protocol as for
gDNA extraction. Extractions were then purified with the OneStep PCR Inhibitor Removal Kit
(Zymo Research, CA, USA).
Sequencing libraries were prepared as follows. RNA samples were treated with DNAse
using the 'Rigorous DNAse treatment' for the Turbo DNA-free kit (AMBION, Life
Technologies, Waltham, MA, USA) and RNA concentration was measured by nucleic acid

quantification in Epoch Microplate Spectrophotometer (BioTek, Winooski, VT, USA). Transfer
RNAs and 5S RNA were then removed using the MEGAclear Kit Purification for Large Scale
Transcription Reactions (AMBION, Life Technologies) following manufacturer instructions.
Absence of tRNA and 5S RNA was verified by running 100 ng of RNA on a 1.5% agarose gel
and RNA concentration was quantified by nucleic acid quantification in Epoch Microplate
Spectrophotometer. Also, presence of gDNA was assessed by PCR using universal bacterial 16S
PCR primers (Forward primer: AGAGTTTGATCCTGGCTCAG, Reverse Primer:
GGTTACCTTGTTACGACTT). The PCR was performed in a final volume of 20 μL: 10 μL of
Q5 polymerase master mix (New England Biolabs), 0.5 μL of forward primer 10 uM, 0.5 μL of
reverse primer 10 uM and 5 μL of non-diluted RNA. PCR products were run on a 1.7% agarose
gel and if gDNA was amplified, another DNAse treatment was performed as well as a new
verification of absence of gDNA.
Ribosomal RNA depletion was performed using the Ribominus Transcriptome Isolation
Kit (Yeast and Bacteria) for the *E. coli* alone samples and using both the Ribominus
Transcriptome Isolation Kit (Yeast and Bacteria) and the RiboMinus Eukaryote Kit v2 for the
mixed *E. coli/Penicillium sp. str. #12* samples (ThermoFisher Scientific). For the *E. coli* alone
samples, each sample was divided into two for treatment and then repooled for RNA recovery
with ethanol precipitation. For the *E. coli/*Penicillium sp. str. #12 samples, an equal volume of
the eukaryotic probe and RiboMinus Bacterial Probe Mix were used to deplete both bacterial and
fungal ribosomal RNA and RNA was recovered with ethanol precipitation. Concentrations after
ribosomal RNA depletion were measured using Qubit RNA HS Assay Kits (Invitrogen). The
RNASeq library was produced using the NEBNextUltraTM RNA Library Prep Kit for Illumina
for purified mRNA or ribosome depleted RNA. We prepared a library with a fragment size of
300 nucleotides and used the 10 μM NEBNext Multiplex Oligos for Illumina (Set 1, NEB
#E7335) lot 0091412 and the NEBNext multiplex Oligos for Illumina (Set 2, NEB #E7500) lot
0071412. We performed PCR product purification with 0.8X Agencourt AMPure XP Beads.
Library samples were quantified with Qubit DNA HS Assay Kits before the quality and fragment
size were validated by TapeStation (HiSensD1000 ScreenTape). Library samples were pooled at
a concentration of 15 nM each and were sequenced on HiSeq4000 (50 bp, single-end).
TapeStation assays and sequencing were performed by the IGM Genomics Center at the
University of California San Diego.
Following sequencing, reads were mapped to the concatenated genome of *E. coli* K12
BW25113[109] and *Penicillium* sp. str. #12 using Geneious version R9 9.1.8
(http://www.geneious.com). Only the reads that uniquely mapped to a single location on the *E.*
coli genome section were kept. *E. coli* expression analysis was performed using the following R
packages: Rsamtools (R package version 2.0.3), GenomeInfoDb (R package version 1.20.0),
GenomicFeatures[110] (R package version 1.36.4), GenomicAlignments[110] (R package version
1.20.1), GenomicRanges[110] (R package version 1.36.1) and DESeq2[111] (R package version
1.20.1). We followed the workflow described by Love et al. and performed the differential
expression analysis using the package DESeq2. Differentially expressed genes between two
conditions were selected with an adjusted p-value lower than 5% (Benjamini-Hochberg
correction for multiple testing) and an absolute log2 of fold change equal to or greater than 1.5.

Construction of *E. coli* mutants and visual interaction assays

Visual assays for hydroxamate siderophore stimulation:

Antibiotic assay discs (Whatman) were placed on CCA medium pH 7 with .005% tetrazolium
chloride (an indicator of cellular respiration) and 20 µl of water, 10 µM coprogen or ferrichrome
(EMC Microcollections GmbH) solutions (in water) were slowly pipetted onto the disc and
allowed to absorb. 2.5 µl of *E. coli* 37°C overnight LB cultures of *E. coli* K12 BW25113 WT,
Δ*fepA,* Δ*fepC,* Δ*fhuE,* Δ*fhuA,* Δ*fepA*Δ*fhuE,* Δ*fepC*Δ*fhuE,* or Δ*fepA*Δ*fhuA* mutants[107] and were
spotted next to the discs. Double mutants were constructed as described below. Plates were left
at room temperature until red color developed.

Visual assays for fungal stimulation of bacterial mutants:

Fungal spores were inoculated on CCA pH 7 with .005% tetrazolium chloride. After a week of
fungal growth at room temperature (cheese fungal isolates) or 30°C (gut yeast) , 2.5 µl of *E. coli*
37°C overnight LB cultures were spotted at increasing distances from the fungal front. Plates
were left at room temperature until red color developed. *A. fumigatus* isolate AF293 was
received from Nancy Keller, University of Wisconsin-Madison. *M. pachydermatis* was originally
isolated from the ear of a dog in Sweden (ATCC14522).

Creation of Δ*fepA*Δ*fhuE* and Δ*fepC*Δ*fhuE*:

Chemically competent cells for Δ*fepA* or Δ*fepC* mutants were created. An overnight culture of
Δ*fepA* or Δ*fepC* mutants was diluted 1:100 and grown at 37°C until OD 0.4-0.6. The culture was
placed on ice for 20 minutes and then centrifuged at 4°C for 10 minutes at 6000 rpm to collect
the cells. Supernatant was removed and cells were resuspended in half the previous volume of
pre-cooled 0.1M CaCl2. After being left on ice for 30 minutes, centrifugation was repeated and
supernatant was removed before resuspension in a quarter of the original volume of pre-cooled
0.1M CaCl2/15% glycerol. Cells were aliquoted and stored at -80°C until transformation. These
cells were transformed with the pKD46 plasmid [112], recovered at 30°C and plated on LB plates
with 100 µg/mL Ampicillin and grown at 30°C. Overnight cultures were started from individual
colonies for creation of electrocompetent cells. Overnight cultures of Δ*fepC*-pkD46 or Δ*fepA*-
pkD46 were diluted 1:100 in fresh LB- 100 µg/mL Ampicillin and grown at 30°C until an OD of
0.1. 20 ul of fresh 1 M L-arabinose were added and growth was continued at 30°C until OD 0.4-
0.6. Cells were then chilled on ice for 15 minutes and then centrifuged for ten minutes at 4000
rcf 4°C. Cells were resuspended in 1 mL of ice water and centrifuged for ten minutes at 4000 rcf
at 4°C. Cells were resuspended in 0.5 mL of ice water and centrifuged for ten minutes at 4000 rcf
4°C. Cells were resuspended in 50 uL of ice water and kept on ice until transformation. The
chloramphenicol resistance cassette was amplified from the pKD3 plasmid [112] using the custom
primers: FhuEcatF
(CAGATGGCTGCCTTTTTTACAGGTGTTATTCAGAATTGATACGTGCCGGTAATGGCG
CGCCTTACGCCCC) and FhuEcatR
(CCTCCTCCGGCATGAGCCTGACGACAACATAAACCAAGAGATTTCAAATGCTGGGC
CAACTTTTGGCGAA) and the following PCR conditions (i) 98°C - 30 sec, (ii) 30 cycles of:
98°C – 10 s; 70°C – 20 s; 72°C – 30 s, (iii) 72°C – 5 min. Amplification was performed on 4 ng
of pKD3 plasmid using Q5 High-Fidelity 2X Master Mix (New England Bioloab). The PCR
product was digested for 1 hour with the restriction enzymes DpnI and ClaI at 37°C and then the
PCR product was run on a 1% agarose gel. The PCR product was extracted using the QIAquick
Gel Extraction Kit (Qiagen) and then dialyzed for 4 hours with TE buffer. 1.5 ul of dialyzed PCR
product was used to transform the electrocompetent Δ*fepC*-pkD46 or Δ*fepA*-pkD46 cells. After
2 hours of recovery in SOC medium with 1mM arabinose at 37°C, the transformation was plated
on LB with 50 µg/mL kanamycin and chloramphenicol. Transformants were confirmed to be

Δ*fhuE* with Eton Bioscience Inc. sequencing of the chloramphenicol cassette.

Creation of Δ*fepA*Δ*fhuA:*

Creation was done as with Δ*fepA*Δ*fhuE,* except that the chloramphenicol resistance cassette was
amplified from pKD3 [112] using FhuAcatF (ATCATTCTCGTTTACGTTATCATTCACTTT
ACATCAGAGATATACCAATGAATGGCGCGCCTTACGCCCCAATGGCGCGCCTTACG
CCCC) and FhuAcatR
(GCACGGAAATCCGTGCCCCAAAAGAGAAATTAGAAACGGAAGGTTGCGGTCTGGG
CCAACTTTTGGCGAACTGGGCCAACTTTTGGCGAA) custom primers.

Penicillium sp. str. #12 genome sequencing, assembly, and annotation

Genomic DNA was extracted from *Penicillium* sp. str. #12 using the genomic DNA extraction
protocol described above. High molecular weight DNA (average 16 Kb) was sequenced on the
Oxford Nanopore MinION with a R.9.5 flow cell using 1D^2 sequencing adaptors from kit SQK-
LSK308 (Oxford Nanopore Technologies, Oxford, United Kingdom). Raw data was basecalled
using guppy 3.3.0 (Oxford Nanopore Technologies, Oxford, United Kingdom)(guppy_basecaller
-config dna_r9.5_450bps.cfg --fast5_out) for 1D basecalls and these were used to also obtain
higher accuracy 1D^2 basecalls (guppy_basecaller_1d2 -i 1Dbasecall/workspace/ --config
dba_r9.5_450bps_1d2_raw.cfg --index_file 1Dbasecall/sequencing_summary.txt) These reads
were assembled by canu 1.8[113] and polished by racon 1.4.3[114] four times and by pilon 1.23[115]
once. The final assembly is 38 Mbp and consists of 52 contigs.

Then, *Penicillium* sp. str. #12 genome annotations were obtained by combining genomic and
transcriptomic information from RNASeq. To obtain the gene expression profile of *Penicillium*
sp. str. #12, approximately 700,000 WT *Penicillium* sp. str. #12 spores were inoculated in
triplicate on 10% CCA pH 7. After 3 days, the biofilms were harvested for RNA extraction and
washed with 1mL of RNAprotect. RNA was extracted and RNA-Seq libraries were prepared as
described above with the following modification: Ribosomal RNA depletion was performed
using the RiboMinus Eukaryote Kit v1 and RNA was recovered with ethanol precipitation. After
sequencing, the RNASeq reads from these *Penicillium* sp. str. #12 alone cultures were
concatenated with RNASeq reads from the previously described *E. coli/Penicillium* sp. str. #12
coculture conditions that uniquely mapped to a single location on the *Penicillium* sp. str. #12
genome. The full set of transcriptomic reads were then used as input into the FunGAP annotation
pipeline and 77 million of these reads mapped[116]. This pipeline predicted 13261 protein-coding
genes in the *Penicillium* sp. str. #12 genome. Interproscan[117] was used within the FunGAP
pipeline for function prediction of genes.

Creation and confirmation of *laeA* deletion in *Penicillium* sp. str. #12

Deletion cassette design strategy: In order to knockout *laeA* in *Penicillium* sp. str. #12, the
isolate was first screened for hygromycin and phleomycin resistance, two antibiotics that are
commonly employed by our group as selectable markers. *Penicillium* sp. str. #12 showed a
confirmed sensitivity to both antibiotics. A three round PCR deletion strategy was used to
replace the *laeA* ORF with the *hph* gene, whose expression confers selection on hygromycin[118].
The schematic representation of the *laeA* gene replacement with the hph gene is depicted in
Supplementary Figure 8. The deletion cassette (5'flank- hph- 3'flank) was constructed using

three sequential PCR reactions. In the first PCR round, about 1kb genomic sequence that flank
either the 5' or 3' end of the *laeA* ORF were amplified from strains 12 using respectively the
primer sets P12_KOlaeA_5' F (CTCCGTTGGGCCCTCAC) and 5'R
(GCAATTTAACTGTGATAAACTACCGCATTAAAGCTGTTGATATCGGCAATCAATCA
ATG) or P12_KOlaeA_3'F
(GGTGGGCCTTGACATGTGCAGCCGGTGGAGCGGCGCCTGGTGAATCCTACCCACAT
GG) and 3'R (CGTTGGGAGGAAAAGCTTCTGCG). The *hph* gene was amplified from
plasmid pUCH2-8 using primers hph_F (AGCTTTAATGCGGTAGTTTATCACAG) and hph_R
(CTCCACCGGCTGCACATGTC). A second PCR reaction was performed to assemble by
homologous recombination the three individual fragments from the first round PCR. The
deletion cassettes were finally amplified using the nested primer set, P12_KOlaeA_NestedF
(CAGACGGTCCGCATCCCG) and P12_KOlaeA_NestedR
(GGTCCAGGTGCAGTAGTACTG).
806
Fungal transformation: To generate the deletion strains, a protoplast-mediated transformation
protocol routinely used by our group was employed and improved to achieve a successful
protoplasting of these *Penicillium* sp. strains. Briefly, 109 fresh spores from each strain are
cultured in 500 mL of LMM for 12 h under 25 °C and 280 rpm. Newly born hyphae were
harvested by centrifugation at 8000 rpm for 15 min and hydrolyzed in a mixture of 30 mg Lysing
Enzyme from Trichoderma (Sigma-Aldrich) and 20 mg Yatalase (Fisher Scientific) in 10 mL of
Osmotic Medium. The quality of protoplast was monitored under the microscope after four hours
of shaking under 28 °C and 80 rpm. The protoplast mixture was later overlaid with 10 mL of
trapping buffer and centrifuged for 15 min under 4 oC and 5000 rpm. Protoplast were collected
from the interface, overlaid with an equal volume of STC (1M sorbitol, 10 mM Tris-HCL and 10
mM CaCl2) and decanted by centrifugation at 6000 rpm for 8 min. The protoplast pellet was
resuspended in 500 µL STC and used for transformation.
After 5 days of incubation at 25 °C, colonies grown on SMM plates supplemented with
hygromycin were subjected to a second round of selection on hygromycin plates. In total, 25
hygromycin-resistant transformants were isolated after a rapid screening procedure on SMM
supplemented with hygromycin. Single-spored transformants were later tested for proper
homologous recombination at the ORF locus by PCR and Southern blot analysis.
824
Gene-deletion strain confirmation: The correct replacement of the *laeA* with the *hph* gene was
first verified by PCR analysis of genomic DNA derived from the transformant strains using
primer set P12_laeA_F (CACAATGGCTGAACACTCTCGG) and P12_laeA_R
(GGGATATGGAGCATCGAAGTTGC), that amplify the *laeA* ORF. About 12% (3/25) of the
monoconidial lines generated from primary transformants of stains 12 were PCR-positive for the
absence of the *laeA* ORF. The positive deletion strains were further checked for a single insertion
of the deletion cassette by Southern blot analysis and revealed single-site integration of the
deletion cassette in one transformant (Supplementary Figure 8). Probes corresponding to the 5'
and 3' flanks of the *laeA* gene in each strain were labeled using [α32P] dCTP (PerkinElmer,
USA) following the manufacturer's instructions.
835
RNA-Seq analysis of WT and Δ*laeA Penicillium sp. str. #12*
To characterize the effect of the *laeA* deletion on the *Penicillium* sp. str. #12 gene expression
profile, we performed RNA-Seq analysis for Δ*laeA Penicillium* sp. str. #12. As for WT

Penicillium sp. str. #12, 700,000 Δ*laeA Penicillium* sp. str. #12 spores were inoculated in
triplicate on 10% CCA pH 7 and biofilms were harvested after 3 days. Harvest, RNA extraction
and library preparation were performed identically to WT *Penicillium* sp. str. #12. Then,
Penicillium sp. str. #12 and Δ*laeA* differential expression analysis was performed as described
for *E. coli/Penicillium* sp. str. #12 above. To look for enrichment of functions in the set of
differentially expressed genes, we input the protein sequences of the genes into the gene-list
enrichment function of KOBAS 3.0[119]. Sequences were searched against the Gene Ontology
(GO) database[120,121] using *A. fumigatus* as a reference for GO assignment before conducting a
hypergeometric test with Benjamini-Hochberg correction. Functions with a corrected p-value
<.05 were considered enriched.

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Library, strain background (*Escherichia coli* K12)	Keio collection	PMID: 16738554	CGSC, RRID:SCR_0 02303	Collection of 3,818 *E. coli* knockout strains
Library, strain background (Escherichia coli K12)	Keio_ML9	PMID: 25968644		RB-TnSeq library of *E. coli* K12 BW25113 (152,018 pooled insertion mutants)
Library, strain background (*Pseudomonas psychrophila*)	JB418_ECP1	PMID: 30211673		RB-TnSeq library generated in the *P. psychrophila* JB418 strain isolated from cheese (272,329 pooled insertion mutants)
Strain, strain background (*Escherichia coli* K12)	Keio ME9062	PMID: 16738554	CGSC#: 7636	Parent strain of the Keio collection mutants
Penicillium sp. str. SAM3		Danisco - CHOOZIT	PC SAM 3 LYO 10D	Industrial starter for cheese production
P. psychrophila	*Pseudomonas psychrophila* JB418	PMID: 30211673		Strain isolated from Robiola due latti cheese
Candida sp. str. 135E		This paper		Strain isolated from chees
Debaryomyces sp. str. 135B		This paper		Strain isolated from cheese
Penicillium sp. str. #12		This paper		Strain isolated from cheese
Penicillium sp. str. RS17		This paper		Strain isolated from cheese
Scopulariopsis sp. str. JB370.		This paper		Strain isolated from cheese

Scopulariopsis sp. str. 165-5		This paper		Strain isolated from cheese
Fusarium sp. str. 554A		This paper		Strain isolated from cheese
Aspergillus fumigatus	*A. fumigatus* str. AF293	Nancy Keller, University of Wisconsin-Madison		
Malassezia pachydermatis	*M. pachydermatis* str. CBS 1879	ATCC14522		

ACKNOWLEDGMENTS

The authors would like to thank: the Arkin lab and the Deutschbauer lab at UC-Berkeley for the *E. coli* Keio_ML9 library, Kristen Jepsen at the IGM Genomics Center at the University of California San Diego for assistance with sequencing, Dr. Sergey Kryazhimskiy (UCSD) for his input on RB-TnSeq data processing, Cong Dinh (UCSD) for assistance with fungal genome assembly, William Bushnell for assistance with fitness validation experiments, Dr. Sinem Beyhan for advice on fungal genome annotation, Drs. Joe and Kit Pogliano (UCSD) for the use of their microscope, Lisa Marotz (UCSD) for assistance with non-cheese yeasts, and members of the Dutton lab, especially Brooke Anderson, for constructive comments on the manuscript.

FUNDING SOURCES
This work was supported by NSF grant MCB-1817887: Chemical and genetic mechanisms driving microbial interactions in a model microbiome (R.D. and L.S.), the UCSD Center for Microbiome Innovation (E.P.), the UCSD Ruth Stern Award (E.P.), NIH Institutional Training grant 5 T32 GM 7240-40 (E.P.), and NIH R01GM112739-01 (N.P.K.).

REFERENCES

1. Laforest-Lapointe, I. & Arrieta, M.-C. Microbial Eukaryotes: a Missing Link in Gut

   Microbiome Studies. *mSystems* **3**, (2018).

2. Forbes, J. D., Bernstein, C. N., Tremlett, H., Van Domselaar, G. & Knox, N. C. A Fungal

   World: Could the Gut Mycobiome Be Involved in Neurological Disease? *Front. Microbiol.*

   9, 3249 (2018).

3.   Huseyin, C. E., O'Toole, P. W., Cotter, P. D. & Scanlan, P. D. Forgotten fungi-the gut

mycobiome in human health and disease. *FEMS Microbiol. Rev.* **41**, 479–511 (2017).

4.   Bergelson, J., Mittelstrass, J. & Horton, M. W. Characterizing both bacteria and fungi

improves understanding of the Arabidopsis root microbiome. *Sci. Rep.* **9**, 24 (2019).

5.   Huffnagle, G. B. & Noverr, M. C. The emerging world of the fungal microbiome. *Trends*

Microbiol. **21**, 334–341 (2013).

6.   Soverini, M. *et al.* HumanMycobiomeScan: a new bioinformatics tool for the

characterization of the fungal fraction in metagenomic samples. *BMC Genomics* **20**, 496

(2019).

7.   Bradford, L. L. & Ravel, J. The vaginal mycobiome: A contemporary perspective on fungi

in women's health and diseases. *Virulence* **8**, 342–351 (2017).

8.   Marsh, A. J., O'Sullivan, O., Hill, C., Ross, R. P. & Cotter, P. D. Sequence-based analysis

of the bacterial and fungal compositions of multiple kombucha (tea fungus) samples. *Food*

Microbiol. **38**, 171–178 (2014).

9.   De Filippis, F., Laiola, M., Blaiotta, G. & Ercolini, D. Different Amplicon Targets for

Sequencing-Based Studies of Fungal Diversity. *Appl. Environ. Microbiol.* **83**, (2017).

10.   Fan, D. *et al.* Activation of HIF-1α and LL-37 by commensal bacteria inhibits Candida

albicans colonization. *Nat. Med.* **21**, 808–814 (2015).

11.   Jiang, T. T. *et al.* Commensal Fungi Recapitulate the Protective Benefits of Intestinal

Bacteria. *Cell Host Microbe* **22**, 809–816.e4 (2017).

12.   Wheeler, M. L. *et al.* Immunological Consequences of Intestinal Fungal Dysbiosis. *Cell*

Host Microbe **19**, 865–873 (2016).

13.   Peleg, A. Y., Hogan, D. A. & Mylonakis, E. Medically important bacterial–fungal

interactions. *Nat. Rev. Microbiol.* **8**, 340–349 (2010).

14. Mason, K. L. *et al.* Interplay between the gastric bacterial microbiota and Candida albicans

during postantibiotic recolonization and gastritis. *Infect. Immun.* **80**, 150–158 (2012).

15. Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E. E. & van der Heijden, M. G. A. Fungal-

bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat.*

Commun. **10**, 4841 (2019).

16. Durán, P. *et al.* Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival.

Cell **175**, 973–983.e14 (2018).

17. Tourneroche, A. *et al.* Bacterial–Fungal Interactions in the Kelp Endomicrobiota Drive

Autoinducer-2 Quorum Sensing. *Frontiers in Microbiology* vol. 10 (2019).

18. Wolfe, B. E., Button, J. E., Santarelli, M. & Dutton, R. J. Cheese Rind Communities Provide

Tractable Systems for In Situ and In Vitro Studies of Microbial Diversity. *Cell* **158**, 422–

433 (2014).

19. Wolfe, B. E. & Dutton, R. J. Fermented foods as experimentally tractable microbial

ecosystems. *Cell* **161**, 49–55 (2015).

20. May, A. *et al.* Kombucha: a novel model system for cooperation and conflict in a complex

multi-species microbial ecosystem. *PeerJ* **7**, e7565 (2019).

21. Morin, M., Pierce, E. C. & Dutton, R. J. Changes in the genetic requirements for microbial

interactions with increasing community complexity. *Elife* **7**, (2018).

22. Zhang, Y., Kastman, E. K., Guasto, J. S. & Wolfe, B. E. Fungal networks shape dynamics of

bacterial dispersal and community assembly in cheese rind microbiomes. *Nat. Commun.* **9**,

336 (2018).

23. Cleary, J. L., Kolachina, S., Wolfe, B. E. & Sanchez, L. M. Coproporphyrin III Produced by

the Bacterium Glutamicibacter arilaitensis Binds Zinc and Is Upregulated by Fungi in

Cheese Rinds. *mSystems* **3**, (2018).

24. Wetmore, K. M. *et al.* Rapid Quantification of Mutant Fitness in Diverse Bacteria by

Sequencing Randomly Bar-Coded Transposons. *MBio* **6**, (2015).

25. Rubin, B. E. *et al.* The essential gene set of a photosynthetic organism. *Proceedings of the*

National Academy of Sciences **112**, E6634–E6643 (2015).

26. Price, M. N. *et al.* Mutant phenotypes for thousands of bacterial genes of unknown function.

Nature **557**, 503–509 (2018).

27. Cole, B. J. *et al.* Genome-wide identification of bacterial plant colonization genes. *PLoS*

Biol. **15**, e2002860 (2017).

28. Hallen-Adams, H. E. & Suhr, M. J. Fungi in the healthy human gastrointestinal tract.

Virulence **8**, 352–358 (2017).

29. Frąc, M., Hannula, S. E., Bełka, M. & Jędryczka, M. Fungal Biodiversity and Their Role in

Soil Health. *Front. Microbiol.* **9**, 707 (2018).

30. Richards, T. A., Jones, M. D. M., Leonard, G. & Bass, D. Marine fungi: their ecology and

molecular diversity. *Ann. Rev. Mar. Sci.* **4**, 495–522 (2012).

31. Frey-Klett, P. *et al.* Ectomycorrhizal symbiosis affects functional diversity of rhizosphere

fluorescent pseudomonads. *New Phytol.* **165**, 317–328 (2005).

32. Lopez-Medina, E. *et al.* Candida albicans Inhibits Pseudomonas aeruginosa Virulence

through Suppression of Pyochelin and Pyoverdine Biosynthesis. *PLoS Pathog.* **11**,

e1005129 (2015).

33. Nguyen, D. D. *et al.* Indexing the Pseudomonas specialized metabolome enabled the

discovery of poaeamide B and the bananamides. *Nat Microbiol* **2**, 16197 (2016).

34. Choi, K.-H., Lee, H., Lee, S., Kim, S. & Yoon, Y. Cheese Microbial Risk Assessments - A

Review. *Asian-australas. J. Anim. Sci.* **29**, 307–314 (2016).

35. Perrin, F. *et al.* Quantitative Risk Assessment of Haemolytic and Uremic Syndrome Linked

to O157:H7 and Non-O157:H7 Shiga-Toxin Producing Escherichia coli Strains in Raw Milk

Soft Cheeses : Quantitative Risk Assessment of HUS Linked to Pathogenic STEC in

Cheese. *Risk Anal.* **35**, 109–128 (2015).

36. Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed

models. *Bioinformatics* **19**, 1572–1574 (2003).

37. Jukes, T. H. & Cantor, C. R. Evolution of protein molecules. *Mammalian protein*

metabolism (1969).

38. Cosetta, C. M. & Wolfe, B. E. Deconstructing and Reconstructing Cheese Rind

Microbiomes for Experiments in Microbial Ecology and Evolution. *Curr. Protoc. Microbiol.*

56, e95 (2020).

39. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of

intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

40. Farris, C., Sanowar, S., Bader, M. W., Pfuetzner, R. & Miller, S. I. Antimicrobial peptides

activate the Rcs regulon through the outer membrane lipoprotein RcsF. *J. Bacteriol.* **192**,

4894–4903 (2010).

41. Laubacher, M. E. & Ades, S. E. The Rcs phosphorelay is a cell envelope stress response

activated by peptidoglycan stress and contributes to intrinsic antibiotic resistance. *J.*

Bacteriol. **190**, 2065–2074 (2008).

42. Majdalani, N., Heck, M., Stout, V. & Gottesman, S. Role of RcsF in signaling to the Rcs

phosphorelay pathway in Escherichia coli. *J. Bacteriol.* **187**, 6770–6778 (2005).

43. Yethon, J. A., Heinrichs, D. E., Monteiro, M. A., Perry, M. B. & Whitfield, C. Involvement

of waaY, waaQ, andwaaP in the Modification of Escherichia coliLipopolysaccharide and

Their Role in the Formation of a Stable Outer Membrane. *J. Biol. Chem.* **273**, 26310–26316

(1998).

44. Bociek, K. *et al.* Lipopolysaccharide Phosphorylation by the WaaY Kinase Affects the

Susceptibility of Escherichia coli to the Human Antimicrobial Peptide LL-37. *J. Biol. Chem.*

290, 19933–19941 (2015).

45. Lofton, H., Pränting, M., Thulin, E. & Andersson, D. I. Mechanisms and fitness costs of

resistance to antimicrobial peptides LL-37, CNY100HL and wheat germ histones. *PLoS One*

8, e68875 (2013).

46. Nonejuie, P., Burkart, M., Pogliano, K. & Pogliano, J. Bacterial cytological profiling rapidly

identifies the cellular pathways targeted by antibacterial molecules. *Proc. Natl. Acad. Sci. U.*

S. A. **110**, 16169–16174 (2013).

47. Laich, F., Fierro, F. & Martín, J. F. Production of penicillin by fungi growing on food

products: identification of a complete penicillin gene cluster in Penicillium griseofulvum

and a truncated cluster in Penicillium verrucosum. *Appl. Environ. Microbiol.* **68**, 1211–1219

(2002).

48. Staggs, C. G., Sealey, W. M., McCabe, B. J., Teague, A. M. & Mock, D. M. Determination

of the biotin content of select foods using accurate and sensitive HPLC/avidin binding. *J.*

Food Compost. Anal. **17**, 767–776 (2004).

49. Streit, W. R. & Entcheva, P. Biotin in microbes, the genes involved in its biosynthesis, its

biochemical role and perspectives for biotechnological production. *Appl. Microbiol.*

Biotechnol. **61**, 21–31 (2003).

50. Hall, C. & Dietrich, F. S. The reacquisition of biotin prototrophy in Saccharomyces

cerevisiae involved horizontal gene transfer, gene duplication and gene clustering. *Genetics*

177, 2293–2307 (2007).

51. Bonham, K. S., Wolfe, B. E. & Dutton, R. J. Extensive horizontal gene transfer in cheese-

associated bacteria. *Elife* **6**, (2017).

52. Kastman, E. K. *et al.* Biotic Interactions Shape the Ecological Distributions of

Staphylococcus Species. *MBio* **7**, (2016).

53. Monnet, C., Back, A. & Irlinger, F. Growth of aerobic ripening bacteria at the cheese

surface is limited by the availability of iron. *Appl. Environ. Microbiol.* **78**, 3185–3192

(2012).

54. Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining

pipeline. *Nucleic Acids Res.* **47**, W81–W87 (2019).

55. Hantke, K. Identification of an iron uptake system specific for coprogen and rhodotorulic

acid in Escherichia coli K12. *Mol. Gen. Genet.* **191**, 301–306 (1983).

56. Fecker, L. & Braun, V. Cloning and expression of the fhu genes involved in iron(III)-

hydroxamate uptake by Escherichia coli. *J. Bacteriol.* **156**, 1301–1314 (1983).

57. Sauer, M., Hantke, K. & Braun, V. Ferric-coprogen receptor FhuE of Escherichia coli:

processing and sequence common to all TonB-dependent outer membrane receptor proteins.

J. Bacteriol. **169**, 2044–2049 (1987).

58. Matzanke, B. F., Anemüller, S., Schünemann, V., Trautwein, A. X. & Hantke, K. FhuF, part

of a siderophore-reductase system. *Biochemistry* **43**, 1386–1392 (2004).

59. Pain, A. *et al.* Insight into the genome of Aspergillus fumigatus: analysis of a 922kb region

encompassing the nitrate assimilation gene cluster. *Fungal Genet. Biol.* **41**, 443–453 (2004).

60. Chryssanthou, E., Broberger, U. & Petrini, B. Malassezia pachydermatis fungaemia in a

neonatal intensive care unit. *Acta Paediatr.* **90**, 323–327 (2007).

61. Prohic, A., Jovovic Sadikovic, T., Krupalija-Fazlic, M. & Kuskunovic-Vlahovljak, S.

Malassezia species in healthy skin and in dermatological conditions. *Int. J. Dermatol.* **55**,

494–504 (2016).

62. Haas, H. Fungal siderophore metabolism with a focus on Aspergillus fumigatus. *Nat. Prod.*

Rep. **31**, 1266–1276 (2014).

63. Park, M., Cho, Y.-J., Lee, Y. W. & Jung, W. H. Understanding the Mechanism of Action of

the Anti-Dandruff Agent Zinc Pyrithione against Malassezia restricta. *Sci. Rep.* **8**, 12086

(2018).

64. Gründlinger, M. *et al.* Fungal siderophore biosynthesis is partially localized in peroxisomes.

Mol. Microbiol. **88**, 862–875 (2013).

65. Wu, G. *et al.* Genus-Wide Comparative Genomics of Malassezia Delineates Its Phylogeny,

Physiology, and Niche Adaptation on Human Skin. *PLoS Genet.* **11**, e1005614 (2015).

66. Triana, S. *et al.* Draft Genome Sequence of the Animal and Human Pathogen Malassezia

pachydermatis Strain CBS 1879. *Genome Announc.* **3**, (2015).

67. Perrin, R. M. *et al.* Transcriptional regulation of chemical diversity in Aspergillus fumigatus

by LaeA. *PLoS Pathog.* **3**, e50 (2007).

68. Bok, J. W. & Keller, N. P. LaeA, a regulator of secondary metabolism in Aspergillus spp.

Eukaryot. Cell **3**, 527–535 (2004).

69. Kosalková, K. *et al.* The global regulator LaeA controls penicillin biosynthesis,

pigmentation and sporulation, but not roquefortine C synthesis in Penicillium chrysogenum.

Biochimie **91**, 214–225 (2009).

70. Bräse, S., Encinas, A., Keck, J. & Nising, C. F. Chemistry and biology of mycotoxins and

related fungal metabolites. *Chem. Rev.* **109**, 3903–3990 (2009).

71. Luckner, M. [On the synthesis of quinoline alkaloids in plants. 2. Fermentative conversion

of the penicillin alkaloids cyclopenin and cyclopenol to viridicatin and viridicatol]. *Eur. J.*

Biochem. **2**, 74–78 (1967).

72. Ma, Y.-M. *et al.* A new isoquinolone alkaloid from an endophytic fungus R22 of Nerium

indicum. *Nat. Prod. Res.* **31**, 951–958 (2017).

73. Dalsgaard, P. W. *et al.* Atlantinone A, a Meroterpenoid Produced by Penicillium ribeum and

Several Cheese Associated Penicillium Species. *Metabolites* **2**, 214–220 (2012).

74. Wang, X. *et al.* Chemical epigenetics alters the secondary metabolite composition of guttate

excreted by an atlantic-forest-soil-derived Penicillium citreonigrum. *J. Nat. Prod.* **73**, 942–

948 (2010).

75. Tomoda, H. *et al.* Pyripyropenes, novel ACAT inhibitors produced by Aspergillus

fumigatus. IV. Structure elucidation of pyripyropenes M to R. *J. Antibiot.* **49**, 292–298

(1996).

76. Patti, G. J. *et al.* A view from above: cloud plots to visualize global metabolomic data. *Anal.*

Chem. **85**, 798–804 (2013).

77. Peters, B. M., Jabra-Rizk, M. A., O'May, G. A., Costerton, J. W. & Shirtliff, M. E.

Polymicrobial interactions: impact on pathogenesis and human disease. *Clin. Microbiol.*

Rev. **25**, 193–213 (2012).

78. Shirtliff, M. E., Peters, B. M. & Jabra-Rizk, M. A. Cross-kingdom interactions: Candida

albicans and bacteria. *FEMS Microbiol. Lett.* **299**, 1–8 (2009).

79. Scherlach, K., Graupner, K. & Hertweck, C. Molecular bacteria-fungi interactions: effects

1056  on environment, food, and medicine. *Annu. Rev. Microbiol.* **67**, 375–397 (2013).

1057 80. Wargo, M. J. & Hogan, D. A. Fungal—bacterial interactions: a mixed bag of mingling

1058  microbes. *Curr. Opin. Microbiol.* **9**, 359–364 (2006/8).

1059 81. Leclair, L. W. & Hogan, D. A. Mixed bacterial-fungal infections in the CF respiratory tract.

1060  *Med. Mycol.* **48 Suppl 1**, S125–32 (2010).

1061 82. Kim, D. *et al.* Candida albicans stimulates Streptococcus mutans microcolony development

1062  via cross-kingdom biofilm-derived metabolites. *Sci. Rep.* **7**, 41332 (2017).

1063 83. de Boer, W. & Folman, L. B. Living in a fungal world: impact of fungi on soil bacterial

1064  niche development. *FEMS microbiology* (2005).

1065 84. Johansson, J. F., Paul, L. R. & Finlay, R. D. Microbial interactions in the mycorrhizosphere

1066  and their significance for sustainable agriculture. *FEMS Microbiol. Ecol.* **48**, 1–13 (2004).

1067 85. Tarkka, M. T., Sarniguet, A. & Frey-Klett, P. Inter-kingdom encounters: recent advances in

1068  molecular bacterium-fungus interactions. *Curr. Genet.* **55**, 233–243 (2009).

1069 86. Moree, W. J. *et al.* Interkingdom metabolic transformations captured by microbial imaging

1070  mass spectrometry. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 13811–13816 (2012).

1071 87. Clancy, A. *et al.* Evidence for siderophore-dependent iron acquisition in group B

1072  streptococcus. *Mol. Microbiol.* **59**, 707–721 (2006).

1073 88. Jin, B. *et al.* Iron acquisition systems for ferric hydroxamates, haemin and haemoglobin in

1074  Listeria monocytogenes. *Mol. Microbiol.* **59**, 1185–1198 (2006).

1075 89. Sheldon, J. R. & Heinrichs, D. E. Recent developments in understanding the iron acquisition

1076  strategies of gram positive pathogens. *FEMS Microbiol. Rev.* **39**, 592–630 (2015).

1077 90. Arias, A. A. *et al.* Growth of desferrioxamine-deficient Streptomyces mutants through

1078  xenosiderophore piracy of airborne fungal contaminations. *FEMS Microbiol. Ecol.* **91**,

(2015).

91. Mishra, R. P. N. *et al.* Staphylococcus aureus FhuD2 is involved in the early phase of

staphylococcal dissemination and generates protective immunity in mice. *J. Infect. Dis.* **206**,

1041–1049 (2012).

92. Li, H. *et al.* The outer mucus layer hosts a distinct intestinal microbial niche. *Nat. Commun.*

6, 8292 (2015).

93. Rocha, E. R. & Krykunivsky, A. S. Anaerobic utilization of Fe(III)-xenosiderophores

among Bacteroides species and the distinct assimilation of Fe(III)-ferrichrome by

Bacteroides fragilis within the genus. *Microbiologyopen* **6**, (2017).

94. Emri, T. *et al.* Towards high-siderophore-content foods: optimisation of coprogen

production in submerged cultures of Penicillium nalgiovense. *J. Sci. Food Agric.* **93**, 2221–

2228 (2013).

95. Ong, S. A. & Neilands, J. B. Siderophores in microbially processed cheese. *J. Agric. Food*

Chem. **27**, 990–995 (1979).

96. David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature*

505, 559–563 (2014).

97. Rehner, S. A. & Samuels, G. J. Molecular systematics of the Hypocreales: a teleomorph

gene phylogeny and the status of their anamorphs. *Can. J. Bot.* **73**, 816–823 (1995).

98. Glass, N. L. & Donaldson, G. C. Development of primer sets designed for use with the PCR

to amplify conserved genes from filamentous ascomycetes. *Appl. Environ. Microbiol.* **61**,

1323–1330 (1995).

99. Dunnett, C. W. A Multiple Comparison Procedure for Comparing Several Treatments with a

Control. *J. Am. Stat. Assoc.* **50**, 1096–1121 (1955).

100. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. (2009).

101. Cleary, J. L., Luu, G. T., Pierce, E. C., Dutton, R. J. & Sanchez, L. M. BLANKA: an

Algorithm for Blank Subtraction in Mass Spectrometry of Complex Biological Samples. *J.*

Am. Soc. Mass Spectrom. **30**, 1426–1434 (2019).

102. Mohimani, H. *et al.* Dereplication of microbial metabolites through database search of mass

spectra. *Nature Communications* vol. 9 (2018).

103. Mohimani, H. *et al.* Dereplication of peptidic natural products through database search of

mass spectra. *Nat. Chem. Biol.* **13**, 30–37 (2017).

104. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular

interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

105. Tang, Y., Horikoshi, M. & Li, W. ggfortify: Unified Interface to Visualize Statistical Result

of Popular R Packages. *The R Journal* vol. 8 (2016).

106. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.*

47, D309–D314 (2019).

107. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

108. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

109. Grenier, F., Matteau, D., Baby, V. & Rodrigue, S. Complete Genome Sequence of

Escherichia coli BW25113. *Genome Announc.* **2**, (2014).

110. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS Comput.*

Biol. **9**, e1003118 (2013).

111. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

112. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 6640–6645

(2000).

113. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

114. Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome assembly

from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

115. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

116. Min, B., Grigoriev, I. V. & Choi, I.-G. FunGAP: Fungal Genome Annotation Pipeline using

evidence-based gene model evaluation. *Bioinformatics* **33**, 2936–2937 (2017).

117. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

118. Lim, F. Y., Sanchez, J. F., Wang, C. C. C. & Keller, N. P. Toward awakening cryptic

secondary metabolite gene clusters in filamentous fungi. *Methods Enzymol.* **517**, 303–324

(2012).

119. Xie, C. *et al.* KOBAS 2.0: a web server for annotation and identification of enriched

pathways and diseases. *Nucleic Acids Res.* **39**, W316–22 (2011).

120. The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing

strong. *Nucleic Acids Res.* **47**, D330–D338 (2019).

121. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

Consortium. *Nat. Genet.* **25**, 25–29 (2000).

— 0.04

1178
1179
1180
1181 **Figure 1: Fungal interaction partners span the phylogenetic and morphological diversity of**
1182 **the cheese ecosystem.** Phylogenetic tree based on large subunit rRNA of the cheese fungi used
1183 as interaction partners in this study. The tree was built using Bayesian phylogenetic inference
1184 with MrBayes[36] and the Jukes and Cantor substitution model[37]. Branch labels display posterior
1185 probability.

many genes have either positive (blue) or negative (orange) interaction fitness. **a,** *E. coli.* **b,** *P. psychrophila.*

Figure 3: Bacterial genes with differential fitness in the presence of fungi. **a,** Network of *E. coli* (left) or *P. psychrophila* (right) genes with an interaction fitness based on RB-TnSeq. Each orange node represents a fungal partner and is labeled as follows: Fungal partner (Number of genes with interaction fitness; Number of genes with interaction fitness unique to this condition). Each green node represents a bacterial gene. They are colored by the number of fungal conditions in which this gene has an interaction fitness as shown in the legend below. They are sized by average strength of interaction fitness across partners. **b,** UpSet[39] plots showing the overlaps, or intersections, of *E. coli* (left) or *P. psychrophila* (right) gene sets with interaction fitness across fungal partners. These UpSet plots are conceptually similar to Venn Diagrams. The connected circles indicate which fungal conditions are included in the intersection, and the size of the intersection (the number of genes that have an interaction fitness in all the highlighted conditions) is displayed in the main bar chart. Intersections <5 genes are not shown. For example, in the *E. coli* panel, 30 genes have an interaction fitness with all partners (all fungi circles are connected), while 22 other genes have an interaction fitness with *Penicillium* sp. str. #12 and with *Penicillium* sp. str. SAM3 (only *Penicillium* sp. str. #12 *and Penicillium* sp. str. SAM3 circles are connected). **c,** PCA of the raw fitness values for all *E. coli* (left) or *P. psychrophila* (right) genes with an interaction fitness in at least one fungal condition.

Figure 4: Functional analysis of the bacterial genes associated with interaction fitness.
Comparison of *E. coli* (left) or *P. psychrophila* (right) gene fitness values alone compared to
fitness values with a fungal partner, colored by COG category and sized by the conservation of
effect among fungal partners (1-8 fungal species).

1274

1275
1276
Figure 5: Bacterial cytological profiling of *E. coli* grown in a mixed biofilm with *Penicillium*
sp. str. #12 and *Penicillium* sp. str. SAM3 on CCA plates. The phenotype of *E. coli* grown
with these fungi is similar to that seen when *E. coli* is exposed to antibiotics targeting cell wall
biosynthesis. DAPI dye stains DNA and FM4-64 dye stains bacterial membranes. Scale bars
represent 2 μm.

1301
1302
Figure 6: Utilization of fungal siderophores by *E. coli*. a, RB-TnSeq fitness values for *fep*
operon genes in alone or interactive conditions, showing an increase in fitness in the presence of
fungal species. NS- not significant. **b,** Colony forming units of WT and Δ*fep* mutants after 7 days
of 1:1 competitive growth on CCA. Competitions were performed either alone or with

Penicillium sp. str. #12 or *Scopulariopsis* sp. str. JB370. Asterisk indicates significantly different
growth in the presence of a fungus relative to growth alone (t-test p-value < 0.05). **c,** *E. coli* iron-
related genes upregulated in the presence of *Penicillium* sp. str. #12. Significance cutoff made at
abs($\log_2$(fold-change)) >1.5 and adjusted p-value <0.05. **d,** Fungal siderophores identified by
mass spectrometry. Insets show the node that represents the desferrichrome fragmentation
pattern depicted. Coprogen B and ferrichrome were found by matching fragmentation patterns to
library spectra. Both identifications were confirmed using retention time and fragmentation
matching to a purchased standard. **e,** Visual assays of *Δfep* mutant growth with purified
siderophores coprogen and ferrichrome. **f,** Visual assays of *E. coli* mutant growth at varying
distances from cheese fungi, *A. fumigatus* (soil, human pathogen), and *M. pachydermatis* (skin
commensal).

1353

Figure 7: Fungal metabolite production impacts bacterial-fungal interactions. a,
Differential expression of WT *Penicillium* sp. str. #12 relative to Δ*laeA* after three days of
growth on CCA. Labeled on the volcano plot are the number of genes with a $\log_2$FC of >1.5
(green) or <-1.5 (red) and adjusted p-value of <0.05. **b,** The metabolomics data analysis platform
XCMS[76] was used to compare features detected by LC-MS analyses of Δ*laeA* and WT
Penicillium sp. str. #12 extracts. Features of higher abundance in WT relative to Δ*laeA* are
depicted as green nodes on the top of the mirror plot and features of lower abundance in WT
relative to Δ*laeA* are depicted as red nodes on the bottom. Node radius is proportional to the fold
change of the detected features and color intensity is dependent on p-value. The graph displays
only those features with a p value less than or equal to 0.05, fold change higher than or equal to
10, *m/z* between 200 and 2000 Da, and intensity higher than 500. **c,** *E. coli* genes with significant
interaction fitness with Δ*laeA* and WT *Penicillium* sp. str. #12. Each dot represents a gene, with
colored dots indicating genes with interaction fitness. X and Y values (alone on x-axis and
+fungal partner on y-axis) indicate gene fitness values in each condition, and the numbers in the
lower right hand corner indicate how many genes have either positive (blue) or negative (orange)
interaction fitness. Venn diagrams display the overlap of these gene sets. **d,** Liquid CAS assay of
supernatants from blank control CCA medium, Δ*laeA,* or WT *Penicillium* sp. str. #12. Asterisk
indicates significantly different siderophore production (t-test p-value < 0.05).

SUPPLEMENTAL FIGURE AND TABLE LEGENDS

Supplementary Figure 1: **Impacts of fungal species on bacterial growth after 7 days of coculture on cheese curd agar, pH 7**. CFU: colony forming units. * represents a significant difference in bacterial growth in the presence of the fungal partner relative to alone (Dunnett's test, p-value <.05).

Supplementary Figure 2: **Impacts of bacterial species on fungal growth after 7 days of coculture on cheese curd agar, pH 7**. Spore counts were used as a proxy for fungal growth. * represents a significant difference in fungal growth in the presence of the bacterial partner relative to alone (Dunnett's test, p-value <.05).

Supplementary Figure 3: RB-TnSeq assay workflow. Characterized pooled bacterial mutant libraries were grown in a biofilm either alone or in a mixed biofilm with a fungal partner. After seven days of growth, mutant abundances were compared to the starting library abundances for each condition. Changes in barcode abundances were used to calculate gene fitness values. Genes with fitness values that differed significantly between co-culture and alone conditions (significant interaction fitness) were identified as potentially relevant to fungal interaction.

Supplementary Figure 4: COG categories of genes with interaction fitness. Number of genes with interaction fitness falling into each COG category for *E. coli* (left) or *P. psychrophila* (right).

Supplementary Figure 5: Bacterial Cytological Profiling of △*tolC E.coli* treated with known antibiotic compounds on cheese curd agar. DAPI dye stains DNA and FM4-64 dye stains bacterial membranes. SYTOX green stains nucleic acids but cannot penetrate live cells.

Supplementary Figure 6: **Siderophore production by filamentous molds.** Liquid CAS assay was performed on filtered and concentrated fungal supernatants from three replicates grown in liquid cheese for 12 days. Row A) 1-3: Liquid cheese control 4-6: *Penicillium* SAM3. Row B) 1-3: *Debaryomyces* 135B 4-6: *Penicillium* #12. Row C) 1-3: *Candida* 135E. 4-6: *Penicillium* RS17. Row D) 1-3: *Scopulariopsis* 165-5 Row E) 1-3: *Scopulariopsis* JB370. Row F) 1-3: *Fusarium* 554A. % Siderophore units calculated as $[(A_r - A_s)/(A_r)]*100$, where A_r is the absorbance of the cheese curd agar supernatant blank and A_s is the absorbance of the sample. N=3, error bars show standard deviation.

Supplementary Figure 7: **Fitness defect of *fep* mutants on iron-limiting CCA.** Visual assays of *E. coli* mutant growth spotted alone on CCA pH 7.

Supplementary Figure 8. Deletion of *laeA* gene in *Penicillium* sp. str. #12. A. Schematic representation of the genetic construct for *laeA* deletion in *P. solitum* strain 12. The construct is constituted of the *hph* gene conferring resistance to hygromycin. The positions of the restriction enzyme cutting sites are shown on the map B. Southern blot analyses of genomic DNA from the WT and the Δ*laeA* strains. Ten micrograms of total DNA from each strain was digested with the appropriate enzymes and subjected to southern blot analysis using respectively the 5' flank fragment (blue) and the 3'fragment (grey) as probes. The 1 kb DNA ladder from New England Biolabs was used to determine the size of the expected bands.

Supplementary Data 1: RB-TnSeq fitness values for *E. coli* grown with fungal partners compared

1418 **to alone.**
1419
1420
1421 **Supplementary Data 2: RB-TnSeq fitness values for *P. psychrophila* grown with fungal partners**
1422 **compared to alone.**
1423
1424 **Supplementary Data 3: Genes with significant interaction fitness for *E. coli* grown with fungal**
1425 **partners compared to alone.**
1426
1427 **Supplementary Data 4: Genes with significant interaction fitness for *P. psychrophila* grown with**
1428 **fungal partners compared to alone.**
1429
1430 **Supplementary Data 5: Intersection lists for *E. coli* genes with interaction fitness across all**
1431 **conditions.**
1432
1433 **Supplementary Data 6: Intersection lists for *P. psychrophila* genes with interaction fitness across all**
1434 **conditions.**
1435
1436 **Supplementary Data 7: Functional enrichment results for *E. coli* genes with interaction fitness**
1437 **across all conditions.**
1438
1439 **Supplementary Data 8: Functional enrichment results for *P. psychrophila* genes with interaction**
1440 **fitness across all conditions.**
1441
1442 **Supplementary Data 9: RNA-Seq differential expression analysis for *E. coli* grown with *Penicillium***
1443 **sp. str. #12 (*E. coli* perspective, alone condition as reference).**
1444
1445 **Supplementary Data 10: Iron-related genes that are differentially expressed by *E. coli* when grown**
1446 **with *Penicillium* sp. str. #12 relative to *E. coli* growth alone.**
1447
1448 **Supplementary Data 11: Metabolite production by Δ*laeA* and WT *Penicillium* sp. str. #12. Δ*laeA***
1449 **used as the reference condition.**
1450
1451 **Supplementary Data 12: RB-TnSeq fitness values for *E. coli* grown with Δ*laeA* or WT *Penicillium***
1452 **sp. str. #12 compared to alone.**
1453
1454 **Supplementary Data 13: RNA-Seq differential expression analysis for Δ*laeA* vs. WT *Penicillium* sp.**
1455 **str. (WT used as the reference condition).**
1456
1457 **Supplementary Data 14: Functional enrichment results for genes differentially expressed in Δ*laeA***
1458 **vs. WT *Penicillium* #12.**
1459
1460 **Supplementary Data 15: Overlap of *E. coli* or *P. psychrophila* genes with interaction fitness.**

1466
1467