From iron to antibiotics: Identification of conserved bacterial-fungal interactions across diverse partners

Emily C. Pierce[1], Manon Morin[1], Jessica C. Little[2], Roland B. Liu[1], Joanna Tannous[3], Nancy P. Keller[3,4,5], Benjamin E. Wolfe[6], Kit Pogliano[1], Laura M. Sanchez[2], and Rachel J. Dutton[1,7]

1. Division of Biological Sciences, University of California, San Diego, La Jolla, California, USA
2. Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois, USA
3. Department of Medical Microbiology & Immunology, University of Wisconsin-Madison, Madison, Wisconsin, USA
4. Department of Bacteriology, University of Wisconsin – Madison, Madison, Wisconsin, USA
5. Food Research Institute, University of Wisconsin – Madison, Madison, Wisconsin, USA
6. Department of Biology, Tufts University, Medford, MA, USA
7. Center for Microbiome Innovation, Jacobs School of Engineering, University of California San Diego, La Jolla, United States

ABSTRACT

Microbial interactions are major determinants in shaping microbiome structure and function. Although fungi are found across diverse microbiomes, the mechanisms through which fungi interact with other species remain largely uncharacterized. In this work, we explore the diversity of ways in which fungi can impact bacteria by characterizing interaction mechanisms across 16 different bacterial-fungal pairs, involving 8 different fungi and 2 bacteria (*Escherichia coli* and *Pseudomonas psychrophila*). Using random barcode transposon-site sequencing (RB-TnSeq), we identified a large number of bacterial genes and pathways important in fungal interaction contexts. Within each interaction, fungal partners elicit both antagonistic and beneficial effects. Using a panel of phylogenetically diverse fungi allowed us to identify interactions that were conserved across all species. Our data show that all fungi modulate the availability of iron and biotin, suggesting that these may represent conserved bacterial-fungal interactions. Several fungi also appear to produce previously uncharacterized antibiotic compounds. Generating a mutant in a master regulator of fungal secondary metabolite production showed that fungal metabolites are key shapers of bacterial fitness profiles during interactions. This work demonstrates a diversity of mechanisms through which fungi are able to interact with bacterial species. In addition to many species-specific effects, there appear to be conserved interaction mechanisms which may be important across microbiomes.

INTRODUCTION

Despite awareness that fungi have an immense capacity to produce biologically active metabolites and to reshape ecosystems, fungi are frequently overlooked in microbiome studies in favor of a bacterial-centric focus[1–3]. Recently, fungi and other microeukaryotes have received increased attention in sequencing-based studies[4–9], and there is growing interest in exploring the role fungi and bacterial-fungal interactions play in environmental and host-associated

microbiome[10–17]. However, the broader patterns and diversity of bacterial-fungal interaction
mechanisms have been challenging to characterize given the abiotic and biotic complexity of
many microbiomes. While specific interaction mechanisms have been elucidated for particular
pairwise bacterial-fungal associations, little is known about the conservation of these
mechanisms for other bacterial-fungal pairs and thus about general and widespread interaction
mechanisms that could be a major factor in shaping microbiomes.
Fermented foods have been developed as experimentally tractable systems to study multi-
kingdom microbial communities[18–20]. Cheese rind biofilms, in particular, represent an ideal
system to investigate the diversity and conservation of bacterial-fungal interactions due to the
presence of culturable bacterial and fungal species representing diverse phyla. Previous work has
shown that biofilms that form on the surface of cheese are composed of an average of six
bacterial and three fungal genera[18]. These genera are frequently found in natural and human
environments, suggesting that interactions in this system may resemble interspecies interactions
in other microbiomes. Communities of isolated fungal and bacterial species from cheese can be
reassembled to investigate how these members interact in co-culture[21]. Prior work in this system
has demonstrated that fungi in rind biofilms can have both strong positive and negative impacts
on bacterial growth and that specialized metabolites can impact these interactions[18,19,22,23].
To systematically characterize the potential effects of fungal species on bacteria within
this system, we have employed an interdisciplinary approach. Specifically, we used the high-
throughput genetic screen RB-TnSeq[24], RNA-Seq, bacterial cytological profiling, and
metabolomics to investigate the diversity of mechanisms through which eight diverse fungal
species impact two bacteria (*Escherichia coli* or a cheese-associated *Pseudomonas*
psychrophila). A previous study used RB-TnSeq in the cheese rind microbiome system to
identify interspecies interactions and to characterize the occurrence of higher-order interactions
in a microbial community[21]. RB-TnSeq has been used in other systems to identify genes
important for host colonization, to determine essential gene sets, and to characterize genes of
unknown function[25–27].
Here, to infer potential mechanisms of bacterial-fungal interactions, we have developed
customized experimental and computational RB-TnSeq pipelines to identify genes impacting
bacterial fitness in a mixed biofilm with a fungal partner. We observed a diversity of gene
functions that affect bacterial fitness in the presence of a fungal partner. Additionally, we found
conservation among the impacts of yeasts and filamentous molds on bacteria; a key example of
this is the widespread effect on iron metabolism in bacteria, which is mediated by the acquisition
of fungal siderophores. Furthermore, we observed similar effects when we expanded our analysis
to include soil and skin fungi, suggesting that this mechanism is relevant not only within cheese
rind biofilms, but also in other systems. Consistent with the understanding that fungal species
have diverse metabolic repertoires, we find several examples of species-specific effects including
the apparent production of antimicrobial compounds by filamentous fungal species. Analysis of a
fungal mutant defective in secondary metabolite production revealed changes in the bacterial
response compared to the wild type (WT) strain, including genes involved in siderophore uptake
and antibiotic resistance. This work provides new perspectives on the biology and mechanisms
of bacterial-fungal interactions and highlights the key roles of fungi in microbiomes.
84
85   **RESULTS**

Selection of Bacterial-Fungal Interaction Partners

To represent fungal diversity within the cheese environment, we selected a panel of 8 species of commonly found yeasts and filamentous fungi (Figure 1). The selected fungal species from cheese rind microbiomes, which represent five different genera and three fungal classes, include two yeasts, *Candida* sp. str. 135E and *Debaryomyces* sp. str. 135B, and filamentous fungi *Penicillium* sp. str. #12, *Penicillium* sp. str. SAM3, *Penicillium* sp. str. RS17, *Fusarium* sp. str. 554A, *Scopulariopsis* sp. str. JB370, and *Scopulariopsis* sp. str. 165-5. These fungal genera were all present at >1% average abundance in 137 geographically diverse cheese microbiomes analyzed in a previous study of rind diversity[18]. These genera are also found in the human gut mycobiome[28], in the soil[29], and in marine environments[30].

The bacterial interaction partners selected were two species of Gammaproteobacteria, *Pseudomonas psychrophila* str. JB418 and *Escherichia coli*. We decided to focus on Proteobacteria for this work, as they are common inhabitants of cheese rind communities from diverse geographic locations and have been shown to be responsive to the presence of fungi in experimental community conditions[18] (Supplementary Figure 1 and Supplementary Figure 2). *P. psychrophila* is a native and relatively uncharacterized cheese community member isolated from a Robiola due latti cheese rind. *Pseudomonas* species are of interest not only in cheese, but also in human and soil environments[27,31–33]. *E. coli* was also included as a bacterial partner in this study to take advantage of the vast genetic resources available for this organism. While *E. coli* is not a common member of cheese rind communities, it is relevant both as a causative agent of foodborne illness in cheese and other foods, and as a commensal member of the human gut microbiome that encounters species from consumed fermented foods[34,35].

Characterization of bacterial genes with differential fitness in the presence of fungal partners

Using a pooled library of barcoded transposon-insertion mutants, RB-TnSeq[24] experiments and analyses generate a fitness value for each gene, reflecting the importance of a gene for survival in the experimental condition. Here, we used RB-TnSeq to identify bacterial mutants that have a differential fitness in the presence of a fungal partner compared to growth alone. To do this, we created a modified experimental and computational pipeline that allowed us to measure and quantitatively compare fitness values across multiple conditions. Previous applications of RB-TnSeq have been designed for intra-condition fitness comparisons, allowing the quantitative comparison of fitness values of different genes within a given condition. However, they were not developed for quantitative inter-condition comparisons, which allow the quantitative comparison of fitness values of the same genes between two different conditions. Specifically, additional normalization steps were needed for inter-condition comparisons. Our updated pipeline includes custom R scripts (available at https://github.com/DuttonLab/RB-TnSeq-Microbial-interactions) which provide step-by-step data visualization to follow the progress of data transformation from raw number of reads to normalized fitness values.

Pooled *P. psychrophila*[21] or *E. coli*[24] RB-TnSeq mutant libraries were grown as a lawn for seven days on solid cheese curd agar (CCA) plates[38] either alone or mixed evenly with one of the eight fungal species (Supplementary Figure 3). Mutant abundances at T0 (inoculation) and day 7 were measured via barcode sequencing and differences in barcode abundance were used to calculate gene fitness values within each condition (Supplementary Data 1 and Supplementary Data 2). To identify genes related to interactions, we looked for mutants whose fitness values

132   were significantly different in a "with fungus" condition versus "alone" condition (p<0.05)
133   (Figure 2). We hereafter refer to these differences as an interaction fitness. In some cases, the
134   presence of a fungus increases the fitness of a mutant (positive interaction fitness), whereas in
135   others the fitness of a mutant is decreased (negative interaction fitness).
136
In total, we found 453 *E. coli* and 692 *P. psychrophila* genes whose disruption leads to
fitness alteration in the presence of at least one of the fungal partners used in this study
(Supplementary Data 3 and Supplementary Data 4). This represents an average of 163 ± 24 *E.*
coli genes per fungal condition and 290 ± 32 *P. psychrophila* genes per fungal condition that
have interaction fitness (Figure 3a). These gene sets represent around ten percent of the genes of
each bacterium. For *E. coli*, interaction fitness values range from -5.66 to 5.72, and for *P.*
psychrophila, -6.18 to 6.02. These values are consistent with previously reported fitness values
in other experimental systems, where fitness values greater than 2 and less than -2 are generally
thought to represent strong fitness effects[24].
146
147   **Comparison of interaction fitness across fungal partners**
148
To assess the specificity of interactions, we evaluated the intersections of gene sets across
the entire set of fungal interaction conditions (Figure 3b, Supplementary Data 5 and
Supplementary Data 6). Many genes show differential fitness only in the presence of specific
fungal partners. For *E. coli*, 40 percent of the genes with interaction fitness are specific to a
single fungus (n=171), and for *P. psychrophila,* 32 percent (n=219). We also identified a number
of genes that had interaction fitness across the entire set of partners (n=30 for *E. coli*, and n=65
for *P. psychrophila*). In addition, around 40 percent (n=269) of the interaction-related genes for
P. psychrophila and 30 percent (n=137) for *E. coli* were common to at least four of the eight
fungal interaction conditions. For both *E. coli* and *P. psychrophila,* growth with *Penicillium* sp.
str. #12 and *Penicillium* sp. str. SAM3 results in a large number of the same genes with
significant interaction fitness (Figure 3a and Figure 3b). These species also do not cluster with
the other fungi in Principal Component Analysis (PCA) of the raw fitness values for all *E. coli*
(left) or *P. psychrophila* (right) genes having interaction fitness in at least one fungal condition
(Fig 3c).
163
164   **Mechanisms of fungal impacts on bacterial gene fitness**
165
To identify potential mechanisms underlying bacterial-fungal interactions, we examined
the functional distribution of genes associated with interaction fitness. Three functional themes
common to multiple fungi used in this study were identified using Clusters of Orthologous Genes
(COG) categorization and functional enrichment analysis (Figure 4, Supplementary Data 7 and
Supplementary Data 8, Supplementary Figure 4) and analysis of conservation of the effect across
fungal species. These interactions include antimicrobial stress on the bacterial cell envelope,
competition for biotin, and provision of bioavailable iron.

Penicillium sp. str. #12 and *Penicillium* sp. str. SAM3 induce bacterial envelope stress

Penicillium sp. str. #12 and *Penicillium* sp. str. SAM3 consistently shared impacts on
bacterial mutant fitness, as seen by their large number of network connections for both bacteria
(Figure 3a and Figure 3b). For *E. coli,* of 116 total genes shared, 22 of these genes are shared

only between these two fungi; for *P. psychrophila,* of 187 total genes shared, 19 of these genes
are shared only between these two fungi. The gene set shared by these fungi suggests that these
two fungal species are producing antibiotic molecules. For *E. coli*, this overlapping gene set
includes the genes encoding for the stress protein CspC, beta-lactam antibiotic resistance
membrane protein Blr, and the MdtK multidrug efflux pump. Mutants of these genes displayed
decreased fitness in the presence of these fungi. This is also supported by the gene set specific to
Penicillium sp. str. #12/*E.coli,* which includes a regulator of the EmrA multidrug efflux pump
and three gene components of the Rcs regulator of capsule synthesis system (*rcsA, rcsC, rcsD*).
The Rcs system can respond to damage to the cell envelope and is induced, for example, by
membrane-active antimicrobial peptides[40], peptidoglycan damage caused by inhibition of
penicillin-binding proteins[41], and mutations in lipopolysaccharide biosynthesis genes[42].
Additionally, three genes in the *waa* operon (*waaY, waaP, waaQ)* affecting the modification of
the heptose region of the core LPS, which plays a crucial role in membrane stability[43], have
increased fitness in the presence of *Penicillium* sp. str. #12. WaaY activity is dependent on WaaP
and WaaQ[43]. Inactivation of *waaY* has previously been shown to increase *E. coli*'s resistance to
antimicrobial peptide LL-37, due to reduced binding of the peptide to the cell surface[44], and it
has also been shown to increase resistance of *Salmonella typhimurium* to antimicrobial
peptides[45].
When grown in co-culture with *P. psychrophila,* these two fungi induce an interaction
fitness for universal stress protein A, whose mRNA is stabilized by the protein encoded by *cspC,*
one of the genes with mutants impacted in *E. coli.* With *Penicillium* sp. str. #12, three *P.*
psychrophila genes involved in LPS/lipid A/outer membrane biogenesis (*msbA, arnB, waaG*)
also had interaction fitness. While the *P. psychrophila* genome is not as well documented as *E.*
coli's, these data are consistent with the *E. coli* data and with the production of antibiotics by
these fungi.
To investigate fungal antibiotic activity that could be related to the observed changes in
bacterial gene fitness, we modified bacterial cytological profiling (BCP) protocols for use on *in*
vitro cheese biofilms[46]. This microscopy technique was previously developed to rapidly
determine the bacterial cellular pathway targeted by antibiotic compounds. We grew wild type
(WT) or Δ*mdtK E. coli* alone or in a mixed biofilm with *Penicillium* sp. str. #12 or *Penicillium*
sp. str. SAM3 on CCA plates for seven days. Δ*mdtK E. coli* was chosen due to the RB-TnSeq
fitness defect of this mutant specifically in the presence of these *Penicillium.* We performed BCP
on bacterial cells either co-cultured in the mixed biofilms or after 2 days of growth on CCA
plates with known antibiotic compounds as reference controls (Supplementary Figure 5, Figure
5). Microscopy showed a strong change in cell morphology for both WT and Δ*mdtK E. coli*
when grown with *Penicillium* compared to growth alone (Figure 5). When cultured with these
fungi, *E coli* cells exhibit a rounded phenotype, consistent with a reduction in cell wall integrity
and reminiscent of cells treated with antibiotics that target cell wall biosynthesis such as
mecillinam and amoxicillin. Δ*mdtK* cells are strongly affected and have spheroplasted, indicative
of the complete loss of structural integrity. Neither of these two fungal strains are known
producers of penicillin, and analysis of the *Penicillium* sp. str. #12 draft genome failed to detect
penicillin biosynthesis gene clusters[47]. However, these BCP results are consistent with our RB-
TnSeq data and suggest that these two fungal strains are inducing stress on the bacterial cell
envelope through an undetermined mechanism that may involve novel antimicrobials.

Fungi increase bacterial need for biotin biosynthesis

Pseudomonas interaction fitness genes suggested competition for environmental biotin between *P. psychrophila* and all of the studied fungal partners. Biotin, found in both milk and cheese[48], is present in our cheese curd agar medium at 73 nmol/mg and represents an essential cofactor for enzymes involved in key cellular functions like amino acid metabolism and lipid synthesis[49]. Three *P. psychrophila* genes associated with biotin biosynthesis mutants (*bioB, bioD, bioF*) have a negative interaction fitness in the presence of all eight fungi. An additional three genes are associated with a negative interaction fitness in the presence of seven fungi (*bioA, bioC, bioH*) (Figure 4). The biotin biosynthesis pathway was also significantly enriched with fungal partners in functional enrichment analysis of interaction fitness gene sets (Supplementary Data 8). Additionally, differential expression analysis from RNA-Seq of *E. coli* grown either alone or in the presence of *Penicillium* sp. str. #12 showed that *bioA, bioB, bioC, bioD,* and *bioF* were all significantly upregulated in the presence of the fungus with an average fold change of 4.4 (Supplementary Data 9). This highlights an increased need for bacterial biotin synthesis, which again supports that fungi and bacteria are consistently competing for available biotin in the medium, or potentially that bacteria have higher biotin requirements in the presence of fungi. Although some species of fungi are capable of biotin biosynthesis, others are capable of only partial synthesis or are incapable[50]. Notably, this interaction appears to be conserved across both bacteria and all fungi.

Fungi increase iron availability for bacterial community members

Because iron is essential for bacterial growth and cheese is an iron-limited environment, with free iron levels measured to be approximately 3 ppm, microbial species growing on cheese require iron chelators with specific transport systems to sustain their growth[21,51–53]. Our RB-TnSeq fitness data revealed that *E. coli* mutants defective in enterobactin transport grow better in the presence of all fungal partners than they do alone (Figure 4, Figure 6a). These genes include members of the *fep* operon (*fepC, fepG, fepA*), which encodes enterobactin transport functions, and *exbD,* which encodes a component of an iron-siderophore transport complex. In most fungal conditions, mutants in an enterobactin esterase encoded by *fes* in addition to iron-siderophore related transport genes *fepB, fepD, tonB* and *exbB* also grow better in the presence of a fungal partner. When grown with any fungal partner, ferric-enterobactin transport is significantly enriched among all the functions associated with genes that have an interaction fitness (Supplementary Data 7).

For *P. psychrophila,* we noticed two genes that have a strong fitness defect alone, but positive interaction fitness when any fungus is present. These genes, Ga0212129_114259 (avg. interaction fitness 3.3) and Ga0212129_114260 (avg. interaction fitness 5.3) are annotated as uncharacterized protein DUF3649 and as an uncharacterized iron-regulated membrane protein. The protein encoded by Ga0212129_114259 is 99% identical to an iron transporter from *Pseudomonas fragi* (NCBI Protein Accession WP_133145017.1). Moreover, immediately upstream of these two genes we find a ferric enterobactin receptor (fepA) and the PfeR-PfeS two-component regulatory system required for production of the ferric enterobactin receptor, suggesting that this region may be involved in siderophore uptake. Again, positive interaction fitness for these likely siderophore-uptake associated genes suggests that possibly more iron is available for *P. psychrophila* when fungi are present.

To validate the RB-TnSeq results related to the Fep system genes, we performed
competitive mutant fitness assays with a 1:1 ratio of WT *E. coli* and Δ*fepC*, Δ*fepA,* and Δ*fepG*
mutants either alone or in the presence of *Scopulariopsis* sp. str. JB370 or *Penicillium* sp. str.
#12. When no fungus is present, there is a clear growth defect for Δ*fep* mutants. After 7 days,
both Δ*fepA* and Δ*fepG* mutants grew significantly better with both fungi, whereas there was no
difference in wild-type growth with or without fungus. The Δ*fepC* mutant grew significantly
better with *Penicillium* sp. str. #12 (Figure 6b).
Additionally, differential expression analysis from RNA-Seq of *E. coli* grown on CCA
for three days either alone or in the presence of *Penicillium* sp. str. #12 revealed 34 genes (out of
a total of 348 significantly upregulated genes) involved in iron acquisition that are specifically
upregulated in the presence of the fungus (Supplementary Data 10). Notably, we find genes
associated with the uptake machinery for hydroxamate siderophores (including *fhuA* and *fhuE*),
which are commonly produced by fungi. We also observe upregulation of enterobactin
biosynthesis and uptake, suggesting that *E. coli*, even in the presence of fungi, still produces and
utilizes its native siderophore, enterobactin (Figure 6c).
All filamentous molds in this study, but not yeast, produce siderophores that were
detectable by liquid Chrome Azurol S (CAS) assay (Supplementary Figure 6). Given the positive
CAS assays, we next sought to identify the siderophores produced by these fungal species. We
performed liquid chromatography mass spectrometry (LC-MS and LC-MS/MS) for all fungal
species. This metabolomics data showed evidence of the hydroxamate fungal siderophores
coprogen and ferrichrome in *Fusarium* and *Penicillium* species (Figure 6d). Although not
detected in these extracts, *Scopulariopsis* sp. str. JB370 is predicted to make dimethylcoprogen
based on antiSMASH analysis of the draft genome[54].
We hypothesized that the increased fitness of enterobactin *fep* mutants was due to the
uptake of fungal siderophores through an alternate pathway. Indeed, we found that coprogen and
ferrichrome can rescue the growth defect of Δ*fepA* and Δ*fepC* on CCA (Figure 6e). The *E. coli*
Fhu system has previously been shown to allow uptake of hydroxamate-type siderophores[55–58].
In *E. coli,* ferrichrome uptake is known to be mediated by the outer membrane receptor FhuA,
and coprogen uptake is mediated by the outer membrane receptor FhuE. Mutants of *fhuA* or *fhuE*
alone do not show a growth defect on CCA, likely because the enterobactin system is intact.
Thus, to specifically examine fungal siderophore uptake, we constructed mutants of *fhuA* or *fhuE*
in an enterobactin-uptake defective background. Δ*fepA*Δ*fhuA,* Δ*fepA*Δ*fhuE,* and Δ*fepC*Δ*fhuE*
were grown on CCA in the presence or absence of fungal siderophores (Figure 6e). Combined
loss of enterobactin uptake and *fhuA* eliminates the alleviation seen with ferrichrome, whereas
loss of either *fhuA* or *fhuE* in the Δ*fepA* background seems to eliminate the alleviation seen with
coprogen. This suggests that *E. coli* requires *fhuA* for ferrichrome uptake, and both *fhuA* and
fhuE for coprogen uptake.
To validate that the fungi used in this study are alleviating *E. coli's* need for enterobactin
through the same mechanism, equal volumes of WT or Δ*fep*/Δ*fhu* mutants were spotted on CCA
at varying distances from a pre-cultured fungal front (Figure 6f, Supplementary Figure 7).
Growth of Δ*fepA* and Δ*fepC* is restored closest to the fungal fronts of all filamentous molds, but
not yeast species. For *Penicillium* sp. str. #12, *Scopulariopsis* sp. str. JB370, and *Scopulariopsis*
sp. str. 165-5, this effect is lost in the Δ*fepA*Δ*fhuE* and Δ*fepC*Δ*fhuE* double mutant, suggesting it
is *fhuE*-dependent. For *Fusarium* sp. str. 554A and *Penicillium* sp. str. RS17*,* it is both *fhuE* and
fhuA-dependent. For *Penicillium* sp. str. SAM3, loss of *fhuA* decreases but does not eliminate
alleviation. Thus, we can conclude that near a fungal partner, *E. coli* is likely to use and benefit

from fungal hydroxamate siderophores that are taken up by the FhuA and FhuE uptake systems
independently of the enterobactin uptake system.
Given that iron limitation is a common challenge across many environments, we wanted
to examine whether fungal species from other ecosystems could also be producing siderophores
accessible to neighboring bacterial species. We performed similar assays with *Aspergillus*
fumigatus, a soil-dwelling filamentous ascomycete that was originally isolated from the lung
tissue of a patient who had aspergillosis[59]. This assay was also performed for *Malassezia*
pachydermatis, a yeast commensal resident on animal skin. *M. pachydermatis* is also sometimes
found on human skin and can act as an opportunistic pathogen; this species has caused
bloodstream infections in hospitalized neonates[60,61]. Interestingly, $\Delta fepA\Delta fhuE$ and $\Delta fepC\Delta fhuE$
mutants are rescued next to *A. fumigatus*, whereas the $\Delta fepA\Delta fhuA$ mutant lacking the
ferrichrome receptor was not, suggesting that *A. fumigatus* produces a siderophore capable of
being imported through FhuA (Figure 6f). *A. fumigatus* is known to produce the extracellular
siderophores fusarinine C and triacetylfusarinine C, which are not known to be imported via the
Fhu system, and the intracellular siderophore ferricrocin, which is similar to ferrichrome but not
expected to be excreted[62]. Ultimately, it is unclear what siderophore is responsible for the effects
by *A. fumigatus*. We see a similar effect using *M. pachydermatis,* suggesting that bacteria are
able to utilize siderophores from a yeast species using the Fhu system (Figure 6f). *Malassezia*
restricta and *Malassezia globosa* have previously been found to possess genes for siderophore
biosynthesis[63,64]. We performed AntiSMASH[54] analysis on a previously published genome of
this *Malassezia pachydermatis* strain, and were able to identify a NRPS biosynthetic gene cluster
containing a ferrichrome peptide synthetase[54,65,66]. In sum, our results suggest that diverse fungi
can reduce bacterial dependence on their own siderophores by secreting xenosiderophores.

Loss of a fungal secondary metabolite regulator alters the profile of interaction fitness

334
The cases above show that bacterial gene fitness can be impacted by the production of
fungal secondary metabolites, including antimicrobial compounds and siderophores. Previous
studies have shown that fungal secondary metabolite production is regulated by the master
regulator, LaeA[67]. Loss of LaeA in *Aspergillus* spp., *Fusarium oxysporum,* and *Penicillium*
chrysogenum is associated with loss of secondary metabolite production[68,69]. To test the impact
of alterations in fungal metabolite production on bacterial gene fitness, we generated a $\Delta laeA$
mutant in *Penicillium* sp. str. #12.
To assess the impact of the *laeA* knockout on fungal metabolites in *Penicillium* sp. str.
#12, we performed RNA-Seq and liquid chromatography with mass spectrometry (LC-MS) on
the WT and *ΔlaeA* mutant. Fourteen percent of the genome (1925 out of 13261 genes) was
differentially expressed between WT and *ΔlaeA* (Figure 7a, Supplementary Data 11). This is
consistent with previous findings in *A. fumigatus* that LaeA influenced expression of around 10
percent of the fungal genome[67]. GO term enrichment analysis identified melanin, organic
hydroxy compound, phenol-containing compound, pigment, sterigmatocystin, depsipeptide,
lactone, mycotoxin, and organic heteropentacyclic compound biosynthesis as pathways
overrepresented in in the set of 1070 genes more expressed in WT (Supplementary Data 12).
Among the genes more expressed in WT, we find four genes in a nonribosomal peptide
synthetase cluster region predicted by AntiSMASH[54]; these four genes have homology to *sid2,*
sidF, sidH, and *sitT,* genes associated with siderophore biosynthesis and transport in

Aspergillus[62]. This gene cluster also includes genes with homology to *sidJ* and *mirB,* providing
further evidence that this region has a siderophore-related function. Five genes in a predicted
gene cluster encoding production of serinocyclin A and four genes in a predicted cluster
encoding production of nidulanin A are also more highly expressed in WT than in *ΔlaeA*. All
together, this confirms that deletion of *laeA* is likely to reduce production of fungal secondary
metabolites including siderophores. Among the 855 genes more expressed in *ΔlaeA,* GO term
enrichment identified amino acid and nucleoside metabolism, fungal cell wall, and ion transport
(Supplementary Data 12), highlighting a possible reorganization of *Penicillium* sp. str. #12
metabolism in the absence of *laeA*.
LC-MS comparison of the two extracts showed differential production of many
metabolites, 94 of which have a >10-fold change between the two (Supplementary Data 13). Of
these, 93 are less abundant in the *ΔlaeA* mutant, which is consistent with the loss of secondary
metabolite production in the *ΔlaeA* mutant (Figure 7b). Some of these metabolites matched with
known secondary metabolites and their analogs based on mass and fragmentation pattern.
Namely, atlantinone A and cyclopenol were found to be produced by the WT *Penicillium* sp. str.
#12 in >10-fold higher quantity than the *ΔlaeA* mutant, and pyripyropene O was putatively
identified and produced in >2-fold higher quantity. Cyclopenol is an alkaloid that is described as
a mycotoxin in the benzodiazepine class[70] and serves as an biosynthetic intermediate for
viridicatol[71], which has been described as having antibacterial activity against *Staphylococcus*
aureus[72]. Small amounts of viridicatol were indeed found in extracts with high quantities of
cyclopenol. Atlantinone A is a meroterpenoid that is derived from the same biosynthetic pathway
as other mixed polyketide-terpenoids such as andrastins and citrehybridones produced by various
Penicillium and *Aspergillus* species[73]. Both pyripyropene O and atlantinone A have tested
negative for antimicrobial activity when screened against a panel of bacteria, including an *E. coli*
strain[74,75]. The ecological role of these metabolites remains undetermined. In combination with
our RNA-Seq analysis, these data highlight an important diminution of secondary metabolite
production in the *ΔlaeA* strain.
We next performed RB-TnSeq experiments with the *Penicillium* sp. str. #12 Δ*laeA*
mutant to determine whether this single fungal mutation leads to changes in the genes needed for
bacterial fitness. While some interaction fitnesses are preserved between Δ*laeA* and WT, loss of
LaeA appears to significantly change the profile of interactions (Figure 7c). As LaeA is predicted
to impact siderophore production, we would expect that *E. coli* enterobactin uptake mutants
would not be as strongly impacted by the presence of the Δ*laeA* mutant. Although we see
positive interaction fitness of *fepA*, *fepB*, *fepC* and *fepG* genes when E. coli is grown with WT
Penicillium sp. str. #12 relative to *E. coli* growth alone, we do not see positive interaction fitness
of *fep* genes with Δ*laeA Penicillium* sp. str. #12 (Supplementary Data 14). Indeed, liquid CAS
assays demonstrate that this mutant produces less siderophores than WT on cheese media (Figure
7d). Together, these data support the hypothesis that loss of a fungal secondary metabolite
regulator corresponds to changes in bacterial interaction fitness effects.
Our RB-TnSeq results, combined with the antibiotic assay BCP, suggested that
antibiotic activity by *Penicillium* sp. str. #12 causes damage to *E. coli*'s cell envelope. As the
laeA mutant should have a decreased ability to produce antimicrobial compounds, we examined
whether there were changes in the fitness of genes related to responses to antibiotics. The *mdtK*
efflux pump gene, which was previously seen to have negative interaction fitness when grown
with WT *Penicillium* sp. str. #12, no longer has an interaction fitness with Δ*laeA*, suggesting that
the Δ*laeA* mutant has decreased antibiotic production. Although the exact nature of the

antimicrobials produced by this strain is unknown, these results are consistent with previous
studies, which show that LaeA regulates antibiotic production in other fungal species, and with
our metabolite analysis, which showed a decrease in production of many metabolites in Δ*laeA*.
Overall, these findings suggest that a single fungal gene can strongly impact the bacterial genes
needed for fitness and in particular point out that fungal specialized metabolite production may
play a large role in determination of bacterial fitness in bacterial-fungal interactions.

DISCUSSION

Fungi have been shown to strongly impact bacterial neighbors in diverse systems, from
soil to polymicrobial infections[77–85]. To characterize the effects of fungi from a simple
microbiome on bacteria, we used a high-throughput genetic screen, RB-TnSeq, to identify
bacterial genes relevant to fungal interactions. We observed a diversity of interactions, in terms
of direction (+/-), strength, and mechanism. These mechanisms include effects on important
cellular pathways such as biotin synthesis and antibiotic resistance. Our work demonstrates that
there is both a large diversity of bacterial-fungal interactions as well as conservation of key
interaction mechanisms across different fungi.
One of the strongest and most widespread bacterial-fungal interactions that we observed
suggests that fungal species can dramatically modulate access to iron through the provision of
fungal siderophores, such as ferrichrome and coprogen. Although it has long been known that
bacteria grown in isolation are able to uptake purified fungal siderophores[55,56], our results
demonstrate that this exchange takes place between bacteria and fungi growing in a biofilm and
that this exchange can have impacts on the competitive fitness of bacteria.
Hydroxamate siderophore receptors homologous to FhuE and FhuA are widespread in
Proteobacteria, suggesting that this interkingdom siderophore transfer may play an important role
in altering metal availability in diverse systems. *Pseudomonas aeruginosa* has previously been
shown to produce phenazine metabolites that led to upregulation of extracellular siderophore
production by *A. fumigatus*[86]. Fhu uptake systems have also been identified in the Gram-positive
bacterial pathogens *Streptococcus agalactiae* and *Listeria monocytogenes*, and growth of non-
siderophore producing mutants of *Streptomyces coelicolor* was restored by the presence of
siderophores from airborne contaminant fungi[87–90]. Additionally, hydroxamate siderophore
uptake systems have been shown to impact *Staphylococcus aureus* fitness in a murine infection
model[91]. Another study has shown that the presence of a siderophore-producing, cheese-
associated *Scopulariopsis* produced siderophores and caused downregulation of siderophore
production by *Staphylococcus equorum*[52]. *Bacteroides fragilis,* a human gut symbiont, is also
able to use ferrichrome to grow in iron-limiting conditions, and *fhu* genes are expressed by *E.*
coli in colonic mucus[92,93]. Some fungi consumed as a part of fermented foods have been shown
to survive digestive system transit, and fermented foods are known to contain fungal
siderophores[94,95], which could be a source of fungal siderophores in the gut in addition to
potential siderophore production by gut-resident species[94–96]. All together, these results suggest
that bacterial use of fungal siderophores may be a widespread mechanism of interkingdom
interaction. Considering the crucial role of iron acquisition and metabolism in microbial survival,
this conserved interaction mechanism could be a determinant in shaping environmental and host-
associated microbiomes.
This study provides new insight into the wide range of fungal impacts on bacteria that
can occur even in a simple microbiome. Our data also suggest that even among Proteobacteria,

fungal impacts may be diverse. Of the 692 *P. psychrophila* genes and 453 *E. coli* genes that had
an interaction fitness with at least one fungal partner, only 58 were homologous based on
BLAST (Supplementary Data 15). Further, many genes with interaction fitness do not have a
known function. We anticipated that by looking for fungal impacts on *E. coli,* we could leverage
the vastly superior genetic information available for this species. However, even in this well-
characterized organism, 43% of genes identified as having interaction fitness are annotated as
hypothetical or putative. For *P. psychrophila,* 29% of genes with interaction fitness are
hypothetical proteins. This highlights that many genes involved in interspecies interactions might
yet to be characterized, and that studying microbes in the context of their interactions with other
species, and not just in monoculture, provides an avenue for uncovering new areas of biology.
454
455

MATERIALS AND METHODS

Fungal Ribosomal RNA Sequencing

Genomic DNA was extracted with phenol-chloroform (pH 8) from cultures of the eight cheese
fungal species used in this study. For each extraction: 125 µL of 425–600 µm acid-washed beads
and 125 µL of 150–212 µm acid-washed beads were poured in a screw-capped 2 mL tube. 500
µL of 2X buffer B (200 mM NaCl, 20 mM EDTA) and 210 µL of SDS 20% were added to the
tube containing fungal material and 500 µL of Phenol:Chloroform (pH 8). Cells were lysed by
vortexing the tubes for 2 min at maximum speed. Aqueous and organic phases were separated by
centrifugation at 4°C, 8,000 RPM for 3 min and 450 µL of the aqueous phase (upper phase) were
recovered in a 1.5 mL Eppendorf tube. 45 µL of sodium acetate 3M and 450 µL of ice cold
isopropanol were added before incubating the tubes at −80°C for 10 min. The tubes were then
centrifuged for 5 min at 4°C at 13,000 RPM. The pellet was then washed in 750 µL of 70% ice
cold ethanol and re-suspended in 50 µL of DNase/RNase free water. Following DNA extraction,
LROR (ACCCGCTGAACTTAAGC) and LR6 (CGCCAGTTCTGCTTACC)[97] primers were
used to amplify the large subunit of the ribosomal RNA and for *Penicillium* species, Bt2a
(GGTAACCAAATCGGTGCTGCTTTC) Bt2b (ACCCTCAGTGTAGTGACCCTTGGC)[98]
primers were used to amplify the beta-tubulin gene. PCR was performed in a final volume of 50
µL: 25 µL of Q5 polymerase master mix (New England Biolabs), 2.5 µL of the forward primer
at 10 µM, 2.5 µL of the reverse primer at 10 µM, 100 ng of genomic DNA, and water using the
following PCR programs: LSU-(i) 98°C - 30 s, (ii) 35 cycles of: 98°C – 10 s; 52°C – 30 s; 72°C
– 1.5 min, (iii) 72°C – 5 min; Beta-tubulin-(i) 98°C - 30 s, (ii) 35 cycles of: 98°C – 10 s; 57°C –
30 s; 72°C – 1 min, (iii) 72°C – 5 min. PCR products were purified with the QIAquick PCR
purification kit (Qiagen) and sequenced using the forward and reverse primer by Eton Bioscience
Inc. (San Diego, USA). Consensus sequences from forward and reverse sequencing reactions
were aligned using Geneious version R9 9.1.8 (http://www.geneious.com). The MrBayes[36]
plugin for Geneious was used to build the phylogenetic tree with the following parameters:
Substitution model- JC69; Rate variation- gamma; Outgroup- *Fusarium* sp. str. 554A; Gamma
categories-4; Chain Length- 1100000; Subsampling freq- 200; Heated chains-4; Burn-in length-
100000; Heated chain temp- 0.2; Random seed-1160; Unconstrained branch lengths- 1, 0.1, 1, 1.
FigTree v1.4.4 was used for visualization (https://github.com/rambaut/figtree/releases).
486

Bacterial-Fungal Growth Assays

We aimed to inoculate 60,000 bacterial cells alone or with 6,000 fungal spores per well on 10%
CCA medium[38] adjusted to pH 7 in a 96-well plate. Each bacterial or bacterial-fungal assay was
done in triplicate. After 7 days of growth, the entire well was harvested and homogenized in
PBS1X-Tween 0.05% prior to dilution and plating on LB with 20 µg/ml cycloheximide (for
bacterial counts) or milk plate count agar with 50 µg/ml chloramphenicol (for fungal spore
counts). Counts were done at inoculation and after harvest. Final growth counts were then
compared in co-culture condition relative to growth alone to identify interaction impacts.
Significant growth impacts were determined based on Dunnett's test[99], p-value < 0.05. Plots
were made with R package ggplot2 3.2.1[100].

Microbial Culturing for LC-MS/MS extraction

All fungal cultures were grown on plate count agar milk salt (PCAMS; 1 g/L whole milk
powder, 1 g/L dextrose, 2.5 g/L yeast extract, 5 g/L tryptone, 10 g/L sodium chloride, 15 g/L
agar). Plates were kept at room temperature and spores were harvested at 7 days of growth (or
after sporulation was observed) for subsequent experiments. Spores harvested from fungi were
put into PBS and normalized to an OD_{600} of 0.1 for working stock.

Extraction of cultures

Three biological replicates of each condition were plated (distinct samples) and extracted from
solid agar. For extraction from solid agar plates, 5 µL of fungal working stock were spotted onto
10% CCA medium adjusted to pH 7. Following 7 days of growth, agar was removed from the
Petri dish and placed into 50 mL falcon tubes. Acetonitrile (10 mL) was added to each tube and
all were sonicated for 30 minutes. All falcon tubes were centrifuged and liquid was removed
from the solid agar pieces and transferred to 15 mL falcon tubes. The 15 mL falcon tubes
containing liquid were then centrifuged and liquid was again removed from any residual solid
debris and transferred to glass scintillation vials. These liquid extractions were then dried *in*
vacuo. Dried extracts were weighed and diluted with MeOH to obtain 1 mg/mL solutions which
were stored at -20°C until analysis via LC-MS/MS.

LC-MS/MS data collection

High resolution LC-MS and LC-MS/MS data were collected on a Bruker impact II qTOF in
positive mode with the detection window set from 50 to 1500 Da, on a 2.1x150mm C18 cortecs
UPLC column with a flow rate of 0.5mL/min for a gradient of 10-100% ACN with 0.1% formic
acid over 16 minutes. For each sample, 10 μL of a 1mg/mL solution was injected. The ESI
conditions were set with the capillary voltage at 4.5 kV. For MS^2, dynamic exclusion was used,
and the top nine precursor ions from each MS^1 scan were subjected to collision energies scaled
according to mass and charge state for a total of nine data dependent MS^2 events per MS^1. MS^2
data for pooled biological replicates is deposited under MassIVE accession number
MSV000085070. MS^1 and MS^2 data for *ΔlaeA* and WT *Penicillium* sp. str. #12 is deposited
under MassIVE accession number MSV000085054 and was collected under identical conditions
on a Bruker compact qTOF.

Molecular Networking

For all extractions, all precursor *m/z*'s that were found in solvent and agar controls (based on
both retention time and mass tolerance) were removed prior to input into GNPS using the
BLANKA algorithm.[101] A molecular network (
https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=464b331ef9d54de9957d23b4f9b9db14) was
created using the online workflow at GNPS. The data was filtered by removing all MS/MS peaks
within +/- 17 Da of the precursor *m/z*. MS/MS spectra were window filtered by choosing only
the top 6 peaks in the +/- 50Da window throughout the spectrum. The data was then clustered
with MS-Cluster with a parent mass tolerance of .02 Da and a MS/MS fragment ion tolerance of
.02 Da to create consensus spectra. Further, consensus spectra that contained less than 2 spectra
were discarded. A network was then created where edges were filtered to have a cosine score
above 0.7 and more than 6 matched peaks. Further edges between two nodes were kept in the
network if and only if each of the nodes appeared in each other's respective top 10 most similar
nodes. The spectra in the network were then searched against GNPS' spectral libraries. The
library spectra were filtered in the same manner as the input data. All matches kept between
network spectra and library spectra were required to have a score above 0.7 and at least 6
matched peaks. Solvent and agar control files were also loaded into the networks in order to
perform removal based on fragmentation patterns. All nodes with precursor masses less than
200Da were also removed. The extensive background and low *m/z* Da removal was done to more
accurately reflect the metabolomic profiles of fungal genera in an attempt to represent only true
metabolites. The DEREPLICATOR was used to annotate MS/MS spectra[102,103]. The molecular
networks were visualized using Cytoscape software[104].

RB-TnSeq Assays

All RB-TnSeq assays were performed on 10% CCA medium adjusted to pH 7. Prior to
inoculation, one aliquot of each library was thawed and inoculated into 25 mL of liquid LB-
kanamycin (50 µg/mL). Once the culture reached mid-log phase (OD = 0.6–0.8), 5 mL of that
pre-culture was pelleted and stored at −80°C for the T0 reference in the fitness analysis. The
remaining cells were used to inoculate the fitness assay conditions. For each RB-TnSeq fitness
assay, we aimed to inoculate 7,000,000 cells of the bacterial library (on average 50 cells per
insertion mutant). For fitness assays including a fungal partner, 700,000 fungal cells were
inoculated based on spore counts. Pre-cultured cells were washed in PBS1X-Tween 0.05%,
mixed with appropriate volumes of quantified fungal spore stocks, and then inoculated by
spreading evenly on a 100 mm petri dish containing 10% CCA medium, pH 7. For each
condition, assays were performed in triplicate (3 distinct samples). After seven days, each plate
was flooded with 1.5 mL of PBS1X-Tween 0.05% and cells were scraped off, taking care to not
disturb the CCA. The liquid was then transferred into a 1.5 mL microfuge tube and cells were
pelleted by centrifugation. After removing the supernatant, the cells were washed in 1 mL of
RNAprotect solution (Qiagen, Hilden, Germany), pelleted and stored at −80°C until gDNA
extraction. Genomic DNA was extracted with phenol-chloroform (pH 8) using the same protocol
used for fungal gDNA extraction described above. Samples were stored at −80°C until further
analysis.
After gDNA extraction, extracts containing Penicillium sp. str. #12 DNA were purified
using the OneStep PCR Inhibitor Removal Kit (Zymo Research, CA, USA). Then, the 98°C
BarSeq PCR as described in Wetmore *et al.*, 2015[24] was used to amplify only the barcoded

region of the transposons. PCR was performed in a final volume of 50 µL: 25 µL of Q5
polymerase master mix (New England Biolabs, MA, USA), 10 µL of GC enhancer buffer (New
England Biolabs), 2.5 µL of the common reverse primer (BarSeq_P1 – Wetmore et al., 2015) at
10 µM, 2.5 µL of a forward primer from the 96 forward primers (BarSeq_P2_ITXXX) at 10 µM
and either 200 ng of gDNA for alone conditions, or 2 µg of gDNA for fungal interaction
conditions. For *E. coli* analysis, we performed 84 PCRs (T0 sample and 28 harvest samples in
triplicate) involving 28 different multiplexing indexes. For *P. psychrophila* JB418 analysis, we
performed 84 PCRs (T0 sample and 28 harvest samples in triplicate) involving 28 different
multiplexing indexes. We used the following PCR program: (i) 98°C - 4 min, (ii) 30 cycles of:
98°C – 30 s; 55°C – 30 s; 72°C – 30 s, (iii) 72°C – 5 min. After the PCR, for both *E. coli* and *P.*
psychrophila, 10 µL of each of the PCR products were pooled together to create the BarSeq
sequencing library and 200 µL of the pooled library were purified using the MinElute
purification kit (Qiagen, Germany). The final elution of the BarSeq library was performed in 30
µL in DNase and RNase free water. The BarSeq libraries were then quantified using Qubit
dsDNA HS assay kit (Invitrogen, CA, USA) and sequenced on HiSeq4000 (75 bp, single-end
reads), by the IGM Genomics Center at the University of California San Diego. The sequencing
depth for each condition varied between 6.1 and 11.7 million reads for *E. coli* and 5.8 and 13.3
million reads for *P. psychrophila.*

RB-TnSeq Data Processing

Custom R scripts were used to determine the average fitness scores for each gene across three
RB-TnSeq assay replicates. These scripts are available at https://github.com/DuttonLab/RB-
TnSeq-Microbial-interactions. The Readme document provides an in-depth explanation of all
data processing steps performed in these scripts. In brief, insertion mutants that did not have a
sufficient T0 count (3) in each condition or that are not centrally inserted (10-90% of gene) were
removed from analysis. Counts determined by Wetmore *et al.*[24] scripts were then normalized to a
reference gene (*allB* in *E. coli* and closest protein BLAST match Ga0212129_11710 in *P.*
psychrophila) to be able to compare across conditions and to account for differences in
sequencing depth. These genes do not have a strong fitness effect in any condition based on
former fitness determination developed by Wetmore *et al.*, 2015[24]. Un-normalized strain fitness
was then calculated per insertion mutant as the log2 of the ratio of the corrected counts in the
condition and the corrected counts in the T0 sample. Un-normalized gene fitness and variance
was next calculated by averaging insertion mutants within a gene. These values were then
normalized based on the position of the gene along the chromosome, as well as by the mean of
the data distribution. These steps were all done on individual replicates. Next, the average gene
fitness and associated variance were calculated using the weighted average of the fitness values
across the three different replicates. Then, for each condition, fitness values are corrected by the
mean of the replicate means (the replicate means being the mean values used to center fitness
value within a replicate). Finally, based on the assumption that most genes should have no fitness
effect, we corrected gene fitness values in each condition by the mode (peak of the density
distribution). Final fitness values were then compared for fungal interaction conditions compared
to bacterial alone conditions and comparisons associated with a p-value lower than 5% were
considered significant interaction fitness (alpha parameter=0.05 in
2conditions_FitnessComparison.R code). Networks of fitness values were visualized in
Cytoscape v. 3.5.1[104] and PCA plots were made with R package ggplot2 3.2.1[100] and ggfortify

0.4.7[105]. COG category mapping of *E. coli* and *P. psychrophila* protein sequences was done by
eggNOG-mapper v2[106] and visualized with R package ggplot2 3.2.1[100].

Bacterial Cytological Profiling

Approximately 7,000,000 WT *E. coli* K-12 strain BW25113 or Keio collection *mdtK* mutant
cells[107] were inoculated alone or co-inoculated with 700,000 *Penicillium* sp. str. #12 or
Penicillium sp. str. SAM3 spores on 10% CCA pH 7. After 7 days of growth, 1 mL of T-Base
buffer was added to the surface of the biofilms, and biofilms were scraped into the buffer. For
co-culture conditions, the sample was filtered through a 0.5 μm filter to specifically remove
fungal material. 2 μL of concentrated dye mix (1 μL 1 mg/mL FM4-64, 1 μL 2 mg/mL DAPI in
48 μL T-Base) were added to 20 μL of filtrate. Dye-filtrate mix was spotted onto agarose-LB
pads (1% Agarose, 20% LB liquid medium, 80% ddH$_2$O) and imaged by fluorescence and phase
contrast. Control compound references on CCA medium were obtained by spotting 30 μL of 5x,
10x, 25x, and 100x MIC dilutions of antibiotics onto quadrants on CCA medium pH 7 plates,
allowing to dry, and spread-plating 200 μL of log-phase (OD 0.1) *E. coli* cultures. After two
days of growth, cells near the edge of the zone of inhibition on appropriate dilution spots were
resuspended in 10 μL of prediluted dye mix (1 μL 1 mg/mL FM4-64, 1 μL 2 mg/mL DAPI in
998 μL T-Base) and spotted onto agarose-LB pads and imaged as described above. Resulting
images were deconvolved using Deltavision SoftWorx software (Applied Precision, Inc., WA,
USA), analyzed using Fiji[108], and assembled in Adobe Photoshop (Adobe, CA, USA).

CCA medium biotin quantification

Biotin quantification of CCA medium was performed on three replicate samples by Creative
Proteomics (NY, USA) as follows: 100 mg of each sample was homogenized in water (10
μL/mg) for 1 min three times with the aid of 5-mm metal balls on a MM400 mill mixer.
Methanol at 10 μL/mg was then added. Water-soluble vitamins were extracted by vortex-mixing
for 2 min and sonication in a water batch for 5 min. After centrifugation, the clear supernatants
were cleaned up by solid-phase extraction on a Strata-X (60 mg/mL) cartridge. The eluted
fractions containing water-soluble vitamins were collected, pooled and then dried under a gentle
nitrogen gas flow in a nitrogen evaporator. The residues were dissolved in 200 μL of 10%
methanol. Twenty microliter aliquots were injected to run UPLC-MRM/MS with the use of a
C18 UPLC column and with (+) ion detection and (-) ion detection, respectively. Calibration
curves were prepared by injection of serially-diluted mixed standard solutions of water-soluble
vitamins. Concentrations of detected vitamins were calculated by interpolating the linear
calibration curves.

Δ*fep* mutant competitive growth assays

Approximately 600,000 WT and Δ*fepA,* Δ*fepC,* or Δ*fepG* Keio collection[107] mutant cells were
inoculated at a 1:1 ratio either alone or co-inoculated with approximately 60,000 *Penicillium* sp.
str. #12 or *Scopulariopsis* sp. str. JB370 spores on 10% CCA pH 7 in a 96-well plate in four
replicates each (4 distinct samples). After seven days of growth, the entire well was harvested
and homogenized in PBS1X-Tween 0.05% prior to dilution and plating on LB with 20 μg/ml
cycloheximide (total bacterial counts) or with 20 μg/ml cycloheximide and 50 μg/ml kanamycin
(bacterial mutant counts). Final growth counts were then compared in co-culture condition

relative to growth alone to identify interaction impacts. Significant growth impacts were

determined by significantly different growth in the presence of a fungus relative to growth alone

based on a two-sided two-sample equal variance t-test p-value < 0.05. Plots made with R

package ggplot2 3.2.1[100].

RNA-Seq and differential expression analysis of *E. coli* with *Penicillium* sp. str. #12

Approximately 7,000,000 *E. coli* cells were inoculated in triplicate (3 distinct samples) either

alone or with approximately 700,000 *Penicillium* sp. str. #12 spores on 10% CCA pH 7. After 3

days, the biofilms were harvested for RNA extraction and washed with 1 mL of RNAprotect.

RNA was extracted by a phenol-chloroform extraction (pH 8) using the same extraction protocol

as for gDNA extraction. Extractions were then purified with the OneStep PCR Inhibitor

Removal Kit (Zymo Research, CA, USA).

Sequencing libraries were prepared as follows. RNA samples were treated with DNase

using the 'Rigorous DNase treatment' for the Turbo DNA-free kit (AMBION, Life

Technologies, Waltham, MA, USA) and RNA concentration was measured by nucleic acid

quantification in Epoch Microplate Spectrophotometer (BioTek, Winooski, VT, USA). Transfer

RNAs and 5S RNA were then removed using the MEGAclear Kit Purification for Large Scale

Transcription Reactions (AMBION, Life Technologies) following manufacturer instructions.

Absence of tRNA and 5S RNA was verified by running 100 ng of RNA on a 1.5% agarose gel

and RNA concentration was quantified by nucleic acid quantification in Epoch Microplate

Spectrophotometer. Also, presence of gDNA was assessed by PCR using universal bacterial 16S

PCR primers (Forward primer: AGAGTTTGATCCTGGCTCAG, Reverse Primer:

GGTTACCTTGTTACGACTT). The PCR was performed in a final volume of 20 µL: 10 µL of

Q5 polymerase master mix (New England Biolabs), 0.5 µL of forward primer 10 uM, 0.5 µL of

reverse primer 10 uM and 5 µL of non-diluted RNA. PCR products were run on a 1.7% agarose

gel and if gDNA was amplified, another DNase treatment was performed as well as a new

verification of absence of gDNA.

Ribosomal RNA depletion was performed using the RiboMinus Transcriptome Isolation

Kit (Yeast and Bacteria) for the *E. coli* alone samples and using both the RiboMinus

Transcriptome Isolation Kit (Yeast and Bacteria) and the RiboMinus Eukaryote Kit v2 for the

mixed *E. coli/Penicillium* sp. str. #12 samples (ThermoFisher Scientific). For the *E. coli* alone

samples, each sample was divided into two for treatment and then repooled for RNA recovery

with ethanol precipitation. For the *E. coli/Penicillium* sp. str. #12 samples, an equal volume of

the eukaryotic probe and RiboMinus Bacterial Probe Mix were used to deplete both bacterial and

fungal ribosomal RNA and RNA was recovered with ethanol precipitation. Concentrations after

ribosomal RNA depletion were measured using Qubit RNA HS Assay Kits (Invitrogen). The

RNA-Seq library was produced using the NEBNext Ultra RNA Library Prep Kit for Illumina for

purified mRNA or ribosome-depleted RNA. We prepared a library with a fragment size of 300

nucleotides and used the 10 µM NEBNext Multiplex Oligos for Illumina (Set 1, NEB #E7335)

lot 0091412 and the NEBNext multiplex Oligos for Illumina (Set 2, NEB #E7500) lot 0071412.

We performed PCR product purification with 0.8X Agencourt AMPure XP Beads. Library

samples were quantified with Qubit DNA HS Assay Kits before the quality and fragment size

were validated by TapeStation (HiSensD1000 ScreenTape). Library samples were pooled at a

concentration of 15 nM each and were sequenced on HiSeq4000 (50 bp, single-end). TapeStation

assays and sequencing were performed by the IGM Genomics Center at the University of
California San Diego.
Following sequencing, reads were mapped to the concatenated genome of *E. coli* K12
BW25113[109] and *Penicillium* sp. str. #12 using Geneious version R9 9.1.8
(http://www.geneious.com). Only the reads that uniquely mapped to a single location on the *E.*
coli genome section were kept. *E. coli* expression analysis was performed using the following R
packages: Rsamtools (R package version 2.0.3), GenomeInfoDb (R package version 1.20.0),
GenomicFeatures[110] (R package version 1.36.4), GenomicAlignments[110] (R package version
1.20.1), GenomicRanges[110] (R package version 1.36.1) and DESeq2[111] (R package version
1.20.1). We followed the workflow described by Love *et al*. and performed the differential
expression analysis using the package DESeq2. Differentially expressed genes between two
conditions were selected with an adjusted p-value lower than 5% (Benjamini-Hochberg
correction for multiple testing) and an absolute log2 of fold change equal to or greater than 1.5.

Construction of *E. coli* mutants and visual interaction assays

Visual assays for hydroxamate siderophore stimulation:

Antibiotic assay discs (Whatman) were placed on CCA medium pH 7 with .005% tetrazolium
chloride (an indicator of cellular respiration) and 20 µL of water, 10 µM coprogen or ferrichrome
(EMC Microcollections GmbH) solutions (in water) were slowly pipetted onto the disc and
allowed to absorb. 2.5 µL of 37°C overnight LB cultures of *E. coli* K12 BW25113 WT, Δ*fepA,*
Δ*fepC,* Δ*fhuE,* Δ*fhuA,* Δ*fepA*Δ*fhuE,* Δ*fepC*Δ*fhuE,* or Δ*fepA*Δ*fhuA* mutants[107] were spotted next
to the discs. Double mutants were constructed as described below. Plates were left at room
temperature until red color developed.

Visual assays for fungal stimulation of bacterial mutants:

Fungal spores were inoculated on CCA pH 7 with .005% tetrazolium chloride. After fungal
growth at room temperature (cheese fungal isolates) or 30°C (*A. fumigatus* and *M.*
pachydermatis), 2.5 µL of *E. coli* 37°C overnight LB cultures were spotted at increasing
distances from the fungal front. Plates were left at room temperature until red color developed. *A.*
fumigatus isolate AF293 was received from Nancy Keller, University of Wisconsin-Madison. *M.*
pachydermatis was originally isolated from the ear of a dog in Sweden (ATCC14522).

Creation of Δ*fepA*Δ*fhuE* and Δ*fepC*Δ*fhuE*:

Chemically competent cells for Δ*fepA* or Δ*fepC* mutants were created. An overnight culture of
Δ*fepA* or Δ*fepC* mutants was diluted 1:100 and grown at 37°C until OD 0.4-0.6. The culture was
placed on ice for 20 minutes and then centrifuged at 4°C for 10 minutes at 6000 rpm to collect
the cells. Supernatant was removed and cells were resuspended in half the previous volume of
pre-cooled 0.1M CaCl$_2$. After being left on ice for 30 minutes, centrifugation was repeated and
supernatant was removed before resuspension in a quarter of the original volume of pre-cooled
0.1M CaCl$_2$/15% glycerol. Cells were aliquoted and stored at -80°C until transformation. These
cells were transformed with the pKD46 plasmid [112], recovered at 30°C and plated on LB plates
with 100 µg/mL Ampicillin and grown at 30°C. Overnight cultures were started from individual
colonies for creation of electrocompetent cells. Overnight cultures of Δ*fepC*-pkD46 or Δ*fepA*-
pkD46 were diluted 1:100 in fresh LB- 100 µg/mL Ampicillin and grown at 30°C until an OD of
0.1. 20 µL of fresh 1 M L-arabinose were added and growth was continued at 30°C until OD 0.4-
0.6. Cells were then chilled on ice for 15 minutes and then centrifuged for ten minutes at 4000

rcf 4°C. Cells were resuspended in 1 mL of ice water and centrifuged for ten minutes at 4000 rcf

at 4°C. Cells were resuspended in 0.5 mL of ice water and centrifuged for ten minutes at 4000 rcf

4°C. Cells were resuspended in 50 µL of ice water and kept on ice until transformation. The

chloramphenicol resistance cassette was amplified from the pKD3 plasmid [112] using the custom

primers: FhuEcatF

(CAGATGGCTGCCTTTTTTACAGGTGTTATTCAGAATTGATACGTGCCGGTAATGGCG

CGCCTTACGCCCC) and FhuEcatR

(CCTCCTCCGGCATGAGCCTGACGACAACATAAACCAAGAGATTTCAAATGCTGGGC

CAACTTTTGGCGAA) and the following PCR conditions (i) 98°C - 30 sec, (ii) 30 cycles of:

98°C – 10 s; 70°C – 20 s; 72°C – 30 s, (iii) 72°C – 5 min. Amplification was performed on 4 ng

of pKD3 plasmid using Q5 High-Fidelity 2X Master Mix (New England Biolabs). The PCR

product was digested for 1 hour with the restriction enzymes DpnI and ClaI at 37°C and then the

PCR product was run on a 1% agarose gel. The PCR product was extracted using the QIAquick

Gel Extraction Kit (Qiagen) and then dialyzed for 4 hours with TE buffer. 1.5 µL of dialyzed

PCR product was used to transform the electrocompetent Δ*fepC*-pkD46 or Δ*fepA*-pkD46 cells.

After 2 hours of recovery in SOC medium with 1mM arabinose at 37°C, the transformation was

plated on LB with 50 mg/mL kanamycin and chloramphenicol. Transformants were confirmed

to be Δ*fhuE* with Eton Bioscience Inc. sequencing of the chloramphenicol cassette.

Creation of Δ*fepA*Δ*fhuA*:

Creation was done as with Δ*fepA*Δ*fhuE*, except that the chloramphenicol resistance cassette was

amplified from pKD3[112] using FhuAcatF (ATCATTCTCGTTTACGTTATCATTCACTTT

ACATCAGAGATATACCAATGAATGGCGCGCCTTACGCCCCAATGGCGCGCCTTACG

CCCC) and FhuAcatR

(GCACGGAAATCCGTGCCCCAAAAGAGAAATTAGAAACGGAAGGTTGCGGTCTGGG

CCAACTTTTGGCGAACTGGGCCAACTTTTGGCGAA) custom primers.

Penicillium sp. str. #12 genome sequencing, assembly, and annotation

Genomic DNA was extracted from *Penicillium* sp. str. #12 using the genomic DNA extraction

protocol described above. High molecular weight DNA (average 16 Kb) was sequenced on the

Oxford Nanopore MinION with a R.9.5 flow cell using $1D^2$ sequencing adaptors from kit SQK-

LSK308 (Oxford Nanopore Technologies, Oxford, United Kingdom). Raw data was basecalled

using guppy 3.3.0 (Oxford Nanopore Technologies, Oxford, United Kingdom)(guppy_basecaller

-config dna_r9.5_450bps.cfg --fast5_out) for 1D basecalls and these were used to also obtain

higher accuracy 1D^2 basecalls (guppy_basecaller_1d2 -i 1Dbasecall/workspace/ --config

dba_r9.5_450bps_1d2_raw.cfg --index_file 1Dbasecall/sequencing_summary.txt) These reads

were assembled by canu 1.8[113] and polished by racon 1.4.3[114] four times and by pilon 1.23[115]

once. The final assembly is 38 Mbp and consists of 52 contigs.

Penicillium sp. str. #12 genome annotations were obtained by combining genomic and

transcriptomic information from RNA-Seq. To obtain the gene expression profile of *Penicillium*

sp. str. #12, approximately 700,000 WT *Penicillium* sp. str. #12 spores were inoculated in

triplicate on 10% CCA pH 7. After 3 days, the biofilms were harvested for RNA extraction and

washed with 1mL of RNAprotect. RNA was extracted and RNA-Seq libraries were prepared as

described above with the following modification: Ribosomal RNA depletion was performed

using the RiboMinus Eukaryote Kit v1 and RNA was recovered with ethanol precipitation. After

sequencing, the RNA-Seq reads from these *Penicillium* sp. str. #12 alone cultures were
concatenated with RNA-Seq reads from the previously described *E. coli/Penicillium* sp. str. #12
co-culture conditions that uniquely mapped to a single location on the *Penicillium* sp. str. #12
genome. The full set of transcriptomic reads were then used as input into the FunGAP annotation
pipeline and 77 million of these reads mapped[116]. This pipeline predicted 13261 protein-coding
genes in the *Penicillium* sp. str. #12 genome. Interproscan[117] was used within the FunGAP
pipeline for function prediction of genes. This Whole Genome Shotgun project has been
deposited at DDBJ/ENA/GenBank under the accession JAASRZ000000000. The version
described in this paper is version JAASRZ010000000.

Creation and confirmation of *laeA* deletion in *Penicillium* sp. str. #12

Deletion cassette design strategy: In order to knockout *laeA* in *Penicillium* sp. str. #12, the
isolate was first screened for hygromycin and phleomycin resistance. *Penicillium* sp. str. #12
showed a confirmed sensitivity to both antibiotics. A three round PCR deletion strategy was used
to replace the *laeA* ORF with the *hph* gene, whose expression confers selection on
hygromycin[118]. The schematic representation of the *laeA* gene replacement with the hph gene is
depicted in Supplementary Figure 8. The deletion cassette (5'flank- hph- 3'flank) was
constructed using three sequential PCR reactions. In the first PCR round, about 1 Kb genomic
sequence flanking either the 5' or 3' end of the *laeA* ORF were amplified using the primer sets
P12_KOlaeA_5' F (CTCCGTTGGGCCCTCAC) and 5'R
(GCAATTTAACTGTGATAAACTACCGCATTAAAGCTGTTGATATCGGCAATCAATCA
ATG) or P12_KOlaeA_3'F
(GGTGGGCCTTGACATGTGCAGCCGGTGGAGCGGCGCCTGGTGAATCCTACCCACAT
GG) and 3'R (CGTTGGGAGGAAAAGCTTCTGCG) respectively. The *hph* gene was amplified
from plasmid pUCH2-8 using primers hph_F (AGCTTTAATGCGGTAGTTTATCACAG) and
hph_R (CTCCACCGGCTGCACATGTC). A second PCR reaction was performed to assemble
by homologous recombination the three individual fragments from the first round PCR. The
deletion cassettes were finally amplified using the nested primer set, P12_KOlaeA_NestedF
(CAGACGGTCCGCATCCCG) and P12_KOlaeA_NestedR
(GGTCCAGGTGCAGTAGTACTG).

Fungal transformation: To generate the deletion strains, a protoplast-mediated transformation
protocol was employed. Briefly, 109 fresh spores were cultured in 500 mL of liquid minimal
medium (LMM) for 12 h under 25°C and 280 rpm. Newly born hyphae were harvested by
centrifugation at 8000 rpm for 15 min and hydrolyzed in a mixture of 30 mg Lysing Enzyme
from Trichoderma (Sigma-Aldrich) and 20 mg Yatalase (Fisher Scientific) in 10 mL of Osmotic
Medium (1.2 M $MgSO_4$, 10 mM NaPB, pH 5.8). The quality of protoplast was monitored under
the microscope after four hours of shaking under 28 °C and 80 rpm. The protoplast mixture was
later overlaid with 10 mL of trapping buffer (0.6 M sorbitol, 100 mM Tris-HCl pH 7.0) and
centrifuged for 15 min under 4°C and 5000 rpm. Protoplasts were collected from the interface,
overlaid with an equal volume of STC (1.2 M sorbitol, 10 mM Tris-HCL pH 7.5,10 mM $CaCl_2$)
and decanted by centrifugation at 6000 rpm for 8 min. The protoplast pellet was resuspended in
500 µL STC and used for transformation. After 5 days of incubation at 25 °C, colonies grown on
stabilized minimal medium (SMM) plates supplemented with hygromycin were subjected to a

second round of selection on hygromycin plates. In total, 25 hygromycin-resistant transformants
were isolated after a rapid screening procedure on SMM supplemented with hygromycin. Single-
spored transformants were later tested for proper homologous recombination at the ORF locus by
PCR and Southern blot analysis.
833
Gene-deletion strain confirmation: The correct replacement of the *laeA* with the *hph* gene was
first verified by PCR analysis of genomic DNA derived from the transformant strains using
primer set P12_laeA_F (CACAATGGCTGAACACTCTCGG) and P12_laeA_R
(GGGATATGGAGCATCGAAGTTGC) that amplify the *laeA* ORF. About 12% (3/25) of the
monoconidial lines generated from primary transformants of *Penicillium* sp. str. #12 were PCR-
positive for the absence of the *laeA* ORF. The positive deletion strains were further checked for a
single insertion of the deletion cassette by Southern blot analysis and revealed single-site
integration of the deletion cassette in one transformant (Supplementary Figure 8). Probes
corresponding to the 5' and 3' flanks of the *laeA* gene in each strain were labeled using [α32P]
dCTP (PerkinElmer, USA) following the manufacturer's instructions.
844
RNA-Seq analysis of WT and Δ*laeA Penicillium* sp. str. #12

To characterize the effect of the *laeA* deletion on the *Penicillium* sp. str. #12 gene expression
profile, we performed RNA-Seq analysis for Δ*laeA Penicillium* sp. str. #12. As for WT
Penicillium sp. str. #12, 700,000 Δ*laeA Penicillium* sp. str. #12 spores were inoculated in
triplicate (3 distinct samples) on 10% CCA pH 7 and biofilms were harvested after 3 days.
Harvest, RNA extraction and library preparation were performed identically to WT *Penicillium*
sp. str. #12. Then, *Penicillium* sp. str. #12 and Δ*laeA* differential expression analysis was
performed as described for *E. coli/Penicillium* sp. str. #12 above. To look for enrichment of
functions in the set of differentially expressed genes, we input the protein sequences of the genes
into the gene-list enrichment function of KOBAS 3.0[119]. Sequences were searched against the
Gene Ontology (GO) database[120,121] using *A. fumigatus* as a reference for GO assignment before
conducting a hypergeometric test with Benjamini-Hochberg correction. Functions with a
corrected p-value <.05 were considered enriched.
858
Data Availability
Sequence data that support the findings of this study (RB-TnSeq, RNA-Seq) have been deposited
in the NCBI SRA database with BioProject PRJNA624168. Mass spectrometry data is available
in the MassIVE database under accession numbers MSV000085070 and MSV000085054. The
GNPS molecular network is available at
https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=464b331ef9d54de9957d23b4f9b9db14. The
Whole Genome Shotgun project for *Penicillium* sp. str. #12 has been deposited at
DDBJ/ENA/GenBank under the accession JAASRZ000000000 in BioProject PRJNA612335.
The version described in this paper is version JAASRZ010000000. In addition to these sources,
source data used to create figures 2,3,4, 6, and 7 is available in the Supplementary Data provided
with the paper.
870
871　Code Availability
872　The R scripts developed for processing RB-TnSeq data described in this manuscript are available
873　at https://github.com/DuttonLab/RB-TnSeq-Microbial-interactions along with usage instructions.

874   The perl scripts needed for initial processing of RB-TnSeq data published in Wetmore *et al.*
875   2015[24] are available at https://bitbucket.org/berkeleylab/feba/src/master/.
876
Availability of biological materials
All unique materials, including described fungal strains isolated from cheese, the *P. psychrophila*
JB418 strain and RB-TnSeq library, *laeA Penicillium* sp. str. #12 deletion mutant, and *E. coli*
siderophore uptake double mutants, are readily available from the authors upon request. The *E.*
coli RB-TnSeq library and Keio strains can be requested from the groups that created these
resources (PMID references provided). *Penicillium* sp. str. SAM3 is commercially available from
Danisco.
884

Resource Sources			
Resource	**Source or reference**	**Identifiers**	**Additional information**
Keio collection, strain background *Escherichia coli* K12	PMID: 16738554	CGSC, RRID:SCR_002303	Collection of 3,818 *E. coli* knockout strains
Keio_ML9 library, strain background *Escherichia coli* K12	PMID: 25968644		RB-TnSeq library of *E. coli* K12 BW25113 (152,018 pooled insertion mutants)
JB418_ECP1 library, strain background *Pseudomonas psychrophila* JB418	PMID: 30211673		RB-TnSeq library generated in the *P. psychrophila* JB418 strain isolated from cheese (272,329 pooled insertion mutants)
Keio ME9062, strain background *Escherichia coli* K12	PMID: 16738554	CGSC#: 7636	Parent strain of the Keio collection mutants
Penicillium sp. str. SAM3	Danisco - CHOOZIT	PC SAM 3 LYO 10D	Industrial starter for cheese production
Pseudomonas psychrophila JB418	PMID: 30211673		Strain isolated from Robiola due latti cheese
Candida sp. str. 135E	This paper		Strain isolated from cheese
Debaryomyces sp. str. 135B	This paper		Strain isolated from cheese
Penicillium sp. str. #12	This paper		Strain isolated from cheese
Penicillium sp. str. RS17	This paper		Strain isolated from cheese
Scopulariopsis sp. str. JB370	This paper		Strain isolated from cheese
Scopulariopsis sp. str. 165-5	This paper		Strain isolated from cheese
Fusarium sp. str. 554A	This paper		Strain isolated from cheese
Aspergillus fumigatus str. AF293	Nancy Keller, University of Wisconsin-Madison		
Malassezia pachydermatis str. CBS 1879	ATCC14522		

ACKNOWLEDGMENTS

The authors would like to thank: the Arkin lab and the Deutschbauer lab at UC-Berkeley for the
E. coli Keio_ML9 library, Kristen Jepsen at the IGM Genomics Center at the University of
California San Diego for assistance with sequencing, Dr. Sergey Kryazhimskiy (UCSD) for his
input on RB-TnSeq data processing, Cong Dinh (UCSD) for assistance with fungal genome
assembly, William Bushnell for assistance with fitness validation experiments, Dr. Sinem
Beyhan for advice on fungal genome annotation, Lisa Marotz (UCSD) for assistance with non-
cheese yeasts, and members of the Dutton lab, especially Brooke Anderson, for constructive
comments on the manuscript.
896
897 **FUNDING SOURCES**
898
This work was supported by National Institutes of Health grants T32-AT007533 (J.C.L.), F31-
AT010418 (J.C.L.), National Institutes of Health grant R01-AI117712 (R.B.L), National Science
Foundation grant MCB-1817955 (L.M.S.), NSF grant MCB-1817887 (R.J.D. and L.M.S.), the
UCSD Center for Microbiome Innovation (E.C.P.), the UCSD Ruth Stern Award (E.C.P.), NIH
Institutional Training grant 5 T32 GM 7240-40 (E.C.P.), and National Institutes of Health grant
R01GM112739-01 (N.P.K.).

REFERENCES

1. Laforest-Lapointe, I. & Arrieta, M. C. Microbial Eukaryotes: a Missing Link in Gut

Microbiome Studies. *mSystems* **3**, (2018).

2. Forbes, J. D., Bernstein, C. N., Tremlett, H., Van Domselaar, G. & Knox, N. C. A Fungal

World: Could the Gut Mycobiome Be Involved in Neurological Disease? *Front. Microbiol.*

9, 3249 (2018).

3. Huseyin, C. E., O'Toole, P. W., Cotter, P. D. & Scanlan, P. D. Forgotten fungi-the gut

mycobiome in human health and disease. *FEMS Microbiol. Rev.* **41**, 479–511 (2017).

4. Bergelson, J., Mittelstrass, J. & Horton, M. W. Characterizing both bacteria and fungi

improves understanding of the Arabidopsis root microbiome. *Sci. Rep.* **9**, 24 (2019).

5. Huffnagle, G. B. & Noverr, M. C. The emerging world of the fungal microbiome. *Trends*

Microbiol. **21**, 334–341 (2013).

6. Soverini, M. *et al.* HumanMycobiomeScan: a new bioinformatics tool for the

characterization of the fungal fraction in metagenomic samples. *BMC Genomics* **20**, 496

(2019).

7. Bradford, L. L. & Ravel, J. The vaginal mycobiome: A contemporary perspective on fungi

in women's health and diseases. *Virulence* **8**, 342–351 (2017).

8. Marsh, A. J., O'Sullivan, O., Hill, C., Ross, R. P. & Cotter, P. D. Sequence-based analysis

of the bacterial and fungal compositions of multiple kombucha (tea fungus) samples. *Food*

Microbiol. **38**, 171–178 (2014).

9. De Filippis, F., Laiola, M., Blaiotta, G. & Ercolini, D. Different Amplicon Targets for

Sequencing-Based Studies of Fungal Diversity. *Appl. Environ. Microbiol.* **83**, (2017).

10. Fan, D. *et al.* Activation of HIF-1α and LL-37 by commensal bacteria inhibits Candida

albicans colonization. *Nat. Med.* **21**, 808–814 (2015).

11. Jiang, T. T. *et al.* Commensal Fungi Recapitulate the Protective Benefits of Intestinal

Bacteria. *Cell Host Microbe* **22**, 809–816.e4 (2017).

12. Wheeler, M. L. *et al.* Immunological Consequences of Intestinal Fungal Dysbiosis. *Cell*

Host Microbe **19**, 865–873 (2016).

13. Peleg, A. Y., Hogan, D. A. & Mylonakis, E. Medically important bacterial–fungal

interactions. *Nat. Rev. Microbiol.* **8**, 340–349 (2010).

14. Mason, K. L. *et al.* Interplay between the gastric bacterial microbiota and Candida albicans

during postantibiotic recolonization and gastritis. *Infect. Immun.* **80**, 150–158 (2012).

15. Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E. E. & van der Heijden, M. G. A. Fungal-

bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat.*

Commun. **10**, 4841 (2019).

16. Durán, P. *et al.* Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival.

Cell **175**, 973–983.e14 (2018).

17. Tourneroche, A. *et al.* Bacterial–Fungal Interactions in the Kelp Endomicrobiota Drive

Autoinducer-2 Quorum Sensing. *Frontiers in Microbiology* vol. 10 (2019).

18. Wolfe, B. E., Button, J. E., Santarelli, M. & Dutton, R. J. Cheese Rind Communities Provide

Tractable Systems for In Situ and In Vitro Studies of Microbial Diversity. *Cell* **158**, 422–

433 (2014).

19. Wolfe, B. E. & Dutton, R. J. Fermented foods as experimentally tractable microbial

ecosystems. *Cell* **161**, 49–55 (2015).

20. May, A. *et al.* Kombucha: a novel model system for cooperation and conflict in a complex

multi-species microbial ecosystem. *PeerJ* **7**, e7565 (2019).

21. Morin, M., Pierce, E. C. & Dutton, R. J. Changes in the genetic requirements for microbial

interactions with increasing community complexity. *Elife* **7**, (2018).

22. Zhang, Y., Kastman, E. K., Guasto, J. S. & Wolfe, B. E. Fungal networks shape dynamics of

bacterial dispersal and community assembly in cheese rind microbiomes. *Nat. Commun.* **9**,

336 (2018).

23. Cleary, J. L., Kolachina, S., Wolfe, B. E. & Sanchez, L. M. Coproporphyrin III Produced by

the Bacterium Glutamicibacter arilaitensis Binds Zinc and Is Upregulated by Fungi in

Cheese Rinds. *mSystems* **3**, (2018).

24. Wetmore, K. M. *et al.* Rapid Quantification of Mutant Fitness in Diverse Bacteria by

Sequencing Randomly Bar-Coded Transposons. *MBio* **6**, (2015).

25. Rubin, B. E. *et al.* The essential gene set of a photosynthetic organism. *Proceedings of the*

National Academy of Sciences **112**, E6634–E6643 (2015).

26. Price, M. N. *et al.* Mutant phenotypes for thousands of bacterial genes of unknown function.

Nature **557**, 503–509 (2018).

27. Cole, B. J. *et al.* Genome-wide identification of bacterial plant colonization genes. *PLoS*

Biol. **15**, e2002860 (2017).

28. Hallen-Adams, H. E. & Suhr, M. J. Fungi in the healthy human gastrointestinal tract.

Virulence **8**, 352–358 (2017).

29. Frąc, M., Hannula, S. E., Bełka, M. & Jędryczka, M. Fungal Biodiversity and Their Role in

Soil Health. *Front. Microbiol.* **9**, 707 (2018).

30. Richards, T. A., Jones, M. D. M., Leonard, G. & Bass, D. Marine fungi: their ecology and

molecular diversity. *Ann. Rev. Mar. Sci.* **4**, 495–522 (2012).

31. Frey-Klett, P. *et al.* Ectomycorrhizal symbiosis affects functional diversity of rhizosphere

fluorescent pseudomonads. *New Phytol.* **165**, 317–328 (2005).

32. Lopez-Medina, E. *et al.* Candida albicans Inhibits Pseudomonas aeruginosa Virulence

through Suppression of Pyochelin and Pyoverdine Biosynthesis. *PLoS Pathog.* **11**,

e1005129 (2015).

33. Nguyen, D. D. *et al.* Indexing the Pseudomonas specialized metabolome enabled the

discovery of poaeamide B and the bananamides. *Nat Microbiol* **2**, 16197 (2016).

34. Choi, K.-H., Lee, H., Lee, S., Kim, S. & Yoon, Y. Cheese Microbial Risk Assessments - A

Review. *Asian-australas. J. Anim. Sci.* **29**, 307–314 (2016).

35. Perrin, F. *et al.* Quantitative Risk Assessment of Haemolytic and Uremic Syndrome Linked

to O157:H7 and Non-O157:H7 Shiga-Toxin Producing Escherichia coli Strains in Raw Milk

Soft Cheeses : Quantitative Risk Assessment of HUS Linked to Pathogenic STEC in

Cheese. *Risk Anal.* **35**, 109–128 (2015).

36. Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed

987  models. *Bioinformatics* **19**, 1572–1574 (2003).

988 37. Jukes, T. H. & Cantor, C. R. Evolution of protein molecules. *Mammalian protein*

989  *metabolism* (1969).

990 38. Cosetta, C. M. & Wolfe, B. E. Deconstructing and Reconstructing Cheese Rind

991  Microbiomes for Experiments in Microbial Ecology and Evolution. *Curr. Protoc. Microbiol.*

992  **56**, e95 (2020).

993 39. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of

994  intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

995 40. Farris, C., Sanowar, S., Bader, M. W., Pfuetzner, R. & Miller, S. I. Antimicrobial peptides

996  activate the Rcs regulon through the outer membrane lipoprotein RcsF. *J. Bacteriol.* **192**,

997  4894–4903 (2010).

998 41. Laubacher, M. E. & Ades, S. E. The Rcs phosphorelay is a cell envelope stress response

999  activated by peptidoglycan stress and contributes to intrinsic antibiotic resistance. *J.*

1000  *Bacteriol.* **190**, 2065–2074 (2008).

1001 42. Majdalani, N., Heck, M., Stout, V. & Gottesman, S. Role of RcsF in signaling to the Rcs

1002  phosphorelay pathway in Escherichia coli. *J. Bacteriol.* **187**, 6770–6778 (2005).

1003 43. Yethon, J. A., Heinrichs, D. E., Monteiro, M. A., Perry, M. B. & Whitfield, C. Involvement

1004  of waaY, waaQ, and waaP in the Modification of Escherichia coli Lipopolysaccharide and

1005  Their Role in the Formation of a Stable Outer Membrane. *J. Biol. Chem.* **273**, 26310–26316

1006  (1998).

1007 44. Bociek, K. *et al.* Lipopolysaccharide Phosphorylation by the WaaY Kinase Affects the

1008  Susceptibility of Escherichia coli to the Human Antimicrobial Peptide LL-37. *J. Biol. Chem.*

1009  **290**, 19933–19941 (2015).

45. Lofton, H., Pränting, M., Thulin, E. & Andersson, D. I. Mechanisms and fitness costs of

resistance to antimicrobial peptides LL-37, CNY100HL and wheat germ histones. *PLoS One*

8, e68875 (2013).

46. Nonejuie, P., Burkart, M., Pogliano, K. & Pogliano, J. Bacterial cytological profiling rapidly

identifies the cellular pathways targeted by antibacterial molecules. *Proc. Natl. Acad. Sci. U.*

S. A. **110**, 16169–16174 (2013).

47. Laich, F., Fierro, F. & Martín, J. F. Production of penicillin by fungi growing on food

products: identification of a complete penicillin gene cluster in Penicillium griseofulvum

and a truncated cluster in Penicillium verrucosum. *Appl. Environ. Microbiol.* **68**, 1211–1219

(2002).

48. Staggs, C. G., Sealey, W. M., McCabe, B. J., Teague, A. M. & Mock, D. M. Determination

of the biotin content of select foods using accurate and sensitive HPLC/avidin binding. *J.*

Food Compost. Anal. **17**, 767–776 (2004).

49. Streit, W. R. & Entcheva, P. Biotin in microbes, the genes involved in its biosynthesis, its

biochemical role and perspectives for biotechnological production. *Appl. Microbiol.*

Biotechnol. **61**, 21–31 (2003).

50. Hall, C. & Dietrich, F. S. The reacquisition of biotin prototrophy in Saccharomyces

cerevisiae involved horizontal gene transfer, gene duplication and gene clustering. *Genetics*

177, 2293–2307 (2007).

51. Bonham, K. S., Wolfe, B. E. & Dutton, R. J. Extensive horizontal gene transfer in cheese-

associated bacteria. *Elife* **6**, (2017).

52. Kastman, E. K. *et al.* Biotic Interactions Shape the Ecological Distributions of

Staphylococcus Species. *MBio* **7**, (2016).

53. Monnet, C., Back, A. & Irlinger, F. Growth of aerobic ripening bacteria at the cheese

surface is limited by the availability of iron. *Appl. Environ. Microbiol.* **78**, 3185–3192

(2012).

54. Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining

pipeline. *Nucleic Acids Res.* **47**, W81–W87 (2019).

55. Hantke, K. Identification of an iron uptake system specific for coprogen and rhodotorulic

acid in Escherichia coli K12. *Mol. Gen. Genet.* **191**, 301–306 (1983).

56. Fecker, L. & Braun, V. Cloning and expression of the fhu genes involved in iron(III)-

hydroxamate uptake by Escherichia coli. *J. Bacteriol.* **156**, 1301–1314 (1983).

57. Sauer, M., Hantke, K. & Braun, V. Ferric-coprogen receptor FhuE of Escherichia coli:

processing and sequence common to all TonB-dependent outer membrane receptor proteins.

J. Bacteriol. **169**, 2044–2049 (1987).

58. Matzanke, B. F., Anemüller, S., Schünemann, V., Trautwein, A. X. & Hantke, K. FhuF, part

of a siderophore-reductase system. *Biochemistry* **43**, 1386–1392 (2004).

59. Pain, A. *et al.* Insight into the genome of Aspergillus fumigatus: analysis of a 922kb region

encompassing the nitrate assimilation gene cluster. *Fungal Genet. Biol.* **41**, 443–453 (2004).

60. Chryssanthou, E., Broberger, U. & Petrini, B. Malassezia pachydermatis fungaemia in a

neonatal intensive care unit. *Acta Paediatr.* **90**, 323–327 (2007).

61. Prohic, A., Jovovic Sadikovic, T., Krupalija-Fazlic, M. & Kuskunovic-Vlahovljak, S.

Malassezia species in healthy skin and in dermatological conditions. *Int. J. Dermatol.* **55**,

494–504 (2016).

62. Haas, H. Fungal siderophore metabolism with a focus on Aspergillus fumigatus. *Nat. Prod.*

Rep. **31**, 1266–1276 (2014).

63. Park, M., Cho, Y.-J., Lee, Y. W. & Jung, W. H. Understanding the Mechanism of Action of

the Anti-Dandruff Agent Zinc Pyrithione against Malassezia restricta. *Sci. Rep.* **8**, 12086

(2018).

64. Gründlinger, M. *et al.* Fungal siderophore biosynthesis is partially localized in peroxisomes.

Mol. Microbiol. **88**, 862–875 (2013).

65. Wu, G. *et al.* Genus-Wide Comparative Genomics of Malassezia Delineates Its Phylogeny,

Physiology, and Niche Adaptation on Human Skin. *PLoS Genet.* **11**, e1005614 (2015).

66. Triana, S. *et al.* Draft Genome Sequence of the Animal and Human Pathogen Malassezia

pachydermatis Strain CBS 1879. *Genome Announc.* **3**, (2015).

67. Perrin, R. M. *et al.* Transcriptional regulation of chemical diversity in Aspergillus fumigatus

by LaeA. *PLoS Pathog.* **3**, e50 (2007).

68. Bok, J. W. & Keller, N. P. LaeA, a regulator of secondary metabolism in Aspergillus spp.

Eukaryot. Cell **3**, 527–535 (2004).

69. Kosalková, K. *et al.* The global regulator LaeA controls penicillin biosynthesis,

pigmentation and sporulation, but not roquefortine C synthesis in Penicillium chrysogenum.

Biochimie **91**, 214–225 (2009).

70. Bräse, S., Encinas, A., Keck, J. & Nising, C. F. Chemistry and biology of mycotoxins and

related fungal metabolites. *Chem. Rev.* **109**, 3903–3990 (2009).

71. Luckner, M. [On the synthesis of quinoline alkaloids in plants. 2. Fermentative conversion

of the penicillin alkaloids cyclopenin and cyclopenol to viridicatin and viridicatol. *Eur. J.*

Biochem. **2**, 74–78 (1967).

72. Ma, Y.-M. *et al.* A new isoquinolone alkaloid from an endophytic fungus R22 of Nerium

indicum. *Nat. Prod. Res.* **31**, 951–958 (2017).

73. Dalsgaard, P. W. *et al.* Atlantinone A, a Meroterpenoid Produced by Penicillium ribeum and

Several Cheese Associated Penicillium Species. *Metabolites* **2**, 214–220 (2012).

74. Wang, X. *et al.* Chemical epigenetics alters the secondary metabolite composition of guttate

excreted by an atlantic-forest-soil-derived Penicillium citreonigrum. *J. Nat. Prod.* **73**, 942–

948 (2010).

75. Tomoda, H. *et al.* Pyripyropenes, novel ACAT inhibitors produced by Aspergillus

fumigatus. IV. Structure elucidation of pyripyropenes M to R. *J. Antibiot.* **49**, 292–298

(1996).

76. Patti, G. J. *et al.* A view from above: cloud plots to visualize global metabolomic data. *Anal.*

Chem. **85**, 798–804 (2013).

77. Peters, B. M., Jabra-Rizk, M. A., O'May, G. A., Costerton, J. W. & Shirtliff, M. E.

Polymicrobial interactions: impact on pathogenesis and human disease. *Clin. Microbiol.*

Rev. **25**, 193–213 (2012).

78. Shirtliff, M. E., Peters, B. M. & Jabra-Rizk, M. A. Cross-kingdom interactions: Candida

albicans and bacteria. *FEMS Microbiol. Lett.* **299**, 1–8 (2009).

79. Scherlach, K., Graupner, K. & Hertweck, C. Molecular bacteria-fungi interactions: effects

on environment, food, and medicine. *Annu. Rev. Microbiol.* **67**, 375–397 (2013).

80. Wargo, M. J. & Hogan, D. A. Fungal—bacterial interactions: a mixed bag of mingling

microbes. *Curr. Opin. Microbiol.* **9**, 359–364 (2006/8).

81. Leclair, L. W. & Hogan, D. A. Mixed bacterial-fungal infections in the CF respiratory tract.

Med. Mycol. **48 Suppl 1**, S125–32 (2010).

82. Kim, D. *et al.* Candida albicans stimulates Streptococcus mutans microcolony development

via cross-kingdom biofilm-derived metabolites. *Sci. Rep.* **7**, 41332 (2017).

1102 83. de Boer, W. & Folman, L. B. Living in a fungal world: impact of fungi on soil bacterial

1103   niche development. *FEMS microbiology* (2005).

1104 84. Johansson, J. F., Paul, L. R. & Finlay, R. D. Microbial interactions in the mycorrhizosphere

1105   and their significance for sustainable agriculture. *FEMS Microbiol. Ecol.* **48**, 1–13 (2004).

1106 85. Tarkka, M. T., Sarniguet, A. & Frey-Klett, P. Inter-kingdom encounters: recent advances in

1107   molecular bacterium-fungus interactions. *Curr. Genet.* **55**, 233–243 (2009).

1108 86. Moree, W. J. *et al.* Interkingdom metabolic transformations captured by microbial imaging

1109   mass spectrometry. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 13811–13816 (2012).

1110 87. Clancy, A. *et al.* Evidence for siderophore-dependent iron acquisition in group B

1111   streptococcus. *Mol. Microbiol.* **59**, 707–721 (2006).

1112 88. Jin, B. *et al.* Iron acquisition systems for ferric hydroxamates, haemin and haemoglobin in

1113   Listeria monocytogenes. *Mol. Microbiol.* **59**, 1185–1198 (2006).

1114 89. Sheldon, J. R. & Heinrichs, D. E. Recent developments in understanding the iron acquisition

1115   strategies of gram positive pathogens. *FEMS Microbiol. Rev.* **39**, 592–630 (2015).

1116 90. Arias, A. A. *et al.* Growth of desferrioxamine-deficient Streptomyces mutants through

1117   xenosiderophore piracy of airborne fungal contaminations. *FEMS Microbiol. Ecol.* **91**,

1118   (2015).

1119 91. Mishra, R. P. N. *et al.* Staphylococcus aureus FhuD2 is involved in the early phase of

1120   staphylococcal dissemination and generates protective immunity in mice. *J. Infect. Dis.* **206**,

1121   1041–1049 (2012).

1122 92. Li, H. *et al.* The outer mucus layer hosts a distinct intestinal microbial niche. *Nat. Commun.*

1123   **6**, 8292 (2015).

1124 93. Rocha, E. R. & Krykunivsky, A. S. Anaerobic utilization of Fe(III)-xenosiderophores

among Bacteroides species and the distinct assimilation of Fe(III)-ferrichrome by

Bacteroides fragilis within the genus. *Microbiologyopen* **6**, (2017).

94. Emri, T. *et al.* Towards high-siderophore-content foods: optimisation of coprogen

production in submerged cultures of Penicillium nalgiovense. *J. Sci. Food Agric.* **93**, 2221–

2228 (2013).

95. Ong, S. A. & Neilands, J. B. Siderophores in microbially processed cheese. *J. Agric. Food*

Chem. **27**, 990–995 (1979).

96. David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut microbiome. *Nature*

505, 559–563 (2014).

97. Rehner, S. A. & Samuels, G. J. Molecular systematics of the Hypocreales: a teleomorph

gene phylogeny and the status of their anamorphs. *Can. J. Bot.* **73**, 816–823 (1995).

98. Glass, N. L. & Donaldson, G. C. Development of primer sets designed for use with the PCR

to amplify conserved genes from filamentous ascomycetes. *Appl. Environ. Microbiol.* **61**,

1323–1330 (1995).

99. Dunnett, C. W. A Multiple Comparison Procedure for Comparing Several Treatments with a

Control. *J. Am. Stat. Assoc.* **50**, 1096–1121 (1955).

100.Wickham, H. ggplot2: Elegant Graphics for Data Analysis. (2009).

101.Cleary, J. L., Luu, G. T., Pierce, E. C., Dutton, R. J. & Sanchez, L. M. BLANKA: an

Algorithm for Blank Subtraction in Mass Spectrometry of Complex Biological Samples. *J.*

Am. Soc. Mass Spectrom. **30**, 1426–1434 (2019).

102.Mohimani, H. *et al.* Dereplication of microbial metabolites through database search of mass

spectra. *Nature Communications* vol. 9 (2018).

103.Mohimani, H. *et al.* Dereplication of peptidic natural products through database search of

mass spectra. *Nat. Chem. Biol.* **13**, 30–37 (2017).

104. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular

interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

105. Tang, Y., Horikoshi, M. & Li, W. ggfortify: Unified Interface to Visualize Statistical Result

of Popular R Packages. *The R Journal* vol. 8 (2016).

106. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.*

47, D309–D314 (2019).

107. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

108. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

109. Grenier, F., Matteau, D., Baby, V. & Rodrigue, S. Complete Genome Sequence of

Escherichia coli BW25113. *Genome Announc.* **2**, (2014).

110. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS Comput.*

Biol. **9**, e1003118 (2013).

111. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

112. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 6640–6645

(2000).

113. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

114. Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome assembly

from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

115. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection

and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

116. Min, B., Grigoriev, I. V. & Choi, I.-G. FunGAP: Fungal Genome Annotation Pipeline using

evidence-based gene model evaluation. *Bioinformatics* **33**, 2936–2937 (2017).

117. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

118. Lim, F. Y., Sanchez, J. F., Wang, C. C. C. & Keller, N. P. Toward awakening cryptic

secondary metabolite gene clusters in filamentous fungi. *Methods Enzymol.* **517**, 303–324

(2012).

119. Xie, C. *et al.* KOBAS 2.0: a web server for annotation and identification of enriched

pathways and diseases. *Nucleic Acids Res.* **39**, W316–22 (2011).

120. The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing

strong. *Nucleic Acids Res.* **47**, D330–D338 (2019).

121. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

Consortium. *Nat. Genet.* **25**, 25–29 (2000).

SUPPLEMENTAL FIGURE AND TABLE LEGENDS
1189
1190 **Supplementary Figure 1**: **Impacts of fungal species on bacterial growth after 7 days of co-**
1191 **culture on cheese curd agar, pH 7**. CFU: colony forming units. N=3, error bars show standard
1192 deviation and black point is the mean. * represents a significant difference in bacterial growth in
1193 the presence of the fungal partner relative to alone (two-sided Dunnett's test, p-value <.05).
1194
1195 **Supplementary Figure 2**: **Impacts of bacterial species on fungal growth after 7 days of co-**
1196 **culture on cheese curd agar, pH 7**. For filamentous fungi, spore counts were used as a proxy
1197 for fungal CFUs. N=3, error bars show standard deviation and black point is the mean. *
1198 represents a significant difference in fungal growth in the presence of the bacterial partner

1199 relative to alone (two-sided Dunnett's test, p-value <.05).

1200

Supplementary Figure 3: RB-TnSeq assay. Characterized pooled bacterial mutant libraries

were grown in a biofilm either alone or in a mixed biofilm with a fungal partner. After seven

days of growth, mutant abundances were compared to the starting library abundances for each

condition. Changes in barcode abundances were used to calculate gene fitness values. Genes

with fitness values that differed significantly between co-culture and alone conditions

(significant interaction fitness) were identified as potentially relevant to fungal interaction.

1207

1208 **Supplementary Figure 4: COG categories of genes with interaction fitness.** Number of genes

1209 with interaction fitness falling into each COG category for *E. coli* (left) or *P. psychrophila*

1210 (right).

1211

1212 **Supplementary Figure 5: Bacterial Cytological Profiling of △*tolC E.coli* treated with**

1213 **known antibiotic compounds on cheese curd agar.** DAPI dye stains DNA and FM4-64 dye

1214 stains bacterial membranes. SYTOX green stains nucleic acids but cannot penetrate live cells.

1215 Scale bars represent 2 μm.

1216

Supplementary Figure 6: **Siderophore production by filamentous molds.** Liquid CAS assay

was performed on filtered and concentrated fungal supernatants from three replicates grown in

liquid cheese for 12 days. Row A) 1-3: Liquid cheese control 4-6: *Penicillium* SAM3. Row B) 1-

3: *Debaryomyces* 135B 4-6: *Penicillium* #12. Row C) 1-3: *Candida* 135E. 4-6: *Penicillium*

RS17. Row D) 1-3: *Scopulariopsis* 165-5 Row E) 1-3: *Scopulariopsis* JB370. Row F) 1-3:

Fusarium 554A. % Siderophore units calculated as $[(A_r - A_s)/(A_r)]*100$, where A_r is the

absorbance of the cheese curd agar supernatant blank and A_s is the absorbance of the sample.

N=3, error bars show standard deviation and black point is the mean.

1225

1226 **Supplementary Figure 7**: **Fitness defect of *fep* mutants on iron-limiting CCA.** Visual assays

1227 of *E. coli* mutant growth spotted alone on CCA pH 7.

1228

Supplementary Figure 8. Deletion of *laeA* gene in *Penicillium* sp. str. #12. A. Schematic

representation of the genetic construct for *laeA* deletion in *Penicillium* sp. str. #12. The construct

is constituted of the *hph* gene conferring resistance to hygromycin. The positions of the

restriction enzyme cutting sites are shown on the map. B. Southern blot analyses of genomic

DNA from the WT and the Δ*laeA* strains. Ten micrograms of total DNA from each strain was

digested with the appropriate enzymes and subjected to Southern blot analysis using respectively

the 5' flank fragment (blue) and the 3'fragment (grey) as probes. The 1 Kb DNA ladder from

New England Biolabs was used to determine the size of the expected bands. The blot image was

cropped to place the confirmed mutant adjacent to the positive control. The transformants that

were confirmed to not have the correct insertion were not included in the figure.

1239

Supplementary Data 1: RB-TnSeq fitness values for *E. coli* grown with fungal partners compared to alone.

Supplementary Data 2: RB-TnSeq fitness values for *P. psychrophila* grown with fungal partners compared to alone.

Supplementary Data 3: Genes with significant interaction fitness for *E. coli* grown with fungal partners.

Supplementary Data 4: Genes with significant interaction fitness for *P. psychrophila* grown with fungal partners.

Supplementary Data 5: Intersection lists for *E. coli* genes with interaction fitness across all conditions.

Supplementary Data 6: Intersection lists for *P. psychrophila* genes with interaction fitness across all conditions.

Supplementary Data 7: Functional enrichment results for *E. coli* genes with interaction fitness across all conditions.

Supplementary Data 8: Functional enrichment results for *P. psychrophila* genes with interaction fitness across all conditions.

Supplementary Data 9: RNA-Seq differential expression analysis for *E. coli* grown with *Penicillium* sp. str. #12 (*E. coli* perspective, alone condition as reference).

Supplementary Data 10: Iron-related genes that are differentially expressed by *E. coli* when grown with *Penicillium* sp. str. #12 relative to *E. coli* growth alone.

Supplementary Data 11: Metabolite production by Δ*laeA* and WT *Penicillium* sp. str. #12. Δ*laeA* used as the reference condition.

Supplementary Data 12: RB-TnSeq fitness values for *E. coli* grown with Δ*laeA* or WT *Penicillium* sp. str. #12 compared to alone.

Supplementary Data 13: RNA-Seq differential expression analysis for Δ*laeA* vs. WT *Penicillium* sp. str. #12 (WT used as the reference condition).

Supplementary Data 14: Functional enrichment results for genes differentially expressed in Δ*laeA* vs. WT *Penicillium* sp. str. #12.

Supplementary Data 15: Overlap of *E. coli* or *P. psychrophila* genes with interaction fitness.

1285
1286

1287
Figure 1: Fungal interaction partners span the phylogenetic and morphological diversity of
the cheese ecosystem. Phylogenetic tree based on large subunit rRNA of the cheese fungi used
as interaction partners in this study. The tree was built using Bayesian phylogenetic inference
with MrBayes[36] and the Jukes and Cantor substitution model[37]. Branch labels display posterior
probability.

and Y values indicate gene fitness values in each condition (alone on x-axis versus grown with a fungal partner on y-axis), and the colored numbers in the lower right hand corner indicate how many genes have either positive (blue) or negative (orange) interaction fitness. **a,** *E. coli.* **b,** *P. psychrophila.*

1355
1356
Figure 3: Bacterial genes with differential fitness in the presence of fungi. **a,** Network of *E.*
coli (left) or *P. psychrophila* (right) genes with an interaction fitness based on RB-TnSeq. Each

orange node represents a fungal partner and is labeled as follows: Fungal partner (Number of genes with interaction fitness; Number of genes with interaction fitness unique to this condition). Each green node represents a bacterial gene. Green nodes are shaded by the number of fungal conditions in which this gene has an interaction fitness as shown in the legend below and are sized by average strength of interaction fitness across partners. **b,** UpSet[39] plots showing the overlaps, or intersections, of *E. coli* (left) or *P. psychrophila* (right) gene sets with interaction fitness across fungal partners. These UpSet plots are conceptually similar to Venn Diagrams. The connected circles indicate which fungal conditions are included in the intersection, and the size of the intersection (the number of genes that have an interaction fitness in all the highlighted conditions) is displayed in the main bar chart. Intersections <5 genes are not shown. For example, in the *E. coli* panel, 30 genes have an interaction fitness with all partners (all fungi circles are connected), while 22 other genes have an interaction fitness with *Penicillium* sp. str. #12 and with *Penicillium* sp. str. SAM3 (only *Penicillium* sp. str. #12 *and Penicillium* sp. str. SAM3 circles are connected). **c,** PCA of the raw fitness values for all *E. coli* (left) or *P. psychrophila* (right) genes with an interaction fitness in at least one fungal condition.

1405
1406
Figure 4: Functional analysis of the bacterial genes associated with interaction fitness.
Comparison of *E. coli* (left) or *P. psychrophila* (right) gene fitness values alone compared to
fitness values with a fungal partner, colored by COG category and sized by the conservation of
effect among fungal partners (1-8 fungal species). Genes discussed in the text are labeled with
the gene name.

1429
1430
Figure 5: Bacterial cytological profiling of *E. coli* grown in a mixed biofilm with *Penicillium* sp. str. #12 and *Penicillium* sp. str. SAM3 on CCA plates. The phenotype of *E. coli* grown with these fungi is similar to that seen when *E. coli* is exposed to antibiotics targeting cell wall biosynthesis. DAPI dye stains DNA and FM4-64 dye stains bacterial membranes. Scale bars represent 2 μm.

Figure 6: Utilization of fungal siderophores by *E. coli*. a, RB-TnSeq fitness values for *fep*
operon genes in alone or interactive conditions, showing an increase in fitness in the presence of
fungal species. NS- not significant. **b,** Colony forming units of WT and Δ*fep* mutants after 7 days
of 1:1 competitive growth on CCA. Competitions were performed either alone or with

Penicillium sp. str. #12 or *Scopulariopsis* sp. str. JB370. N=4, error bars show standard deviation. Asterisk indicates significantly different growth in the presence of a fungus relative to growth alone (two-sided two-sample equal variance t-test p-value < 0.05). **c,** *E. coli* iron-related genes upregulated in the presence of *Penicillium* sp. str. #12. Significance cutoff made at abs($\log_2$(fold-change)) >1.5 and adjusted p-value <0.05. **d,** Fungal siderophores identified by mass spectrometry. Inset on the left shows the node that represents the desferrichrome fragmentation pattern depicted while network on the right represents coprogen-related molecules. Coprogen B and ferrichrome were found by matching fragmentation patterns to library spectra. Both identifications were confirmed using retention time and fragmentation matching to a purchased standard. **e,** Visual assays of *Δfep* mutant growth with purified siderophores coprogen and ferrichrome. **f,** Visual assays of *E. coli* mutant growth at varying distances from pre-cultured cheese fungi, *A. fumigatus* (soil, human pathogen), and *M. pachydermatis* (skin commensal). Growth performed on CCA containing tetrazolium chloride (red growth indicator).

1508
Figure 7: Fungal metabolite production impacts bacterial-fungal interactions. a,
Differential expression of WT *Penicillium* sp. str. #12 relative to Δ*laeA* after three days of
growth on CCA. Labeled on the volcano plot are the number of genes with a $\log_2$FC of >1.5
(green) or <-1.5 (red) and adjusted p-value of <0.05. **b,** The metabolomics data analysis platform
XCMS[76] was used to compare features detected by LC-MS analyses of Δ*laeA* and WT
Penicillium sp. str. #12 extracts. Features of higher abundance in WT relative to Δ*laeA* are
depicted as green nodes on the top of the mirror plot and features of lower abundance in WT
relative to Δ*laeA* are depicted as red nodes on the bottom. Node radius is proportional to the fold
change of the detected features and color intensity is dependent on p-value. The graph displays
only those features with a p-value less than or equal to 0.05, fold change higher than or equal to
10, *m/z* between 200 and 2000 Da, and intensity higher than 500. **c,** *E. coli* genes with significant
interaction fitness with Δ*laeA* and WT *Penicillium* sp. str. #12. Each dot represents a gene, with
colored dots indicating genes with interaction fitness. X and Y values (alone on x-axis and
+fungal partner on y-axis) indicate gene fitness values in each condition, and the numbers in the
lower right hand corner indicate how many genes have either positive (blue) or negative (orange)
interaction fitness. Venn diagrams display the overlap of these gene sets. **d,** Liquid CAS assay of
supernatants from blank control CCA medium, Δ*laeA,* or WT *Penicillium* sp. str. #12. N=3, error
bars show standard deviation from the mean. Asterisk indicates significantly different
siderophore production (two-sided two-sample equal variance t-test p-value 0.009).